

This is a repository copy of *Summary vision screening data*: *Albania*.

White Rose Research Online URL for this paper: http://eprints.whiterose.ac.uk/147025/

Version: Published Version

Monograph:

Mazzone, P. orcid.org/0000-0003-0944-8031, Griffiths, H. orcid.org/0000-0003-4286-5371 and Carlton, J. orcid.org/0000-0002-9373-7663 (2018) Summary vision screening data: Albania. Report. Vision Screening Country Reports . EUScreen

©2019 EUScreen. For reuse permissions, please contact the publisher.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.









Summary Vision Screening Data: Albania

Produced as part of Work Package 3

Paolo Mazzone¹, Dr Helen Griffiths², Dr Jill Carlton³

- 1. Research Assistant, School of Health and Related Research, University of Sheffield, United Kingdom (UK)
- 2. Senior Research Fellow, School of Health and Related Research, University of Sheffield, United Kingdom (UK)
- 3. Senior Lecturer, Academic Unit of Ophthalmology and Orthoptics, University of Sheffield, United Kingdom (UK)

Information provided by Dr. Alketa Tandili (University of Medicine, Tirana)

20th December 2018

Disclaimer: This is a summary report representing the responses from a country representative working within eye care services of the country reported. This report does not represent conclusions made by the authors, and is the product of professional research conducted for the EUSCREEN study. It is not meant to represent the position or opinions of the EUSCREEN study or its Partners. The information cannot be fully verified by the authors and represent only the information supplied by the country representatives.

This project has received funding from the European Union's Horizon 2020 research and innovation programme under Grant Agreement No 733352





Contents

Summary Vision Screening Data: Albania

1	Glossary of Terms: Vision Screening				
2	2 Abbreviations				
3	Population and Healthcare Overview				
4	V	ision Screening Commissioning and Guidance	3		
5	So	creening programme	4		
	5.1	Vision screening - Preterm babies	4		
	5.2	Vision screening - Birth to 3 months	4		
	5.3	Vision screening - 3 months to 36 months	4		
	5.4	Vision screening - 36 months to 7 years	4		
6	Α	utomated Screening	8		
7	Provision for Visually Impaired				
8	K	nowledge of existing screening programme	10		
	8.1	Prevalence/Diagnosis	10		
	8.2	Coverage	10		
	8.3	Screening Evaluation	10		
	8.3	Treatment success	10		
9	C	osts of vision screening in children	11		
	9.1	Cost of vision screening	11		
	9.2	Cost of treatment for amblyopia	11		
	9.3	Cost of Treatment for strabismus	11		
	9.4	Cost of treatment for cataract	11		
1	0	References	12		





1 Glossary of Terms: Vision Screening

Abnormal test	A test result where a normal "pass" response could not be		
result	detected under good conditions. The result on screening		
	equipment may indicate "no response," "fail," or "refer."		
Attendance rate	The proportion of all those invited for screening that are tested and receive a result: • Invited for screening includes all those that are offered the screening test. • Tested and receive a result could be a "pass" or "referral to		
	diagnostic assessment". Attendance rate provides information on the willingness of families to participate in screening.		
Compliance with referral (percentage)	The percentage of those who are referred from screening to a diagnostic assessment that actually attend the diagnostic assessment.		
	Percentage of compliance provides information on the willingness of families to attend the diagnostic assessment after referral from screening.		
Coverage	The proportion of those eligible for screening that are tested and receive a result:		
	 Eligible for screening includes those within the population that are covered under the screening or health care programme. Tested and receive a result could be a "pass" or "refer to diagnostic assessment". 		
	Factors such as being offered screening, willingness to participate, missed screening, ability to complete the screen, and ability to document the screening results will influence the coverage.		
False negatives	The percentage of children with a visual deficit (defined by the target condition) that receive a result of "pass" during screening. Example: If 100 children with visual deficit are screened, and 1 child passes the screening, the percentage of false negatives is 1%.		
False positives	The percentage of children with normal vision that are referred from screening to a diagnostic assessment.		
Guidelines	Recommendations or instructions provided by an authoritative body on the practice of screening in the country or region.		





Vision screening professional	A person qualified to perform vision screening, according to the practice in the country or region.
Inconclusive test result	A test result where a normal "pass" response could not be detected due to poor test conditions or poor cooperation of the child.
Invited for screening	Infants/children and their families who are offered screening.
Outcome of vision screening	An indication of the effectiveness or performance of screening, such as a measurement of coverage rate, referral rate, number of children detected, etc.
Untreated amblyopia	Those children who have not received treatment for amblyopia due to missed screening or missed follow-up appointment.
Persistent amblyopia	Amblyopia that is missed by screening, or present after the child has received treatment.
Positive predictive value	The percentage of children referred from screening who have a confirmed vision loss.
	For example, if 100 babies are referred from screening for diagnostic assessment and 10 have normal vision and 90 have a confirmed visual defect, the positive predictive value would be 90%.
Prevalence	The percentage or number of individuals with a specific disease or condition. Prevalence can either be expressed as a percentage or as a number out of 1000 individuals within the same demographic.
Programme	An organised system for screening, which could be based nationally, regionally or locally.
Protocol	Documented procedure or sequence for screening, which could include which tests are performed, when tests are performed, procedures for passing and referring, and so forth.
Quality assurance	A method for checking and ensuring that screening is functioning adequately and meeting set goals and benchmarks.
Referral criteria	A pre-determined cut-off boundary for when a child should be retested or seen for a diagnostic assessment.
Risk babies / Babies at-risk	All infants that are considered to be at-risk or have risk-factors for vision defects/ophthalmic pathology according to the screening programme.
	Two common risk factors are admission to the neonatal-intensive care unit (NICU) or born prematurely. However, other risk factors for visual defects may also be indicated in the screening programme.







Sensitivity	The percentage of children with visual defects that are identified via the screening program. For example, if 100 babies with visual defects are tested, and 98 of these babies are referred for diagnostic assessment and 2 pass the screening, the sensitivity is 98%.
Specificity	The percentage of children with normal vision that pass the screening. For example, if 100 babies with normal vision are tested, and 10 of these babies are referred for diagnostic assessment and 90 pass the screening, the specificity is 90%.
Target condition	The visual defect you are aiming to detect via the screening programme.
Well, healthy babies	Infants who are <i>not</i> admitted into the NICU or born prematurely (born after a gestation period of less than 37 weeks).





Summary Vision Screening Data: Albania

2 Abbreviations

AAO – American Academy of Ophthalmology

GDP – Gross Domestic Product

ICO – International Council of Ophthalmology

NICU – Neonatal-intensive care unit

PPP – Purchasing Power Parity

WHO – Word Health Organisation





3 Population and Healthcare Overview

The population of Albania is estimated at 2,800,000 (World Bank, 2018a) and a birth rate estimated at 12 births/1,000 population (World Bank, 2018b). The change in population and birth rate from 1960 to 2017 is shown in Figure 1, graphs A and B respectively.


Albania had a reported population density of 105 people per square kilometre in 2017 and this has risen from 61 people per square kilometre in 1961 (World Bank, 2018c). In terms of healthcare facilities, the total density of hospitals in 2013 was 0.72 per 100,000 population (WHO, 2016a). Infant mortality in 2017 is estimated at 8 deaths/1,000 live births (World Bank 2018d).


The average life expectancy in Albania is estimated at 78 years (World Bank, 2018e), with a death rate of 7 deaths/1,000 population in 2016 (World Bank, 2018f). Albania has a gross national income per capita (PPP int. \$, 2013) of \$10,000 (WHO, 2016b). The estimated total expenditure on health per capita in 2014 was \$615 (Intl \$) and the total expenditure on health in 2014 as percentage of GDP was 5.9% (WHO, 2016b).





Figure 1: Change in the Total Population and Birth Rate in Albania between 1960 and 2017





Source: Information sourced from World Bank (2018)





4 Vision Screening Commissioning and Guidance

Vision screening in Albania is funded through national health insurance and is not embedded into general preventative child healthcare screening. There are no guidelines for ophthalmologists to follow however, they are guided by the American Academy of Ophthalmology (AAO) and International Council of Ophthalmology (ICO) guidelines. The content of the Albanian vision screening programme is decided upon by the Ministry of Health and was implemented nationally in 2013, with no changes since its commencement.

Vision screening is conducted by ophthalmologists, nurses, or paediatricians. No other general health care professionals have been identified that do not currently screen, but could do so with additional training. There is no specific training provided for vision screening; ophthalmologists undertake a 4-year residency where they acquire the skills and knowledge to perform such tasks.

Vision screening is conducted in hospitals or primary service. It is not known how often the vision screening programme is reviewed, who decides upon revisions, how revisions are funded, or how the revisions take place.

There are no methods imposed by the government regarding quality monitoring of the vision screening programme, but there has been some PhD research concerning the programme, the details of this research have not been shared and therefore the content is not known. There has been no cost-effectiveness analysis conducted pertaining to the vision screening programme, nor studies on the effectiveness of vision screening in Albania.





5 Screening programme

In Albania, retinopathy of prematurity, congenital eye defects and amblyopia are the target conditions screened for by vision screening. The health care professionals delivering vision screening, venue for screening and tests used vary depending on the age of the child as shown in Tables 1, 2 and 3 respectively. Specific details of the screening offered within each age group are described more fully in sections 5.1 to 5.4 below.

5.1 Vision screening - Preterm babies

Preterm babies up to the age of 3 months are screened using an eye inspection. This vision screening is conducted by an ophthalmologist in either a maternity clinic or an eye clinic.

5.2 Vision screening - Birth to 3 months

Well, healthy babies up to the age of 3 months are screened using an eye inspection and red reflex examination. These tests are conducted by an ophthalmologist or a nurse in either a hospital or primary service. Paediatricians or ophthalmologists perform the fundus red reflex examination to diagnose white pupil. The timing and number of eye screenings offered during this 3 month period is not clearly defined and it is not known after how many abnormal or inconclusive tests a baby is referred for further diagnostic examination.

5.3 Vision screening - 3 months to 36 months

Vision screening is conducted on children aged 3 months to 36 months by an ophthalmologist at an eye clinic. It is not clear what vision screening tests are performed in children aged 3 months to 36 months. It is known that a visual acuity measurement is conducted at the age of 2 years, although the method of measurement is not clear and the sequence of testing during this period or the referral criteria are not known.

5.4 Vision screening - 36 months to 7 years

Children aged 36 months to 7 years are screened by either an ophthalmologist or a paediatrician in a primary care clinic, using an eye inspection and an unspecified 'other' test. The timing and frequency of vision screening at any other age between 36 months and 7 years is not known and it is not known after how many abnormal or inconclusive tests a child is referred for further diagnostic examination. Visual acuity is measured at age 3 to 4 years by an ophthalmologist, using Lea Symbols, HOTV, Keeler, Snellen, Allen Cards or Numbers. Visual acuity is repeated at ages 6 to 7 years. It is not known how many abnormal or inconclusive tests necessitate a child to be referred for further diagnostic examination.





 Table 1: Healthcare professionals who conduct vision screening in each age group

Table 1	Paediatrician	Ophthalmologist	Nurse
Preterm babies	×	✓	×
0 to 3 months	✓	✓	√
3 to 36 months	×	✓	×
3 to 7 years	✓	✓	×





 Table 2: Vision screening tests used in vision screening for each age group

Table 2	Eye inspection	Red Reflex	Visual Acuity Measurement	Other
Preterm babies	√	×	×	×
0 to 3 months	✓	✓	×	×
3 to 36 months	×	×	~	×
3 to 7 years	×	×	✓	~





 Table 3: Location of vision screening for each age group

Table 3	Maternity	Eye Clinic	Hospital	Primary Care Clinic
Preterm babies	✓	✓	×	×
0 to 3 months	×	×	√	√
3 to 36 months	×	✓	×	×
3 to 7 years	×	×	×	✓





6 Automated Screening

Automated vision screening is achieved using handheld, portable devices designed to detect presence of refractive error in infants from 6 months of age. It provides objective results and is used to detect amblyopic risk factors. This differs from other methods used to screen children for amblyopia which focus on detection of the actual condition and the resulting visual loss.

Automated vision screening is used in Albania, but only in the private sector (therefore not every child who attends vision screening has automated vision screening). The content of this private sector vision screening is not clear. These devices cost approximately 5,000 Euros, however, the maintenance costs and the number of years before a device is scheduled to be replaced are not known.





7 Provision for Visually Impaired

In Albania, there is one school for blind children, it is located in Tirana and accommodates around 60 pupils. There are specific lessons concerning Braille writing and this school is supported by the Albanian government. The costs per child for these schools and whether or not there is any special support for visually impaired children who attend regular primary school is not known.





8 Knowledge of existing screening programme

8.1 Prevalence/Diagnosis

There is no information provided pertaining to the prevalence and diagnosis of vision disorders in children aged up to 7 years.

8.2 Coverage

All children commencing school at the age of 6 years are invited and must attend vision screening. There is no information provided relating to the coverage and treatment success of vision disorders in children aged up to 7 years. The coverage and subsequent attendance rates of vision screening are not known.

8.3 Screening Evaluation

There is no information provided pertaining to the screening outcomes of vision disorders in children aged up to 7 years.

8.3 Treatment success

Ophthalmologists prescribe glasses to children under the age of 7 years; other treatments include patching and cataract surgery. All eligible children are offered treatment. There is no information provided pertaining to treatment outcomes.





9 Costs of vision screening in children

In Albania, vision screening is free of charge for parents and there is no financial reward for attending and no penalty for not attending a vision screening appointment.

9.1 Cost of vision screening

There is no information provided relating to the costs of performing vision screening in children aged up to 7 years.

9.2 Cost of treatment for amblyopia

There is not information provided relating to the costs of treatment for amblyopia in children aged up to 7 years.

9.3 Cost of Treatment for strabismus

There is not information provided relating to the costs of treatment for strabismus in children aged up to 7 years.

9.4 Cost of treatment for cataract

There is not information provided relating to the costs of treatment for cataract in children aged up to 7 years.





10 References

The World Bank (2018a). *Population, total | Data*. [online] Available at: https://data.worldbank.org/indicator/SP.POP.TOTL [Accessed 26 Nov. 2018].

The World Bank. (2018b). *Birth rate, crude (per 1,000 people) | Data*. [online] Available at: https://data.worldbank.org/indicator/SP.DYN.CBRT.IN [Accessed 26 Nov. 2018].

The World Bank. (2018c). *Population density (people per sq. km of land area) | Data*. [online] Available at: https://data.worldbank.org/indicator/EN.POP.DNST [Accessed 26 Nov. 2018].

The World Bank. (2018d). *Mortality rate, infant (per 1,000 live births) | Data*. [online] Available at: https://data.worldbank.org/indicator/SP.DYN.IMRT.IN [Accessed 26 Nov. 2018].

The World Bank. (2018e). *Life expectancy at birth, total (years) | Data*. [online] Available at: https://data.worldbank.org/indicator/SP.DYN.LE00.IN [Accessed 26 Nov. 2018].

The World Bank. (2018f). *Death rate, crude (per 1,000 people) | Data*. [online] Available at: https://data.worldbank.org/indicator/SP.DYN.CDRT.IN [Accessed 26 Nov. 2018].

World Health Organisation (WHO). (2016a). Health Infrastructure - Data by country. [ONLINE] Available at: http://apps.who.int/gho/data/view.main.30000. [Accessed 26 November 2018].

World Health Organisation (WHO). (2016b). Countries, Albania. [ONLINE] Available at: http://www.who.int/countries/alb/en/. [Accessed 26 November 2018].