

Maximum cardiac performance of Antarctic fishes that lack haemoglobin and myoglobin: exploring the effect of warming on nature's natural knockouts

Stuart Egginton¹ , Michael Axelsson², Elizabeth L. Crockett³, Kristin M. O'Brien⁴ and Anthony P. Farrell^{5,*}

¹School of Biomedical Sciences, University of Leeds, Leeds LS2 9JT, UK.

²Department of Biological and Environmental Sciences, University of Gothenburg, Gothenburg SE-405 30, Sweden.

³Department of Biological Sciences, Ohio University, Athens, OH OH45701, USA.

⁴Institute of Arctic Biology, University of Alaska Fairbanks, Fairbanks, AK AK99775, USA.

⁵Department of Zoology, University of British Columbia, Vancouver, BC V6T 124, Canada.

*Corresponding author: Department of Zoology, University of British Columbia, Vancouver, BC, Canada. Email: tony.farrell@ubc.ca

Antarctic notothenioids, some of which lack myoglobin (Mb) and/or haemoglobin (Hb), are considered extremely stenothermal, which raises conservation concerns since Polar regions are warming at unprecedented rates. Without reliable estimates of maximum cardiac output (\dot{Q}), it is impossible to assess their physiological scope in response to warming seas. Therefore, we compared cardiac performance of two icefish species, *Chionodraco rastrospinosus* (Hb⁻Mb⁺) and *Chaenocephalus aceratus* (Hb⁻Mb⁻), with a related notothenioid, *Notothenia coriiceps* (Hb⁺Mb⁺) using an *in situ* perfused heart preparation. The maximum \dot{Q} , heart rate (f_H), maximum cardiac work (W_C) and relative ventricular mass of *N. coriiceps* at 1°C were comparable to temperate-water teleosts, and acute warming to 4°C increased f_H and W_C , as expected. In contrast, icefish hearts accommodated a higher maximum stroke volume (V_S) and maximum \dot{Q} at 1°C, but their unusually large hearts had a lower f_H and maximum afterload tolerance than *N. coriiceps* at 1°C. Furthermore, maximum V_S , maximum \dot{Q} and f_H were all significantly higher for the Hb⁻Mb⁺ condition compared with the Hb⁻Mb⁻ condition, a potential selective advantage when coping with environmental warming. Like *N. coriiceps*, both icefish species increased f_H at 4°C. Acutely warming *C. aceratus* increased maximum \dot{Q} , while *C. rastrospinosus* (like *N. coriiceps*) held at 4°C for 1 week maintained maximum \dot{Q} when tested at 4°C. These experiments involving short-term warming should be followed up with long-term acclimation studies, since the maximum cardiac performance of these three Antarctic species studied seem to be tolerant of temperatures in excess of predictions associated with global warming.

Key words: Cardiac output, cardiac work, heart rate, icefishes, thermal adaptation, warming

Editor: Anne Todgham

Received 3 December 2018; Revised 18 June 2019; Editorial Decision 26 June 2019; Accepted 27 June 2019

Cite as: Egginton S, Axelsson M, Crockett EL, O'Brien KM, Farrell AP (2019) Maximum cardiac performance of Antarctic fishes that lack haemoglobin and myoglobin: exploring the effect of warming on nature's natural knockouts. *Conserv Physiol* 7(1): coz049; doi:10.1093/conphys/coz049.

Introduction

A remarkable trait among vertebrates is the evolutionary loss of haemoglobin (Hb) and myoglobin (Mb) expression within the Channichthyid family of Antarctic notothenioid fishes (Holeton 1970). Mutations leading to the Hb⁻ state occurred once during evolution of icefishes, ~8.5 MYA (Near, 2004), subsequent to cooling of the Southern Ocean. Loss of Mb expression occurred four times during radiation of the Channichthyids (Sidell *et al.*, 1997), resulting in 6 of the 16 species of icefishes lacking Mb in their heart (Sidell *et al.*, 1997). Thus, the Hb⁻Mb⁺ and Hb⁻Mb⁻ states are unique to icefishes (Sidell and O'Brien, 2006), although loss of Mb expression is a trait shared by a few temperate teleosts (Grove & Sidell, 2002; Macqueen *et al.*, 2014) whose hearts can function adequately provided blood PO₂ is sufficient (Bailey *et al.*, 1990).

The Hb⁻ and Mb⁻ condition is thought to have restricted Channichthyid distribution primarily to the thermally stable offshore waters of Antarctica, which hover near -1.8°C below the surface year-round. While cold water may be key to survival of stenothermal species by ensuring high levels of dissolved oxygen, and presumably low oxygen requirements, the Western Antarctic Peninsula region is warming at an unprecedented rate. Thus, endemic species possessing modest thermal plasticity (e.g. Bilyk *et al.*, 2018) may be especially vulnerable to short-term extreme temperature fluctuations, as well as long-term climate warming (Clarke *et al.*, 2007; Beers *et al.*, 2010). Given the limited options for altered geographic distribution, documenting scope for performance provides critical insight into capacity to accommodate shifts in environmental temperature, and hence the potential for longer-term adaptations necessary to allow stock conservation within the current ecological niche.

In terms of physiological performance, cardiac performance is viewed by some as playing a critical role in setting the upper thermal limits of temperate-zone fishes (Wang and Overgaard, 2007; Portner and Farrell, 2008; Farrell, 2009, 2016; Eliason *et al.*, 2011), though others have challenged this contention (Ern *et al.*, 2016). Indeed, extensive adaptational remodelling of the icefish cardiovascular system is apparent for life in the frigid, but stable Southern Ocean. For example, the 90% decrease in oxygen-carrying capacity associated with the Hb-free blood of icefishes (Ruud, 1954) has been compensated for by a thin-walled ventricle with relative mass 10-times larger than other benthic teleosts (Tota *et al.*, 1991; Farrell and Jones, 1992; Axelsson, 2005; Farrell and Smith, 2017) that can eject a very large stroke volume (V_S) and produce an exceptionally high cardiac output (\dot{Q}) for an animal living at ~0°C (Hemmingsen and Douglas, 1977). The concern, however, is that a very large routine \dot{Q} needed to offset a low blood oxygen-carrying capacity might leave little cardiac scope for activity or facing environmental challenges such as warming.

Earlier studies with icefishes have reported on the maximum \dot{Q} (an integrative measure limiting fish activity),

but its response to warming is poorly documented. For example, maximum \dot{Q} in *Chionodraco hamatus* (Hb⁻Mb⁺) is impressive at 3°C: ~300 ml min⁻¹ kg⁻¹ (Tota *et al.*, 1991; Acierno *et al.*, 1997). This maximum performance index is best measured with a working perfused heart given the difficulty of inducing a maximum performance of icefish *in vivo* (see Joyce *et al.*, 2018b). To determine cardiac scope also requires reliable routine \dot{Q} values, but these vary considerably, as pointed out by Joyce *et al.*, (2018b), who also reported routine \dot{Q} for *Chaenocephalus aceratus* (Hb⁻Mb⁻) at 0.8°C that was lower than previously reported and increased with both moderate activity and with acute warming. By comparison, routine \dot{Q} is just 6 ml min⁻¹ kg⁻¹ at ~0°C in *N. coriiceps* (Hb⁺Mb⁺), which has an haematocrit of ~16% (Egginton, 1997; Joyce *et al.*, 2018b), although other red-blooded notothenioids of different ecotype and size have a higher routine \dot{Q} (Axelsson, 2005). Moreover, *N. coriiceps* can increase routine oxygen uptake, \dot{Q} , V_S and f_H appreciably after warm-acclimation (Egginton and Campbell, 2016; Joyce *et al.*, 2018a). Other cardiovascular adaptations of icefishes include wide-bore arteries and capillaries that reduce overall systemic vascular resistance to just 10–20% that of Hb⁺ fishes, and lower arterial blood pressure (Egginton, 1997) and cardiac afterload (Egginton *et al.*, 2002). Thus, mass-specific cardiac work (W_C) in icefishes is estimated to range between 0.35 and 2.3 mW g⁻¹ (Axelsson, 2005), at the lower end of the range for teleost fishes (Farrell and Smith, 2017).

Therefore, indications exist that icefishes may have a reasonable cardiac scope and a moderate capacity to respond to warming, but without robust estimates of maximum \dot{Q} for an Hb⁻Mb⁻ icefish we cannot test these possibilities. Indeed, maximum cardiac performance has never been assessed in *C. aceratus*. Therefore, to better understand how loss of Mb and Hb expression might affect vulnerability to ocean warming, we hypothesized that warming impairs maximum cardiac performance of icefishes, especially when compared with the heart of a species possessing Hb and Mb. To test this hypothesis, we compared maximum cardiac performance at 1°C and 4°C for three Antarctic notothenioids that naturally express different levels of Hb and Mb. We reasoned that the loss of facilitated oxygen transport, either by systemic convection (Hb) or cellular diffusion (Mb), would limit maximum cardiac capacity. In comparing two icefish species that are Hb⁻ the influence of Mb would be revealed, and the influence of Hb evident by comparison with a red-blooded nototheniid.

Material and methods

Collection of animals

Adult *Chaenocephalus aceratus* (Hb⁻Mb⁻), *Chionodraco rastrospinosus* (Hb⁻Mb⁺) and *Notothenia coriiceps* (Hb⁺Mb⁺) were collected around the Western Antarctic Peninsula during the austral fall and winter of 2015 (see O'Brien *et al.*, 2018). They were held at Palmer Station in ambient seawater temperature (0 ± 1°C) for a maximum of 3 weeks before

experimentation. *N. coriiceps* were fed cubes of fish muscle every other day, but not 24–48 h prior to surgery. The icefishes did not feed in captivity; they are known to maintain a lower metabolic rate than other notothenioids and likely grow sporadically, feeding at relatively broad but seasonally dependent time intervals (Johnston and Battram, 1993; Campbell *et al.*, 2008).

The *in situ* heart preparation

We adopted a well-established *in situ* working, perfused heart preparation that has been used across a wide variety of fish species, including Arctic fish at low temperatures (e.g. Farrell *et al.*, 1988, 2013). This preparation retains full anatomical integrity of the heart and generates a maximum \dot{Q} equivalent to that observed *in vivo*. Output pressure can be raised to determine the maximum pressure-generating ability and maximum W_C (Farrell *et al.*, 1991). All experiments were approved by the University of Alaska, Fairbanks Institutional Animal Use and Care Committee (570217-18).

Briefly, anaesthetized fish (Egginton and Campbell, 2016) were maintained on a surgical operating sling by continuously irrigating the gills with ice-chilled seawater containing MS-222 (50 mg l⁻¹). Heparin (~100 i.u. kg⁻¹ in saline) was injected into the caudal vessels to prevent blood clots. A mid-line ventral incision, occlusion of the gut arterial blood supply with umbilical tape and removal of the intestines provided access to the hepatic veins, all but one of which were ligated. A saline-filled input cannula [connected to a Marriot bottle containing physiological perfusate (see below) bubbled with 100% oxygen] was advanced into the sinus venosus *via* the remaining vein and secured in place. Air bubbles sucked into the heart when this vein was cut were prevented from entering gill vessels (which would restrict \dot{Q}) by severing the ventral aorta at the isthmus and anterior to the pericardium prior to the cut. The Marriot bottle height was adjusted so that \dot{Q} was not excessive and the exceptionally thin-walled cardiac chambers were not over filled. Next, the output cannula was inserted into the severed ventral aorta and secured. The ducts of Cuvier were then occluded so that the heart only received perfusate; this procedure also crushed all nerves supplying the heart. Within ~20 min the heart preparation was transferred to and fully immersed in a saline bath (250 mM NaCl) housed inside a 4°C environmental room. The input cannula was then connected to a pressure head that delivered oxygenated perfusate at a precise cardiac filling pressure (P_i). The bath and the reservoirs containing oxygenated perfusate were maintained at a constant temperature (either 1°C or 4°C) with a Neslab RTE recirculating water bath (ThermoFisher Scientific, Waltham, MA, USA).

Initial \dot{Q} was nominally set at ~13 ml min⁻¹ kg⁻¹ body mass by adjusting P_i (typically < 1 cm H₂O and sometimes sub-ambient; 1 cm H₂O = 0.1 kPa) to set V_S while the heart beat with a myogenic rhythm. A constant pressure head device connected to the output cannula set the initial cardiac afterload (P_o) at a nominal ~40 cm H₂O for *N. coriiceps*

and ~20 cm H₂O for both icefish species. Mean dorsal aortic blood pressure is higher for *N. coriiceps* (18.7–42.8 cm H₂O) than *C. aceratus* (12.3–23.4 cm H₂O) (Egginton, 1997), but information on *in vivo* ventral aortic blood pressure to set their initial P_o is lacking. Regardless, maximum P_o was directly measured during the protocol by incremental increases until maximum \dot{Q} began to fail, suggesting this was a reasonable approach, especially since we repeated this response several times for each preparation. These stable perfusion conditions were maintained for at least 10 min prior to assessing maximum cardiac performance. The pericardium was opened at the end of the experiment to ensure proper placement of input and output cannulae and weigh the heart.

Assessing maximum cardiac performance *in situ*

Protocols to generate maximum V_S , maximum \dot{Q} , maximum P_o and a maximum cardiac work (W_C) are detailed elsewhere (e.g. Farrell *et al.*, 1988). Briefly, incremental increases in P_i with a myogenic heartbeat yielded a maximum V_S and a maximum \dot{Q} . All preparations gave a robust Starling response, similar to other teleost working heart preparations, with V_S being most sensitive to a P_i < 3 cm H₂O; smaller incremental changes in \dot{Q} occurred up to 5 cm H₂O and occasionally up to 8 cm H₂O. (Note: maximum V_S could be higher *in vivo* if cardiac vagal tonus lowered f_H , but maximum \dot{Q} would not necessarily be greater.) Maximum P_o was determined at maximum \dot{Q} by raising P_o in 5–10 cm H₂O increments until maximum W_C was attained, at which point any increase in P_o was matched by an equivalent decrease in \dot{Q} . While hearts can generate a higher P_o , but not without disproportionately decreasing \dot{Q} and W_C , icefishes displayed an interesting phenomenon when P_o was increased beyond the maximum W_C , wherein f_H abruptly halved, V_S doubled and maximum \dot{Q} hardly changed; this situation was immediately reversed by reducing P_o . After maximum cardiac performance had been assessed, routine \dot{Q} and P_o were rapidly restored, which quickly re-established a stable routine \dot{Q} .

Adrenaline can enhance maximum cardiac performance of perfused teleost hearts (Farrell *et al.*, 1986), including Arctic fish at 1°C (Farrell *et al.*, 2013). However, low circulating levels of catecholamines in Antarctic fishes, even after moderate stress that rose substantially only *in extremis* (Whiteley and Egginton, 1999), raised the possibility that no adrenergic stimulation would be needed in the present study. Nevertheless, preliminary experiments showed that *C. aceratus* preparations quickly lost pumping capacity unless they had been stabilized with 50 nM adrenaline in the perfusate, which is a low concentration when compared to the levels induced by stress (~5 μM). Accordingly, our initial perfusion conditions, including during surgery, were standardized to include 50 nM adrenaline. Thus, the initial f_H reported below reflects a tonic adrenergic effect on the myogenic heartbeat. Since the preliminary experiments with all three species showed that adrenaline concentrations > 5 μM elicited no further

stimulation of cardiac performance, maximum cardiac performance was assessed with three levels of catecholamine (50 nM, 0.5 μ M and 5 μ M) to provide insight into the modulatory effects of adrenergic stimulation. The need to test a range of concentrations to establish the effect of different levels of adrenergic stimulation reflects a lack of prior data in the literature on these species, so was an essential prerequisite for a robust experimental design. Up to five concentrations have been tested previously without performance deterioration.

Temperature treatments

The three species were assessed at 1°C, which was near their habitat temperature ($0 \pm 1^\circ\text{C}$). Attempts to more closely match test and exposure temperature in a 4°C environmental chamber produced unwanted ice formation. Acute warming to 4°C of heart preparations from fish acclimated to $0 \pm 1^\circ\text{C}$ were assessed with the experimental bath and perfusate reservoirs maintained at 4°C, but only with *C. aceratus* ($\text{Hb}^- \text{Mb}^-$) and *N. coriiceps* ($\text{Hb}^+ \text{Mb}^+$) because too few *C. rastrispinosus* ($\text{Hb}^- \text{Mb}^+$) were caught. In addition, hearts from both *N. coriiceps* and *C. rastrispinosus* were similarly tested at 4°C after an exposure period of a minimum of 5 days at $4 \pm 0.5^\circ\text{C}$ in 700 l insulated recirculating seawater tanks (4–6 fish per tank); fish were initially held at $0 \pm 1^\circ\text{C}$ for 24 h before increasing the temperature by 1°C daily using 3-kW Electro Titanium inline heaters (Aqualogic, San Diego, CA, USA). *C. aceratus*, which was intolerant of more than 1–2°C rise in temperature, proved difficult to acclimate to 4°C.

Measurements and analysis

Perfusate outflow was measured with an in-line Transonic probe (4.0 mm diameter) and flowmeter (T206, Transonic Systems, Ithaca, NY, USA), gravimetrically calibrated at 1°C and 4°C to adjust for temperature effects on factory-calibrated flow readings. P_i and P_o were measured near the input and output cannulae, respectively, via fluid-filled tubes connected to pressure transducers (DP6100, Peter von Berg, Medizintechnik, Germany) and referenced to saline level in the bath. Recorded pressures were corrected to provide true values using individual calibrations of cannula resistance. Flow and pressure signals were amplified (4CHAMP amplifier, Somedic, AB, Hörby, Sweden) and processed on-line (custom LabView program; National Instruments Sweden AB, Kista, Sweden) to display real time raw values of P_i , P_o , V_S , \dot{Q} and f_H (determined from the pulsatile flow trace) throughout the experiment. W_C , the product of [\dot{Q} (ml min^{-1}) \times ($P_o - P_i$) \times 0.00167] was displayed on-line and used to determine maximum performance when P_o was elevated. Data were analysed off-line for 5–10 stable, consecutive heartbeats for each incremental change in P_i and P_o . W_C was corrected for ventricular mass (mW g^{-1} ventricle mass), and \dot{Q} was corrected for body mass ($\text{ml min}^{-1} \text{kg}^{-1}$ body mass).

Statistical analyses

All data are presented as mean \pm standard error (s.e.m.), analysed using SPSS v.23 (SPSS Inc., Chicago, IL, USA). Intraspecific comparisons (temperature \times adrenaline), or interspecific comparisons for similar exposure and test temperatures (species \times temperature), used a two-way ANOVA with Tukey's Kramer *post-hoc* tests. A one-way Welch's ANOVA was used to compare differences between 1°C-acclimated animals tested at 1°C and 4°C, and between 4°C-exposed animals tested at 4°C; the dose-response to adrenaline was tested by one-way repeated measures ANOVA with Sidák *post-hoc* tests at each test temperature. Relative ventricular mass (RVM) was transformed by Lngamma function to achieve normality for statistical comparisons. Statistical significance was assigned to $P \leq 0.05$.

Chemicals

The physiological saline used for perfusate (pH 8.1 at 1°C) contained 250 mM NaCl, 2.5 mM KCl, 0.9 mM $\text{MgSO}_4 \cdot 6\text{H}_2\text{O}$, 2.5 mM $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 5.6 mM glucose, 3.9 mM TES free acid and 6.1 mM TES free base. Tricaine methane sulfonate (MS-222) and adrenaline bitartrate were from Sigma-Aldrich (St. Louis, USA).

Results

Morphometry

At ambient (holding) temperature, RVM (Table 1) was 30% larger in *C. rastrispinosus* ($\text{Hb}^- \text{Mb}^+$) than for *C. aceratus* ($\text{Hb}^- \text{Mb}^-$) ($P < 0.05$). RVM of both *C. aceratus* (3-fold) and *C. rastrispinosus* (4-fold) were significantly ($P < 0.05$) larger than *N. coriiceps* ($\text{Hb}^+ \text{Mb}^+$), which was more typical of temperate teleosts. RVM values for *C. rastrispinosus* and *N. coriiceps* did not change significantly after a period of exposure to 4°C.

Maximum cardiac performance tested at 1°C

C. aceratus ($\text{Hb}^- \text{Mb}^-$)

Initial f_H ($19.0 \pm 0.8 \text{ min}^{-1}$) was unresponsive to further adrenergic stimulation (Fig. 1), which improved some indices of maximum cardiac performance. For example, 5 μ M adrenaline generated a maximum V_S of $4.6 \pm 0.5 \text{ ml kg}^{-1}$, a maximum \dot{Q} of $80.2 \pm 11.6 \text{ ml min}^{-1} \text{kg}^{-1}$ and a maximum W_C of $1.3 \pm 0.3 \text{ mW g}^{-1}$. By comparison, maximum V_S ($3.8 \pm 0.4 \text{ ml kg}^{-1}$) with 50 nM adrenaline was significantly ($P < 0.05$) lower, but maximum \dot{Q} ($70.0 \pm 9.1 \text{ ml min}^{-1} \text{kg}^{-1}$) and maximum W_C ($1.1 \pm 0.3 \text{ mW g}^{-1}$) were unchanged.

C. rastrispinosus ($\text{Hb}^- \text{Mb}^+$)

Initial f_H ($26.0 \pm 0.5 \text{ min}^{-1}$) was also unresponsive to further adrenergic stimulation (Fig. 2) that, as with *C. aceratus*, improved some indices of maximum cardiac performance.

Table 1: Morphometric details of fish used for the perfused heart experiments (mean \pm s.e.m.)

	Acclim. temp ($^{\circ}$ C)	Test temp ($^{\circ}$ C)	N	Body mass (g)	Total length (cm)	Relative atrial mass (%BM)	Relative ventricular mass (%BM)
<i>C. aceratus</i>	1	1	8	1054 \pm 123 ^a	52 \pm 2 ^a	0.107 \pm 0.011 ^a	0.304 \pm 0.011 ^a
	1	4	7	676 \pm 62 ^b	44 \pm 1 ^b	0.077 \pm 0.005 ^a	0.253 \pm 0.011 ^a
<i>C. rastrispinosus</i>	1	1	6	478 \pm 57 ^b	40 \pm 3 ^b	0.148 \pm 0.013 ^b	0.395 \pm 0.029 ^b
	4	4	5	350 \pm 79 ^c	36 \pm 2 ^c	0.077 \pm 0.014 ^a	0.364 \pm 0.030 ^a
<i>N. coriiceps</i>	1	1	7	1093 \pm 87 ^a	40 \pm 1 ^b	0.027 \pm 0.002 ^c	0.098 \pm 0.003 ^c
	1	4	6	1102 \pm 142 ^a	41 \pm 2 ^b	0.024 \pm 0.002 ^c	0.097 \pm 0.007 ^c
	4	4	6	1217 \pm 102 ^a	42 \pm 1 ^b	0.025 \pm 0.004 ^c	0.089 \pm 0.003 ^c

Acclim, acclimation; temp, temperature; BM, body mass. A dissimilar letter indicates significant difference ($P < 0.05$), tested by one-way ANOVA.

For example, 5 μ M adrenaline generated a maximum V_S of 5.8 ± 1.1 ml kg^{-1} , a maximum \dot{Q} of 126 ± 23 ml min^{-1} kg^{-1} and a maximum W_C of 1.2 ± 0.3 mW g^{-1} . Maximum V_S (4.8 ± 1.0 ml kg^{-1}) and maximum \dot{Q} (109 ± 22 ml min^{-1} kg^{-1}) with 50 nM adrenaline was significantly ($P < 0.05$) lower, but maximum W_C (1.0 ± 0.3 mW g^{-1}) was unchanged.

N. coriiceps (Hb⁺Mb⁺)

In contrast to icefishes, 5 μ M adrenaline significantly decreased initial f_H (26.0 ± 0.9 to 23.5 ± 1.0 min^{-1} ; $P < 0.05$, Fig. 3), but maximum V_S (1.9 ± 0.1 ml kg^{-1}), maximum \dot{Q} (40.0 ± 2.3 ml min^{-1} kg^{-1}) and maximum W_C (2.8 ± 0.3 mW g^{-1}) were not significantly different compared with 50 nM adrenaline.

Interspecific comparisons of maximum performance with 5 μ M adrenaline at 1 $^{\circ}$ C

f_H was significantly ($P < 0.05$) lower in *C. aceratus* (Hb⁻Mb⁻) than in either *C. rastrispinosus* (Hb⁻Mb⁺) or *N. coriiceps* (Hb⁺Mb⁺), whose f_H was not significantly different from each other (Fig. 4). Maximum V_S and maximum \dot{Q} were highest for *C. rastrispinosus*, intermediate for *C. aceratus* and lowest for *N. coriiceps*. *N. coriiceps* generated a significantly ($P < 0.05$) and substantially higher maximum P_o and maximum W_C than both icefish species, which were similar to each other in this regard.

Response of maximum cardiac performance on acute warming to 4 $^{\circ}$ C

C. aceratus (Hb⁻Mb⁻)

Both f_H (26.3 ± 1.1 min^{-1}) and maximum \dot{Q} (98.3 ± 11.6 ml min^{-1} kg^{-1}) measured at 4 $^{\circ}$ C with 5 μ M adrenaline were significantly ($P < 0.05$) higher when compared to 1 $^{\circ}$ C (Fig. 1). Even so, maximum V_S (5.0 ± 0.4 ml kg^{-1}) and maximum W_C (1.7 ± 0.2 mW g^{-1}) did not change significantly. By comparison, maximum V_S was significantly ($P < 0.05$) lower

(4.0 ± 0.5 ml kg^{-1}) with 50 nM adrenaline, similar to the situation at 1 $^{\circ}$ C, but the initial f_H , maximum \dot{Q} and maximum W_C were unchanged.

N. coriiceps (Hb⁺Mb⁺)

Both f_H (31.7 ± 1.4 min^{-1}) and maximum W_C (4.1 ± 0.3 mW g^{-1}), but not maximum V_S (1.6 ± 0.2 ml kg^{-1}) or maximum \dot{Q} (48 ± 4 ml min^{-1} kg^{-1}), with 5 μ M adrenaline were significantly ($P < 0.05$) higher with acute warming to 4 $^{\circ}$ C when compared with 1 $^{\circ}$ C (Fig. 3). The heart could also generate a significantly higher maximum P_o (76 cm H₂O vs. 53 cm H₂O; Fig. 4). Adrenergic stimulation had no significant effects on maximum W_C , maximum V_S or maximum \dot{Q} at 4 $^{\circ}$ C, but initial f_H (33.8 ± 2.1 min^{-1}) was significantly ($P < 0.05$) decreased by a small amount.

Interspecific comparisons with acute warming to 4 $^{\circ}$ C:

Similar to 1 $^{\circ}$ C, f_H remained significantly ($P < 0.05$) lower in *C. aceratus* than in *N. coriiceps*, while maximum V_S and \dot{Q} remained significantly ($P < 0.05$) higher for *C. aceratus* than *N. coriiceps* (Fig. 4). Again, *N. coriiceps* (Hb⁺Mb⁺) generated a significantly ($P < 0.05$) higher maximum P_o and maximum W_C than *C. aceratus* (Hb⁻Mb⁻).

Maximum cardiac performance after 5 days exposure to 4 $^{\circ}$ C

C. rastrispinosus (Hb⁻Mb⁺)

When 4 $^{\circ}$ C-exposed fish were tested at 4 $^{\circ}$ C, the initial f_H (37.3 ± 3.3 min^{-1}) was significantly ($P < 0.05$) higher compared with fish acclimated to 1 $^{\circ}$ C and tested at 1 $^{\circ}$ C (26.0 ± 0.5 min^{-1} ; Fig. 2). While 5 μ M adrenaline produced a small, significant ($P < 0.05$) decrease in f_H (33.8 ± 3.4 min^{-1}) at 4 $^{\circ}$ C when compared with 50 nM adrenaline, maximum V_S and maximum \dot{Q} increased significantly ($P < 0.05$) by a small amount without significantly changing maximum W_C . Even so, neither maximum V_S (6.0 ± 0.7 ml kg^{-1}), maximum \dot{Q} (175 ± 19 ml min^{-1} kg^{-1}), nor maximum W_C




Figure 1: Maximum cardiac performance of the *in situ* perfused heart of *C. aceratus* (Hb⁻ Mb⁻) acclimated to 1°C as a function of adrenaline concentration in the perfusate, tested at two temperatures. A significant difference between adrenaline concentrations is indicated by dissimilar numbers; an asterisk indicates a significant difference between the performance at 1°C ($n = 8$) and 4°C ($n = 7$) for a given adrenaline concentration.

(1.3 ± 0.2 mW g⁻¹) were significantly ($P > 0.05$) different at 4°C when compared with fish tested at 1°C.

***N. coriiceps* (Hb⁺ Mb⁺)**

When tested at 4°C, the initial f_H (36.0 ± 1.0 min⁻¹) was significantly ($P < 0.05$) higher when compared with fish tested at 1°C (Fig. 3). Stimulation with 5 μM adrenaline had no significant effects on maximum W_C , maximum V_S or maximum \dot{Q} , but produced a small, significant ($P < 0.05$) decrease in f_H when compared with 50 nM adrenaline. While maximum W_C (4.6 ± 0.3 mW g⁻¹) with 5 μM adrenaline was significantly ($P < 0.05$) higher after a period of exposure to 4°C compared




Figure 2: Maximum cardiac performance of the *in situ* perfused heart of *C. rastrispinosus* (Hb⁻ Mb⁺) acclimated to 1°C and after an acclimation period at 4°C as a function of adrenaline concentration in the perfusate and tested at two temperatures (1°C and 4°C; $n = 6$ and 5, respectively). Significant differences are identified as Fig. 1.

with fish acclimated to 1°C, neither maximum V_S nor maximum \dot{Q} were different ($P > 0.05$). Moreover, none of the maximum performance parameters were different ($P > 0.05$) when 4°C-exposed fish tested at 4°C were compared with fish acclimated to 1°C and acutely warmed to 4°C (Fig. 4).

Interspecific comparisons with a period of exposure to 4°C

While 4°C-exposed *N. coriiceps* had a significantly ($P < 0.05$) higher maximum W_C and maximum P_o than 4°C-exposed *C. rastrispinosus*, maximum \dot{Q} and maximum V_S were significantly ($P < 0.05$) lower (Fig. 4). In general, cardiac performance at 4°C was similar whether acutely exposed




Figure 3: Maximum cardiac performance of the *in situ* perfused heart of *N. coriiceps* (Hb^+Mb^+) as a function of the adrenaline concentration in the perfusate. Fish were acclimated either to 1°C and tested at 1°C ($n = 7$) and 4°C ($n = 6$), or exposed to 4°C for an acclimation period and tested at 4°C ($n = 6$). Significant differences are identified as Fig. 1.

to this temperature change or measured after a period of exposure to 4°C.

Discussion

Maximum cardiac performance

We provide the first estimate of maximum cardiac pumping capacity for an icefish that lacks Mb, and the first comparison of maximum *in situ* cardiac performance among notothenioids differing in their expression of Hb and Mb. This comprehensive study of maximum cardiac function of notothenioid fishes now allows a better assessment of the likely scope for acclimation during Southern Ocean warming. While we show that *N. coriiceps* had a robust cardiac




Figure 4: A comparison of the maximum cardiac performance of *in situ* perfused hearts of *C. aceratus* (Hb^-Mb^- ; solid square), *C. rastrospinosus* (Hb^-Mb^+ ; solid triangle) and *N. coriiceps* (Hb^+Mb^+ ; solid circle) as a function of test temperature, using 5 μM adrenaline in the perfusate. Dashed lines connect fish tested at their acclimation temperature (either 1°C or 4°C), solid lines connect fish acclimated to 1°C and acutely warmed to 4°C for testing. A significant difference is indicated by dissimilar numbers between species when tested at 1°C, by dissimilar lower letters between species when tested at 4°C and by an asterisk between performance at 1°C and 4°C for each species.

performance at all of the temperature conditions we tested, the extremely large hearts associated with loss of Hb expression in icefishes incur a cost to maximum W_C , which was considerably lower than in *N. coriiceps*. While the low vascular resistance of icefish (e.g. in *C. hamatus* it is only 10–20% that of Hb⁺ fishes; *Acierno et al.*, 1997) will offset this to an extent, low W_C may still limit scope for activity despite an unusually high maximum \dot{Q} near 0°C. In this regard, the additional loss of Mb in *C. aceratus* may create a more precarious situation than *C. rastrispinosus* in a warmer world. Specifically, the maximum \dot{Q} values recorded here are in the lower range of those previously published for icefishes (70–300 ml min⁻¹ kg⁻¹; *Axelsson*, 2005). Moreover, a functional consequence for the evolutionary loss of Mb may be our novel finding of a significantly higher maximum \dot{Q} at 1°C for *C. rastrispinosus* (Hb⁻Mb⁺; 128 ml min⁻¹ kg⁻¹) than for *C. aceratus* (Hb⁻Mb⁻; 80 ml min⁻¹ kg⁻¹). Routine \dot{Q} *in vivo* is 27 ml min⁻¹ kg⁻¹ and maximum \dot{Q} 80 ml min⁻¹ kg⁻¹ for *C. aceratus* at 1°C (*Joyce et al.*, 2018b), suggesting that cardiac factorial scope could be as high as 3 (i.e. 80/27) and similar to other teleosts (*Farrell and Smith*, 2017). However, routine \dot{Q} *in vivo* for *N. coriiceps* at 1°C is 7 ml min⁻¹ kg⁻¹, while the maximum \dot{Q} is 40 ml min⁻¹ kg⁻¹, suggesting a higher cardiac factorial scope of 5.7. These comparisons suggest that the red-blooded *N. coriiceps* has the potential to withstand a warming climate better than icefishes, where conservation efforts within the Southern Ocean should be focused.

Species comparisons of maximum performance

In vitro measurements previously showed that \dot{Q} in icefishes at -0.5°C ($V_S \sim 3.5$ ml kg⁻¹) declined appreciably when afterload was increased to > 30 cm H₂O (*Acierno et al.*, 1997; cf. maximum P_o in the present study ~ 35 cm H₂O). Interestingly, poisoning Mb with nitrite in *C. rastrispinosus* (Hb⁻Mb⁺) significantly reduced W_C to 0.8 mW g⁻¹ ($P_o = 29$ cm H₂O) from 1.1 mW g⁻¹ ($P_o = 31$ cm H₂O), whereas W_C in *C. aceratus* (Hb⁻Mb⁻) was unaffected and remained at 1.4 mW g⁻¹ with a similar afterload (*Acierno et al.*, 1997). Unfortunately, this study did not assess maximum \dot{Q} , but the present findings suggest that *C. aceratus* may have been working near maximum, unlike *C. rastrispinosus*. Previous *in vitro* estimates of maximum \dot{Q} in an icefish are limited to those for *C. hamatus* (Hb⁻Mb⁺) at 3°C (200–300 ml min⁻¹ kg⁻¹ and maximum V_S of 8–11.5 ml kg⁻¹; *Tota et al.*, 1991). Our *in situ* values for maximum \dot{Q} at 4°C were considerably lower for *C. rastrispinosus* (Hb⁻Mb⁺; 175 ml min⁻¹ kg⁻¹) and still lower in *C. aceratus* (Hb⁻Mb⁻; 98 ml min⁻¹ kg⁻¹); whether this reflects an important species variation or methodological differences is unclear. However, an *in situ* heart preparation is likely a superior preparation to placing a heart in an artificial pericardium, as potential damage during isolation is avoided. As observed, icefish hearts are easily overstretched by excessive filling pressure even when the pericardium is intact, resulting in a spontaneous decrease in f_H . In addition,

in vitro heart perfusions place an input cannula in the atrium at the sino-atrial junction, potentially impairing activity of pacemaker cells located there (*Haverinen and Vornanen*, 2007; *Farrell and Smith*, 2017); intrinsic f_H for isolated heart preparations was far more variable for *C. aceratus* at -0.5°C (22–30 min⁻¹; *Acierno et al.*, 1997) and *C. hamatus* at 0.1°C (19.5–26.3 min⁻¹; *Tota et al.*, 1991) compared with the stable initial f_H of our study. Notably, neither *Tota et al.* (1991) nor *Acierno et al.* (1997) used adrenaline in their *in vitro* studies, although it is plausible that catecholamines released from cardiac tissues could have accumulated in recycled perfusate.

Both *C. hamatus* and *C. rastrispinosus* have a much higher maximum V_S compared with *C. aceratus*, a difference that may be related to Mb expression facilitating a somewhat higher cardiac afterload, while a 30% larger relative ventricular mass and a $\sim 60\%$ higher maximum \dot{Q} at 1°C (*C. rastrispinosus* vs. *C. aceratus*; above) are also of potential benefit. Although both these species have similar values of maximum P_o and W_C , Mb has a much higher oxygen affinity than Hb and enhances both oxygen storage and diffusion rate within cells. Other species differences among notothenioids also might affect maximal cardiac performance, e.g. icefish hearts lack mitochondrial creatine kinase (*O'Brien et al.*, 2014) and energy charge of *N. coriiceps* hearts is typically higher than *C. aceratus* (*O'Brien et al.*, 2018). In addition, the Mb⁺ state (independent of Hb loss) is associated with a 22% higher initial f_H at 1°C (26 min⁻¹ in *C. rastrispinosus* vs. 19 min⁻¹ in *C. aceratus*), a difference also seen at 4°C and between *N. coriiceps* and *C. aceratus*. Because the sino-atrial node was intact in the *in situ* preparation, we are confident that f_H was driven by a steady pacemaker activity, justifying our novel suggestion of a species difference in initial f_H among icefishes. A similar species difference was observed *in vivo* after blockade of cardiac vagal tone (24 min⁻¹ for *N. coriiceps* and 14.2 min⁻¹ for *C. aceratus*) (*Joyce et al.*, 2018b), while initial *in situ* f_H for *C. aceratus* (19.0 min⁻¹) compares well with *in vivo* f_H measured immediately following activity (17.6 min⁻¹; *Joyce et al.*, 2018b). Of potential value in setting resting metabolic rate, and hence scope for activity, comparisons of initial f_H values are confounded by the tonic level (50 nM) of adrenaline used to stabilize the *in situ* heart preparation; f_H of *C. aceratus* may simply be less sensitive to tonic adrenergic stimulation at concentrations < 50 nM than *C. rastrispinosus* (both species were unresponsive to further additions of adrenaline). In contrast, with a similar preparation, *M. scorpius* f_H increased by 22% at 1°C and by 27% at 6°C with maximal adrenergic stimulation, stimulatory effects that were fully reversed by β -adrenergic blockade (*Farrell et al.*, 2013).

Maximum W_C of *N. coriiceps* at 1°C was impressive when compared with those of temperate-water teleosts at much warmer temperatures (*Farrell and Jones*, 1992; *Axelsson*, 2005; *Farrell and Smith*, 2017), maintaining maximum \dot{Q} against a comparable afterload. Indeed, it shows a remarkably similar maximum cardiac performance

to *M. scorpius* (Hb⁺Mb⁺) from Greenland, with comparable adaptive response observed on acute warming (Farrell *et al.*, 2013). Despite a lower maximum V_S and \dot{Q} , *N. coriiceps* hearts generated more than twice the relative work output compared with both *C. rastrispinosus* and *C. aceratus*, in part due to an enhanced pressure-generating ability compared with icefish hearts, as reported earlier (Acierno *et al.*, 1997). However, with the much larger ventricle of icefishes, absolute maximum cardiac work output for a 1 kg representative fish was similar for all three species. Whether this translates to a similar resilience to environmental change will depend on the origin of physiological challenges to be overcome.

Fish hearts are exquisitely sensitive to cardiac filling pressure (venous return), and perfused hearts can generate an output even with a sub-ambient P_i (*C. hamatus*: Tota *et al.*, 1991; *C. aceratus*: Acierno *et al.*, 1997 and present study; *C. rastrispinosus* and *N. coriiceps*: present study). However, neither icefish species examined here displayed the sensitivity to preload shown by *C. hamatus* (Tota *et al.*, 1991), which increased V_S up to 11.5 ml kg⁻¹ (\dot{Q} up to 300 ml min⁻¹ kg⁻¹) with just a 0.25 cm H₂O (≈ 0.025 kPa) increase in P_i , although why this difference exists is unclear. Adrenergic stimulation can increase sensitivity to P_i (Farrell and Jones, 1992), as seen in the present study and with *M. scorpius* (Farrell *et al.*, 2013), by increasing maximum V_S . This effect may be particularly important for icefishes where high concentrations of nitric oxide resulting from lack of Hb conversion to nitrate (Beers *et al.*, 2010) may reduce adrenergic-stimulated contractility *in vivo* (Rastaldo *et al.*, 2007). The importance of adrenergic stimulation for cardiac contractility has been demonstrated in notothenioids: maximum tension of isolated ventricular strips from *C. aceratus* increased by 47% with adrenaline concentrations similar to those of the present study, while *N. coriiceps* maximum tension increased by a remarkable 377% with 1 mM adrenaline (Skov *et al.*, 2009). These positive inotropic effects were more marked and species-specific than the modest stimulatory effects of adrenaline on V_S or \dot{Q} (20–30% increases) seen *in situ*, so may reflect a potentially powerful role for tonic adrenergic stimulation of contractility not fully realized *in vivo*. Differences in adrenergic sensitivity could reflect species difference as well as thermal effects: adrenaline responsiveness increased at 4°C in *N. coriiceps* (present study) and at 6°C in *M. scorpius* (Farrell *et al.*, 2013), although a dependency on tonic adrenergic stimulation existed in rainbow trout (*Oncorhynchus mykiss*) hearts at 5°C, but not at higher temperatures (Graham and Farrell, 1989). Ventricular strips of both *C. aceratus* and *N. coriiceps* developed alternans at high-pacing frequencies, an impairment exacerbated by adrenaline (Skov *et al.*, 2009). A protective effect may be provided by the modest negative chronotropic effect of adrenaline, seen in both *N. coriiceps* and *C. rastrispinosus* at 4°C (present study), perhaps due to a stimulation of cardiac α -adrenoceptors or a prolongation of the ventricular action potential (Farrell and Jones, 1992). Clearly, the subtle differences in adrenergic stimulation of the heart that exist among polar fish species need further study to

better understand their autonomic cardiac regulation, as sympathovagal balance helps define a species ecotype (Campbell *et al.*, 2009).

Temperature effects

The present study revealed a clear (30–40%) increase in f_H when notothenioid species were tested at 4°C, independent of whether the fish had been acutely warmed or experienced an exposure period at 4°C. Temperature has a variety of direct cardiovascular effects, but perhaps the most important and universal is an increase in f_H during acute warming (e.g. Sandblom and Axelsson, 2007; Farrell, 2009; Eliason *et al.*, 2011; Keen and Gamperl, 2012). Notothenioids are clearly no exception in this regard, e.g. isolated *C. hamatus* hearts warmed from 0.6°C to 5.8°C, increased f_H by 21% to 24 min⁻¹ (Tota *et al.*, 1991) and *C. aceratus* doubled f_H to 36 min⁻¹ when exposed for 48 h to 10°C (Hemmingsen *et al.*, 1972). Routine f_H and \dot{Q} both increased with acute warming from -1°C to 8°C in another Antarctic fish, *Pagothenia borchgrevinki* (Franklin *et al.*, 2007). However, its scope to increase f_H , V_S and \dot{Q} following burst activity all decreased with increasing temperature because the increase in maximum cardiac capacity did not match the greater routine oxygen uptake. In the present study, both *C. aceratus* and *N. coriiceps* increased f_H by a similar amount when acutely warmed. Notably, only *N. coriiceps* significantly increased W_C (by 64%), largely through an increase in maximum P_{O_2} , while *C. aceratus* maintained maximum V_S and maximum P_{O_2} , but increased maximum \dot{Q} . Interestingly, the cardiac response to warming in *N. coriiceps* was similar to *M. scorpius*, which is considered a temperature generalist (Franklin *et al.*, 2007) and maintained scope for \dot{Q} with forced activity over a 10°C acute warming, whereas two closely related Arctic sculpins with a more restricted range could not do so (Franklin *et al.*, 2013). These comparisons suggest that the ability to tolerate ocean warming may differ even among red-blooded polar species, as well as among icefishes.

The period of exposure used here was short relative to the slow responses known for some Antarctic fishes (Franklin *et al.*, 2007; Franklin and Seebacher, 2009; Egginton and Campbell, 2016), even though they involved only small temperature changes. Therefore, conclusions concerning cardiac thermal acclimation for *C. rastrispinosus* and *N. coriiceps* must be treated with caution. For example, acclimation of *P. borchgrevinki* (Hb⁺Mb⁺) to 4°C for 4–5 weeks reduced peak \dot{Q} after exercise at -1°C (factorial scope of 1.4 vs. 2.6 for -1°C acclimated fish), while peak \dot{Q} was maintained on acute exposure to 4°C and 8°C (Franklin *et al.*, 2007). Thermal compensation maintained cardiac performance at the new acclimation temperature and allowed scope for \dot{Q} change up to 8°C, but at the expense of reduced cardiac performance when acutely cooled to -1°C (Franklin *et al.*, 2007). For *N. coriiceps* maximum performance with acute warming and exposure to 4°C were not statistically different; in contrast, thermal tolerance increased by 1.2°C after just 7 days at 4°C

(Bilyk and DeVries, 2011). Like acute warming of *C. aceratus*, *C. rastrispinosus* maintained maximum V_S , maximum \dot{Q} and maximum P_o after exposure at 4°C (present study). Clearly, a more extensive exploration of thermal acclimation effects in polar fishes is warranted, which may allow the capacity for different niche selection to be predicted, although facilities in remote areas remain a challenge for such detailed studies.

It is possible that over evolutionary time icefishes could accommodate warming as the Patagonian icefish *Champscephalus esox* apparently has done, where surface water temperature is 10–12°C over its summer range. In addition, some notothenioids have escaped the stable cold waters of the Southern Ocean and successfully colonized warmer temperate water. Stranded following a northerly extension of the ice front the black cod, *Notothenia angustata*, has established a viable population around the southern coast of New Zealand.

Concluding remarks

We provide robust estimates of maximal cardiac performance in two species of Channichthyid icefishes lacking facilitated systemic oxygen transport. While exhibiting the expected rise in heart rate on warming, our results indicate a potential to accommodate some degree of climatic change, which would have significant bearing on species survival in a warming world. While the loss of Hb is associated with large ventricles that deliver high cardiac output, retention of Mb allows additional improvements in cardiac function. Although both icefish species increased cardiac pumping capacity when warmed to 4°C, their hearts were unable to tolerate a high afterload, suggesting limited capacity to cope with stressful situations where sympathetic drive may increase peripheral resistance. However, during acute warming with activity *C. aceratus* increased *in vivo* \dot{Q} 3-fold and vascular conductance 5-fold, suggesting that short-term cardiovascular compensation is possible (Joyce *et al.*, 2018b). In contrast, a closely related species expressing both Hb and Mb was capable of higher maximum work capacity, an enhanced performance that may convey greater resilience to near-future ocean warming. Although these data demonstrate a limited capacity to withstand more frequently occurring short-term thermal extremes, and perhaps the potential to accommodate a gradual warming of the Southern Ocean, long-term temperature acclimation experiments are required to understand interspecific limits for stock conservation.

This novel information allows greater insight into plasticity of fish species previously considered to be living perilously close to their upper thermal limits as a consequence of unique physiological constraints. The issue is of some urgency, as they inhabit ocean regions around Antarctica that are experiencing the most dramatic effect of global warming; compelling reasons why we need to understand the ecophysiology of such species before conservation efforts become untenable. In addition, there has been relatively little research on ther-

mal resilience of such important predators in the simplified food chain, and we provide the most reliable cross-specific comparison on maximal cardiac performance, a key factor in ecological fitness, to date.

Acknowledgments

We gratefully acknowledge the outstanding support from the Masters and crew of the ARSV *Laurence M. Gould* and the staff at the U.S. Antarctic Research Station, Palmer Station. We thank Elizabeth Evans for assistance with experiments and fish care, Amanda Biederman with fish care and William Joyce for helpful discussions. S.E. and A.P.F. contributed to conception of the study; A.P.F. and S.E. collected field data; A.P.F. conducted statistical analysis, curated the data and drafted the manuscript; A.P.F., E.L.C. and K.M.O. provided resources and administered the project; all authors participated in funding acquisition, contributed to manuscript editing and gave final approval for publication.

Funding

Supported by operating grants from National Science Foundation (ANT 1341602 to E.L.C. and ANT 1341663 to K.M.O.) and the Natural Sciences and Engineering Council of Canada. S.E. acknowledges support from the University of Leeds, and M.A. from the Swedish Research Council. A.P.F. holds a Canada Research Chair (Tier I) and is supported by NSERC Canada.

Supplementary material

Supplementary dataset supporting this article is available at Dryad: doi:10.5061/dryad.83vc5.

References

- Acierno R, Agnisola C, Tota B, Sidell BD (1997) Myoglobin enhances cardiac performance in antarctic icefish species that express the protein. *Am J Physiol* 273: R100–R106.
- Axelsson M (2005) The circulatory system and its control. In AP Farrell, JF Steffensen, eds, *Polar Fishes* Vol 22. Academic Press, New York, pp. 239–280
- Bailey JR, Sephton DH, Driedzic WR (1990) Oxygen uptake by isolated perfused fish hearts with differing myoglobin concentrations under hypoxic conditions. *J Mol Cell Cardiol* 22: 1125–1134.
- Beers JM, Borley KA, Sidell BD (2010) Relationship among circulating hemoglobin, nitric oxide synthase activities and angiogenic poise in red- and white-blooded Antarctic notothenioid fishes. *Comp Biochem Physiol A Mol Integr Physiol* 156: 422–429.
- Bilyk KT, DeVries AL (2011) Heat tolerance and its plasticity in Antarctic fishes. *Comp Biochem Physiol A Mol Integr Physiol* 158: 382–390.

- Bilyk KT, Vargas-Chacoff L, Cheng C-H (2018) Evolution in chronic cold: varied loss of cellular response to heat in Antarctic notothenioid fish. *BMC Evol Biol* 18: 143 s12862-018-1254-6.
- Campbell HA, Davison W, Fraser K, Peck L, Egginton S (2009) Heart rate and ventilation in Antarctic fishes are largely determined by ecotype. *J Fish Biol* 74: 535–552.
- Campbell HA, Fraser KP, Bishop CM, Peck LS, Egginton S (2008) Hibernation in an antarctic fish: on ice for winter. *PLoS One* 3: e1743.
- Clarke A, Murphy EJ, Meredith MP, King JC, Peck LS, Barnes DK, Smith RC (2007) Climate change and the marine ecosystem of the western Antarctic peninsula. *Philos Trans R Soc Lond B Biol Sci* 362: 149–166.
- Egginton S (1997) A comparison of the response to induced exercise in red- and white-blooded Antarctic fishes. *J Comp Physiol B Biochem Syst Environ Physiol* 167: 129–134.
- Egginton S, Campbell HA (2016) Cardiorespiratory responses in an Antarctic fish suggest limited capacity for thermal acclimation. *J Exp Biol* 219: 1283–1286.
- Egginton S, Skilbeck C, Hoofd L, Calvo J, Johnston IA (2002) Peripheral oxygen transport in skeletal muscle of Antarctic and sub-Antarctic notothenioid fish. *J Exp Biol* 205: 769–779.
- Eliason EJ, Clark TD, Hague MJ, Hanson LM, Gallagher ZS, Jeffries KM, Gale MK, Patterson DA, Hinch SG, Farrell AP (2011) Differences in thermal tolerance among sockeye salmon populations. *Science* 332: 109–112.
- Ern R, Norin T, Gamperl AK, Esbaugh AJ (2016) Oxygen dependence of upper thermal limits in fishes. *J Exp Biol* 219: 3376–3383.
- Farrell AP (2009) Environment, antecedents and climate change: lessons from the study of temperature physiology and river migration of salmonids. *J Exp Biol* 212: 3771–3780.
- Farrell AP, Altimiras J, Franklin CE, Axelsson M (2013) Niche expansion of the shorthorn sculpin (*Myoxocephalus scorpius*) to Arctic waters is supported by a thermal independence of cardiac performance at low temperature. *Can J Zool-Rev Can de Zool* 91: 573–580.
- Farrell AP, Johansen JA, Graham MS (1988) The role of the pericardium in cardiac performance of the trout (*Salmo gairdneri*). *Physiol Zool* 61: 213–221.
- Farrell AP, Johansen JA, Suarez RK (1991) Effects of exercise-training on cardiac performance and muscle enzymes in rainbow trout, *Oncorhynchus mykiss*. *Fish Physiol Biochem* 9: 303–312.
- Farrell AP, Jones DR (1992) The heart. In WS Hoar, DJ Randall, AP Farrell, eds, *Fish Physiology* Vol 12. Academic Press Inc, San Diego, pp. 1–88.
- Farrell AP, MacLeod KR, Chancey B (1986) Intrinsic mechanical properties of the perfused rainbow trout heart and the effects of catecholamines and extracellular calcium under control and acidotic conditions. *J Exp Biol* 125: 319–345.
- Farrell AP, Smith F (2017) Cardiac form, function and physiology. In KA Gamperl, TE Gillis, AP Farrell, CJ Brauner, eds, *The Cardiovascular System: Morphology, Control and Function* Vol 36A. Academic Press Inc, San Diego, pp. 155–264.
- Franklin CE, Davison W, Seebacher F (2007) Antarctic fish can compensate for rising temperatures: thermal acclimation of cardiac performance in *Pagothenia borchgrevinki*. *J Exp Biol* 210: 3068–3074.
- Franklin CE, Farrell AP, Altimiras J, Axelsson M (2013) Thermal dependence of cardiac function in arctic fish: implications of a warming world. *J Exp Biol* 216: 4251–4255.
- Franklin CE, Seebacher F (2009) Adapting to climate change. *Science* 323: 876–877 author reply 877.
- Graham MS, Farrell AP (1989) The effect of temperature acclimation and adrenaline on the performance of a perfused trout heart. *Physiol Zool* 62: 38–61.
- Grove TJ, Sidell BD (2002) Myoglobin deficiency in the hearts of phylogenetically diverse temperate-zone fish species. *Can J Zool* 80: 893–901.
- Haverinen J, Vornanen M (2007) Temperature acclimation modifies sinoatrial pacemaker mechanism of the rainbow trout heart. *Am J Physiol Regul Integr Comp Physiol* 292: R1023–R1032.
- Hemmingsen EA, Douglas EL (1977) Respiratory and circulatory adaptations to the absence of hemoglobin in Chaenichthyid fishes. In GA Llano, ed, *Adaptations within Antarctic Ecosystems*. Smithsonian Institution, Washington, DC, pp. 479–487.
- Hemmingsen EA, Douglas EL, Johansen K, Millard RW (1972) Aortic blood flow and cardiac output in the hemoglobin-free fish *Chaenocephalus aceratus*. *Comp Biochem Physiol A Comp Physiol* 43: 1045–1051.
- Holeton GF (1970) Oxygen uptake and circulation by a hemoglobinless Antarctic fish (*Chaenocephalus aceratus* Lönnberg) compared with three red-blooded Antarctic fish. *Comp Biochem Physiol* 34: 457–471.
- Johnston IA, Battram J (1993) Feeding energetics and metabolism in Demersal fish species from Antarctic, temperate and tropical environments. *Marine Biology* 115: 7–14.
- Joyce W, Axelsson M, Egginton S, Farrell AP, Crockett EL, O'Brien KM (2018a) The effects of thermal acclimation on cardio-respiratory performance in an Antarctic fish (*Notothenia coriiceps*). *Conserv Physiol* 6: doi:10.1093/conphys/coy069.
- Joyce W, Egginton S, Farrell AP, Crockett EL, O'Brien KM, Axelsson M (2018b) Exploring nature's natural knockouts: *In vivo* cardiorespiratory performance of Antarctic icefish during warming. *J Exp Biol* 221: 1–12. doi:10.1242/jeb.183160.
- Keen AN, Gamperl AK (2012) Blood oxygenation and cardiorespiratory function in steelhead trout (*Oncorhynchus mykiss*) challenged with an acute temperature increase and zatebradine-induced bradycardia. *J Therm Biol* 37: 201–210.
- Macqueen DJ, Garcia de la Serrana D, Johnston IA (2014) Cardiac myoglobin deficit has evolved repeatedly in teleost fishes. *Biol Lett* 10: 20140304.

- Near TJ (2004) Estimating divergence times of notothenioid fishes using a fossil-calibrated molecular clock. *Antarctic Science* 16: 37–44.
- O'Brien KM, Rix AS, Egginton S, Farrell AP, Crockett EL, Schlauch K, Woolsey R, Hoffman M, Merrima S (2018) Mitochondrial metabolism may limit cardiac performance during warming in Antarctic notothenioid fishes. *J Exp Biol* 221: 1–11. doi:10.1242/jeb.177816.
- O'Brien KM, Mueller IA, Orczewska JI, Dullen KR, Ortego M (2014) Hearts of some Antarctic fishes lack mitochondrial creatine kinase. *Comp Biochem Physiol A Mol Integr Physiol* 178: 30–36.
- Portner HO, Farrell AP (2008) Physiology and climate change. *Science* 322: 690–692.
- Rastaldo R, Pagliaro P, Cappello S, Penna C, Mancardi D, Westerhof N, Losano G (2007) Nitric oxide and cardiac function. *Life Sci* 81: 779–793.
- Ruud JT (1954) Vertebrates without erythrocytes and blood pigment. *Nature* 173: 848–850.
- Sandblom E, Axelsson M (2007) Venous hemodynamic responses to acute temperature increase in the rainbow trout (*Oncorhynchus mykiss*). *Am J Physiol Regul Integr Comp Physiol* 292: R2292–R2298.
- Sidell BD, O'Brien KM (2006) When bad things happen to good fish: the loss of hemoglobin and myoglobin expression in Antarctic icefishes. *J Exp Biol* 209: 1791–1802.
- Sidell BD, Vayda ME, Small DJ, Moylan TJ, Londraville RL, Yuan ML, Rodnick KJ, Eppley ZA, Costello L (1997) Variable expression of myoglobin among the hemoglobinless Antarctic icefishes. *Proc Natl Acad Sci USA* 94: 3420–3424.
- Skov PV, Bushnell P, Tirsgaard B, Steffensen JF (2009) The role of adrenaline as a modulator of cardiac performance in two Antarctic fishes. *Polar Biology* 32: 215–223.
- Tota B, Acierno R, Agnisola C (1991) Mechanical performance of the isolated and perfused heart of the haemoglobinless Antarctic icefish *Chionodraco hamatus* (Lönnberg) - effects of loading conditions and temperature. *Philos Trans R Soc Lond B Biol Sci* 332: 191–198.
- Wang T, Overgaard J (2007) Ecology. The heartbreak of adapting to global warming. *Science* 315: 49–50.
- Whiteley NM, Egginton S (1999) Antarctic fish have limited capacity for catecholamine synthesis. *J Exp Biol* 202: 3623–3629.