



This is a repository copy of *Imagining disasters in the era of climate change : is Japan's seawall a new Maginot Line?*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/148388/>

Version: Published Version

Article:

Matanle, P. orcid.org/0000-0002-8600-3856, Littler, J. and Slay, O. (2019) Imagining disasters in the era of climate change : is Japan's seawall a new Maginot Line? *The Asia-Pacific Journal: Japan Focus*, 17 (13). 1. pp. 1-29. ISSN 1557-4660

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Imagining Disasters in the Era of Climate Change: Is Japan's Seawall a New Maginot Line?

Peter Matanle, Joel Littler and Oliver Slay

Abstract

Following the Great East Japan earthquake, tsunami and nuclear disaster of 11 March 2011, the Japanese government began constructing a series of 440 seawalls along the north-eastern coast of Honshu. Cumulatively measuring 394.2km, they are designed to defend coastal communities against tsunami that frequently strike the region. We present a case study of the new seawall in Tarō, Iwate Prefecture, which had previously constructed massive sea defences in the wake of two tsunami in 1896 and 1933, which were subsequently destroyed in 2011. We ask whether the government has properly *imagined* the next disaster for the era of climate change and, therefore, whether its rationale for Tarō's new seawall is sufficient. We argue that the government has implemented an incremental strengthening of Tarō's existing tsunami defence infrastructure. Significantly, this does not anticipate global warming driven sea level rise, which is accelerating, and which requires transformational adaptation. This continues a national pattern of disaster preparedness and response established in the early 20th century, which resulted in the failure to imagine the 2011 tsunami. We conclude by recalling the lessons of France's Maginot Line and invoke the philosophy of Tanaka Shōzō, father of Japan's modern environmental movement, who urged Japanese to adjust to the flow (*nagare*) of nature, rather than defend against it, lest they are undone by the force of its backflow (*gyakuryū*).

Keywords: Tsunami; Disaster; Reconstruction; Infrastructure; Japan; Climate change; Global warming; PTHA; Maginot Line; Transformational adaptation

Man is a part of nature, and his war against nature is inevitably a war against himself.
Rachel Carson, 1963.

Introduction

Following the Great East Japan earthquake, tsunami and nuclear disaster of 11 March 2011, in 2012 the Japanese government began constructing a series of 440 seawalls along the north-eastern coast of Honshu. Cumulatively measuring 394.2 km (Table 1), they connect with natural formations along the coast of the three eastern prefectures of Tōhoku - Iwate, Miyagi and Fukushima - to form a single barrier against destructive tsunami that frequently strike the region.¹ In 2011, for example, the majority of casualties and damage were due not to the earthquake, but the tsunami that struck approximately forty minutes later, as existing defences were overwhelmed, and communities swept away (Box 1). Nationwide a total of 18,440 people have been declared dead or missing, with 121,852 buildings totally destroyed and

281,042 half-destroyed (NPA, 2016).²

We present a case study of the new tsunami defences taking shape in Tarō in Iwate Prefecture,³ a town which had previously constructed 2.4 kilometres of seawall of between 10 and 13.7 metres in height in the wake of two tsunamis in 1896 and 1933, which was destroyed in 2011. Tarō's example is significant because its experience prior to 2011 informs post-2011 reconstruction plans and their rationales in Tōhoku. Though, as we argue below, climate change⁴ is a present-day reality worldwide and all coastal regions need to consider whether and how they adapt to the particular risks facing their communities.

Despite the accumulation of research into the 2011 disaster in Japan, perhaps due to Tarō's small size and merger into Miyako City, with the notable exceptions of Smits' (2014: 51-52, 174-75), Strusińska-Correia's (2017), and Takezawa's (2016: xix, 178) mentions of the town within the context of larger studies, there has been no detailed scholarly analysis in English about this town's experience of tsunami. Moreover, few studies in English provide a close examination of specific local infrastructures; and those researchers who do examine local structures rarely frame their descriptions within a broader contextual analysis. Furthermore, although excellent overview analyses of post-2011 tsunami defence infrastructure in Japan are available (See: Strusińska-Correia, 2017 and Suppasri *et al*, 2016), we know of no case study in English which situates post-tsunami settlement reconstruction and defence along the Tōhoku coastline within the context of Japan's historical development and experience with tsunami, and which simultaneously addresses the twin 21st century challenges of depopulation and climate change, in particular global warming driven sea level rise.

Below we provide a detailed empirical study of the history of Tarō's tsunami experiences, an

analysis of the reconstruction of the town and its seawall since 2011, and we situate these within Tōhoku's and Japan's 20th and 21st century contexts. Significantly, we take careful note of the consequences of global warming driven sea level rise for presenting a new contextual environment requiring transformative adaptation in tsunami disaster risk reduction.

Our study employs a multi-disciplinary and multi-methods approach for triangulation, depth, and rigour, including inductive and deductive information gathering and cycling, in accordance with case study analysis for sustainability research in critical human geography (Yin, 2004; Fahy & Rau, 2013; Graham, 1999; Madsen & Adriansen, 2004). Our research is therefore significant for informing resilience building in post-growth settings worldwide in an era of climate change, wherever coastal communities are under threat of tsunami inundation. The study includes:

- Systematic review of central and local government planning documentation,
- Historical near-Earth composite imagery in combination with ground level photographic observation for longitudinal verification and accuracy,
- Participant observation involving five site visits to Tōhoku between 2011 and 2017 - the most recent including ten days driving the length of the eastern Tōhoku coastline - to observe settlement and defence reconstruction in various locales, interact with local residents, and experience post-tsunami local life.
- Interviews and information checks with Japanese local and central government officials and geological scientists.

In this research we inquire into the necessity of the reconstructed tsunami defences at Tarō, given that Tarō's population is shrinking, residential zones are being relocated to higher ground, and escape routes improved. Indeed,

as we shall see, the wall may be counterproductive to efficient and rapid escape to higher ground. Second, we ask how effective these defences will be against future tsunami. All but one of the manmade tsunami defences built in Tōhoku failed in 2011, with just Fudai Village's massive wall and gate, which lie inland and are thus more effective, surviving intact while allowing overtopping, resulting in just one death (Normile, 2012; Suppasri *et al*, 2013; Takezawa, 2016). Significantly, none of the government's published post-2011 recommendations imagine the potential scale of future tsunami by including global warming driven sea level rise as a transformative change in the local risk environment. We ask, therefore, whether the government's rationale for the wall is sufficient and whether it has properly *imagined* the next disaster.

Although DeWit (2014; 2016a; 2016b) argues persuasively for Japan's success with post-Fukushima energy resilience and climate change mitigation,⁵ with respect to coastal adaptation in rural locales we argue below that Japan is yet to adopt a necessarily transformational approach suitable for an era of depopulation, global warming, and increasing severity of 'mega disasters'. This is despite the fact that the government has stated its intention to do so in its National Spatial Strategy and through being a signatory to the Paris Agreement of December 2015 (MLIT, 2014 & 2015; UNFCCC, 2017). Consequently, we contend that the seawall represents an incremental improvement on pre-2011 defences. The wall is thus more aptly understood as a continuation of the 20th century Japanese Construction State and its habitual deployment of concrete infrastructure to correct for an assumed dangerous and defective nature (See: Bird, 2013; Takezawa, 2016) We conclude by arguing against preparing for the past by failing to imagine the future, and we invoke the philosophy of Tanaka Shōzō, father of Japan's modern environmental movement, who urged Japanese to live within

and adjust to the flow (*nagare*) of nature, rather than fight against it, lest they are undone by the force of its backflow (*gyakuryū*) (Stolz, 2007; 2014; Tanaka, 2004 & 2005).

Before describing the historical context of Tarō's decision to build its wall, we need to clarify three issues. First, we are as concerned as anyone about the levels of greenhouse gasses (GHG) in the Earth's atmosphere, which passed 415ppm of carbon dioxide concentration for the first time in human history on 12 May 2019 (Scripps Institution, 2019). Japan bears a heavy responsibility both as the world's third largest economy and its accumulated historical emissions as the earliest industrialised country in Asia. Nevertheless, this article is about coastal adaptation to the actual or expected outcomes of climate change, and discussion of mitigation measures is beyond our scope. Second, we do not wish to single out Japan as exceptionally unresponsive to climate change. Rather, nearly all countries are doing considerably less than necessary to acknowledge, prepare for, and adapt to sea level rise, among many other outcomes of global warming. Third, our research is rooted in the perspectives of critical human geography and we do not offer alternatives or solutions, which again would be beyond our scope and a distraction from our focus on the reconstruction of Tarō and its tsunami defences.

Table 1 Wall heights and lengths in Iwate, Miyagi and Fukushima Prefectures, pre- and post-2011.

Prefecture Length of coast	Wall Heights	Length of wall km (Percentage of Coastline) Pre-2011	Length of wall km (Percentage of Coastline) Planned/Under Construction Post-2011
Iwate 710 km of coastline	Less than 5 m	16.4 km (2%)	1.1 km (<1%)
	5 to 10 m	41.3 km (6%)	35.8 km (5%)
	10 m +	11.1 km (2%)	45.9 km (6%)
	Total	68.8 km (10%)	82.8 km (12%)
Miyagi 830 km of coastline	Less than 5 m	108.6 km (13%)	92.6 km (11%)
	5 to 10 m	45.5 km (6%)	143.2 km (17%)
	10 m +	0 km (0%)	3.6 km (1%)
	Total	154.1 km (19%)	239.4 km (29%)
Fukushima 160 km of coastline	Less than 5 m	3.5 km (2%)	0.6 km (<1%)
	5 to 10 m	66.8 km (42%)	71.4 km (45%)
	10 m +	0 km (0%)	0 km (0%)
	Total	70.3 km (44%)	72.0 km (45%)
Total (1,700 km)		293.2 km (17.2%)	394.2 km (23.2%)

Source: Data compiled by the authors from NASC (2014).



See the film here
(<https://tinyurl.com/y2es7ck6>).

Box 1: Surveying the extent of tsunami damage and reconstruction in Tarō and Tōhoku.

Below are four short films made using near Earth remote sensing composite imagery from Google Earth Pro. They survey Tarō and the coastlines of Iwate, Miyagi and Fukushima prefectures, featuring selected towns and topographic features. Yellow, red, and purple lines are 10m, 15m, and 20m contours marking elevation above sea level, using K. Tani’s GIS contour data (see here (http://ktgis.net/tohoku_data/index_e.html)). Please click on the thumbnail image to view each film.

Film 1. Tarō - (3m10s)

Tarō’s location as a ria town on Iwate’s coast - Coastal defences - Overview from various angles showing damage and reconstruction challenges - Main seawall - Port area - New residential zone - Agricultural, education, and industrial zones - Solar farm - Parks and amenities zone - Zoom out.

Film 2. Iwate Prefecture - (4m33s).

Aerial view of Iwate’s ria coastline, with significant settlements - Miyako, the long ria, small settlements, and main town - Ongoing reconstruction of residential zones & infrastructure - Kamaishi and its tsunami barrier - Rikuzentakata and its seawall - Reconstruction of Rikuzentakata - Zoom out.



See the film here
(<https://tinyurl.com/y5wrj9dg>).

Film 3. Miyagi Prefecture - (3m12s)

Aerial view of Miyagi coast - Zoom to Ishinomaki City - Ridge protecting inland areas - Sendai Plain - Coastal plain south through Natori & Watari - Long seawall - Zoom out.



See the film here
 (<https://tinyurl.com/y6kvamkc>).

Film 4. Fukushima Prefecture - (3m04s)

Fukushima coastline - Long and straight with settlements separated by mountains - Minamisoma - Seawall - Namie & Futaba - Fukushima Daiichi Nuclear Power Plant - Okuma & Tomioka - Fukushima Daini NPP - Zoom out.



See the film here
 (<https://tinyurl.com/y334jq5t>).

Tarō and Sanriku on 11 March 2011

Tarō is located in the northern part of the Sanriku region of Iwate and Miyagi Prefectures (Box 1); located approximately 130km west of the Japan Trench tectonic subduction zone. The area is famous for its picturesque ria coastline

(Photo 1),⁶ with settlements built on narrow alluvial fans nestled between steep forested mountainsides at the head of each ria. Tarō is one such town.

Photos 1-4,⁷ Click on the photos for an expanded slideshow of all images and tables used in this article. Click on the hyperlinks for a Google Earth view.

- Photo 1 (<https://tinyurl.com/y6ns4sn9>): The entrance to Tarō's ria, showing a bank of tetrapods, reconstructed since 2011. Tsunami amplification due to the ria's topography can be deduced from this image.
- Photo 2 (<https://tinyurl.com/y3ajdaoa>): The 'Tsunami Prevention Town Declaration' (*Tsunami bōsai no machi sengen*) inscribed in granite and placed at the entrance to Tarō's administrative offices, post-2011.
- Photo 3 (<https://tinyurl.com/y52cfb93>): The Tarō Fishing Coop port tower, with yellow markers for the 1896, 1933 & 2011 tsunami, displaying 17.3m for the latter. For scale, Peter Matanle is under the markers.
- Photo 4 (<https://tinyurl.com/yydfzrjs>): The Tarō Kankō Hotel from where Satoshi Itō viewed the advancing tsunami. The hotel is preserved in its post-tsunami state as a visitor centre, memorial, and warning.






Sanriku is a known tsunami disaster zone. Despite efforts to discourage settlement in vulnerable locations, regular disaster drills, and the construction of massive concrete coastal defence infrastructure, loss of life occurs repeatedly when tsunami strike. Indeed, humanity has yet to prove capable of defending against tsunami of the size and power that struck Tōhoku on 11 March 2011. The disaster was notable due to many towns having constructed advanced sea defences, which may have provided a false sense of security, since all but those in Fudai were either destroyed or

damaged (Furuta & Seino, 2016; Onishi, 2011). Stretching from northern Iwate to the Ōtsuchi Peninsula in Miyagi, the ria topography itself contributes to disaster severity by funnelling advancing tsunami, raising their height and amplifying their power (Hayashi, 2012).

Tarō was its own municipality until 2005, when it and neighbouring Niisato Village were absorbed into Miyako City under the ‘Heisei gappei’ municipal mergers.⁸ The population of Tarō has been reduced by more than half since the 1960s, with accelerating ageing and depopulation (Figure 1). Miyako too is shrinking, losing more than a quarter since 1960. Tarō shows the classic characteristics of Japan’s shrinking regions, and the recent acceleration in depopulation is not only due to the tsunami but the town’s increasing death rate as ageing intensifies, a decreasing birth rate as the number of adults of child-bearing age declines and low fertility continues,⁹ and out-migration by younger adults seeking educational, employment, and social opportunities in urban areas (See: Matanle, 2006; Matanle & Rausch *et al*, 2011; MIC, 2018). Significantly, in 2008 the Japanese population as a whole began shrinking, rendering expectations for growth in Tarō unrealistic, and post-tsunami long-term reconstruction planning challenging; since under a conventional reconstruction scenario new housing and infrastructure may in future decades become underutilised as the town continues to shrink.

Figure 1. Left: Total population of Tarō (left hand scale) and Miyako (including Tarō; right hand scale), 1945-2015. Right: Population pyramid for Tarō, 2010.




Data source: MIC, 2018. Graphs created by authors.

Older people are especially vulnerable to disasters (McMillan, 2014; Ngo, 2012). During Hurricane Katrina in New Orleans and the Gulf coast nursing home residents were unable to evacuate independently and triage protocols prevented treatment for chronic conditions. There, 71% of deceased victims were over 60 years old and 47% over 75, exceeding their 15% of the resident population (Klein and Nagel, 2007). Similarly, in 2010, 30.9% of Miyako’s and 33.0% of Tarō’s populations were aged 65 or over, comfortably qualifying as hyper-aged communities, defined as 21% or more of the population over 65 (Coulmas, 2007: 5). In Tōhoku, of the 13,135 people assigned a cause of death by 11 April 2011, 65.2% were 60 or over and 92.5% had drowned (Japan Times, 2011). Mainly for mobility reasons many older people were unable to reach safety and we can assume that many disabled - who are often older - also fell victim. Tarō and other hyper-aged communities in coastal Tōhoku are therefore especially vulnerable to tsunami.

Ageing and depopulation are global phenomena, and Japan is in the vanguard in Asia. Indeed, nearly half of the world’s countries are now experiencing below-replacement fertility, and more than 30 countries have shrinking populations (GBD 2017 Causes of Death Collaborators, 2018). China, South Korea, and Taiwan are rapidly ageing and expect to begin depopulating soon. Even relatively youthful Anglophone societies

with higher rates of immigration such as New Zealand are experiencing ageing and depopulation processes sub-nationally and in disaster prone areas (Jackson and Brabyn, 2017; Jackson and Felmingham, 2002; McMillan, 2014). All these countries are located on the Pacific Ring of Fire and experience frequent tectonic events. Beyond building resilience at home, therefore, Japan’s significance lies in how disasters and their human responses inform other countries as they enter their own post-growth pathways, particularly in disaster-prone regions of East and Southeast Asia (Matanle, 2013). Then there is climate change and its transformative contribution to the dynamics of human-environmental interactions.

Figure 2. Google Earth image showing the ria coastline of Sanriku, the Japan Trench (dark blue, centre), and major undersea (M8.0+) earthquakes in the region’s history.



Source: NOAA, 2019.

NOAA (2019)¹⁰ lists 134 known fatal¹¹ tsunami in Iwate since the Mw8.6 Jōgan earthquake in 869CE, which killed an estimated 1,000 people. Of these, 85 had parent earthquakes in the Japan Trench. Eleven of the most powerful are shown in Figure 2. Among NOAA’s listed tsunami, 17 note runup occurring at Tarō, beginning with the 1611CE Mw8.1 Keichō Sanriku earthquake, whose tsunami advanced 1.2km inland with a maximum run-up of 25m, and killed approximately 5,000 people nationwide (NOAA, 2019). This earthquake

occurred six years after the Keichō Nankaidō earthquake, whose tsunami killed more than 5,000 people in central Honshu, and just two and a half months after the Aizu earthquake, which killed approximately 3,700 people in central Tōhoku. The unpredictability of earthquakes is underlined by Smits (2014) in his detailed history of earthquakes in Japan, and reinforced by Lay (2015), who demonstrates a surge in the number of great earthquakes worldwide in the decade 2004-14, by 265% compared to 1900-2004.

Hence, although earthquakes along the Japan Trench can occur in a regular sequence, unexpected combinations of events also occur. Drawing on the example of 2011, which included a serious accident at the Fukushima Daiichi nuclear power plant, in the 21st century it is reasonable to assume that powerful earthquakes and tsunami are likely to intersect with other risk vectors - climate change and industrial infrastructure - to generate compound techno-environmental events of great complexity and severity (Matanle, 2011).

Of the 27,122 people killed in the 1896 Meiji Sanriku tsunami, Tarō lost 1,867, or 83.1% of the town’s population, and all registered buildings were destroyed (Table 2) (Yamashita, 2003; Hayashi 2012). The 1933 Shōwa Sanriku tsunami also had a devastating impact, destroying 500 of 559 buildings, and killing 32% of Tarō’s 2,773 population (Yamashita, 2003). Hence, in 1934 the townspeople began to build a huge defensive seawall.

Completed in 1978, Tarō’s pre-2011 physical tsunami defence infrastructure was mainly comprised of three walls forming an X shape (Figure 3). This was in addition to port breakwaters and a bank of concrete tetrapods at the mouth of the ria facing east towards the Japan Trench for advance protection (Figure 3; Photo 1). In total 2,433m of seawall at a height of 10.0-13.7m was constructed between 1934 and 1978, forming what was considered the most comprehensive tsunami barrier in Japan; this was superseded in 2009 by the world’s deepest breakwater at Kamaishi (USD1.5 billion), which also failed in 2011 (Ogasawara and Sakai, 2012; Yamashita, 2003).

Due to their overwhelming stature dominating the surrounding town, Tarō’s seawalls became known as the *Nihon no Banri no Chōjō* [The Great Wall of Japan] (Nikkei, 2011), and the town earned the nickname *Tsunami Tarō* [Tsunami Boy] (Inoue, 2013). On the seventieth anniversary of the Shōwa Sanriku tsunami in 2003, Tarō Town Office published its ‘Tsunami Prevention Town Declaration’ (*Tsunami bōsai no machi sengen*), which described the walls as a symbol of the resilience of Tarō in defending against tsunami (Yamashita, 2003). The declaration acknowledged the fear unleashed, and declared the walls representing the townspeople’s defiant courage (*chōsen suru yūki*) (Photo 2). Though the walls became symbolic of Tarō’s identity and pride, in 2011 they did not protect the town as effectively as those who drafted the declaration had hoped (Figures 3 & 4).

Table 2. Tsunami effects on building damage and mortality in Tarō.

Tsunami	Date	Population	Fatalities	Percentage	Buildings	Destroyed	Percentage
Meiji Sanriku	15 June 1896	2248	1867	83.1%	345	345	100%
Shōwa Sanriku	3 March 1933	2773	911	32.5%	559	500	89.4%
Great E. Japan	11 March 2011	4302	166	3.9%	1467	979	66.7%

Source: Hayashi, 2012.

Figure 3. Google Earth Pro historical image of Tarō on 20 July 2009, with pre-2011 seawalls superimposed. Image compiled by the authors.



Figure 4. Google Earth Pro historical image of Tarō on 14 March 2011, showing the extent of damage from the tsunami. The image is marked with 'A', 'B', 'C', and 'D'.



Prior to the construction of Tarō's pre-2011 walls the two most destructive recent tsunami had been the 14.6m Meiji Sanriku and 10.1m Showa Sanriku tsunami (Hayashi, 2012). Subsequent to beginning construction, four tsunami had struck Tarō between 1934 and 2011, causing minimal damage (Hayashi, 2012; Inoue, 2013). This included the 1960 Mw9.5 Valdivia earthquake near Chile, whose tsunami killed 53 people in Ōfunato and 41 in Minamisanriku, Iwate. Since the walls prevented damage from relatively common smaller tsunami, they were assumed to be sufficient.

The Failure of Tarō's Seawalls

Shortly after 3.00pm on 11 March 2011 the tsunami broke through the tetrapods, destroyed the port breakwaters, and overcame all three seawalls, demolishing the newer second seawall (Figures 3 & 4 - Area D), killing 166 people and demolishing 979 buildings. The tsunami's officially recorded height in Tarō was

15.75m, 1.1m higher than in 1896, and around 5.3m higher than the existing seawalls. The Tarō Fishing Cooperative's port tower registers the tsunami reaching 17.3m (Photo 3), or 6.9m higher than the wall. The first wall survived structurally intact; but massive overtopping caused damage to buildings behind (Figures 3 & 4 - Area A).

The main reason for the infrastructure failure was over-confidence in predicting the maximum magnitude of future earthquakes as well as the height and strength of associated tsunami. Seventy-four per cent of seawalls in Tōhoku were built on assumptions of an Mw8 earthquake triggered tsunami, not the Mw9.0 quake of 2011 (Cyranoski, 2012). Despite this, Davis *et al* (2012) suggest that 3/11 was a 'missed opportunity' for disaster preparedness, because a powerful undersea quake with a giant tsunami occurring within the foreseeable future had been anticipated. In 2001, they had identified with a 70% certainty that an Mw8+ earthquake would occur. Furthermore, Satake *et al* (2007) assumed a 99% probability of an Mw8.1-8.3 earthquake occurring within 30 years. Neither of these studies, however, anticipated the Mw9.0+ event. Indeed, Smits (2014: 22-23) argues that the change from using the Richter scale to moment magnitude as a method for measuring earthquake strength from the late-1970s onwards, and historical earthquakes' subsequent re-classification, may in part have contributed to the lack of studies predicting a Mw9.0+ earthquake for the Japan Trench region.

Secondly, initial warnings underestimated the tsunami's size (Nakahara, 2011) (Table 3), and residents' behavior during the preceding minutes suggests the seawalls gave a false sense of security. Three minutes after the earthquake, Iwate Prefecture announced a 3m+ tsunami warning. This was standard for Mw8.0+ earthquakes and, according to drills, residents should have evacuated. Then, 44 minutes after the earthquake, the announced

height was upgraded to 10m+. Possibly this did not provide enough evacuation time for some of those who had been slow respond to the initial warning. Significantly, the Japan Meteorological Association’s observed wave heights were recorded using floating buoys, some of which ceased data collection, having been damaged by the very tsunami they were designed to measure (Ozaki, 2011).

Table 3. Seawall heights, announced tsunami heights, and observed tsunami and run-up heights in selected locations in Iwate Prefecture on 11 March 2011 (Metres).

City/Town/Village in Iwate Prefecture	Seawall heights	Announced height 14:49 JST (3 minutes post-earthquake)	Announced height 15:30 JST (44 minutes post-earthquake)	Observed tsunami height (JMA tide gauge)	Observed run-up height
Hirano, Kujikita	12	3	10+		12
Kuji minami	12	3	10+		14.5
Noda Bay	12	3	10+		21.4
Fudai	15.5	3	10+		18.4
Tarō	10–13.7	3	10+		16.3
Miyako Bay	8.0–8.5	3	10+	8.5+	11.6
Yamada Bay	6.6	3	10+		10.9
Otsuchi Bay	6.4	3	10+		15.1
Kamaishi Bay	4.0–6.1	3	10+	4.2+	10.1
Ōfunato Bay	3.0–3.5	3	10+		10.4
Ōno Bay	4.8–8.5	3	10+		16.6
Hirota Bay	4.95–6.5	3	10+		18.3

Source: JMA, 2011a; JMA, 2011b; AIJ, 2012; Ozaki, 2011.

Thirdly, eyewitness Itō Satoshi, owner of the Tarō Kankō Hotel, who observed the tsunami from the fourth floor of his hotel (Photo 4), recounts people milling around behind the walls, apparently unaware of the danger, despite broadcasted warnings. He suggested that, although they had felt the earthquake, the high walls prevented them from seeing the approaching wave, nor were they rushing to evacuate due to their sense of safety behind the walls. Itō called out to warn them and filmed the wave. According to Itō, the areas immediately behind the seawalls had the highest mortality, compared to other areas where people had seen the advancing wave and quickly moved to higher ground along evacuation routes (Shifleen, 2015).

Although the proportion of casualties in 2011 was lower than in the two most serious recent tsunami, at 3.9% of the town’s population

(Table 2), suggesting that the wall saved many lives, additional factors may also have contributed to reduced mortality.

- Advanced construction techniques using concrete and steel frames for new buildings.
- Retrofitting older buildings for earthquake safety.
- The population in 2011 had increased by 92% and 55% since 1896 and 1933, and there were 325% more buildings than in 1896. Many newer structures were located on higher ground inland of Zones A, B and C (Figures 3 & 4) and were unaffected.
- Tarō’s pre-2011 investment in soft-infrastructure was considerable (Yamashita, 2003). This included coastal forests; disaster radio, satellite receivers, and observation cameras; evacuation drills and disaster memorials; evacuation sites and routes; hosting a coastal municipality tsunami summit, etc.

Further, an alternative explanation is emerging from Tōhoku of the relative effectiveness of hard defence infrastructure. Aldrich (2019; Aldrich & Sawada, 2015) compared mortality for different communities along the entire coast of the three most affected prefectures and found that social capital stocks were a more significant predictor, and that differential mortality rates were also due to the pace of evacuation. Importantly, his research shows that social capital is also a better predictor for post-disaster survival and recovery, with communities possessing higher levels of capital recovering more rapidly (2019). Finally, the failure of seawall 2 (Figure 3) may have had a negative impact on survival. It is sensible to assume tsunami defences are more effective if they remain standing. The collapsing wall probably caused a water surge, much like a dam failure, accelerating the wave’s velocity and increasing its destructive power (Kanda, 2016).

Tarō's experience with seawalls is significant because the townspeople's decision to build what was considered long-term protection was in response to the 1896 and 1933 disasters and is a cautionary example for the post-2011 decision to build a comprehensive barrier along the coastline of Iwate, Miyagi, and Fukushima prefectures. Among the additional 100km of seawall nearing completion in Tōhoku, 49.5 km of which is over 10m in height, the new defences at Tarō will be 14.7m high (Reconstruction Agency, 2015), 2.6m lower than the height of the 2011 tsunami as measured on the Fishing Cooperative's port tower. Assuming the continuation of current conditions and the successful experience of Fudai Village in 2011 with overtopping, the new seawall will likely save considerable life and damage. We now proceed to analysing the new defences and their emerging 21st century context.

Reconstructing Tarō and its Tsunami Defences

Tarō

On 1 June 2011 Miyako City began formulating its basic reconstruction plan and, after a series of open meetings on matters such as housing, education, health, transportation, and industry, presented a basic plan to citizens on 14 October. Citizens had four days to respond before the City Office finalised the plan on 31 October. Since then the City Office has regularly consulted residents about their reactions to current progress and delivery, though the extent to which citizen voices are genuinely incorporated into plans, or whether local authorities have sufficiently communicated their rationales to citizens, is contested.¹² Indeed, residents in Tarō and along the Tōhoku coast expressed concern that they wanted a greater say in post-tsunami reconstruction planning, which is corroborated by Strusińska-Correia (2017).

Figure 5. Composite image of Tarō (9 January 2016) with reconstruction plans superimposed.



Source: Image prepared by the authors using Google Earth Pro and Miyako City (2011: 5).

Figure 5 shows a Google Earth image of Tarō taken on 9 January 2016 with the town's basic reconstruction plan superimposed. Noteworthy aspects are:

- Formerly forested mountainside located behind cliffs (north-east yellow) has been appropriated for housing and amenities, including a hospital (Photos 5 & 6).
- A second residential housing zone is under construction to the north-west (yellow). Slightly higher than the port area, this is being raised by up to three metres. Some housing will be taller structures with rooftop evacuation. West of this (brown) will be a school, amenities, and park space doubling as an evacuation zone (Photo 7).
- A third residential zone is under construction in a forested mountainous area to the south, out of range of the image, and connected to the town by road. It too is protected by cliffs.
- Low-lying areas in the town centre are reserved for parks, sports pitches, amenities (brown; Photo 8), marine industrial and business facilities, a solar farm (purple; Photo 9), retail facilities (pink; Photo 10), and transportation (red, dotted lines; Photo 11). There will be no

housing here and signposted evacuation routes are under construction.

- Areas along the river valley stretching westwards from the south will be a railway station zone (red); information and communications technology zone (purple); education zone around a senior high school (brown); and an agricultural zone (green). These areas are either on higher ground or located inland.

As the photos show, there is much to do and it is possible the plans are unrealistically costly in reconstructing what is already an ageing and shrinking community which in future decades, and in all likelihood, will not require all of the housing, services, and infrastructure being built under the present plans (Matanle, 2013).

Photos 5-11 (Clockwise from the top). Click on the photos for a slideshow and hyperlinks for a Google Earth view.



Google Earth view
(<https://tinyurl.com/y4vlpsnl>)



Google Earth view

(<https://tinyurl.com/y24m7xmw>)



Google Earth view
(<https://tinyurl.com/y3s715xp>)



Google Earth view
(<https://tinyurl.com/y6rzny95>)



Google Earth view
(<https://tinyurl.com/y419ju8u>)



Google Earth view
(<https://tinyurl.com/y3r5cxzo>)



Google Earth view
(<https://tinyurl.com/y5zzo8b5>)

Concrete seawalls are built for primary defence against storm surges, tsunami and coastal erosion, and may be used in conjunction with breakwaters, natural reefs, sand beaches, lagoons, forests, canals, levees, manmade banks, and road embankments (Tokida & Tanimoto, 2014; Koshimura *et al*, 2014). Built-up areas close to the coast often rely on a seawall alone, or a harbour with breakwaters. Tetrapods are also deployed either onshore or near-shore. Throughout Tōhoku on 3/11, including Tarō, each of these suffered damage or destruction (Mikami *et al*, 2012). Although Strusińska-Correia (2017) analyses the government's overall tsunami defence strategy for Tōhoku in great detail, including mention of Tarō, it is worth reviewing this issue before

proceeding to an analysis of the importance of addressing the challenges from the perspective of climate change.


Tarō's Seawalls

Tokida and Tanimoto (2014) examined combined sea defences' effectiveness, suggesting that the flow and force of the tsunami can be reduced most effectively with a seawall, forested embankment, and manmade canal. This incorporates the government's new two-level tsunami classification system, introduced in response to the failure of 2011 (Kanda, 2016; MLIT, 2011; Raby *et al*, 2015; Shibayama *et al*, 2013). Level 1 events are considered tsunami protection events and assumed to occur every 50-60 to 150-160 years, with inundation depths of less than 7-10m. Level 2 are evacuation events of 10m+ inundation depth and assumed to occur every few hundred to a few thousand years apart. The 2011 tsunami is classified Level 2, resembling the 869CE Jōgan tsunami (Minoura *et al*, 2001; Shibayama *et al*, 2013).

Under this classification, post-2011 defences are designed to contain Level 1 events and allow Level 2 events to overtop walls without destroying them, as happened in Fudai in 2011, with overtopping seawater being collected in a land side canal running parallel (Raby *et al*, 2015). In addition, the specifications prevent land-side scouring caused by retreating waters pulling seaward, allowing subsequent waves to flow through destroyed defences. The wall's slope is 2:1 on both sides, meaning a 10m wall, with a 3m flat top is altogether 43m across (Figure 6), which is devastating for sensitive coastal ecologies. Government literature addresses this but frames the choice as a question of positioning rather than expediency (Bird, 2013; MLIT, 2011; NACS-J, 2013). The original pre-2011 X-shaped walls constructed in Tarō closely resemble this design but were too small and lacked the land-side canal. Under these new recommendations the defences at

Tarō are therefore being modified with an inner port wall of 14.7m and the original wall partially reconstructed and strengthened to a height of 10-12m (Photos 12-15; Miyako City, 2011), which will provide a base of more than 40m width. In addition, the harbor breakwaters and near-shore tetrapods are being strengthened, replaced and expanded in size and scope, giving the town four lines of concrete defence, as opposed to three in 2011. Concrete defences in Tōhoku have an anticipated effective lifespan of approximately 80-100 years, or until 2100-2120.

Figure 6. Cross section of the post-3/11 government recommended concrete sea wall design.



Source: Adapted from MLIT, 2011; NASC, 2014.

internal breakwater will obscure ocean views from all sea level or near sea level locations in Tarō.



Photos 12-15 (July 2017). Click on the photos for a slideshow and hyperlinks for a Google Earth view.

- Photo 12 (<https://tinyurl.com/y6x3ddj3>): Cross-sectional view through the central gate in the main seawall at Tarō. The lighter concrete atop the darker pre-2011 wall is a recent addition. Click here for a Google Earth view.
- Photo 13 (<https://tinyurl.com/y2fxgq8m>): Atop the main seawall, plus its landward slope.
- Photo 14 (<https://tinyurl.com/y4kgs3d5>): The new internal breakwater under construction at the waterside fishing port zone.
- Photo 15 (<https://tinyurl.com/y36jl4y3>): Taken from atop the main seawall, with debris from the wall's destruction in 2011 still evident, showing that the



(Un)Imagining the Future? Climate Change and Tsunami Size

The reduction in Taro’s death rate from 83.9% of its population in 1896, through 32.5% in 1933, to 3.9% in 2011 (Table 2), indicates significant improvements in disaster risk reduction, and in combining physical sea defences with soft-infrastructure. Nevertheless, questions arise as to whether, by constructing defences to prevent a recurrence of the previous disaster, the authorities are adequately imagining and preparing for a changing future. Indeed, there is already evidence of a public trust deficit in the effectiveness of the reconstruction (Aldrich, 2017; Aldrich & Sawada, 2015; Bird, 2013; Strusińska-Correia, 2017).

In the aftermath of the 2011 disaster, the word *sôteigai* (‘beyond expectations’; ‘unimaginable’) was deployed by the government and Tokyo Electric Power Company to explain their failure to prevent the nuclear disaster (Samuels, 2013, 104). Significantly, the Japanese government acknowledges the relationship between anthropogenic climate change and rising sea levels. Japan is a member of the United Nations Framework Convention on Climate Change (UNFCCC), was host to COP3, which produced the Kyoto Protocol, and is signatory to the 2015 COP21 Paris Agreement, ratified into Japanese law on 8 November 2016 (UNFCCC, 2017), legally committing itself to climate mitigation and adaptation. Here we ask whether authorities are adequately ‘imagining’ the next tsunami.

Table 4. Historically recorded fatal tsunami affecting Iwate Prefecture with parent earthquakes originating in/near Japan (Data for deaths and damage shows totals for Japan, including Iwate Prefecture).

Year	Earthquake Location	Earthquake and Tsunami Parameters					Tsunami Effects					
		Latitude	Longitude	Parent Earthquake Magnitude	Maximum Water Height	No. of Runups	Deaths	Injuries	Damage (US\$Mill)	Houses Destroyed	Houses Damaged	
869	SANRIKU	38.5	143.8	8.6	ND	5	1000					
1611	SANRIKU	39	144.5	8.1	25	11	5000					
1616	SANRIKU	39	144.5	7.5	ND	4	200					
1793	SANRIKU	38.5	144	8.3	4.5	8	720				100	
1847	SANRIKU	ND	ND	ND	ND	4	335					
1856	SE HOKKAIDO	40.5	143.5	8	6	33	26				193	238
1896	SANRIKU	39.5	144	8.3	38.2	72	27122	9247			11000	
1933	SANRIKU	39.224	144.622	8.4	29	294	3022				6000	
1952	SE HOKKAIDO	42.15	143.85	8.1	6.54	218	33					
1968	E HONSHU	40.6	143.2	8.2	6	329	52					529
2011	NE HONSHU	38.297	142.372	8.1	38.9	6271	18453	6152	220085.5	127511		273796

Adapted: NGDC/WDS Global Historical Tsunami Database, NOAA Website. Search: 08/03/2017.

Between 869 and 2017 an earthquake in the Japan Trench leading to a fatal tsunami in Iwate Prefecture has occurred on average every 104 years (Table 4). Five have resulted in more than 1,000 deaths, killing approximately 55,963 people; on average, they occurred every 296 years. However, the timing of the 1896, 1933 and 2011 tsunami shows that earthquakes involving mass fatalities are not temporally equidistant (See: Lay, 2015; Smits, 2014). Hence, while it could be argued the 2011 earthquake was unexpected, it was not unimaginable. It was the fourth most powerful on Earth since modern records began in 1900, came less than seven years after the Mw9.1 Indian Ocean earthquake whose tsunami killed nearly 230,000 people, and two other Mw9.0+ earthquakes have occurred in the Pacific Rim since 1960.¹³

Before 2011, Rahmstorf (2007) estimated sea level will rise 0.5-1.4m by 2100 compared with 1990 and warned that rises could be greater. Even so, estimates of the rate of sea level rise are regularly revised, usually upwards, as knowledge of ice sheet melting and analogous paleoclimatic conditions accumulates and scenario modelling improves. While disagreement remains over the rate of sea level rise, it is occurring, and likely more rapidly than at any period in the Earth’s history (Hansen, 2009). Indeed, IPCC AR4 estimated a global average rise of 0.18-0.59m by 2090-99

compared with 1980-99; seven years later AR5 estimates a rise of 0.26-0.82m by 2081-2100 relative to 1986-2005 (IPCC, 2007: 8; 2014: 13).

Some argue that IPCC projections are too cautious. Schaefer *et al* (2014) state that '[C]limate projections in ... (AR5) ... do not adequately account for emissions from thawing permafrost'. Hansen *et al* (2016) assert that a global average 2°C surface temperature increase may result in rises 'reaching several meters over a timescale of 50-150 years', due to 'amplifying feedback processes'; concluding that humanity is experiencing 'a global emergency' (2016: 3801). One of these is the potential for ice-cliff instability contributing more than a metre extra to current sea level rise predictions by 2100 (DeConto & Pollard, 2016). Furthermore, although the 'Paris Agreement's central aim is to ... pursue efforts to limit the temperature increase even further to 1.5 degrees' (UNFCCC, 2017), assessments of its achievability are pessimistic (Reilly *et al*, 2015; Rogelj *et al*, 2016). Recent studies confirm that ice-melt and sea level rise are accelerating (IMBIE Team, 2018; Nerem *et al*, 2018), and that IPCC projections underestimate future change.

The IPCC (2014) anticipates that current GHG emissions trajectories require significant coastal remodelling worldwide this century, and the Paris Agreement mandates adaptation and mitigation into national laws (UNFCCC, 2017). In the USA this includes new infrastructure, land use modifications, and repair and retrofitting of coastal facilities (EPA, 2017). Recently adaptation there has begun to include nature-based features (NNBFs) within a mix of hard, soft, and green infrastructure (Hill, 2015; Bridges, 2015).¹⁴

Climate change adaptation is usually incremental. Hinkel *et al*'s (2018) research in six locations worldwide concurs with this in shorter timescales; and it is sensible to assume

that ongoing incremental change may accumulate into a transformational adaptation. Nevertheless, where human systems may be suddenly overwhelmed, where longer timescales are involved, or where incrementalism is insufficient, transformational adaptation may be required. Kates *et al* (2012) argue that these 'may be difficult to implement because of uncertainties about climate change risks and adaptation benefits, the high costs of transformational actions, and institutional and behavioural actions that tend to maintain existing resource systems and policies'; what Hinkel *et al* (2018) refer to as social conflicts, economic and financial barriers, and technological limits. They describe transformational adaptations as those 'adopted at a much larger scale or intensity, those that are truly new to a particular region or resource system, and those that transform places and shift locations', and they can be responsive or anticipatory. Moreover, they set two conditions for implementation; whether there is large vulnerability in certain regions, populations, or resource systems; and whether climate change threatens to overwhelm even robust human systems.

The eastern coast of Tōhoku fulfils Kates *et al*'s criteria for transformative adaptation. Ageing and depopulation are increasing community vulnerability and, because these settlements are permanently at risk from tsunami, climate change driven sea level rise has potential to contribute to suddenly overwhelming defence infrastructure. Consequently, it is reasonable to ask whether and how government planning anticipates sea level rise in settlement (re)construction in Tōhoku in preparation for the next giant tsunami.

We analysed 181 post-3/11 basic reconstruction plans, plus updates and supplementary documentation, from the websites of the Reconstruction Agency; Iwate, Miyagi and Fukushima Prefectures; and all 36 coastal municipalities in eastern Tōhoku (Table 5).

Using NVivo 11 we coded for Japanese language terms and synonyms¹⁵ for climate change, global warming, greenhouse effect, sea level rise, seawall, and tidal breakwater, looking for mentions in combination to find whether authorities were incorporating climate change driven sea level rise into reconstruction planning.

We found one mention of climate change driven sea level rise in reference to protecting Matsushima Bay’s natural beauty and ecology. There were 96 mentions of global warming and synonyms in 19 documents, with the most numerous in Miyagi and Iwate Prefectures’ plans. None associated climate change with sea level rise, and there were no references to sea level rise as an additional hazard for tsunami defence infrastructure.

Table 5. References to seawalls and climate change driven sea level rise in Japanese reconstruction planning documents.

Japanese	English	Sources	Refs.	Document	Type of Activity (No. of mentions)
防波堤	Tidal breakwater	143	1414	Multiple	Multiple activities.
防波堤	Breakwater	65	269	Multiple	Multiple activities.
海面上昇	Sea level rise	1	1	Miyagi Pref. (2017: 172)	Matsushima Bay’s scenic beauty & ecology.
温暖化	Warming	1	1	Kamaishi City (2011: 51)	Forest restoration.
温暖化	Warming	1	5	Kuji City (2011: 58)	Activities to prevent global warming.
温暖化	Warming	1	1	Yamamoto Town (2012: 39)	Activities to prevent global warming.
温暖化	Warming	1	1	Natori Town (2011: 33)	Activities to prevent global warming.
温暖化	Warming	1	1	Matsushima Town (2011: 16)	Matsushima Bay’s scenic beauty & ecology.
温暖化	Warming	1	2	Ishinomaki City (2011: 93)	Towards a low carbon society. (2).
温暖化	Warming	1	1	Shinchi Town (2012: 25)	Reducing in energy consumption.
温暖化	Warming	1	1	Naraha Town (2016: 124)	Towards a low carbon society.
温暖化	Warming	1	7	Iwate Prefecture (2017: 117)	Smart community creation and global warming countermeasures (7).
温暖化	Warming	1	58	Miyagi Pref. (2017: Passim)	Sustainable agriculture etc. (34), Education (9), Resources (12), Housing (2), Conservation (1).
温暖化	Warming	1	4	Miyagi Pref. (2011: Passim)	Energy use, conservation (2), Forest thinning (1), Global warming prevention (1).
温暖化	Warming	1	1	Fukushima Prefecture (2011: 32)	Renewable energy industries (1).
温暖化	Warming	1	1	Central Govt Basic Law	Environmental protection (1).
温室効果	Greenhouse effect	1	2	Sendai City (2011: 41)	Reducing GHGs (2).
温室効果	Greenhouse effect	1	1	Naraha Town (2016: 124)	Towards a low carbon society.
温室効果	Greenhouse effect	1	1	Iwate Prefecture (2011: 97)	Towards a low carbon society.
温室効果	Greenhouse effect	1	5	Miyagi Pref. (2017: 455)	Carbon Offsetting, GHG emissions reduction (4).
温室効果	Greenhouse effect	1	1	Fukushima Pref. (2011: 35)	GHG emissions reductions.
気候変動	Climate change	1	1	Okuma Town (2012: 34)	Agricultural support.
異常気象	Abnormal weather	1	1	Matsushima Town (2011: 16)	Matsushima Bay’s scenic beauty & ecology.

Sources: Central, prefectural and local government websites.

Although combined scenario modelling of sea level rise and tsunami risks is still rare, the scientific literature now takes it seriously. Gica and Reynolds (2012) demonstrate an increased risk of coastal inundation at Midway from combined sea level rise and Mw8+ tsunami

occurring in the northwestern Pacific. And Ramanamurthy *et al* (2005: 1740) advise that:

... for long-term human settlement planning, safe elevations along with distances from the coast or vulnerability lines for human settlement, especially in low-lying areas need to be determined taking in to account other future probabilistics such as occurrence of epicentre of tsunamigenic earthquake ... anticipated sea level rise due to global warming ... etc.

Probabalistic Tsunami (Multi-)Hazard Assessment (PTHA) is an important development in disaster risk reduction science, with most studies appearing after the 2004 Indian Ocean tsunami. Its purpose is to understand tsunami hazard to inform risk reduction activities by generating hazard maps or curves which estimate ‘the probability of exceeding specific levels of tsunami intensity metrics (e.g., runup or maximum inundation heights) (Grezio *et al*, 2017: 1158). Despite some criticism of its validity (Mulargia, Stark & Geller, 2017; Stein, Geller & Liu, 2012), PTHA is widely used for tsunami defence planning because it is globally applicable to tsunami with differing generating sources and return periods, and can incorporate epistemic variation, such as forecasts for sea level rise (Grezio *et al*, 2017; Dall’Osso *et al*, 2014). Even so, ‘tsunami risk assessments have mostly failed to take in to consideration future sea level rise associated with enhanced anthropogenic climate change. As such, future tsunami may well be even worse than our current best estimates as inundation may change markedly as sea level rises’ (Dominey-Howes & Goff, 2013).

Although PTHA for eastern Tōhoku have been conducted for retrospective modelling and methodological development (Goda *et al*, 2015; De Risi & Goda, 2016) to date we are unable to locate studies where climate change driven sea level rise has been included as a risk variable.

Moreover, four officials from the Reconstruction Agency confirmed in an interview that the government had not included climate change driven sea level rise among its variables for advising prefectures and local governments in Tōhoku on coastal defence infrastructure 'because sea level is only rising very slowly'. This begs the question whether Japanese authorities should build estimates for sea level rise into their scenario modelling and/or reconstruction and tsunami resilience planning in Tōhoku.

Internationally, acknowledgement of climate change as an additional risk for tsunami vulnerability is increasing (See: Hinkel *et al*, 2018). For example, Dall'Osso *et al* (2014) suggest that Sydney's 'existing exposure to earthquake-generated tsunamis is relatively low but would increase significantly under higher sea level conditions caused by combinations of tide and/or sea level rise'. In New Zealand, Tonkin and Taylor Ltd.'s consultancy report for Wellington City Council (Tonkin & Taylor Ltd., 2013: 63) assumes increased risk exposure and that '[I]ncreased sea levels may also mean greater water depths closer to the shore that can increase near shore wave heights and may also modify tsunami impacts' (2013: 10). In 2004 Otago Regional Council collected LiDAR data which 'enabled a detailed hydrodynamic model study to be undertaken of the risk of tsunami and storm-related inundation for the entire region, including an assessment of the potential effects of future sea level rise' (NZME, 2009: 15). Indeed, New Zealand's government is moving from deterministic perspectives of disasters as single events, towards an approach that understands disasters within a dynamic system of sudden events and long-term processes (NZME, 2009). As an ageing and shrinking society prone to tsunami, Tōhoku suits this approach.

Among PTHA performed in regions in Asia with similar physical characteristics to Tōhoku that

do include sea level rise, Li *et al* (2018) conducted 1,886 scenarios for Macao, which is vulnerable to tsunami generated by megathrust earthquakes originating in the Manila Trench. Their results indicate a sea level rise of '0.5m (by 2060) and 1m (by 2100) would dramatically increase the frequency of tsunami-induced flooding incidences by a factor of 1.2 to 2.4 and 1.5 to 4.7, respectively'. Li *et al* also found that 'inundation depths increase significantly', noting depths of up to '0.2 to 0.5m greater than the 0.5 and 1m increased sea level'. Their 'worst case scenario' modelled a Mw9.3 rupture across three earthquake zones along the Manila Trench, describing its similarity to the 2011 Tōhoku earthquake, showing extensive inundation. Finally, Li *et al* emphasise that they use conservative estimates for sea level rise - that current research suggests a 1.5m rise by 2100 for Macau - and that they used mean high-water as their tidal variable, stressing that coastlines with a larger tidal range might have a vulnerability greater than their scenarios suggest. The ria coastline of Iwate Prefecture is one such coastline.

The Tarō Fishing Cooperative's port tower measures the Shōwa Sanriku, Meiji Sanriku, and 2011 tsunami at 10.0, 15.0, and 17.3m. Tarō has therefore experienced two, possibly three, Level 2 tsunami causing mass fatalities within 124 years. The Japanese authorities have not included estimates for the effects of climate change on sea levels and future tsunami height and frequency, despite scientific research emerging that sea level rise places coastal communities at elevated risk. It is possible that, not only will Tarō experience an increased incidence of tsunami, including Level 2 tsunami, within the next 50-100 years, but that earthquakes which in the past would have caused a Level 1 tsunami of 7-10m may in future create Level 2 events greater than 10m. In turn, Level 2 tsunami likely may be higher than current estimates for safe overtopping of sea defences, increasing the risk of a repeat of 2011 when the majority of tsunami defence

infrastructure was ineffective, with disastrous consequences for vulnerable ageing and shrinking communities like Tarō. Using Li *et al* (2018) and the Tarō fishing cooperative's measures as a base, it is possible to anticipate that a Mw9.0 earthquake as occurred in 2011, with the addition of 1.5m in sea level within the next 100 years, would generate a 19+m Tsunami in Tarō, overtopping the highest point of the new sea defences by more than 4m.

Concluding Discussion

The defences nearing completion in Tarō are stronger than previously. In the event of a Level 1 tsunami they will likely prevent considerable damage and save lives. Tarō's new layout is also safer and some transformational adaptation to living space is being implemented, but this is not in response to climate change. Notwithstanding, we have shown that it is not unimaginable for sea level to rise such that some Level 1 tsunami events will become Level 2, and for Level 2 events to become large enough potentially to overwhelm defences, just as the 'unimaginable' tsunami of 11 March 2011 overwhelmed what were then the strongest defences ever built in Tarō and Kamaishi. Consequently, we argue that the government's rationale for the construction of the wall is potentially insufficient, given the accumulated risks of another powerful undersea earthquake occurring in the region, global warming driven sea level rise changing the dynamics of tsunami in unpredictable ways, and coastal communities will be considerably different in their size and composition in the decades to come. Nevertheless, important though the social capital perspective is in demanding the development of systems of mutual aid in local communities at risk of disaster, it is necessarily one part of implementing a disaster prevention strategy that responds to the whole spectrum of risks involved in complex techno-environmental

events and acknowledges the need for transformational adaptation to the impacts of climate change.

At the start of this article we argued that discussions of climate change mitigation are beyond our scope. In many respects, however, mitigation is necessarily part of any truly transformational adaptation strategy since, for example, climate change requires radical decarbonisation in response. Indeed, does not transformational adaptation therefore require a reconfiguration of the social and economic system beyond removing fossil fuels towards a deprioritisation of growth in favour of resilience and sustainability? Hence, instead of understanding depopulation as a problem in search of more people, even a social crisis, isn't it time that Japan looks to the opportunities it presents in imagining a different future - a 'depopulation dividend' (Matanle, 2017)?

In April 2011, then Prime Minister Kan Naoto appointed the Reconstruction Design Council to produce a 'grand vision' for Tōhoku's reconstruction which would not be 'stuck within a traditional framework' and would be 'highly regarded in history'. Moreover, plans should have local ownership and result in creative outcomes, should emphasise 'harmony among nature, human beings and technology', provide 'clean energy' solutions, and they should inspire hope for the future (Prime Minister's Office, 2011). This was a call for transformational adaptation both of coastal communities in Tōhoku, where incrementalism had been insufficient in preventing mass fatalities, as well as Japan's energy system towards one that is less centralised, is ecologically sustainable, and which would be resilient to disasters. Five months later Kan was forced to resign.

Ever since the Meiji period (1868-1912), the use of concrete infrastructure to resolve contradictions between the logic of modern development and the perceived constraints of

nature has been deeply institutionalised in Japan. In response to the remodelling of the Tone and Watarase River basins as a result of the Ashio Copper Mine disaster (1880), the founder of Japan's environmental movement, Tanaka Shōzō, criticised the government's lack of imagination by arguing that it was not nature that was defective, and which needed correcting with the application of modern technocratic rationalism, but modern developmentalism, which placed humanity outside nature and, as others have also argued, at war with it (Carson, 1962: 7; Foster *et al*, 2010; Marcuse, 1972: 52). In so doing Tanaka placed humanity within nature and urged modern Japanese to adjust to its flow (*nagare*), rather than attempt to defend against it, lest they are undone by the force of its backflow (*gyakuryū*) (Littler, 2017; Stolz, 2007; 2014; Tanaka, 2004 & 2005).

Through the ages a common strategic error has been to prepare for the past by failing to imagine the future. European schoolchildren are taught France's catastrophic mistake in constructing the Maginot line in the 1930s, for example, which was built to defend against the type of assault by Germany that had occurred in 1914, but resulted in the encirclement and near annihilation of the allied forces at Dunkirk, and effectively allowed France's defeat and occupation. Nearly a century later and facing the twin 21st century challenges of depopulation and global warming, Japan is potentially in the vanguard of a transformation to a sustainable post-growth order in Asia (See: DeWit, 2015; Matanle, 2017). However, by adopting an incremental approach to disaster prevention in Tōhoku and its reluctance to imagine the consequences of climate change, the Japanese state demonstrates that it is yet to seize that opportunity. Instead, the Construction State (Feldhoff, 2002; McCormack, 2002) and its logic of modern developmentalism as a defensive bulwark against the assumed destructive encroachments of nature remains in place.

Sea-levels will likely rise more rapidly and higher than much of the science on climate change at present suggests. Recent discoveries in the arctic permafrost show thawing 'exceeding modelled future thaw depths for 2090 under IPCC RCP 4.5' (Farquharson *et al*, 2019), in other words 70 years in advance of that projected by the United Nations official documentation on which are based the international treaties that Japan has signed into law and relies upon to help guide its adaptation policy. Consequently, even the most highly regarded coastal defence projects, such as those in the Netherlands and Venice, may have shorter lives than publicly acknowledged due to the rapid onset of climate change and its impacts on human-environmental relationships.

It is disconcerting for us to observe the lack of acknowledgement of sea level rise among governments and engineers, whilst planning projects intended to last for more than 50 years. Humanity has a limited time window to mitigate and adapt to the impacts of global warming. We have known for many decades, if not a century or more, that environmental systems are sensitive to human activity, and that positive feedback loops can exacerbate and accelerate changes already under way. At the time that Tanaka Shōzō was campaigning to save his village, Yanaka, from the Japanese government's remodelling of the Kanto watershed, and building on the 19th century work of Fourier, Pouillet, Foote, and Tyndall, in 1896 Swedish scientist Svante Arrhenius made the first quantitative prediction of global warming under a doubling of atmospheric carbon dioxide (Arrhenius, 1896). In the 21st century, as younger people imagine a much-altered future for themselves (Thunberg, 2019), we would all do well to heed Tanaka's words.

Acknowledgements

The authors would like to thank the Japan Foundation Endowment Committee for

fieldwork funding for this research, and Professor Kenji Tani (<http://en.saitama-u.ac.jp/research/researchers/dr-kenji-tani/>) of Saitama University for use of topographic data from his website (<http://ktgis.net/>).

References

- AIJ (Architectural Institute of Japan). 2012. *Hisai 3 ken no kaigan teibōkō no settei jōkyō* [Information on the plans of the coastal embankment heights in the disaster hit 3 prefectures], Tokyo: Architectural Institute of Japan. Online (<https://www.aij.or.jp/jpn/databox/2012/20111115.pdf>). (Japanese).
- Aldrich, D.P. 2017. Trust deficit: Japanese communities and the challenge of rebuilding Tohoku, *Japan Forum*, 29 (1), 39-52.
- Aldrich, D.P. 2019. *Black Wave: How Networks and Governance Shaped Japan's 3/11 Disasters*, Chicago, IL: Chicago University Press.
- Aldrich, D.P. & Sawada, Y. 2015. The physical and social determinants of mortality in the 3.11 tsunami, *Social Science and Medicine*, 124, 66-75.
- Aldrich, D.P., Lipsky, P.Y. & McCarthy, M.M. 2019. Japan's Opportunity to Lead, *Nature Climate Change*, 9 (July), 492.
- Arrhenius, S. 1896. On the Influence of Carbonic Acid in the Air upon the Temperature of the Ground, *Philosophical Magazine and Journal of Science*, 5 (41) 251, 237-276. Online at the Royal Society of Chemistry website (https://www.rsc.org/images/Arrhenius1896_tcm18-173546.pdf).
- Bird, W., 2013. Post-Tsunami Japan's Push to Rebuild Coast in Concrete, *The Asia-Pacific Journal: Japan Focus*, 11 (2), 1. Online (<https://apjff.org/2013/11/21/Winifred-Bird/3945/article.html>).
- Bridges, T.S. *et al.* 2015. *Use of Natural and Nature-based Features for Coastal Resilience*. Washington, DC: US Army Corps of Engineers. Online (<http://www.nad.usace.army.mil/Portals/40/docs/NACCS/NNBF%20FINAL.pdf>).
- Carrington, D. 2019. Why the Guardian is changing the language it uses about the environment, *The Guardian*. 17 May. Online (<https://www.theguardian.com/environment/2019/may/17/why-the-guardian-is-changing-the-language-it-uses-about-the-environment>).
- Carson, R. 1962. *Silent Spring*. New York: Houghton Mifflin.
- Coulmas, F. 2007. *Population Decline and Ageing in Japan: The Social Consequences*, London: Routledge.
- Cyranoski, D. 2012. After the Deluge: Japan is Rebuilding its Coastal Cities to Protect People from the Biggest Tsunamis, *Nature*, 483, 141-143.
- Dall'Osso, F., Dominey-Howes D, Moore C., Summerhayes S., & Withycombe G. 2014. The exposure of Sydney (Australia) to earthquake-generated tsunamis, storms and sea level rise: a probabilistic multi-hazard approach, *Scientific Reports*, 4, 7401, DOI:10.1038/srep07401.
- Davis, C., Kelis-Borok, V., Kossobokov, V. and Soloviev, A. 2012. Advance prediction of the March 11, 2011 Great East Japan Earthquake: A missed opportunity for disaster preparedness, *International Journal of Disaster Risk Reduction*, 1, 17-32.
- DeConto, R.M, and Pollard, D. 2016. Contribution of Antarctic to Past and Future Sea level Rise, *Nature*, 531, 591-597.

- De Risi, R. & Goda, K. 2016. Probabilistic Earthquake-Tsunami Multi-Hazard Analysis: Application to the Tohoku Region, Japan, *Frontiers in Built Environment*, 2 (19), 1-19.
- DeWit, A. 2014. Japan's Resilient, Decarbonizing and Democratic Smart Communities, *The Asia-Pacific Journal: Japan Focus*, 12 (50), 3. Online (<https://apjif.org/2014/12/50/Andrew-DeWit/4236.html>).
- DeWit, A. 2015. Japan's Bid to Become a World Leader in Renewable Energy, *The Asia-Pacific Journal: Japan Focus*, 13 (40), 2. Online (<https://apjif.org/-Andrew-DeWit/4385>).
- DeWit, A. 2016a. Japan's "National Resilience" and the Legacy of 3-11, *The Asia-Pacific Journal: Japan Focus*, 14 (6), 1. Online (<https://apjif.org/2016/06/DeWit.html>).
- DeWit, A. 2016b. Hioki's Smart Community and Japan's Structural Reform, *The Asia-Pacific Journal: Japan Focus*, 14 (15), 10. Online (<https://apjif.org/2016/15/DeWit.html>).
- Dominey-Howes, D. & Goff, J. 2013. Tsunami Risk Management in Pacific Island Countries and Territories (PICTs): Some Issues, Challenges and Ways Forward, *Pure and Applied Geophysics*, 170, 1397-1413.
- EPA (United States Environmental Protection Agency). 2017. Climate Change Adaptation Resource Center (ARC-X). Online. Available from here (<https://www.epa.gov/arc-x>).
- Farquharson, L.M., Romanovsky, V.E., Cable, W.L., Walker, D.A., Kokelj, S.V., & Nicolsky, D. 2019. Climate change drives widespread and rapid thermokarst development in very cold permafrost in the Canadian High Arctic, *Geophysical Research Letters*, 10 June. DOI: 10.1029/2019GL082187.
- Fahy, F. & Rau, H. 2013. Researching Complex Sustainability Issues: Reflections on Current Challenges and Future Developments. In: F. Fahy and H. Rau (eds.) *Methods of Sustainability Research in the Social Sciences*, London: Sage, 193-208.
- Feldhoff, T. 2002. Japan's Construction Lobby Activities - Systemic Stability and Sustainable Regional Development, *ASIEN*, 84, 34-42.
- Foster, J.B., Clark, B., & York, R. 2010. *The Ecological Rift: Capitalism's War on the Earth*, London: Monthly Review Press.
- Furuta, N. & Seino, S. 2016. Progress and Gaps in Eco-DRR Policy and Implementation After the Great East Japan Earthquake. In F.G. Renaud, K. Sudmeier-Rieux, M. Estrella & U. Nehren (eds.) *Ecosystem-Based Disaster Risk Reduction and Adaptation in Practice*, Switzerland: Springer, 295-314.
- GBD 2017 Causes of Death Collaborators. 2018. Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980-2017: a systematic analysis for the Global Burden of Disease Study 2017, *The Lancet*, 392: 1736-1788.
- Gica, E., & Reynolds, M. 2012. Incorporating Tsunami Projections to Sea Level Rise Vulnerability Assessments - A Case Study for Midway Atoll, American Geophysical Union, Fall Meeting 2012, abstract #NH23C-07, December.
- Goda, K., Li, S., Mori, N., & Yasuda, T., 2015. Probabilistic Tsunami Damage Assessment Considering Stochastic Source Models: Application to the 2011 Tohoku Earthquake, *Coastal Engineering Journal*, 57 (3), DOI: 10.1142/S0578563415500151.
- Graham, E. 1999. Breaking Out: The Opportunities and Challenges of Multi-Method Research in Population Geography, *The Professional Geographer*, 51 (1), 76-89.

- Grezio, A. *et al* (2017) Probabilistic Tsunami Hazard Analysis: Multiple Sources and Global Applications, *Reviews of Geophysics*, 55, 1158-1198.
- Hansen, J. 2009. *Storms of My Grandchildren: The Truth About the Coming Climate Catastrophe and Our Last Chance to Save Humanity*, London: Bloomsbury Press.
- Hansen, J.E. *et al*. 2016. Ice Melt, Sea Level Rise and Superstorms: Evidence from Paleoclimate Data, Climate Modeling, and Modern Observations that 2°C Global Warming Could be Dangerous, *Atmospheric Chemistry and Physics*, 16, 3761-3812.
- Hayashi, N. 2012. Jishin tsunami ni yoru Tarō-chō no higai [The effects of the earthquake and tsunami on Tarō Town], *Tōkyō Rigaku Daigaku, Tsujimoto Kenkyūshitsu [Tokyo Science University, Tsujimoto Research Centre]*, 5109421. (Japanese).
- Hill, K. 2015. Coastal Infrastructure: a Typology for the Next Century of Adaptation to Sea-level Rise, *Frontiers in Ecology and the Environment*, 13 (9), 468-476.
- Hinkel, J. *et al*. 2018. The Ability of Societies to Adapt to Twenty-First-Century Sea-level Rise, *Nature Climate Change*, 8 (7), 570-578.
- IPCC (Intergovernmental Panel on Climate Change). 2007. *Climate Change 2007: Synthesis Report*, Summary for Policymakers, Geneva: IPCC. Online (https://www.ipcc.ch/pdf/assessment-report/ar4/syr/ar4_syr_spm.pdf).
- IPCC (Intergovernmental Panel on Climate Change). 2014. *Climate Change 2014: Synthesis Report*, Summary for Policymakers. Online. Available from here (https://www.ipcc.ch/pdf/assessment-report/ar5/syr/AR5_SYR_FINAL_SPM.pdf).
- IMBIE Team (The). 2018. Mass balance of the Antarctic Ice Sheet from 1992 to 2017, *Nature*, 558, 219-222.
- Inoue, T. 2013. Tarō chō ni okeru bōchōtei to shumun no shibōritsu [The effect of the seawall on the death rate of the residents of Tarō-chō], *Tōkyō Rigaku Daigaku, Tsujimoto Kenkyūshitsu* [Tokyo University of Science, Tsujimoto Research Centre], 5109354. (Japanese).
- Jackson, N. & Brabyn, L. 2017. The Mechanisms of Subnational Population Growth and Decline in New Zealand 1976-2013, *Policy Quarterly*, 13 (Supplementary Issue), 22-36.
- Jackson, N. & Felmingham, B. 2002. As the Population Clock Winds Down: Indicative Effects of Population Ageing in Australia's States and Territories, *Journal of Population Research*, 19 (2), 97-117.
- JMA. 2011a. Tsunami jōhō: tsunami tōtatsu yosō jikoku to yosōsareru tsunami no takasa ni kansuru johō [Tsunami Information: Estimated Tsunami arrival time and Height] (Issued at 14:46 JST, 11 March 2011). Tokyo, Japan: Japan Meteorological Agency. Online (http://www.jma.go.jp/jp/tsunami/info_04_2011_0311145026.html). (Japanese).
- Japan Times. 2011. 90% of Disaster Casualties Drowned. *Japan Times Online*, 21 April. Online (<http://www.japantimes.co.jp/news/2011/04/21/national/90-of-disaster-casualties-drowned/>).
- JMA. 2011b. Tsunami jōhō: tsunami tōtatsu yosō jikoku to yosōsareru tsunami no takasa ni kansuru johō [Tsunami Information: Estimated Tsunami arrival time and Height] (Issued at 15:31 JST, 11 March 2011). Tokyo, Japan: Japan Meteorological Agency. Online (http://www.jma.go.jp/jp/tsunami/info_04_2011_0311153109.html). (Japanese).
- Kanda, J. 2016. Consideration for Effective Height of Sea Walls against Tsunami, *Structure and Infrastructure Engineering*, 12 (4), 484-489.

- Kates, R.W., Travis, W.R., & Wilbanks, T.J. 2012. Transformational Adaptation when Incremental Adaptations to Climate Change are Insufficient, *Proceedings of the National Academy of Sciences*, 109 (19), 7156-7161.
- Klein, K.R. & Nagel, N.E. 2007. Mass Medical Evacuation: Hurricane Katrina and Nursing Experiences at the New Orleans Airport, *Disaster Management & Response*, 5 (2), 56-61.
- Koshimura, S., Hayashi S. & Gokon, H. 2014. The Impact of the 2011 Tohoku Earthquake Tsunami Disaster and Implications to the Reconstruction, *Soils and Foundations*, 54 (4), 560-572.
- Lay, T. 2015. The Surge of Great Earthquakes from 2004 to 2011, *Earth and Planetary Science Letters*, 409, 133-146.
- Li, L., Switzer, A.D., Wang, Y., Chan, C-H., Qiu, Q. & Weiss, R. 2018. A modest 0.5-m rise in sea level will double the tsunami hazard in Macau, *Science Advances*, 4 (8), DOI: 10.1126/sciadv.aat1180.
- Littler, J., 2017. *The Making of an Alternative Religion in Late-Meiji Japan (1899-1912), A Study of Arai Ōsui (1846-1922)*, Unpublished MSc Dissertation, University of Oxford.
- Madsen, L.M. & Adriansen, H.K. 2004. Understanding the Use of Rural Space: The Need for Multi-Methods, *Journal of Rural Studies*, 20 (4), 485-497.
- Marcuse, H. 1994 [1972]. Ecology and Revolution. In C. Merchant (ed.) *Ecology: Key Concepts in Critical Theory*, Atlantic Highlands, NJ: Humanities Press.
- Matanle, P. 2006. Organic Sources for the Revitalization of Rural Japan: The Craft Potters of Sado, *Japanstudien*, 18, 149-180.
- Matanle, P. 2011. The Great East Japan Earthquake, Tsunami and Nuclear Meltdown: Towards the (Re)construction of a Safe, Sustainable, and Compassionate Society in Japan's Shrinking Regions, *Local Environment*, 16 (9), 823-847.
- Matanle, P. 2013. Post-disaster recovery in ageing and declining communities: the Great East Japan disaster of 11 March 2011, *Geography*, 98 (2), 68-76.
- Matanle, P. 2014. Ageing and Depopulation in Japan: Understanding the Consequences for East and Southeast Asia in the 21st Century. In H. Dobson (ed.) *East Asia in 2013: A Region in Transition*, White Rose East Asia Centre and Foreign and Commonwealth Office Briefing Papers, Sheffield: WREAC: 30-35.
- Matanle, P. 2017. Towards an Asia-Pacific 'Depopulation Dividend' in the 21st Century: Regional Growth and Shrinkage in Japan and New Zealand, *The Asia-Pacific Journal: Japan Focus*, 15 (6), 5. Online (<http://apjjf.org/2017/06/Matanle.html>).
- Matanle, P., Rausch, A.S., with the Shrinking Regions Research Group. 2011. *Japan's Shrinking Regions in the 21st Century: Contemporary Responses to Depopulation and Socioeconomic Decline*. Amherst: Cambria Press.
- McCormack, G. 2002. Breaking the Iron Triangle, *New Left Review*, 5-23.
- McMillan, R. 2014. A Disaster Waiting to Happen? Elderly People and Population Ageing in at Risk Locations. Presentation to the Australian Population Association, Hobart, 3-5 December, Online (https://www.academia.edu/16307991/A_disaster_waiting_to_happen_Elderly_people_and_population_ageing_in_at_risk_locations).
- MIC (Ministry of Internal Affairs and Communications). 2018. Japan Statistical Yearbook 2018, Statistics Bureau. Online (<http://www.stat.go.jp/english/data/nenkan/67n>

enkan/index.html).

Mikami, T., Shibayama, T., Esteban, M. & Matsumaru, R. 2012. Field Survey of the 2011 Tohoku Earthquake and Tsunami in Miyagi and Fukushima Prefectures, *Coastal Engineering Journal*, 54 (1), 1-26.

Minoura, K., Imamura, F., Sugawara, D., Kono, Y. & Iwashita, T. 2001. The 869 Jogan Tsunami Deposit and Recurrence Interval of Large-scale Tsunami on the Pacific Coast of Northeast Japan, *Journal of Natural Disaster Science*, 23, 83-88.

Miyako City. 2011. Tarō chiku fukkō machizukuri keikaku [Tarō District reconstruction and town-making plan], Miyako City Office, 21 December. Online (http://www.city.miyako.iwate.jp/data/open/cnt/3/713/1/03-06siryo4_image.pdf). (Japanese).

MLIT (Ministry of Land, Infrastructure and Transportation). 2011. *Kasen, kaigan kōzōbutsu no fukkyū ni okeru kaikan hairyo no tebiki* [For the restoration of waterways and coasts, guidance on considering the scenery]. Tokyo, Japan: Ministry of Land, Infrastructure and Transportation. (Japanese).

MLIT (Ministry of Land, Infrastructure and Transportation). 2014. Grand Design of National Spatial Development towards 2050, Japan Creation of a country generating diverse synergies among regions. Tokyo, Japan: Ministry of Land, Infrastructure and Transportation. Online (<http://www.mlit.go.jp/common/001088248.pdf>).

MLIT (Ministry of Land, Infrastructure and Transportation). 2015. National Spatial Strategy (National Plan), August 2015. Tokyo, Japan: Ministry of Land, Infrastructure and Transportation. Online (<http://www.mlit.go.jp/common/001127196.pdf>).

Mulargia, F., Stark, P.B. & Geller, R.J. 2017. Why is Probabilistic Seismic Hazard Analysis (PSHA) Still Used? *Physics of the Earth and Planetary Interiors*, 264 (March), 63-75.

Muramatsu, H., Ando, A., Igarashii, H. & Akatani, R. 1991. The construction of tsunami control embankments and the enlargement of an urban area - In the case of Taro Town located on Sanriku coast at Iwate prefecture which is often attacked by tsunamis, *Historical Studies in Civil Engineering*, 11, 85-94.

NACS-J (Nature Conservation Society of Japan). 2013. *Kono mama de ii-no-ka? Bōchōtei keikaku* [Is this way OK? Seawall plans], *Shizen Hogo*, 534, 2-14. (Japanese).

NASC (National Association of Sea Coast). 2014. *Kaigan wo meguru genjō to kadai* [Current Issues and Situation of the Coastline]. Tokyo, Japan: National Association of Sea Coast. Online (http://www.kaigan.or.jp/business/pdf/kaiganmeguru_h26.pdf). (Japanese).

Nakahara, S. 2011. Lessons Learnt from the Recent Tsunami in Japan: Necessity of Epidemiological Evidence to Strengthen Community-based Preparation and Emergency Response Plans, *Injury Prevention*, 17 (6), 361-364.

Nerem, R.S., Beckley, B.D., Fasullo, J.T., Hamlington, B.D., Masters, D. & Mitchum, G.T. 2018, Climate-change-driven Accelerated Sea level Rise Detected in the Altimeter Era, *Proceedings of the National Academy of Sciences*, 115 (9), 2022-2025.

New York Times (2019) Where the Climate Change Action is, 19 June. Online (<https://www.nytimes.com/2019/06/14/opinion/climate-change-states-trump.html>).

Ngo, E.B. 2012. Elderly People and Disaster. In B. Wisner, J.C. Gaillard, & I. Kelman (eds) *The Routledge Handbook of Hazards and Disaster*

- Risk Reduction*, London: Routledge, 413-423.
- Nikkei Shimbun. 2011. Banri no chōjō ha nokotta ga... [The Great Wall of Japan remains, but...], *Nihon Keizai Shimbun*, 4 April. Online (http://www.nikkei.com/article/DGXNASFK01023_R00C11A4000000/). (Japanese).
- NOAA (National Oceanic and Atmospheric Administration). 2019. NGDC/WDS Global Historical Tsunami Database. Online (https://www.ngdc.noaa.gov/hazard/tsu_db.shtml).
- Normile, D. 2012. One Year After the Devastation, Tohoku Designs Its Renewal, *Science*, 335 (6073), 1164-1166.
- NPA (National Police Agency). 2016. Heisei 23-nen (2011-nen) Tōhoku chihō Taiheiyō oki no higai jōkyō to keisatsu sochi [Police Report of the Damage Situation off the Pacific Coast of Tōhoku]. NPA, 9 December. Online (<http://www.npa.go.jp/archive/keibi/biki/higaijokyo.pdf>). (Japanese).
- NZME (New Zealand Ministry for the Environment). 2009. *Preparing for Coastal Change: A Guide for Local Government in New Zealand*, Ministry for the Environment, ME 907, Wellington: New Zealand Government.
- Ogasawara, T. & Sakai, S. 2012. Characteristics on Damages of 2011 Tohoku Tsunami on Iwate Coast, *Japanese Journal of Multiphase Flow*, 26 (1), 28-35. (Japanese).
- Onishi, N. 2011. In Japan, Seawall Offered a False Sense of Security, *New York Times*. Online (<http://www.nytimes.com/2011/04/02/world/asia/02wall.html>).
- Ozaki, T. 2011. Outline of the 2011 off the Pacific Coast of Tohoku Earthquake (Mw 9.0) Tsunami Warnings, Advisories and Observations, *Earth Planets Space*, 63, 827-830.
- Prime Minister's Office. 2011. Press briefing at the Prime Minister's Office for Members of the Foreign Press, Prime Minister's Office, 14 April. Online (<http://japan.kantei.go.jp/incident/pdf/foreign-press-briefing-20110414-script-e.pdf>).
- Raby, A., Macabuag, J., Pomonis, A., Wilkinson, S., & Rossetto, T. 2015. Implications of the 2011 Great East Japan Tsunami on Sea Defence Design, *International Journal of Disaster Risk Reduction*, 14 (4), 332-346.
- Rahmstorf, S. 2007. A Semi-Empirical Approach to Projecting Future Sea level Rise, *Science*, 315 (5810), 368-370.
- Ramanamurthy, M.V., Sundaramoorthy, S., Pari, Y. & Rao V.R. 2005. Inundation of Sea Water in Andaman and Nicobar Islands and Parts of Tamil Nadu Coast during 2004 Sumatra Tsunami, *Current Science*, 88 (11), 1736-1740.
- Rausch, A.S. 2012. A Framework for Japan's New Municipal Reality: Assessing the Heisei *gappei* Mergers, *Japan Forum*, 24 (2), 185-204.
- Reconstruction Agency. 2015. *Kōgyō keikaku (Iwate-Ken, Miyako-Shi) (Reconstruction plan, Iwate Prefecture, Miyako City)*, Tokyo, Japan: Reconstruction Agency. Online (http://www.reconstruction.go.jp/topics/main-cat1/sub-cat1-3/20150731_Iwate07Miyako.pdf). (Japanese).
- Reilly, J., *et al.* 2015. Energy & Climate Outlook: Perspectives from 2015, MIT Joint Program on the Science and Policy of Global Change. Online (<https://globalchange.mit.edu/sites/default/files/newsletters/files/2015%20Energy%20%26%20Climate%20Outlook.pdf>).
- Rogelj, J., *et al.* 2016. Paris Agreement Climate Proposals Need a Boost to Keep Warming Well Below 2°C, *Nature*, 534, 631-639.

- Samuels, R.J. 2013 Japan's Rhetoric of Crisis: Prospects for Change after 3.11., *Journal of Japanese Studies*, 39 (1), 97-120.
- Satake, K., Sawai, Y., Shishikura, M., Okamura, Y., Namegaya, Y., & Yamaki, S. (2007) Tsunami source of the unusual AD 869 earthquake off Miyagi, Japan, inferred from tsunami deposits and numerical simulation of inundation [online]. *American Geophysical Union*, Fall Meeting 2007, abstract no. T31G-03. Available from: <http://adsabs.harvard.edu/abs/2007AGUFM.T31G..03S> [Accessed 30 March 2016].
- Schaefer, K., Lantuit, H., Romanovsky, V.E., Schuur, E.A.G., & Witt, R. 2014. The Impact of the Permafrost Carbon Feedback on Global Climate, *Environmental Research Letters*, 9 (8). Online (<http://iopscience.iop.org/article/10.1088/1748-9326/9/8/085003/meta>).
- Scripps Institution. 2019. The Keeling Curve Hits 415 PPM. Scripps Institution of Oceanography Youtube Channel. Online (<https://www.youtube.com/watch?v=Z43FQCSg4Ow>).
- Shibayama, T. *et al.* 2013. Classification of Tsunami and Evacuation Areas, *Natural Hazards*, 67, 365-386.
- Shifleen, A. (2015) The 'Great Wall' of Japan - a Shield against Tsunamis. *Haveeru Online*, 2 October. Archived by Sanriku Fukkō, Deutsch-Japanisches Synergie Forum (DJSF). Online (<https://sanrikufukkou.com/2015/11/20/the-great-wall-of-japan-a-shield-against-tsunamis/>).
- Smits, G. 2014. *When the Earth Roars: Lessons from the History of Earthquakes in Japan*, Lanham, MD: Rowman & Littlefield.
- Stein, S., Geller, R.J., & Liu, M. 2012. Why earthquake hazard maps fail and what to do about it, *Tectonophysics*, 562-563 (August), 1-25.
- Stolz, R. 2007. Remake Politics, Not Nature: Tanaka Shozo's Philosophies of 'Poison' and 'Flow' and Japan's Environment, *Asia Pacific Journal: Japan Focus*, 5 (1) 2 January. Online (<https://apjff.org/-Robert-Stolz/2331/article.html>).
- Stolz, R. 2014. *Bad Water: Nature, Pollution and Politics in Japan, 1870-1950*, Durham, NC: Duke University Press.
- Strusińska-Correia, A. 2017. Tsunami mitigation in Japan after the 2011 Tōhoku Tsunami, *International Journal of Disaster Risk Reduction*, 22, 397-411.
- Suppasri, A., Shuto, N., Imamura, F., Koshimura, S., Mas, E. & Yalciner, A.C. 2013. Lessons Learned from the 2011 Great East Japan Tsunami: Performance of Tsunami Countermeasures, Coastal Buildings, and Tsunami Evacuation in Japan, *Pure and Applied Geophysics*, 170, 993-1018.
- Suppasri, A. *et al.* 2016. Improvement of Tsunami Countermeasures Based on Lessons from The 2011 Great East Japan Earthquake and Tsunami — Situation After Five Years, *Coastal Engineering Journal*, 58 (4), DOI: 10.1142/S0578563416400118.
- Takezawa, S. (2016) *The Aftermath of the 2011 East Japan Earthquake and Tsunami: Living among the Rubble*, Lanham, MD: Lexington Books.
- Tanaka, S. 2004 & 2005. *Tanaka Shōzō Bunshū* (Collected Writings of Shozo Tanaka) (Vols. 1 & 2). M. Yui & Y. Komatsu (eds.), Tokyo: Iwanami Shoten.
- Tokida, K. & Tanimoto, R. 2014. Lessons for Countermeasures Using Earth Structures against Tsunami Obtained in the 2011 Off the Pacific Coast of Tohoku Earthquake, *Soils and Foundations*, 54 (4), 523-543.
- Tonkin & Taylor Ltd., 2013. *REPORT:*

- Wellington City Council: Sea Level Rise Options Analysis*, Wellington: Wellington City Council. Online (<http://wellington.govt.nz/~media/services/environment-and-waste/environment/files/61579-wcc-sea-level-rise-options.pdf>).
- Thunberg, G. 2019. *No One is Too Small to Make a Difference*, London: Penguin.
- UNFCCC (United Nations Framework Convention on Climate Change). 2017. Paris Agreement - Status of Ratification, 21 April. Online (http://unfccc.int/paris_agreement/items/9444.php).
- Yamashita, F. 2003. Sanriku kaigan, Tarōchō ni okeru 'tsunami bōsai no machi sengen' to daibōchōtei no ryakushi [Sanriku Coast, Tarō Town. A short history of the huge seawall in the town declared tsunami-proof], *Rekishi Tsunami*, 19, 165-171. (Japanese).
- Yin, R.K. 2004. *The Case Study Anthology*, London: Sage.

Peter Matanle is Senior Lecturer and Director of Research and Innovation at the School of East Asian Studies, University of Sheffield, UK. His research and teaching interests are in the social and cultural geography of the Asia-Pacific, focusing on Japan.

Contact: p.matanle@sheffield.ac.uk

(<https://apjif.org/mailto:p.matanle@sheffield.ac.uk>)University Profile Page

(<https://www.sheffield.ac.uk/seas/staff/japanese/matanle>)Personal Website

(<http://www.peter-matanle.net/>)Publications and Talks

(<https://sheffield.academia.edu/PeterMatanle>)Twitter: @matapet

Joel Littler is a lecturer at the Pridi Banomyong International College, Thammasat University, Thailand, and a graduate of the School of East Asian Studies, University of Sheffield (BA, Japanese Studies, 2016), and Pembroke College, University of Oxford (MSc with Distinction, Japanese Studies, 2017).

Contact: joel.l@pbic.tu.ac.th (<https://apjif.org/mailto:joel.l@pbic.tu.ac.th>).

Oliver Slay is a PhD candidate in the Department of Bioscience and Chemistry, Sheffield Hallam University, UK. He is a graduate of the School of East Asian Studies, University of Sheffield (BA, Japanese and Korean Studies, 1993). After following a 20-year career in IT administration and software development (bespoke GIS/mapping software), he completed a BSc and MSc (Hallam) in Molecular and Cell Biology.

Notes

¹ Japan's whole eastern coastline is undergoing redevelopment of its tsunami defence infrastructure (NASC, 2014). Tōhoku (lit. East-North - 東北) includes the six northernmost prefectures of Honshu: Aomori, Akita, Iwate, Yamagata, Miyagi, and Fukushima.

² See Matanle (2011) for a description of the disaster and its immediate aftermath.

³ As of writing in 2018, these structures are nearing completion.

⁴ There is a long-running semantic debate as to the correct terminology to describe the

greenhouse effect and the Earth's climate. 'Climate change' and 'global warming' are both contested terms, though they are currently the most commonly used. Increasingly, terms such as 'climate breakdown', 'climate emergency' and 'climate crisis' are being used to refer to the former, and 'global heating' for the latter. In this article we use 'climate change' and 'global warming', though we prefer the newer terms. We do so from the perspective of not wanting semantic discussions to distract from the core message of our research. See Carrington (2019) for a deeper discussion of these terms' usage.

⁵ Recently Aldrich *et al* (2019) have been rather less enthusiastic about Japan's success with climate change mitigation.

⁶ A ria coastline comprises several roughly parallel inlets separated by mountain ranges. Unlike fjords, which are U-shaped inundated valleys scoured by glaciation, rias are unglaciated river valleys that subsequently flooded when the ice retreated and sea-level rose at the end of the last ice age. Consequently, they often have low-lying and sheltered alluvial fans at their head which are attractive for human settlement. Other examples include the western coast of Galicia, Spain, and the Marlborough Sounds at the northern end of South Island, New Zealand.

⁷ All photographs in this article © Peter Matanle.

⁸ See Rausch (2012) for a summary analysis of the Heisei municipal mergers.

⁹ The crude birth rate measures the average number of children born per 1,000 population, and the total fertility rate the average number of children born per woman over her lifetime. Currently in Japan the birth rate is falling, because the total number of women giving birth has dropped precipitously, even as the fertility rate has risen slightly.

¹⁰ National Oceanic and Atmospheric Administration.

¹¹ Defined as a tsunami causing at least one fatality in Japan.

¹² Information on public meetings and published planning documentation produced by Miyako City Office can be found on the City's website here (<http://www.city.miyako.iwate.jp/higashi.html>). Information specific to the reconstruction of Tarō District is archived by Miyako City here (<http://www.city.miyako.iwate.jp/index/sosiki/taro.html>).

¹³ 1960 Valdivia, Chile (Mw9.5); 1964 Prince William Sound, Alaska (Mw9.2).

¹⁴ Despite the Trump administration's disavowal of the 2015 Paris Agreement, and its policy of undoing the previous Obama administration's actions on climate change, considerable activity in climate mitigation and adaptation is accelerating at state and municipal levels, among some federal institutions, and some private and third sector organisations (See, for example, New York Times, 2019).

¹⁵ Weblio Thesaurus (<http://thesaurus.weblio.jp/>) was used to generate the following terms: Warming: 温暖化, 温室効果; Climate change: 気候変動, 気候変更, 気候変化, 異常気象; Sea level rise: 海面上昇, 水面上昇, 海水準変動, 海面水位変化, 海面変化; Sea wall: 防潮堤, 防波堤, 海壁, 波除, 波戸, 波止.