



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/148409/>

Version: Accepted Version

---

**Article:**

Tovo, A, Formentin, M, Suweis, S et al. (2019) Inferring macro-ecological patterns from local presence/absence data. *Oikos*, 128 (11). pp. 1525-1677. ISSN: 0030-1299

<https://doi.org/10.1111/oik.06754>

---

© 2019 The Authors. *Oikos* © 2019 Nordic Society *Oikos*. This is the peer reviewed version of the following article: Tovo, A. , Formentin, M. , Suweis, S. , Stivanello, S. , Azaele, S. and Maritan, A. (2019), Inferring macro-ecological patterns from local presence/absence data. *Oikos*, 128: 1641-1652, which has been published in final form at <https://doi.org/10.1111/oik.06754>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.


 OIKOS

## Inferring macro-ecological patterns from local presence/absence data

Journal:	<i>Oikos</i>
Manuscript ID	OIK-06754
Wiley - Manuscript type:	Research
Keywords:	biodiversity patterns, species-abundance distribution, species-accumulation curve, upscaling biodiversity patterns, spatial ecology
Abstract:	<p>1. Biodiversity provides support for life, vital provisions, regulating services and has positive cultural impacts. It is therefore important to have accurate methods to measure biodiversity, in order to safeguard it when we discover it to be threatened. For practical reasons, biodiversity is usually measured at fine scales whereas diversity issues (e.g. conservation) interest regional or global scales. Moreover, biodiversity may change across spatial scales. It is therefore a key challenge to be able to translate local information on biodiversity into global patterns.</p> <p>2. Many databases give no information about the abundances of a species within an area, but only its occurrence in each of the surveyed plots. In this paper, we introduce an analytical framework to infer species richness and abundances at large spatial scales in biodiversity-rich ecosystems when species presence/absence information is available on various scattered samples (i.e. upscaling).</p> <p>3. This framework is based on the scale-invariance property of the negative binomial. Our approach allows to infer and link within a unique framework important and well-known biodiversity patterns of ecological theory, such as the Species Accumulation Curve (SAC) and the Relative Species Abundance (RSA) as well as a new emergent pattern, which is the Relative Species Occupancy (RSO).</p> <p>4. Our estimates are robust and accurate, as confirmed by tests performed on both in silico-generated and real forests. We demonstrate the accuracy of our predictions using data from two well-studied forest stands. Moreover, we compared our results with other popular methods proposed in the literature to infer species richness from presence-absence data and we showed that our framework gives better estimates. It has thus important applications to biodiversity research and conservation practice.</p>

## Research

### 0 Inferring macro-ecological patterns from local presence/absence data 61

5 Anna Tovo, Marco Formentin, Samir Suweis, Samuele Stivanello, Sandro Azaele and Amos Maritan 65

10 A. Tovo (<https://orcid.org/0000-0003-3834-3749>) ✉ ([anna.tovo@unipd.it](mailto:anna.tovo@unipd.it)), S. Suweis and A. Maritan, *Dipto di Fisica e Astronomia 'Galileo Galilei', Istituto Nazionale di Fisica Nucleare, Univ. di Padova, Via Marzolo 8, IT-35131 Padova, Italy.* – M. Formentin and S. Stivanello, *Dipto di Matematica 'Tullio Levi-Civita', Univ. di Padova, Padova, Italy.* – S. Azaele, *Dept of Applied Mathematics, School of Mathematics, Univ of Leeds, Leeds, UK.* 75

#### Oikos

00: 1–12, 2019

20 doi: 10.1111/oik.06754

Editor-in-Chief: Dries Bonte

Accepted 21 June 2019

Biodiversity provides support for life, vital provisions, regulating services and has positive cultural impacts. It is therefore important to have accurate methods to measure biodiversity, in order to safeguard it when we discover it to be threatened. For practical reasons, biodiversity is usually measured at fine scales whereas diversity issues (e.g. conservation) interest regional or global scales. Moreover, biodiversity may change across spatial scales. It is therefore a key challenge to be able to translate local information on biodiversity into global patterns. 80

Many databases give no information about the abundances of a species within an area, but only its occurrence in each of the surveyed plots. In this paper, we introduce an analytical framework (implemented in a ready-to-use R code) to infer species richness and abundances at large spatial scales in biodiversity-rich ecosystems when species presence/absence information is available on various scattered samples (i.e. upscaling). 90

This framework is based on the scale-invariance property of the negative binomial. Our approach allows to infer and link within a unique framework important and well-known biodiversity patterns of ecological theory, such as the species accumulation curve (SAC) and the relative species abundance (RSA) as well as a new emergent pattern, which is the relative species occupancy (RSO). 95

Our estimates are robust and accurate, as confirmed by tests performed on both in silico-generated and real forests. We demonstrate the accuracy of our predictions using data from two well-studied forest stands. Moreover, we compared our results with other popular methods proposed in the literature to infer species richness from presence to absence data and we showed that our framework gives better estimates. It has thus important applications to biodiversity research and conservation practice. 100

Keywords: biodiversity patterns, spatial ecology, species–abundance distribution, species–accumulation curve, upscaling biodiversity patterns 110

#### Introduction 115

The problem of inferring total biodiversity when only scattered samples are observed is a long-standing problem. In the 1940s, the British chemist and naturalist A. S. Corbet spent two years in Malaya to trap butterflies (Corbet 1941). For every species 121



[www.oikosjournal.org](http://www.oikosjournal.org)

© 2019 The Authors. Oikos © 2019 Nordic Society Oikos

Note: This manuscript has undergone open peer review, accessible here: <https://ecology.peercommunityin.org/public/rec?id=23>.

he saw, he noted down how many individuals of that species he trapped. When Corbet returned to England, he showed the table to its colleague R. A. Fisher and asked him how many new species he would trap if he returned to Malaya for another couple of years. The father of statistics was only the first to tackle the problem of species estimation (Fisher et al. 1943), which since then has found large applications in different scientific fields, from ecology (Bunge and Fitzpatrick 1993, Colwell and Coddington 1994, Chao and Bunge 2002) to bioscience (Hughes et al. 2001, Ionita-Laza et al. 2009, Locey and Lennon 2016), leading to the development of a myriad of estimators (Good and Toulmin 1956, Chao and Chiu 2016, Orlitsky et al. 2016).

Indeed, although ecological drivers crucial for conservations act at large scales, biodiversity is typically monitored at limited spatial scales (Alonso et al. 2008, Bertuzzo et al. 2016). Extrapolating species richness from the local to the whole ecosystem scale is not straightforward, because it is not additive as a function of the area. As a result, a huge number of biodiversity estimators have been proposed in ecological literature (Bunge and Fitzpatrick 1993, Brose et al. 2003, Colwell et al. 2004, Mao and Colwell 2005, Ulrich and Ollik 2005, Wang and Lindsay 2005, Shen and He 2008, Bunge et al. 2012, Kunin et al. 2018). Their commonest limitation is to have a limited application range (local/regional-scale extrapolations), and to be sensitive to the trees' spatial distribution (Plotkin et al. 2000, Carrara et al. 2012, Azaele et al. 2016), sample coverage and sampling methods (Chao et al. 2009).

Many analytical methods have been proposed to upscale species richness using as input the local relative species abundance distribution (RSA) (Harte et al. 2009, Ter Steege et al. 2013, Slik et al. 2015, Tovo et al. 2017), i.e. the list of the species present at the sampled scale along with the proportion of individuals belonging to each of them. For example, estimates of biodiversity at large scales have been performed using log-series as the RSA (Fisher et al. 1943). The log-series distribution is often used to describe RSA patterns in many different ecological communities, characterised by high biodiversity (Azaele et al. 2016). Thanks to the availability and reliability of the species abundance data in forests (given by systematic and periodic field campaigns and high detectability of species), this method has been typically applied to tropical forests. In particular, it has been used to estimate the

species richness of the Amazonia (Ter Steege et al. 2013) and the global tropical tree species richness (Slik et al. 2015).

These methods have been proved to typically perform better than non-parametric estimators of biodiversity (Chao 2005). In contrast with the former, non-parametric approaches do not assume a specific family of probability distributions. In particular, non-parametric methods do not make any assumption on the RSA distribution and they thus perform no fit of empirical patterns, rather they only take into account rare species, which are intuitively assumed to carry all the needed information on the undetected species in a sample.

Nevertheless, all the aforementioned methods need abundance data in order to infer biodiversity at larger scale. However, in many open-access databases (e.g. species abundance data obtained from metagenomics (Thompson et al. 2017)) this information is highly imprecise, if available at all. Indeed, there are lots of datasets which give only information about the presence or absence of a species in different surveyed plots, without specifying the number of individuals within them. Some non-parametric approaches have been generalized to infer species richness from this presence to absence data (Chao 2005, Chao and Chiu 2016). Table 2 summarizes the most popular estimators and for each one details the predicted biodiversity as a function of the input data. However, most of them have the strong limitation that they do not have an explicit dependence of the observation scale, leading to poor estimates of the number of species at the global scales. The only estimator which takes into account the ratio between the surveyed area and the global one is the one introduced by Chao (Chao 2005, Chao et al. 2009, Chao and Chiu 2016) and denoted here as  $Chao_{wor}$  (Table 2). This method takes into account the number of species detected in one sample only and those detected in exactly two samples observed at the sample scale to infer the total species richness at the whole forest scale. However, it has been shown that Chao's method, although giving reliable species estimates, it does not properly capture the empirical species accumulation curve (SAC) (Tovo et al. 2017), which describes how the number of species changes across spatial scales. In absence of spatial correlation, it is equivalent to another macro-ecological pattern of interest which is the species area relation (SAR).

Moreover, both parametric and non-parametric methods proposed in the literature do not give any insights on the

Table 1. Predictive error for three generated forest (characterized by a log-normal and a negative binomial RSA) having individuals distributed according to a high clustering Thomas process and at random. Tests were performed by sampling a fraction  $p=0.05$  of each forest and by applying our framework (P/A columns) and the abundance-based method (RSA columns) to predict the true number of species  $S$  (5000 for the LN forest and 4974 for the NB forest). For each estimated  $S_{pred}$ , the average relative percentage error  $(S_{pred} - S)/S \times 100$  between the true number of species and the predicted one is shown together with the corresponding standard deviation. Results are relative to 100 iterations.

Forest RSA	Spatial distribution			
	Random		Thomas	
	P/A	RSA	P/A	RSA
Log-normal	$3.1 \pm 0.51$	$7.6 \pm 0.52$	$2.5 \pm 1.8$	$7.2 \pm 3.1$
Negative binomial	$-0.50 \pm 0.34$	$-0.52 \pm 0.28$	$-0.81 \pm 1.6$	$-0.60 \pm 1.7$

0 Table 2. Summary table of the most popular biodiversity estimators for presence/absence data. In formulas,  $M^*$  is the total number of sam- 61  
 5 pled cells. See Chao (2005), Chao and Chiu (2016) for more details about non-parametric methods.

Estimator	Predicted $S$	Details	
NB	$S^* \frac{1 - (1 - \xi)^r}{1 - (1 - \hat{\xi}_p)^r}$	$(\xi, r)$ NB parameters at $p = 1$ $(\hat{\xi}_p, r)$ NB parameters at $p^*$	65
Chao <sub>2</sub>	$S^* + \begin{cases} \frac{M^* - 1}{M^*} \frac{Q_1^2}{2Q_2} & Q_2 > 0 \\ \frac{M^* - 1}{M^*} \frac{Q_1(Q_1 - 1)}{2} & Q_2 = 0 \end{cases}$	$Q_i =$ number of species detected in $i$ plot at the scale $p^*$	70
iChao <sub>2</sub>	$S_{\text{Chao}_2} + \frac{M^* - 3}{4M^*} \frac{Q_3}{Q_4} \max\left(Q_1 - \frac{(M^* - 3)Q_2Q_3}{2(M^* - 1)Q_4}, 0\right)$	$S_{\text{Chao}_1} = S$ predicted by Chao <sub>1</sub> method	75
Chao <sub>wor</sub>	$S^* + \frac{Q_1^2}{\frac{M^*}{M^* - 1} 2Q_2 + \frac{p^*}{1 - p^*} Q_1}$		80
Jackknife <sub>1</sub>	$S^* + \frac{M^* - 1}{M^*} Q_1$		
Jackknife <sub>2</sub>	$S^* + \frac{2M^* - 3}{M^*} Q_1 - \frac{(M^* - 2)^2}{M^*(M^* - 1)} Q_2$		85
Turing	$S_{\text{abun}}^* + \frac{S_{\text{rare}}^*}{\hat{C}_{\text{rare}}}$	$S_{\text{abun}}^* = \sum_{n > 10} Q_n$ $S_{\text{rare}}^* = \sum_{n=1}^{10} Q_n$ $\hat{C}_{\text{rare}} = 1 - Q_1 / \sum_{n=1}^{10} nQ_n$	90
ICE	$S_{\text{Turing}} + \frac{Q_1}{\hat{C}_{\text{rare}}} \hat{\gamma}_{\text{rare}}^2$	$\hat{\gamma}_{\text{rare}}^2 = \max\{\gamma - 1, 0\}$ , where $\gamma = \frac{S_{\text{rare}}^* C_{\text{rare}}}{\hat{C}_{\text{rare}} (C_{\text{rare}} - 1)} \frac{\sum_{n=1}^{10} n(n-1)Q_n}{(\sum_{n=1}^{10} nQ_n)(\sum_{n=1}^{10} nQ_n - 1)}$	95
		$C_{\text{rare}} =$ number of samples containing at least one rare species	

40 species abundance at both local or larger scales. Indeed the  
 45 problem of relating occupancy data with information on  
 50 species abundance is a relevant issue in theoretical ecology  
 (He and Gaston 2000, Royle and Nichols 2003, Elith et al.  
 2006). In particular, given the information on the presence  
 or absence of a species in different scattered plots, one would  
 like to infer its population size or, more generally, the RSA  
 distribution of the forest.

55 In this paper, we present a general analytical framework  
 to extrapolate species richness and other relevant biodiversity  
 patterns (e.g. RSA, SAC) at the whole forest scale from local  
 information on species presence/absence. Our framework  
 exploits the form-invariance property of the negative binomial  
 (NB) distribution. Such a distribution emerges as the  
 long time behavior distribution of a birth and death stochastic  
 dynamics, accounting for effective immigration and/or intra-  
 specific interactions (Volkov et al. 2007, Azaele et al. 2016,  
 Tovo et al. 2017). Crucially, the functional form of a negative  
 binomial does not change when sampling different fractions

100 of areas. This property allows for an analytical expression  
 for how parameters of the distribution change across scales.  
 Form-invariance under different sampling efforts is at the  
 core of our approach, however our method can be applied  
 any time the dependence of the distribution on the size of the  
 105 sampled area can be calculated exactly.

We will find an analytical relation between the NB RSA  
 at a given spatial scale and the SAC. Thanks to this function,  
 starting from the empirical SAC constructed at the sample  
 scale from the local presence to absence data (Eq. 13), we will  
 be able to:

1. infer species richness at larger scales, thus the SAC up to  
 the whole forest scale;
2. obtain information on species abundances in order to  
 115 construct the RSA at both local and global scales;
3. introduce and infer the relative species occupancy (RSO),  
 i.e. the distribution of the occurrences (number of occu-  
 pied cells) across species, at both local and global scales.  
 This biodiversity pattern is a prediction of our modelling

60

framework, can be measured empirically and may be of ecological relevance as it proxies the distribution of species ranges (the area where a particular species can be found) in the ecosystem.

We tested our framework on in-silico generated forests and on the two well-studied tropical forests of Barro Colorado Island and Pasoh. We finally compared the global estimates with the abundance-based method proposed in Tovo et al. (2017).

Before illustrating the details of our approach, we want to highlight differences and similarities between the present work and Tovo et al. (2017). Both papers are based on the form-invariance property of the negative binomial distribution but, instead of using population estimates at local scales (Tovo et al. 2017), here we require only the knowledge of species' occurrences at multiple local scales. In other words, the loss of information at one local scale (i.e. for each sample we know if a species is present, but not the number of its individuals) is balanced by the presence-absence information on multiple local scales. Such a generalization of Tovo et al. (2017) is useful when empirical datasets provide information only on the presence/absence of species. We will show that this will be enough to infer population's distribution as well.

## Material and methods

### Theoretical framework

We denote as  $P(n|1)$  the relative species abundance, – i.e. the probability that a species has exactly  $n$  individuals – at the whole forest scale (here 1 refers to the whole forest). Note that  $P(n|1)$  should be defined only for  $n \geq 1$ , because  $S$  is the total number of species actually present in the forest, assuming that the area was exhaustively surveyed with no missing species.

Here the RSA at the scale  $p=1$ , is postulated to be proportional to a negative binomial distribution (NB) (He and Gaston 2003, He and Hubbell 2003, Tovo et al. 2017),  $\mathcal{P}(n|r, \xi)$  with parameters  $r > 0$  and  $0 \leq \xi < 1$ :

$$P(n|1) = c(r, \xi) \mathcal{P}(n|r, \xi), \quad \text{for } n \geq 1 \quad (1)$$

with

$$\mathcal{P}(n|r, \xi) = \binom{n+r-1}{n} \xi^n (1-\xi)^r, \quad c(r, \xi) = \frac{1}{1-(1-\xi)^r}$$

where  $c(r, \xi)$  is the normalisation constant. Notice that since  $n \geq 1$ , the sum  $\sum_{n \geq 1} \mathcal{P}(n|r, \xi) < 1$  and that is why we need a normalizing factor, taking into account only species with non-zero abundance, which is different from the usual NB normalization. It may be worth to mention here that classically, for a NB distribution, one has  $r \in \mathbb{N}$  whereas in our framework  $r \in \mathbb{R}^+$ . Such a distribution can be derived

as the steady-state RSA of a simple birth and death stochastic dynamics (He and Gaston 2003, He and Hubbell 2003, Tovo et al. 2017), where  $r$ , known as the clustering coefficient, models the effects due to immigration events and/or intraspecific interactions, and  $\xi$  is the ratio between the birth and death rate of a species.

Let us now consider a sub-sample of area  $a$  of the whole forest and define  $p = a/A$  the sample scale. Assuming that the local RSA is not affected by spatial correlations and/or strong environmental gradients, the conditional probability that a species has  $k$  individuals in the smaller area  $a = pA$ , given that it has total abundance  $n$  in the whole forest of area  $A$  is given by the binomial distribution

$$P_{\text{binom}}(k|n, p) = \binom{n}{k} p^k (1-p)^{n-k}, \quad k = 0, \dots, n$$

It is worth highlighting that this is where we use the 'well-mixed' (or mean-field) hypothesis. This assumption can be tested in the data by looking at the beta-diversity and RSA patterns. If correlation lengths are of the same scale as the system linear size, and the RSA of sub-samples displays the same functional shape, then we can assume that no strong spatial constraints affect the abundance species distribution.

With this information in hand, it can be proved (Tovo et al. 2017, Supplementary material Appendix 1) that, under the hypothesis that the RSA has a negative binomial form, the RSA at scale  $p$ ,  $P(k|p)$ , is again proportional to a negative binomial, for  $k \geq 1$ , with rescaled parameter  $\xi_p$  and the same  $r$ :

$$P(k|p) = \begin{cases} c(r, \xi) \times \mathcal{P}(k|r, \xi_p) & k \geq 1 \\ 1 - c(r, \xi) / c(r, \xi_p) & k = 0 \end{cases} \quad (2)$$

with

$$\xi_p = \frac{p\xi}{1-\xi(1-p)} \quad (3)$$

A RSA with the property of having the same functional form at different scales is said to be form-invariant.

The form-invariant property allows for simple formula describing how birth and death ratios at two different spatial scales are related. Indeed, given the parameters  $r$  and  $\xi_{p^*}$  of the RSA at the sampling scale  $p^*$ , we can get the value of  $\xi$  by inverting Eq. 3:

$$\xi = \frac{\xi_{p^*}}{p^* + \xi_{p^*}(1-p^*)} \quad (4)$$

Using (3) to eliminate  $\xi$  from the last equation, one gets the following relation for the parameter  $\xi$  at the two scales  $p$  and  $p^*$

$$\xi_p = \frac{p\xi_p}{p^* + \xi_p \cdot (p - p^*)} \quad (5)$$

Let now determine the relation between the total number of species at the whole scale  $p=1$ ,  $S$ , and the total number of species surveyed at a local scale  $p$ ,  $S_p$ . For the sampling scale  $p^*$ , in the following, we will use the notation  $S^* \equiv S_{p^*}$ . Note that, denoting with  $S^*(k)$  the number of species having  $k$  individuals at the scale  $p^*$ , one can estimate  $P(k=0|p^*)$  and  $P(k|p^*)$  as follows

$$P(k=0|p^*) \approx (S - S^*) / S \quad (6)$$

$$P(k|p^*) \approx S^*(k) / S \quad (7)$$

Thus, the total number of species in the whole forest, in terms of the data on the surveyed sub-plot is given by

$$S \stackrel{\text{eq.(6)}}{\approx} \frac{S^*}{1 - P(k=0|p^*)} \quad (8)$$

$$\stackrel{\text{eq.(2)}}{=} S^* \frac{1 - (1 - \xi)^r}{1 - (1 - \xi_p)^r}$$

where  $\xi$  is given by Eq. 4.

In general, given two scales  $p$  and  $p^*$ , one has the following relation between the number of species at the scale  $p$ , the one at  $p^*$  and the RSA parameters ( $r, \xi_p$ ) at the scale  $p^*$ :

$$S_p \approx S^* \frac{1 - (1 - U(p|p^*, \xi_p))^r}{1 - (1 - \xi_p)^r} \quad (9)$$


where

$$U(p|p^*, \xi_p) \stackrel{\text{def}}{=} \frac{p\xi_p}{p^* + \xi_p \cdot (p - p^*)} \quad (10)$$

The proposed method is, under the 'well mixed' hypothesis, general and not limited to tropical forests.

In the sequel we illustrate how, within the theoretical framework developed so far, it is possible to use presence-absence information on various samples to infer  $r, \xi$  and then  $S$ . To start with, let us suppose we surveyed  $M^*$  cells of the same area  $a$ . This assumption is not essential to our approach to species estimation at the global scale. It simplifies computations and its implementation but can be removed. Suppose we have presence-absence information on each of  $M^*$  cells. This implies we know  $S_{p_k}$  (i.e. the number of species at scale  $p_k$ ) for  $p_k = ka/A, k = 1, \dots, M^*$ . From Eq. 9 we obtain

#### DATA AT THE LOCAL SCALE $p^*$



#### PREDICTION AT THE GLOBAL SCALE $p=1$




Figure 1. Schematic presentation of our theoretical upscaling framework. It consists of three steps. (A) We start from a dataset in the form of a binary matrix giving information on the presence or absence of  $S^*$  species within each of the  $M^*$  surveyed plots. (B) We perform the best fit of the empirically SAC computed via Eq. 13. (C) Using the best-fit parameters obtained in (B) and using our upscaling Eq. 8, 9 and 12, we predict the species richness  $S$  of the whole forest and three important macro-ecological patterns: the SAC, the RSA and the RSO.

$$S_{p_k} \approx S^* \frac{1 - \left(1 - U(p_k | p^*, \xi_{p^*})\right)^r}{1 - (1 - \xi_{p^*})^r} \quad (10)$$

$$= S^* \frac{1 - \left(1 - U(\tilde{p}_k | 1, \xi_{p^*})\right)^r}{1 - (1 - \xi_{p^*})^r}$$

where  $\tilde{p}_k = p_k / p^* = k / M^*$  is the fraction of sub-sampled cells. In the last equality of Eq. 10 we use that

$$U(p | p^*, \xi_{p^*}) = U\left(\frac{p}{p^*} | 1, \xi_{p^*}\right) \text{ which is obtained from Eq. 9}$$

factorizing  $p^*$ .

The latter equation states that the function of  $p$  on the righthand side of Eq. 9 takes the value  $S_{p_k}$  at  $p_k$  for  $k=1, \dots, M^*$ . For  $M^* \gg 1$ , these information allow for a robust estimate of the two unknown parameters  $\xi_{p^*}$  and  $r$ . Therefore from the empirical values of  $S_{p_k}$  one can get the parameters  $r$  and  $\xi_{p^*}$  shaping the RSA at the sample scale  $p^*$ . From these, one can estimate the  $\xi$  parameter by using Eq. 4 to predict both the number of species at the global scale  $S$  via Eq. 8, the RSA through Eq. 1 and the SAC by using Eq. 9.

Another important pattern which we can predict with our framework is the relative species occurrence (RSO) distribution,  $Q(v|M, 1)$ , which gives the probability that a species occupies  $v$  cells at the global scale, given that the forest can be tiled in  $M$  equal-sized cells of area  $a$ . The latter assumption is essential to our derivation of RSO formulae (Eq. 11, 12). Also notice the difference between  $M$  and  $M^*$ : in our notation  $M$  is the number of cells at the global scale whereas  $M^*$  refers to the fraction  $p^*$ .

RSO pattern is of ecological relevance as it gives information on the fraction of species that occupy the same amount of area of the ecosystem. For example, if the RSO distribution displayed a small variance unimodal shape, then it means that most of species have similar species ranges. On the other hand if  $Q(v|M, 1)$  follows a power law behaviour it indicates a strong heterogeneities in the species ranges.

In order to find an expression for it, we firstly need the probability,  $Q_{\text{occ}}(v|n, M, 1)$ , that a species occupies  $v$  over  $M$  cells at the global scale, given that it has abundance  $n$ . Under the hypothesis of absence of spatial correlation, this is given

by an hyper-geometric distribution. Indeed there are  $\binom{M}{v}$  possibilities to choose the  $v$  filled cells,  $\binom{n-1}{v-1}$  possibilities to distribute  $n$  species among  $v$  cells so that no cell is empty, and  $\binom{n+M-1}{M-1}$  ways to distribute  $n$  species in  $M$  cells allowing empty bins. See for example W. Feller, Introduction to probability theory and its applications, Chapter 2. Thus we compute

$$Q_{\text{occ}}(v|n, M, 1) = \frac{\binom{M}{v} \binom{n-1}{v-1}}{\binom{n+M-1}{M-1}} \quad (11)$$

The RSO distribution  $Q(v|M, 1)$  can thus be obtained by marginalizing with respect to the abundance  $n$ :

$$Q(v|M, 1) = \sum_{n=v}^{\infty} Q_{\text{occ}}(v|n, M, 1) P(n|1) \quad (12)$$

where  $P(n|1)$  is the global RSA given by Eq. 1. This series cannot be calculated analytically for arbitrary values of the parameters; nevertheless, it has some regimes which are physically important and can be investigated in more detail. For instance, when  $\xi \approx 1$ ,  $P(n|1)$  can be approximated by a gamma distribution and, for  $0 < r < 1$  and  $v, M \gg 1$  such that  $v/M \ll 1$ , one can show that  $Q(v|M, 1) \propto v^{r-1}$ . Since for most forest plots  $r \ll 1$ , when  $M$  and  $v$  are sufficiently large we expect  $Q(v|M, 1) \propto cv^{-1}$ , where  $c$  depends on  $M$ ,  $r$  and  $\xi$ . This prediction is supported by the empirical data we have studied as shown in Fig. 3.

## Implementation of the framework

Our analytical framework has been implemented into a ready-to-use R-code (see Data availability section) and consists of the following steps (Fig. 1).

- First, given a set of scattered samples, list the species in it.

In formulae, sample  $C = \{c_1, \dots, c_{M^*}\}$ ,  $M^* \geq 2$ , cells covering a fraction  $p^*$  of the whole forest in which  $S^*$  species are observed. To each cell  $c_i$ , associate a vector  $\Omega(c_i) = \{\omega_1^i, \dots, \omega_{S^*}^i\}$ , with  $\omega_s^i \in \{0, 1\}$ ,  $s \in \{1, \dots, S^*\}$ ,  $i \in \{1, \dots, M^*\}$ . The entry  $\omega_s^i$  of vector  $\Omega(c_i)$  gives information on the presence/absence of the species  $s$  in the cell  $c_i$  – i.e.  $\omega_s^i = 1$  if species  $s$  is present in cell  $c_i$ ,  $\omega_s^i = 0$  otherwise.

- Compute the empirical species–accumulation curve as follows. From now on, let us suppose that all the  $M^*$  cells are of equal size  $a$ . This assumption does not affect the general framework but it simplifies the computation of the SAC. Call  $A$  the area of the whole forest, so that  $p^* = M^* a / A$ . At each sub-sampling scale  $p_k = k a / A$ , with  $k \in \{1, \dots, M^*\}$ , compute the average number of observed species as

$$S_{\text{emp}}(p_k) = \frac{1}{\binom{M^*}{k}} \sum_{I \subseteq \{1, \dots, M^*\}} \sum_{|I|=k} \sum_{s=1}^{S^*} 1\left(\sum_{i \in I} \omega_s^i \geq 1\right) \quad (13)$$

where  $1(\mathcal{E})$  is the indicator function, which equals one when the random event  $\mathcal{E}$  happens and it is zero otherwise.

In words: for every scale  $p_k$ , one should compute the empirical average of the number of the species observed in all subsets of  $k$  cells. Since computing all subsets of  $k$  cells among  $M^*$  is numerically expensive for large  $M^*$ , in the analyses we computed the average among 100 randomly chosen subsets. Note that computing the species accumulation curve through the empirical average of the number of species in  $k$  random selected cells, we are neglecting any spatial information. Let us stress once again that null or small spatial correlation is required for a rigorous derivation of our estimates.

- Fit the empirical species accumulation curve with the theoretical equation

$$S(p_k) = S^* \frac{1 - \left(1 - U(\tilde{p}_k | 1, \xi_{p^*})\right)^r}{1 - \left(1 - \xi_{p^*}\right)^r} \quad (14)$$

and obtain the parameters  $(r, \xi_{p^*})$  which best describe the empirical curve  $S_{\text{emp}}(p_k)$ . These are the parameters of the NB relative species abundance distribution at the sample scale  $p^*$ . This protocol allows us to capture some spatial effects in the effective parameters.

- As showed in Tovo et al. (2017), under the hypotheses of absence of strong spatial correlations due to both

inter-specific or intra-specific interactions, strong environmental gradients and abundances distributed according to a negative binomial at the whole forest scale, the RSA distributions at different scales have the same functional form of the RSA at the scale  $p^*$ , and only the values of the parameter  $\xi$  changes as a function of the scale. Thus we obtain an analytical form of the upscaled RSA at any scale  $p$  given we know it at scale  $p^*$  in term of the equation  $\xi(p|\xi_{p^*}) = U(p|p^*, \xi_{p^*})$ , relating  $\xi_p = \xi(p)$  to  $p$ ,  $p^*$  and  $\xi_{p^*} = \xi(p^*)$ . Therefore, using the RSA parameters at scale  $p^*$  and the upscaling equations (see below), we can predict the total number of species,  $S$ , at the whole forest scale,  $p = 1$ .

- The key feature of the method is the possibility, given only presence/absence data, to connect and infer different biodiversity patterns at the global scale. Indeed, we can predict, in addition to the SAC, the RSO, the cell occupancy distribution and the RSA, the abundance proportions of the  $S$  species present at  $p = 1$ .

### Data availability

All data are publicly available. The Pasoh and Barro Colorado Island datasets are provided by the Center of Tropical Research Science of the Smithsonian Tropical Research Institute (<<https://stri.si.edu/>>). R codes are available at




Figure 2. Test from different scales for Pasoh and BCI. For each forest we sub-sample a fraction  $p$  of  $p^*$  of the available spatial cells and apply different popular upscaling methods based on presence/absence data (Table 2) and our method to predict the true number of species,  $S^*$  (dashed line), observed in our data. While our (P/A NB) and Chao<sub>war</sub> methods do converge at  $S_{p^*}$  as  $p$  goes to  $p^*$ , all the others have a monotonically increasing behaviour due to the independence, in their predictions, of the scale  $p^*$ . We can see that for both rainforests, our method outperforms all the others. Bottom panels show the relative percentage error  $(S_{\text{pred}} - S^*)/S^* \times 100$  obtained with our framework between the predicted number of species  $S_{\text{pred}}$  and  $S^*$ . We find that the method underestimated the true number of species of at most 5%. The larger the sample area, the smaller the relative error.




Figure 3. Test on ecological macro-patterns for Pasoh and BCI. For each forest we sub-sample a fraction  $p=0.1$  of the available spatial cells and apply our framework to predict three important ecological pattern at the largest scale at which we have information,  $p^*$ . In the first row we see the prediction for the SAC curve, which describes how the number of observed species increases with the sampled area, from  $p=0.1$  to 1, corresponding to  $p^*$  in these tests. In the second row we plot the cumulative empirical RSA, the distribution of abundances across species against the framework prediction in logarithmic scale. Finally, in the third and fourth rows we test the ability of the model to capture the empirical RSO, i.e. the distribution of the occurrences (number of occupied cells) across species in logarithmic scale (third row panel shows the cumulative distribution). In figures, predicted patterns are in the form of confidence intervals obtained from the SAC fitting errors on the  $r$  and  $\xi_p$  parameters. For both forests, all the three patterns result to be well described by our framework.

<<https://github.com/annatovo/Inferring-macro-ecological-patterns-from-local-species-occurrences>>.

## Results

### Tests on in-silico databases

We test our presence/absence upscaling method on four computer generated forests without and with spatial correlations. Indeed, we expect that in the first case our framework will give more accurate estimates, and we wish to test how the introduction of correlations affect the reliability of our results.

0 As RSA we choose a negative binomial (NB forest) of  
 parameters  $r=0.8$  and  $\xi=0.999$  and a log-normal (LN forest)  
 with parameters  $\mu=5$  and  $\sigma=1$ . Once generated the abun-  
 5 dance of every species ( $S=4974$  for the NB forest and 5000  
 for the LN forest), we distribute the individuals within the  
 forest area, here set equal to a square of  $4900 \times 4900$  units,  
 according to two different processes: at random or according  
 to a modified Thomas process (Tovo et al. 2016, 2017, Tovo  
 and Favretti 2017) with a clustering radius of 15 units.

10 We then divide each forest generated as described above  
 into  $M=98 \times 98$  units cells and compute the  $M \times S$  presence/  
 absence matrix, thus forgetting the information about the  
 species distribution. Finally, we sub-sample the 5% of the  
 15 cells (corresponding to a fraction  $p=0.05$  of the total forest  
 area) and apply our method to infer the total number of spe-  
 cies in each of the four in-silico forests.

We also compared our results to those obtained by  
 accounting for the data on species populations with an  
 abundance-based upscaling framework developed and tested  
 in Tovo et al. (2017). In the case of the NB forest, the two  
 20 methods performed very well for both the random and the  
 clumped distribution (i.e. individuals distributed on the  
 space according to a Thomas point process) with an average  
 prediction error below 1% in absolute value (Table 1). In  
 the Thomas distributed forests, the error increased, although  
 25 remaining around 3% for the presence/absence method and  
 around 7.5% for the abundance-based one (using maxi-  
 mum likelihood methods. The latter percentage error can be  
 improved using calibrated statistical method for the single  
 fit). Thus, with respect to the degree of individuals' cluster-  
 30 ing, the new framework seems to give more robust estimates  
 than the second one. This is due to two main reasons: 1) for  
 the presence-absence case, we fit the empirical SAC, which  
 has a very smooth functional shape, and it is easy to describe  
 through our analytical SAC. On the other hand, the RSA  
 displays a more complex and variable shape and thus fitting it  
 with the NB is a more delicate task (indeed we find sensible  
 35 differences on the accuracy using different statistical methods  
 for the fit); 2) binary data on which the empirical SAC is  
 based are less sensitive to sampling fluctuations.

### 45 Tests on real databases

We finally test our method on sub-samples taken from two  
 empirical forest data for which we have informations on both  
 species occurrence and abundances. In particular we extract  
 abundances of tree species observed in 50 ha of rainforests  
 50 from Pasoh (Malaysia) and Barro Colorado Island (Panama)  
 together with the spatial locations of each of their individual.

55 Firstly, we divide both forest data into a grid consisting  
 of  $M=800$  equal-sized cells of area  $625 \text{ m}^2$  and we derive the  
 $M \times S^*$  presence/absence matrix for the  $S^*$  observed species  
 ( $S^*=927$  for Pasoh forest and 301 for BCI). We then sub-  
 sample species occurrence for different fractions  $0 < p < p^*$  of  
 the cells and apply our framework to infer the number of  
 species and other biodiversity patterns (RSA, RSO and SAC)

61 at the corresponding largest empirically-observable scale  $p^*$ ,  
 for which we know the ground truth.

65 We compared our results on species richness obtained only  
 from presence to absence data with the most popular non-  
 parametric indicators proposed in the literature (Chao 2005,  
 Chao and Chiu 2016), which are summarized in Table 2.  
 We found that our method outperforms all the others for  
 both BCI and Pasoh forests. We also remark that all these  
 70 methods have the further limitation that they can only infer  
 the total species richness, without allowing for an estimate of  
 the abundances' and occurrences' distributions, i.e the shape  
 of the RSA and the RSO.

75 Indeed, as shown in Fig. 3, from the local presence to  
 absence data, we can reconstruct, among the SAC, the RSA  
 at the whole tropical forest scale, thus relating species occur-  
 rence data with information on the abundances. In particular  
 we can see that the inferred RSA are statistically comparable  
 with the empirical ones obtained by using all the information  
 80 on species' abundances which we deleted before applying our  
 method.

85 Another biodiversity pattern that we can infer from our  
 framework is the RSO. As shown in Fig. 3, we find that,  
 as for the RSA, this pattern seems to have a universal form  
 which can be well described and correctly inferred through  
 our neutral approach. Also, our finding suggests that, when  
 spatial effects are negligible, the RSO distribution has a wide  
 range of values in which it is well approximated by a universal  
 90 power law, regardless of the details of the populations' dynam-  
 ics. One may assume that this latter is driven by a simple sto-  
 chastic process with constant per capita birth and death rates.  
 Such a slow decay of  $Q(v|M, 1)$  indicates that species in real  
 systems exhibit huge variations in their occurrences, which  
 95 may be weakly correlated to species' habitat preferences or  
 environmental heterogeneities. We should expect strong  
 asymmetries among their occurrences: for instance, if we tile  
 up a landscape into  $M=1000$  elementary cells, then about  
 a third of all species should live in less than 1% of them;  
 100 whereas about 1.5% of the species should be found in more  
 than 90% of the total cells (Fig. 3).

105 We highlight that the SAC (green line), the cumulative  
 RSA (red line) and cumulative RSO (blue line) predicted pat-  
 terns in Fig. 3 have not been obtained through the fit of some  
 parameters, but they have been analytically predicted through  
 our upscaling Eq. 1, 9 and 12. The only fitting occurs at the  
 scale  $p=0.1p^*$  using the empirical SAC to parametrize Eq.  
 13. In other words, by fitting species occurrence data at the  
 110 sample scale, our framework allows to estimate: 1) the RSA at  
 the sample scale; 2) the SAC, the RSA and the RSO at larger  
 scales. We provide an open source R code that performs the  
 above estimates giving as input only the presence-absence  
 matrix data.

115 After testing our model on controlled computer generated  
 data and real forest sub-samples, we apply our framework  
 to predict the species richness of the two tropical forests.  
 Moreover, we compare our results to those obtained with the

61  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
121




Figure 4. SAC predicted for Pasoh and BCI using abundance method versus presence/absence method. Using all the available data for both tropical forests, we compare the prediction for the SAC curve obtained by the abundance method (Tovo et al. 2017) with the results obtained with the presence/absence framework presented here. At the whole forests' scale  $p=1$ , the two predictions are  $3\sigma$  compatible ( $S_{\text{Pasoh}}^{\text{abund}} = 1193 \pm 36$ ,  $S_{\text{Pasoh}}^{\text{p/a}} = 1260 \pm 22$ ,  $S_{\text{BCI}}^{\text{abund}} = 366 \pm 15$ ,  $S_{\text{BCI}}^{\text{p/a}} = 359 \pm 2$ ).

upscaling framework based on RSA pattern previously developed and tested in Tovo et al. (2017) by our group.

We therefore predict, through the presence/absence method, the species richness at the whole forest scale ( $p=1$ ) for BCI and Pasoh tropical forests. Figure 4 shows the prediction of the overall (and unknown) SAC for a scale ranging from 50 to 14 000 hectares for the Pasoh ( $p^* \approx 0.0036$ ) and to 1560 for the BCI ( $p^* \approx 0.032$ ). The blue curves represent the prediction obtained only using presence-absence data whereas red curves are the SAC inferred by exploiting also the information about species' population through the abundance-based method (Tovo et al. 2017).

We find that the two methods give comparable results for both the databases, a confirm of the robustness of the theoretical framework.

## Discussion

In this work we proposed and tested a novel rigorous statistical framework to upscale ecological biodiversity patterns from local information on species occurrence data. Different upscaling approaches have been proposed in ecological literature (Good and Toulmin 1956, Bunge and Fitzpatrick 1993, Wang and Lindsay 2005, Harte et al. 2009, Azaele et al. 2015, Slik et al. 2015, Chao and Chiu 2016, Orlitsky et al. 2016, Ter Steege et al. 2017, Ter Steege et al. 2017). However, to the best of our knowledge, they have not been generalized to the case of binary data. The present paper provides a generalization of the method recently proposed in Tovo et al. (2017) to presence-absence information. In Tovo et al. (2017) species abundance distributions at one given scale was required to make predictions at global scale, whereas the present approach allows to extract abundance distributions at any scale from species occurrence data in multiple small scale samples. The underlying hypotheses that we need in order to perform these estimates is that the RSA at a given scale is a negative binomial distribution, a RSA that arises naturally as the steady-state species abundance distributions for ecosystems undergoing simple birth and death dynamics (Volkov et al. 2005, Azaele et al. 2016). The negative binomial is a simple

and versatile distribution that depending on its parameters can display an interior mode or log-series like behaviour, i.e. it can accommodate different RSA shapes. Therefore we can use the same RSA function to reproduce different ecosystems' RSA, as those typically observed in real ecosystems (Chave 2004, Magurran 2005, Chave et al. 2006, Chisholm 2007, Volkov et al. 2007, Magurran 2013, Matthews and Whittaker 2014, Kessler et al. 2015, Azaele et al. 2016). Even more generally, by using mixtures of negative binomials – a case for which our framework still works – we can fit more complex RSA shapes (Tovo et al. 2017).

Furthermore, we introduce a new descriptor/measure of biodiversity within an ecological community, the RSO, which describes the distribution of species occurrences in scattered plots. The RSO distribution displays a fat tail, indicating that many species typically occupies only few scattered plots, while only very few species are pervasive and are found in most of the plot. Our prediction is that this property is not particular for the dataset here considered, rather it is another emergent patterns (Suweis et al. 2013, Azaele et al. 2016) pervasive in highly biodiverse ecosystems. Our framework gives directly all parameters of the RSO by solely fitting the SAC curve, through which one can obtain the  $r$  and  $\xi$  parameters, which well describe both the RSA and the RSO distributions at all spatial scales of interest.

Expanding the ability to upscale species richness and obtain abundance distributions from presence to absence data is of fundamental importance in many contexts, where abundance information are not available or trustable. This is particularly true for microbial or marine (e.g. plankton) ecological data obtained from metagenomics (Menzel et al. 2016) and 16S ribosomal gene sequences (Soergel et al. 2012). The use of sequence-based taxonomic classification of environmental microbes has exploded in recent years (Soergel et al. 2012, De Vargas et al. 2015, Menzel et al. 2016, Thompson et al. 2017) and these approaches are becoming a standard method for characterizing the biodiversity of both prokaryotes and eukaryotes (De Vargas et al. 2015). Thanks to advance in high throughput sequencing we begin to be able quantifying the vast number of microbes in our environments, expanding our knowledge on microbial diversity (Thompson et al.

0 2017). However, large fractions of the sequence reads remain  
 unclassified (Menzel et al. 2016) and also species abundance  
 estimated have a very high uncertainty (Thompson et al.  
 2017). Thus, being able to estimated species richness and  
 5 abundance distributions from species occurrence data may  
 lead to a big step-forward in the taxonomic classification of  
 microbial ecosystems.

To summarize, this flexible analytical method provides,  
 from local presence/absence information, robust estimates  
 10 of species richness and important macro-ecological patterns  
 of biodiversity (SAC, RSA, RSO), as tested in both in-silico  
 generated and two rainforests. The method may be applied to  
 any database in the form of a binary matrix, where presence/  
 15 absence features (tree species in our case) are detected across  
 different samples.

*Acknowledgments* – SS acknowledges STARS grant 2018 from  
 University of Padova. AM was supported by Excellence Project  
 20 2017 of the Cariparo Foundation. This preprint has been reviewed  
 and recommended by Peer Community in Ecology (<https://dx.doi.org/10.24072/pci.ecology.100009>).

*Competing interests* – The authors of this preprint declare that they  
 have no financial conflict of interest with the content of this article.  
 25 Samir Suweis is recommender at PCI Ecology.

*Author contributions* – AT and MF contributed equally to this paper.  
 AT and MF carried out the numerical simulations, the data analysis  
 and performed the figures. All the authors carried out analytical  
 30 calculations. All the authors contributed to other aspects of the  
 paper and the writing of the manuscript.

## References

35 Alonso, D. et al. 2008. The implicit assumption of symmetry and  
 the species abundance distribution. – *Ecol. Lett.* 11: 93–105.  
 Azale, S. et al. 2015. Towards a unified descriptive theory for spatial  
 ecology: predicting biodiversity patterns across spatial scales.  
 40 – *Methods Ecol. Evol.* 6: 324–332.  
 Azale, S. et al. 2016. Statistical mechanics of ecological systems:  
 neutral theory and beyond. – *Rev. Mod. Phys.* 88: 035003.  
 Bertuzzo, E. et al. 2016. Geomorphic controls on elevational gra-  
 dients of species richness. – *Proc. Natl Acad. Sci. USA* 113:  
 1737–1742.  
 45 Brose, U. et al. 2003. Estimating species richness: sensitivity to  
 sample coverage and insensitivity to spatial patterns. – *Ecology*  
 84: 2364–2377.  
 Bunge, J. and Fitzpatrick, M. 1993. Estimating the number of  
 species: a review. – *J. Am. Stat. Assoc.* 88: 364–373.  
 50 Bunge, J. et al. 2012. Estimating population diversity with catchall.  
 – *Bioinformatics* 28: 1045–1047.  
 Carrara, F. et al. 2012. Dendritic connectivity controls biodiversity  
 patterns in experimental metacommunities. – *Proc. Natl Acad. Sci. USA* 109:  
 5761–5766.  
 55 Chao, A. 2005. Species estimation and applications. – *Encyclopedia*  
*of Statistical Sciences*.  
 Chao, A. and Bunge, J. 2002. Estimating the number of species in  
 a stochastic abundance model. – *Biometrics* 58: 531–539.  
 Chao, A. and Chiu, C.-H. 2016. Species richness: estimation and  
 60 comparison. – *Wiley StatsRef: Statistics Reference Online*.

Chao, A. et al. 2009. Sufficient sampling for asymptotic minimum  
 species richness estimators. – *Ecology* 90: 1125–1133.  
 Chave, J. 2004. Neutral theory and community ecology. – *Ecol.*  
*Lett.* 7: 241–253.  
 Chave, J. et al. 2006. Comparing models of species abundance. – 65  
*Nature* 441: E1.  
 Chisholm, R. A. 2007. Sampling species abundance distributions:  
 resolving the veil-line debate. – *J. Theor. Biol.* 247: 600–607.  
 Colwell, R. K. and Coddington, J. A. 1994. Estimating terrestrial  
 biodiversity through extrapolation. – *Phil. Trans. Biol. Sci.* 70<sup>2</sup>  
 101–118.  
 Colwell, R. K. et al. 2004. Interpolating, extrapolating and compar-  
 ing incidence-based species accumulation curves. – *Ecology* 85:  
 2717–2727.  
 Corbet, A. S. 1941. The distribution of butterflies in the Malay  
 Peninsula (lepid.). – *Physiol. Entomol.* 16: 101–116.  
 75 De Vargas, C. et al. 2015. Eukaryotic plankton diversity in the  
 sunlit ocean. – *Science* 348: 1261605.  
 Elith, J. et al. 2006. Novel methods improve prediction of species  
 distributions from occurrence data. – *Ecography* 29: 129–151.  
 80 Fisher, R. A. et al. 1943. The relation between the number of spe-  
 cies and the number of individuals in a random sample of an  
 animal population. – *J. Anim. Ecol.* 42–58.  
 Good, I. J. and Toulmin, G. H. 1956. The number of new species,  
 and the increase in population coverage, when a sample is  
 increased. – *Biometrika* 43: 45–63.  
 85 Harte, J. et al. 2009. Biodiversity scales from plots to biomes with  
 a universal species–area curve. – *Ecol. Lett.* 12: 789–797.  
 He, F. and Gaston, K. J. 2000. Estimating species abundance from  
 occurrence. – *Am. Nat.* 156: 553–559.  
 90 He, F. and Gaston, K. J. 2003. Occupancy, spatial variance and the  
 abundance of species. – *Am. Nat.* 162: 366–375.  
 He, F. and Hubbell, S. P. 2003. Percolation theory for the distribu-  
 tion and abundance of species. – *Phys. Rev. Lett.* 91: 198103.  
 Hughes, J. B. et al. 2001. Counting the uncountable: statistical  
 approaches to estimating microbial diversity. – *Appl. Environ.*  
*Microbiol.* 67: 4399–4406.  
 95 Ionita-Laza, I. et al. 2009. Estimating the number of unseen vari-  
 ants in the human genome. – *Proc. Natl Acad. Sci. USA* 106:  
 5008–5013.  
 Kessler, D. et al. 2015. Neutral dynamics with environmental noise:  
 age-size statistics and species lifetimes. – *Phys. Rev. E* 92:  
 022722.  
 100 Kunin, W. E. et al. 2018. Upscaling biodiversity: estimating the  
 species–area relationship from small samples. – *Ecol. Monogr.*  
 88: 170–187.  
 105 Locey, K. J. and Lennon, J. T. 2016. Scaling laws predict global  
 microbial diversity. – *Proc. Natl Acad. Sci. USA* 113:  
 5970–5975.  
 Magurran, A. E. 2005. Species abundance distributions: pattern or  
 process? – *Funct. Ecol.* 19: 177–181.  
 110 Magurran, A. E. 2013. Measuring biological diversity. – *Wiley*.  
 Mlao, C. X. and Colwell, R. K. 2005. Estimation of species rich-  
 ness: mixture models, the role of rare species and inferential  
 challenges. – *Ecology* 86: 1143–1153.  
 115 Mlatthews, T. J. and Whittaker, R. J. 2014. Neutral theory and the  
 species abundance distribution: recent developments and pros-  
 pects for unifying niche and neutral perspectives. – *Ecol. Evol.*  
 4: 2263–2277.  
 Mlenzel, P. et al. 2016. Fast and sensitive taxonomic classification  
 for metagenomics with Kaiju. – *Nat. Commun.* 7: 11257.  
 121

- 0 Orlitsky, A. et al. 2016. Optimal prediction of the number of  
Q4 unseen species. – *Proc. Natl Acad. Sci. USA* 201607774. 61
- Plotkin, J. B. et al. 2000. Species–area curves, spatial aggregation  
and habitat specialization in tropical forests. – *J. Theor. Biol.*  
207: 81–99. 65
- 5 Royle, J. A. and Nichols, J. D. 2003. Estimating abundance from  
repeated presence–absence data or point counts. – *Ecology* 84:  
777–790.
- Shen, T.-J. and He, F. 2008. An incidence-based richness estimator  
for quadrats sampled without replacement. – *Ecology* 89:  
2052–2060. 70
- 10 Slik, J. W. F. et al. 2015. An estimate of the number of tropical tree  
species. – *Proc. Natl Acad. Sci. USA* 112: 7472–7477.
- Soergel, D. A. et al. 2012. Selection of primers for optimal taxo-  
nomic classification of environmental 16s rRNA gene sequences.  
– *ISME J.* 6: 1440. 75
- 15 Suweis, S. et al. 2013. Emergence of structural and dynamical prop-  
erties of ecological mutualistic networks. – *Nature* 500: 449.
- Ter Steege, H. et al. 2013. Hyperdominance in the Amazonian tree  
flora. – *Science* 342: 1243092.
- 20 Ter Steege, H. et al. 2017. Estimating species richness in hyper-  
diverse large tree communities. – *Ecology* 98: 1444–1454.
- Supplementary material (available online as Appendix oik-  
06754 at <[www.oikosjournal.org/appendix/oik-06754](http://www.oikosjournal.org/appendix/oik-06754)>). 85
- 25 Appendix 1.
- 30
- 35
- 40
- 45
- 50
- 55
- 60
- Thompson, L. R. et al. 2017. A communal catalogue reveals earth's  
multiscale microbial diversity. – *Nature* 551: 457–463.
- Tovo, A. and Favretti, M. 2017. The distance decay of similarity in  
tropical rainforests. A spatial point processes analytical formula-  
tion. – *Theor. Popul. Biol.* 120: 78–89. 65
- Tovo, A. et al. 2016. Application of optimal data-based binning  
method to spatial analysis of ecological datasets. – *Spat. Stat.*  
16: 137–151.
- Tovo, A. et al. 2017. Upscaling species richness and abundances in  
tropical forests. – *Sci. Adv.* 3: e1701438. 70
- Ulrich, W. and Ollik, M. 2005. Limits to the estimation of species  
richness: the use of relative abundance distributions. – *Divers.*  
*Distrib.* 11: 265–273.
- Volkov, I. et al. 2005. Density dependence explains tree species  
abundance and diversity in tropical forests. – *Nature* 438:  
658–61. 75
- Volkov, I. et al. 2007. Patterns of relative species abundance in rain-  
forests and coral reefs. – *Nature* 450: 45–49.
- Wang, J.-P. Z. and Lindsay, B. G. 2005. A penalized nonparamet-  
ric maximum likelihood approach to species richness estima-  
tion. – *J. Am. Stat. Assoc.* 100: 942–959. 80

**Author Queries**

*JOB NUMBER: 6754*

JOURNAL: OIK\_OIK

- Q1 Please provide the in-text citation for the Figure 2.
- Q2 Please provide the volume number for the reference 'Colwell and Coddington 1994'.
- Q3 Please provide the volume number for the reference 'Fisher et al. 1943'.
- Q4 Please provide the volume number for the reference 'Orlitsky et al. 2016'.

For Review Only