

This is a repository copy of *Linear response theory for Gogny interaction*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/148490/>

Version: Accepted Version

Article:

davesne, dany, Navarro, Jesus and Pastore, Alessandro orcid.org/0000-0003-3354-6432
(2019) Linear response theory for Gogny interaction. Acta Physica Polonica B. ISSN 0587-4254

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Linear response theory for Gogny interaction

D. DAVESNE^{a,b}, J. NAVARRO^c, A. PASTORE^d

^a Université de Lyon, Université Lyon 1, 43 Bd. du 11 Novembre 1918, F-69622
Villeurbanne cedex, France

^b CNRS-IN2P3, UMR 5822, Institut de Physique Nucléaire de Lyon

^c IFIC (CSIC-Universidad de Valencia), Apartado Postal 22085,
E-46.071-Valencia, Spain

^d Department of Physics, University of York, Heslington, York, YO10 5DD,
United Kingdom

We present the formalism of the linear response theory in symmetric nuclear matter for a finite-range central interaction including zero-range spin-orbit and tensor components.

PACS numbers: 21.30.Fe 21.60.Jz 21.65.-f 21.65.Mn

1. Introduction

Symmetric nuclear matter (SNM) is a spin-saturated system composed of an equal amount of neutrons and protons with no finite-size effect. Although it may appear as an highly ideal system, the study of its excitation modes provides us with several useful guidelines on other systems as atomic nuclei and neutron stars (NS).

In Ref. [1], we have presented in great detail the formalism of Linear Response (LR) theory for a Skyrme functional including both spin-orbit and tensor terms [2]. By studying the response function of the system to an external probe, we have been able to identify the critical densities and momenta at which instabilities occur in the system [3] and relate them to the appearance of finite-size instabilities in nuclei [4, 5]. We have then provided in Refs. [6, 7] a simple method based on LR in SNM to avoid such instabilities directly at the level of the fitting procedure.

The response functions of SNM, or more generally of asymmetric nuclear matter [8], have also direct application to study neutrino opacities in NS [9, 10, 11] which in turn have important effects on NS cooling [12].

In the present article, we present an extension of the formalism of LR theory to the case of finite-range interactions with spin-orbit and tensor

terms. The basic formalism has been discussed in Ref. [13], for the case of a central part of the Gogny interaction [14]. The authors of Ref. [15] have already discussed the LR of a Gogny interaction using continued fraction (CF) approximation [16]. The main inconvenient of such a method is that most of the integrations required to obtain the response functions need to be performed numerically via Monte-Carlo samplings. Moreover, in Ref. [15], the spin-orbit term has been neglected, which is only valid for low-transfer momenta, see discussion in Ref. [3]. The current formalism, being based on partial wave decomposition [17], offers us the opportunity of including all terms of the interaction. Compared to Ref. [15], our formalism requires more detailed analytical derivations of all matrix elements in the different channels, but it is numerically less expensive.

The article is organised as follows: in Sec. 2 we briefly introduce the LR formalism using multipolar expansion, while in Sec. 3, we illustrate our results for the case of a Gogny interaction. We present our conclusions in Sec. 4.

2. Formalism

The response function of the system is obtained by integrating, over the momentum, the RPA propagator [1, 18]. The latter is the solution of the Bethe-Salpeter equation in each spin (S), spin-projection (M) and isospin (I) channel, for brevity called $\alpha \equiv (S, M, I)$ in this paper. It reads

$$G_{\text{RPA}}^{(\alpha)}(q, \omega, \mathbf{k}_1) = G_{\text{HF}}(q, \omega, \mathbf{k}_1) + G_{\text{HF}}(q, \omega, \mathbf{k}_1) \sum_{\alpha'} \int \frac{d^3 \mathbf{k}_2}{(2\pi)^3} V_{ph}^{(\alpha, \alpha')}(q, \mathbf{k}_1, \mathbf{k}_2) G_{\text{RPA}}^{(\alpha')}(q, \omega, \mathbf{k}_2), \quad (1)$$

where q is the transferred momentum and $\mathbf{k}_1, \mathbf{k}_2$ is the momentum of the particle-hole pair and ω the transferred energy. $V_{ph}^{(\alpha, \alpha')}(q, \mathbf{k}_1, \mathbf{k}_2)$ represents the residual particle-hole interaction. In the case of a Skyrme interaction [1, 3, 18, 19], Eq. (1) can be solved analytically using a system of symbolic equations. For the case of a finite-range interaction as Gogny, this is not possible. We have thus adopted the technique presented in Ref. [13] and performed a multipolar expansion as

$$G_{\text{HF}}(q, \omega, \mathbf{k}_1) = \sum_l G_l^{\text{HF}}(q, \omega, k_1) Y_{l0}(\cos \theta_1) \quad (2)$$

$$V_{ph}^{(\alpha, \alpha')}(q, \mathbf{k}_1, \mathbf{k}_2) = \sum_{lm, l'm'} V_{lm; l'm'}^{(\alpha, \alpha')}(k_1, k_2) Y_{lm}(\hat{k}_1) Y_{l'm'}^*(\hat{k}_2) \quad (3)$$

$$G_{\text{RPA}}^{(\alpha)}(q, \omega, \mathbf{k}_1) = \sum_{lm} G_{lm}^{(\alpha)}(q, \omega, k_1) Y_{lm}(\hat{k}_1) \quad (4)$$

where Y_{lm} is the usual spherical harmonic. We then obtain the multipolar expansion of Eq. (1) as

$$G_{lm}^{(\alpha)}(k_1) = \delta_{m,0} G_l^{\text{HF}}(k_1) + \sum_{\alpha'} \sum_{l'm'} \int \frac{dk_2 k_2^2}{(2\pi)^3} M_{lm, l'm'}^{(\alpha, \alpha')}(k_1, k_2) G_{l'm'}^{(\alpha')}(k_2), \quad (5)$$

where, for simplicity, we have dropped the explicit dependence on q and ω . The matrix elements $M_{lm, l'm'}^{(\alpha, \alpha')}$ are defined as

$$M_{lm, l'm'}^{(\alpha, \alpha')}(k_1, k_2) = \sum_{l_1 l_2} \left[\frac{(2l_2 + 1)(2l_1 + 1)}{4\pi(2l + 1)} \right]^{1/2} C_{l_1 0 l_2 0}^{l 0} C_{l_2 m, l_1 0}^{l m} \times G_{l_1}^{\text{HF}}(k_1) V_{lm; l'm'}^{(\alpha, \alpha')}(k_1, k_2) \quad (6)$$

The notation used here simplifies to the one of Ref. [13] in the case of a central interaction for which $l = l'$ and $m = m'$. Eq. (5) actually represents a system of coupled integral equations. To solve them, we discretise the integrals over a uniform mesh over the range $k \in [0, q + k_F]$ as done in Ref. [13] and then invert the matrix of the system. In principle, the size of the system is infinite, since a finite range interaction contains all multipoles. However, as discussed in Ref. [13], the contribution of higher order multipoles becomes smaller and smaller. We have thus introduced a cut-off parameter L_{Max} to limit the number of equations we have to solve. In Sec. (3), we discuss the convergence of our results with L_{Max} .

3. Results

To test the quality of our results, we start by considering the case of a Skyrme force with tensor term, namely T44 [2]. Since the Skyrme interaction is a simple combination of S and P waves [20], the partial wave expansion has a natural truncation. Notice that additional caution should

be taken when considering an explicit tensor contribution : since the tensor term t_e couples S and D waves, we have to take $L_{Max} = 2$. Beyond $L = 2$ the other contributions are zero, due to the particular form of the Skyrme force.

In Fig. 1, we compare the response functions obtained with the method given in Eq. (5) (dashed lines) with those obtained by using the technique described in Ref. [1]. Both calculations are done at saturation density and transferred momentum $q = k_F$. The results stay on top of each other : the small deviations are only due to the discretisation on a finite grid of Eq. (5). This example clearly proves the validity of our method based on multipolar expansion and we will now apply it to the case of finite-range interactions.




Fig. 1. (Colors online) Response functions $-\text{Im}\chi^{(\alpha)}/\pi$ in SNM calculated at saturation density and transferred moment $q = k_F$ for Skyrme T44 interaction. The small differences between solid and dashed lines are the consequence of the method (discretisation) used to solve Bethe-Salpeter equations. See text for details.

In Figs. 2-3, we present the response functions in the different spin-isospin channels for different values of the maximum angular momentum L_{Max} in the case of Gogny D1S interaction [14]. The response functions are calculated at $\rho = 0.16 \text{ fm}^{-3}$ and at two different values of transferred momentum $q = 0.1k_F, q = k_F$. The Gogny D1S interaction is not equipped with a tensor term, thus the spin-orbit term only induces a splitting of the different projections of spin $M = 0, \pm 1$. Since the transferred momentum is still quite small, such a splitting is quite negligible as discussed in Ref. [15]. We clearly observe that $L_{Max} = 2$ is enough to obtain a reasonable description of the response function, confirming the results of Ref. [13].




Fig. 2. (Colors online) Response function $-\text{Im}\chi^{(\alpha)}/\pi$ in SNM calculated at $\rho = 0.16 \text{ fm}^{-3}$ and transferred momentum $q = 0.5k_F$ for Gogny D1S interaction as a function of cut-off in angular momentum expansion L_{Max} .




Fig. 3. (Colors online) Same as Fig.2, but for $q = k_F$.

Finally we compare in Fig. 4 our calculations obtained using Eq. (1) with a cut-off of $L_{Max} = 2$ with the ones of Ref. [15], based on Continued Fraction (CF) approximation [16]. Following Ref. [15], the terms included in the calculations are the central part of the Gogny D1S interaction. The density is $\rho = 0.174 \text{ fm}^{-3}$ and the transferred momentum $q = k_F$. Without tensor or spin-orbit terms, there is no longer a splitting in the different M projections of the total spin S . For such a reason we present only results with $M = 0$. We can see that both methods are in excellent agreement, thus proving the validity of our results.

It is also interesting to compare the $S = 0$ channels from Fig. 4 and Fig. 3




Fig. 4. (Colors online) Response function for the central term of the D1S Gogny interaction. The calculations are done at $\rho = 0.174 \text{ fm}^{-3}$ and $q = k_F$. See text for details.

since we can observe a non-negligible difference of the low-energy part of the isoscalar response function. To understand the role of spin-orbit in this channel, it is useful to observe the expression of the response function given in Ref. [3]: due to the spin-orbit term the interaction terms of the $S = 1$ do contribute in the $S = 0$ channel and the term mixing the two channels is typically proportional to the transferred momentum to the power of 4. As a consequence when the transferred momentum is small we can neglect such a term, but for large transferred momenta this term starts playing an important role.

A very important aspect of the LR formalism is the detection of poles in the response functions. In Fig. 5, we considered the example of the Gogny D1S interaction and showed the evolution of the response function in the channel (0,0,1) for different values of transferred momentum. The calculations have been performed at $\rho = 1.5\rho_0$, ρ_0 being the saturation density of the Gogny D1S interaction because it is known (see Ref. [15]) that at this particular value of density, D1S exhibits a pole at $q_c = 2.53 \text{ fm}^{-1}$. Effectively, we can observe the shape evolution of the response function when q approaches the critical value q_c , showing the stability of our numerical approach and the ability of the formalism to detect instabilities.




Fig. 5. (Colors online) Response functions for the Gogny D1S interaction the channel (0,0,1) and different values of the transferred momentum. See text for details.

4. Conclusions

In this paper, we have presented the formalism of the LR theory for finite-range interactions using multipolar expansion of the Bethe-Salpeter equation. We have tested our formalism against the results of the LR formalism given in Ref. [1] for the case of Skyrme T44 [2]. In particular, thanks to the multipolar expansion, we clearly observe the tensor coupling between S and D waves. As a further benchmark of the formalism, we have compared the result for the case of a Gogny interaction with Ref. [15]. In this case the results are in excellent agreement as well. The advantage of our technique (compared to Ref. [15]) is the explicit inclusion of spin-orbit term in the residual interaction. As explicitly shown, this may lead to non-negligible effects on the response function since it induces an extra coupling between the $S = 0$ and $S = 1$ channels.

Finally, we have explored how the current formalism describes the presence of poles in the response function. The goal is to continue in that direction so that we can include directly a test to detect instabilities in the fitting procedure itself. This can be important in the context of future Gogny-like parametrisation devoted to nuclear astrophysics by instance [21, 22].

Acknowledgments

The work of J.N. has been supported by grant FIS2017-84038-C2-1-P, Mineco (Spain). The work of A.P. is supported by the UK Science and Tech-

nology Facilities Council under Grants No. ST/L005727 and ST/M006433.

REFERENCES

- [1] A. Pastore, D. Davesne, J. Navarro, *Phys. Reports* **563**, 1 (2015).
- [2] T. Lesinski, M. Bender, K. Bennaceur, T. Duguet, J. Meyer, *Phys. Rev. C* **76**, 014312 (2007).
- [3] A. Pastore, D. Davesne, Y. Lallouet, M. Martini, K. Bennaceur, J. Meyer, *ibid.* **85**, 054317 (2012).
- [4] V. Hellemans, A. Pastore, T. Duguet, K. Bennaceur, D. Davesne, J. Meyer, M. Bender, P.-H. Heenen, *ibid.* **88**, 064323 (2013).
- [5] A. Pastore, D. Tarpanov, D. Davesne, J. Navarro, *ibid.* **92**, 024305 (2015).
- [6] A. Pastore, D. Davesne, K. Bennaceur, J. Meyer, V. Hellemans, *Phys. Scripta* **T154**, 014014 (2013).
- [7] P. Becker, D. Davesne, J. Meyer, J. Navarro, A. Pastore, *Phys. Rev. C* **96**, 044330 (2017).
- [8] D. Davesne, A. Pastore, J. Navarro, *ibid.* **89**, 044302 (2014).
- [9] N. Iwamoto, C.J. Pethick, *Phys. Rev. D* **25**, 313 (1982).
- [10] A. Pastore, M. Martini, V. Buridon, D. Davesne, K. Bennaceur, J. Meyer, *Phys. Rev. C* **86**, 044308 (2012).
- [11] A. Pastore, M. Martini, D. Davesne, J. Navarro, S. Goriely, N. Chamel, *ibid.* **90**, 025804 (2014).
- [12] P.S. Shternin, D.G. Yakovlev, C.O. Heinke, W.C.G. Ho, D.J. Patnaude, *Mon. Not. R. Astron. Soc. Lett.* **412**, L108 (2011).
- [13] J. Margueron, Nguyen Van Giai, J. Navarro, *Phys. Rev. C* **72**, 034311 (2005).
- [14] J. Dechargé, D. Gogny, *ibid.* **21**, 1568 (1980).
- [15] A. De Pace, M. Martini, *ibid.* **94**, 024342 (2016).
- [16] J. Margueron, J. Navarro, Nguyen Van Giai, P. Schuck, *ibid.* **77**, 064306 (2008).
- [17] D. Davesne, J. Meyer, A. Pastore, J. Navarro, *Phys. Scr.* **90**, 114002 (2015).
- [18] C. Garcia-Recio, J. Navarro, L.L. Salcedo, Nguyen Van Giai, *Ann. Phys.* **214**, 293 (1992).
- [19] P. Becker, D. Davesne, J. Meyer, A. Pastore, J. Navarro, *J. Phys. G: Nucl. Part. Phys.* **42**, 034001 (2015).
- [20] D. Davesne, P. Becker, A. Pastore, J. Navarro, *Ann. Phys.* **375**, 288 (2016).
- [21] X. Viñas, C. Gonzalez-Boquera, M. Centelles, LM. Robledo, C. Mondal, *arXiv preprint arXiv:1810.07469* (2018).
- [22] M. Martini, A. De Pace, K. Bennaceur, *arXiv preprint arXiv:1806.02080* (2018).