This is a repository copy of *Dispersal limitations and historical factors determine the biogeography of specialized terrestrial protists*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/148666/>

Version: Accepted Version

Article:

Singer, David, Mitchell, Edward A D, Payne, Richard John et al. (1 more author) (2019) Dispersal limitations and historical factors determine the biogeography of specialized terrestrial protists. *Molecular Ecology*. ISSN 0962-1083

<https://doi.org/10.1111/mec.15117>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Dispersal limitations and historical factors determine the biogeography of specialized terrestrial protists

David Singer^{1,2}, Edward A.D. Mitchell^{1,3}, Richard J. Payne⁴, Quentin Blandenier^{1,5}, Clément Duckert¹, Leonardo D. Fernández^{1,6}, Bertrand Fournier⁷, Cristián E. Hernández⁸, Gustaf Granath⁹, Håkan Rydin⁹, Luca Bragazza^{10,11,12}, Natalia G. Koronatova¹³, Irina Goia¹⁴, Lorna I. Harris¹⁵, Katarzyna Kajukalo¹⁶, Anush Kosakyan¹⁷, Mariusz Lamentowicz¹⁶, Natalia P. Kosykh¹³, Kai Vellak¹⁸, Enrique Lara^{1,5}

Corresponding author: David Singer, Department of Zoology, Institute of Biosciences, Rua do Matão, Tv. 14, n101, São Paulo, SP - 05508-090, Brazil, tel: 0041792182426, email: david.singer.bio@outlook.com

Affiliations:

¹Laboratory of Soil Biodiversity, Institute of Biology, University of Neuchâtel, Rue Emile-Argand 11, CH-2000 Neuchâtel, Switzerland

²Department of Zoology, Institute of Biosciences, University of São Paulo, 05508-090, Brazil

³Jardin Botanique de Neuchâtel, Chemin du Perthuis-du-Sault 58, CH-2000 Neuchâtel, Switzerland

⁴Environment, University of York, Heslington, York, YO10 5DD, United Kingdom

⁵Real Jardín Botánico, CSIC, Plaza de Murillo 2, 28014 Madrid, Spain

⁶Centro de Investigación en Recursos Naturales y Sustentabilidad (CIRENYS), Universidad Bernardo O'Higgins, Avenida Viel 1497, Santiago, Chile

⁷Community and Quantitative Ecology Laboratory, Department of Biology, Concordia University, 7141 Sherbrooke Street West, Montreal, QC H4B 1R6, Canada

⁸Centro de Investigación en Recursos Naturales y Sustentabilidad (CIRENYS), Universidad Bernardo O'Higgins, Avenida Viel 1497, Santiago, Chile

⁹Department of Ecology and Genetics, Evolutionary Biology Centre, Uppsala University, Norbyvägen 18D, SE-752 36 Uppsala, Sweden.

¹⁰WSL Swiss Federal Institute for Forest, Snow and Landscape Research, Site Lausanne, Station 2, CH-1015 Lausanne, Switzerland

¹¹Laboratory of Ecological Systems (ECOS), Ecole Polytechnique Fédérale de Lausanne (EPFL), School of Architecture, Civil and Environmental Engineering (ENAC), Station 2, CH-1015 Lausanne, Switzerland

¹²Department of Life Science and Biotechnologies, University of Ferrara, Corso Ercole I d'Este 32, I-44121 Ferrara, Italy

¹³Laboratory of Biogeocenology, Institute of Soil Science and Agrochemistry, Siberian Branch of Russian Academy of Sciences, Ak. Lavrent'ev ave., 8/2, Novosibirsk, 630090 Russia

¹⁴Babeş-Bolyai University, Faculty of Biology and Geology, Department of Taxonomy and Ecology, 42 Republicii Street, RO-400015, Cluj-Napoca, Romania

¹⁵School of Geography and Earth Sciences, McMaster University, General Sciences Building, Hamilton, Ontario, L8S 4K1, Canada

¹⁶Laboratory of Wetland Ecology and Monitoring, Faculty of Geographical and Geological Sciences and Department of Biogeography and Paleoecology, Adam Mickiewicz University, Poznań, Poland

¹⁷Institute of Parasitology, Biology Center, Czech Academy of Sciences, Branisovska 1160/31, 37005 Czeske Budejovice, Czech Republic

¹⁸Institute of Ecology and Earth Sciences, Natural History Museum, University of Tartu, Lai St 40, Tartu 51005, Estonia

Abstract

Recent studies show that soil eukaryotic diversity is immense and dominated by microorganisms. However, it is unclear to what extent the processes that shape the distribution of diversity in plants and animals also apply to microorganisms. Major diversification events in multicellular organisms have often been attributed to long-term climatic and geological processes, but the impact of such processes on protist diversity has received much less attention as their distribution has often been believed to be largely cosmopolitan. Here, we quantified phylogeographic patterns in *Hyalosphenia papilio*, a large testate amoeba restricted to Holarctic *Sphagnum*-dominated peatlands, to test if the current distribution of its genetic diversity can be explained by historical factors or by the current distribution of suitable habitat. Phylogenetic diversity was higher in Western North America, corresponding to the inferred geographical origin of the *H. papilio* complex, and was lower in Eurasia despite extensive suitable habitat. These results suggest that patterns of phylogenetic diversity and distribution can be explained by the history of Holarctic *Sphagnum* peatland range expansions and contractions in response to Quaternary glaciations that promoted cladogenetic range evolution, rather than the contemporary distribution of suitable habitats. Species distributions were positively correlated with climatic niche breadth, suggesting that climatic tolerance is key to dispersal ability in *H. papilio*. This implies that, at least for large and specialized terrestrial microorganisms, propagule dispersal is slow enough that historical processes may contribute to their diversification and phylogeographic patterns and may partly explain their very high overall diversity.

Keywords

Hyalosphenia papilio, phylogeography, *Sphagnum* peatland, protists, Distribution, Holarctic

Introduction

The question of whether the same rules structure the diversity of all eukaryotes, micro and – macroscopic alike, has been the subject of a heated debate since the early 2000's. The classical paradigm is that “everything is everywhere, but, the environment selects” (Baas-Becking, 1934). Defenders of this paradigm have argued that geographic barriers are ineffective in preventing dispersal of microbes (Fenchel, 2005; Finlay, 1998). Other researchers, while accepting that some microbes do indeed occur worldwide, have argued that others are clearly restricted to certain regions: the ‘moderate endemism’ model (Foissner, 1999). This argument is based particularly on a limited number of so-called “biogeographic flagship species”, with conspicuous morphology.

The application of barcoding to protists has brought new nuance to the debate (Pawlowski et al., 2012). Single-cell DNA barcoding studies (Pawlowski et al., 2012) of individual “morphospecies” are now revealing the existence of numerous “cryptic” biological species (Singer et al., 2018). Barcoding studies have now demonstrated geographically limited distributions in soil (Ryšánek, Hřčková, & Škaloud, 2015) freshwater (Škaloud et al., 2019) and marine organisms (Santoferrara, Rubin, & Mcmanus, 2018), although cases of cosmopolitan distribution have also been reported (Geisen, Fiore-Donno, Walochnik, & Bonkowski, 2014; Šlapeta, López-García, & Moreira, 2005). The development of microbial phylogeography, combining biogeography and molecular phylogeny, in turn allows the evaluation of possible drivers of diversity patterns, and comparison to those known to drive plant and animal diversity (Martiny et al., 2006).

Among terrestrial microorganisms, testate amoebae are particularly useful models for phylogeographical studies. Testate amoebae are conspicuous and relatively easy to identify

and are also large enough to be isolated individually for DNA barcoding. Many species have narrow ecological tolerances and thus can only colonize specific, often geographically discontinuous habitats (Singer et al., 2018). Furthermore, some species have well-documented distribution and ecology. A good example is *Hyalosphenia papilio*, a widely recorded and morphologically distinctive testate amoeba taxon (Fig. 1). Of particular interest for phylogeographic studies is that, based on single cell barcoding and the variable molecular marker Cytochrome c Oxidase subunit I (COI), *H. papilio* is known to represent a species complex of at least twelve lineages (Heger, Mitchell, & Leander, 2013).

Hyalosphenia papilio is found exclusively in Holarctic *Sphagnum*-dominated peatlands (Amesbury et al., 2018, 2016) and it is known to be absent from similar southern hemisphere sites despite extensive study (Fernández, Lara, & Mitchell, 2015; Smith, Bobrov, & Lara, 2007). *Sphagnum*-dominated peatlands are comparatively young ecosystems, dating back to the expansion of boreal and subarctic environments near the Pliocene (Shaw et al., 2010). *Sphagnum* is an ecosystem engineer that modifies habitats by increasing soil wetness and decreasing pH and available nutrient content, producing decay-resistant litter rich in phenols and sphagnum (van Breemen, 1995) and hosting very distinctive prokaryotic, algal and fungal communities (Kostka et al., 2016; Mutinová, Neustupa, Bevilacqua, & Terlizzi, 2016). Thus, *Sphagnum* represents a highly selective habitat for macro and microorganisms. This explains why *Sphagnum*-dominated ecosystems are species-poor and these same factors are likely to also drive evolutionary adaptations in testate amoebae (Kosakyan et al., 2016; Singer et al., 2018). Hence, it is likely that this taxon does not pre-date the radiation of peat-forming *Sphagnum* species between 17 and 7 Mya (Shaw et al., 2010). Large extents of *Sphagnum* comparable to modern *Sphagnum*-dominated peatlands probably appeared during the late Miocene/early Pliocene, concomitantly with global cooling, i.e. between 7 and 5.5 Mya

(Herbert et al., 2016). While *Sphagnum* occurs at low as well as high latitudes it is only a dominant component of peatlands in higher latitudes (e.g. Tierra-del-Fuego and the boreal zone of the Holarctic). The taxonomic richness of the genus is low in the Southern Hemisphere high latitudes and high in the Northern Hemisphere high latitudes which correspond to its inferred origin (Shaw, Carter, Aguero, da Costa, & Cowl, 2019).

Holarctic *Sphagnum*-dominated peatlands have experienced considerable changes in their extent due to the repeated advances and retreats of ice sheets during the Quaternary. Many of the largest areas covered by peatlands today were under ice during the last glacial maximum (e.g. Fennoscandia, boreal Canada), while peatlands may have persisted in others (e.g. Pacific coast of Canada)(Treat et al., 2019). These successive glacial expansions and contractions are known to have shaped genetic diversity in multicellular taxa (Schönswetter, Stehlik, Holderegger, & Tribsch, 2005), whose dispersal is assumed to be slow in contrast with eukaryotic microorganisms (Bahram et al., 2016). If, like plants and animals, protist dispersal is relatively slow, the genetic structure of their populations will bear traces of such range expansions and contractions. The origin of taxa can potentially be inferred and the timing of phylogenetic events estimated based on molecular clocks (Arbogast et al., 2006; Arbogast, Edwards, Wakeley, Beerli, & Slowinski, 2002). On the contrary, fast dispersal in protists would blur any such signature and taxonomic or phylogenetic diversity would tend to be distributed randomly and peak in areas with largest extent of favourable habitats (Forest, Colville, & Cowling, 2018).

It follows that the phylogeographical pattern of a given taxon, here *H. papilio*, can be used to test two alternative hypotheses: 1) Dispersal is low and/or slow enough so that traces of glacial cycles are reflected in its extant diversity. The highest diversity, and the likely geographical origin of *H. papilio* would be expected to occur in refugia corresponding to the

margins of ice sheets during Last Glacial Maximum where *Sphagnum* peatlands could survive. 2) Dispersal is high and/or fast, and diversity would be expected to be maximal where the largest expanses of *Sphagnum* peatlands are found today (e.g. Western Siberia). Empirical evidence demonstrates a relationship between testate amoeba shell size and geographic range (Wilkinson, 2001). Population genetics analyses (Lara, Heger, Scheihing, & Mitchell, 2011) and modelling (Wilkinson, Koumoutsaris, Mitchell, & Bey, 2011) show a decline in dispersal potential for testate amoebae and theoretical organisms of smaller sizes (ca. 60 μm) than that reported for *H. papilio* (size range 90-175 μm). The ability of entering a dormant stage (cysts) which can withstand desiccation and other stresses is considered to be a key dispersal trait in protists (Geisen et al., 2018); however, such structures have never been reported in *H. papilio*. We therefore predict that the first hypothesis is more likely to be supported.

Material and Methods

Dataset preparation

We retrieved all 360 existing COI gene sequences of *H. papilio* from GenBank, together with information on the origin of the cells from four studies (Gomaa et al., 2014; Heger et al., 2013; Kosakyan et al., 2012; Oliverio, Lahr, Grant, & Katz, 2015). In addition, we isolated 57 single cells of *H. papilio* from *Sphagnum* samples collected at 13 new sites, targeting under-sampled regions to compile a global dataset (Supplementary Table 1). Briefly, the cells were washed three times in autoclaved distilled water before DNA extraction, which was performed using the guanidine thiocyanate-base protocol (Duckert et al., 2018). Amplifications of COI gene fragments were performed in two steps: a first PCR was

undertaken with the general COI primers LCO1490 and HCO2198 (Folmer, Black, Hoeh, Lutz, & Vrijenhoek, 1994), which was followed by a nested PCR using, *H. papilio*-specific primers HPcoiF and HPcoiR (Gomaa et al., 2014). The first DNA amplification profiles consisted of an initial denaturation step for 3 min at 95°C, followed by 39 cycles of 15 sec of denaturation at 95°C, 15 sec of annealing at 43°C and 1 min of elongation at 72°C with an additional final elongation step at 72°C for 10 min. The procedure for the second PCR profile was the same except that the annealing temperature was increased to 55 °C. Sequencing was carried out using a BigDye197 Terminator Cycle Sequencing Ready Reaction Kit (Applied Biosystems) and analysed with an ABI-3130XL DNA sequencer (Applied Biosystems). Sequences were deposited in GenBank with the following accession numbers: MK823130- MK823186. COI sequences were edited and aligned (ClustalW algorithm (Thompson, Gibson, & Higgins, 2002)) using Bioedit (V.7.2.3 (Hall, 1999)). The final dataset including published and new sequences consisted of 418 sequences from 61 sites (Supplementary Table 1).

Lineage delineation

We delimited genetic lineages following the approach described in Heger (2013). Briefly, to obtain a general overview of the existing lineages, we first constructed the phylogenetic tree based on the matrix of the unique sequences (haplotypes) among the 418 considered in this study. The sequence lengths of the dataset vary from 430 to 620 bp (depending on the primers used to barcode the isolated cells). We constructed both a Maximum Likelihood (ML) and a Bayesian tree, with the RAxML algorithm (Stamatakis, Hoover, Rougemont, & Renner, 2008) and MrBayes v. 3.2.6 (Ronquist & Huelsenbeck, 2003), using in both cases a GTR + Γ model. We then tested if the haplotypes were distributed randomly by comparing our observed distribution with a null model obtained by haplotypes randomly attributed to

lineages (10000 replicates). The tree root was placed between two major clades (clade I contains the lineage L, K, M, J and clade II contains the lineages C, DE, F, B, A, G, H, I) that appeared well supported in earlier works (Heger et al., 2013; Kosakyan et al., 2012). Bi-partition support values were evaluated with 1,000 bootstrap replicates. Bayesian MCMC analysis was carried out with two simultaneous chains and 50,000,000 generations. Trees were sampled every 1,000 generation and the burn-in was set at 25%. The trees were rooted internally based on the topology of trees obtained in earlier works (Heger et al., 2013; Kosakyan et al., 2016), which showed two major clades with maximum support; we rooted the tree between these two clades. As both trees were congruent, we presented only the ML tree and used the Bayesian analysis to evaluate the nodes posterior probabilities (pp). We used three independent methods of lineage delimitation to compare our assignments with those of Heger 2013: 1) Automatic Barcode Gap Discovery (ABGD) (Fontaneto, Flot, & Tang, 2015; Puillandre, Lambert, Brouillet, & Achaz, 2011) using the ABGD web-server <http://www.wabi.snv.jussieu.fr/public/abgd/abgdweb.html>, 2) The sequence divergences using the Kimura 2-parameter (Kimura, 1980; Nasonova, Smirnov, Fahrni, & Pawlowski, 2010) using the “ape” package (V3.2 (Paradis, Claude, & Strimmer, 2004)) in R (V3.0.1. (R. Core Team, 2014)), and 3) GMYC analysis performed with the SPLITS package, version 1.0-19 (Fujisawa & Barraclough, 2013) coded in R, version 3.1.2 (R. Core Team, 2014).

Haplotype and lineage network

Haplotypes (defined as genetic units separated by at least a single mutation) were assigned to the previously determined lineages. Haplotype networks were constructed using minimum spanning network analysis as implemented in the software PopART (V1.7 (Leigh & Bryant, 2015), Supplementary Table 2). Four main geographical zones (Eastern North America,

Western North America, Europe and Asia) were defined to highlight the distribution of the haplotypes.

Historical Biogeography

We first tested whether the observed patterns could be due to chance or to a sampling bias by calculating the observed beta diversity following (Legendre & De Cáceres, 2013). We then tested if the observed beta diversity was higher than expected by chance. We simulated beta diversity values under null expectations and compared them to observed beta diversity values to obtain p-values and standardized effect sizes. Simulated beta diversity values were calculated using the same approach (Legendre & De Cáceres, 2013) on a permuted site by species matrix. Permutations were conducted using the permatswap algorithm of the R-package “vegan” (Oksanen, Blanchet, & Kindt, 2015) which preserves column sums. This allows us to randomly attribute species to station while preserving species total abundance.

In order to determinate whether species distribution areas were correlated with ecological tolerance, we determined the climatic niche breadth for each species using the tolerance index (Dolédec, Chessel, & Gimaret-Carpentier, 2000), with the R package “ade4” (Dray & Dufour, 2007). This index estimates niche breadth based on environmental tolerance (i.e. climate) (Hurlbert, 1978; Thuiller, 2004) using the dispersion of geographic cells that contain the target species in the climatic multivariate space. Low values of the index suggest narrow tolerance while high values correspond to generalists. These indices were inferred based on geographical coordinates for each occurrence (Supplementary Table 1) and interpolated climate data sets (Bioclim, 19 variables) that were generated at 2.5 arcmin resolution from meteorological data (Hijmans, Cameron, Parra, Jones, & Jarvis, 2005). We then estimated distribution areas of the species based on the number of plots where a given species was

observed (i.e. a rough estimate of the spatial range) and plotted it against the estimated climatic niche breadths.

To evaluate the evolutionary events that may explain the current distribution of *H. papilio* (e.g. dispersal, extinction, range-switching, sympatry, vicariance and founder effect), we estimated the ancestral distribution and the frequency of event counts in each of 1000 Biogeographic Stochastic Mapping (BSM) analyses, using the BioGeoBEARS package (Matzke, 2013) in R (V3.0.1. (R. Core Team, 2014)). BioGeoBEARS allows for the estimation of ancestral geographic ranges on dated phylogeny, comparing several models of range evolution. We used the DEC model (Ree, Smith, & Baker, 2008) with two free parameters: "d" (dispersal rate) and "e" (extinction rate), and a fixed cladogenetic model (cladogenetic event allowed: vicariance, sympatric-subset speciation, and sympatric range-copying). We also used a DEC model with an extra parameter, "j", which represents the founder-event speciation, where the new species "jumps" to a range outside of the ancestral range (DEC + j model). The comparison of these two models was performed using Akaike Information Criterion (AIC). The age of the nodes of the rescaled BEAST Tree of *H. papilio* was estimated by constraining its root to 7 Mya, which corresponds to the documented origin of *Sphagnum* peatlands (Stenøien, Shaw, Shaw, Hassel, & Gunnarsson, 2010), as all known lineages of *H. papilio* are restricted to these environments. This type of approach has been used to identify the geographical origin of multicellular taxa of different ages, dispersion strategies and lifestyles, including hyacinthoid monocots (Ali, Yu, Pfosser, & Wetschnig, 2012), chameleons (Tolley, Townsend, & Vences, 2013) and bees (Trunz, Packer, Vieu, Arrigo, & Praz, 2016), and is used here, to our knowledge, for the first time in microorganisms. This approach allowed us to test the two hypotheses: If the frequency of event counts in each of 1000 BSM sustain low frequency of dispersal events, related to other

biogeographic events, the first hypothesis is supported (1.- Dispersal is low and/or slow enough so that traces of glacial cycles are reflected in its extant diversity), and otherwise, the second hypothesis is supported (2.- Dispersal is high and/or fast, and diversity would be expected to be maximal where the largest expanses of *Sphagnum* peatlands are found today).

Results

Lineage delineation and diversity

The 418 mitochondrial COI sequences of *H. papilio* revealed the existence of 13 or 14 distinct lineages (Fig. 2). The Kimura 2-parameter test suggested the existence of 14 lineages based on a threshold of $\geq 1\%$ sequence divergence. The Generalized Mixed Yule Coalescent (GMYC) method yielded 13 lineages (lower and upper confidence intervals: 10 and 29 lineages, respectively; $P = 0.046$) based on single threshold methods. Finally, the Automatic Barcode Gap Discovery (ABGD) method identified 13 lineages, using a distinctive barcoding gap of 7%. One of these lineages, called here "M" has not been previously recorded. This lineage was recovered from localities not included in previous studies (i.e. (Gomaa et al., 2014; Heger et al., 2013; Kosakyan et al., 2012; Oliverio et al., 2015)). It was supported by all analyses, although the Kimura 2-parameter test suggested dividing it into two (Fig. 2).

Phylogenetic reconstruction

Sequences from the previously overlooked lineage M diverged from all others ($pp = 1$) and branched as a sister group to lineages J and K (Fig. 2). Only a single haplotype was retrieved from Lineage E, and five from lineage D (*sensu* Heger et al. (2013)). Here again the genetic

divergence was low (i.e. at most six nucleotides difference between the sequence of lineage E, and the five sequences of lineage D, all of which were separated by a single nucleotide (Fig. 3)).

Haplotype network

The haplotype network (Fig. 3) showed that some lineages (B, H, and L) were composed of only a single haplotype, whereas others included several haplotypes, independently of the number of individual cells barcoded. Some lineages were relatively rare (e.g. B, L and M with seven, two, and seven individuals respectively) whilst others were extensively recorded (e.g. lineage A was identified more than a hundred times). Null model analyses show that such a pattern is not expected under random assembly of lineages (Supplementary Fig. 1).

Spatial patterns of phylogenetic richness

We found that the observed beta diversity was significantly higher than expected by chance ($SES = 1.71$; $p=0.99$), showing a strong spatial structuring of diversity. We also found a strong positive correlation between niche breadth as estimated using Dolédec tolerance indices and distribution areas ($R^2 = 0.75$; $P=0.001$). We also found that lineages differed in their climatic niches with some lineages preferring colder and drier conditions (Supplementary Fig. 2, lineage H) and others preferring warmer conditions with abundant precipitation (Supplementary Fig. 2, lineage I).

The geographical distribution of phylogenetic richness showed a clear contrast (Fig 1, 3). Only four lineages (A, C, J and G) were recovered from all of Eurasia, five from Eastern

North America (A, F, K, J and M), and nine from Western North America (six in Alaska and five in the Pacific Northwest, only two being shared between these two regions). Thus, regional as well as overall diversity and diversity turnover were all higher in North America than in Eurasia.

The distribution of the different lineages (Fig. 3) suggests that several haplotypes are specific to certain geographical areas (B, DE, H, I and L occur only in Western North America, while K and M occurred only in Eastern North America), whereas others were geographically widespread (e.g. J is found throughout the Holarctic realm). Null model analyses show that such a pattern is not expected under random assembly of lineages (Supplementary Fig. 1). This structure in lineage distribution suggests that geographic dispersal has occurred comparatively slowly, allowing it to be recovered with a genetic marker such as mtCOI used for species-level delineation in this group of organisms (Kosakyan et al., 2012).

Origin of lineages and evaluation of the diversification processes

The AIC selection of biogeographic models implemented in BioGeoBEARS indicated that a DEC model was the best-supported (Supplementary Table 3). Based on this model, the most likely ancestral areas for *H. papilio* are in Western North America (Fig. 4). The dispersal summary extracted from the 1,000 BSM's maps showed that most of the dispersal events occurred from Western North America and Asia to the other biogeographic areas, and from Asia to Europe (Supplementary Table 4). The results of the ancestral area estimation and number of dispersal events analyses showed that the most frequent process during the historical biogeography of *H. papilio* was narrow sympatry (i.e. when the ancestral range contains one area, and both daughter lineages inherit that area), followed by a low frequency

of dispersal events (range expansion) (Supplementary Fig. 1). The importance of vicariance and founder events were comparatively limited (Supplementary Fig. 1).

Discussion

Diversity and geographical distribution of the lineages and haplotypes

The species complex *H. papilio* is represented by at least 13 lineages in the Holarctic region, one of which had not been previously described. Although it is possible that some lineages remain to be discovered, our globally extensive sampling retrieved only one additional lineage (M), suggesting that we have now captured most of the group's diversity. The genetic distances determined by our taxon delineating approaches are consistent with the barcoding gap (<4%) used to discriminate species in other related testate amoebae lineages (e.g. genus *Nebela*, Hyalospheniidae). The above-mentioned lineages were defined as species under multiple and independent concepts, including ecological, morphological and evolutionary (Singer et al., 2018). This might imply that the lineages retrieved in the present study can all be considered as separate species (Kosakyan, Gomaa, Mitchell, Heger, & Lara, 2013; Kosakyan et al., 2012; Singer, Kosakyan, Pillonel, Mitchell, & Lara, 2015; Singer et al., 2018). The accuracy of a species tree built on a single locus may be still questioned, especially in the case of recent radiations, as the existence of several caveats (like sequencing pseudogenes, ongoing hybridization processes) cannot be ruled out and may distort the tree's topology. In Amoebozoa, COI has been chosen as the most accurate marker notably because of its sensitivity and lack of intra-individual variability (Nassonova et al., 2010) and we therefore consider it reliable.

Lineages of *H. papilio* show different distribution patterns over the Holarctic realm. Four lineages (J, A, C, G) were found in several regions with contrasted climates (Fig. 1)

suggesting that they have a greater ecological tolerance. This is corroborated by the strong correlation between climatic niche breadth and estimated distribution ranges (Fig. 5), suggesting that colonization capacity is constrained by specific tolerance to climates. If these distributional patterns reflect evolutionary adaptation to long-distance dispersal, it would then imply that the required physiological/lifestyle adaptations to long range migration have appeared independently at least four times in the history of the *H. papilio* species complex (Fig. 4).

The existence of restricted distributions is even clearer at the haplotype level. Of the 74 total *H. papilio* haplotypes, only seven (9.2%) were present in two zones, two in three zones (2.6%) and no single haplotype was found in all four zones. This indicates that even widespread lineages (e.g. lineage J) show high infra-specific genetic structuring, which suggests limited gene flow among sites, and thus, geographical isolation (Fernández, Hernández, Schiaffino, Izaguirre, & Lara, 2017; Lara, Heger, Scheihing, & Mitchell, 2010).

Hypothesis 2, that diversity is maximal where the largest expanses of *Sphagnum* peatlands are found today cannot be supported by our data and analyses. Under this hypothesis, highest diversity would be expected in regions such as Western Siberia where peatlands are at their most extensive and cover more than 20% of the landscape (Peregon, Maksyutov, Kosykh, & Mironycheva-Tokareva, 2008). However, only four “far travelled” lineages were found in the entire of Eurasia, as compared to 13 in North America. This is despite larger overall area, more extensive *Sphagnum* peatland extent and an extensive range of climatic conditions. Six lineages, 50% more than the entire of Eurasia, were found only in Alaska. In contrast, most genetic diversity seems to be located along the Western North American coast, a region where peatlands are typically small and scattered today. This fact, together with the strong spatial

patterns in lineage distribution observed (Fig. 1, 4) advocates against our hypothesis 2 (fast dispersal).

Geographic origin and influence of historical events

All Eurasian lineages identified were also present in North America, while several lineages were restricted to North America. This observation alone suggests an American origin for *H. papilio*. Our ancestral range reconstruction corroborates this inference, placing the most probable origin of the *H. papilio* complex in Western North America (Fig. 4).

Dating speciation events is difficult in testate amoebae as their lineages cannot be morphologically distinguished (Mulot et al., 2017); testate amoeba shell records in peat are rare before the Holocene. Nevertheless, it is still possible to infer a time window for the radiation of the lineages indirectly based on the very strict habitat specificity of this taxon. All lineages of *H. papilio* thus-far identified are restricted to *Sphagnum* peatlands. It is therefore reasonable to assume that this highly adapted taxon evolved within these ecosystems. The oldest fossils of genus *Hyalosphenia* were described from the Triassic (*H. baueri* 220 MYA) (Schönborn, Dörfelt, Foissner, Krienitz, & Schäffer, 1999). *H. baueri* shares some traits like an “indistinctly vase shape” and the presence of an organic lip surrounding the aperture with *H. papilio*. However, it is far from clear that both taxa are directly related. Firstly, it has been shown that the genus *Hyalosphenia* is paraphyletic, as *H. papilio* and *H. elegans* are only distantly related (Lahr et al., 2019; Lara, Heger, Ekelund, Lamentowicz, & Mitchell, 2008). Furthermore, a rough calculation can rule out the possibility of a very old age for *H. papilio*. “Standard” coxI (estimated for animals) mutation rates are typically in the range of a few percent per million year (Ho & Lo, 2013; Papadopoulou, Anastasiou, & Vogler, 2010), sometimes much higher (Ney, Frederick, & Schul, 2018). The most divergent *H. papilio*

sequences are separated by roughly 10%, which implies that, in order for the deepest branching in the complex (see Fig. 4) to be 100 MYA old, the mutation rate would need to be of 0.01% / MYA. This is far below all rates known to date, and even lower than the mutation rate of cnidarians which are known for their extremely slow evolving mitochondria (Park et al., 2012). By contrast, to obtain an age of 7 million years the mutation rate would need to be 1.3% / MYA, which is similar to the mutation rate of many arthropods and thus more parsimonious than the alternative.

During the Pleistocene, large areas of North America were intermittently covered by ice although ice-free refugia remained. The area of *Sphagnum* peatlands likely repeatedly expanded during inter-glacial periods and contracted in response to glacial periods when ice masses covered most of the landscape (Shaw et al., 2014). This period is also coetaneous with most cladogenesis in the *H. papilio* phylogenetic history, which suggests a series of speciation events by cladogenetic range evolution which may have occurred during interglacial periods (Fig. 4). Indeed, our analyses show that at least 8 out of 12 cladogenesis events occurred during the Pleistocene, immediately after the 2.5 MYA boundary (Fig. 4).

This hypothesis is also in line with the fact that the BioGeoBears analyses designated narrow sympatry or the inheritance of the ancestral area of a range by both daughter lineages, as a key process explaining the distribution of *H. papilio* lineages. At the onset of Quaternary glaciations (2.58 Mya), one lineage probably existed in Eastern and two in Western North America (Fig. 4). While the first lineage probably survived south of the ice sheet, where conditions were wet enough to allow the development of peatlands (Shaw et al., 2010), the two others were most likely confined to refugia in Western North America.

412 The location of these refugia is known to have shaped the distribution of plants (Eidesen et
413 al., 2013) and animals (Klüttsch, Manseau, Anderson, Sinkins, & Wilson, 2017). In particular,
414 Eastern Beringia (today Alaska and Yukon Territory) was wet enough to support the growth
415 of *Sphagnum* mosses and *Sphagnum* peatlands (Shaw et al., 2013, 2014). These peatlands
416 allowed the survival of associated organisms, likely including the lineages of *H. papilio*. In
417 contrast, Western Beringia (today far eastern Russia) was too dry to support large expanses of
418 *Sphagnum* peatlands (Shaw et al., 2013, 2014) and likely constituted a barrier for the
419 migration of *H. papilio* westwards. Our data suggest that the colonization of the Palaearctic
420 region occurred recently, possibly after the last glaciation (Fig. 1). Western Siberia, which
421 was a cold desert during the Last Glacial Maximum became covered with peatlands after
422 11000 BP (Velichko, Timireva, Kremenetski, MacDonald, & Smith, 2011) and could have
423 constituted a bridge that facilitated the invasion of the Western Palaearctic by “far travelled”
424 lineages of *H. papilio*. Interestingly, a similar pattern has been suggested for the species
425 *Sphagnum angermanicum* (Stenøien et al., 2010).

426 The present-day distribution of lineages and the local palaeogeographical context designates
427 Eastern Beringia or the Pacific Coast as the most probable origin for all extant *H. papilio*
428 lineages. The higher diversity of *H. papilio* haplotypes in North America as compared to
429 Europe mirrors the higher diversity of vascular plants (Earl Latham & Ricklefs, 1993;
430 Svenning, 2003), and both were likely similarly driven by glaciations. The phylogeographic
431 history of *H. papilio*, used here as a convenient model taxon for protists lacking specialised
432 morphological adaptation for dispersal, thus highlights the importance of historical processes
433 in explaining the distribution of extant microbial diversity.

434 Thus, following a dispersal event, sympatric diversification could indeed have played a major
435 role in shaping the current phylogeography of *H. papilio* (Supplementary Fig. 1). It remains to

be determined if the case of *H. papilio* is representative for free-living microorganisms in general. *H. papilio* is large by microbial standards; testate amoebae mostly range between 20 and 200µm and many other protists and most fungi and prokaryotes are smaller. *H. papilio* is also restricted to *Sphagnum* mosses, which, although widespread across the Holarctic, nevertheless constitute a very specific habitat. More generalist, smaller species and/or species possessing structures adapted to dispersal (e.g. fruiting bodies as in many other Amoebozoa, (Shadwick, Spiegel, Shadwick, Brown, & Silberman, 2009) may show patterns which agree better with the second hypothesis. Elucidating the historical processes shaping the diversity of protists with different dispersal strategies, and comparing patterns with better known macroscopic organisms will open the way to understanding the processes of diversification that produced the immense diversity existing today.

Acknowledgements

This work was funded by the Swiss NSF (310003A_143960 & 31003A_163254) and intramural project 201730E063 (CSIC) to E.L., the Swiss NSF (P2NEP3_178543) to D.S., the Swedish Research Council (VR) (2015-05174) to G.G. and H.R., FONDECYT (11170927) and UBO/VRIP (170201) to L.D.F., FONDECYT (1170815) to C.E.H and the Russian Foundation for Basic Research, N 16-55-16007 to N.G.K.. We thank Indrek Hiiesalu for help during fieldwork, Boris Droz and Christopher Niewoehner (University of North California) and Matt McGlone (Manaaki Whenua / Landcare Research, Lincoln, NZ) for useful comments on the manuscript.

References

- Ali, S. S., Yu, Y., Pfosser, M., & Wetschnig, W. (2012). Inferences of biogeographical histories within subfamily Hyacinthoideae using S-DIVA and Bayesian binary MCMC analysis implemented in RASP (Reconstruct Ancestral State in Phylogenies). *Annals of Botany*, 109(1), 95–107. doi: 10.1093/aob/mcr274
- Amesbury, M. J., Booth, R. K., Roland, T. P., Bunbury, J., Clifford, M. J., Charman, D. J., ... Hughes, P. D. M. (2018). Towards a Holarctic synthesis of peatland testate amoeba ecology: Development of a new continental-scale palaeohydrological transfer function for North America and comparison to European data. *Quaternary Science Reviews*, 201, 483–500.
- Amesbury, M. J., Swindles, G. T., Bobrov, A., Charman, D. J., Holden, J., Lamentowicz, M., ... Payne, R. J. (2016). Development of a new pan-European testate amoeba transfer function for reconstructing peatland palaeohydrology. *Quaternary Science Reviews*, 152, 132–151.
- Arbogast, B. S., Drovetski, S. V., Curry, R. L., Boag, P. T., Seutin, G., Grant, P. R., ... Anderson, D. J. (2006). The origin and diversification of Galapagos mockingbirds. *Evolution*, 60(2), 370–382.
- Arbogast, B. S., Edwards, S. V., Wakeley, J., Beerli, P., & Slowinski, J. B. (2002). Estimating divergence times from molecular data on phylogenetic and population genetic timescales. *Annual Review of Ecology and Systematics*, 33(1), 707–740.
- Baas-Becking, L. G. M. (1934). *Geobiologie; of inleiding tot de milieukunde*. WP Van Stockum & Zoon NV.
- Bahram, M., Kohout, P., Anslan, S., Harend, H., Abarenkov, K., & Tedersoo, L. (2016). Stochastic distribution of small soil eukaryotes resulting from high dispersal and drift in a local environment. *The ISME Journal*, 10(4), 885.
- Dolédec, S., Chessel, D., & Gimaret-Carpentier, C. (2000). Niche separation in community analysis: a new method. *Ecology*, 81(10), 2914–2927.
- Dray, S., & Dufour, A.-B. (2007). The ade4 package: implementing the duality diagram for ecologists. *Journal of Statistical Software*, 22(4), 1–20.
- Duckert, C., Blandenier, Q., Kupferschmid, F. A. L., Kosakyan, A., Mitchell, E. A. D., Lara, E., & Singer, D. (2018). En garde! Redefinition of *Nebela militaris* (Arcellinida, Hyalospheniidae) and erection of *Alabasta* gen. nov. *European Journal of Protistology*, 66, 156–165.
- Earl Latham, R., & Ricklefs, R. E. (1993). Global patterns of tree species richness in moist forests: energy-diversity theory does not account for variation in species richness. *Oikos*, 67(2), 325–333. doi: 10.2307/3545479
- Eidesen, P. B., Ehrich, D., Bakkestuen, V., Alsos, I. G., Gilg, O., Taberlet, P., & Brochmann, C. (2013). Genetic roadmap of the Arctic: plant dispersal highways, traffic barriers and capitals of diversity. *New Phytologist*, 200(3), 898–910. doi: 10.1111/nph.12412
- Fenchel, T. (2005). Cosmopolitan microbes and their ‘cryptic’ species. *Aquatic Microbial Ecology*, 41(1), 49–54.
- Fernández, L. D., Hernández, C. E., Schiaffino, M. R., Izaguirre, I., & Lara, E. (2017). Geographical distance and local environmental conditions drive the genetic population structure of a freshwater microalga (Bathycoccaceae; Chlorophyta) in Patagonian lakes. *FEMS Microbiology Ecology*, 93(10). doi: 10.1093/femsec/fix125
- Fernández, L. D., Lara, E., & Mitchell, E. A. D. (2015). Checklist, diversity and distribution of testate amoebae in Chile. *European Journal of Protistology*, 51(5), 409–424.
- Finlay, B. J. (1998). The global diversity of protozoa and other small species. *International Journal for Parasitology*, 28(1), 29–48.
- Foissner, W. (1999). Protist diversity: estimates of the near-imponderable. *Protist*, 150(4), 363–368.
- Folmer, O., Black, M., Hoeh, W., Lutz, R., & Vrijenhoek, R. (1994). DNA primers for amplification of mitochondrial cytochrome c oxidase subunit 1 from diverse metazoan invertebrates.

- Molecular Marine Biology and Biotechnology*, (5). Retrieved from <http://www.vliz.be/en/imis?refid=64543>
- Fontaneto, D., Flot, J.-F., & Tang, C. Q. (2015). Guidelines for DNA taxonomy, with a focus on the meiofauna. *Marine Biodiversity*, 45(3), 433–451. doi: 10.1007/s12526-015-0319-7
- Forest, F., Colville, J. F., & Cowling, R. M. (2018). Evolutionary diversity patterns in the Cape flora of South Africa. In *Phylogenetic Diversity* (pp. 167–187). Springer.
- Fujisawa, T., & Barraclough, T. G. (2013). Delimiting species using single-locus data and the generalized mixed yule coalescent approach: a revised method and evaluation on simulated data sets. *Systematic Biology*, 62(5), 707–724. doi: 10.1093/sysbio/syt033
- Geisen, S., Fiore-Donno, A. M., Walochnik, J., & Bonkowski, M. (2014). *Acanthamoeba* everywhere: high diversity of *Acanthamoeba* in soils. *Parasitology Research*, 113(9), 3151–3158.
- Geisen, S., Mitchell, E. A. D., Adl, S., Bonkowski, M., Dunthorn, M., Ekelund, F., ... Lara, E. (2018). Soil protists: a fertile frontier in soil biology research. *FEMS Microbiology Reviews*, 42(3), 293–323. doi: 10.1093/femsre/fuy006
- Gomaa, F., Kosakyan, A., Heger, T. J., Corsaro, D., Mitchell, E. A. D., & Lara, E. (2014). One alga to rule them all: unrelated mixotrophic testate amoebae (Amoebozoa, Rhizaria and Stramenopiles) share the same symbiont (Trebouxioophyceae). *Protist*, 165(2), 161–176. doi: 10.1016/j.protis.2014.01.002
- Hall, T. A. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series*, 41, 95–98.
- Heger, T. J., Mitchell, E. A. D., & Leander, B. S. (2013). Holarctic phylogeography of the testate amoeba *Hyalosphenia papilio* (Amoebozoa: Arcellinida) reveals extensive genetic diversity explained more by environment than dispersal limitation. *Molecular Ecology*, 22(20), 5172–5184. doi: 10.1111/mec.12449
- Herbert, T. D., Lawrence, K. T., Tzanova, A., Cleaveland Peterson, L., Caballero-Gill, R., & Kelly, C. S. (2016). Late Miocene global cooling and the rise of modern ecosystems. *Nature Geoscience*, 9(11), 843–847. doi: 10.1038/ngeo2813
- Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, 25(15), 1965–1978.
- Ho, S. Y. W., & Lo, N. (2013). The insect molecular clock. *Australian Journal of Entomology*, 52(2), 101–105.
- Hurlbert, S. H. (1978). The measurement of niche overlap and some relatives. *Ecology*, 59(1), 67–77.
- Kimura, M. (1980). A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, 16(2), 111–120.
- Klüttsch, C. F. C., Manseau, M., Anderson, M., Sinkins, P., & Wilson, P. J. (2017). Evolutionary reconstruction supports the presence of a Pleistocene Arctic refugium for a large mammal species. *Journal of Biogeography*, 44(12), 2729–2739. doi: 10.1111/jbi.13090
- Kosakyan, A., Gomaa, F., Mitchell, E. A. D., Heger, T. J., & Lara, E. (2013). Using DNA-barcoding for sorting out protist species complexes: A case study of the *Nebela tinctoria-collaris-bohemica* group (Amoebozoa; Arcellinida, Hyalospheniidae). *European Journal of Protistology*, 49(2), 222–237. doi: 10.1016/j.ejop.2012.08.006
- Kosakyan, A., Heger, T. J., Leander, B. S., Todorov, M., Mitchell, E. A. D., & Lara, E. (2012). COI barcoding of Nebelid testate amoebae (Amoebozoa: Arcellinida): extensive cryptic diversity and redefinition of the Hyalospheniidae Schultze. *Protist*, 163(3), 415–434. doi: 10.1016/j.protis.2011.10.003
- Kosakyan, A., Lahr, D. J. G., Mulot, M., Meisterfeld, R., Mitchell, E. A. D., & Lara, E. (2016). Phylogenetic reconstruction based on COI reshuffles the taxonomy of hyalosphenid shelled

(testate) amoebae and reveals the convoluted evolution of shell plate shapes. *Cladistics*, 32(6), 606–623. doi: 10.1111/cla.12167

Kostka, J. E., Weston, D. J., Glass, J. B., Lilleskov, E. A., Shaw, A. J., & Turetsky, M. R. (2016). The *Sphagnum* microbiome: new insights from an ancient plant lineage. *New Phytologist*, 211(1), 57–64.

Lahr, D. J. G., Kosakyan, A., Lara, E., Mitchell, E. A. D., Morais, L., Porfirio-Sousa, A. L., ... Kang, S. (2019). Phylogenomics and morphological reconstruction of Arcellinida testate amoebae highlight diversity of microbial eukaryotes in the Neoproterozoic. *Current Biology*.

Lara, E., Heger, T. J., Ekelund, F., Lamentowicz, M., & Mitchell, E. A. D. (2008). Ribosomal RNA genes challenge the monophyly of the Hyalospheniidae (Amoebozoa: Arcellinida). *Protist*, 159(2), 165–176.

Lara, E., Heger, T. J., Scheihing, R., & Mitchell, E. A. D. (2010). COI gene and ecological data suggest size-dependent high dispersal and low intra-specific diversity in free-living terrestrial protists (Euglyphida: *Assulina*). *Journal of Biogeography*, 38(4), 640–650. doi: 10.1111/j.1365-2699.2010.02426.x

Lara, E., Heger, T. J., Scheihing, R., & Mitchell, E. A. D. (2011). COI gene and ecological data suggest size-dependent high dispersal and low intra-specific diversity in free-living terrestrial protists (Euglyphida: *Assulina*). *Journal of Biogeography*, 38(4), 640–650.

Legendre, P., & De Cáceres, M. (2013). Beta diversity as the variance of community data: dissimilarity coefficients and partitioning. *Ecology Letters*, 16(8), 951–963.

Leigh, J. W., & Bryant, D. (2015). popart: full-feature software for haplotype network construction. *Methods in Ecology and Evolution*, 6(9), 1110–1116. doi: 10.1111/2041-210X.12410

Martiny, J. B. H., Bohannan, B. J. M., Brown, J. H., Colwell, R. K., Fuhrman, J. A., Green, J. L., ... Kuske, C. R. (2006). Microbial biogeography: putting microorganisms on the map. *Nature Reviews Microbiology*, 4(2), 102.

Matzke, N. J. (2013). *Probabilistic historical biogeography: new models for founder-event speciation, imperfect detection, and fossils allow improved accuracy and model-testing* (UC Berkeley). Retrieved from <https://escholarship.org/uc/item/8227p52c>

Mulot, M., Marcisz, K., Grandgirard, L., Lara, E., Kosakyan, A., Robbroek, B. J. M., ... Mitchell, E. A. D. (2017). Genetic determinism vs. phenotypic plasticity in protist morphology. *Journal of Eukaryotic Microbiology*, 64(6), 729–739. doi: 10.1111/jeu.12406

Mutinová, P. T., Neustupa, J., Bevilacqua, S., & Terlizzi, A. (2016). Host specificity of epiphytic diatom (Bacillariophyceae) and desmid (Desmidiaceae) communities. *Aquatic Ecology*, 50(4), 697–709.

Nassonova, E., Smirnov, A., Fahrni, J., & Pawlowski, J. (2010). Barcoding amoebae: comparison of SSU, ITS and COI genes as tools for molecular identification of naked lobose amoebae. *Protist*, 161(1), 102–115. doi: 10.1016/j.protis.2009.07.003

Ney, G., Frederick, K., & Schul, J. (2018). A Post-pleistocene Calibrated Mutation Rate from Insect Museum Specimens. *PLoS Currents*, 10.

Oksanen, J., Blanchet, F. G., & Kindt, R. (2015). *Vegan: community ecology package. R package version 2.3-0*.

Oliverio, A. M., Lahr, D. J. G., Grant, J., & Katz, L. A. (2015). Are microbes fundamentally different than macroorganisms? Convergence and a possible case for neutral phenotypic evolution in testate amoeba (Amoebozoa: Arcellinida). *Royal Society Open Science*, 2(12), 150414. doi: 10.1098/rsos.150414

Papadopoulou, A., Anastasiou, I., & Vogler, A. P. (2010). Revisiting the insect mitochondrial molecular clock: the mid-Aegean trench calibration. *Molecular Biology and Evolution*, 27(7), 1659–1672.

Paradis, E., Claude, J., & Strimmer, K. (2004). APE: Analyses of Phylogenetics and Evolution in R language. *Bioinformatics*, 20(2), 289–290. doi: 10.1093/bioinformatics/btg412

- Park, E., Hwang, D.-S., Lee, J.-S., Song, J.-I., Seo, T.-K., & Won, Y.-J. (2012). Estimation of divergence times in cnidarian evolution based on mitochondrial protein-coding genes and the fossil record. *Molecular Phylogenetics and Evolution*, 62(1), 329–345.
- Pawlowski, J., Audic, S., Adl, S., Bass, D., Belbahri, L., Berney, C., ... de Vargas, C. (2012). CBOL protist working group: barcoding eukaryotic richness beyond the animal, plant, and fungal kingdoms. *PLOS Biology*, 10(11), e1001419. doi: 10.1371/journal.pbio.1001419
- Peregón, A., Maksyutov, S., Kosykh, N. P., & Mironycheva-Tokareva, N. P. (2008). Map-based inventory of wetland biomass and net primary production in western Siberia. *Journal of Geophysical Research: Biogeosciences*, 113(G1).
- Puillandre, N., Lambert, A., Brouillet, S., & Achaz, G. (2011). ABGD, Automatic Barcode Gap Discovery for primary species delimitation. *Molecular Ecology*, 21(8), 1864–1877. doi: 10.1111/j.1365-294X.2011.05239.x
- R. Core Team. (2014). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. 2013. ISBN 3-900051-07-0.
- Ree, R. H., Smith, S. A., & Baker, A. (2008). Maximum likelihood inference of geographic range evolution by dispersal, local extinction, and cladogenesis. *Systematic Biology*, 57(1), 4–14. doi: 10.1080/10635150701883881
- Ronquist, F., & Huelsenbeck, J. P. (2003). MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics*, 19(12), 1572–1574. doi: 10.1093/bioinformatics/btg180
- Ryšánek, D., Hřčková, K., & Škaloud, P. (2015). Global ubiquity and local endemism of free-living terrestrial protists: phylogeographic assessment of the streptophyte alga *Klebsormidium*. *Environmental Microbiology*, 17(3), 689–698.
- Santoferrara, L. F., Rubin, E., & Mcmanus, G. B. (2018). Global and local DNA (meta) barcoding reveal new biogeography patterns in tintinnid ciliates. *Journal of Plankton Research*, 40(3), 209–221.
- Schönborn, W., Dörfelt, H., Foissner, W., Krienitz, L., & Schäffer, U. (1999). A fossilized microcenosis in Triassic amber. *Journal of Eukaryotic Microbiology*, 46(6), 571–584.
- Schönswetter, P., Stehlik, I., Holderegger, R., & Tribsch, A. (2005). Molecular evidence for glacial refugia of mountain plants in the European Alps. *Molecular Ecology*, 14(11), 3547–3555.
- Shadwick, L. L., Spiegel, F. W., Shadwick, J. D. L., Brown, M. W., & Silberman, J. D. (2009). *Eumycetozoa*=*Amoebozoa*?: SSUrDNA phylogeny of protosteloid slime molds and its significance for the amoebozoan supergroup. *PloS One*, 4(8), e6754.
- Shaw, A. J., Carter, B. E., Agüero, B., da Costa, D. P., & Crowl, A. A. (2019). Range change evolution of peat mosses (*Sphagnum*) within and between climate zones. *Global Change Biology*, 25(1), 108–120.
- Shaw, A. J., Devos, N., Cox, C. J., Boles, S. B., Shaw, B., Buchanan, A. M., ... Seppelt, R. (2010). Peatmoss (*Sphagnum*) diversification associated with Miocene Northern Hemisphere climatic cooling? *Molecular Phylogenetics and Evolution*, 55(3), 1139–1145. doi: 10.1016/j.ympev.2010.01.020
- Shaw, A. J., Golinski, G. K., Clark, E. G., Shaw, B., Stenøien, H. K., & Flatberg, K. I. (2013). Intercontinental genetic structure in the amphi-Pacific peatmoss *Sphagnum miyabeaenum* (Bryophyta: Sphagnaceae). *Biological Journal of the Linnean Society*, 111(1), 17–37. doi: 10.1111/bij.12200
- Shaw, A. J., Shaw, B., Stenøien, H. K., Golinski, G. K., Hassel, K., & Flatberg, K. I. (2014). Pleistocene survival, regional genetic structure and interspecific gene flow among three northern peat-mosses: *Sphagnum inexpectatum*, *S. orientale* and *S. miyabeaenum*. *Journal of Biogeography*, 42(2), 364–376. doi: 10.1111/jbi.12399
- Singer, D., Kosakyan, A., Pilonel, A., Mitchell, E. A. D., & Lara, E. (2015). Eight species in the *Nebela collaris* complex: *Nebela gimlii* (Arcellinida, Hyalospheniidae), a new species described from a

- Swiss raised bog. *European Journal of Protistology*, 51(1), 79–85. doi: 10.1016/j.ejop.2014.11.004
- Singer, D., Kosakyan, A., Seppey, C. V. W., Pillonel, A., Fernández, L. D., Fontaneto, D., ... Lara, E. (2018). Environmental filtering and phylogenetic clustering correlate with the distribution patterns of cryptic protist species. *Ecology*, 99(4), 904–914. doi: 10.1002/ecy.2161
- Škaloud, P., Škaloudová, M., Doskočilová, P., Kim, J. Im., Shin, W., & Dvořák, P. (2019). Speciation in protists: Spatial and ecological divergence processes cause rapid species diversification in a freshwater chrysophyte. *Molecular Ecology*.
- Šlapeta, J., López-García, P., & Moreira, D. (2005). Global dispersal and ancient cryptic species in the smallest marine eukaryotes. *Molecular Biology and Evolution*, 23(1), 23–29.
- Smith, H. G., Bobrov, A., & Lara, E. (2007). Diversity and biogeography of testate amoebae. In *Protist Diversity and Geographical Distribution* (pp. 95–109). Springer.
- Stamatakis, A., Hoover, P., Rougemont, J., & Renner, S. (2008). A rapid bootstrap algorithm for the RAxML web servers. *Systematic Biology*, 57(5), 758–771. doi: 10.1080/10635150802429642
- Stenøien, H. K., Shaw, A. J., Shaw, B., Hassel, K., & Gunnarsson, U. (2010). North American origin and recent European establishments of the Amphi-Atlantic peat moss *Sphagnum Angermanicum*. *Evolution*, 65(4), 1181–1194. doi: 10.1111/j.1558-5646.2010.01191.x
- Svenning, J.-C. (2003). Deterministic Plio-Pleistocene extinctions in the European cool-temperate tree flora. *Ecology Letters*, 6(7), 646–653. doi: 10.1046/j.1461-0248.2003.00477.x
- Thompson, J. D., Gibson, T. J., & Higgins, D. G. (2002). Multiple sequence alignment using ClustalW and ClustalX. *Current Protocols in Bioinformatics*, 2.3.1–2.3.22.
- Thuiller, W. (2004). Patterns and uncertainties of species' range shifts under climate change. *Global Change Biology*, 10(12), 2020–2027.
- Tolley, K. A., Townsend, T. M., & Vences, M. (2013). Large-scale phylogeny of chameleons suggests African origins and Eocene diversification. *Proceedings of the Royal Society of London B: Biological Sciences*, 280(1759), 20130184. doi: 10.1098/rspb.2013.0184
- Treat, C. C., Kleinen, T., Broothaerts, N., Dalton, A. S., Dommain, R., Douglas, T. A., ... Brovkin, V. (2019). Widespread global peatland establishment and persistence over the last 130,000 y. *Proceedings of the National Academy of Sciences*, 116(11), 4822. doi: 10.1073/pnas.1813305116
- Trunz, V., Packer, L., Vieu, J., Arrigo, N., & Praz, C. J. (2016). Comprehensive phylogeny, biogeography and new classification of the diverse bee tribe Megachilini: Can we use DNA barcodes in phylogenies of large genera? *Molecular Phylogenetics and Evolution*, 103, 245–259. doi: 10.1016/j.ympev.2016.07.004
- van Breemen, N. (1995). How *Sphagnum* bogs down other plants. *Trends in Ecology & Evolution*, 10(7), 270–275. doi: 10.1016/0169-5347(95)90007-1
- Velichko, A. A., Timireva, S. N., Kremenetski, K. V., MacDonald, G. M., & Smith, L. C. (2011). West Siberian Plain as a late glacial desert. *Quaternary International*, 237(1–2), 45–53.
- Wilkinson, D. M. (2001). What is the upper size limit for cosmopolitan distribution in free-living microorganisms? *Journal of Biogeography*, 28(3), 285–291.
- Wilkinson, D. M., Koumoutsaris, S., Mitchell, E. A. D., & Bey, I. (2011). Modelling the effect of size on the aerial dispersal of microorganisms. *Journal of Biogeography*, 39(1), 89–97. doi: 10.1111/j.1365-2699.2011.02569.x

Data Accessibility

The 57 DNA sequences of COI gene of *Hyalophenia papilio* are available in in GenBank with the following accession numbers: MK823130- MK823186.

Author contributions

D.S., E.A.D.M. and E.L. designed the experiments; D.S., E.A.D.M., G.G., H.R., L.B., N.G.K, I.G., L.I.H., K.K., M.L., N.P.K., R.J.P. and K.V., collected the samples; D.S., Q.B., C.D., L.D.F., B.F., C.E.H. and E.L. analysed the data; D.S., E.A.D.M., R.J.P. and E.L. wrote the first version of the manuscript, which was then edited by all co-authors.

Conflict of interest

The material in this manuscript is original research, has not been previously published and has not been submitted for publication elsewhere while under consideration for Molecular Ecology. We would also like that only the online version appears in colour, as we took special care in building our figures in a way that they can be read in black and white as well. We have no conflict of interest in this research.

Figures

Fig. 1) Holarctic distribution of *Hyalosphenia papilio* lineages. Each circle corresponds to a sampling site where the lineage has been detected. Lineage codes correspond to phylogenetic groups, as identified in Heger et al. 2013. I and II present a detailed representation of Beringia area. Inset: Light microscopy image of *Hyalosphenia papilio*. The pyriform outline corresponds to the shell that protects the single-cell body of the organism and its endosymbiotic microalgae (green dots)

Fig. 2) Maximum Likelihood (ML) and Bayesian concatenated phylogenetic tree from 76 unique sequences of *Hyalosphenia papilio* isolated from *Sphagnum* peatlands across the Holarctic realm. Numbers along branches represent, respectively, bootstraps obtained by ML and posterior probabilities as calculated with Bayesian analyses. Trees were rooted internally based on the topology of trees obtained in earlier works (Kosakyan et al. 2016, Heger et al. 2013), which showed two major clades with maximum support; we rooted the tree between these two clades. The tree also represents the different lineages obtained with the ABGD and GMYC analyses and the Kimura 2-parameter test.

Fig. 3) Median joining haplotype network of cytochrome oxidase subunit 1 (COI) gene of *Hyalosphenia papilio* from *Sphagnum* peatlands in the Holarctic realm. Grey boxes and letters represent the different lineages identified in the present study. Colours indicate geographical regions (legend: bottom right inset). Circle sizes are proportional to the number of sequenced single cells of *H. papilio* within each haplotype. Cross lines show the number of mutational steps between haplotypes.

Fig. 4) Biogeographical analysis of *H. papilio* from *Sphagnum* peatlands across the Holarctic realm using BioGeoBEARS. The four biogeographical areas are: Eastern North America (in blue), Western North America (in red), Europe (in purple) and Asia (in green). Pie charts at nodes indicate support for each area. The tips are labelled with present-day distributions. The secondary colours indicate range combinations of the tip ranges.

Fig. 5) Relationship of lineage climatic niche breadth to lineage range size (estimated as the number of locations where a given lineage is present) of *Hyalosphenia papilio*. Climatic niche breadth was estimating from Bioclim variables using the tolerance index of Doledec et al. (2000)