



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/148851/>

Version: Accepted Version

Article:

Hosie, Annmarie, Siddiqi, Najma, Featherstone, Imogen et al. (2019) Inclusion, characteristics and outcomes of people requiring palliative care in studies of non-pharmacological interventions for delirium: A systematic review. *Palliative Medicine*. 269216319853487. pp. 878-899. ISSN: 0269-2163

<https://doi.org/10.1177/0269216319853487>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Palliative Medicine

Inclusion, characteristics and outcomes of people requiring palliative care in studies of non-pharmacological interventions for delirium: a systematic review

Journal:	<i>Palliative Medicine</i>
Manuscript ID	PMJ-18-0287.R2
Manuscript Type:	Review Article
Date Submitted by the Author:	12-Apr-2019
Complete List of Authors:	<p>Hosie, Annmarie; University of Technology Sydney, IMPACCT (Improving Palliative, Aged and Chronic Care through Clinical Research and Translation)</p> <p>Siddiqi, Najma; University of York</p> <p>Featherstone, Imogen; University of York</p> <p>Johnson, Miriam; University of Hull, Wolfson Palliative Care Research Centre</p> <p>Lawlor, Peter; University of Ottawa, Palliative Care; Bruyere Continuing Care, Palliative Care</p> <p>Bush, Shirley; 3. University of Hull</p> <p>Amgarth-Duff, Ingrid; University of Technology Sydney, IMPACCT (Improving Palliative, Aged and Chronic Care through Clinical Research and Translation)</p> <p>Edwards, Layla; University of Technology Sydney, IMPACCT (Improving Palliative, Aged and Chronic Care through Clinical Research and Translation)</p> <p>Cheah, Seong Leang; University of Technology Sydney, IMPACCT (Improving Palliative, Aged and Chronic Care through Clinical Research and Translation)</p> <p>Phillips, Jane; University of Technology, Sydney, IMPACCT (Improving Palliative, Aged and Chronic Care through Clinical Research and Translation)</p> <p>Agar, Meera; University of Technology Sydney, IMPACCT (Improving Palliative, Aged and Chronic Care through Clinical Research and Translation)</p>
Keywords:	clinical trial, delirium, hospice, inpatient, palliative care, review
Abstract:	<p>Background Delirium is common, distressing, serious and under-researched in specialist palliative care settings.</p> <p>Objectives To examine whether people requiring palliative care were included in non-pharmacological delirium intervention studies in inpatient settings, how they were characterised, and what their outcomes were.</p> <p>Design Systematic review (PROSPERO 2017 CRD42017062178).</p> <p>Data sources</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	<p>Systematic search in March 2017 for non-pharmacological delirium intervention studies in adult inpatients. Database search terms were 'delirium', 'hospitalisation', 'inpatient', 'palliative care', 'hospice', 'critical care', 'geriatrics'. Scottish Intercollegiate Guidelines Network methodological checklists guided risk of bias assessment.</p> <p>Results</p> <p>The 29 included studies were conducted between 1994-2015 in diverse settings in 15 countries (9136 participants, mean age 76.5 years [SD 8.1], 56% women). Most studies tested multicomponent interventions (n=26) to prevent delirium (n=19). Three-quarters of the 29 included studies (n=22) excluded various groups of people requiring palliative care; however, inclusion criteria, participant diagnoses, illness severity and mortality indicated their presence in almost all studies (n=26). Of these, 21 studies did not characterise participants requiring palliative care or report their specific outcomes (72%), four reported outcomes for older people with frailty, dementia, cancer and comorbidities, and one was explicitly focused on people receiving palliative care. Study heterogeneity and limitations precluded definitive determination of intervention effectiveness and only allowed interpretations of feasibility for people requiring palliative care. Acceptability outcomes (intervention adverse events and patients' subjective experience) were rarely reported overall.</p> <p>Conclusion</p> <p>Non-pharmacological delirium interventions have frequently excluded and under-characterised people requiring palliative care and infrequently reported their outcomes.</p>



Inclusion, characteristics and outcomes of people requiring palliative care in studies of non-pharmacological interventions for delirium: a systematic review

Authors

Annmarie Hosie,¹ Najma Siddiqi,² Imogen Featherstone,² Miriam Johnson,³ Peter G. Lawlor,³ Shirley H. Bush,³ Ingrid Amgarth-Duff,¹ Layla Edwards,¹ Seong Leang Cheah,¹ Jane Phillips,¹ Meera Agar¹

Institutions

1. University of Technology Sydney, IMPACCT, Bld. 10, Level 3, 235-253 Jones St, Ultimo, Australia
2. University of York, Heslington, YO10 5DD, United Kingdom
3. University of Hull, Hull, HU6 7RX, United Kingdom
4. University of Ottawa, Department of Medicine, 43 Bruyère Street, Ottawa K1N 5C8, Canada

Corresponding Author

Dr Annmarie Hosie
Phone: +61 2 9514 4404
Email: annmarie.hosie@uts.edu.au

Document details

Number of tables: 3
Number of supplementary tables: 2
Number of figures: 5
Number of references: 97
Word count: 5623

Key words

Clinical trial, Delirium, Hospice, Inpatient, Palliative Care, Review

Inclusion, characteristics and outcomes of people requiring palliative care in studies of non-pharmacological interventions for delirium: a systematic review

Abstract

Background

Delirium is common, distressing, serious and under-researched in specialist palliative care settings.

Objectives

To examine whether people requiring palliative care were included in non-pharmacological delirium intervention studies in inpatient settings, how they were characterised, and what their outcomes were.

Design

Systematic review (PROSPERO 2017 CRD42017062178).

Data sources

Systematic search in March 2017 for non-pharmacological delirium intervention studies in adult inpatients. Database search terms were 'delirium', 'hospitalisation', 'inpatient', 'palliative care', 'hospice', 'critical care', 'geriatrics'. Scottish Intercollegiate Guidelines Network methodological checklists guided risk of bias assessment.

Results

The 29 included studies were conducted between 1994-2015 in diverse settings in 15 countries (9136 participants, mean age 76.5 years [SD 8.1], 56% women). Most studies tested multicomponent interventions (n=26) to prevent delirium (n=19). Three-quarters of the 29 included studies (n=22) excluded various groups of people requiring palliative care; however, inclusion criteria, participant diagnoses, illness severity and mortality indicated their presence in almost all studies (n=26). Of these, 21 studies did not characterise

1
2
3
4 participants requiring palliative care or report their specific outcomes (72%), four reported
5
6 outcomes for older people with frailty, dementia, cancer and comorbidities, and one was
7
8 explicitly focused on people receiving palliative care. Study heterogeneity and limitations
9
10 precluded definitive determination of intervention effectiveness and only allowed
11
12 interpretations of feasibility for people requiring palliative care. Acceptability outcomes
13
14 (intervention adverse events and patients' subjective experience) were rarely reported
15
16 overall.

17 18 **Conclusion**

19
20 Non-pharmacological delirium interventions have frequently excluded and under-
21
22 characterised people requiring palliative care and infrequently reported their outcomes.
23

24 25 **Key words**

26
27 Clinical trial, Delirium, Hospice, Inpatient, Palliative Care, Review
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Key statements

What is already known about the topic?

- Delirium is a distressing and serious neurocognitive condition that frequently occurs for patients in palliative care inpatient settings.
- In contrast to other hospital settings, there is limited evidence to guide non-pharmacological intervention to prevent and treat delirium in palliative care inpatient settings.

What this paper adds

- This review found that various groups of people requiring palliative care were excluded from three-quarters of non-pharmacological delirium intervention studies in inpatient settings; despite this, they were present in most studies and their outcomes were reported in five.
- Non-pharmacological delirium interventions appear feasible for people requiring palliative care yet there is no definitive evidence they are effective or acceptable for this inpatient group.

Implications for research

- Phase II and III randomised controlled trials of non-pharmacological interventions to prevent and treat delirium are needed in specialist palliative care settings.
- Adaptations to future trials of non-pharmacological delirium interventions in other inpatient settings are needed to promote representative study populations and allow outcomes for sub-groups of people requiring palliative care to be reported.
- Additional outcomes related to patient and family subjective experience, goals of care and quality of life would enhance the relevance of delirium intervention research in inpatient settings where people are cared for at the end of life.

Introduction

Delirium is a serious complication of medical illness and hospitalisation.¹ The condition is characterised by acute disturbances to attention, awareness and cognition, has multifactorial aetiology, and variously affects memory, language and visuospatial ability, orientation and perception.² Affected persons often experience feelings of fear, humiliation, confusion and disconnection from others.^{3, 4} Family members' experience distress when delirium causes sudden changes in behaviour and decline in the person they love.^{5, 6} Patients who experience an episode of delirium during hospitalisation experience many poorer outcomes, including being more likely to die.^{1, 7, 8}

Delirium most often occurs in people with older age, advanced or severe illness and/or pre-existing cognitive impairment. Hospital-wide, one in five patients have delirium,⁹ with occurrence higher again in intensive, geriatric and palliative care units.^{1, 6} Studies of delirium epidemiology in palliative care inpatient units that screened patients at least daily reported incidence of 33–45% and prevalence of 58–88% in those who died.

Development of delirium guidelines^{1, 10–13} policy¹⁴ and international advocacy¹⁵ indicate growing awareness of the seriousness and prevalence of delirium and importance of evidence based care for hospitalised patients.^{16, 17} There now is sufficient evidence to implement non-pharmacological interventions for delirium in certain hospital settings.^{18, 19} For example, reviews of studies of multicomponent interventions addressing physical and cognitive activity, sleep, hearing, vision and hydration, as in the original Hospital Elder Life Program (HELP) study,²⁰ reported reduction in incident delirium in older hospitalised patients.^{21–23} Reduction in length of hospital stay and improvement in return to independent living were also demonstrated.²²

In contrast, guideline recommendations for non-pharmacological interventions as the first approach to prevent and treat delirium during advanced illness and at the end of life are not evidence based.^{10, 17, 24} A recent scoping review reported the need to generate evidence to inform clinical care in palliative care settings and populations, for non-pharmacological interventions in particular.²⁵ Poorer outcomes with antipsychotics,²⁶ and over-sedation

1
2
3
4 when benzodiazepines were given for agitated delirium,²⁷ in two recent trials in specialist
5 palliative care settings also highlight the need to establish 'drug-free' ways to prevent and
6 relieve the difficulties of delirium at the end of life.
7
8
9

10 In response, the authors established the '*Studies to Understand and Improve Delirium Care*
11 *in Palliative Settings*' international collaborative (SUNRISE) to generate high-quality delirium
12 research in palliative care. We identified the need to inform our future trials in palliative
13 care through a review of studies of non-pharmacological interventions for delirium
14 conducted in a wide range of inpatient settings. This wide review was premised on our
15 clinical experience and knowledge that many hospitalised patients have advanced and/or
16 serious illness, frailty and high comorbidity and consequently much in common with
17 patients in specialist palliative care settings, especially those in intensive care, medical and
18 geriatric units where rates of delirium are similarly high.^{28, 29} Based on this premise, our
19 specific objectives were to examine whether people requiring palliative care were included
20 in non-pharmacological delirium intervention studies in various inpatient settings, how
21 these participants were characterised, and whether the non-pharmacological interventions
22 were effective, feasible and/or acceptable for them.
23
24
25
26
27
28
29
30
31
32
33
34
35

36 **Methods**

37 **Design**

38
39
40 Systematic review of the literature, following the Preferred Reporting Items for Systematic
41 Reviews and Meta-Analyses (PRISMA).³⁰
42
43
44
45

46 **Participants/settings**

47
48 Adults (≥ 18 years) receiving inpatient hospital or hospice care. In this review, we refer to
49 'hospice' as an inpatient facility with the primary function to provide specialist palliative
50 care to people with life-threatening illness, and analogous to a palliative care inpatient unit.
51
52
53

54 **Search strategy**

55
56 In March 2017, we searched MEDLINE, Embase, CINAHL, PsycINFO, Cochrane Library and
57 Web of Science with the following search terms in MEDLINE: Delirium AND Hospitalization
58
59
60

1
2
3
4 OR Inpatient OR Hospice AND Palliative Care OR Critical Care OR Geriatrics. Filters in
5
6 Medline were: 1. Study types: clinical study, clinical trial all, comparative study, controlled
7
8 clinical trial, meta-analysis, multicenter study, pragmatic clinical trial, randomised controlled
9
10 trial, systematic review; 2. Peer reviewed journal and 3. Published from 1980 onward (when
11
12 delirium was first included in the American Psychological Association (APA) Diagnostic and
13
14 Statistical Manual (DSM-III).³¹ We tailored search terms and filters to subsequent databases.
15
16 We examined reference lists of relevant systematic reviews and meta-analyses identified in
17
18 the search for additional eligible studies.

19 20 **Inclusion/exclusion criteria**

21
22 Studies included were primary reports of prospective trials (i.e. studies of an intervention
23
24 with a comparator); with a primary objective to prevent or treat delirium through non-
25
26 pharmacological intervention/s in adult patients in hospital or hospice unit settings; a
27
28 primary outcome of delirium incidence, severity or duration; published in English in a peer-
29
30 reviewed journal.

31
32 Studies excluded were reports of interventions for alcohol withdrawal delirium only;
33
34 systematic reviews and meta-analyses of non-pharmacological delirium intervention/s;
35
36 studies where the primary outcome was not participants' delirium status (e.g. process or
37
38 cost effectiveness outcomes, validation of delirium tools); studies that did not use diagnostic
39
40 criteria or a tool with established psychometric properties to measure delirium; protocols;
41
42 and ongoing studies.

43 44 **Study selection, data extraction and management**

45
46 We imported search results into Endnote X7 software, removed duplicates and exported
47
48 results to Covidence,³² where three reviewers [IAD, LE, AH] independently applied eligibility
49
50 criteria to titles and abstracts. Reviewers compared decisions about inclusion, documented
51
52 reasons for exclusions at full text review and resolved discrepancies through discussion. Two
53
54 reviewers [IAD, LE] extracted data according to the template for intervention description
55
56 and replication (TIDieR) checklist and guide³³ into an Excel V15.28 spreadsheet. The lead
57
58 reviewer [AH] contributed guidance, oversight and independent data checking.
59
60

Risk of bias assessment

Two reviewers [LE and AH] independently assessed each study for selection, performance, detection, attrition, confounding, and reporting biases according to the Scottish Intercollegiate Guidelines Network (SIGN) Methodology Checklists³⁴ for controlled trials and cohort studies. Discrepancies were resolved through discussion.

Outcomes

To identify our sample of interest (i.e. people requiring palliative care), we examined study inclusion and exclusion criteria, participant diagnoses (including severity or staging), and mortality. We assessed eligibility criteria and diagnoses against the Gold Standards Framework Proactive Identification Guidance (GSF PIG), a clinical tool to help identify people likely to need additional supportive (i.e. palliative) care in the last 12 months of life.²⁹ According to the GSF PIG, these people are those with life-threatening conditions, including illnesses that are advanced, progressive, incurable and/or likely to cause acute crises; frailty and co-morbidities; and sudden catastrophic events.

Previous reviews have reported effectiveness outcomes of non-pharmacological delirium interventions for the entire study sample;²¹⁻²³ whereas this review focused on effectiveness, feasibility and acceptability outcomes of our sample of interest. We examined effectiveness according to each study's primary outcome and any sub-group analysis for our sample of interest. We assessed feasibility by examining intervention characteristics, adherence and study modifications, and acceptability through intervention-related adverse effects and patient, family, clinician and volunteer subjective experiences of the interventions.

Synthesis and analysis plan

We generated tables, text and graphs to report study characteristics, participants, interventions and outcomes. Data transformation and descriptive numerical analyses were performed using Excel. We planned to perform subgroup meta-analysis using Review Manager Analyses software³⁵ of intervention effectiveness if we could definitively distinguish our sample of interest and if interventions and comparators and measures were comparable.

Updated search

We updated the search prior to publication in February 2019 and identified two new eligible papers published after the original search date.^{36, 37} There were not incorporated as neither paper altered the conclusions of the review.

Results

The original database search strategy generated 4300 records. After removing 35 duplicates and 4169 records through title and abstract screening, we reviewed 69 full text papers and excluded 48. We included another eight through reference list searching, resulting in 29 papers reporting 29 studies for inclusion (Figure 1).

(Insert Figure 1 PRISMA Flow Diagram here)³⁰

Study characteristics

The 29 studies were conducted between 1994-2015 across 15 countries: six in the US,^{38 39 40 41 42 20} three each in Australia,^{43 44 45} Belgium,⁴⁶⁻⁴⁸ and Canada,⁴⁹⁻⁵¹ two each in the Netherlands^{52, 53} and Sweden,^{54, 55} and one each in Chile,⁵⁶ France,⁵⁷ Ireland,⁵⁸ Italy,⁵⁹ Japan,⁶⁰ Korea,⁶¹ Singapore,⁶² Spain,⁶³ UK,⁶⁴ and the US/Canada⁶⁵ (Table 1).

Study designs were before/after studies (one with an additional concurrent arm⁶²) (n=11),^{38-41, 44, 45, 48, 52, 57, 62, 64} randomised controlled trials (RCTs) (n=10),^{42, 43, 47, 50, 51, 53, 55, 56, 61, 65} non-randomised controlled trials (one with matched participants²⁰) (n=5),^{20, 46, 54, 59, 63} a quasi-experimental study,⁴⁹ a pilot randomised controlled trial,⁶⁰ and a comparative time series study.⁵⁸

Services and settings were medical (n=10),^{20, 41, 43, 50-52, 54, 56, 59, 63} geriatric (n=7),^{39, 44, 55, 57, 59, 62, 63} medical and/or surgical intensive care (n=6),^{38, 40, 47, 58, 61, 64} perioperative hip fracture (n=6),^{42, 45, 46, 48, 55, 65} other perioperative (n=3)^{52, 53, 60} and palliative care and hospice units (n=1),⁴⁹ with eight studies involving more than one service.^{47, 52, 55, 58, 59, 61, 63, 64}

Most studies tested multicomponent interventions (n=26) and aimed to prevent delirium in non-delirious participants (n=19) (Table1). Six studies aimed to prevent and treat delirium^{38, 42, 48, 54, 55, 65} three were treatment only studies.^{50, 51, 62}

(Insert Table 1 Study characteristics here)

Code: **P** Prevention, **T** Treatment, **RCT** Randomised controlled trial, **CTS** Comparative time series, **B/A** Before–after study, **CT** Controlled trial, **ICU** Intensive Care Unit, **NR** Not reported, **MC** Medical Centre, **M** Multicomponent, **PRCT** Pilot RCT, **QE** Quasi-experimental, **S** Single component

^Δ Numbers signify the domains in which the study was assessed as having **low risk of bias**: 1 = Representative study sample; 2 = Concealed allocation; 3 = Random sequence generation; 4 = Blinded participants and intervention personnel; 5 = Blinded outcome assessors; 6 = Valid, reliable delirium measures; 7 = <20% sample lost to analysis; 8 = Intention to treat analysis; 9 = Confounders accounted for; 10 = Only *a priori* outcomes reported

Risk of bias assessment

Almost all studies were assessed as having at least one form of selection bias (n=28). The exception was a Chilean RCT of a prevention intervention where family members delivered cognition, vision, and hearing strategies to patients.⁵⁶ Eight RCTs minimised internal selection bias through concealed allocation and random sequence generation.^{42, 43, 47, 50, 53, 55, 56, 65} There was high risk of performance bias in all studies, due to the inherent difficulty of blinding non-pharmacological interventions: one RCT blinded participants to their allocation,⁴³ and no studies blinded those performing the intervention. Detection bias was minimised in eight studies through blinded outcome assessors combined with valid, reliable delirium measures.^{42, 43, 47, 50, 51, 53-55} Although we excluded studies that did not use delirium diagnostic criteria and/or a tool with established psychometric properties, three studies were assigned uncertain risk of detection bias due to retrospective assignment of delirium status when the Confusion Assessment Method (CAM) was not performed during weekends in two before-after studies;^{41, 45} and when delirium incidence was reported as a mean rather than a proportion in a comparative time series study.⁵⁸ High risk of detection bias was assigned to a delirium prevention study in seven Canadian palliative care units due use of a screening tool, the Confusion Rating Scale (CRS) to measure delirium incidence, severity and duration.⁴⁹ Six studies had low risk of attrition bias.^{42, 43, 50, 55, 56, 63} Confounders were accounted for in 17 studies.^{20, 39, 40, 42, 46, 47, 49-52, 54, 55, 58, 59, 61, 63, 65} Most studies (n=22) reported only *a priori* outcomes,^{20, 38-43, 46, 48, 50-53, 55-57, 60-65} which we assessed by information provided within the papers rather than through examination of trial registrations or published protocols, which no studies cited.

Figure 2 presents overall risk of bias in the included studies.

(Insert Figure 2 Overall risk of bias graph here)

Figure 2: Overall risk of bias

Participant characteristics

There were 9136 participants (n=4553 in intervention arms) across the 29 studies (Table 2). Sample sizes ranged from 15-1516 participants. Participants' overall mean age (excluding two studies that reported only age range or median)^{48, 58} was 76.5 years (SD \pm 8.1). There were more female participants (56%) than male.

Inclusion and exclusion criteria

Inclusion

Most studies (n=19) included only patients aged 65 years and older (Table 2). Twenty-two studies included consecutively admitted patients, 16 of which further required certain physical conditions to be present (e.g. frailty,⁵³ hip fracture,^{42, 45, 46, 48, 55, 65} anemia,⁶⁵ dementia,⁵⁷ advanced cancer,⁴⁹ oesophageal cancer surgery⁶⁰). Five studies included only patients with risk factors for delirium^{20, 44, 52, 56, 63} and three included only patients with delirium.^{50, 51, 62} The study in seven Canadian palliative care units (hereafter termed the 'palliative care unit study') included only patients with terminal cancer, defined as dying within 90 days of admission which was retrospectively identified. This study had the largest sample of all the studies (n=1516, 17%).⁴⁹

Exclusion

Twenty-three studies excluded patients on various diagnostic grounds (Table 2). These were stroke or cerebrovascular accident (CVA), moderate-severe traumatic brain injury or coma (n=9);^{20, 38, 43, 48, 50, 51, 59, 62, 63} dementia or prior cognitive impairment (n=7);^{20, 38, 44, 47, 61, 63, 64} psychiatric disorders or alcohol/drug addiction (n=6);^{43, 44, 59-61, 64} pathological fracture (n=4);^{46, 48, 55, 65} poly- or multiple trauma (n=2);^{46, 48} neurologic diagnosis/neurosurgical (n=2);^{61, 64} metastatic cancer or referred to oncology services (n=2);^{42, 50} and others (myocardial infarction,⁶⁵ severe renal failure,⁵⁵ brain concussion,⁴⁸ history of sleep pathology,⁶⁴ severe rheumatoid arthritis⁵⁵ and severe hip osteoarthritis⁵⁵).

1
2
3
4 Fifteen studies excluded patients with impaired verbal communication or who were
5 otherwise unable to complete study assessments.^{20, 38, 39, 41, 43, 46, 48, 50, 52, 53, 59, 61-63, 65} Six
6 studies excluded patients with hearing and/or visual impairment.^{46-48, 60, 61, 64} Three studies
7 excluded patients isolated for infection.^{43, 61, 62}

8
9
10
11
12 Eight studies explicitly excluded patients expected to die, using the following terms:
13 “terminal diagnosis” or “terminal illness” (n=3),^{20, 58, 62} “terminal condition and receiving
14 comfort care” (on the basis that they were unlikely to benefit from the intervention)
15 (n=2),^{41, 44} “life expectancy less than six months”,⁴⁶ “metastatic cancer or comorbid illnesses
16 likely to reduce life expectancy to less than six months”,⁴² “death expected within 24
17 hours”,⁴³ and “dangerously ill”.⁶²

18
19
20
21
22
23 Two studies excluded participants who subsequently died from analysis;^{51, 52} a delirium
24 prevention study in elderly care wards excluded 70% of 2162 admitted patients due to being
25 “too unwell”;³⁹ and another study excluded patients with “deterioration of condition”.⁶⁰

26
27
28
29
30 The palliative care unit study excluded patients who lived for more than 90 days or who
31 were still alive at discharge from analysis.⁴⁹

32 33 34 **Characterisation of study participants**

35 36 ***Participant diagnoses and illness severity***

37
38 Reporting of participant diagnoses, illness severity and mortality varied. Fifteen studies
39 reported co-morbid diagnoses, twelve reported only primary diagnoses, and four reported
40 both (Table 2). Diagnoses’ categorisation varied. Studies rarely reported staging of
41 diagnoses (n=2).^{49, 56}

42
43
44
45
46
47 Sixteen studies reported illness severity for all participants using at least one measure, more
48 often as a mean or median score (n=15)^{20, 38, 43, 46, 47, 50, 52, 55, 56, 59, 61-65} than by categorising
49 participants proportionally (n=6).^{20, 39, 42, 47, 52, 62} The most frequently used illness severity
50 measures were the Charlson Comorbidity Index (CCI)⁶⁶ (n=7; reported as: mean scores 2.3 -
51 3.3,^{46, 50, 62} median 2,^{43, 56} and scores of four or greater 19.8%³⁹ and 39%⁴²); and the Acute
52 Physiology And Chronic Health Evaluation II (APACHE II)⁶⁷ (n=6; mean scores 11.3-15.6).^{20, 43,}
53
54
55
56
57
58
59
60 Two ICU studies that measured illness severity using the APACHE II reported

1
2
3
4 means scores (13.9-14.6)^{61, 64} approximately equivalent to or less than those reported in
5 three of four medical and geriatric unit studies (14.1-15.6).^{20, 43, 59} Other measures were the
6 Cumulative Illness Rating Scale–Geriatrics (CIRS-G) comorbidity and severity indexes,^{52, 59, 68}
7 American Society of Anesthesiologists physical status classification (ASA)^{52, 65, 69}; Blessed
8 Dementia Rating Scale;^{20, 42, 70} Modified Early Warning Score (MEWS);^{39, 71} Sepsis-related
9 Organ Failure (SOFA);^{47, 72} Risk, Injury, Failure, Loss of kidney function, and End-stage kidney
10 disease (RIFLE);^{47, 73} Simplified Acute Physiology Score (SAPS 3);^{47, 74} and the 15-item
11 Geriatric Depression Scale (GDS-15).^{55, 75}

12
13 Three studies reporting comorbidities without the CCI reported: i. 60% had more than two
14 comorbidities (diabetes, COPD, hypertension, myocardial infarction, other cardiovascular
15 disorders, neurological disorders, cerebrovascular disorders, hearing and vision problems,
16 memory problems in daily life, psychiatric disorders or musculoskeletal disorders);⁵³ ii. 72%
17 had at least one co-morbid illness;⁴⁷ and iii. a mean of 2.7 comorbidities.⁶³

18
19 Almost every diagnosis, sensory deficit and impairment that were exclusion criteria in some
20 studies were reported in the overall study population (Table 2). For example, more studies
21 reported participants with dementia or cognitive impairment (n=13)^{39, 41-43, 46, 48-51, 55-57, 65}
22 than excluded people with these conditions (n=7).^{20, 38, 44, 47, 61, 63, 64}

23 24 25 **Participant mortality**

26
27 Mortality was reported in 19 studies at nine different time points, ranging from ‘in the
28 ICU’^{38, 40} to one year post intervention.^{46, 55, 62} In these studies, 2090 participants died
29 (23%). Mortality rates ranged from 1% in three studies^{20, 47, 61} to 100% in the palliative care
30 unit study.⁴⁹ There were 596 participant deaths in 18 studies reporting mortality that were
31 conducted outside of a specialist palliative care setting (12%). High mortality was reported
32 for intervention and control medical unit participants at eight weeks (22.1% vs 19.37%;⁵⁰
33 33% vs 37% for delirious patients⁵¹) and at six months (33.8% vs 30.9%);³⁹ and at one year
34 for older (≥65) traumatic hip fracture patients (33% vs 22%).⁴⁶ Six of the 19 studies
35 measuring mortality excluded patients expected to die; despite this exclusion criteria, 134
36 participants of these six studies subsequently died during the study period (12%).^{20, 44, 46, 52,}
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

of which reported significantly less mortality in intervention cohorts compared to controls,^{54, 61} with one (at day 7 in hospital) not sustained at 30 days in hospital.⁶¹

Study approaches to people requiring palliative care

By examination of these data we identified only one study that explicitly reported outcomes for participants requiring palliative care (n=1).⁴⁹ Using the GSF PIG, we identified another four studies that reported primary or sub-group effectiveness outcomes for older people likely to require palliative care due to the presence of frailty, dementia, cancer and comorbidity.^{42, 52, 53, 57} All five studies were of interventions to prevent delirium. Twenty-two studies either explicitly or resultantly excluded groups of people requiring palliative care (76%).^{20, 38, 39, 41-44, 46-48, 50-52, 55, 58-65} Yet, through our interpretation of reported diagnoses, levels of comorbidity, frailty and mortality (signified with a † in Table 2), we also identified that people requiring palliative care were present in 21 studies that did not specifically characterise them or report their outcomes (72%).^{20, 38-41, 43, 45-48, 50, 51, 54-56, 58, 61-63, 65} Figure 3 presents these findings diagrammatically.

(Insert Table 2: Study inclusion and exclusion criteria and participant diagnoses, illness severity and mortality)

* Statistically significant difference ^Δ Intervention and control participants combined ^α Rounded to nearest whole number [†] Interpreted as indicating need for palliative care

Illness severity measures: Higher scores represent higher illness severity. **AIS** Abbreviated Injury Scale, **APACHE II** Acute Physiology and Chronic Health Evaluation (scores 0-71), **ASA** American Society of Anesthesiologists physical status classification (I: normal healthy patient - VI: a declared brain-dead patient whose organs are being removed for donor purposes), **BDRS** Blessed Dementia Rating (scores 0-28, cut-off for impairment > 4), **CCI** Charlson Comorbidity Index (scores 0-37), **CIRS-G** Cumulative Illness Rating Scale–Geriatrics (scores 0-56), **CSI** Clinical severity of illness (1 (mild) - 9 (moribund)), **GCS** Glasgow Coma Scale (scores 3-15; scores 3-8 = coma), **GDS-15** 15-item Geriatric Depression Scale (≥6 = suggests depression and need for assessment, ≥11 = depression/severe depression), **ISS** Injury Severity Score (scores 1-75), **MEWS** Modified Early Warning Score (score ≥5 is statistically linked to increased likelihood of death or admission to an intensive care unit), **RIFLE** (Risk, Injury, Failure, Loss of kidney function, and End-stage kidney disease categories), **SAPS 3** Simplified acute physiology score (scores 0-217), **SOFA** Sepsis-related Organ Failure Assessment (scores 0 to 24). **Other abbreviations:** **ADL** activities of daily living, **AF** atrial fibrillation, **Ca** cancer, **CAM** Confusion Assessment Method, **CAM-ICU** Confusion Assessment Method for the Intensive Care Unit, **CI** cognitive impairment, **CKD** Chronic Kidney Disease, **COPD** chronic obstructive pulmonary disease, **CVD** cardiovascular disease, **CVA** cerebrovascular accident, **DM** diabetes mellitus, **ED** Emergency Department, **GI** gastrointestinal, **HF** heart failure, **hrs** hours, **HT** hypertension, **I/C** intervention/control, **ICU** intensive care unit, **IHD** ischaemic heart disease, **IQR** interquartile range, **MI** Myocardial Infarction, **MMSE** Mini-Mental State Examination, **NR** not reported, **O/A** on admission, **OP** osteoporosis, **PVD** peripheral vascular disease, **RASS** Richard Agitation Sedation Scale, **RF** renal failure, **SD** standard deviation, **SICU** surgical intensive care unit, **TIA** transient ischemic attack, **UTI** urinary tract infection.

Figure 3: Study approaches to people requiring palliative care

NB: Combined percentages do not add up to 100% as studies simultaneously excluded and reported people requiring palliative care.

Effectiveness outcomes for people requiring palliative care

Overall, 20 studies reported that the intervention was effective according to at least one primary outcome (69%)^{20, 38-42, 44-46, 51, 54-57, 59-64}

The one study explicitly focused on preventing delirium in people receiving palliative care (patients with terminal cancer) tested three primarily nurse-delivered intervention components for orientation, informing family, and assessment of medication risk factors plus querying physicians about changes to medication and found no statistically significant difference in delirium incidence ($p=0.66$, OR 0.94), severity or duration between intervention and control sites.⁴⁹

Outcomes of four participant groups that we identified as requiring palliative care were as follows:

A delirium prevention RCT of a geriatric liaison intervention of comprehensive assessment, cognitive and physical activity, hearing, vision and nutrition for frail elderly cancer patients undergoing surgery for a solid tumour reported no significant difference between delirium incidence in intervention and control groups (9.4% vs. 14.3%, OR: 0.63, 95% CI: 0.29–1.35).⁵³

A delirium prevention RCT of geriatric consultation and multiple care components (Table 3) performed analysis for the sub-group of people with dementia within a perioperative hip fracture population, reporting a non-statistically significant reduction in delirium incidence in the intervention group compared with control ($n=13$ (62%) vs. $n=20$ (69%) $p=0.6$).⁴²

A delirium prevention before-after study of orientation and communication strategies for patients with dementia in a French acute geriatric unit reported a 66% relative risk reduction of delirium in the after cohort (RRR 0.34 IC 95% 0.15–0.78).⁵⁷

A delirium prevention before-after study of delirium/cognitive screening, comprehensive assessment, cognitive and physical activity, nutrition, falls prevention, medication review and staff education in frail elderly surgical patients found no statistically significant

1
2
3
4 difference in incidence of delirium (11% vs 10% $p=0.945$) or cognitive decline (15% vs 12%
5
6 $p=0.431$) in the before group compared with the after.⁵²
7

8 Due to heterogeneity of study interventions and measures (Table 2), we did not conduct
9
10 meta-analysis of these effectiveness outcomes.
11
12

13 **Types of outcomes measured in the included studies**

14
15 Twenty-nine different outcomes were measured overall (Figure 4). There were nine
16
17 different primary outcomes, the most frequent was delirium incidence ($n=21$).^{20, 38, 39, 41-46,}
18
19 ^{48, 49, 52, 53, 56-61, 63, 64} Of the 22 different secondary outcomes, the most often reported was
20
21 length of ICU or hospital stay ($n=16$).^{38-43, 45, 46, 48, 50, 51, 53-56, 62}
22

23 **Figure 4: Types and rates of outcomes measured**

24 **Feasibility outcomes**

25 **Intervention components**

26
27 The interventions consisted of 24 components overall, varying in type and number per study
28
29 (1-15) ($M=6.3$ $SD \pm 4$) (Table 3). The type and number of strategies in each component also
30
31 varied (data not reported here).
32
33

34
35 Eighteen multicomponent *prevention* interventions most often included strategies for:
36
37 cognitive activity ($n=15$);^{20, 39, 41, 43-45, 49, 52, 53, 56, 57, 59, 61, 63, 64} physical activity ($n=9$);^{20, 39, 43, 45, 52,}
38
39 ^{53, 59, 63, 64} hearing ($n=9$);^{20, 39, 41, 44, 45, 53, 56, 61, 63} vision ($n=9$);^{20, 39, 41, 44, 45, 53, 56, 61, 63} sleep-wake
40
41 cycle preservation ($n=8$);^{20, 40, 41, 45, 59, 61, 63, 64} nutrition ($n=8$);^{41, 44, 45, 52, 53, 59, 61, 63} staff
42
43 education;^{39-41, 52, 59, 61, 63, 64} and hydration ($n=7$).^{20, 41, 44, 45, 59, 61, 63} These components largely
44
45 correspond to those of the original HELP study.²⁰
46
47

48 Components of six delirium *prevention and treatment* interventions were more diverse
49
50 ($M=8$), with staff education ($n=4$),^{38, 48, 54, 55} staff changes ($n=3$)^{48, 54, 55} and strategies for pain
51
52 ($n=3$) most often included.^{42, 48, 55}
53

54 The most common components of three delirium *treatment* interventions were cognitive
55
56 activity ($n=3$);^{50, 51, 62} physical activity ($n=3$);^{50, 51, 62} hearing ($n=3$);^{50, 51, 62} vision ($n=3$);^{50, 51, 62}
57
58
59
60

1
2
3
4 sleep-wake cycle preservation (n=2);^{50, 62} and environmental changes (e.g. adjustments to
5
6 lighting and noise) (n=2).^{51, 62}
7

8 Overall, multicomponent interventions also often included: family involvement (e.g.
9 information/education about delirium, increased family presence) (n=9);^{38, 45, 49-51, 56, 58, 59, 61}
10 environmental strategies (n=9);^{38, 40, 42, 45, 50, 51, 61, 62, 64} and comprehensive assessment
11 (n=8).^{42, 46, 50-53, 55, 61} Pharmacological strategies (e.g. medication reviews and alerts,
12 protocols for pain and sedation strategies) were present in half of the multicomponent
13 interventions (n=12).^{38, 40-42, 45, 48, 49, 52, 55, 59, 61, 64}
14
15
16
17
18
19

20 The three *single component* interventions were blood transfusion,⁶⁵ bright light therapy⁶⁰
21 and earplugs at night.⁴⁷
22
23

24 The five studies that reported effectiveness outcomes for our sample of interest addressed:
25 cognitive activity,^{42, 49, 52, 53, 57} physical activity,^{42, 52, 53} comprehensive assessment,^{42, 52, 53}
26 hearing,^{42, 53} vision,^{42, 53} nutrition,^{42, 52, 53} pharmacological strategies,^{42, 49, 52} environmental
27 strategies,^{42, 57} family involvement,⁴⁹ staff changes,⁴² pain,⁴² medical complications,⁴²
28 oxygen,⁴² staff education,⁵² falls prevention⁵² delirium/cognitive screening⁵² and
29 bladder/bowel function⁴² (Figure 5).
30
31
32
33
34
35

36 **Intervention delivery**

37 Specialist geriatric clinicians or teams led almost half of the interventions, as either
38 consultants or directly (n=12).^{20, 39, 42, 45, 46, 50-53, 55, 62, 63} Interventions were delivered by
39 interdisciplinary teams (n=5),^{20, 38, 45, 46, 55} physicians and nurses (n=3),⁴⁹⁻⁵¹ nurses alone
40 (n=3),^{47, 48, 61} physician, nurse and physiotherapists (n=2),^{59, 64} family members (n=2),^{56, 58}
41 and physiotherapists and allied health assistants (n=1).⁴³ Volunteers delivered part or all of
42 the intervention in four studies.^{20, 41, 44, 52} Three studies did not report who delivered the
43 intervention, including two single component interventions.^{57, 60, 65} Most studies (n=25)
44 wholly or partly tailored components and strategies to patients' individual needs (Table 3).
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

(Insert Table 3 Intervention characteristics here)

Abbreviations: **ICU** Intensive Care Unit, **NR** Not reported, **OA** On admission, **OT** Occupational therapist, **PT** physiotherapist **Component codes:** **C** Cognitive activity, **E** Physical activity, **H** Hearing, **V** Vision, **P** Sleep-wake cycle preservation, **W** Hydration, **S** Staff education, **F** Family involvement, **N** Nutrition, **Pa** Pain, **O** Oxygen, **L** Falls prevention, **G** Staff changes, **B** Bladder/bowel, **J** Environment/lighting/noise, **CA** Comprehensive assessment, **BT** Blood transfusion, **Z** Address medical complications, **PE** Patient education, **Ph** Physiological monitoring, **K** Pharmacological

Figure 5: Types and rates of intervention components, including for sample of interest

Code: **P** Delirium prevention studies, **P & T** Combined delirium prevention and treatment studies, **T** Delirium treatment studies, **S of I** Sample of interest

Adherence

Eighteen studies reported adherence to intervention strategies, using different methods and levels of detail (Supplementary Table 1). Almost all reported less than 100% adherence, with the exception reporting 100% geriatric nurse compliance with the “semi-structured protocol”.⁶² The palliative care unit study reported 89.7% overall adherence to the study protocol.⁴⁹ Three studies compared adherence to intervention strategies with that for control participants and reported that usual care sometimes mirrored some intervention strategies.^{46, 50, 55} Two studies reported reasons for non-adherence: pharmacological sedation, coma and absence of a relative in the palliative care unit study;⁴⁹ and patient refusal or unavailability, staff member unavailability, and medical contraindications in the original HELP study.²⁰

Study modifications

Two studies modified the intervention in the pilot phase, including changes in staff education and practice change materials,³⁹ and providing study information to family on admission rather than waiting until the researcher was present.⁵⁸ The palliative care unit study modified the primary outcome measure by not substituting the CRS for the CAM, due to CAM completion delays and poor completion rates (39% of participants) caused by “the challenge of conducting the baseline CAM structured interview in the last days or hours of life”.⁴⁹

Acceptability outcomes

Adverse effects

No study reported systematic processes of attribution of adverse effects to the intervention. Ten studies reported adverse events related to hospitalisation rather than the intervention^{38, 44, 51-53, 55, 56, 62, 63} (Figure 4 and Supplementary Table 2). No study reported statistically significant increased adverse events for intervention participants. Two simply stated that the intervention had no adverse effects.^{20, 43} A RCT,⁵⁵ a controlled trial⁶³ and a before-after study⁶² reported statistically significant reductions in some adverse events for intervention participants: falls,⁵⁵ physical restraint,^{62, 63} pressure ulcers,^{55, 62} infection,^{55, 62} and sleeping problems.⁵⁵

Patient, family, clinician and volunteer subjective experiences

No study reported patients' subjective experiences of intervention (Supplementary Table 3). A survey of family caregiver's satisfaction with intervention and care for delirious patients within one study found most experienced moderate to high levels of distress (72%), believed that the environment and intervention helped the patient's recovery (86.6% and 83.5%, respectively) and that the three most useful strategies were activity, reality orientation and thrice-daily mobilisation.⁶² A before-after study of older medical and surgical inpatients interviewed patients, volunteers, nurses and physicians and analysed meeting minutes to assess intervention integration in clinical practice, with minimal details of analysis methods and findings provided.⁵² Four studies reported multidisciplinary and expert input during intervention development.^{39, 40, 45, 61} The RCT of nocte earplugs in the ICU stated in the discussion that some participants preferred not to use them in order to remain "in direct contact with their environment".⁴⁷ No studies reported measures of patient distress related to delirium (Figure 5).

Discussion

This review found that studies of non-pharmacological delirium interventions frequently excluded and under-characterised people requiring palliative care and subsequently their outcomes were infrequently reported. The likely reasons for these findings and how to address barriers to inclusion and report characteristics and outcomes for this patient subgroup in future research are discussed.

With regards to inclusion, this review identified a selection bias against people requiring palliative care through exclusion of people expected to die (using various prognoses and terminology) and also of those with greater acuity or severity of illness, particular diagnoses and with cognitive, sensory and/or communication impairments. These exclusions were rarely explained or justified and often seemingly arbitrary. Nor were they consistent across the studies. These exclusions represented people highly at risk of delirium.^{1, 6} This finding reflects those of other reviews that identified selection biases against people with dementia in geriatric research⁷⁶ and older people from clinical trials in oncology.⁷⁷ Reasons for the exclusions identified in our review are likely to be multifactorial. One factor may be assumptions that the interventions were not appropriate, possible or likely to be effective for people requiring palliative care, as indicated by the two studies which justified the exclusion of patients with a terminal condition and requiring comfort care as that the intervention would be unlikely to benefit them.^{41, 44} Historical views that people requiring palliative care are separate from the wider hospital population, rather than universally and always within it, may also have influenced some exclusion decisions.^{78, 79} There are also pragmatic challenges to ensuring informed consent and completion of study measures when patients are frail, expected to decline, and have pre-existing cognitive and sensory impairments. However, these challenges may be overcome through a variety of research strategies, such as proxy, opt out and advance consent methods, early contact, tailoring of interventions and messaging to patients and family, adequate research resources and governance, and cluster designs.^{26, 27, 80, 81} Outcomes such as days alive without delirium or coma may promote the inclusion of more severely ill patients in future delirium research, because consciousness is pre-requisite for delirium measurement.^{40, 82} Brief and

1
2
3
4 observational delirium diagnostic tools that can be validly used with patients with cognitive,
5 sensory and communication impairment are also available.⁸³⁻⁸⁶
6
7

8
9 Despite most studies excluding certain groups of people likely to require palliative care,
10 unsurprisingly we identified that they were indeed present, despite under-reporting of their
11 characteristics and outcomes. One reason it was difficult to retrospectively distinguish this
12 sub-group for the purposes of ascertaining their outcomes in this review was that the
13 studies reported participants' diagnoses, illness severity, and mortality using different
14 measures, time-points, methods and degrees of detail. While identifying inpatients with
15 palliative care needs in hospital is challenging and an uncertain science, both retrospectively
16 and prospectively, there are ways it can be achieved.^{29, 87-89} In addition to the GSF PIG which
17 we applied in this review,²⁹ the Palliative Care Needs Assessment Tool⁹⁰ and the Supportive
18 and Palliative Care Indicators Tool (SPICT)⁹¹ can be used to identify which patients are at
19 risk of deteriorating and dying in hospital. Other studies suggest identifying patients with
20 palliative care needs through the presence of life-threatening illness coupled with receipt
21 and acceptance of care focused on supporting quality of life,⁸⁷ or retrospectively through the
22 use of death registration data.⁸⁹ Such methods could be used in delirium intervention
23 research in settings where mortality is high, such as intensive care units, where existing
24 illness severity measures focus on acuity and intensity of intervention and, despite having
25 some prognostic utility, were never designed to identify a patient's need for palliative
26 care.^{67, 74, 92}
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

42 Better characterisation of people requiring palliative care in studies of non-pharmacological
43 interventions for delirium would help to build evidence of their outcomes in two ways.

44
45
46 Firstly, it would allow sub-group analysis to be performed and reported in future studies in
47 any setting. This is important because the question of whether delirium can be prevented
48 and treated through non-pharmacological means was not definitively determined for this
49 population by this review. The multi-site palliative care unit study had the greatest number
50 of participants and reported good overall adherence to the study protocol;⁴⁹ however, the
51 only core care domain²¹ was cognitive activity, consisting of nurses orientating patients to
52 time, person and place each shift. Planned use of the CAM was not accomplished and the
53
54
55
56
57
58
59
60

1
2
3
4 substituted screening tool⁹³ was not a validated tool for delirium diagnosis or severity. All
5 participants had 'terminal' cancer and died within 90 days of admission and thereby were
6 not fully representative of people requiring inpatient palliative care, including in specialist
7 units⁴⁹, not all of whom will die within three months. Only one geriatric unit study of people
8 with dementia reported reduced delirium incidence; however, the report lacked detail,
9 including of analysis methods.⁵⁷ The remaining three studies were focused on frail elderly
10 patients with cancer, dementia, hip fracture and comorbidity following surgery,^{42, 52, 53} a
11 level of intervention and delirium risk that many people requiring palliative care would not
12 undergo. Lastly, we found no studies focused on non-pharmacological interventions to treat
13 delirium in people requiring palliative care, highlighting the particular need to research this
14 challenging area of clinical care.

15
16
17
18
19
20
21
22
23
24
25 Secondly, better recognition of people requiring palliative care in delirium research would
26 heighten awareness of their needs, which in turn would instigate consideration of outcomes
27 that are most meaningful to them. Reducing delirium incidence, duration and severity,
28 length of admission and mortality were the foci of the included studies. While worthwhile
29 aims at any point in the illness trajectory, more person-centred outcomes, such as reduction
30 of delirium related distress, and improved patient and family caregiver experiences of
31 decision making, respect and dignity, and quality of life, were almost completely absent.^{5, 94}
32 Measuring these additional outcomes, which are highly valued by patients and family
33 caregivers,^{95, 96} would enrich this field of research and is achievable through both
34 quantitative and qualitative methods. Adverse effects of intervention will also be important
35 to systematically measure in future trials, given that patients with cognitive impairment
36 have increased vulnerability to harm in hospital.⁹⁷

37
38
39
40
41
42
43
44
45
46
47
48 Lastly, this review allowed an interpretation that many intervention components are
49 feasible for people requiring palliative care by virtue of their delivery to elderly, frail and/or
50 critically ill patients with and without delirium in the included studies. However, we suggest
51 that studies measuring the feasibility of components of the original HELP intervention²⁰ that
52 were subsequently found to be effective for other older hospitalised patients i.e. those
53 targeting cognitive and physical activity, sleep, hearing, vision and hydration,²¹⁻²³ are
54 necessary next steps within delirium research programs in specialist palliative care settings.

Limitations and strengths

A limitation of this review was that only English language studies were included. Another limitation was that our retrospective identification of people requiring palliative care was not sufficient to definitively determine outcomes for this patient sub-group; therefore this review makes only recommendations for future research, not clinical practice. A strength of this review was our systematic approach adhering to PRISMA.³⁰

Conclusion

This review found that studies of non-pharmacological delirium interventions have excluded, poorly characterised and infrequently reported outcomes for people requiring palliative care. Based on these findings, we suggest new approaches to generate evidence for delirium interventions in this important patient sub-group. First, randomised controlled trials of non-pharmacological interventions to prevent and treat delirium are needed in specialist palliative care settings, with feasibility studies required before trials of effectiveness. Interventions should be based on those implemented and proven effective for older patients, especially those who were frail and severely ill with comorbidity. Second, broad inclusion criteria, justified exclusions, and tailoring in future inpatient trials of non-pharmacological delirium interventions would promote representative study populations. Third, systematic characterisation of the sub-groups of people requiring palliative care would allow their specific outcomes to be reported. Last, additional outcomes related to patient and family subjective experience, goals of care and quality of life would enhance the relevance of delirium research in inpatient settings where people are cared for at the end of life.

Acknowledgements

We thank Dr E. Wesley Ely for his insights in ICU illness severity measures and constructive feedback on the initial manuscript, and a University of Technology Sydney Health Futures Research Development Fund grant which supported this research.

References

1. National Institute for Health and Clinical Excellence (NICE) National Clinical Guideline centre. Delirium: diagnosis, prevention and management, <https://www.nice.org.uk/guidance/cg103> (2010, accessed March 2 2019).
2. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5)*. Arlington, VA: American Psychiatric Publisher, 2013.
3. O'Malley G, Leonard M, Meagher D, et al. The delirium experience: a review. *Journal of Psychosomatic Research* 2008; 65: 223-228. 2008/08/19. DOI: 10.1016/j.jpsychores.2008.05.017.
4. Van Rompaey B, Van Hoof A, van Bogaert P, et al. The patient's perception of a delirium: A qualitative research in a Belgian intensive care unit. *Intensive and Critical Care Nursing* 2016; 32: 66-74. DOI: 10.1016/j.iccn.2015.03.002.
5. Finucane AM, Lugton J, Kennedy C, et al. The experiences of caregivers of patients with delirium, and their role in its management in palliative care settings: an integrative literature review. *Psycho-Oncology* 2017; 26: 291-300. DOI: 10.1002/pon.4140.
6. Hosie A, Davidson PM, Agar M, et al. Delirium prevalence, incidence, and implications for screening in specialist palliative care inpatient settings: A systematic review. *Palliative Medicine* 2013; 27: 486-498. DOI: 10.1177/0269216312457214.
7. Australian Commission on Safety and Quality in Health Care. Evidence for the safety and care of patients with a cognitive impairment in acute care settings: a rapid review. Sydney: ACSQHC, 2013.
8. Leslie DL, Marcantonio ER, Zhang Y, et al. One-year health care costs associated with delirium in the elderly population. *Archives Internal Medicine* 2008; 168: 27-32. 2008/01/16. DOI: 10.1001/archinternmed.2007.4.
9. Ryan DJ, O'Regan NA, Caoimh RÓ, et al. Delirium in an adult acute hospital population: predictors, prevalence and detection. *BMJ Open* 2013; 3. DOI: 10.1136/bmjopen-2012-001772.
10. Canadian Coalition for Seniors' Mental Health. Guideline on the Assessment and Treatment of Delirium in Older Adults at the End of Life. Toronto: Canadian Coalition for Seniors' Mental Health (CCSMH), 2010.
11. Australian Commission on Safety and Quality in Health Care. A better way to care: Safe and high-quality care for patients with cognitive impairment (dementia and delirium) in hospital - Actions for consumers. Sydney: ACSQHC, 2014.
12. Barr J, Fraser GL, Puntillo K, et al. Clinical practice guidelines for the management of pain, agitation, and delirium in adult patients in the intensive care unit. *Critical Care Medicine* 2013; 41: 263-306. DOI: 10.1097/CCM.0b013e3182783b72.
13. Bush SH, Marchington KL, Agar M, et al. Quality of clinical practice guidelines in delirium: a systematic appraisal. *BMJ Open* 2017; 7. DOI: 10.1136/bmjopen-2016-013809.
14. Australian Commission on Quality and Safety in Healthcare. Delirium Clinical Care Standard, <https://www.safetyandquality.gov.au/our-work/clinical-care-standards/delirium-clinical-care-standard/> (2017, accessed March 9 2019).
15. iDelirium, <http://www.idelirium.org/> (2018, accessed March 9 2019).
16. Lawlor PG, Davis DHJ, Ansari M, et al. An Analytic Framework for Delirium Research in Palliative Care Settings: Integrated Epidemiological, Clinician-Researcher and Knowledge User Perspectives. *Journal of Pain and Symptom Management* 2014; 48: 159-175. DOI: 10.1016/j.jpainsymman.2013.12.245.
17. Palliative Care Expert Group. *Therapeutic Guidelines: Palliative Care (Version 4)*. Melbourne: Therapeutic Guidelines Ltd, 2016.

18. Abraha I, Rimland JM, Trotta F, et al. Non-Pharmacological Interventions to Prevent or Treat Delirium in Older Patients: Clinical Practice Recommendations The SENATOR-ONTOP Series. *Journal of Nutrition, Health and Aging* 2016; 20: 927-936. 2016/10/30. DOI: 10.1007/s12603-016-0719-9.
19. Devlin JW, Skrobik Y, Gélinas C, et al. Clinical Practice Guidelines for the Prevention and Management of Pain, Agitation/Sedation, Delirium, Immobility, and Sleep Disruption in Adult Patients in the ICU. *Critical Care Medicine* 2018; 46: e825-e873. DOI: 10.1097/ccm.0000000000003299.
20. Inouye SK, Bogardus STJ, Charpentier PA, et al. A Multicomponent Intervention to Prevent Delirium in Hospitalized Older Patients. *New England Journal of Medicine* 1999; 340: 669-676. DOI: 10.1056/nejm199903043400901.
21. Hshieh TT, Yue J, Oh E, et al. Effectiveness of multicomponent nonpharmacological delirium interventions: a meta-analysis. *JAMA Internal Medicine* 2015; 175: 512-520. 2015/02/03. DOI: 10.1001/jamainternmed.2014.7779.
22. Siddiqi N, Harrison JK, Clegg A, et al. Interventions for preventing delirium in hospitalised non-ICU patients. *Cochrane Database of Systematic Reviews* 2016. DOI: 10.1002/14651858.CD005563.pub3.
23. Abraha I, Trotta F, Rimland JM, et al. Efficacy of Non-Pharmacological Interventions to Prevent and Treat Delirium in Older Patients: A Systematic Overview. The SENATOR project ONTOP Series. *PLOS ONE* 2015; 10: e0123090. DOI: 10.1371/journal.pone.0123090.
24. Bush SH, Lawlor PG, Ryan K, et al. Delirium in adult cancer patients: ESMO Clinical Practice Guidelines†. *Annals of Oncology* 2018; 29: iv143–iv165. DOI: 10.1093/annonc/mdy147.
25. Lawlor PG, Rutkowski NA, MacDonald AR, et al. A scoping review to map empirical evidence regarding key domains and questions in the clinical pathway of delirium in palliative care. *Journal of Pain and Symptom Management* 2018; 57: 661-681. DOI: 10.1016/j.jpainsymman.2018.12.002.
26. Agar M, Lawlor PG, Quinn S, et al. Efficacy of oral risperidone, haloperidol, or placebo for symptoms of delirium among patients in palliative care: A randomized clinical trial. *JAMA Internal Medicine* 2017; 177: 34-42. DOI: 10.1001/jamainternmed.2016.7491.
27. Hui D, Frisbee-Hume S, Wilson A, et al. Effect of Lorazepam With Haloperidol vs Haloperidol Alone on Agitated Delirium in Patients With Advanced Cancer Receiving Palliative Care: A Randomized Clinical Trial. *JAMA* 2017; 318: 1047-1056. DOI: 10.1001/jama.2017.11468.
28. To THM, Greene AG, Agar M, et al. A point prevalence survey of hospital inpatients to define the proportion with palliation as the primary goal of care and the need for specialist palliative care. *Internal Medicine Journal* 2011; 41: 430-433. DOI: 10.1111/j.1445-5994.2011.02484.x.
29. National Gold Standards Framework Centre in End of Life Care. Gold Standards Framework Proactive Identification Guidance (GSF PIG), <http://www.goldstandardsframework.org.uk/pig> (2016).
30. Moher D, Liberati A, Tetzlaff J, et al. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med* 2009; 6: e1000097. DOI: 10.1371/journal.pmed.1000097.
31. American Psychiatric Association. DSM-III Diagnostic and Statistical Manual of Mental Disorders. In: Association AP, (ed.). 3rd ed. Washington DC: American Psychiatric Association, 1980.
32. Covidence systematic review software, Veritas Health Innovation. Melbourne, Australia. Available at www.covidence.org.
33. Hoffmann TC, Glasziou PP, Boutron I, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ* 2014; 348. DOI: 10.1136/bmj.g1687.
34. Healthcare Improvement Scotland. Scottish Intercollegiate Guidelines Network Methodology Checklists, <http://www.sign.ac.uk/checklists-and-notes.html> (accessed June 7 2018).
35. Review Manager (RevMan) [Computer program]. Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

36. Hamzhepour H, Valiee S, Azad Majedi M, et al. The Effect of Care Plan Based on Roy Adaptation Model on the Incidence and Severity of Delirium in Intensive Care Unit Patients: A Randomised Controlled Trial. *Journal of Clinical and Diagnostic Research* 2018; 12: LC21-LC25. November 1,2018. DOI: 10.7860/jcdr/2018/36366.12256.
37. Johnson K, Fleury J and McClain D. Music intervention to prevent delirium among older patients admitted to a trauma intensive care unit and a trauma orthopaedic unit. *Intensive and Critical Care Nursing* 2018; 47: 7-14. DOI: 10.1016/j.iccn.2018.03.007.
38. Bryczkowski SB, Lopreiato MC, Yonclas PP, et al. Delirium prevention program in the surgical intensive care unit improved the outcomes of older adults. *Journal of Surgical Research* 2014; 190: 280-288. 2014/03/29. DOI: 10.1016/j.jss.2014.02.044.
39. Holt R, Young J and Heseltine D. Effectiveness of a multi-component intervention to reduce delirium incidence in elderly care wards. *Age and Ageing* 2013; 42: 721-727. DOI: 10.1093/ageing/aft120.
40. Kamdar BB, King LM, Collop NA, et al. The effect of a quality improvement intervention on perceived sleep quality and cognition in a medical ICU. *Critical Care Medicine* 2013; 41: 800. DOI: 10.1097/CCM.0b013e3182746442.
41. Zaubler TS, Murphy K, Rizzuto L, et al. Quality improvement and cost savings with multicomponent delirium interventions: replication of the hospital elder life program in a community hospital. *Psychosomatics* 2013; 54: 219-226. DOI: 10.1016/j.psych.2013.01.010.
42. Marcantonio ER, Flacker JM, Wright RJ, et al. Reducing delirium after hip fracture: a randomized trial. *Journal of the American Geriatrics Society* 2001; 49: 516-522. DOI: 10.1046/j.1532-5415.2001.49108.x.
43. Jeffs KJ, Berlowitz DJ, Grant S, et al. An enhanced exercise and cognitive programme does not appear to reduce incident delirium in hospitalised patients: a randomised controlled trial. *BMJ Open* 2013; 3: e002569. DOI: 10.1136/bmjopen-2013-002569.
44. Caplan GA and Harper EL. Recruitment of volunteers to improve vitality in the elderly: the REVIVE study. *Internal Medicine Journal* 2007; 37: 95-100. DOI: 10.1111/j.1445-5994.2007.01265.x.
45. Wong Tin Niam DM, Bruce JJ and Bruce DG. Quality project to prevent delirium after hip fracture. *Australasian Journal on Ageing* 2005; 24: 174-177. DOI: 10.1111/j.1741-6612.2005.00108.x.
46. Deschodt M, Braes T, Flamaing J, et al. Preventing delirium in older adults with recent hip fracture through multidisciplinary geriatric consultation. *Journal of the American Geriatrics Society* 2012; 60: 733-739. DOI: 10.1111/j.1532-5415.2012.03899.x.
47. Van Rompaey B, Elseviers MM, Drom W, et al. The effect of earplugs during the night on the onset of delirium and sleep perception: a randomized controlled trial in intensive care patients. *Critical Care* 2012; 16: R73. DOI: 10.1016/j.iccn.2015.03.002.
48. Milisen K, Foreman MD, Abraham IL, et al. A Nurse-Led Interdisciplinary Intervention Program for Delirium in Elderly Hip-Fracture Patients. *Journal of the American Geriatrics Society* 2001; 49: 523-532. DOI: 10.1046/j.1532-5415.2001.49109.x.
49. Gagnon P, Allard P, Gagnon B, et al. Delirium prevention in terminal cancer: assessment of a multicomponent intervention. *Psycho-Oncology* 2012; 21: 187-194. DOI: 10.1002/pon.1881.
50. Cole MG, McCusker J, Bellavance F, et al. Systematic detection and multidisciplinary care of delirium in older medical inpatients: a randomized trial. *Canadian Medical Association Journal* 2002; 167: 753-759.
51. Cole MG, Primeau FJ, Bailey RF, et al. Systematic intervention for elderly inpatients with delirium: a randomized trial. *Canadian Medical Association Journal* 1994; 151: 965-970.
52. Bakker FC, Persoon A, Bredie SJ, et al. The CareWell in Hospital program to improve the quality of care for frail elderly inpatients: results of a before-after study with focus on surgical patients. *American Journal of Surgery* 2014; 208: 735-746. DOI: 10.1016/j.amjsurg.2014.04.009.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
53. Hempenius L, Slaets JPJ, van Asselt D, et al. Outcomes of a geriatric liaison intervention to prevent the development of postoperative delirium in frail elderly cancer patients: Report on a multicentre, randomized, controlled trial. *PloS One* 2013; 8: e64834. DOI: 10.1371/journal.pone.0064834.
54. Lundström M, Edlund A, Karlsson S, et al. A multifactorial intervention program reduces the duration of delirium, length of hospitalization, and mortality in delirious patients. *Journal of the American Geriatrics Society* 2005; 53: 622-628. DOI: 10.1111/j.1532-5415.2005.53210.x.
55. Lundström M, Olofsson B, Stenvall M, et al. Postoperative delirium in old patients with femoral neck fracture: a randomized intervention study. *Aging Clinical and Experimental Research* 2007; 19: 178-186.
56. Martinez FT, Tobar C, Beddings CI, et al. Preventing delirium in an acute hospital using a non-pharmacological intervention. *Age and Ageing* 2012; 41: 629-634. DOI: 10.1093/ageing/afs060.
57. Andro M, Comps E, Estivin S, et al. Prevention of delirium in demented hospitalized patients. *European Journal of Internal Medicine* 2012; 23: 124-125. 2012/01/31. DOI: 10.1016/j.ejim.2011.05.011.
58. Black P, Boore JRP and Parahoo K. The effect of nurse-facilitated family participation in the psychological care of the critically ill patient. *Journal of Advanced Nursing* 2011; 67: 1091-1101. DOI: 10.1111/j.1365-2648.2010.05558.x.
59. Bo M, Martini B, Ruatta C, et al. Geriatric ward hospitalization reduced incidence delirium among older medical inpatients. *American Journal of Geriatric Psychiatry* 2009; 17: 760-768. DOI: 10.1097/JGP.0b013e3181a315d5.
60. Taguchi T, Yano M and Kido Y. Influence of bright light therapy on postoperative patients: a pilot study. *Intensive & Critical Care Nursing* 2007; 23: 289-297. DOI: 10.1016/j.iccn.2007.04.004.
61. Moon K-J and Lee S-M. The effects of a tailored intensive care unit delirium prevention protocol: A randomized controlled trial. *International Journal of Nursing Studies* 2015; 52: 1423-1432. DOI: 10.1016/j.ijnurstu.2015.04.021.
62. Chong MS, Chan M, Tay L, et al. Outcomes of an innovative model of acute delirium care: the Geriatric Monitoring Unit (GMU). *Clinical Interventions in Aging* 2014; 9: 603-612. DOI: 10.2147/CIA.S60259.
63. Vidan MT, Sanchez E, Alonso M, et al. An intervention integrated into daily clinical practice reduces the incidence of delirium during hospitalization in elderly patients. *Journal of the American Geriatrics Society* 2009; 57: 2029-2036. DOI: 10.1111/j.1532-5415.2009.02485.x.
64. Patel J, Baldwin J, Bunting P, et al. The effect of a multicomponent multidisciplinary bundle of interventions on sleep and delirium in medical and surgical intensive care patients. *Anaesthesia* 2014; 69: 540-549. DOI: 10.1111/anae.12638.
65. Gruber-Baldini AL, Marcantonio E, Orwig D, et al. Delirium outcomes in a randomized trial of blood transfusion thresholds in hospitalized older adults with hip fracture. *Journal of the American Geriatrics Society* 2013; 61(8): 1286-95. DOI: 10.1111/jgs.12396.
66. Charlson ME, Pompei P, Ales K, et al. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *Journal of Chronic Disease* 1987; 40: 373-383.
67. Knaus WA, Draper EA, Wagner DP, et al. APACHE II: a severity of disease classification system. *Critical Care Medicine* 1985; 13: 818-829.
68. Miller MD, Paradis CF, Houck PR, et al. Rating chronic medical illness burden in geropsychiatric practice and research: application of the Cumulative Illness Rating Scale. *Psychiatry Research* 1992; 41: 237-248. 1992/03/01. DOI: 10.1016/0165-1781(92)90005-N.
69. Saklad M. Grading of patients for surgical procedures. *Anesthesiology: The Journal of the American Society of Anesthesiologists* 1941; 2: 281-284.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
70. Blessed G, Tomlinson BE and Roth M. The Association Between Quantitative Measures of Dementia and of Senile Change in the Cerebral Grey Matter of Elderly Subjects. *British Journal of Psychiatry* 1968; 114: 797-811. DOI: 10.1192/bjp.114.512.797.
71. Subbe CP, Kruger M, Rutherford P, et al. Validation of a modified Early Warning Score in medical admissions. *QJM: An International Journal of Medicine* 2001; 94: 521-526. DOI: 10.1093/qjmed/94.10.521.
72. Vincent J-L, Moreno R, Takala J, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. *Intensive Care Medicine* 1996; 22: 707-710.
73. Bellomo R, Ronco C, Kellum JA, et al. Acute renal failure—definition, outcome measures, animal models, fluid therapy and information technology needs: the Second International Consensus Conference of the Acute Dialysis Quality Initiative (ADQI) Group. *Critical Care* 2004; 8: R204. DOI: 10.1186/cc2872.
74. Metnitz PGH, Moreno RP, Almeida E, et al. SAPS 3—From evaluation of the patient to evaluation of the intensive care unit. Part 1: Objectives, methods and cohort description. *Intensive Care Medicine* 2005; 31: 1336-1344. DOI: 10.1007/s00134-005-2762-6.
75. Yesavage JA and Sheikh JI. 9/Geriatric depression scale (GDS) recent evidence and development of a shorter version. *Clinical Gerontologist* 1986; 5: 165-173. DOI: 10.1300/J018v05n01_09.
76. Taylor JS, DeMers SM, Vig EK, et al. The disappearing subject: exclusion of people with cognitive impairment and dementia from geriatrics research. *Journal of the American Geriatric Society* 2012; 60: 413-419. DOI: 10.1111/j.1532-5415.2011.03847.x.
77. Singh H, Kanapuru B, Smith C, et al. FDA analysis of enrollment of older adults in clinical trials for cancer drug registration: A 10-year experience by the U.S. Food and Drug Administration. *Journal of Clinical Oncology* 2017; 35: 10009-10009. DOI: 10.1200/JCO.2017.35.15_suppl.10009.
78. Hosie A, Agar M, Lobb E, et al. Improving delirium recognition and assessment for people receiving inpatient palliative care: a mixed methods meta-synthesis. *International Journal of Nursing Studies* 2017; 75: 123-129. DOI: 10.1016/j.ijnurstu.2017.07.007.
79. Knaul FM, Farmer PE, Krakauer EL, et al. Alleviating the access abyss in palliative care and pain relief—an imperative of universal health coverage: the Lancet Commission report. *The Lancet* 2017; 39(10128): 1391-1454. DOI: 10.1016/S0140-6736(17)32513-8.
80. Boland J, Currow DC, Wilcock A, et al. A systematic review of strategies used to increase recruitment of people with cancer or organ failure into clinical trials: implications for palliative care research. *Journal of Pain and Symptom Management* 2015; 49: 762-772.e765. DOI: 10.1016/j.jpainsymman.2014.09.018.
81. Hanson LC, Bull J, Wessell K, et al. Strategies to support recruitment of patients with life-limiting illness for research: the Palliative Care Research Cooperative Group. *Journal of Pain and Symptom Management* 2014; 48: 1021-1030. DOI: 10.1016/j.jpainsymman.2014.04.008.
82. Girard TD, Exline MC, Carson SS, et al. Haloperidol and Ziprasidone for Treatment of Delirium in Critical Illness. *New England Journal of Medicine* 2018; 379: 2506-2516. DOI: 10.1056/NEJMoa1808217.
83. Marcantonio ER, Ngo LH, O'Connor M, et al. 3D-CAM: Derivation and Validation of a 3-Minute Diagnostic Interview for CAM-Defined Delirium. *Annals of Internal Medicine* 2014; 161: 554-561. DOI: 10.7326/m14-0865.
84. Ely EW, Margolin R, Francis J, et al. Evaluation of delirium in critically ill patients: validation of the Confusion Assessment Method for the Intensive Care Unit (CAM-ICU). *Critical Care Medicine* 2001; 29: 1370-1379.
85. Han JH, Wilson A, Vasilevskis EE, et al. Diagnosing Delirium in Older Emergency Department Patients: Validity and Reliability of the Delirium Triage Screen and the Brief Confusion Assessment

- 1
2
3
4
5 Method. *Annals of Emergency Medicine* 2013; 62: 457-465. DOI:
6 10.1016/j.annemergmed.2013.05.003.
7 86. Hospital Elder Life Program. Confusion Assessment Method (Short CAM),
8 <http://www.hospitalelderlifeprogram.org/delirium-instruments/short-cam/> (2015, accessed April 12
9 2019).
- 10 87. Van Mechelen W, Aertgeerts B, De Ceulaer K, et al. Defining the palliative care patient: A
11 systematic review. *Palliative Medicine* 2012; 27: 197-208. DOI: 10.1177/0269216311435268.
- 12 88. Downar J, Goldman R, Pinto R, et al. The “surprise question” for predicting death in seriously
13 ill patients: a systematic review and meta-analysis. *Canadian Medical Association Journal* 2017;
14 189(13): E484-E493. DOI: 10.1503/cmaj.160775.
- 15 89. Murtagh FEM, Bausewein C, Verne J, et al. How many people need palliative care? A study
16 developing and comparing methods for population-based estimates. *Palliative Medicine* 2013; 28(1):
17 49-58. DOI: 10.1177/0269216313489367.
- 18 90. Waller A, Girgis A, Currow D, et al. Development of the Palliative Care Needs Assessment
19 Tool (PC-NAT) for use by multi-disciplinary health professionals. *Palliative Medicine* 2008; 22(8): 956-
20 964. DOI: 10.1177/0269216308098797.
- 21 91. Hight G, Crawford D, Murray SA, et al. Development and evaluation of the Supportive and
22 Palliative Care Indicators Tool (SPICT): a mixed-methods study. *BMJ Supportive & Palliative Care*
23 2014; 4(3): 285. DOI: 10.1136/bmjspcare-2013-000488.
- 24 92. Moreno RP, Metnitz PGH, Almeida E, et al. SAPS 3—From evaluation of the patient to
25 evaluation of the intensive care unit. Part 2: Development of a prognostic model for hospital
26 mortality at ICU admission. *Intensive Care Medicine* 2005; 31(10): 1345-1355. DOI: 10.1007/s00134-
27 005-2763-5.
- 28 93. Williams MA, Ward SE and Campbell EB. Confusion: Testing versus Observation. *Journal of*
29 *Gerontological Nursing* 1988; 14(1): 25-30. DOI: 10.3928/0098-9134-19880101-08.
- 30 94. Pandharipande PP and Ely EW. Humanizing the Treatment of Hyperactive Delirium in the
31 Last Days of Life. *JAMA* 2017; 318(11): 1104-1105. DOI: 10.1001/jama.2017.11466.
- 32 95. Virdun C, Lockett T, Davidson PM, et al. Dying in the hospital setting: A systematic review of
33 quantitative studies identifying the elements of end-of-life care that patients and their families rank
34 as being most important. *Palliative Medicine* 2015; 29(9): 774-796. 2015/04/30. DOI:
35 10.1177/0269216315583032.
- 36 96. Virdun C, Lockett T, Lorenz K, et al. Dying in the hospital setting: A meta-synthesis identifying
37 the elements of end-of-life care that patients and their families describe as being important.
38 *Palliative Medicine* 2016; 31(7): 687-601. DOI: 10.1177/0269216316673547.
- 39 97. Reynish EL, Hapca SM, De Souza N, et al. Epidemiology and outcomes of people with
40 dementia, delirium, and unspecified cognitive impairment in the general hospital: prospective cohort
41 study of 10,014 admissions. *BMC Medicine* 2017; 15: 140. 2017/07/28. DOI: 10.1186/s12916-017-
42 0899-0.
- 43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Inclusion, characteristics and outcomes of people requiring palliative care in studies of non-pharmacological interventions for delirium: a systematic review

Abstract

Background

Delirium is common, distressing, serious and under-researched in specialist palliative care settings.

Objectives

To examine whether people requiring palliative care were included in non-pharmacological delirium intervention studies in inpatient settings, how they were characterised, and what their outcomes were.

Design

Systematic review (PROSPERO 2017 CRD42017062178).

Data sources

Systematic search in March 2017 for non-pharmacological delirium intervention studies in adult inpatients. Database search terms were 'delirium', 'hospitalisation', 'inpatient', 'palliative care', 'hospice', 'critical care', 'geriatrics'. Scottish Intercollegiate Guidelines Network methodological checklists guided risk of bias assessment.

Results

The 29 included studies were conducted between 1994-2015 in diverse settings in 15 countries (9136 participants, mean age 76.5 years [SD 8.1], 56% women). Most studies tested multicomponent interventions (n=26) to prevent delirium (n=19). Three-quarters of the 29 included studies (n=22) excluded various groups of people requiring palliative care; however, inclusion criteria, participant diagnoses, illness severity and mortality indicated their presence in almost all studies (n=26). Of these, 21 studies did not characterise

1
2
3
4 participants requiring palliative care or report their specific outcomes (72%), four reported
5
6 outcomes for older people with frailty, dementia, cancer and comorbidities, and one was
7
8 explicitly focused on people receiving palliative care. Study heterogeneity and limitations
9
10 precluded definitive determination of intervention effectiveness and only allowed
11
12 interpretations of feasibility for people requiring palliative care. Acceptability outcomes
13
14 (intervention adverse events and patients' subjective experience) were rarely reported
15
16 overall.

17 18 **Conclusion**

19
20 Non-pharmacological delirium interventions have frequently excluded and under-
21
22 characterised people requiring palliative care and infrequently reported their outcomes.
23

24 25 **Key words**

26
27 Clinical trial, Delirium, Hospice, Inpatient, Palliative Care, Review
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Key statements

What is already known about the topic?

- Delirium is a distressing and serious neurocognitive condition that frequently occurs for patients in palliative care inpatient settings.
- In contrast to other hospital settings, there is limited evidence to guide non-pharmacological intervention to prevent and treat delirium in palliative care inpatient settings.

What this paper adds

- This review found that various groups of people requiring palliative care were excluded from three-quarters of non-pharmacological delirium intervention studies in inpatient settings; despite this, they were present in most studies and their outcomes were reported in five.
- Non-pharmacological delirium interventions appear feasible for people requiring palliative care yet there is no definitive evidence they are effective or acceptable for this inpatient group.

Implications for research

- Phase II and III randomised controlled trials of non-pharmacological interventions to prevent and treat delirium are needed in specialist palliative care settings.
- Adaptations to future trials of non-pharmacological delirium interventions in other inpatient settings are needed to promote representative study populations and allow outcomes for sub-groups of people requiring palliative care to be reported.
- Additional outcomes related to patient and family subjective experience, goals of care and quality of life would enhance the relevance of delirium intervention research in inpatient settings where people are cared for at the end of life.

Introduction

Delirium is a serious complication of medical illness and hospitalisation.¹ The condition is characterised by acute disturbances to attention, awareness and cognition, has multifactorial aetiology, and variously affects memory, language and visuospatial ability, orientation and perception.² Affected persons often experience feelings of fear, humiliation, confusion and disconnection from others.^{3, 4} Family members' experience distress when delirium causes sudden changes in behaviour and decline in the person they love.^{5, 6} Patients who experience an episode of delirium during hospitalisation experience many poorer outcomes, including being more likely to die.^{1, 7, 8}

Delirium most often occurs in people with older age, advanced or severe illness and/or pre-existing cognitive impairment. Hospital-wide, one in five patients have delirium,⁹ with occurrence higher again in intensive, geriatric and palliative care units.^{1, 6} Studies of delirium epidemiology in palliative care inpatient units that screened patients at least daily reported incidence of 33–45% and prevalence of 58–88% in those who died.

Development of delirium guidelines^{1, 10–13} policy¹⁴ and international advocacy¹⁵ indicate growing awareness of the seriousness and prevalence of delirium and importance of evidence based care for hospitalised patients.^{16, 17} There now is sufficient evidence to implement non-pharmacological interventions for delirium in certain hospital settings.^{18, 19} For example, reviews of studies of multicomponent interventions addressing physical and cognitive activity, sleep, hearing, vision and hydration, as in the original Hospital Elder Life Program (HELP) study,²⁰ reported reduction in incident delirium in older hospitalised patients.^{21–23} Reduction in length of hospital stay and improvement in return to independent living were also demonstrated.²²

In contrast, guideline recommendations for non-pharmacological interventions as the first approach to prevent and treat delirium during advanced illness and at the end of life are not evidence based.^{10, 17, 24} A recent scoping review reported the need to generate evidence to inform clinical care in palliative care settings and populations, for non-pharmacological interventions in particular.²⁵ Poorer outcomes with antipsychotics,²⁶ and over-sedation

1
2
3
4 when benzodiazepines were given for agitated delirium,²⁷ in two recent trials in specialist
5 palliative care settings also highlight the need to establish 'drug-free' ways to prevent and
6 relieve the difficulties of delirium at the end of life.
7
8
9

10 In response, the authors established the '*Studies to Understand and Improve Delirium Care*
11 *in Palliative Settings*' international collaborative (SUNRISE) to generate high-quality delirium
12 research in palliative care. We identified the need to inform our future trials in palliative
13 care through a review of studies of non-pharmacological interventions for delirium
14 conducted in a wide range of inpatient settings. This wide review was premised on our
15 clinical experience and knowledge that many hospitalised patients have advanced and/or
16 serious illness, frailty and high comorbidity and consequently much in common with
17 patients in specialist palliative care settings, especially those in intensive care, medical and
18 geriatric units where rates of delirium are similarly high.^{28, 29} Based on this premise, our
19 specific objectives were to examine whether people requiring palliative care were included
20 in non-pharmacological delirium intervention studies in various inpatient settings, how
21 these participants were characterised, and whether the non-pharmacological interventions
22 were effective, feasible and/or acceptable for them.
23
24
25
26
27
28
29
30
31
32
33
34
35

36 **Methods**

37 **Design**

38
39
40 Systematic review of the literature, following the Preferred Reporting Items for Systematic
41 Reviews and Meta-Analyses (PRISMA).³⁰
42
43
44

45 **Participants/settings**

46
47 Adults (≥ 18 years) receiving inpatient hospital or hospice care. In this review, we refer to
48 'hospice' as an inpatient facility with the primary function to provide specialist palliative
49 care to people with life-threatening illness, and analogous to a palliative care inpatient unit.
50
51
52
53

54 **Search strategy**

55
56 In March 2017, we searched MEDLINE, Embase, CINAHL, PsycINFO, Cochrane Library and
57 Web of Science with the following search terms in MEDLINE: Delirium AND Hospitalization
58
59
60

1
2
3
4 OR Inpatient OR Hospice AND Palliative Care OR Critical Care OR Geriatrics. Filters in
5
6 Medline were: 1. Study types: clinical study, clinical trial all, comparative study, controlled
7
8 clinical trial, meta-analysis, multicenter study, pragmatic clinical trial, randomised controlled
9
10 trial, systematic review; 2. Peer reviewed journal and 3. Published from 1980 onward (when
11
12 delirium was first included in the American Psychological Association (APA) Diagnostic and
13
14 Statistical Manual (DSM-III).³¹ We tailored search terms and filters to subsequent databases.
15
16 We examined reference lists of relevant systematic reviews and meta-analyses identified in
17
18 the search for additional eligible studies.

19 20 **Inclusion/exclusion criteria**

21
22 Studies included were primary reports of prospective trials (i.e. studies of an intervention
23
24 with a comparator); with a primary objective to prevent or treat delirium through non-
25
26 pharmacological intervention/s in adult patients in hospital or hospice unit settings; a
27
28 primary outcome of delirium incidence, severity or duration; published in English in a peer-
29
30 reviewed journal.

31
32 Studies excluded were reports of interventions for alcohol withdrawal delirium only;
33
34 systematic reviews and meta-analyses of non-pharmacological delirium intervention/s;
35
36 studies where the primary outcome was not participants' delirium status (e.g. process or
37
38 cost effectiveness outcomes, validation of delirium tools); studies that did not use diagnostic
39
40 criteria or a tool with established psychometric properties to measure delirium; protocols;
41
42 and ongoing studies.

43 44 **Study selection, data extraction and management**

45
46 We imported search results into Endnote X7 software, removed duplicates and exported
47
48 results to Covidence,³² where three reviewers [IAD, LE, AH] independently applied eligibility
49
50 criteria to titles and abstracts. Reviewers compared decisions about inclusion, documented
51
52 reasons for exclusions at full text review and resolved discrepancies through discussion. Two
53
54 reviewers [IAD, LE] extracted data according to the template for intervention description
55
56 and replication (TIDieR) checklist and guide³³ into an Excel V15.28 spreadsheet. The lead
57
58 reviewer [AH] contributed guidance, oversight and independent data checking.
59
60

Risk of bias assessment

Two reviewers [LE and AH] independently assessed each study for selection, performance, detection, attrition, confounding, and reporting biases according to the Scottish Intercollegiate Guidelines Network (SIGN) Methodology Checklists³⁴ for controlled trials and cohort studies. Discrepancies were resolved through discussion.

Outcomes

To identify our sample of interest (i.e. people requiring palliative care), we examined study inclusion and exclusion criteria, participant diagnoses (including severity or staging), and mortality. We assessed eligibility criteria and diagnoses against the Gold Standards Framework Proactive Identification Guidance (GSF PIG), a clinical tool to help identify people likely to need additional supportive (i.e. palliative) care in the last 12 months of life.²⁹ According to the GSF PIG, these people are those with life-threatening conditions, including illnesses that are advanced, progressive, incurable and/or likely to cause acute crises; frailty and co-morbidities; and sudden catastrophic events.

Previous reviews have reported effectiveness outcomes of non-pharmacological delirium interventions for the entire study sample;²¹⁻²³ whereas this review focused on effectiveness, feasibility and acceptability outcomes of our sample of interest. We examined effectiveness according to each study's primary outcome and any sub-group analysis for our sample of interest. We assessed feasibility by examining intervention characteristics, adherence and study modifications, and acceptability through intervention-related adverse effects and patient, family, clinician and volunteer subjective experiences of the interventions.

Synthesis and analysis plan

We generated tables, text and graphs to report study characteristics, participants, interventions and outcomes. Data transformation and descriptive numerical analyses were performed using Excel. We planned to perform subgroup meta-analysis using Review Manager Analyses software³⁵ of intervention effectiveness if we could definitively distinguish our sample of interest and if interventions and comparators and measures were comparable.

Updated search

We updated the search prior to publication in February 2019 and identified two new eligible papers published after the original search date.^{36, 37} There were not incorporated as neither paper altered the conclusions of the review.

Results

The original database search strategy generated 4300 records. After removing 35 duplicates and 4169 records through title and abstract screening, we reviewed 69 full text papers and excluded 48. We included another eight through reference list searching, resulting in 29 papers reporting 29 studies for inclusion (Figure 1).

(Insert Figure 1 PRISMA Flow Diagram here)³⁰

Study characteristics

The 29 studies were conducted between 1994-2015 across 15 countries: six in the US,^{38 39 40 41 42 20} three each in Australia,^{43 44 45} Belgium,⁴⁶⁻⁴⁸ and Canada,⁴⁹⁻⁵¹ two each in the Netherlands^{52, 53} and Sweden,^{54, 55} and one each in Chile,⁵⁶ France,⁵⁷ Ireland,⁵⁸ Italy,⁵⁹ Japan,⁶⁰ Korea,⁶¹ Singapore,⁶² Spain,⁶³ UK,⁶⁴ and the US/Canada⁶⁵ (Table 1).

Study designs were before/after studies (one with an additional concurrent arm⁶²) (n=11),^{38-41, 44, 45, 48, 52, 57, 62, 64} randomised controlled trials (RCTs) (n=10),^{42, 43, 47, 50, 51, 53, 55, 56, 61, 65} non-randomised controlled trials (one with matched participants²⁰) (n=5),^{20, 46, 54, 59, 63} a quasi-experimental study,⁴⁹ a pilot randomised controlled trial,⁶⁰ and a comparative time series study.⁵⁸

Services and settings were medical (n=10),^{20, 41, 43, 50-52, 54, 56, 59, 63} geriatric (n=7),^{39, 44, 55, 57, 59, 62, 63} medical and/or surgical intensive care (n=6),^{38, 40, 47, 58, 61, 64} perioperative hip fracture (n=6),^{42, 45, 46, 48, 55, 65} other perioperative (n=3)^{52, 53, 60} and palliative care and hospice units (n=1),⁴⁹ with eight studies involving more than one service.^{47, 52, 55, 58, 59, 61, 63, 64}

Most studies tested multicomponent interventions (n=26) and aimed to prevent delirium in non-delirious participants (n=19) (Table1). Six studies aimed to prevent and treat delirium^{38, 42, 48, 54, 55, 65} three were treatment only studies.^{50, 51, 62}

(Insert Table 1 Study characteristics here)

Code: **P** Prevention, **T** Treatment, **RCT** Randomised controlled trial, **CTS** Comparative time series, **B/A** Before–after study, **CT** Controlled trial, **ICU** Intensive Care Unit, **NR** Not reported, **MC** Medical Centre, **M** Multicomponent, **PRCT** Pilot RCT, **QE** Quasi-experimental, **S** Single component

^Δ Numbers signify the domains in which the study was assessed as having **low risk of bias**: 1 = Representative study sample; 2 = Concealed allocation; 3 = Random sequence generation; 4 = Blinded participants and intervention personnel; 5 = Blinded outcome assessors; 6 = Valid, reliable delirium measures; 7 = <20% sample lost to analysis; 8 = Intention to treat analysis; 9 = Confounders accounted for; 10 = Only *a priori* outcomes reported

Risk of bias assessment

Almost all studies were assessed as having at least one form of selection bias (n=28). The exception was a Chilean RCT of a prevention intervention where family members delivered cognition, vision, and hearing strategies to patients.⁵⁶ Eight RCTs minimised internal selection bias through concealed allocation and random sequence generation.^{42, 43, 47, 50, 53, 55, 56, 65} There was high risk of performance bias in all studies, due to the inherent difficulty of blinding non-pharmacological interventions: one RCT blinded participants to their allocation,⁴³ and no studies blinded those performing the intervention. Detection bias was minimised in eight studies through blinded outcome assessors combined with valid, reliable delirium measures.^{42, 43, 47, 50, 51, 53-55} Although we excluded studies that did not use delirium diagnostic criteria and/or a tool with established psychometric properties, three studies were assigned uncertain risk of detection bias due to retrospective assignment of delirium status when the Confusion Assessment Method (CAM) was not performed during weekends in two before-after studies;^{41, 45} and when delirium incidence was reported as a mean rather than a proportion in a comparative time series study.⁵⁸ High risk of detection bias was assigned to a delirium prevention study in seven Canadian palliative care units due use of a screening tool, the Confusion Rating Scale (CRS) to measure delirium incidence, severity and duration.⁴⁹ Six studies had low risk of attrition bias.^{42, 43, 50, 55, 56, 63} Confounders were accounted for in 17 studies.^{20, 39, 40, 42, 46, 47, 49-52, 54, 55, 58, 59, 61, 63, 65} Most studies (n=22) reported only *a priori* outcomes,^{20, 38-43, 46, 48, 50-53, 55-57, 60-65} which we assessed by information provided within the papers rather than through examination of trial registrations or published protocols, which no studies cited.

Figure 2 presents overall risk of bias in the included studies.

(Insert Figure 2 Overall risk of bias graph here)

Figure 2: Overall risk of bias

Participant characteristics

There were 9136 participants (n=4553 in intervention arms) across the 29 studies (Table 2). Sample sizes ranged from 15-1516 participants. Participants' overall mean age (excluding two studies that reported only age range or median)^{48, 58} was 76.5 years (SD \pm 8.1). There were more female participants (56%) than male.

Inclusion and exclusion criteria

Inclusion

Most studies (n=19) included only patients aged 65 years and older (Table 2). Twenty-two studies included consecutively admitted patients, 16 of which further required certain physical conditions to be present (e.g. frailty,⁵³ hip fracture,^{42, 45, 46, 48, 55, 65} anemia,⁶⁵ dementia,⁵⁷ advanced cancer,⁴⁹ oesophageal cancer surgery⁶⁰). Five studies included only patients with risk factors for delirium^{20, 44, 52, 56, 63} and three included only patients with delirium.^{50, 51, 62} The study in seven Canadian palliative care units (hereafter termed the 'palliative care unit study') included only patients with terminal cancer, defined as dying within 90 days of admission which was retrospectively identified. This study had the largest sample of all the studies (n=1516, 17%).⁴⁹

Exclusion

Twenty-three studies excluded patients on various diagnostic grounds (Table 2). These were stroke or cerebrovascular accident (CVA), moderate-severe traumatic brain injury or coma (n=9);^{20, 38, 43, 48, 50, 51, 59, 62, 63} dementia or prior cognitive impairment (n=7);^{20, 38, 44, 47, 61, 63, 64} psychiatric disorders or alcohol/drug addiction (n=6);^{43, 44, 59-61, 64} pathological fracture (n=4);^{46, 48, 55, 65} poly- or multiple trauma (n=2);^{46, 48} neurologic diagnosis/neurosurgical (n=2);^{61, 64} metastatic cancer or referred to oncology services (n=2);^{42, 50} and others (myocardial infarction,⁶⁵ severe renal failure,⁵⁵ brain concussion,⁴⁸ history of sleep pathology,⁶⁴ severe rheumatoid arthritis⁵⁵ and severe hip osteoarthritis⁵⁵).

1
2
3
4 Fifteen studies excluded patients with impaired verbal communication or who were
5 otherwise unable to complete study assessments.^{20, 38, 39, 41, 43, 46, 48, 50, 52, 53, 59, 61-63, 65} Six
6 studies excluded patients with hearing and/or visual impairment.^{46-48, 60, 61, 64} Three studies
7 excluded patients isolated for infection.^{43, 61, 62}

8
9
10
11
12 Eight studies explicitly excluded patients expected to die, using the following terms:
13 “terminal diagnosis” or “terminal illness” (n=3),^{20, 58, 62} “terminal condition and receiving
14 comfort care” (on the basis that they were unlikely to benefit from the intervention)
15 (n=2),^{41, 44} “life expectancy less than six months”,⁴⁶ “metastatic cancer or comorbid illnesses
16 likely to reduce life expectancy to less than six months”,⁴² “death expected within 24
17 hours”,⁴³ and “dangerously ill”.⁶²

18
19
20
21
22
23 Two studies excluded participants who subsequently died from analysis;^{51, 52} a delirium
24 prevention study in elderly care wards excluded 70% of 2162 admitted patients due to being
25 “too unwell”;³⁹ and another study excluded patients with “deterioration of condition”.⁶⁰

26
27
28
29
30 The palliative care unit study excluded patients who lived for more than 90 days or who
31 were still alive at discharge from analysis.⁴⁹

32 33 34 **Characterisation of study participants**

35 36 ***Participant diagnoses and illness severity***

37
38 Reporting of participant diagnoses, illness severity and mortality varied. Fifteen studies
39 reported co-morbid diagnoses, twelve reported only primary diagnoses, and four reported
40 both (Table 2). Diagnoses’ categorisation varied. Studies rarely reported staging of
41 diagnoses (n=2).^{49, 56}

42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Sixteen studies reported illness severity for all participants using at least one measure, more
often as a mean or median score (n=15)^{20, 38, 43, 46, 47, 50, 52, 55, 56, 59, 61-65} than by categorising
participants proportionally (n=6).^{20, 39, 42, 47, 52, 62} The most frequently used illness severity
measures were the Charlson Comorbidity Index (CCI)⁶⁶ (n=7; reported as: mean scores 2.3 -
3.3,^{46, 50, 62} median 2,^{43, 56} and scores of four or greater 19.8%³⁹ and 39%⁴²); and the Acute
Physiology And Chronic Health Evaluation II (APACHE II)⁶⁷ (n=6; mean scores 11.3-15.6).^{20, 43,}
^{59, 61, 63, 64} Two ICU studies that measured illness severity using the APACHE II reported

1
2
3
4 means scores (13.9-14.6)^{61, 64} approximately equivalent to or less than those reported in
5 three of four medical and geriatric unit studies (14.1-15.6).^{20, 43, 59} Other measures were the
6 Cumulative Illness Rating Scale–Geriatrics (CIRS-G) comorbidity and severity indexes;^{52, 59, 68}
7 American Society of Anesthesiologists physical status classification (ASA)^{52, 65, 69}; Blessed
8 Dementia Rating Scale;^{20, 42, 70} Modified Early Warning Score (MEWS);^{39, 71} Sepsis-related
9 Organ Failure (SOFA);^{47, 72} Risk, Injury, Failure, Loss of kidney function, and End-stage kidney
10 disease (RIFLE);^{47, 73} Simplified Acute Physiology Score (SAPS 3);^{47, 74} and the 15-item
11 Geriatric Depression Scale (GDS-15).^{55, 75}

12
13 Three studies reporting comorbidities without the CCI reported: i. 60% had more than two
14 comorbidities (diabetes, COPD, hypertension, myocardial infarction, other cardiovascular
15 disorders, neurological disorders, cerebrovascular disorders, hearing and vision problems,
16 memory problems in daily life, psychiatric disorders or musculoskeletal disorders);⁵³ ii. 72%
17 had at least one co-morbid illness;⁴⁷ and iii. a mean of 2.7 comorbidities.⁶³

18
19 Almost every diagnosis, sensory deficit and impairment that were exclusion criteria in some
20 studies were reported in the overall study population (Table 2). For example, more studies
21 reported participants with dementia or cognitive impairment (n=13)^{39, 41-43, 46, 48-51, 55-57, 65}
22 than excluded people with these conditions (n=7).^{20, 38, 44, 47, 61, 63, 64}

23 24 25 **Participant mortality**

26
27 Mortality was reported in 19 studies at nine different time points, ranging from ‘in the
28 ICU’^{38, 40} to one year post intervention.^{46, 55, 62} In these studies, 2090 participants died
29 (23%). Mortality rates ranged from 1% in three studies^{20, 47, 61} to 100% in the palliative care
30 unit study.⁴⁹ There were 596 participant deaths in 18 studies reporting mortality that were
31 conducted outside of a specialist palliative care setting (12%). High mortality was reported
32 for intervention and control medical unit participants at eight weeks (22.1% vs 19.37%;⁵⁰
33 33% vs 37% for delirious patients⁵¹) and at six months (33.8% vs 30.9%);³⁹ and at one year
34 for older (≥65) traumatic hip fracture patients (33% vs 22%).⁴⁶ Six of the 19 studies
35 measuring mortality excluded patients expected to die; despite this exclusion criteria, 134
36 participants of these six studies subsequently died during the study period (12%).^{20, 44, 46, 52,}
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

of which reported significantly less mortality in intervention cohorts compared to controls,^{54, 61} with one (at day 7 in hospital) not sustained at 30 days in hospital.⁶¹

Study approaches to people requiring palliative care

By examination of these data we identified only one study that explicitly reported outcomes for participants requiring palliative care (n=1).⁴⁹ Using the GSF PIG, we identified another four studies that reported primary or sub-group effectiveness outcomes for older people likely to require palliative care due to the presence of frailty, dementia, cancer and comorbidity.^{42, 52, 53, 57} All five studies were of interventions to prevent delirium. Twenty-two studies either explicitly or resultantly excluded groups of people requiring palliative care (76%).^{20, 38, 39, 41-44, 46-48, 50-52, 55, 58-65} Yet, through our interpretation of reported diagnoses, levels of comorbidity, frailty and mortality (signified with a † in Table 2), we also identified that people requiring palliative care were present in 21 studies that did not specifically characterise them or report their outcomes (72%).^{20, 38-41, 43, 45-48, 50, 51, 54-56, 58, 61-63, 65} Figure 3 presents these findings diagrammatically.

(Insert Table 2: Study inclusion and exclusion criteria and participant diagnoses, illness severity and mortality)

* Statistically significant difference ^Δ Intervention and control participants combined ^α Rounded to nearest whole number [†] Interpreted as indicating need for palliative care

Illness severity measures: Higher scores represent higher illness severity. **AIS** Abbreviated Injury Scale, **APACHE II** Acute Physiology and Chronic Health Evaluation (scores 0-71), **ASA** American Society of Anesthesiologists physical status classification (I: normal healthy patient - VI: a declared brain-dead patient whose organs are being removed for donor purposes), **BDRS** Blessed Dementia Rating (scores 0-28, cut-off for impairment > 4), **CCI** Charlson Comorbidity Index (scores 0-37), **CIRS-G** Cumulative Illness Rating Scale–Geriatrics (scores 0-56), **CSI** Clinical severity of illness (1 (mild) - 9 (moribund)), **GCS** Glasgow Coma Scale (scores 3-15; scores 3-8 = coma), **GDS-15** 15-item Geriatric Depression Scale (≥6 = suggests depression and need for assessment, ≥11 = depression/severe depression), **ISS** Injury Severity Score (scores 1-75), **MEWS** Modified Early Warning Score (score ≥5 is statistically linked to increased likelihood of death or admission to an intensive care unit), **RIFLE** (Risk, Injury, Failure, Loss of kidney function, and End-stage kidney disease categories), **SAPS 3** Simplified acute physiology score (scores 0-217), **SOFA** Sepsis-related Organ Failure Assessment (scores 0 to 24). **Other abbreviations:** **ADL** activities of daily living, **AF** atrial fibrillation, **Ca** cancer, **CAM** Confusion Assessment Method, **CAM-ICU** Confusion Assessment Method for the Intensive Care Unit, **CI** cognitive impairment, **CKD** Chronic Kidney Disease, **COPD** chronic obstructive pulmonary disease, **CVD** cardiovascular disease, **CVA** cerebrovascular accident, **DM** diabetes mellitus, **ED** Emergency Department, **GI** gastrointestinal, **HF** heart failure, **hrs** hours, **HT** hypertension, **I/C** intervention/control, **ICU** intensive care unit, **IHD** ischaemic heart disease, **IQR** interquartile range, **MI** Myocardial Infarction, **MMSE** Mini-Mental State Examination, **NR** not reported, **O/A** on admission, **OP** osteoporosis, **PVD** peripheral vascular disease, **RASS** Richard Agitation Sedation Scale, **RF** renal failure, **SD** standard deviation, **SICU** surgical intensive care unit, **TIA** transient ischemic attack, **UTI** urinary tract infection.

Figure 3: Study approaches to people requiring palliative care

NB: Combined percentages do not add up to 100% as studies simultaneously excluded and reported people requiring palliative care.

Effectiveness outcomes for people requiring palliative care

Overall, 20 studies reported that the intervention was effective according to at least one primary outcome (69%)^{20, 38-42, 44-46, 51, 54-57, 59-64}

The one study explicitly focused on preventing delirium in people receiving palliative care (patients with terminal cancer) tested three primarily nurse-delivered intervention components for orientation, informing family, and assessment of medication risk factors plus querying physicians about changes to medication and found no statistically significant difference in delirium incidence ($p=0.66$, OR 0.94), severity or duration between intervention and control sites.⁴⁹

Outcomes of four participant groups that we identified as requiring palliative care were as follows:

A delirium prevention RCT of a geriatric liaison intervention of comprehensive assessment, cognitive and physical activity, hearing, vision and nutrition for frail elderly cancer patients undergoing surgery for a solid tumour reported no significant difference between delirium incidence in intervention and control groups (9.4% vs. 14.3%, OR: 0.63, 95% CI: 0.29–1.35).⁵³

A delirium prevention RCT of geriatric consultation and multiple care components (Table 3) performed analysis for the sub-group of people with dementia within a perioperative hip fracture population, reporting a non-statistically significant reduction in delirium incidence in the intervention group compared with control ($n=13$ (62%) vs. $n=20$ (69%) $p=0.6$).⁴²

A delirium prevention before-after study of orientation and communication strategies for patients with dementia in a French acute geriatric unit reported a 66% relative risk reduction of delirium in the after cohort (RRR 0.34 IC 95% 0.15–0.78).⁵⁷

A delirium prevention before-after study of delirium/cognitive screening, comprehensive assessment, cognitive and physical activity, nutrition, falls prevention, medication review and staff education in frail elderly surgical patients found no statistically significant

1
2
3
4 difference in incidence of delirium (11% vs 10% $p=0.945$) or cognitive decline (15% vs 12%
5
6 $p=0.431$) in the before group compared with the after.⁵²
7

8 Due to heterogeneity of study interventions and measures (Table 2), we did not conduct
9
10 meta-analysis of these effectiveness outcomes.
11
12

13 **Types of outcomes measured in the included studies**

14
15 Twenty-nine different outcomes were measured overall (Figure 4). There were nine
16
17 different primary outcomes, the most frequent was delirium incidence (n=21).^{20, 38, 39, 41-46,}
18
19 ^{48, 49, 52, 53, 56-61, 63, 64} Of the 22 different secondary outcomes, the most often reported was
20
21 length of ICU or hospital stay (n=16).^{38-43, 45, 46, 48, 50, 51, 53-56, 62}
22

23 **Figure 4: Types and rates of outcomes measured**

24 **Feasibility outcomes**

25 **Intervention components**

26
27 The interventions consisted of 24 components overall, varying in type and number per study
28
29 (1-15) (M=6.3 SD ± 4) (Table 3). The type and number of strategies in each component also
30
31 varied (data not reported here).
32
33

34
35 Eighteen multicomponent *prevention* interventions most often included strategies for:
36
37 cognitive activity (n=15);^{20, 39, 41, 43-45, 49, 52, 53, 56, 57, 59, 61, 63, 64} physical activity (n=9);^{20, 39, 43, 45, 52,}
38
39 ^{53, 59, 63, 64} hearing (n=9);^{20, 39, 41, 44, 45, 53, 56, 61, 63} vision (n=9);^{20, 39, 41, 44, 45, 53, 56, 61, 63} sleep-wake
40
41 cycle preservation (n=8);^{20, 40, 41, 45, 59, 61, 63, 64} nutrition (n=8);^{41, 44, 45, 52, 53, 59, 61, 63} staff
42
43 education;^{39-41, 52, 59, 61, 63, 64} and hydration (n=7).^{20, 41, 44, 45, 59, 61, 63} These components largely
44
45 correspond to those of the original HELP study.²⁰
46
47

48 Components of six delirium *prevention and treatment* interventions were more diverse
49
50 (M=8), with staff education (n=4),^{38, 48, 54, 55} staff changes (n=3)^{48, 54, 55} and strategies for pain
51
52 (n=3) most often included.^{42, 48, 55}
53

54 The most common components of three delirium *treatment* interventions were cognitive
55
56 activity (n=3);^{50, 51, 62} physical activity (n=3);^{50, 51, 62} hearing (n=3);^{50, 51, 62} vision (n=3);^{50, 51, 62}
57
58
59
60

1
2
3
4 sleep-wake cycle preservation (n=2);^{50, 62} and environmental changes (e.g. adjustments to
5 lighting and noise) (n=2).^{51, 62}
6
7

8 Overall, multicomponent interventions also often included: family involvement (e.g.
9 information/education about delirium, increased family presence) (n=9);^{38, 45, 49-51, 56, 58, 59, 61}
10 environmental strategies (n=9);^{38, 40, 42, 45, 50, 51, 61, 62, 64} and comprehensive assessment
11 (n=8).^{42, 46, 50-53, 55, 61} Pharmacological strategies (e.g. medication reviews and alerts,
12 protocols for pain and sedation strategies) were present in half of the multicomponent
13 interventions (n=12).^{38, 40-42, 45, 48, 49, 52, 55, 59, 61, 64}
14
15
16
17
18
19

20 The three *single component* interventions were blood transfusion,⁶⁵ bright light therapy⁶⁰
21 and earplugs at night.⁴⁷
22
23

24 The five studies that reported effectiveness outcomes for our sample of interest addressed:
25 cognitive activity,^{42, 49, 52, 53, 57} physical activity,^{42, 52, 53} comprehensive assessment,^{42, 52, 53}
26 hearing,^{42, 53} vision,^{42, 53} nutrition,^{42, 52, 53} pharmacological strategies,^{42, 49, 52} environmental
27 strategies^{42, 57} family involvement,⁴⁹ staff changes,⁴² pain,⁴² medical complications,⁴²
28 oxygen,⁴² staff education,⁵² falls prevention⁵² delirium/cognitive screening⁵² and
29 bladder/bowel function⁴² (Figure 5).
30
31
32
33
34
35

36 **Intervention delivery**

37
38 Specialist geriatric clinicians or teams led almost half of the interventions, as either
39 consultants or directly (n=12).^{20, 39, 42, 45, 46, 50-53, 55, 62, 63} Interventions were delivered by
40 interdisciplinary teams (n=5),^{20, 38, 45, 46, 55} physicians and nurses (n=3),⁴⁹⁻⁵¹ nurses alone
41 (n=3),^{47, 48, 61} physician, nurse and physiotherapists (n=2),^{59, 64} family members (n=2),^{56, 58}
42 and physiotherapists and allied health assistants (n=1).⁴³ Volunteers delivered part or all of
43 the intervention in four studies.^{20, 41, 44, 52} Three studies did not report who delivered the
44 intervention, including two single component interventions.^{57, 60, 65} Most studies (n=25)
45 wholly or partly tailored components and strategies to patients' individual needs (Table 3).
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

(Insert Table 3 Intervention characteristics here)

Abbreviations: **ICU** Intensive Care Unit, **NR** Not reported, **OA** On admission, **OT** Occupational therapist, **PT** physiotherapist **Component codes:** **C** Cognitive activity, **E** Physical activity, **H** Hearing, **V** Vision, **P** Sleep-wake cycle preservation, **W** Hydration, **S** Staff education, **F** Family involvement, **N** Nutrition, **Pa** Pain, **O** Oxygen, **L** Falls prevention, **G** Staff changes, **B** Bladder/bowel, **J** Environment/lighting/noise, **CA** Comprehensive assessment, **BT** Blood transfusion, **Z** Address medical complications, **PE** Patient education, **Ph** Physiological monitoring, **K** Pharmacological

Figure 5: Types and rates of intervention components, including for sample of interest

Code: **P** Delirium prevention studies, **P & T** Combined delirium prevention and treatment studies, **T** Delirium treatment studies, **S of I** Sample of interest

Adherence

Eighteen studies reported adherence to intervention strategies, using different methods and levels of detail (Supplementary Table 1). Almost all reported less than 100% adherence, with the exception reporting 100% geriatric nurse compliance with the “semi-structured protocol”.⁶² The palliative care unit study reported 89.7% overall adherence to the study protocol.⁴⁹ Three studies compared adherence to intervention strategies with that for control participants and reported that usual care sometimes mirrored some intervention strategies.^{46, 50, 55} Two studies reported reasons for non-adherence: pharmacological sedation, coma and absence of a relative in the palliative care unit study;⁴⁹ and patient refusal or unavailability, staff member unavailability, and medical contraindications in the original HELP study.²⁰

Study modifications

Two studies modified the intervention in the pilot phase, including changes in staff education and practice change materials,³⁹ and providing study information to family on admission rather than waiting until the researcher was present.⁵⁸ The palliative care unit study modified the primary outcome measure by not substituting the CRS for the CAM, due to CAM completion delays and poor completion rates (39% of participants) caused by “the challenge of conducting the baseline CAM structured interview in the last days or hours of life”.⁴⁹

Acceptability outcomes

Adverse effects

No study reported systematic processes of attribution of adverse effects to the intervention. Ten studies reported adverse events related to hospitalisation rather than the intervention^{38, 44, 51-53, 55, 56, 62, 63} (Figure 4 and Supplementary Table 2). No study reported statistically significant increased adverse events for intervention participants. Two simply stated that the intervention had no adverse effects.^{20, 43} A RCT,⁵⁵ a controlled trial⁶³ and a before-after study⁶² reported statistically significant reductions in some adverse events for intervention participants: falls,⁵⁵ physical restraint,^{62, 63} pressure ulcers,^{55, 62} infection,^{55, 62} and sleeping problems.⁵⁵

Patient, family, clinician and volunteer subjective experiences

No study reported patients' subjective experiences of intervention (Supplementary Table 3). A survey of family caregiver's satisfaction with intervention and care for delirious patients within one study found most experienced moderate to high levels of distress (72%), believed that the environment and intervention helped the patient's recovery (86.6% and 83.5%, respectively) and that the three most useful strategies were activity, reality orientation and thrice-daily mobilisation.⁶² A before-after study of older medical and surgical inpatients interviewed patients, volunteers, nurses and physicians and analysed meeting minutes to assess intervention integration in clinical practice, **with minimal details of analysis methods and findings provided.**⁵² Four studies reported multidisciplinary and expert input during intervention development.^{39, 40, 45, 61} The RCT of nocte earplugs in the ICU stated in the discussion that some participants preferred not to use them in order to remain "in direct contact with their environment".⁴⁷ No studies reported measures of patient distress related to delirium (Figure 5).

Discussion

This review found that studies of non-pharmacological delirium interventions frequently excluded and under-characterised people requiring palliative care and subsequently their outcomes were infrequently reported. The likely reasons for these findings and how to address barriers to inclusion and report characteristics and outcomes for this patient subgroup in future research are discussed.

With regards to inclusion, this review identified a selection bias against people requiring palliative care through exclusion of people expected to die (using various prognoses and terminology) and also of those with greater acuity or severity of illness, particular diagnoses and with cognitive, sensory and/or communication impairments. These exclusions were rarely explained or justified and often seemingly arbitrary. Nor were they consistent across the studies. These exclusions represented people highly at risk of delirium.^{1, 6} This finding reflects those of other reviews that identified selection biases against people with dementia in geriatric research⁷⁶ and older people from clinical trials in oncology.⁷⁷ Reasons for the exclusions identified in our review are likely to be multifactorial. One factor may be assumptions that the interventions were not appropriate, possible or likely to be effective for people requiring palliative care, as indicated by the two studies which justified the exclusion of patients with a terminal condition and requiring comfort care as that the intervention would be unlikely to benefit them.^{41, 44} Historical views that people requiring palliative care are separate from the wider hospital population, rather than universally and always within it, may also have influenced some exclusion decisions.^{78, 79} There are also pragmatic challenges to ensuring informed consent and completion of study measures when patients are frail, expected to decline, and have pre-existing cognitive and sensory impairments. However, these challenges may be overcome through a variety of research strategies, such as proxy, opt out and advance consent methods, early contact, tailoring of interventions and messaging to patients and family, adequate research resources and governance, and cluster designs.^{26, 27, 80, 81} Outcomes such as days alive without delirium or coma may promote the inclusion of more severely ill patients in future delirium research, because consciousness is pre-requisite for delirium measurement.^{40, 82} Brief and

1
2
3
4 observational delirium diagnostic tools that can be validly used with patients with cognitive,
5 sensory and communication impairment are also available.⁸³⁻⁸⁶
6
7

8
9 Despite most studies excluding certain groups of people likely to require palliative care,
10 unsurprisingly we identified that they were indeed present, despite under-reporting of their
11 characteristics and outcomes. One reason it was difficult to retrospectively distinguish this
12 sub-group for the purposes of ascertaining their outcomes in this review was that the
13 studies reported participants' diagnoses, illness severity, and mortality using different
14 measures, time-points, methods and degrees of detail. While identifying inpatients with
15 palliative care needs in hospital is challenging and an uncertain science, both retrospectively
16 and prospectively, there are ways it can be achieved.^{29, 87-89} In addition to the GSF PIG which
17 we applied in this review,²⁹ the Palliative Care Needs Assessment Tool⁹⁰ and the Supportive
18 and Palliative Care Indicators Tool (SPICT)⁹¹ can be used to identify which patients are at
19 risk of deteriorating and dying in hospital. Other studies suggest identifying patients with
20 palliative care needs through the presence of life-threatening illness coupled with receipt
21 and acceptance of care focused on supporting quality of life,⁸⁷ or retrospectively through the
22 use of death registration data.⁸⁹ Such methods could be used in delirium intervention
23 research in settings where mortality is high, such as intensive care units, where existing
24 illness severity measures focus on acuity and intensity of intervention and, despite having
25 some prognostic utility, were never designed to identify a patient's need for palliative
26 care.^{67, 74, 92}
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

42 Better characterisation of people requiring palliative care in studies of non-pharmacological
43 interventions for delirium would help to build evidence of their outcomes in two ways.
44
45

46 Firstly, it would allow sub-group analysis to be performed and reported in future studies in
47 any setting. This is important because the question of whether delirium can be prevented
48 and treated through non-pharmacological means was not definitively determined for this
49 population by this review. The multi-site palliative care unit study had the greatest number
50 of participants and reported good overall adherence to the study protocol;⁴⁹ however, the
51 only core care domain²¹ was cognitive activity, consisting of nurses orientating patients to
52 time, person and place each shift. Planned use of the CAM was not accomplished and the
53
54
55
56
57
58
59
60

1
2
3
4 substituted screening tool⁹³ was not a validated tool for delirium diagnosis or severity. All
5 participants had 'terminal' cancer and died within 90 days of admission and thereby were
6 not fully representative of people requiring inpatient palliative care, including in specialist
7 units⁴⁹, not all of whom will die within three months. Only one geriatric unit study of people
8 with dementia reported reduced delirium incidence; however, the report lacked detail,
9 including of analysis methods.⁵⁷ The remaining three studies were focused on frail elderly
10 patients with cancer, dementia, hip fracture and comorbidity following surgery,^{42, 52, 53} a
11 level of intervention and delirium risk that many people requiring palliative care would not
12 undergo. Lastly, we found no studies focused on non-pharmacological interventions to treat
13 delirium in people requiring palliative care, highlighting the particular need to research this
14 challenging area of clinical care.

15
16
17
18
19
20
21
22
23
24
25 Secondly, better recognition of people requiring palliative care in delirium research would
26 heighten awareness of their needs, which in turn would instigate consideration of outcomes
27 that are most meaningful to them. Reducing delirium incidence, duration and severity,
28 length of admission and mortality were the foci of the included studies. While worthwhile
29 aims at any point in the illness trajectory, more person-centred outcomes, such as reduction
30 of delirium related distress, and improved patient and family caregiver experiences of
31 decision making, respect and dignity, and quality of life, were almost completely absent.^{5, 94}
32 Measuring these additional outcomes, which are highly valued by patients and family
33 caregivers,^{95, 96} would enrich this field of research and is achievable through both
34 quantitative and qualitative methods. Adverse effects of intervention will also be important
35 to systematically measure in future trials, given that patients with cognitive impairment
36 have increased vulnerability to harm in hospital.⁹⁷

37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Lastly, this review allowed an interpretation that many intervention components are
feasible for people requiring palliative care by virtue of their delivery to elderly, frail and/or
critically ill patients with and without delirium in the included studies. However, we suggest
that studies measuring the feasibility of components of the original HELP intervention²⁰ that
were subsequently found to be effective for other older hospitalised patients i.e. those
targeting cognitive and physical activity, sleep, hearing, vision and hydration,²¹⁻²³ are
necessary next steps within delirium research programs in specialist palliative care settings.

Limitations and strengths

A limitation of this review was that only English language studies were included. Another limitation was that our retrospective identification of people requiring palliative care was not sufficient to definitively determine outcomes for this patient sub-group; therefore this review makes only recommendations for future research, not clinical practice. A strength of this review was our systematic approach adhering to PRISMA.³⁰

Conclusion

This review found that studies of non-pharmacological delirium interventions have excluded, poorly characterised and infrequently reported outcomes for people requiring palliative care. Based on these findings, we suggest new approaches to generate evidence for delirium interventions in this important patient sub-group. First, randomised controlled trials of non-pharmacological interventions to prevent and treat delirium are needed in specialist palliative care settings, with feasibility studies required before trials of effectiveness. Interventions should be based on those implemented and proven effective for older patients, especially those who were frail and severely ill with comorbidity. Second, broad inclusion criteria, justified exclusions, and tailoring in future inpatient trials of non-pharmacological delirium interventions would promote representative study populations. Third, systematic characterisation of the sub-groups of people requiring palliative care would allow their specific outcomes to be reported. Last, additional outcomes related to patient and family subjective experience, goals of care and quality of life would enhance the relevance of delirium research in inpatient settings where people are cared for at the end of life.

Acknowledgements

We thank Dr E. Wesley Ely for his insights in ICU illness severity measures and constructive feedback on the initial manuscript, and a University of Technology Sydney Health Futures Research Development Fund grant which supported this research.

References

1. National Institute for Health and Clinical Excellence (NICE) National Clinical Guideline centre. Delirium: diagnosis, prevention and management, <https://www.nice.org.uk/guidance/cg103> (2010, accessed March 2 2019).
2. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5)*. Arlington, VA: American Psychiatric Publisher, 2013.
3. O'Malley G, Leonard M, Meagher D, et al. The delirium experience: a review. *Journal of Psychosomatic Research* 2008; 65: 223-228. 2008/08/19. DOI: 10.1016/j.jpsychores.2008.05.017.
4. Van Rompaey B, Van Hoof A, van Bogaert P, et al. The patient's perception of a delirium: A qualitative research in a Belgian intensive care unit. *Intensive and Critical Care Nursing* 2016; 32: 66-74. DOI: 10.1016/j.iccn.2015.03.002.
5. Finucane AM, Lugton J, Kennedy C, et al. The experiences of caregivers of patients with delirium, and their role in its management in palliative care settings: an integrative literature review. *Psycho-Oncology* 2017; 26: 291-300. DOI: 10.1002/pon.4140.
6. Hosie A, Davidson PM, Agar M, et al. Delirium prevalence, incidence, and implications for screening in specialist palliative care inpatient settings: A systematic review. *Palliative Medicine* 2013; 27: 486-498. DOI: 10.1177/0269216312457214.
7. Australian Commission on Safety and Quality in Health Care. Evidence for the safety and care of patients with a cognitive impairment in acute care settings: a rapid review. Sydney: ACSQHC, 2013.
8. Leslie DL, Marcantonio ER, Zhang Y, et al. One-year health care costs associated with delirium in the elderly population. *Archives Internal Medicine* 2008; 168: 27-32. 2008/01/16. DOI: 10.1001/archinternmed.2007.4.
9. Ryan DJ, O'Regan NA, Caoimh RÓ, et al. Delirium in an adult acute hospital population: predictors, prevalence and detection. *BMJ Open* 2013; 3. DOI: 10.1136/bmjopen-2012-001772.
10. Canadian Coalition for Seniors' Mental Health. Guideline on the Assessment and Treatment of Delirium in Older Adults at the End of Life. Toronto: Canadian Coalition for Seniors' Mental Health (CCSMH), 2010.
11. Australian Commission on Safety and Quality in Health Care. A better way to care: Safe and high-quality care for patients with cognitive impairment (dementia and delirium) in hospital - Actions for consumers. Sydney: ACSQHC, 2014.
12. Barr J, Fraser GL, Puntillo K, et al. Clinical practice guidelines for the management of pain, agitation, and delirium in adult patients in the intensive care unit. *Critical Care Medicine* 2013; 41: 263-306. DOI: 10.1097/CCM.0b013e3182783b72.
13. Bush SH, Marchington KL, Agar M, et al. Quality of clinical practice guidelines in delirium: a systematic appraisal. *BMJ Open* 2017; 7. DOI: 10.1136/bmjopen-2016-013809.
14. Australian Commission on Quality and Safety in Healthcare. Delirium Clinical Care Standard, <https://www.safetyandquality.gov.au/our-work/clinical-care-standards/delirium-clinical-care-standard/> (2017, accessed March 9 2019).
15. iDelirium, <http://www.idelirium.org/> (2018, accessed March 9 2019).
16. Lawlor PG, Davis DHJ, Ansari M, et al. An Analytic Framework for Delirium Research in Palliative Care Settings: Integrated Epidemiological, Clinician-Researcher and Knowledge User Perspectives. *Journal of Pain and Symptom Management* 2014; 48: 159-175. DOI: 10.1016/j.jpainsymman.2013.12.245.
17. Palliative Care Expert Group. *Therapeutic Guidelines: Palliative Care (Version 4)*. Melbourne: Therapeutic Guidelines Ltd, 2016.

18. Abraha I, Rimland JM, Trotta F, et al. Non-Pharmacological Interventions to Prevent or Treat Delirium in Older Patients: Clinical Practice Recommendations The SENATOR-ONTOP Series. *Journal of Nutrition, Health and Aging* 2016; 20: 927-936. 2016/10/30. DOI: 10.1007/s12603-016-0719-9.
19. Devlin JW, Skrobik Y, Gélinas C, et al. Clinical Practice Guidelines for the Prevention and Management of Pain, Agitation/Sedation, Delirium, Immobility, and Sleep Disruption in Adult Patients in the ICU. *Critical Care Medicine* 2018; 46: e825-e873. DOI: 10.1097/ccm.0000000000003299.
20. Inouye SK, Bogardus STJ, Charpentier PA, et al. A Multicomponent Intervention to Prevent Delirium in Hospitalized Older Patients. *New England Journal of Medicine* 1999; 340: 669-676. DOI: 10.1056/nejm199903043400901.
21. Hshieh TT, Yue J, Oh E, et al. Effectiveness of multicomponent nonpharmacological delirium interventions: a meta-analysis. *JAMA Internal Medicine* 2015; 175: 512-520. 2015/02/03. DOI: 10.1001/jamainternmed.2014.7779.
22. Siddiqi N, Harrison JK, Clegg A, et al. Interventions for preventing delirium in hospitalised non-ICU patients. *Cochrane Database of Systematic Reviews* 2016. DOI: 10.1002/14651858.CD005563.pub3.
23. Abraha I, Trotta F, Rimland JM, et al. Efficacy of Non-Pharmacological Interventions to Prevent and Treat Delirium in Older Patients: A Systematic Overview. The SENATOR project ONTOP Series. *PLOS ONE* 2015; 10: e0123090. DOI: 10.1371/journal.pone.0123090.
24. Bush SH, Lawlor PG, Ryan K, et al. Delirium in adult cancer patients: ESMO Clinical Practice Guidelines†. *Annals of Oncology* 2018; 29: iv143–iv165. DOI: 10.1093/annonc/mdy147.
25. Lawlor PG, Rutkowski NA, MacDonald AR, et al. A scoping review to map empirical evidence regarding key domains and questions in the clinical pathway of delirium in palliative care. *Journal of Pain and Symptom Management* 2018; 57: 661-681. DOI: 10.1016/j.jpainsymman.2018.12.002.
26. Agar M, Lawlor PG, Quinn S, et al. Efficacy of oral risperidone, haloperidol, or placebo for symptoms of delirium among patients in palliative care: A randomized clinical trial. *JAMA Internal Medicine* 2017; 177: 34-42. DOI: 10.1001/jamainternmed.2016.7491.
27. Hui D, Frisbee-Hume S, Wilson A, et al. Effect of Lorazepam With Haloperidol vs Haloperidol Alone on Agitated Delirium in Patients With Advanced Cancer Receiving Palliative Care: A Randomized Clinical Trial. *JAMA* 2017; 318: 1047-1056. DOI: 10.1001/jama.2017.11468.
28. To THM, Greene AG, Agar M, et al. A point prevalence survey of hospital inpatients to define the proportion with palliation as the primary goal of care and the need for specialist palliative care. *Internal Medicine Journal* 2011; 41: 430-433. DOI: 10.1111/j.1445-5994.2011.02484.x.
29. National Gold Standards Framework Centre in End of Life Care. Gold Standards Framework Proactive Identification Guidance (GSF PIG), <http://www.goldstandardsframework.org.uk/pig> (2016).
30. Moher D, Liberati A, Tetzlaff J, et al. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med* 2009; 6: e1000097. DOI: 10.1371/journal.pmed.1000097.
31. American Psychiatric Association. DSM-III Diagnostic and Statistical Manual of Mental Disorders. In: Association AP, (ed.). 3rd ed. Washington DC: American Psychiatric Association, 1980.
32. Covidence systematic review software, Veritas Health Innovation. Melbourne, Australia. Available at www.covidence.org.
33. Hoffmann TC, Glasziou PP, Boutron I, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ* 2014; 348. DOI: 10.1136/bmj.g1687.
34. Healthcare Improvement Scotland. Scottish Intercollegiate Guidelines Network Methodology Checklists, <http://www.sign.ac.uk/checklists-and-notes.html> (accessed June 7 2018).
35. Review Manager (RevMan) [Computer program]. Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

36. Hamzhepour H, Valiee S, Azad Majedi M, et al. The Effect of Care Plan Based on Roy Adaptation Model on the Incidence and Severity of Delirium in Intensive Care Unit Patients: A Randomised Controlled Trial. *Journal of Clinical and Diagnostic Research* 2018; 12: LC21-LC25. November 1,2018. DOI: 10.7860/jcdr/2018/36366.12256.
37. Johnson K, Fleury J and McClain D. Music intervention to prevent delirium among older patients admitted to a trauma intensive care unit and a trauma orthopaedic unit. *Intensive and Critical Care Nursing* 2018; 47: 7-14. DOI: 10.1016/j.iccn.2018.03.007.
38. Bryczkowski SB, Lopreiato MC, Yonclas PP, et al. Delirium prevention program in the surgical intensive care unit improved the outcomes of older adults. *Journal of Surgical Research* 2014; 190: 280-288. 2014/03/29. DOI: 10.1016/j.jss.2014.02.044.
39. Holt R, Young J and Heseltine D. Effectiveness of a multi-component intervention to reduce delirium incidence in elderly care wards. *Age and Ageing* 2013; 42: 721-727. DOI: 10.1093/ageing/aft120.
40. Kamdar BB, King LM, Collop NA, et al. The effect of a quality improvement intervention on perceived sleep quality and cognition in a medical ICU. *Critical Care Medicine* 2013; 41: 800. DOI: 10.1097/CCM.0b013e3182746442.
41. Zaubler TS, Murphy K, Rizzuto L, et al. Quality improvement and cost savings with multicomponent delirium interventions: replication of the hospital elder life program in a community hospital. *Psychosomatics* 2013; 54: 219-226. DOI: 10.1016/j.psym.2013.01.010.
42. Marcantonio ER, Flacker JM, Wright RJ, et al. Reducing delirium after hip fracture: a randomized trial. *Journal of the American Geriatrics Society* 2001; 49: 516-522. DOI: 10.1046/j.1532-5415.2001.49108.x.
43. Jeffs KJ, Berlowitz DJ, Grant S, et al. An enhanced exercise and cognitive programme does not appear to reduce incident delirium in hospitalised patients: a randomised controlled trial. *BMJ Open* 2013; 3: e002569. DOI: 10.1136/bmjopen-2013-002569.
44. Caplan GA and Harper EL. Recruitment of volunteers to improve vitality in the elderly: the REVIVE study. *Internal Medicine Journal* 2007; 37: 95-100. DOI: 10.1111/j.1445-5994.2007.01265.x.
45. Wong Tin Niam DM, Bruce JJ and Bruce DG. Quality project to prevent delirium after hip fracture. *Australasian Journal on Ageing* 2005; 24: 174-177. DOI: 10.1111/j.1741-6612.2005.00108.x.
46. Deschodt M, Braes T, Flamaing J, et al. Preventing delirium in older adults with recent hip fracture through multidisciplinary geriatric consultation. *Journal of the American Geriatrics Society* 2012; 60: 733-739. DOI: 10.1111/j.1532-5415.2012.03899.x.
47. Van Rompaey B, Elseviers MM, Drom W, et al. The effect of earplugs during the night on the onset of delirium and sleep perception: a randomized controlled trial in intensive care patients. *Critical Care* 2012; 16: R73. DOI: 10.1016/j.iccn.2015.03.002.
48. Milisen K, Foreman MD, Abraham IL, et al. A Nurse-Led Interdisciplinary Intervention Program for Delirium in Elderly Hip-Fracture Patients. *Journal of the American Geriatrics Society* 2001; 49: 523-532. DOI: 10.1046/j.1532-5415.2001.49109.x.
49. Gagnon P, Allard P, Gagnon B, et al. Delirium prevention in terminal cancer: assessment of a multicomponent intervention. *Psycho-Oncology* 2012; 21: 187-194. DOI: 10.1002/pon.1881.
50. Cole MG, McCusker J, Bellavance F, et al. Systematic detection and multidisciplinary care of delirium in older medical inpatients: a randomized trial. *Canadian Medical Association Journal* 2002; 167: 753-759.
51. Cole MG, Primeau FJ, Bailey RF, et al. Systematic intervention for elderly inpatients with delirium: a randomized trial. *Canadian Medical Association Journal* 1994; 151: 965-970.
52. Bakker FC, Persoon A, Bredie SJ, et al. The CareWell in Hospital program to improve the quality of care for frail elderly inpatients: results of a before-after study with focus on surgical patients. *American Journal of Surgery* 2014; 208: 735-746. DOI: 10.1016/j.amjsurg.2014.04.009.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
53. Hempenius L, Slaets JPJ, van Asselt D, et al. Outcomes of a geriatric liaison intervention to prevent the development of postoperative delirium in frail elderly cancer patients: Report on a multicentre, randomized, controlled trial. *PloS One* 2013; 8: e64834. DOI: 10.1371/journal.pone.0064834.
54. Lundström M, Edlund A, Karlsson S, et al. A multifactorial intervention program reduces the duration of delirium, length of hospitalization, and mortality in delirious patients. *Journal of the American Geriatrics Society* 2005; 53: 622-628. DOI: 10.1111/j.1532-5415.2005.53210.x.
55. Lundström M, Olofsson B, Stenvall M, et al. Postoperative delirium in old patients with femoral neck fracture: a randomized intervention study. *Aging Clinical and Experimental Research* 2007; 19: 178-186.
56. Martinez FT, Tobar C, Beddings CI, et al. Preventing delirium in an acute hospital using a non-pharmacological intervention. *Age and Ageing* 2012; 41: 629-634. DOI: 10.1093/ageing/afs060.
57. Andro M, Comps E, Estivin S, et al. Prevention of delirium in demented hospitalized patients. *European Journal of Internal Medicine* 2012; 23: 124-125. 2012/01/31. DOI: 10.1016/j.ejim.2011.05.011.
58. Black P, Boore JRP and Parahoo K. The effect of nurse-facilitated family participation in the psychological care of the critically ill patient. *Journal of Advanced Nursing* 2011; 67: 1091-1101. DOI: 10.1111/j.1365-2648.2010.05558.x.
59. Bo M, Martini B, Ruatta C, et al. Geriatric ward hospitalization reduced incidence delirium among older medical inpatients. *American Journal of Geriatric Psychiatry* 2009; 17: 760-768. DOI: 10.1097/JGP.0b013e3181a315d5.
60. Taguchi T, Yano M and Kido Y. Influence of bright light therapy on postoperative patients: a pilot study. *Intensive & Critical Care Nursing* 2007; 23: 289-297. DOI: 10.1016/j.iccn.2007.04.004.
61. Moon K-J and Lee S-M. The effects of a tailored intensive care unit delirium prevention protocol: A randomized controlled trial. *International Journal of Nursing Studies* 2015; 52: 1423-1432. DOI: 10.1016/j.ijnurstu.2015.04.021.
62. Chong MS, Chan M, Tay L, et al. Outcomes of an innovative model of acute delirium care: the Geriatric Monitoring Unit (GMU). *Clinical Interventions in Aging* 2014; 9: 603-612. DOI: 10.2147/CIA.S60259.
63. Vidan MT, Sanchez E, Alonso M, et al. An intervention integrated into daily clinical practice reduces the incidence of delirium during hospitalization in elderly patients. *Journal of the American Geriatrics Society* 2009; 57: 2029-2036. DOI: 10.1111/j.1532-5415.2009.02485.x.
64. Patel J, Baldwin J, Bunting P, et al. The effect of a multicomponent multidisciplinary bundle of interventions on sleep and delirium in medical and surgical intensive care patients. *Anaesthesia* 2014; 69: 540-549. DOI: 10.1111/anae.12638.
65. Gruber-Baldini AL, Marcantonio E, Orwig D, et al. Delirium outcomes in a randomized trial of blood transfusion thresholds in hospitalized older adults with hip fracture. *Journal of the American Geriatrics Society* 2013; 61(8): 1286-95. DOI: 10.1111/jgs.12396.
66. Charlson ME, Pompei P, Ales K, et al. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *Journal of Chronic Disease* 1987; 40: 373-383.
67. Knaus WA, Draper EA, Wagner DP, et al. APACHE II: a severity of disease classification system. *Critical Care Medicine* 1985; 13: 818-829.
68. Miller MD, Paradis CF, Houck PR, et al. Rating chronic medical illness burden in geropsychiatric practice and research: application of the Cumulative Illness Rating Scale. *Psychiatry Research* 1992; 41: 237-248. 1992/03/01. DOI: 10.1016/0165-1781(92)90005-N.
69. Saklad M. Grading of patients for surgical procedures. *Anesthesiology: The Journal of the American Society of Anesthesiologists* 1941; 2: 281-284.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
70. Blessed G, Tomlinson BE and Roth M. The Association Between Quantitative Measures of Dementia and of Senile Change in the Cerebral Grey Matter of Elderly Subjects. *British Journal of Psychiatry* 1968; 114: 797-811. DOI: 10.1192/bjp.114.512.797.
71. Subbe CP, Kruger M, Rutherford P, et al. Validation of a modified Early Warning Score in medical admissions. *QJM: An International Journal of Medicine* 2001; 94: 521-526. DOI: 10.1093/qjmed/94.10.521.
72. Vincent J-L, Moreno R, Takala J, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. *Intensive Care Medicine* 1996; 22: 707-710.
73. Bellomo R, Ronco C, Kellum JA, et al. Acute renal failure—definition, outcome measures, animal models, fluid therapy and information technology needs: the Second International Consensus Conference of the Acute Dialysis Quality Initiative (ADQI) Group. *Critical Care* 2004; 8: R204. DOI: 10.1186/cc2872.
74. Metnitz PGH, Moreno RP, Almeida E, et al. SAPS 3—From evaluation of the patient to evaluation of the intensive care unit. Part 1: Objectives, methods and cohort description. *Intensive Care Medicine* 2005; 31: 1336-1344. DOI: 10.1007/s00134-005-2762-6.
75. Yesavage JA and Sheikh JI. 9/Geriatric depression scale (GDS) recent evidence and development of a shorter version. *Clinical Gerontologist* 1986; 5: 165-173. DOI: 10.1300/J018v05n01_09.
76. Taylor JS, DeMers SM, Vig EK, et al. The disappearing subject: exclusion of people with cognitive impairment and dementia from geriatrics research. *Journal of the American Geriatric Society* 2012; 60: 413-419. DOI: 10.1111/j.1532-5415.2011.03847.x.
77. Singh H, Kanapuru B, Smith C, et al. FDA analysis of enrollment of older adults in clinical trials for cancer drug registration: A 10-year experience by the U.S. Food and Drug Administration. *Journal of Clinical Oncology* 2017; 35: 10009-10009. DOI: 10.1200/JCO.2017.35.15_suppl.10009.
78. Hosie A, Agar M, Lobb E, et al. Improving delirium recognition and assessment for people receiving inpatient palliative care: a mixed methods meta-synthesis. *International Journal of Nursing Studies* 2017; 75: 123-129. DOI: 10.1016/j.ijnurstu.2017.07.007.
79. Knaul FM, Farmer PE, Krakauer EL, et al. Alleviating the access abyss in palliative care and pain relief—an imperative of universal health coverage: the Lancet Commission report. *The Lancet* 2017; 39(10128): 1391-1454. DOI: 10.1016/S0140-6736(17)32513-8.
80. Boland J, Currow DC, Wilcock A, et al. A systematic review of strategies used to increase recruitment of people with cancer or organ failure into clinical trials: implications for palliative care research. *Journal of Pain and Symptom Management* 2015; 49: 762-772.e765. DOI: 10.1016/j.jpainsymman.2014.09.018.
81. Hanson LC, Bull J, Wessell K, et al. Strategies to support recruitment of patients with life-limiting illness for research: the Palliative Care Research Cooperative Group. *Journal of Pain and Symptom Management* 2014; 48: 1021-1030. DOI: 10.1016/j.jpainsymman.2014.04.008.
82. Girard TD, Exline MC, Carson SS, et al. Haloperidol and Ziprasidone for Treatment of Delirium in Critical Illness. *New England Journal of Medicine* 2018; 379: 2506-2516. DOI: 10.1056/NEJMoa1808217.
83. Marcantonio ER, Ngo LH, O'Connor M, et al. 3D-CAM: Derivation and Validation of a 3-Minute Diagnostic Interview for CAM-Defined Delirium. *Annals of Internal Medicine* 2014; 161: 554-561. DOI: 10.7326/m14-0865.
84. Ely EW, Margolin R, Francis J, et al. Evaluation of delirium in critically ill patients: validation of the Confusion Assessment Method for the Intensive Care Unit (CAM-ICU). *Critical Care Medicine* 2001; 29: 1370-1379.
85. Han JH, Wilson A, Vasilevskis EE, et al. Diagnosing Delirium in Older Emergency Department Patients: Validity and Reliability of the Delirium Triage Screen and the Brief Confusion Assessment

- 1
2
3
4
5 Method. *Annals of Emergency Medicine* 2013; 62: 457-465. DOI:
6 10.1016/j.annemergmed.2013.05.003.
7 86. Hospital Elder Life Program. Confusion Assessment Method (Short CAM),
8 <http://www.hospitalelderlifeprogram.org/delirium-instruments/short-cam/> (2015, accessed April 12
9 2019).
- 10 87. Van Mechelen W, Aertgeerts B, De Ceulaer K, et al. Defining the palliative care patient: A
11 systematic review. *Palliative Medicine* 2012; 27: 197-208. DOI: 10.1177/0269216311435268.
- 12 88. Downar J, Goldman R, Pinto R, et al. The “surprise question” for predicting death in seriously
13 ill patients: a systematic review and meta-analysis. *Canadian Medical Association Journal* 2017;
14 189(13): E484-E493. DOI: 10.1503/cmaj.160775.
- 15 89. Murtagh FEM, Bausewein C, Verne J, et al. How many people need palliative care? A study
16 developing and comparing methods for population-based estimates. *Palliative Medicine* 2013; 28(1):
17 49-58. DOI: 10.1177/0269216313489367.
- 18 90. Waller A, Girgis A, Currow D, et al. Development of the Palliative Care Needs Assessment
19 Tool (PC-NAT) for use by multi-disciplinary health professionals. *Palliative Medicine* 2008; 22(8): 956-
20 964. DOI: 10.1177/0269216308098797.
- 21 91. Hight G, Crawford D, Murray SA, et al. Development and evaluation of the Supportive and
22 Palliative Care Indicators Tool (SPICT): a mixed-methods study. *BMJ Supportive & Palliative Care*
23 2014; 4(3): 285. DOI: 10.1136/bmjspcare-2013-000488.
- 24 92. Moreno RP, Metnitz PGH, Almeida E, et al. SAPS 3—From evaluation of the patient to
25 evaluation of the intensive care unit. Part 2: Development of a prognostic model for hospital
26 mortality at ICU admission. *Intensive Care Medicine* 2005; 31(10): 1345-1355. DOI: 10.1007/s00134-
27 005-2763-5.
- 28 93. Williams MA, Ward SE and Campbell EB. Confusion: Testing versus Observation. *Journal of*
29 *Gerontological Nursing* 1988; 14(1): 25-30. DOI: 10.3928/0098-9134-19880101-08.
- 30 94. Pandharipande PP and Ely EW. Humanizing the Treatment of Hyperactive Delirium in the
31 Last Days of Life. *JAMA* 2017; 318(11): 1104-1105. DOI: 10.1001/jama.2017.11466.
- 32 95. Virdun C, Lockett T, Davidson PM, et al. Dying in the hospital setting: A systematic review of
33 quantitative studies identifying the elements of end-of-life care that patients and their families rank
34 as being most important. *Palliative Medicine* 2015; 29(9): 774-796. 2015/04/30. DOI:
35 10.1177/0269216315583032.
- 36 96. Virdun C, Lockett T, Lorenz K, et al. Dying in the hospital setting: A meta-synthesis identifying
37 the elements of end-of-life care that patients and their families describe as being important.
38 *Palliative Medicine* 2016; 31(7): 687-601. DOI: 10.1177/0269216316673547.
- 39 97. Reynish EL, Hapca SM, De Souza N, et al. Epidemiology and outcomes of people with
40 dementia, delirium, and unspecified cognitive impairment in the general hospital: prospective cohort
41 study of 10,014 admissions. *BMC Medicine* 2017; 15: 140. 2017/07/28. DOI: 10.1186/s12916-017-
42 0899-0.
- 43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 Study characteristics

Author, year	Country	Aim	Design	Service/setting	Component	Risk of bias ^Δ
Moon 2015 ⁶¹	Korea	P	RCT	105-bed medical/surgical ICU, general 1049-bed hospital	M	2,6,7,9,10
Bakker 2014 ⁵²	Netherlands	P	BA	Frailty service, two surgical + one internal medicine department, university MC	M	6,7,9,10
Bryczkowski 2014 ³⁸	US	P, T	BA	14-bed surgical ICU, Level I academic urban trauma centre	M	6,10
Chong 2014 ⁶²	Singapore	T	BA + Control	Geriatric Monitoring Unit for acute delirium	M	6,7,10
Patel 2014 ⁶⁴	UK	P	BA	24-bed adult mixed surgical/medical ICU, teaching hospital	M	6,7,10
Gruber-Baldini 2013 ⁶⁵	US & Canada	P, T	RCT	Postoperative hip fracture services in thirteen hospitals	S	2,3,6,7,9,10
Hempenius 2013 ⁵³	Netherlands	P	RCT	Solid tumour perioperative services at university MC, large teaching hospital + community hospital	M	2,3,5,6,7,10
Holt 2013 ³⁹	US	P	BA	Three specialist elderly care wards, general hospital	M	6,7,9,10
Jefferis 2013 ⁴³	Australia	P	RCT	Medical inpatients, secondary referral hospital	M	2,3,4,5,6,7,8,10
Kamdar 2013 ⁴⁰	US	P	BA	Medical ICU, tertiary academic hospital	M	1,6,9,10
Zaubler 2013 ⁴¹	US	P	BA	38-bed general medical floor, 600-bed teaching community hospital	M	7,10
Andro, 2012 ⁵⁷	France	P	BA	Acute geriatric care unit	M	6,10
Deschodt 2012 ⁴⁶	Belgium	P	CT	Perioperative hip fracture in two trauma wards, university hospital	M	6,8,9,10
Gagnon 2012 ⁴⁹	Canada	P	QE	Four hospital palliative care units + three stand-alone hospices	M	7,9
Martinez 2012 ⁵⁶	Chile	P	RCT	Internal medicine ward in an acute hospital	M	1,2,3,6,7,8,10
Van Rompaey 2012 ⁴⁷	Belgium	P	RCT	45-bed medical/surgical/cardiac surgical ICU, 625-bed university hospital	S	2,3,5,6,9
Black 2011 ⁵⁸	Ireland	P	CTS	Seven-bed medical and surgical ICU, inner city public hospital	M	9
Bo 2009 ⁵⁹	Italy	P	CT	Medical and geriatric units, university teaching hospital	M	6,7,9
Vidan 2009 ⁶³	Spain	P	CT	Geriatric and internal medicine units, large public university hospital	M	6,7,8,9,10
Caplan & Harper 2007 ⁴⁴	Australia	P	BA	52 bed geriatric ward, tertiary referral university hospital	M	6,7
Lundström 2007 ⁵⁵	Sweden	P, T	RCT	Postoperative hip fracture service in geriatric orthopaedic + orthopaedic ward, university hospital	M	2,3,5,6,7,8,9,10
Taguchi 2007 ⁶⁰	Japan	P	PRCT	Postoperative oesophageal cancer service, university hospital	S	3,6,10

Author, year	Country	Aim	Design	Service/setting	Component	Risk of bias ^Δ
Lundström 2005 ⁵⁴	Sweden	P, T	CCT	Two hospital wards, general internal medicine department	M	1,5,6,9
Wong 2005 ⁴⁵	Australia	P	BA	Hip-fracture service in orthopaedic ward, 460-bed metropolitan teaching hospital	M	1
Cole 2002 ⁵⁰	Canada	T	RCT	Five medical units, 400-bed university-affiliated primary acute care facility	M	2,3,5,6,7,8,9,10
Marcantonio 2001 ⁴²	US	P, T	RCT	Perioperative hip fracture service, academic tertiary MC	M	2,3,5,6,7,8,9,10
Milisen 2001 ⁴⁸	Belgium	P, T	BA	Perioperative hip fracture service, emergency room + two trauma units, urban academic MC centre	M	6,10
Inouye 1999 ²⁰	US	P	CT, matched	200 bed general-medicine service, 800-bed urban teaching hospital	M	6,8,9,10
Cole 1994 ⁵¹	Canada	T	RCT	Medical inpatients of university-affiliated, primary acute 400-bed hospital	M	1,5,6,8,9,10

Code: P Prevention, T Treatment, RCT Randomised controlled trial, CTS Comparative time series, B/A Before–after study, CT Controlled trial, ICU Intensive Care Unit, NR Not reported, MC Medical Centre, M Multicomponent, PRCT Pilot RCT, QE Quasi-experimental, S Single component

^Δ Numbers signify the domains in which the study was assessed as having **low risk of bias**: 1 = Representative study sample; 2 = Concealed allocation; 3 = Random sequence generation; 4 = Blinded participants and intervention personnel; 5 = Blinded outcome assessors; 6 = Valid, reliable delirium measures; 7 = <20% sample lost to analysis; 8 = Intention to treat analysis; 9 = Confounders accounted for; 10 = Only *a priori* outcomes reported

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Table 2 Study inclusion and exclusion criteria and participant diagnoses, illness severity and mortality

Author, year	Inclusion	Exclusion	Sample size ^Δ	Age, Mean ^Δ	Male % ^{Δα}	Diagnoses % ^{Δα}	Illness severity, mean or proportion ^Δ	Mortality I vs C n (%)
Moon 2015 ⁶¹	≥ 18, ability to understand study purpose and/or provide consent independently or via proxy, ICU admission ≥48 hours	Persistent RASS - 4 or -5, severe visual and auditory problems preventing CAM-ICU, serious psychiatric or neurologic diagnosis, Korean MMSE score ≤23, [†] on isolation ward with infection; death or discharge day admitted, inability for CAM-ICU when patient violent with RASS +3 or +4	134	69.7	48	Infection 25	APACHE II 13.9	7-day in-hospital 1 (1.7%) vs 9 (14.3%)* [†] 30-day in-hospital 4 (6.7%) vs 11 (17.5%) [†]
Bakker 2014 ⁵²	≥70, admitted to surgical or medical departments > 48 hrs, at risk of delirium, falls, malnutrition, physical decline, frail [†]	Illness, language barriers, other speciality, refusal, logistic missing, delirium O/A, refusal, died in hospital, [†] died <3 months [†]	386	77	56	NR	ASA: I and II 25.5% III and IV 74.5% [†] CIRS-G comorbidity 13	During admission 5 (3%) vs 3 (2%) [†] Three-month 8 (4%) vs 11 (6%) [†]
Bryczkowski 2014 ³⁸	>50, admitted to SICU ≥ 24 hours	Moderate-severe traumatic brain injury (AIS score 3) [†] , transfer from jail or in police custody, h/o dementia. [†] Participants with unobtainable or undocumented	123	66.5	58	NR	GCS 14.6 ISS 16.5 (Trauma patients)	In-SICU 2 (3%) vs 4 (7%) ^{††}

Author, year	Inclusion	Exclusion	Sample size ^Δ	Age, Mean ^Δ	Male % ^{Δα}	Diagnoses % ^{Δα}	Illness severity, mean or proportion ^Δ	Mortality I vs C n (%)
		delirium status excluded from analysis						
Chong 2014 ⁶²	> 65, admitted to geriatric medicine department, assessed to have delirium (either O/A or incident delirium during hospital stay) according to CAM	Medical illness needing special monitoring, dangerously ill [†] coma, terminal illness, [†] uncommunicative or severe aphasia, [†] severe combative behaviour with high risk of harm, contra-indication to bright light therapy, respiratory or contact precautions, GMU admission refused by family, patient or physician-in-charge	320	84.3	33	NR	CCI 2.7 [†] Severity of Illness Index: Level 1 0.9% Level 2 84.3% Level 3 14.8% [†]	During admission 4 (1.7%) vs 0 [†] 6-month 20 (12%) [†] 12-month 14 (20.5%) (Intervention cohort only) [†]
Patel 2014 ⁶⁴	> 18, ≥1 night in ICU	Pre-existing sleep pathology, severe visual or hearing impairment, alcohol addiction or illicit drug abuse, h/o CI, [†] discharge from ICU this admission, neurosurgical, delirium during study, received	338	60.3	52	NR	APACHE II 14.6	NR

Author, year	Inclusion	Exclusion	Sample size ^Δ	Age, Mean ^Δ	Male % ^{Δα}	Diagnoses % ^{Δα}	Illness severity, mean or proportion ^Δ	Mortality I vs C n (%)
		sedative medications within 24 hrs						
Gruber-Baldini 2013 ⁶⁵	≥ 50, surgical repair of hip fracture, haemoglobin <10 g/dL <3 days after surgery, CVD or CVD risk factors	Unable to walk without human assistance prior to hip fracture, declined blood transfusion, multiple trauma, [†] pathologic hip fracture, [†] acute MI <30 days prior to randomisation, prior enrolment, symptoms of anaemia, actively bleeding at time of potential randomisation, non-English speaking	139	81.5	26	Hip fracture & CVD/CVD risk factors 100 Comorbid: Dementia 32 ^{††} /AF 32/Chronic lung disease 21/DM 20/ Hearing problems 18/Visual problems 12/Stroke/TIA 12/ Ca 11/Alcohol abuse or withdrawal 11/Parkinson's 3 [†]	ASA 2.9	NR
Hempenius 2013 ⁵³	>65, undergoing elective surgery for solid tumour, frail according to >3 on Groningen Frailty Indicator [†] at outpatient departments at participating centres	Unable to fill in questionnaire, unable to complete the study protocol and follow-up schedule before inclusion (e.g. logistical reasons or if extra hospital visits too burdensome)	297	77.5	36	>2 comorbidities 60 (diabetes, COPD, HT, MI, other CVD, hearing and vision problems, memory problems, neurological, cerebrovascular, psychiatric or musculoskeletal disorders)/ RF 3 [†]	Surgery Minor 25.9% Intermediate 21.9% Major 52.2%	In-hospital 7.9% vs 3% (n=14 overall) [†]
Holt 2013 ³⁹	Admitted with acute medical illness from ED directly by GP to	Prevalent delirium, too unwell to be assessed (in opinion of clinical staff), [†]	362	85.4	42	CI (<24 MMSE) 58/Hearing impairment 59 Visual impairment 31	CCI ≥ 4 19.8 [†] Severe illness (>3	In-hospital 17 (11.2%) vs 23 (11.0%) [†]

Author, year	Inclusion	Exclusion	Sample size ^Δ	Age, Mean ^Δ	Male % ^{Δα}	Diagnoses % ^{Δα}	Illness severity, mean or proportion ^Δ	Mortality I vs C n (%)
	participating geriatric wards	unable to communicate verbally in English, consent not obtained <24 hours O/A					MEWS) 0.4% [†]	Six-month post-discharge 50 (33.8%) vs 64 (30.9%) [†]
Jeffs 2013 ⁴³	≥ 65, admitted to medical unit in the study area, in hospital <48 hours	Severe dysphasia making communication impossible, death expected <24 hrs, [†] infection control isolation, contraindication to mobilisation, admission to Stroke Unit or critical care, planned admission <48 hours, major psychiatric diagnosis, prior enrolment, delirium documented O/A, transfer from other hospital	647	79.4	48	Visually impaired 23 Hearing impaired 20 Premorbid CI 14	APACHE II 14.1 CCI median, IQR) 2 (1–3) [†]	NR
Kamdar 2013 ⁴⁰	≥ 70, admitted from ED to participating medical ICU	None	300	54	52	Respiratory failure 30 [†] /GI 15/ Sepsis (non-pulmonary) 12 [†] /CVD 11/Other 32	NR for whole cohort	ICU 24 (14%) vs 18 (16%) [†] In-hospital 34 (19%) vs 28 (25%) [†]
Zaubler 2013 ⁴¹	≥70, admitted to general medical floor	Not likely to benefit from the interventions i.e. non-verbal, terminal illness and receiving comfort care, [†] refusal	595	82.7	43	CI 91/ Pneumonia 8/UTI 7 Malaise/fatigue 6/ Cellulitis 5/Other 70	NR	NR

Author, year	Inclusion	Exclusion	Sample size ^Δ	Age, Mean ^Δ	Male % ^{Δα}	Diagnoses % ^{Δα}	Illness severity, mean or proportion ^Δ	Mortality I vs C n (%)
Andro 2012 ⁵⁷	≥75 years, admitted to acute geriatric unit, demented, [†] not delirious	None	255	84.8	31	Dementia 100 [†]	NR	NR
Deschodt 2012 ⁴⁶	≥65, verbally testable, admitted to ED with a traumatic hip fracture	No traumatology admission, poly-trauma, premorbid assessment missed, life expectancy < 6 months, [†] refusal, pathological fracture, no surgery, non-native speaker, no admission via ED, hard of hearing	171	80.8	27	Hip fracture 100 Dementia 21 [†]	CCI 2.3	One-year 10 (33%) vs 13 (22%) [†]
Gagnon 2012 ⁴⁹	Admitted to participating palliative care units and hospices [†]	Delirious on admission or within 48 hours of admission, hospitalised < 48 hours and > 90 days, alive at discharge	1516	68.4	46	Terminal cancer 100 [†] Comorbid: Depression 5/Anxiety 1.3/ Bipolar 0.5/Alzheimer's 0.5/ Other dementias 0.8/ Schizophrenia or psychosis 0.6/ Alcoholism 0.5/ Drug dependence 0.05/ Personality disorders 0.2/ Other psychiatric 0.7	NR	90-days 1516 (100%) (all) [†]
Martinez 2012 ⁵⁶	All internal medicine ward patients at risk of delirium i.e. presence of at least one risk	Delirium O/A, no family support, refused consent, admitted to a ward other than general internal medicine, in	287	78.2	63	HF 29/ COPD 22/ Ca 18/ CKD 14/Acute MI 9/ Mild CI 8/ DM with end-organ damage 8/ Dementia [†] 6 /PVD 6/Previous delirium	CCI median (IQR) 2 (1–4) [†]	NR

Author, year	Inclusion	Exclusion	Sample size ^Δ	Age, Mean ^Δ	Male % ^{Δα}	Diagnoses % ^{Δα}	Illness severity, mean or proportion ^Δ	Mortality I vs C n (%)
	factor: >70, previous history of CI, [†] MMSE score <24 prior to admission, alcoholism, metabolic imbalances O/A	a room with > two beds				4/Mild liver disease 4/ Mesenchymopathies 4/Peptic ulcer disease 3/Metastatic Ca 3 [†] /Severe liver disease 2 [†] /Lymphoma 0.7 /Leukaemia 0.4		
Van Rompaey 2012 ⁴⁷	≥18, expected LOS in ICU >24 hours, Dutch or English speaking, GCS score ≥10	Hearing impairment, dementia [†] , confusion or delirium O/A	136	59.5	66	≥ 1 Comorbidity 72 [†]	SOFA score first 24 hrs 7.1 SAPS 3 2.3 Maximal RIFLE: No acute kidney injury 6.3% Risk 6.2% Injury 17.9% Failure 69.6% [†]	During study period 2 (1%) (overall) [†]
Black 2011 ⁵⁸	≥18 years, admitted to medical and surgical ICU, family member willing to participate	Terminal diagnosis [†]	170	>60 73%	NR	NR	NR	12-week 32 (19%) (overall) [†]
Bo 2009 ⁵⁹	≥ 70 years, admitted from ED to participating medical and geriatric units	Delirium during ED stay or O/A, history of primary psychiatric disorder or alcohol abuse, coma, [†] aphasia, [†] intubation,	252	82.5	48	NR	APACHE II 14.3 CIRS comorbidity index 4.1 [†]	NR

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Author, year	Inclusion	Exclusion	Sample size ^Δ	Age, Mean ^Δ	Male % ^{Δα}	Diagnoses % ^{Δα}	Illness severity, mean or proportion ^Δ	Mortality I vs C n (%)
		or stroke OA, [†] language barrier, absence of a caregiver					CIRS Severity index 0.4	
Vidan 2009 ⁶³	≥70, not delirious O/A, at least one of four delirium risk factors (CI, [†] visual impairment, acute disease severity, [†] dehydration), admitted to participating geriatric and internal medicine units	Severe dementia with impaired communication, [†] aphasia, coma, agonic status, expected hospital stay ≤ 48 hours	542	84	43	Comorbid: mean 2.7 Primary O/A: Visual impairment 60/Hearing impairment 55/Infection 43/HF 21 [†]	APACHE II 11.3	In-hospital 10 (5.8%) vs 19 (5.1%) [†]
Caplan and Harper 2007 ⁴⁴	≥70, able to communicate and enrolled O/A to geriatric wards, presence of at least one risk factor for delirium (MMSE < 24, sleep deprivation, ADL, vision or hearing impairment, immobility, dehydration)	Patients who would not receive a benefit (severe dementia (MMSE < 10), psychotic disorder, unable to consent or refusal, terminal condition and receiving comfort care, [†] expected discharge <48 hours) or behavioural or medical condition that may risk volunteers' health and safety	37	84.7	22	Fracture 36/ Infection 32/ Collapse 13	NR	NR vs 1 (5%)

Author, year	Inclusion	Exclusion	Sample size ^Δ	Age, Mean ^Δ	Male % ^{Δα}	Diagnoses % ^{Δα}	Illness severity, mean or proportion ^Δ	Mortality I vs C n (%)
Lundström 2007 ⁵⁵	≥70 years, consecutively admitted to orthopaedic department with femoral neck fracture	<70, severe RA, severe hip osteoarthritis; severe RF [†] , pathological fracture, [†] bedridden before fracture [†]	199	82.2	26	Comorbid: CVD 57/ OP 52/ HT 43/ Impaired hearing 43/Depression 40/Impaired vision 39/Dementia 32 [†] /Stroke 25/DM 20/Wrist fracture 20/Lung diseases 16/ Hip fracture 15	GDS-15 4.9	In-hospital 6 (3%) vs 7 (7%) [†] Four-month 3 (3%) vs 6 (7%) [†] 12-month 7 (8%) vs 5 (7%) [†]
Taguchi 2007 ⁶⁰	Oesophageal Ca surgery, capable of communicating in Japanese	Mental or ophthalmologic disorders, reintubation, medical complications, deterioration of condition [†]	15	57.5	100	Oesophageal Ca 100/ Diabetes 20/HT 9	NR	NR
Lundström 2005 ⁵⁴	≥ 70, consecutively admitted to participating medical wards	<70, refusal to participate	400	80.1	44	Comorbid: DM 33/Stroke 25/Asthma12/ Ca 1/Dementia 4.5 [†] /Epilepsy 4.5 O/A: HF 25/Infection 18/Impaired vision 16/Stroke 11/MI 7/Epilepsy 6/Fever ≥38C 6/UTI 5/Impaired hearing 3	NR	In-hospital (delirious patients) 2 (3.2%) vs 9 (14.5%)* [†]
Wong 2005 ⁴⁵	> 50, osteoporotic hip fracture, admitted to orthopaedic ward during study period	None	99	81.8	26	Operation <24 hrs O/A 78/Vascular disease 41/Chronic lung disease 18/Diabetes 14/Renal impairment 11/Depression or anxiety 6	NR	3-month study period 3 (4.2%) vs Baseline 28-days 2 (7.1%) [†]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Author, year	Inclusion	Exclusion	Sample size ^Δ	Age, Mean ^Δ	Male % ^{Δα}	Diagnoses % ^{Δα}	Illness severity, mean or proportion ^Δ	Mortality I vs C n (%)
Cole 2002 ⁵⁰	≥65, admitted to participating medical units with delirium O/A or < 1 week of admission	Primary diagnosis of stroke, [†] ≥48 hour stay on ICU or cardiac monitoring unit, geriatric or oncology service admission, unable to speak English or French, residence outside Montreal	227	82.4	46	Dementia 58 [†]	CCI 3.3 CSI 5.8	8-week 25 22.1%) vs 22 (19.3%) [†]
Marcantonio 2001 ⁴²	≥ 65, admitted to the participating centre for primary surgical repair of hip fracture	Metastatic ca or other comorbid illnesses likely to reduce life expectancy <6 months [†] , unable to obtain informed consent <24 hours of surgery or 48 hours of admission	126	79	22	Hip fracture 100/Pre-fracture dementia 44 [†]	≥4 CCI 36% ≥4 BDRS score 44%	NR
Milisen 2001 ⁴⁸	Admitted to ED with traumatic fracture of proximal femur, hospitalized in one of the two trauma units <24 hours of surgery, Dutch-speaking, verbally testable	Multiple trauma, [†] brain concussion, pathological fractures, [†] surgery >72 hours after admission, aphasia, [†] blindness, deafness, <9 years of formal education	120	81 (median)	19	Comorbid: Previous operations 58/Cardiac 22/HT 18/ Vision or hearing problems 16/ DM 15/Dementia 15 [†] /Vascular 15/Pulmonary 13/Abdominal 13/Depression 11/Urinary 8/ Associated fracture 3/Other 24	NR	NR

Author, year	Inclusion	Exclusion	Sample size ^Δ	Age, Mean ^Δ	Male % ^{Δα}	Diagnoses % ^{Δα}	Illness severity, mean or proportion ^Δ	Mortality I vs C n (%)
Inouye 1999 ²⁰	≥ 70, admitted to one of three general-medicine units, no delirium O/A, baseline intermediate or high risk for delirium	Unable to participate in interviews (profound dementia precluding verbal communication, [†] language barrier, profound aphasia, [†] intubation or respiratory isolation), coma or terminal illness, [†] hospital stay ≤ 48 hrs, prior enrolment, other (e.g., interviewer or patient unavailable)	852	79.7	39	Primary: Pneumonia 12/Chronic lung disease 12/CHF 11 ^{††} /IHD 8/GI 13/DM or metabolic disorder 5/Ca 3/CVD 3/RF [†] 3/Anaemia 2/Other 32	APACHE II 15.6 BDRS 0.5 >2 Score 11.5%	In-hospital 6 (1.4%) vs 7 (1.6%) [†]
Cole 1994 ⁵¹	≥75, delirious first 24 hrs OA to medical department, English or French speaking, not admitted to ICU or cardiac monitoring unit or referred to oncology or geriatric services, delirious	Primary diagnosis of CVA, [†] not delirious	88	86.1	54	Dementia 56 [†] (intervention cohort)	NR	8-week 14 (33%) vs 17 (37%) [†]

* Statistically significant difference ^Δ Intervention and control participants combined ^α Rounded to nearest whole number [†] Interpreted as indicating need for palliative care

Illness severity measures: Higher scores represent higher illness severity. **AIS** Abbreviated Injury Scale, **APACHE II** Acute Physiology and Chronic Health Evaluation (scores 0-71), **ASA** American Society of Anesthesiologists physical status classification (I: normal healthy patient - VI: a declared brain-dead patient)

1
2
3 whose organs are being removed for donor purposes), **BDRS** Blessed Dementia Rating (scores 0-28, cut-off for impairment > 4), **CCI** Charlson Comorbidity Index
4 (scores 0-37), **CIRS-G** Cumulative Illness Rating Scale—Geriatrics (scores 0-56), **CSI** Clinical severity of illness (1 (mild) - 9 (moribund)), **GCS** Glasgow Coma Scale
5 (scores 3-15; scores 3-8 = coma), **GDS-15** 15-item Geriatric Depression Scale (≥ 6 = suggests depression and need for assessment, ≥ 11 = depression/severe
6 depression), **ISS** Injury Severity Score (scores 1-75), **MEWS** Modified Early Warning Score (score ≥ 5 is statistically linked to increased likelihood of death or
7 admission to an intensive care unit), **RIFLE** (Risk, Injury, Failure, Loss of kidney function, and End-stage kidney disease categories), **SAPS 3** Simplified acute
8 physiology score (scores 0-217), **SOFA** Sepsis-related Organ Failure Assessment (scores 0 to 24). **Other abbreviations:** **ADL** activities of daily living, **AF** atrial
9 fibrillation, **Ca** cancer, **CAM** Confusion Assessment Method, **CAM-ICU** Confusion Assessment Method for the Intensive Care Unit, **CI** cognitive impairment, **CKD**
10 Chronic Kidney Disease, **COPD** chronic obstructive pulmonary disease, **CVD** cardiovascular disease, **CVA** cerebrovascular accident, **DM** diabetes mellitus, **ED**
11 Emergency Department, **GI** gastrointestinal, **HF** heart failure, **hrs** hours, **HT** hypertension, **I/C** intervention/control, **ICU** intensive care unit, **IHD** ischaemic heart
12 disease, **IQR** interquartile range, **MI** Myocardial Infarction, **MMSE** Mini-Mental State Examination, **NR** not reported, **O/A** on admission, **OP** osteoporosis, **PVD**
13 peripheral vascular disease, **RASS** Richard Agitation Sedation Scale, **RF** renal failure, **SD** standard deviation, **SICU** surgical intensive care unit, **TIA** transient
14 ischemic attack, **UTI** urinary tract infection.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Table 3 Intervention characteristics

Author, year	Care components	Tailored	Who delivered
Moon 2015 ⁶¹	C, H, P, V, W, S, F, N, Pa, O, B, J, K, CA, Z	✓	Four researchers, ICU nurses
Bakker 2014 ⁵²	C, E, S, N, L, K, CA	✓	Nurses, physicians, trained volunteers, with training delivered by geriatric team
Bryczkowski 2014 ³⁸	P, S, F, D, J, K, PE	✓	Whole team, research staff, family
Chong 2014 ⁶²	C, E, H, V, P, W, G, J	NR	Trained geriatric nurses, cognitive assessment by consultant geriatrician O/A
Patel 2014 ⁶⁴	C, E, P, S, D, Pa, J, K	✓	Physicians, nurses, PT, senior staff acting as champions
Gruber-Baldini 2013 ⁶⁵	T (blood transfusion)	✓	NR
Hempenius 2013 ⁵³	C, E, H, V, N, CA	✓	Consultant geriatricians and geriatric nurses
Holt 2013 ³⁹	C, E, H, V, S, Pa, B	NR	Specialist nurse, consultant geriatrician, nurse manager, ward staff
Jeffs 2013 ⁴³	C, E	✓	Allied health assistant, PT
Kamdar 2013 ⁴⁰	P, S, J, K	✓	Bedside staff
Zaubler 2013 ⁴¹	C, H, V, P, W, S, N, K	✓	Elder Life Specialists, volunteers
Andro, 2012 ⁵⁷	C, J	NR	NR
Deschodt 2012 ⁴⁶	G, CA	✓	Consultative geriatrician, nurse, social worker, occupational therapist, physiotherapist, bedside staff
Gagnon 2012 ⁴⁹	C, F, K	✓	Bedside and research nurses, physicians
Martinez 2012 ⁵⁶	C, H, V, F	✓	Family members with education by researchers
Van Rompaey 2012 ⁴⁷	P (earplugs)	✗	Nurses
Black 2011 ⁵⁸	F	✓	Nurses, family
Bo 2009 ⁵⁹	C, E, P, W, S, F, N, Pa, K	✓	Physicians, nurses, PT
Vidan 2009 ⁶³	C, E, H, V, P, V, W, S, N	✓	Geriatricians, residents, nurses (including a full-time specialist geriatric nurse)
Caplan & Harper 2007 ⁴⁴	C, H, V, W, N	✓	Volunteers, volunteer coordinator
Lundström 2007 ⁵⁵	E, P, W, S, N, Pa, O, L, G, B, K, CA, Z, Ph	✓	Nurses, PT, OT, dietician, geriatricians; liaison with orthopaedic surgeons, geriatricians and community colleagues for post-hospital care
Taguchi 2007 ⁶⁰	J (bright light)	✓	NR
Lundström 2005 ⁵⁴	S, F, G	✓	All ward staff
Wong 2005 ⁴⁵	C, E, H, V, P, W, F, N, D, Pa, O, B, J, Z, K	✓	Geriatric registrar, geriatric team, nurses, nursing assistants, anaesthetist, pharmacist
Cole 2002 ⁵⁰	C, E, H, V, F, G, J, CA	✓	Geriatric internist, geriatric psychiatrist, study nurse, bedside nurses

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Author, year	Care components	Tailored	Who delivered
Marcantonio 2001 ⁴²	C, E, H, V, W, N, D, Pa, O, G, B, J, CA, Z, K	✓	Geriatrician consulting to orthopaedics team
Milisen 2001 ⁴⁸	S, Pa, G, K	NR	Nurses
Inouye 1999 ²⁰	C, E, H, V, P, W	✓	A geriatric nurse, therapeutic-recreation and two Elder Life Specialists, PT, geriatrician, volunteers
Cole 1994 ⁵¹	C, E, H, V, F, J, CA	✓	Geriatrician, geriatric psychiatrist, liaison nurse

Abbreviations: ICU Intensive Care Unit, NR Not reported, OA On admission, OT Occupational therapist, PT physiotherapist **Component codes:** C Cognitive activity, E Physical activity, H Hearing, V Vision, P Sleep-wake cycle preservation, W Hydration, S Staff education, F Family involvement, N Nutrition, Pa Pain, O Oxygen, L Falls prevention, G Staff changes, B Bladder/bowel, J Environment/lighting/noise, CA Comprehensive assessment, BT Blood transfusion, Z Address medical complications, PE Patient education, Ph Physiological monitoring, K Pharmacological




Figure 1 PRISMA Flow Diagram

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47




Figure 2 Overall risk of bias




Figure 3 Study approaches to people requiring palliative care

NB: Combined percentages do not add up to 100% as studies simultaneously excluded and reported people requiring palliative care.



Figure 4 Types and rates of outcomes




Figure 5 Types and rates of intervention components, including for sample of interest

Code: P Delirium prevention studies, P & T Combined delirium prevention and treatment studies, T Delirium treatment studies, S of I Sample of interest

Author, year	Adherence	Modifications
Moon 2015 ¹	NR	NR
Bakker 2014 ²	<p>[Bold: indicated and delivered by Carewell team; not bold: indicated and delivered by departments. Ranges are % of patients for whom adherence confirmed (lower value) to % of patients for whom adherence likely but not confirmed (upper value)]. Frail patients 100%, CareWell plan 95%, Medication review 94% 91%, Therapeutic activities 92%, Volunteers 84% 46% (46% represents mean % of visits performed of the total which could be performed (data from 49 patients), Information in discharge letter 74% 55%, Orientation points 71% 83–92%, Orientation 60%, Consult physiotherapist 59% 56–68%, Nutrition 53%, Day program activities 50% 52–57%, Day program delirium 48% 46–59%, CGA by geriatrician 42% NA, Registration food intake 40% 74–83%, Medical history by proxy 32%, Consult dietitian 33% 37–40%, Falls sensor 22% 46–48%, Discharge planning 16% 100%, NA, Mobilizing 15%, Multidisciplinary meeting 11% NA.</p>	NR
Bryczkowski 2014 ³	NR	NR
Chong 2014 ⁴	100% compliance to semi-structured protocol by trained geriatric nurses	NR
Patel 2014 ⁵	<p>Overall compliance with interventions > 90%. Noise: Offer earplugs to all patients with RASS > 4 100%, Staff and visitors to speak quietly 96%, Close all doors 96%, Turn monitoring equipment to night mode 23:00-07:00 96%, Reduce volumes on all telephones 23:00-07:00 96%, No non-clinical discussions around patients' bed spaces 92%; Light: Dim main ICU lights 23:00-07:00 100%, Use bedside lighting for patient care 26 100%, Offer eye-masks to all patients with RASS > 4 25 96%; Patient care: Orientate patients regarding time, place and date every eight hours 100%, All patients requiring mechanical ventilation of the lungs to be assessed daily for suitability for sedation hold or trials of spontaneous breathing 100%, Hourly pain scores and prompt action to optimize analgesia 100%, Set appropriate sedation targets once per day (based on RASS) 100%, If patients sleep poorly or have a positive result on the CAM-ICU, perform a medication review within 24 h 96%, Complete care procedures before 23:00 or delay completion until after 08:00 where possible 92%, Group care/procedures where possible 88%, Ensure early mobilization when possible and appropriate 88%.</p>	NR
Gruber-Baldini 2013 ⁶	Number of units of blood transfused: None 4.5%, One unit 40.9%, Two units 36.4%, Three units 12.1%, ≥ four units 6.1%. Total units of blood transfused post-randomization = 115 in 66 participants.	NR
Hempenius 2013 ⁷	NR	NR
Holt 2013 ⁸	Staff attendance at delirium education sessions 70%, Adherence to delirium risk factor modification protocols 27–57%. Protocol adherence highest for reorientation and hydration, lowest for mobility and constipation	During pilot phase, materials for education and practice change (30-min interactive lecture with a handout, a delirium quiz, a poster, reference material and case vignettes) were modified following consultation with local opinion leaders.
Jeffs 2013 ⁹	Therapeutic encounters per day, median 1.4 (0.9–1.8), minutes of therapy per day median 38 (25– 52)	NR

Kamdar 2013 ¹⁰	Checklist item completion rates 86-94%. Patient daytime interventions: Blinds raised 79%, Caffeine avoided after 3pm 54%, Less than 50% of day shift spent napping 45%. Patient nighttime interventions: Room lights dimmed before 10pm 78%, Unnecessary alarms prevented 77%, Room temperature optimized 77%, Pain appropriately controlled 68%, Room curtain closed before 10pm 64%, Warm bath before 10pm 49%, Television off 59%, Medication given per sleep guideline 13%, Soft music offered and accepted 11%, Eye mask offered and accepted 2%, Earplugs offered and accepted 1%. ICU-wide nighttime interventions: Hallway lights dimmed by 10pm 89%, Overhead pages after 10pm: none 15%, 1-3 36%, >3 8%, unknown 41%. Estimated number of nurse interruptions between 10pm-7am: 0-5 interruptions 28%, 6-10 interruptions 21%, >10 interruptions 13%, NR 37%.	NR
Zaubler 2013 ¹¹	NR	NR
Andro, 2012 ¹²	NR	NR
Deschodt 2012 ¹³	No significant differences between groups in care given except intervention participants received more occupational therapy (intervention 69.1%, control 41.6%, $p < 0.001$) and opioid pain medication (intervention 91.5%, control 75.3%, $p=0.004$) than controls. 2011 primary report: Recommendations made for 79 of 94 participants in intervention group (84.0%). No recommendations given to 15 participants because no need for additional advice on top of usual care could be identified. Of 338 recommendations for 79 participants, adherence could not be determined for nine recommendations, leaving 329 recommendations for study (97.3%). The occupational therapist made nine recommendations with unknown adherence, suggesting walking aids or adapted footwear. Mean number of recommendations per participant was 4.3+/- 2.1 (range 1-10). Complete adherence: 56.8%, partial adherence 10.6%. Trauma ward team did not comply with 32.5% of recommendations.	NR
Gagnon 2012 ¹⁴	CRS score assigned: 91.2%. Adherence to CAM: 39%. Overall adherence to study protocols: 89.7%. Pharmacological risk alert: 91.2%, Orientation protocol: 84.5%, Family intervention: 84.1%. Most common reasons for noncompliance with study protocols were pharmacological sedation and coma. Main reason for missed family intervention was absence of a relative.	CAM was not used as an outcome measure as per study protocol as it was obtained in only 39% of patients (due to challenges conducting the baseline CAM structured interview in the last days or hours of life). Delays in completion also rendered results invalid.
Martinez 2012 ¹⁵	NR	NR
Van Rompaey 2012 ¹⁶	"No accidental or intentional removal of the earplugs was reported."	NR
Black 2011 ¹⁷	NR	During pilot phase, there was a change to the study protocol allowing the study information and booklet to be given to families on admission rather than wait until the researcher was present, following feedback from two families.
Bo 2009 ¹⁸	NR	NR
Vidan 2009 ¹⁹	Overall rate of adherence (percentage of actions per days performed in each of the seven targeted intervention domains): 75.7% of patient-days per intervention actions. Highest was mobilization: 91%, lowest was sleep preservation: 50%.	NR

1 2 3 4 5	Caplan & Harper 2007 ²⁰	NR	NR
6 7 8 9 10 11 12	Lundström 2007 ²¹	27 eligible patients excluded due to failed inclusion routines. More documented assessment of underlying causes of delirium in the intervention ward compared with the control ward (2.28±1.25 vs 0.90±0.90, p <0.001), also more documented treatments for underlying causes of delirium (1.69±1.56 vs 0.56±0.98, p <0.001). Delirious control patients more often given sedatives (41.7% vs 15.4%, p =0.008) and opioid drugs on demand (61.7% vs 30.8%, p =0.004) than intervention participants.	NR
13 14	Taguchi 2007 ²²	Bright light therapy started at 3.1+/-1.4 days after surgery and performed over a mean of 2.8+/-0.9 days for 110+/-14 min each day	NR
15 16	Lundström 2005 ²³	NR	NR
17 18 19 20 21 22 23 24	Wong 2005 ²⁴	Daily CAM ratings 97.2%. Geriatric registrar recommendation (average six/patient) 89.9%. Recommendations: Regulation of bladder and bowel function 24%, Early detection/treatment of major complications 22%, Correction of fluid and electrolyte imbalance 14.4%, Discontinuation of unnecessary medications 13.8%, Provision of adequate oxygen delivery 8.5%, Treatment of severe pain 5.4%, Treatment of agitated delirium 4.3%, Use of appropriate environmental stimuli 3.4%, Adequate nutritional intake 2.4%, Early mobilization and rehabilitation 1.8%	NR
25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43	Cole 2002 ²⁵	97% patients received the intervention (mean of 1.4 days after enrolment) as planned. Consultants had a mean of 1.96 contacts (median 1.0, range 1–6) with each patient in the intervention group and made a mean of 6.02 recommendations (median 1.0, range 1–17) per patient, most frequently for medication changes or investigations. Recommendations: Medication changes 73.2%, investigations 69%, other recommendations (e.g., patient supports, mobilization) 20.6%. Study nurse contact with each patient: mean of 11.7 (SD 9.8) (median 8.0, range 1–39), lasting a mean of 35.7 (SD 28.8) minutes (median 30, range 5–240). Mean total time spent with each patient was 418 (SD 282) minutes (median 318, range 90–1315). Four most frequent study nurse activities were assessment and support of patients, and education and support of nursing staff and families. I vs C comparisons: Geriatric or geriatric psychiatry consultation 100% v.18%, Study nurse visit 100% v. 0%, Documentation of delirium by attending physicians 41% v. 27%, p = 0.03, Decreases in medication 66.4% v. 57.9%, p = 0.19, Occupational therapy, recreational therapy or social work consultation 64.6% v. 54.4%, p = 0.12, Emotional support by ward nurses 14.3% v. 9.4%, p = 0.70, Orienting cues by ward nurses 23.2% v. 16.7%, p = 0.22, Personal possessions at the bedside 35.4% v. 22.8%, p = 0.04.	NR
44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60	Marcantonio 2001 ²⁶	Initial geriatrics consultation preoperatively 61%, remainder had initial consultation within 24 hours of surgery. 591 recommendations/mean of 9.5 recommendations per patient (range 3–21). Overall adherence rate by orthopedics team: 77%. Data did not describe overall management; therefore, comparable data were not available for the usual-care group. Consultants did not recommend things that the orthopedists or nurses were already doing; only when something was not being done that they felt should be. Recommended/adhered: Adequate CNS oxygen delivery: Supplemental oxygen to keep saturation >90%, preferably >95% 29%/ 89%, Treatment to raise systolic blood pressure >2/3 baseline or >90 mmHg 4 6%/100%, Transfusion to keep hematocrit >30% 92%/79%, Fluid/electrolyte balance: Treatment to restore serum sodium, potassium, glucose to normal limits (glucose <300 mg/dl, <16.5 mmol/L for diabetics) 37%/96%	NR

	<p>Treat fluid overload or dehydration detected by examination or blood tests 48%/90% Treatment of severe pain: Around-the-clock acetaminophen (1 gram four times daily) 40%/32% Early-stage break-through pain: low-dose subcutaneous morphine, avoid meperidine 21%/62%, Late-stage break-through pain: oxycodone as needed 5%/67%, Elimination of unnecessary medications: Discontinue/minimize benzodiazepines, anticholinergics, antihistamines 68%/83%, Eliminate drug interactions, adverse effects, modify drugs accordingly 21%/54%, Eliminate medication redundancies 13%/63% Regulation of bowel/bladder function: Bowel movement by postoperative day 2 and every 48 hours 68%/57%, D/c urinary catheter by postoperative day 2, screen for retention or incontinence 71%/89%, Skin care program for patients with established incontinence 3%/100% Adequate nutritional intake: Dentures used properly, proper positioning for meals, assist as needed 56%/66%, Supplements: 1 can Ensure,* 3 cans Ensure* for poor oral intake 35%/45%, If unable to take food orally, feed via temporary nasogastric tube 2%/100% Early mobilization and rehabilitation: Out of bed on postoperative day 1 and several hours daily 58%/81% Mobilize/ambulate by nursing staff as tolerated, such as to bathroom 29%/72%, Daily physical therapy; occupational therapy if needed 2%/100% Prevention, early detection, and treatment of major postoperative complications: Myocardial infarction/ischemia—electrocardiogram, cardiac enzymes if needed 34%/81% Supraventricular arrhythmias/atrial fibrillation—appropriate rate control, electrolyte adjustments, anticoagulation 5%/3 100%, Pneumonia/chronic obstructive pulmonary disease - screening, treatment, including chest therapy 44%/67%, Pulmonary embolus -appropriate anticoagulation 50%/100%, Screening for and treatment of urinary tract infection 52%/63% Appropriate environmental stimuli: Appropriate use of glasses and hearing aids 5%/67%, Provision of clock and calendar 0%, If available, use of radio, tape recorder, and soft lighting 0% Treatment of agitated delirium: Appropriate diagnostic workup/management 2%/100%, For agitation, calm reassurance, family presence, and/or sitter 3%/100%, For agitation, if absolutely necessary, low-dose haloperidol 0.25–0.5 mg every 4 hours as needed; if contraindicated, use lorazepam at same dose 19%/83%.</p>	
Milisen 2001 ²⁷	NR	NR
Inouye 1999 ²⁸	<p>Overall rate of adherence (complete and partial adherence, based on patient days) to all intervention protocols: 87%. Orientation protocol 96%, vision protocol 92%, hearing protocol 92%, therapeutic activities 86%, early mobilization 84%, volume repletion 81%, non-pharmacological sleep protocol 71%. Most common reasons for non-adherence included refusal by the patient, lack of availability of patient because of procedures elsewhere in the hospital, medical contraindications, and lack of availability of intervention staff members.</p>	NR
Cole 1994 ²⁹	<p>Initial recommendations made for all 39 patients in the intervention group who were assessed on admission, with 25 follow-up recommendations. Initial recommendations: Investigations (n=4), drug prescriptions (n=3), other (n=7) or a combination (n=25). Follow-up recommendations: investigations (n=1), drug prescriptions (n=1), other (n=3) or a combination (n=20). Number of nurse follow-up notes 0-8 (mean 3); 97% of eligible notes completed. Rates of full compliance with initial recommendations ranged from: Other 96% - investigations 77%. Rates for follow-up recommendations ranged from: Other 91% - investigations 50%.</p>	NR

References

1. Moon K-J and Lee S-M. The effects of a tailored intensive care unit delirium prevention protocol: A randomized controlled trial. *International Journal of Nursing Studies* 2015; 52: 1423-1432. DOI: 10.1016/j.ijnurstu.2015.04.021.
2. Bakker FC, Persoon A, Bredie SJ, et al. The CareWell in Hospital program to improve the quality of care for frail elderly inpatients: results of a before-after study with focus on surgical patients. *American Journal of Surgery* 2014; 208: 735-746. DOI: <http://dx.doi.org/10.1016/j.amjsurg.2014.04.009>.
3. Bryczkowski SB, Lopreiato MC, Yonclas PP, et al. Delirium prevention program in the surgical intensive care unit improved the outcomes of older adults. *The Journal of surgical research* 2014; 190: 280-288. 2014/03/29. DOI: 10.1016/j.jss.2014.02.044.
4. Chong MS, Chan M, Tay L, et al. Outcomes of an innovative model of acute delirium care: the Geriatric Monitoring Unit (GMU). *Clinical Interventions in Aging* 9(2014).
5. Patel J, Baldwin J, Bunting P, et al. The effect of a multicomponent multidisciplinary bundle of interventions on sleep and delirium in medical and surgical intensive care patients. *Anaesthesia* 2014; 69: 540-549. DOI: 10.1111/anae.12638.
6. Gruber-Baldini AL, Marcantonio E, Orwig D, et al. Delirium outcomes in a randomized trial of blood transfusion thresholds in hospitalized older adults with hip fracture. *Journal of the American Geriatrics Society* 61(8)(2013).
7. Hempenius L, Slaets JPJ, van Asselt D, et al. Outcomes of a geriatric liaison intervention to prevent the development of postoperative delirium in frail elderly cancer patients: Report on a multicentre, randomized, controlled trial. *PloS One* 2013; 8: e64834.
8. Holt R, Young J and Heseltine D. Effectiveness of a multi-component intervention to reduce delirium incidence in elderly care wards. *Age and Ageing* 2013; 42: 721-727.
9. Jeffs KJ, Berlowitz DJ, Grant S, et al. An enhanced exercise and cognitive programme does not appear to reduce incident delirium in hospitalised patients: a randomised controlled trial. *BMJ Open* 2013; 3: e002569.
10. Kamdar BB, King LM, Collop NA, et al. The effect of a quality improvement intervention on perceived sleep quality and cognition in a medical ICU. *Critical Care Medicine* 2013; 41: 800.
11. Zaubler TS, Murphy K, Rizzuto L, et al. Quality improvement and cost savings with multicomponent delirium interventions: replication of the hospital elder life program in a community hospital. *Psychosomatics* 2013; 54: 219-226.
12. Andro M, Comps E, Estivin S, et al. Prevention of delirium in demented hospitalized patients. *European journal of internal medicine* 2012; 23: 124-125. 2012/01/31. DOI: 10.1016/j.ejim.2011.05.011.
13. Deschodt M, Braes T, Flamaing J, et al. Preventing delirium in older adults with recent hip fracture through multidisciplinary geriatric consultation. *Journal of the American Geriatrics Society* 60(4)(2012).
14. Gagnon P, Allard P, Gagnon B, et al. Delirium prevention in terminal cancer: assessment of a multicomponent intervention. *Psycho-Oncology* 2012; 21: 187-194. DOI: <http://dx.doi.org/10.1002/pon.1881>.
15. Martinez FT, Tobar C, Beddings CI, et al. Preventing delirium in an acute hospital using a non-pharmacological intervention. *Age and Ageing* 41(5)(2012).
16. Van Rompaey B, Elseviers MM, Drom W, et al. The effect of earplugs during the night on the onset of delirium and sleep perception: a randomized controlled trial in intensive care patients. *Critical Care* 2012; 16: R73. DOI: 10.1186/cc11330.
17. Black P, Boore JRP and Parahoo K. The effect of nurse-facilitated family participation in the psychological care of the critically ill patient. *Journal of Advanced Nursing* 2011; 67: 1091-1101. DOI: 10.1111/j.1365-2648.2010.05558.x.
18. Bo M, Martini B, Ruatta C, et al. Geriatric ward hospitalization reduced incidence delirium among older medical inpatients. *American Journal of Geriatric Psychiatry* 2009; 17: 760-768. Comparative Study.
19. Vidan MT, Sanchez E, Alonso M, et al. An intervention integrated into daily clinical practice reduces the incidence of delirium during hospitalization in elderly patients. *Journal of the American Geriatrics Society* 2009; 57: 2029-2036. DOI: <http://dx.doi.org/10.1111/j.1532-5415.2009.02485.x>.
20. Caplan GA and Harper EL. Recruitment of volunteers to improve vitality in the elderly: the REVIVE study. *Internal Medicine Journal* 2007; 37: 95-100. DOI: 10.1111/j.1445-5994.2007.01265.x.
21. Lundström M, Olofsson B, Stenvall M, et al. Postoperative delirium in old patients with femoral neck fracture: a randomized intervention study. *Aging Clinical and Experimental Research* 2007; 19: 178-186.
22. Taguchi T, Yano M and Kido Y. Influence of bright light therapy on postoperative patients: a pilot study. *Intensive & Critical Care Nursing* 23(5)(2007).
23. Lundström M, Edlund A, Karlsson S, et al. A multifactorial intervention program reduces the duration of delirium, length of hospitalization, and mortality in delirious patients. *Journal of the American Geriatrics Society* 2005; 53: 622-628. DOI: 10.1111/j.1532-5415.2005.53210.x.
24. Wong Tin Niam DM, Bruce JJ and Bruce DG. Quality project to prevent delirium after hip fracture. *Australasian Journal on Ageing* 2005; 24: 174-177. DOI: 10.1111/j.1741-6612.2005.00108.x.
25. Cole MG, McCusker J, Bellavance F, et al. Systematic detection and multidisciplinary care of delirium in older medical inpatients: a randomized trial. *Canadian Medical Association Journal* 167(7)(2002).
26. Marcantonio ER, Flacker JM, Wright RJ, et al. Reducing delirium after hip fracture: a randomized trial. *Journal of the American Geriatrics Society* 2001; 49: 516-522.
27. Milisen K, Foreman MD, Abraham IL, et al. A Nurse-Led Interdisciplinary Intervention Program for Delirium in Elderly Hip-Fracture Patients. *Journal of the American Geriatrics Society* 2001; 49: 523-532.
28. Inouye SK, Bogardus STJ, Charpentier PA, et al. A Multicomponent Intervention to Prevent Delirium in Hospitalized Older Patients. *New England Journal of Medicine* 1999; 340: 669-676. DOI: 10.1056/nejm199903043400901.
29. Cole MG, Primeau FJ, Bailey RF, et al. Systematic intervention for elderly inpatients with delirium: a randomized trial. *Canadian Medical Association Journal* 151(7)(1994).

Author, year	Adverse events
Moon 2015 ¹	Mortality – see Table 3
Bakker 2014 ²	≥1 Complication 45% v. 40%
Bryczkowski 2014 ³	Restraint-free days/30 mean 27 (95% CI 26, 29) v. mean 28 (95% CI 27, 29)
Chong 2014 ⁴	[Before v. Intervention v. Control] Physical restraint use: 44.7% v. 0 v. 23.1*, Chemical restraint use: 72.3% v. 40.3% v. 33.3%, Falls: 2.1% v. 1.3% v. 2.6%, Urinary catheter use: 31.9% v. 29.1% v. 25.6, Pressure ulcer rate: 9.1% v. 4.1% v. 1.3%*, Nosocomial infection: 23.4% v. 10.7% v. 7.7%*
Patel 2014 ⁵	NR
Gruber-Baldini 2013 ⁶	Infections 4.6% vs 4.2%, PE 3.0% vs 0, CHF 1.5% vs 2.8%, Hemorrhaging (>100cc) 9.1% vs 5.6%
Hempenius 2013 ⁷	Postoperative complications: Pulmonary 24.4% vs 20.3%, Neurological 6.3% vs 6.0%, Cardiovascular 31.5% vs 27.8%, Thromboembolic 0.8% vs 0, Bleeding 8.7% vs 4.5%, Wound infection 10.2% vs 9.0%, Wound dehiscence 3.1% vs 3.0%, UTI 6.3% vs 5.3%, Anastomotic leakage 3.9% vs 1.5%, Pressure ulcer 3.9% vs 5.3%, RF 3.9% vs 1.5%, Electrolyte disturbance 11.8% vs 9.0%, Fall 3.1% vs 1.5%, Urinary retention 11.8% vs 9.0%, Ileus/gastroparesis 7.1% vs 10.5%
Holt 2013 ⁸	Mortality – see Table 3
Jeffs 2013 ⁹	"No adverse events were reported."
Kamdar 2013 ¹⁰	Mortality – see Table 3
Zaubler 2013 ¹¹	NR
Andro, 2012 ¹²	NR
Deschodt 2012 ¹³	Mortality – see Table 3
Gagnon 2012 ¹⁴	NR
Martinez 2012 ¹⁵	Falls 0 v. 2.8%
Van Rompaey 2012 ¹⁶	NR
Black 2011 ¹⁷	NR
Bo 2009 ¹⁸	NR
Vidan 2009 ¹⁹	Results reported graphically, no exact figures. Physical restraints approximately 2% v. 10%*, Falls approximately 2% v. 1%
Caplan & Harper 2007 ²⁰	Falls 6.3% vs 19%, Increased unplanned readmissions at 1 month 31.3% vs 19%
Lundström 2007 ²¹	Anemia 86.3% vs 82.3%, Constipation 37.3% vs 48.5%, Decubitus ulcers 8.8% vs 22.1%*, Depression 49.5% vs 54.6%, Diarrhea 21.6% vs 27.1%, HF 5.9% vs 11.6%, Pneumonia 4.9% vs 3.1%, UTI 31.4% vs 51.0%* Other infections, 17.8% vs 17.7%, Sleeping problems 27.5% vs 45.4%*, MI 2% vs 4.1%, Nutritional complications 24.5% v. 38.1%, PE 2% vs 0, Stroke 0 vs 1%, Stomach ulcers 3% vs 4.1%, Urinary retention 15.7% vs 18.6%, Falls 11.8% vs 26.8%*
Taguchi 2007 ²²	NR
Lundström 2005 ²³	Mortality – see Table 3
Wong 2005 ²⁴	NR
Cole 2002 ²⁵	Mortality – see Table 3
Marcantonio 2001 ²⁶	NR
Milisen 2001 ²⁷	Mortality – see Table 3
Inouye 1999 ²⁸	"No adverse effects were associated with the intervention protocols."
Cole 1994 ²⁹	Restraint use 37% v. 29%

References

1. Moon K-J and Lee S-M. The effects of a tailored intensive care unit delirium prevention protocol: A randomized controlled trial. *International Journal of Nursing Studies* 2015; 52: 1423-1432. DOI: 10.1016/j.ijnurstu.2015.04.021.
2. Bakker FC, Persoon A, Bredie SJ, et al. The CareWell in Hospital program to improve the quality of care for frail elderly inpatients: results of a before-after study with focus on surgical patients. *American Journal of Surgery* 2014; 208: 735-746. DOI: <http://dx.doi.org/10.1016/j.amjsurg.2014.04.009>.
3. Bryczkowski SB, Lopreiato MC, Yonclas PP, et al. Delirium prevention program in the surgical intensive care unit improved the outcomes of older adults. *The Journal of surgical research* 2014; 190: 280-288. 2014/03/29. DOI: 10.1016/j.jss.2014.02.044.
4. Chong MS, Chan M, Tay L, et al. Outcomes of an innovative model of acute delirium care: the Geriatric Monitoring Unit (GMU). *Clinical Interventions in Aging* 9(2014).
5. Patel J, Baldwin J, Bunting P, et al. The effect of a multicomponent multidisciplinary bundle of interventions on sleep and delirium in medical and surgical intensive care patients. *Anaesthesia* 2014; 69: 540-549. DOI: 10.1111/anae.12638.
6. Gruber-Baldini AL, Marcantonio E, Orwig D, et al. Delirium outcomes in a randomized trial of blood transfusion thresholds in hospitalized older adults with hip fracture. *Journal of the American Geriatrics Society* 61(8)(2013).
7. Hempenius L, Slaets JPJ, van Asselt D, et al. Outcomes of a geriatric liaison intervention to prevent the development of postoperative delirium in frail elderly cancer patients: Report on a multicentre, randomized, controlled trial. *PloS One* 2013; 8: e64834.
8. Holt R, Young J and Heseltine D. Effectiveness of a multi-component intervention to reduce delirium incidence in elderly care wards. *Age and Ageing* 2013; 42: 721-727.
9. Jeffs KJ, Berlowitz DJ, Grant S, et al. An enhanced exercise and cognitive programme does not appear to reduce incident delirium in hospitalised patients: a randomised controlled trial. *BMJ Open* 2013; 3: e002569.
10. Kamdar BB, King LM, Collop NA, et al. The effect of a quality improvement intervention on perceived sleep quality and cognition in a medical ICU. *Critical Care Medicine* 2013; 41: 800.
11. Zaubler TS, Murphy K, Rizzuto L, et al. Quality improvement and cost savings with multicomponent delirium interventions: replication of the hospital elder life program in a community hospital. *Psychosomatics* 2013; 54: 219-226.
12. Andro M, Comps E, Estivin S, et al. Prevention of delirium in demented hospitalized patients. *European journal of internal medicine* 2012; 23: 124-125. 2012/01/31. DOI: 10.1016/j.ejim.2011.05.011.
13. Deschodt M, Braes T, Flamaing J, et al. Preventing delirium in older adults with recent hip fracture through multidisciplinary geriatric consultation. *Journal of the American Geriatrics Society* 60(4)(2012).
14. Gagnon P, Allard P, Gagnon B, et al. Delirium prevention in terminal cancer: assessment of a multicomponent intervention. *Psycho-Oncology* 2012; 21: 187-194. DOI: <http://dx.doi.org/10.1002/pon.1881>.
15. Martinez FT, Tobar C, Beddings CI, et al. Preventing delirium in an acute hospital using a non-pharmacological intervention. *Age and Ageing* 41(5)(2012).
16. Van Rompaey B, Elseviers MM, Drom W, et al. The effect of earplugs during the night on the onset of delirium and sleep perception: a randomized controlled trial in intensive care patients. *Critical Care* 2012; 16: R73. DOI: 10.1186/cc11330.
17. Black P, Boore JRP and Parahoo K. The effect of nurse-facilitated family participation in the psychological care of the critically ill patient. *Journal of Advanced Nursing* 2011; 67: 1091-1101. DOI: 10.1111/j.1365-2648.2010.05558.x.
18. Bo M, Martini B, Ruatta C, et al. Geriatric ward hospitalization reduced incidence delirium among older medical inpatients. *American Journal of Geriatric Psychiatry* 2009; 17: 760-768. Comparative Study.
19. Vidan MT, Sanchez E, Alonso M, et al. An intervention integrated into daily clinical practice reduces the incidence of delirium during hospitalization in elderly patients. *Journal of the American Geriatrics Society* 2009; 57: 2029-2036. DOI: <http://dx.doi.org/10.1111/j.1532-5415.2009.02485.x>.
20. Caplan GA and Harper EL. Recruitment of volunteers to improve vitality in the elderly: the REVIVE study. *Internal Medicine Journal* 2007; 37: 95-100. DOI: 10.1111/j.1445-5994.2007.01265.x.
21. Lundström M, Olofsson B, Stenvall M, et al. Postoperative delirium in old patients with femoral neck fracture: a randomized intervention study. *Aging Clinical and Experimental Research* 2007; 19: 178-186.
22. Taguchi T, Yano M and Kido Y. Influence of bright light therapy on postoperative patients: a pilot study. *Intensive & Critical Care Nursing* 23(5)(2007).
23. Lundström M, Edlund A, Karlsson S, et al. A multifactorial intervention program reduces the duration of delirium, length of hospitalization, and mortality in delirious patients. *Journal of the American Geriatrics Society* 2005; 53: 622-628. DOI: 10.1111/j.1532-5415.2005.53210.x.
24. Wong Tin Niam DM, Bruce JJ and Bruce DG. Quality project to prevent delirium after hip fracture. *Australasian Journal on Ageing* 2005; 24: 174-177. DOI: 10.1111/j.1741-6612.2005.00108.x.
25. Cole MG, McCusker J, Bellavance F, et al. Systematic detection and multidisciplinary care of delirium in older medical inpatients: a randomized trial. *Canadian Medical Association Journal* 167(7)(2002).
26. Marcantonio ER, Flacker JM, Wright RJ, et al. Reducing delirium after hip fracture: a randomized trial. *Journal of the American Geriatrics Society* 2001; 49: 516-522.
27. Milisen K, Foreman MD, Abraham IL, et al. A Nurse-Led Interdisciplinary Intervention Program for Delirium in Elderly Hip-Fracture Patients. *Journal of the American Geriatrics Society* 2001; 49: 523-532.
28. Inouye SK, Bogardus STJ, Charpentier PA, et al. A Multicomponent Intervention to Prevent Delirium in Hospitalized Older Patients. *New England Journal of Medicine* 1999; 340: 669-676. DOI: 10.1056/nejm199903043400901.
29. Cole MG, Primeau FJ, Bailey RF, et al. Systematic intervention for elderly inpatients with delirium: a randomized trial. *Canadian Medical Association Journal* 151(7)(1994).



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	Title page
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	1-2
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	4-5
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	5
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	1 (registration)
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	6
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	6
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	6
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	6
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	6
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	6-7
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	7
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	NA
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I ² for each meta-analysis).	7



PRISMA 2009 Checklist

Page 1 of 2

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	7
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	7
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	8, Figure 1
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	8, Table 1
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	9
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	10-18
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	NA
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	9, Figure 2
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	NA
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	18-21
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	21
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	22
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	22

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit <http://mc.manuscriptcentral.com/palliative-medicine> or www.prisma-statement.org



PRISMA 2009 Checklist

For Peer Review

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47

PALLIATIVE MEDICINE AUTHOR SUBMISSION CHECKLIST

Please complete this checklist for all papers submitted. Please indicate, very briefly, how this has been addressed. This checklist is a mandatory upload on submission.

Item	Explanation	How this has been addressed (briefly, a sentence will suffice)
Article title	<p>WHY: Because we want readers to find your work.</p> <p>Have you followed our guidelines on writing a good title that will be found by search engines? (E.g. with methods in the title, use of common words for the issue addressed, no country names, and possibly indicating findings). If your study has an acronym is it included in the title?</p>	Yes
Abstract	<p>WHY: Because structured abstracts have more detail for readers and search engines.</p> <p>Have you followed our guidelines on writing your structured abstract? Please remember we have separate abstract structures for original research, reviews and case reports. There should be no abbreviations in the abstract, EXCEPT a study acronym which should be included if you have one. If a trial (or other design formally registered with a database) have you included your registration details?</p>	Yes
Key statements	<p>WHY: Because readers want to understand your paper quickly.</p> <p>Have you included our key statements within the body of your paper (after abstract and before the main text is a good place!) and followed our guidelines for how these are to be written? There are three main headings required, and each may have 1-3 separate bullet points. Please use clear, succinct, single sentence separate bullet points rather than complex or multiple sentences.</p>	Yes
Keywords	<p>WHY: Because MeSH headings mean it is properly indexed.</p> <p>Have you given keywords for your study? We ask that these are current MeSH headings unless there is no suitable heading for use (please give explanation in cover letter). https://meshb.nlm.nih.gov/search</p>	Yes
International relevance	<p>WHY: We have readers from around the world who are interested in your work.</p> <p>Have you contextualised your work for an international audience and explained how your work contributes to an international knowledge base? Avoid drawing from policy from one context only, think</p>	Yes

	how your work could be relevant more widely. Do define terms clearly e.g. hospice has a different meaning in many countries.	
Publishing guidelines	WHY: Because clear and robust reporting helps people interpret your work accurately Have you submitted a completed checklist for a relevant publishing guideline as a supplementary file? http://www.equator-network.org/ These include CONSORT, PRISMA, COREQ checklists, but others may be more relevant for your type of manuscript. If no published checklist exists please create one as a table from the list of requirements in your chosen guideline. If your study design does not have a relevant publishing guideline please review closest matches and use the most appropriate with an explanation.	Yes - PRISMA
Word count	WHY: Because readers want to find the core information quickly. Does your paper adhere to our word count for your article type? Please insert number of words in the box to the right. Remember that tables, figures, qualitative data extracts and references are not included in the word count.	5550
Figures and tables and/or quotations	WHY: Because readers want to find the core information quickly. Have you adhered to our guidelines on the number of tables and figures for your article type? Data (e.g. quotations) for qualitative studies are not included in the word count, and we prefer that they are integrated into the text (e.g. not in a separate table).	Yes
Study registration	WHY: Because this means readers understand how you planned your study Where appropriate have you included details (including reference number, date of registration and URL) of study registration on a database e.g. trials or review database. If your study has a published protocol, is this referenced within the paper?	Yes
Other study publications?	WHY: So readers can understand the full context of your study If there are other publications from this study are these referenced within the body of the paper? Please do not reference papers in preparation or submitted, but in-press publications are acceptable.	NA
Scales, measures or questionnaires	WHY: So readers can understand your paper in the context of this information If your study primarily reports the development or testing of scales/measures or questionnaires have you included a copy of the instrument as a supplementary file?	NA

1		
2		
3		
4	Abbreviations	WHY: Because abbreviations make a paper hard to read, and are easily misunderstood Have you removed all abbreviations from the text except for extremely well known, standard abbreviations (e.g. SI units), which should be spelt out in full first? We do not allow abbreviations for core concepts such as palliative or end of life care.
5		Yes
6		
7		
8		
9		
10	Research ethics and governance approvals for research involving human subjects	WHY: We will only publish ethically conducted research, approved by relevant bodies Have you given full details of ethics/governance/data protection approvals with reference numbers, full name of the committee(s) giving approval and the date of approval? If such approvals are not required have you made it explicit within the paper why they were not required. Are details of consent procedures clear in the paper?
11		NA
12		
13		
14		
15		
16		
17	Date(s) of data collection	WHY: So readers understand the context within which data were collected Have you given the dates of data collection for your study within the body of your text? If your data are over 5 years old you will need to articulate clearly why they are still relevant and important to current practice.
18		Yes
19		
20		
21		
22		
23	Structured discussion	WHY: So readers can find key information quickly Papers should have a structured discussion, with sub headings, summarising the main findings, addressing strengths and limitations, articulating what this study adds with reference to existing international literature, and presenting the implications for practice.
24		Yes
25		
26		
27		
28	Case reports	WHY: So that participants are protected, and its importance made clear If your study is a case report have you followed our clear structure for a case report, including highlighting what research is needed to address the issue raised? Have you made clear what consent was required or given for the publication of the case report? Have you provided evidence of such consent as a supplementary file to the editor?
29		NA
30		
31		
32		
33		
34		
35	Acknowledgements and declarations	WHY: So readers understand the context of the research Have you included a funding declaration according to the SAGE format? Are there acknowledgements to be made? Have you stated where data from the study are deposited and how they may be available to others? Have you conflicts of interest to declare?
36		Yes
37		
38		
39		
40		
41		
42		
43		
44		
45		
46		

Supplementary data and materials	WHY: So the context is clear, but the main paper succinct for the reader Is there any content which could be provided as supplementary data which would appear only in the online version of accepted papers? This could include large tables, full search strategies for reviews, additional data etc.	Yes
References	WHY: So people can easily find work you have referenced Are your references provided in SAGE Vancouver style? You can download this style within Endnote and other referencing software.	Yes
Ownership of work.	Can you assert that you are submitting your original work, that you have the rights in the work, that you are submitting the work for first publication in the Journal and that it is not being considered for publication elsewhere and has not already been published elsewhere, and that you have obtained and can supply all necessary permissions for the reproduction of any copyright works not owned by you.	Yes