

This is a repository copy of *Blood pressure change across pregnancy in white British and Pakistani women: analysis of 1 data from the Born in Bradford cohort*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/149582/>

Version: Accepted Version

Article:

Farrar, Diane, Santorelli, Gillian, Lawlor, Debbie et al. (4 more authors) (2019) Blood pressure change across pregnancy in white British and Pakistani women: analysis of 1 data from the Born in Bradford cohort. Scientific Reports. ISSN 2045-2322

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Blood pressure change across pregnancy in white British and Pakistani women: analysis of data from the Born in Bradford cohort

Diane Farrar, Gillian Santorelli, Debbie A Lawlor, Derek Tuffnell, Trevor A Sheldon, Jane West, Corrie Macdonald-Wallis

Supplementary Table 1 Adjusted mean difference (95% CI) in SBP at eight weeks and change in systolic blood pressure in each period of gestation by hypertensive disorders of pregnancy (no HDP, gestational hypertension and preeclampsia)

Hypertensive disorder	Mean Difference in SBP at 8 wk, mmHg	Mean difference in average SBP change, mmHg/wk			
		8-24 weeks	24-30 weeks	30-36 weeks	≥36 weeks
White British women					
No HDP	0	0	0	0	0
Gestational ht	1.75 (-0.90 to 4.41)	0.11 (-0.11 to 0.32)	0.19 -0.24 to 0.63)	0.35 (-0.08 to 0.77)	-0.11 (-0.78 to 0.55)
Preeclampsia	-3.63 (-5.87 to -1.40)	-0.12 (-0.31 to 0.06)	0.28 -0.09 to 0.66)	-0.36 (-0.72 to 0.01)	-1.20 (-1.76 to -0.64)
Pakistani women					
No HDP	0	0	0	0	0
Gestational ht	1.19 (-2.24 to 4.63)	-0.06 (-0.33 to 0.21)	0.26 (-0.27 to 0.80)	0.93 (0.38 to 1.48)	0.94 (0.03 to 1.87)
Preeclampsia	-4.33 (-7.19 to -1.49)	0.25 (-0.47 to -0.01)	0.11(-0.33 to 0.55)	-0.32 (-0.76 to 0.11)	-0.35 (-1.06 to 0.35)

HDP= hypertensive disease of pregnancy

Adjusted for time to delivery and for maternal pregnancy booking BMI, age, parity, smoking in pregnancy status, education, gestational diabetes and infant gender. Reference category = women with no HDP.

In reference category: white British women (normotensive women), mean SBP at 8 wk (mm Hg) 112.45 (107.84 to 117.07); mean SBP change (mm Hg/wk): 8–24 wk 0.40 (0.01 to 0.78); 24–30 wk -0.37 (-1.14 to 0.41); 30–35⁶ wk 0.89 (0.13 to 1.64); >36 wk 3.67 (2.51 to 4.82)

In reference category: Pakistani women (normotensive women), mean SBP at 8 wk (mm Hg) 110.46 (104.65 to 116.26) mean SBP change (mm Hg/wk): 8–24 wk 0.65 (0.19 to 1.12); 24–30 wk -0.07 (-0.97 to 0.83); 30–36 wk 0.85 (-0.06 to 1.74); >36 wk 2.09 (0.66 to 3.52)

Supplementary Table 2 Adjusted mean difference (95% CI) in DBP at eight weeks and change in diastolic blood pressure in each period of gestation by hypertensive disorders of pregnancy (no HDP (reference), gestational hypertension and preeclampsia)

Hypertensive disorder	Mean Difference in DBP at 8 wk, mmHg	Mean difference in average DBP change, mmHg/wk			
		8-24 weeks	24-30 weeks	30-36 weeks	≥36 weeks
White British women					
No HDP	0	0	0	0	0
Gestational ht	1.62 (-0.27 to 0.30)	0.13 (-0.04 to 0.28)	-0.29 (-0.05 to 0.63)	0.07 (-0.28 to 3.54)	0.19 (-0.36 to 0.7)
Preeclampsia	-2.29 (-3.89 to -0.68)	0.00 (-0.13 to 0.14)	-0.27 (-0.02 to 0.56)	-0.40 (-0.70 to -0.10)	-0.43 (-0.89 to 0.02)
Pakistani women					
No HDP	0	0	0	0	0
Gestational ht	3.11 (0.59 to 5.63)	-0.04 (-0.25 to 0.16)	0.63 (0.22 to 1.03)	0.46 (0.04 to 0.87)	0.51 (-0.22 to 1.25)
Preeclampsia	-1.32 (-3.41 to 0.76)	-0.12 (-0.29 to 0.05)	0.20 (-0.13 to 0.53)	-0.27 (-0.61 to 0.06)	-0.64 (-1.21 to -0.08)

HDP= hypertensive disease of pregnancy

Adjusted for time to delivery and for maternal pregnancy booking BMI, age, parity, smoking in pregnancy status, education, gestational diabetes and gender. Reference category = women without a HDP.

In reference category: white British women (normotensive women), mean DBP at 8 wk (mm Hg) 66.594 (63.27 to 69.90) mean DBP change (mm Hg/wk): 8–24 wk 0.01 (-0.27 to 0.30); 24–30 wk -0.26 (-0.86 to 0.33); 30–36 wk 1.14 (0.54 to 1.75); >36 wk 2.21 (1.27 to 3.16)

In reference category: Pakistani women (normotensive women), mean DBP at 8 wk (mm Hg) 62.14 (57.88 to 66.39) mean DBP change (mm Hg/wk): 8–24 wk 0.38 (0.03 to 0.72); 24–30 wk -0.20 (-0.87 to 0.47); 30–36 wk 0.83 (0.15 to 1.52); >36 wk 2.69 (1.55 to 3.84)


Supplementary Table 3 Average length of gestation by standard deviations of blood pressure at eight weeks gestation and change in blood pressure in each period of pregnancy in the joint model adjusted for maternal covariates

Blood pressure variable	Length of gestation	
	Average length of gestation	95% CI
White British women		
Average SBP and DBP at 8 weeks and average change in SBP and DBP in each period of gestation		
SBP at 8 weeks (1 SD = 7.34mm Hg)		
+ 2 SD	39.87	39.70, 40.04
- 2 SD	39.70	39.50, 39.85
SBP change 8-24 weeks (1 SD = 0.32mm Hg/week)		
+ 2 SD	39.40	38.74, 39.76
- 2 SD	39.99	39.61, 40.67
SBP change 24-30 weeks (1 SD = 0.55mm Hg/week)		
+ 2 SD	40.01	35.03, 42.72
- 2 SD	39.52	33.04, 42.26
SBP change 30-36 weeks (1 SD = 0.66mm Hg/week)		
+ 2 SD	38.08	-147.72, 44.82
- 2 SD	41.06	-106.69, 44.86
SBP change >36 weeks (1 SD = 1.03 mm Hg/week)		
+ 2 SD	38.74	32.67, 41.67
- 2 SD	40.64	36.78, 42.78
DBP at 8 weeks (1 SD = 4.53mm Hg)		
+ 2 SD	39.91	39.69, 40.13
- 2 SD	39.64	39.40, 39.85
DBP change 8-24 weeks (1 SD = 0.12mm Hg/week)		
+ 2 SD	39.26	10.47, 44.13
- 2 SD	40.24	13.51, 44.21
DBP change 24-30 weeks (1 SD = 0.39mm Hg/week)		
+ 2 SD	39.43	31.33, 42.77
- 2 SD	40.09	32.78, 43.00
DBP change 30-36 weeks (1 SD = 0.66mm Hg/week)		
+ 2 SD	39.16	35.90, 41.70
- 2 SD	40.33	38.63, 42.24
DBP change >36 weeks (1 SD = 0.95 mm Hg/week)		
+ 2 SD	39.42	35.89, 41.70
- 2 SD	40.11	36.72, 42.00
Pakistani women		
SBP at 8 weeks (1 SD = 6.99mm Hg)		
+ 2 SD	39.53	39.37, 39.69


- 2 SD	39.73	39.34, 39.88
SBP change 8-24 weeks (1 SD =0.29mm Hg/week)		
+ 2 SD	39.54	38.90, 40.04
- 2 SD	39.73	39.20, 40.28
SBP change 24-30 weeks (1 SD =0.59mm Hg/week)		
+ 2 SD	39.58	36.77, 42.62
- 2 SD	39.52	32.88, 41.50
SBP change 30-36 weeks (1 SD = 0.71mm Hg/week)		
+ 2 SD	39.34	31.16, 42.75
- 2 SD	39.92	32.18, 42.92
SBP change >36 weeks (1 SD = 1.23mm Hg/week)		
+ 2 SD	39.09	36.27, 40.94
- 2 SD	40.13	37.89, 41.70
DBP at 8 weeks (1 SD = 5.04mm Hg)		
+ 2 SD	39.52	39.34, 39.70
- 2 SD	39.74	39.57, 39.91
DBP change 8-24 weeks (1 SD = 0.22mm Hg/week)		
+ 2 SD	39.25	37.28, 39.94
- 2 SD	39.99	39.31, 40.27
DBP change 24-30 weeks (1 SD =0.34mm Hg/week)		
+ 2 SD	39.45	33.65, 42.30
- 2 SD	39.80	34.33, 42.46
DBP change 30-36 weeks (1 SD = 0.57 mm Hg/week)		
+ 2 SD	39.30	35.83, 41.03
- 2 SD	39.95	37.74, 41.86
DBP change >36 weeks (1 SD = 1.11 mm Hg/week)		
+ 2 SD	39.42	37.12, 41.06
- 2 SD	39.84	37.69, 41.35

Predictions are for women in the reference category: women without a HDP, and for average levels of each of the blood pressure variables other than that being varied


Supplementary Figure 1a Predicted trajectories of systolic blood pressure across pregnancy for Pakistani women by maternal early-pregnancy BMI and hypertensive disorder. All trajectories are predicted by the fully adjusted joint model (model 2)




Supplementary Figure 1b Predicted trajectories of systolic blood pressure across pregnancy for white British women by maternal early-pregnancy BMI and hypertensive disorder of pregnancy




Supplementary Figure 2a Predicted trajectories of systolic blood pressure across pregnancy for Pakistani women by maternal age and hypertensive disorder of pregnancy




Supplementary Figure 2b Predicted trajectories of systolic blood pressure across pregnancy for white British women by maternal age and hypertensive disorder of pregnancy




Supplementary Figure 3a Predicted trajectories of systolic blood pressure across pregnancy for Pakistani women by parity and hypertensive disorder of pregnancy




Supplementary Figure 3b Predicted trajectories of systolic blood pressure across pregnancy for white British women by parity and hypertensive disorder of pregnancy




Supplementary Figure 4a Predicted trajectories of systolic blood pressure across pregnancy for Pakistani women by gestational diabetes and hypertensive disorder of pregnancy




Supplementary Figure 4b Predicted trajectories of systolic blood pressure across pregnancy for white British women by gestational diabetes and hypertensive disorder of pregnancy




Supplementary Figure 5a Predicted trajectories of systolic blood pressure across pregnancy for Pakistani women by maternal education and hypertensive disorder of pregnancy



Supplementary Figure 5b Predicted trajectories of systolic blood pressure across pregnancy for white British women by maternal education and hypertensive disorder of pregnancy




Supplementary Figure 6a Predicted trajectories of systolic blood pressure across pregnancy for white British women by smoking during pregnancy and hypertensive disorder of pregnancy




Too few Pakistani women smoked during pregnancy to be able to produce trajectories


Supplementary Figure 7a Predicted trajectories of systolic blood pressure across pregnancy for Pakistani women by infant gender and hypertensive disorder of pregnancy




Supplementary Figure 7b Predicted trajectories of systolic blood pressure across pregnancy for white British women by infant gender and hypertensive disorder of pregnancy




Supplementary Figure 8a Predicted trajectories of diastolic blood pressure across pregnancy for Pakistani women by maternal early-pregnancy BMI and hypertensive disorder of pregnancy




Supplementary Figure 8b Predicted trajectories of diastolic blood pressure across pregnancy for white British women by maternal early-pregnancy BMI and hypertensive disorder of pregnancy




Supplementary Figure 9a Predicted trajectories of diastolic blood pressure across pregnancy for Pakistani women by maternal age and hypertensive disorder of pregnancy order of pregnancy




Supplementary Figure 9b Predicted trajectories of diastolic blood pressure across pregnancy for white British women by maternal age and hypertensive disorder of pregnancy order of pregnancy




Supplementary Figure 10a Predicted trajectories of diastolic blood pressure across pregnancy for Pakistani women by parity and hypertensive disorder of pregnancy




Supplementary Figure 10b Predicted trajectories of diastolic blood pressure across pregnancy for white British women by parity and hypertensive disorder of pregnancy




Supplementary Figure 11a Predicted trajectories of diastolic blood pressure across pregnancy for Pakistani women by gestational diabetes and hypertensive disorder of pregnancy




Supplementary Figure 11b Predicted trajectories of diastolic blood pressure across pregnancy for white British women by gestational diabetes and hypertensive disorder of pregnancy




Supplementary Figure 12a Predicted trajectories of diastolic blood pressure across pregnancy for Pakistani women by maternal education and hypertensive disorder of pregnancy



Supplementary Figure 12b Predicted trajectories of diastolic blood pressure across pregnancy for white British women by maternal education and hypertensive disorder of pregnancy




Supplementary Figure 13a Predicted trajectories of diastolic blood pressure across pregnancy for white British women by smoking during pregnancy and hypertensive disorder of pregnancy



Too few Pakistani women smoked during pregnancy to be able to produce trajectories

Supplementary Figure 14a Predicted trajectories of diastolic blood pressure across pregnancy for Pakistani women by infant gender and hypertensive disorder of pregnancy



Supplementary Figure 14b Predicted trajectories of diastolic blood pressure across pregnancy for white British women by infant gender and hypertensive disorder of pregnancy

