



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/149600/>

Version: Accepted Version

Article:

Fordy, AP (2019) First integrals from conformal symmetries: Darboux–Koenigs metrics and beyond. *Journal of Geometry and Physics*, 145. 103475. ISSN: 0393-0440

<https://doi.org/10.1016/j.geomphys.2019.07.006>

Crown Copyright © 2019 Published by Elsevier B.V. All rights reserved. This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>.

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

First Integrals from Conformal Symmetries: Darboux-Koenigs Metrics and Beyond

Allan P. Fordy, School of Mathematics,
University of Leeds, Leeds LS2 9JT, UK.
e-mail a.p.fordy@leeds.ac.uk

July 23, 2019

Abstract

On spaces of constant curvature, the geodesic equations automatically have higher order integrals, which are just built out of first order integrals, corresponding to the abundance of Killing vectors. This is no longer true for general conformally flat spaces, but in this case there is a large algebra of *conformal* symmetries. In this paper we use these conformal symmetries to build higher order integrals for the geodesic equations. We use this approach to give a new derivation of the Darboux-Koenigs metrics, which have only *one* Killing vector, but two quadratic integrals. We also consider the case of possessing one Killing vector and two *cubic* integrals.

The approach allows the *quantum* analogue to be constructed in a simpler manner.

Keywords: Hamiltonian system; super-integrability; Poisson algebra; conformal algebra; quantum integrability; Darboux-Koenigs metrics.

MSC: 17B63,37J15,37J35,70H06,70H20

1 Introduction

In recent years there has been a burst of activity in the identification and classification of super-integrable systems, both classical and quantum (see the review [11] and references therein). Most of the interest is in Hamiltonians which are in “natural form” (the sum of kinetic and potential energies), with the kinetic energy being *quadratic* in momenta. A non-degenerate kinetic energy is associated with (pseudo-)Riemannian metric, so the leading order term in any integral defines a Killing tensor for this metric. For any Killing tensor, this leading order expression in momenta is itself a first integral for the geodesic equations, since it separately commutes with the kinetic energy. Thus, before even starting to consider which potentials can be added to a kinetic energy in order for the whole system to be completely (or even super-) integrable, we need to establish which *geodesic equations* are themselves completely (or even super-) integrable.

For a manifold with coordinates (q_1, \dots, q_n) , metric coefficients g_{ij} , with inverse g^{ij} , the geodesic equations are Hamiltonian, with *kinetic energy*

$$H = \frac{1}{2} \sum_{i,j=1}^n g^{ij} p_i p_j, \quad \text{where} \quad p_i = \sum_k g_{ik} \dot{q}_k. \quad (1)$$

For a metric with isometries, the infinitesimal generators (Killing vectors) give rise to first integrals, which are *linear* in momenta (Noether constants). When the space is either flat or constant curvature, it possesses the maximal group of isometries, which is of dimension $\frac{1}{2}n(n+1)$. In this case, (1) is actually the second order *Casimir* function of the symmetry algebra (see [6]). Furthermore *all* higher order integrals of the geodesic equations are built out of the above Noether constants by just taking polynomial expressions in them. This is an approach employed in [3, 5] for the quantum and classical cases.

Whilst most of the classification results and examples which occur in applications correspond to flat or constant curvature spaces, there are well known examples of conformally flat spaces (but *not* constant curvature), possessing quadratic invariants, which are clearly not just quadratic expressions in Noether constants. Specifically, there are the metrics found by Koenigs [9], which are described and analysed in [8, 7]. There are other examples of conformally flat spaces (but *not* constant curvature), possessing one Noether constant and a *cubic* integral (classified in [10]), which again *cannot* be represented as a cubic expression in the isometry algebra. The main purpose of this paper is to understand the origin of these integrals in terms of the conformal algebra associated with any conformally flat metric. Since the isometry group is a subgroup of the conformal group for a given metric and since every constant curvature metric is conformally flat, our approach will include the standard construction of higher order integrals in the constant curvature case.

In 2 dimensions, as is well known, the conformal group is *infinite*. For $n \geq 3$ this group is *finite* and has *maximal* dimension $\frac{1}{2}(n+1)(n+2)$, which is achieved for *conformally flat* spaces (which includes *flat* and *constant curvature* spaces). Any two conformally equivalent metrics have the same conformal group, so we can describe this in terms of the corresponding *flat* metric. In flat spaces, the infinitesimal generators consist of n *translations*, $\frac{1}{2}n(n-1)$ *rotations*, 1 *scaling* and n *inversions*, totalling $\frac{1}{2}(n+1)(n+2)$. This algebra is isomorphic to $\mathfrak{so}(n+1,1)$ (see Volume 1, p143, of [1]). Whilst the conformal algebra in two dimensional spaces is *infinite*, there still exists the 6 dimensional subalgebra described above (with $n = 2$). In this paper we only consider 2 dimensional systems, but if this is to be a stepping stone to dealing with higher dimensions, we should only consider this 6 dimensional subalgebra.

In Section 2 we list some useful properties of the 6 dimensional conformal algebra of the standard Euclidean metric in 2 dimensions, giving (initially) its Poisson representation in Cartesian coordinates (x, y) and the table of their Poisson relations. We derive some other coordinate systems, related to commuting pairs, and discuss some 3 dimensional subalgebras, which play a role in our construction.

The main idea of the paper is developed in Section 3, where we consider quadratic *conformal invariants* (together with a linear *invariant*) and *systematically* derive the coefficients corresponding to these being *true invariants*. Inevitably, we are led to the Darboux-Koenigs class of metrics [9] (other than flat and constant curvature cases).

In Section 4, this idea is used to construct a class of metric with a linear integral and a pair of third order integrals. We derive Case 3 of the classification given in [10]. Closure of the Poisson algebra leads to an interesting restriction of this metric.

In Section 5, we consider the *quantum* extension of the above results. For the Darboux-Koenigs class of metrics, the quantum and classical formulae are identical at leading order, with some lower order terms being added to some formulae. However, the quantum version of the system with third order integrals is further restricted to a constant curvature case. The isometries, as a subalgebra of the conformal symmetries, are identified and expressions for the integrals in terms of these are derived. The general problem of extending quadratic and cubic integrals to the quantum regime is discussed in detail in [2].

2 The 2D Euclidean Metric and its Conformal Algebra

Consider metrics which are conformally related to the standard Euclidean metric in 2 dimensions, with Cartesian coordinates (x, y) . The corresponding kinetic energy (1) takes the form

$$H = \psi(x, y)(p_x^2 + p_y^2). \quad (2)$$

As discussed in the introduction, the conformal algebra of this 2D metric is *infinite dimensional*, but we can still write down the $\frac{1}{2}(n+1)(n+2)$ (in this case 6) conformal elements

$$\begin{aligned} X_1 &= p_x, & X_2 &= p_y, & X_3 &= yp_x - xp_y, \\ X_4 &= xp_x + yp_y, & X_5 &= (x^2 - y^2)p_x + 2xyp_y, & X_6 &= 2xyp_x + (y^2 - x^2)p_y. \end{aligned} \quad (3)$$

Table 1: The 6–dimensional conformal algebra

	X_1	X_2	X_3	X_4	X_5	X_6
X_1	0	0	X_2	$-X_1$	$-2X_4$	$-2X_3$
X_2		0	$-X_1$	$-X_2$	$2X_3$	$-2X_4$
X_3			0	0	$-X_6$	X_5
X_4				0	$-X_5$	$-X_6$
X_5					0	0
X_6						0

The elements $\{X_i\}_{i=1}^6$ satisfy the Poisson relations depicted in Table 1.

These conformal elements satisfy

$$\{H, X_i\} = \lambda_i H, \quad (4a)$$

where H is given by (2) and

$$\begin{aligned} \lambda_1 &= \frac{\psi_x}{\psi}, \quad \lambda_2 = \frac{\psi_y}{\psi}, \quad \lambda_3 = \frac{y\psi_x - x\psi_y}{\psi}, \quad \lambda_4 = \frac{x\psi_x + y\psi_y - 2\psi}{\psi}, \\ \lambda_5 &= \frac{(x^2 - y^2)\psi_x + 2xy\psi_y - 4x\psi}{\psi}, \quad \lambda_6 = \frac{2xy\psi_x + (y^2 - x^2)\psi_y - 4y\psi}{\psi}. \end{aligned} \quad (4b)$$

2.1 Some Associated Coordinate Systems

Whenever we have $\{X_i, X_j\} = 0$, then there exists a coordinate system (u, v) for which $X_i = p_u$, $X_j = p_v$. They are defined by

$$\{u, X_i\} = 1, \quad \{u, X_j\} = 0 \quad \text{and} \quad \{v, X_i\} = 0, \quad \{v, X_j\} = 1.$$

In the case of X_1, X_2 , we already have the coordinates (x, y) .

Corresponding to $\{X_3, X_4\} = 0$, we have coordinates (u, v) , given by $u = \arctan\left(\frac{x}{y}\right)$, $v = \frac{1}{2} \log(x^2 + y^2)$. The generating function, $S = e^v(\sin u p_x + \cos u p_y)$, gives

$$x = e^v \sin u, \quad y = e^v \cos u, \quad p_x = e^{-v}(p_u \cos u + p_v \sin u), \quad p_y = e^{-v}(p_v \cos u - p_u \sin u).$$

In these coordinates, we have

$$\begin{aligned} X_1 &= e^{-v}(p_u \cos u + p_v \sin u), \quad X_2 = e^{-v}(p_v \cos u - p_u \sin u), \quad X_3 = p_u, \\ X_4 &= p_v, \quad X_5 = e^v(p_v \sin u - p_u \cos u), \quad X_6 = e^v(p_v \cos u + p_u \sin u). \end{aligned} \quad (5)$$

We later need the conformal factors $\{H, X_i\} = \lambda_i H$, in these coordinates:

$$\begin{aligned} \lambda_1 &= \frac{e^{-v}(\cos u \psi_u + \sin u(\psi_v + 2\psi))}{\psi}, \quad \lambda_2 = \frac{e^{-v}(\cos u(\psi_v + 2\psi) - \sin u \psi_u)}{\psi}, \quad \lambda_3 = \frac{\psi_u}{\psi}, \\ \lambda_4 &= \frac{\psi_v}{\psi}, \quad \lambda_5 = \frac{e^v(\sin u(\psi_v - 2\psi) - \cos u \psi_u)}{\psi}, \quad \lambda_6 = \frac{e^v(\sin u \psi_u + \cos u(\psi_v - 2\psi))}{\psi}. \end{aligned} \quad (6)$$

Corresponding to $\{X_5, X_6\} = 0$, we have coordinates (u, v) , given by $u = -\frac{x}{x^2+y^2}$, $v = -\frac{y}{x^2+y^2}$. The generating function, $S = -\frac{x}{x^2+y^2} p_u - \frac{y}{x^2+y^2} p_v$, gives

$$x = -\frac{u}{u^2+v^2}, \quad y = -\frac{v}{u^2+v^2}, \quad p_x = (u^2-v^2)p_u + 2uvp_v, \quad p_y = 2uvp_u - (u^2-v^2)p_v.$$

In these coordinates, we have

$$\begin{aligned} X_1 &= (u^2-v^2)p_u + 2uvp_v, & X_2 &= 2uvp_u + (v^2-u^2)p_v, & X_3 &= vp_u - up_v, \\ X_4 &= -up_u - vp_v, & X_5 &= p_u, & X_6 &= p_v. \end{aligned} \tag{7}$$

2.2 Involutions and Casimir Functions

The algebra (3) possesses a pair of involutive automorphisms:

$$\iota_1 : (x, y) \mapsto (y, x), \quad \iota_2 : (x, y) \mapsto \left(\frac{-x}{x^2+y^2}, \frac{-y}{x^2+y^2} \right),$$

under which the elements X_i transform as follows:

	X_1	X_2	X_3	X_4	X_5	X_6
$\iota_1 :$	X_2	X_1	$-X_3$	X_4	X_6	X_5
$\iota_2 :$	X_5	X_6	X_3	$-X_4$	X_1	X_2

The action of ι_1 is obvious, while that of ι_2 can be seen by comparing the formulae (3) and (7).

It can be shown that the abstract algebra depicted in Table 1 has 2 quadratic Casimirs:

$$\mathcal{C}_1 = X_3^2 - X_4^2 + X_1X_5 + X_2X_6, \quad \mathcal{C}_2 = 2X_3X_4 + X_2X_5 - X_1X_6,$$

which are multiples of the identity in the matrix (adjoint) representation, but both vanish in this Poisson representation, so are *quadratic constraints* on the Poisson algebra. Under the action of the above involutions we have $\mathcal{C}_1 \mapsto \mathcal{C}_1$ and $\mathcal{C}_2 \mapsto -\mathcal{C}_2$.

2.3 3-dimensional Subalgebras

This conformal algebra contains several 3-dimensional subalgebras:

1. $\mathfrak{g}_{(123)} = \{X_1, X_2, X_3\}$ is the Euclidean algebra with Casimir

$$H_0^{123} = (X_1^2 + X_2^2) = (p_x^2 + p_y^2),$$

corresponding to a flat metric.

2. $\mathfrak{g}_{(563)} = \{X_5, X_6, X_3\}$ is the Euclidean algebra with Casimir

$$H_0^{563} = (X_5^2 + X_6^2) = (x^2 + y^2)^2 (p_x^2 + p_y^2),$$

corresponding to a flat metric.

3. $\mathfrak{g}_{(145)} = \{X_1, X_4, X_5\}$ is the algebra $\mathfrak{sl}(2)$ with Casimir

$$H_0^{145} = (X_4^2 - X_1X_5) = y^2 (p_x^2 + p_y^2),$$

corresponding to a metric with constant (non-zero) curvature.

4. $\mathfrak{g}_{(246)} = \{X_2, X_4, X_6\}$ is the algebra $\mathfrak{sl}(2)$ with Casimir

$$H_0^{246} = (X_4^2 - X_2X_6) = x^2(p_x^2 + p_y^2),$$

corresponding to a metric with constant (non-zero) curvature.

These are clearly conformally equivalent. Under the action of ι_1 , we have

$$\mathfrak{g}_{(123)} \rightarrow \mathfrak{g}_{(123)}, \quad \mathfrak{g}_{(563)} \rightarrow \mathfrak{g}_{(563)}, \quad \mathfrak{g}_{(145)} \leftrightarrow \mathfrak{g}_{(246)},$$

whilst under the action of ι_2 , we have

$$\mathfrak{g}_{(123)} \leftrightarrow \mathfrak{g}_{(563)}, \quad \mathfrak{g}_{(145)} \rightarrow \mathfrak{g}_{(145)}, \quad \mathfrak{g}_{(246)} \rightarrow \mathfrak{g}_{(246)}.$$

3 Geodesic Flows in 2D with Linear and Quadratic Integrals

Here we consider the Hamiltonian (2), with general function $\psi(x, y)$. We see that we cannot expect any of λ_i , given by (4b) (or even for any linear combination $\sum_{i=1}^6 \alpha_i \lambda_i$), to be *zero*. When such a combination is *zero*, we have an *invariant* (a *Killing vector*), not just a conformal invariant (a *conformal Killing vector*). In this section, we consider Hamiltonians (2) which possess exactly *one* such linear invariant.

Remark 3.1 *By a theorem of Darboux and Koenigs, if such a metric possesses at least two Killing vectors, then it possesses three and the space has constant curvature (see [8, 7]).*

In fact, we will be more restrictive, and only consider systems for which one of the *basis elements* X_i is itself an invariant. In this case we can use either the coordinates (x, y) or one of the (u, v) systems of Section 2.1.

A *quadratic conformal invariant* is any expression of the form

$$F = \sum_{i,j=1}^6 \beta_{ij} X_i X_j + \sigma(x, y) H, \quad (8)$$

where β_{ij} is any symmetric matrix of (*constant*) coefficients and $\sigma(x, y)$ is an arbitrary function of (x, y) (or, indeed, some appropriate coordinates (u, v)), which satisfies

$$\{F, H\} = (\mu_1(x, y)p_x + \mu_2(x, y)p_y)H,$$

where $\mu_i(x, y)$ are some functions.

We can ask whether there is a choice of β_{ij} and $\sigma(x, y)$ for which $\mu_i(x, y) \equiv 0$, in which case F is a *quadratic invariant*. In fact, we have more structure. If *both* $K = X_i$ (for some i) *and* F are invariants, then so are $\{F, K\}$, $\{\{F, K\}, K\}$, etc. Simple choices of β_{ij} lead to simple equations for $\psi(x, y)$ and $\sigma(x, y)$ and quickly lead to the 4 solutions found by Darboux and Koenigs. Following [8, 7], we label these D_1 to D_4 . To avoid the flat and constant curvature cases, the term $\sigma(x, y)H$ in (8) is essential. All the Darboux and Koenigs metrics reduce to flat and constant curvature cases when the corresponding parameter vanishes.

Such simple choices can be made by considering the 3 dimensional subalgebras given in Section 2.3. We choose a subalgebra that contains our linear invariant K . To avoid unnecessary calculations we also employ the involutions ι_1 and ι_2 of Section 2.2, which means we only need to consider K to be one of X_2, X_3 and X_4 , and to be elements of the algebras $\mathfrak{g}_{(123)}$ or $\mathfrak{g}_{(246)}$ (see the comments at the end of Section 2.3).

3.1 The Case when $\{H, X_2\} = 0$

In this case we use the coordinates (x, y) , so that $X_2 = p_y$, with

$$H = \varphi(x)(p_x^2 + p_y^2) = \varphi(x)(X_1^2 + X_2^2).$$

Note that, for any function $a(y)$, we have $\{a(y), H\} = 2a'(y)\varphi(x)X_2$.

Since X_2 belongs to two of our subalgebras, there are two subcases.

3.1.1 The Algebra $\mathfrak{g}_{(123)}$, with $\{H, X_2\} = 0$

Noting that $\{X_3, X_2\} = X_1$ and $\{X_1, X_2\} = 0$, we define $F_2 = X_2X_3 + a(y)H$, and then

$$F_1 = \{F_2, X_2\} = X_2X_1 + a'(y)H \quad \text{and} \quad \{F_1, X_2\} = a''(y)H$$

should *both* be first integrals *if* F_2 is. We therefore have

$$\{a''(y)H, H\} = 2a'''(y)\varphi(x)X_2H = 0 \quad \Rightarrow \quad a'''(y) = 0.$$

The condition

$$\{F_2, H\} = 0 \quad \Rightarrow \quad (-\lambda_3 + 2a_y\varphi)X_2H = 0, \quad (9)$$

where $\lambda_3 = \frac{y\varphi_x}{\varphi}$ in this case (see (4b)). This is a *separable equation* for $\varphi(x)$ and $a(y)$, leading to

$$\varphi(x) = \frac{1}{\alpha x + \beta}, \quad a(y) = -\frac{\alpha}{4}y^2.$$

We then find

$$F_1 = \{F_2, X_2\} = X_1X_2 - \frac{\alpha}{2}yH,$$

and

$$\{F_1, X_2\} = -\frac{\alpha}{2}H, \quad \{F_1, F_2\} = 2X_2^3 + \beta HX_2.$$

Of course, in 2 degrees of freedom we can only have 3 functionally independent integrals (including H), so the above 4 integrals satisfy a polynomial constraint:

$$F_1^2 + \alpha F_2H + X_2^4 - \beta HX_2^2 = 0.$$

We have obtained, in this way, the first of the Darboux-Koenigs systems D_1 , which is generally non-constant curvature, but reduces to a flat metric when $\alpha = 0$.

Remark 3.2 *The choice of $\sigma(x, y) = a(y)$ guaranteed the factorisation of equation (9), leading to a single equation for the two functions $\varphi(x)$ and $a(y)$.*

Remark 3.3 (The symmetry $x \leftrightarrow y$) *There is an equivalent version of this example, using X_1 in place of X_2 , which can be obtained by just switching $x \leftrightarrow y$, not forgetting that $X_3 \mapsto -X_3$ under this transformation. This is just the involution ι_1 .*

3.1.2 The Algebra $\mathfrak{g}_{(246)}$, with $\{H, X_2\} = 0$

Noting that $\{X_6, X_2\} = 2X_4$ and $\{X_4, X_2\} = X_2$, we define $F_2 = X_2X_6 + a(y)H$, and then

$$F_1 = \frac{1}{2}\{F_2, X_2\} = X_2X_4 + \frac{1}{2}a'(y)H \quad \text{and} \quad \{F_1, X_2\} = X_2^2 + \frac{1}{2}a''(y)H,$$

the latter being a first integral if $a'''(y) = 0$.

The condition

$$\{F_2, H\} = 0 \quad \Rightarrow \quad (-\lambda_6 + 2a_y\varphi)X_2H = 0, \quad (10)$$

where $\lambda_6 = \frac{2xy\varphi_x - 4y\varphi}{\varphi}$ in this case (see (4b)). This is a *separable equation* for $\varphi(x)$ and $a(y)$, leading to

$$\varphi(x) = \frac{x^2}{a_2 - a_1x^2}, \quad a(y) = a_1y^2.$$

We then find

$$F_1 = \frac{1}{2}\{F_2, X_2\} = X_2X_4 + a_1yH,$$

and

$$\{F_1, X_2\} = a_1 H + X_2^2, \quad \{F_1, F_2\} = -2X_2(F_2 + a_2 H).$$

These elements satisfy the polynomial constraint

$$F_1^2 = (a_1 H + X_2^2)F_2 + a_2 H X_2^2.$$

We have obtained, in this way, the second of the Darboux-Koenigs systems D_2 , which reduces to a constant curvature metric when $a_1 = 0$.

3.2 The Case when $\{H, X_3\} = 0$

In this case we use the coordinates (u, v) , associated with $\{X_3, X_4\} = 0$ and use the formulae (5), so that $X_3 = p_u$, so we consider

$$H = \varphi(v)(p_u^2 + p_v^2) = \varphi(v)(X_3^2 + X_4^2).$$

Note that, for any function $a(u)$, we have $\{a(u), H\} = 2a'(u)\varphi(v)X_3$.

In this case we consider the algebra $\mathfrak{g}_{(123)}$, noting that $\{X_1, X_3\} = X_2$ and $\{X_2, X_3\} = -X_1$. We define $F_1 = X_1 X_3 + a(u)H$, and then

$$F_2 = \{F_1, X_3\} = X_2 X_3 + a'(u)H \quad \text{and} \quad \{F_2, X_3\} = -X_1 X_3 + a''(u)H = -F_1,$$

if $a''(u) = -a(u)$.

The condition

$$\{F_1, H\} = 0 \quad \Rightarrow \quad (-\lambda_1 + 2a_u \varphi)X_3 H = 0, \tag{11}$$

where $\lambda_1 = \frac{e^{-v} \sin u(\varphi_v + 2\varphi)}{\varphi}$ in this case (see (6)). This is a *separable equation* for $\varphi(v)$ and $a(u)$, leading to

$$\varphi(v) = \frac{e^{-v}}{\beta e^v - 2\alpha}, \quad a(u) = \alpha \cos u.$$

We then find

$$F_2 = \{F_1, X_3\} = X_2 X_3 - \alpha \sin u H,$$

and

$$\{F_2, X_3\} = -F_1, \quad \{F_1, F_2\} = \beta X_3 H.$$

These elements satisfy the polynomial constraint

$$F_1^2 + F_2^2 = \beta H X_3^2 + \alpha^2 H^2.$$

We have obtained, in this way, the third of the Darboux-Koenigs systems D_3 , which is generally non-constant curvature, but reduces to a flat metric when $\alpha = 0$.

3.3 The Case when $\{H, X_4\} = 0$

In this case we use the coordinates (u, v) , associated with $\{X_3, X_4\} = 0$ and use the formulae (5), so that $X_4 = p_v$, so we consider

$$H = \varphi(u)(p_u^2 + p_v^2) = \varphi(u)(X_3^2 + X_4^2).$$

Note that, for any function $a(v)$, we have $\{a(v), H\} = 2a'(v)\varphi(u)X_4$.

In this case we consider the algebra $\mathfrak{g}_{(246)}$, noting that $\{X_2, X_4\} = -X_2$ and $\{X_6, X_4\} = X_6$.

We first consider $F_1 = X_2 X_4 + a(v)H$ and the condition

$$\{F_1, H\} = 0 \quad \Rightarrow \quad (-\lambda_2 + 2a_v \varphi)X_4 H = 0, \tag{12}$$

where $\lambda_2 = \frac{e^{-v}(2\varphi \cos u - \sin u \varphi_u)}{\varphi}$ in this case (see (6)). This is a *separable equation* for $a_v = \alpha e^{-v}$, leading to

$$-2 \cos u \varphi + 2\alpha \varphi^2 + \sin u \varphi_u = 0 \quad \Rightarrow \quad \varphi(u) = \frac{\sin^2 u}{\beta - 2\alpha \cos u}.$$

This has given us $F_1 = X_2 X_4 - \alpha e^{-v} H$, with $\{F_1, X_4\} = -F_1$.

The canonical transformation $(v, p_v) \mapsto (-v, -p_v)$ corresponds to the Lie algebraic involution $(X_2, X_4, X_6) \mapsto (-X_6, -X_4, -X_2)$, as well as $H \mapsto H$, leading to

$$F_2 = X_6 X_4 - \alpha e^v H.$$

The remaining Poisson relations are

$$\{F_1, X_4\} = -F_1, \quad \{F_2, X_4\} = F_2, \quad \{F_1, F_2\} = 2X_4(\beta H - 2X_4^2).$$

These elements satisfy the polynomial constraint

$$F_1 F_2 = \alpha^2 H^2 + X_4^2 (X_4^2 - \beta H).$$

We have obtained, in this way, the fourth of the Darboux-Koenigs systems D_4 , which reduces to a constant curvature metric when $\alpha = 0$.

4 Geodesic Flows in 2D with Linear and Cubic Integrals

We again consider a Hamiltonian of the form (2), but now possessing *linear* and *cubic* integrals. Such metrics were classified in [10], where it was found that there are three cases, each with a specific form of cubic integral. Here we use the conformal algebra to build the cubic integral corresponding to their Case 3.

Specifically, we consider the case for which $\{X_2, H\} = 0$, so we use Cartesian coordinates (x, y) and consider the Hamiltonian

$$H = \varphi(x)(p_x^2 + p_y^2) = \varphi(x)(X_1^2 + X_2^2), \quad \text{which satisfies} \quad \{X_2, H\} = 0.$$

We again build a cubic *conformal invariant* out of the *linear elements* X_i and *quadratic elements* of the form $\sigma(x, y)H$ and restrict the coefficients by asking for it to be a *true invariant*.

We mirror the structure found in Sections 3.1.1 and 3.1.2, but can no longer restrict to a single 3 dimensional subalgebra. However, we still use the structure

$$\{X_3, X_2\} = X_1, \quad \{X_1, X_2\} = 0, \quad \{X_6, X_2\} = 2X_4 \quad \text{and} \quad \{X_4, X_2\} = X_2. \quad (13)$$

We build F_1 and F_2 , satisfying

$$\{F_2, X_2\} = F_1 \quad \text{and} \quad \{F_1, X_2\} = X_2(c_0 H + c_1 X_2^2), \quad (14)$$

where this last term is the general cubic integral involving only X_2 and H .

4.1 The Integral F_1

The second of (14) is a linear equation for F_1 , which we can solve:

$$F_1 = c_0 y X_2 H + c_1 X_4 X_2^2 + X_1(c_2 X_2^2 + b(x)H), \quad (15)$$

where the first two terms are just a ‘‘particular solution’’, with the remaining part being in the *kernel* of the linear map $F \mapsto \{F, X_2\}$.

Remark 4.1 *There are 3 other possible elements of the kernel (excluding X_2^3 , which is, itself, a first integral):*

$$X_1^2(k_1X_1 + k_2X_2) + \gamma(x)HX_2.$$

Writing $X_1^2 = \varphi^{-1}H - X_2^2$, the first two terms just redefine c_2 and $b(x)$, modify $\gamma(x)$ and add a multiple of the first integral X_2^3 , so we can remove these two terms. The calculation of $\{F_2, H\} = 0$ then gives $\gamma'(x) = 0$, so this is just a constant multiple of X_2H , so can also be removed. Hence, the formula (15) is the general solution of $\{F_1, X_2\} = X_2(c_0H + c_1X_2^2)$.

With F_1 defined by (15), we find

$$\{F_1, H\} = (2c_0\varphi - c_1\lambda_4 - c_2\lambda_1 - 2b_x\varphi)X_2^2H + (2b_x - \lambda_1b)H^2 = 0,$$

where, in this case,

$$\lambda_1 = \frac{\varphi_x}{\varphi}, \quad \lambda_4 = \frac{x\varphi_x - 2\varphi}{\varphi},$$

leading to

$$b\varphi_x = 2\varphi b_x \quad \text{and} \quad (c_2 + c_1x)\varphi_x = 2(c_1 + c_0\varphi - b_x\varphi)\varphi,$$

giving

$$\varphi = b^2 \quad \text{and} \quad b_x = \frac{b(c_1 + c_0b^2)}{c_2 + c_1x + b^3}. \quad (16)$$

4.2 The Integral F_2

Given this solution for F_1 , we can solve $\{F_2, X_2\} = F_1$ for F_2 (using the relations (13)):

$$F_2 = \frac{1}{2}c_0y^2X_2H + \frac{1}{2}c_1X_6X_2^2 + X_3(c_2X_2^2 + b(x)H) + b_1(x)HX_2, \quad (17)$$

where the last term commutes with X_2 . We then find

$$\{F_2, H\} = \left(2c_0y\varphi - \frac{1}{2}c_1\lambda_6 - c_2\lambda_3\right)X_2^2H - \lambda_3bH^2 + 2b_x\varphi X_1X_2H + 2b_{1x}\varphi X_1X_2H = 0,$$

where, in this case,

$$\lambda_3 = \frac{y\varphi_x}{\varphi}, \quad \lambda_6 = \frac{2y(x\varphi_x - 2\varphi)}{\varphi}.$$

If we write

$$X_3 = yX_1 - xX_2, \quad \varphi X_1^2 = H - \varphi X_2^2,$$

then

$$\left(2c_0\varphi - c_1\left(\frac{x\varphi_x - 2\varphi}{\varphi}\right) - c_2\frac{\varphi_x}{\varphi} - 2b_x\varphi\right)yX_2^2H + \left(2b_x - \frac{b\varphi_x}{\varphi}\right)H^2 + 2(b_{1x} - xb_x)\varphi X_1X_2H = 0.$$

These coefficients give

$$b_{1x} = xb_x, \quad b\varphi_x = 2\varphi b_x \quad \text{and} \quad (c_2 + c_1x + b\varphi)\varphi_x = 2(c_1 + c_0\varphi)\varphi,$$

giving, again, (16) and an additional function $b_1(x)$, obtained by integrating $b_{1x} = xb_x$.

Remark 4.2 *The scalar curvature of this metric is given by $R = 2(bb_{xx} - b_x^2)$.*

Remark 4.3 (Equivalence with Case 3 of [10]) In [10], the metric coefficient $\varphi(x)$ is denoted h_x^2 and for case (iii) of equation (1.2), $h(x)$ satisfies

$$h_x(A_0 h_x^2 - A_1 h + A_2) = A_3 x + A_4.$$

Differentiating this and eliminating $h(x)$ itself, we find

$$h_{xx} = \frac{h_x(A_3 + A_1 h_x^2)}{A_4 + A_3 x + 2A_0 h_x^3},$$

which is exactly our equation (16) for $b = h_x$, with $c_0 = \frac{1}{2}A_1$, $c_1 = \frac{1}{2}A_3$, $c_2 = \frac{1}{2}A_4$ and $A_0 = 1$ (absorbed into the definition of b).

4.3 Closing the Poisson Algebra

We have the relations (14), subject to (16) and $b_{1x} = xb_x$. The Jacobi identity implies

$$\{\{F_1, F_2\}, X_2\} = \{\{F_1, X_2\}, F_2\} = -(c_0 H + 3c_1 X_2^2)F_1 = -\{(c_0 H + 3c_1 X_2^2)F_2, X_2\},$$

so we can write

$$\{F_1, F_2\} + (c_0 H + 3c_1 X_2^2)F_2 = d_1 H^2 X_2 + d_2 H X_2^3 + d_3 X_2^5. \quad (18)$$

Comparing coefficients of $p_x^i p_y^{5-i}$ in this equation, we find d_1 and d_2 are arbitrary, $d_3 = 3c_2^2$ and

$$b_1 = \frac{2c_2 + 2c_1 x + d_1 b + 2c_0 x b^2 - b^3}{2c_0 b}, \quad (19)$$

and $b(x)$ is constrained to satisfy the quartic equation:

$$P_4 = 2c_1 b^4 - 4c_0(c_2 + c_1 x)b^3 + (2c_0 x(2c_2 + c_1 x) + 2c_1 d_1 - c_0 d_2)b^2 - 4c_1(c_2 + c_1 x)b + 2c_0(c_2 + c_1 x)^2 = 0. \quad (20)$$

Remarkably, these are consistent with the formulae (16) and $b_{1x} = xb_x$. Differentiating (19) with respect to x and replacing $b'(x)$ by the formula in (16), gives

$$b_{1x} = \frac{xb(c_1 + c_0 b^2)}{c_2 + c_1 x + b^3},$$

which is just $b_{1x} = xb_x$. Differentiating (20) with respect to x and replacing $b'(x)$ by the formula in (16), gives

$$P_{4x} = \frac{2(c_1 + c_0 b^2)}{c_2 + c_1 x + b^3} P_4 = \frac{2b'}{b} P_4,$$

which allows the solution $P_4 = 0$.

These integrals satisfy the constraint

$$F_1^2 - 2X_2(c_0 H + c_1 X_2^2)F_2 = H^3 - c_2^2 X_2^6 - d_1 X_2^2 H^2 - \frac{1}{2} d_2 H X_2^4. \quad (21)$$

5 The Quantum Case

For a given metric g_{ij} , we have the classical kinetic energy and its quantum analogue (the Laplace-Beltrami operator):

$$H = \frac{1}{2} \sum_{i=0}^n g^{ij} p_i p_j \quad \text{and} \quad L_b f = \sum_{i,j=1}^n g^{ij} \nabla_i \nabla_j f = \sum_{i,j=1}^n \frac{1}{\sqrt{g}} \frac{\partial}{\partial x^j} \left(\sqrt{g} g^{ij} \frac{\partial f}{\partial x^i} \right),$$

where g is the determinant of the matrix g_{ij} . However, there is no such general algorithm for finding the quantum analogue of other quadratic and higher order integrals. Nevertheless, there is an algorithm for finding the quantum analogue of a *first order* integral or conformal invariant, and this will be exploited in this section.

For a metric with isometries, the infinitesimal generators (Killing vectors) are just first order differential operators which commute with the Laplace-Beltrami operator L_b . When the space is either flat or constant curvature, it possesses the maximal group of isometries, which is of dimension $\frac{1}{2}n(n+1)$. In this case, L_b is proportional to the second order *Casimir* function of the symmetry algebra (see [6]).

Recall that if f, g are any functions of (\mathbf{q}, \mathbf{p}) , then the Hamiltonian vector field of f is

$$X_f = \sum_{i=1}^3 (\{q_i, f\} \partial_{q_i} + \{p_i, f\} \partial_{p_i}) \quad \text{and} \quad [X_f, X_g] = -X_{\{f, g\}}. \quad (22)$$

Functions which are linear in momenta define vector fields on configuration space, with coordinates (q_1, \dots, q_n) . For any function on configuration space, $f(q_1, \dots, q_n)$, we have:

$$h(\mathbf{q}, \mathbf{p}) = \sum_{i=1}^n a_i(\mathbf{q}) p_i \quad \Rightarrow \quad L_h f = \{f, h\} = \sum_{i=1}^n a_i(\mathbf{q}) \frac{\partial f}{\partial q_i}, \quad (23)$$

and it follows from (22) that $[L_f, L_g] = -L_{\{f, g\}}$. This is one of the advantages of expressing integrals in terms of conformal (or even “true”) invariants. We can immediately *quantise* any classical system which is built in this way. This idea is used in [3, 4].

5.1 The Conformal Algebra

We use (23) to define $L_i = L_{X_i}$ for each of the elements of the conformal algebra (3). We need expressions in both Cartesian coordinates (x, y) and the coordinates (u, v) , related to $\{X_3, X_4\} = 0$.

In Cartesian coordinates (x, y) , we have

$$\begin{aligned} L_1 &= \partial_x, & L_2 &= \partial_y, & L_3 &= y\partial_x - x\partial_y, & L_4 &= x\partial_x + y\partial_y, \\ L_5 &= (x^2 - y^2)\partial_x + 2xy\partial_y, & L_6 &= 2xy\partial_x + (y^2 - x^2)\partial_y. \end{aligned} \quad (24a)$$

These satisfy the commutation rules $[L_i, L_j] = -L_{\{X_i, X_j\}}$, where $\{X_i, X_j\}$ can be found in Table 1. These operators are conformal Killing vectors of the Laplace-Beltrami operator $L_b = \psi(x, y)(\partial_x^2 + \partial_y^2)$, corresponding to the general H of (2), satisfying $[L_i, L_b] = \lambda_i L_b$, where λ_i are given by (4b).

In coordinates (u, v) , we have

$$\begin{aligned} L_1 &= e^{-v}(\cos u \partial_u + \sin u \partial_v), & L_2 &= e^{-v}(\cos u \partial_v - \sin u \partial_u), & L_3 &= \partial_u, \\ L_4 &= \partial_v, & L_5 &= e^v(\sin u \partial_v - \cos u \partial_u), & L_6 &= e^v(\sin u \partial_u + \cos u \partial_v). \end{aligned} \quad (24b)$$

5.2 The Quantum Darboux-Koenigs Cases

We now carry out this procedure for each of the Darboux-Koenigs cases of Sections 3.1 to 3.3. The only change to the formulae is the symmetrisation of products, together with the addition of a few lower order terms to some formulae.

5.2.1 The Quantum Darboux-Koenigs Case D_1

We just take the formulae directly from Section 3.1.1, giving $L_b = \frac{1}{\alpha x + \beta}(\partial_x^2 + \partial_y^2)$. However, since L_i are operators (not just functions on phase space), we replace a simple product by a *symmetric product*, although

this is unnecessary when two operators commute. We have Killing vector L_2 and define

$$\begin{aligned} F_1 &= L_1 L_2 - \frac{\alpha}{2} y L_b = \partial_x \partial_y - \frac{\alpha y}{2(\alpha x + \beta)} (\partial_x^2 + \partial_y^2), \\ F_2 &= \frac{1}{2} (L_2 L_3 + L_3 L_2) - \frac{\alpha}{4} y^2 L_b = y \partial_x \partial_y - x \partial_y^2 + \frac{1}{2} \partial_x - \frac{\alpha y^2}{4(\alpha x + \beta)} (\partial_x^2 + \partial_y^2), \end{aligned}$$

which satisfy

$$[L_2, F_1] = -\alpha L_b, \quad [L_2, F_2] = F_1, \quad [F_1, F_2] = -2L_2^3 + \beta L_b L_2, \quad F_1^2 + \alpha F_2 L_b + L_2^4 - \beta L_b L_2^2 = 0.$$

5.2.2 The Quantum Darboux-Koenigs Case D_2

We now take the formulae from Section 3.1.2, with Killing vector L_2 , giving:

$$\begin{aligned} L_b &= \frac{x^2}{a_2 - a_1 x^2} (\partial_x^2 + \partial_y^2), \\ F_1 &= \frac{1}{2} (L_2 L_4 + L_4 L_2) + a_1 y L_b = x \partial_x \partial_y + y \partial_y^2 + \frac{1}{2} \partial_y + \frac{a_1 x^2 y}{a_2 - a_1 x^2} (\partial_x^2 + \partial_y^2), \\ F_2 &= \frac{1}{2} (L_2 L_6 + L_6 L_2) + a_1 y^2 L_b = 2xy \partial_x \partial_y + (y^2 - x^2) \partial_y^2 + x \partial_x + y \partial_y + \frac{a_1 x^2 y^2}{a_2 - a_1 x^2} (\partial_x^2 + \partial_y^2), \end{aligned}$$

which satisfy

$$\begin{aligned} [L_2, F_1] &= a_1 L_b + L_2^2, \quad [L_2, F_2] = 2F_1, \quad [F_1, F_2] = L_2 F_2 + F_2 L_2 + 2a_2 L_2 L_b - \frac{1}{2} L_2, \\ F_1^2 &= a_1 F_2 L_b + a_2 L_b L_2^2 + \frac{1}{3} (L_2^2 F_2 + L_2 F_2 L_2 + F_2 L_2^2) + \frac{a_1}{6} L_b + \frac{11}{12} L_2^2. \end{aligned}$$

We see that the *leading order terms* in each equation are identical to those of the classical case, but that some *lower order* terms were added in the last two formulae.

5.2.3 The Quantum Darboux-Koenigs Case D_3

We now take the formulae from Section 3.2, with Killing vector L_3 , giving:

$$\begin{aligned} L_b &= \frac{e^{-v}}{c_1 e^v - 2\alpha} (\partial_u^2 + \partial_v^2), \\ F_1 &= \frac{1}{2} (L_1 L_3 + L_3 L_1) + \alpha \cos u L_b = \frac{1}{2} e^{-v} \left(\cos u (2\partial_u^2 + \partial_v) + \sin u (2\partial_u \partial_v - \partial_u) + \frac{2\alpha \cos u}{c_1 e^v - 2\alpha} (\partial_u^2 + \partial_v^2) \right), \\ F_2 &= \frac{1}{2} (L_2 L_3 + L_3 L_2) - \alpha \sin u L_b = \frac{1}{2} e^{-v} \left(\cos u (2\partial_u \partial_v - \partial_u) - \sin u (2\partial_u^2 + \partial_v) - \frac{2\alpha \sin u}{c_1 e^v - 2\alpha} (\partial_u^2 + \partial_v^2) \right), \end{aligned}$$

which satisfy

$$[L_3, F_1] = F_2, \quad [L_2, F_2] = -F_1, \quad [F_1, F_2] = -c_1 L_3 L_b, \quad F_1^2 + F_2^2 = c_1 L_3^2 L_b + \alpha^2 L_b^2 + \frac{c_1}{4} L_b.$$

Again, we see that the *leading order terms* in each equation are identical to those of the classical case, but a *lower order* term is added to the last formula.

5.2.4 The Quantum Darboux-Koenigs Case D_4

We now take the formulae from Section 3.3, with Killing vector L_4 , giving:

$$\begin{aligned} L_b &= \frac{\sin^2 u}{c_1 - 2\alpha \cos u} (\partial_u^2 + \partial_v^2), \\ F_1 &= \frac{1}{2}(L_2 L_4 + L_4 L_2) - \alpha e^{-v} L_b = \frac{1}{2} e^{-v} \left(\cos u (2\partial_v^2 - \partial_v) - \sin u (2\partial_u \partial_v - \partial_u) - \frac{2\alpha \sin^2 u}{c_1 - 2\alpha \cos u} (\partial_u^2 + \partial_v^2) \right), \\ F_2 &= \frac{1}{2}(L_4 L_6 + L_6 L_4) - \alpha e^v L_b = \frac{1}{2} e^v \left(\cos u (2\partial_v^2 + \partial_v) + \sin u (2\partial_u \partial_v + \partial_u) - \frac{2\alpha \sin^2 u}{c_1 - 2\alpha \cos u} (\partial_u^2 + \partial_v^2) \right), \end{aligned}$$

which satisfy

$$\begin{aligned} [L_4, F_1] &= -F_1, \quad [L_4, F_2] = F_2, \quad [F_1, F_2] = -2c_1 L_4 L_b + 4L_4^3 + \frac{1}{2} L_4, \\ F_1 F_2 + F_2 F_1 &= 2\alpha L_b^2 + 2L_4^4 - 2c_1 L_4^2 L_b - \frac{c_1}{2} L_b + \frac{5}{2} L_4^2. \end{aligned}$$

Again, we see that the *leading order terms* in each equation are identical to those of the classical case, but *lower order terms* are added to the last two formulae.

5.3 The Quantum Case with Third Order Integrals

Here we consider the quantum case of the algebra discussed in Section 4. The strategy will again be to take the formulae *directly* from the classical case. However, this time we find that we are forced to a constant curvature case, so presumably it is necessary to look at a more general conformal invariant. Given that we are reduced to the constant curvature case, we can isolate a 3 dimensional subalgebra of Killing vectors from the 6 dimensional conformal algebra and express F_1 and F_2 in terms of these.

5.3.1 The Operator Algebra

We start with L_b and the operator version of (15):

$$L_b = \varphi(x)(\partial_x^2 + \partial_y^2), \tag{25a}$$

$$\begin{aligned} F_1 &= \frac{c_0}{2}(L_2(yL_b) + yL_b L_2) + \frac{c_1}{2}(L_4 L_2^2 + L_2^2 L_4) \\ &\quad + c_2 L_1 L_2^2 + \frac{1}{2}(b(x)L_b L_1 + L_1(b(x)L_b)), \end{aligned} \tag{25b}$$

where the first and last symmetric sum is required as a consequence of $[L_2, yL_b] \neq 0$ and $[L_1, b(x)L_b] \neq 0$. We then have $[L_2, F_1] = c_0 L_b L_2 + c_1 L_2^3$, and the condition $[L_b, F_1] = 0$ implies

$$\varphi = b^2, \quad b_x = \frac{b(c_1 + c_0 b^2)}{c_2 + c_1 x + b^3} \quad \text{and} \quad b_{xxx} = 0.$$

The two conditions on $b(x)$ imply

$$3b^2(c_1 b - c_0(c_2 + c_1 x))(c_1 + c_0 b^2)(4c_0 b^5 + 8c_1 b^3 - 5c_0(c_2 + c_1 x)b^2 - c_1(c_2 + c_1 x)) = 0.$$

Since the scalar curvature of the corresponding metric is given by $R = 2(bb_{xx} - b_x^2)$, we must choose the quintic factor to be zero, in order to avoid *constant curvature*. However, since $b_{xxx} = 0$ gives $b(x) = \alpha_0 + \alpha_1 x + \alpha_2 x^2$, this quintic quickly leads to $\alpha_2 = \alpha_1 = \alpha_0 = 0$. We are thus led to choosing

$$b(x) = \frac{c_0}{c_1}(c_2 + c_1 x) \quad \Rightarrow \quad R = -2c_0^2. \tag{26}$$

The quantum version of formula (17) for F_2 is

$$F_2 = \frac{c_0}{4}(L_2(yL_b) + yL_bL_2) + \frac{c_1}{4}(L_6L_2^2 + L_2^2L_6) + \frac{c_2}{2}(L_3L_2^2 + L_2^2L_3) + \frac{1}{2}(bL_bL_3 + L_3(bL_b)) + b_1(x)L_bL_2. \quad (27)$$

We have $[L_2, F_2] = F_1$ and requiring $[L_b, F_2] = 0$ implies

$$b_{1x} = xb_x = c_0x \quad \Rightarrow \quad b_1(x) = \frac{1}{2}c_0x^2.$$

The final bracket is

$$[F_1, F_2] = c_0L_bF_2 + \frac{3}{2}c_1(L_2^2F_2 + F_2L_2^2) + d_1L_b^2L_2 + d_2L_bL_2^3 + d_3L_2^5 - \frac{3}{2}c_0c_1L_bL_2 - \frac{7}{2}c_1^2L_2^3, \quad (28)$$

where

$$d_1 = \frac{2c_1^3 - c_0^3c_2^2}{c_0c_1^2}, \quad d_2 = \frac{2c_1^3 - 4c_0^3c_2^2}{c_0^2c_1}, \quad d_3 = -3c_2^2.$$

Comparing this with (18), we see that d_1 and d_2 are now determined and there are some lower order ‘‘corrections’’. The constraint (21) now takes the form

$$F_1^2 - c_0(L_2F_2 + F_2L_2)L_b - c_1(L_2^3F_2 + F_2L_2^3) = L_b^3 - c_2^2L_2^6 + d_1L_b^2L_2^2 + \frac{1}{2}d_2L_bL_2^4 - 3c_0c_1L_bL_2^2 - 4c_1^2L_2^4,$$

where d_1, d_2 have the values given above.

5.3.2 Additional Killing Vectors

Given that the metric has constant curvature, it must have a 3 dimensional isometry group. We can determine the Killing vectors quite simply by considering

$$[K, L_b] = 0, \quad \text{with} \quad K = \sum_{i=1}^6 \alpha_i L_i,$$

where L_i are just the conformal vectors (24a). We find

$$\begin{aligned} K_1 &= L_2 = \partial_y, & K_2 &= c_2L_1 + c_1L_4 = (c_2 + c_1x)\partial_x + c_1y\partial_y, \\ K_3 &= 2c_2L_3 + c_1L_6 = 2(c_2 + c_1x)y\partial_x + (c_1(y^2 - x^2) - 2c_2x)\partial_y \end{aligned} \quad (29)$$

which satisfy

$$[K_1, K_2] = c_1K_1, \quad [K_1, K_3] = 2K_2, \quad [K_2, K_3] = c_1K_3 - 2c_2^2K_1,$$

with Casimir

$$\mathcal{C}_K = c_1(K_1K_3 + K_3K_1) - 2K_2^2 - 2c_2^2K_1^2 = -\frac{c_1^2}{c_0^2}L_b.$$

In terms of these we have

$$\begin{aligned} F_1 &= \frac{c_0^3}{c_1^3}K_2^3 - \frac{c_0^3}{6c_1^2}(K_1K_2K_3 + K_2K_3K_1 + K_3K_1K_2 + K_2K_1K_3 + K_1K_3K_2 + K_3K_2K_1) \\ &\quad + \frac{c_1^3 + c_0^3c_2^2}{2c_1^3}(K_1^2K_2 + K_2K_1^2) + \frac{c_0^3}{3c_1}K_2, \\ F_2 &= \frac{c_0^3}{4c_1^3}(K_2^2K_3 + K_3K_2^2) - \frac{c_0^3}{4c_1^2}(K_1K_3^2 + K_3^2K_1) + \frac{c_1^3 + c_0^3c_2^2}{4c_1^3}(K_1^2K_3 + K_3K_1^2) + \frac{c_0^3}{4c_1}K_3 - \frac{c_0^3c_2^2}{2c_1^2}K_1. \end{aligned}$$

Remark 5.1 *It is not strictly necessary to symmetrise these formulae, but not doing so does add more lower order terms.*

6 Conclusions

For metrics of constant curvature (including flat), all (polynomial in momenta) first integrals of the geodesic equations can be written as polynomial functions of the Noether constants of the isometry algebra.

In this paper, we have considered a similar procedure for building integrals in the *conformally flat* case from polynomial functions of *conformal symmetries*. We considered the standard 6 dimensional algebra of conformal Killing vectors, together with additional quadratic elements of the form $\sigma(x, y)H$. This approach allowed us to “derive”, in a simple way, the 4 known Darboux-Koenigs metrics (with 1 linear and 2 quadratic integrals), as well as one of the cases derived in the classification of [10] (with 1 linear and 2 cubic integrals).

This construction gives us a better understanding of the origin of these quadratic and cubic integrals in the classical case. However, it also gives us a mechanism for building quantum analogues, with very little additional calculation.

In this paper, we only considered the 2 dimensional case, since our purpose was to understand the known results in this case. However, the method is easily extended to higher dimensional metrics, although the calculations are considerably more complicated. In 3 dimensions the constant curvature metrics have 6 dimensional isometry algebras and there exist conformally flat metrics with 3 dimensional isometry algebras, out of which it is possible to construct 3 quadratic invariants. Such an example was given in [5], with a closed Poisson algebra with only 4 independent functions (including the Hamiltonian). No further (independent) integrals can be constructed from the isometry algebra, but perhaps the methods of this paper could give us the fifth integral needed for *maximal super-integrability*.

This paper was only concerned with integrals of the geodesic equations (ie building *Killing tensors*), and not with finding potential functions, consistent with integrability. This problem was completely solved in the case of Darboux-Koenigs metrics in the papers [8, 7].

References

- [1] B.A. Dubrovin, A.T. Fomenko, and S.P. Novikov. *Modern Geometry – Methods and Applications (3 volumes)*. Springer-Verlag, NY, 1984.
- [2] C. Duval and G. Valent. Quantum integrability of quadratic Killing tensors. *J.Math.Phys.*, 46:053516 (22 pages), 2005.
- [3] A.P. Fordy. Quantum super-integrable systems as exactly solvable models. *SIGMA*, 3:025, 10 pages, 2007.
- [4] A.P. Fordy. Classical and quantum super-integrability: From Lissajous figures to exact solvability. *Atom. Nuclei*, 81:832 – 42, 2018. preprint arXiv:1711.10583 [nlin.SI].
- [5] A.P. Fordy and Q. Huang. Poisson algebras and 3D superintegrable Hamiltonian systems. *SIGMA*, 14:022, 37 pages, 2018.
- [6] R. Gilmore. *Lie Groups, Lie Algebras and Some of Their Applications*. Wiley, New York, 1974.
- [7] E.G. Kalnins, J.M. Kress, W. Miller, Jr, and P. Winternitz. Superintegrable systems in Darboux spaces. *J.Math.Phys.*, 44:5811, 2003.
- [8] E.G. Kalnins, J.M. Kress, and P. Winternitz. Superintegrability in a two-dimensional space of nonconstant curvature. *J.Math.Phys.*, 43:970, 2002.
- [9] G. Koenigs. Sur les géodésiques a intégrales quadratiques. In G. Darboux, editor, *Leçons sur la théorie générale des surfaces, Vol. 4*, pages 368–404. Chelsea, New York, 1972.
- [10] V.S. Matveev and V.V. Shevchishin. Two-dimensional superintegrable metrics with one linear and one cubic integral. *J.Geom.Phys.*, 61:1353–77, 2011.

- [11] W. Miller Jr, S. Post, and P. Winternitz. Classical and quantum superintegrability with applications. *J.Phys.A*, 46:423001 (97 pages), 2013.