

This is a repository copy of *Higgs bundles over cell complexes and representations of finitely presented groups*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/149809/>

Version: Accepted Version

---

**Article:**

Daskalopoulos, Georgios, Mese, Chikako and Wilkin, Graeme Peter Desmond (2018) Higgs bundles over cell complexes and representations of finitely presented groups. Pacific Journal of Mathematics. pp. 31-55. ISSN 0030-8730

<https://doi.org/10.2140/pjm.2018.296.31>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# HIGGS BUNDLES OVER CELL COMPLEXES AND REPRESENTATIONS OF FINITELY PRESENTED GROUPS

GEORGIOS DASKALOPOULOS, CHIKAKO MESE, AND GRAEME WILKIN

ABSTRACT. The purpose of this paper is to extend the Donaldson-Corlette theorem to the case of vector bundles over cell complexes. We define the notion of a vector bundle and a Higgs bundle over a complex, and describe the associated Betti, de Rham and Higgs moduli spaces. The main theorem is that the  $SL(r, \mathbb{C})$  character variety of a finitely presented group  $\Gamma$  is homeomorphic to the moduli space of rank  $r$  Higgs bundles over an admissible complex  $X$  with  $\pi_1(X) = \Gamma$ . A key role is played by the theory of harmonic maps defined on singular domains.

## CONTENTS

1. Introduction	1
2. Vector bundles over complexes	3
2.1. Basic Definitions of smooth bundles	3
2.2. Relationship to Higgs bundles	8
2.3. The balancing condition	9
3. Harmonic maps and Higgs bundles	10
3.1. Estimates of harmonic maps	10
3.2. Weighted Sobolev spaces	14
4. Equivalence of moduli spaces	16
4.1. Higgs Moduli Space	16
4.2. The inverse map	18
References	20

## 1. INTRODUCTION

Higgs bundles were first introduced by Hitchin in [21] as a PDE on a vector bundle over a Riemann surface obtained by the dimensional reduction of the anti-self dual equations on  $\mathbb{R}^4$ . Since then, the field has seen a remarkable explosion in different directions most notably the work of Simpson on variations of Hodge structures and applications to Kähler groups (cf. [32], [34]). The work of Donaldson and Corlette ([16], [8]) provided links with the theory of flat bundles and character varieties of groups. Higgs bundles have been generalized over noncompact manifolds (cf. [10], [33], [24], [25]) and singular curves (cf. [5]). The goal of this paper is to push this even further by considering Higgs bundles over more general singular spaces; namely, finite simplicial complexes.

---

2000 *Mathematics Subject Classification.* Primary: 58E20 ; Secondary: 53C07, 58D27 .

G.D. supported by grant number NSF DMS 1608764.

C.M. supported by grant number NSF DMS 1709475.

G.W. supported by grant number R-146-000-200-112 from the National University of Singapore.

As pointed out by Hitchin, Donaldson and Corlette, a key role in the relation between character varieties and Higgs bundles is played by the theory of harmonic maps. Harmonic maps had been used before in the study of representations of Kähler manifold groups starting with the work of Siu [35] (also see [6]) and has seen some remarkable applications in providing new proofs of the celebrated Margulis super rigidity theorem (cf. [23] and all the references therein) and the only existing proof of the rank one superrigidity theorem due to Corlette and Gromov-Schoen (cf. [9] and [20]). But these directions involved showing that the representations are rigid, in contrast with Hitchin's point of view which is to study the moduli space of such representations.

In all the above references, one studies representations of fundamental groups of smooth manifolds rather than arbitrary finitely presented groups. In other words, the domain space of the harmonic map is smooth. Chen and Eells-Fuglede (cf. [7], [17]) developed the theory of harmonic maps from a certain class of singular domains including admissible simplicial complexes. By admissible they mean complexes that are dimensionally homogeneous and locally chainable in order to avoid certain analytic pathologies (see next section for precise definitions). Since any finitely presented group is the fundamental group of an admissible complex, there is no real restriction in considering admissible complexes. The key property of harmonic maps shown in the above references is that they are Hölder continuous but in general they fail to be Lipschitz. In fact, the work of the first two authors shows that Lipschitz harmonic maps often imply that the representations are rigid (cf. [12] and [13]).

The starting point of this paper is a finitely presented group  $\Gamma$  and a two-dimensional admissible complex without boundary  $X$  with fundamental group  $\pi_1(X) \simeq \Gamma$ . We also fix a piecewise smooth vector bundle  $E$  over  $X$  that admits a flat  $SL(r, \mathbb{C})$ -structure. Such bundles are parametrized topologically by the (finitely many) connected components of the  $SL(r, \mathbb{C})$ -character variety of  $\pi_1(X)$ . One can write down Hitchin's equations

$$(1.1) \quad F_A + \psi \wedge \psi = 0$$

$$(1.2) \quad d_A \psi = 0$$

for a sufficiently regular unitary connection  $A$  and Higgs field  $\psi$ . Again, as in the smooth case, the  $SL(r, \mathbb{C})$  connection  $d_A + \psi$  is flat and one can ask what the precise condition is so that the pair  $(d_A, \psi)$  corresponds to a representation  $\rho : \pi_1(X) \rightarrow SL(r, \mathbb{C})$ .

Given a representation  $\rho$  as above, we can associate as in the smooth case a  $\rho$ -equivariant harmonic map from the universal cover  $\tilde{X}$  to the symmetric space  $SL(r, \mathbb{C})/SU(r)$ . The first two authors in [12] studied harmonic maps from simplicial complexes to smooth manifolds and discovered the following crucial properties:

- (1) The harmonic map is smooth away from the codimension 2 skeleton of  $\tilde{X}$ .
- (2) The harmonic map satisfies a balancing condition at the codimension 1 skeleton of  $\tilde{X}$  in the sense that the sum of the normal derivatives vanishes identically.
- (3) The harmonic map blows up in a controlled way at the codimension 2 skeleton of  $\tilde{X}$ .

All the above properties are described precisely in Theorem 3.3. This allows us to prove that the derivative of the harmonic map belongs in an appropriate weighted Sobolev space  $L^2_{1,\delta}$  (cf. Proposition 4.5). The definition of weighted Sobolev spaces is given in Section 3.2. Finally, the main theorem describing the correspondence between equivalence

classes of balanced Higgs pairs of class  $L_{1,\delta}^2$  and representations is given in Section 4 (cf. Theorem 4.3).

We would like to end this introduction with a brief discussion of some motivation and future applications of this paper that we will explore elsewhere (cf. for example [14]). Note that, with the exception of [5], the theory of Higgs bundles on singular varieties is not very well understood. For example, one of the important questions about fundamental groups of singular projective varieties is whether fundamental groups of normal varieties behave more like the ones of smooth manifolds, or in the other extreme, there are very few restrictions on them (cf. [1] and [26]). This question will be addressed in detail in [14], and the connection with the results of this paper is as follows: By [17] Example 8.9, an  $n$ -dimensional normal projective variety  $X$  admits a bilipschitz triangulation with its singular set as a sub-complex of dimension at most  $n - 2$ . Furthermore,  $X$  is admissible in the sense of Definition 2.2. Thus, studying harmonic maps on  $X$ , or more generally constructing moduli spaces of bundles on  $X$  could imply restrictions on fundamental groups as in [6] and [34]. This is carried out in detail in [14], but the key analytic estimates are Theorem 3.3 and Proposition 3.5 of this paper. These results can be utilized to describe the behavior of the pluriharmonic map, or equivalently the Higgs bundle near the singularities of the variety. Notice that pluriharmonicity reduces the analytic estimates needed to 2-dimensions, hence the motivation of studying only 2-dimensions in this paper.

## 2. VECTOR BUNDLES OVER COMPLEXES

### 2.1. Basic Definitions of smooth bundles.

**Definition 2.1.** (cf. [27]) Let  $\mathbf{E}^N$  be an  $N$ -dimensional affine space. A *cell* of dimension  $i$  is a non-empty, open, convex, bounded subset in some  $\mathbf{E}^i \subset \mathbf{E}^N$ . We will use the notation  $\sigma^i$  to denote a cell of dimension  $i$  and call  $\mathbf{E}^i$  the *extended plane* defined by  $\sigma^i$ . A *locally finite convex cell complex* or simply a *complex*  $X$  in  $\mathbf{E}^N$  is a locally finite collection  $\mathcal{F} = \{\sigma\}$  of disjoint cells in  $\mathbf{E}^N$  such that for any  $\sigma \in \mathcal{F}$  its closure  $\bar{\sigma}$  is a union of cells in  $\mathcal{F}$ . The *dimension of a complex*  $X$  is the maximum dimension of a cell in  $X$ .

For example, a simplicial complex is a cell complex whose cells are all simplices.

**Definition 2.2.** A connected complex  $X$  of dimension  $n$  is said to be *admissible* (cf. [7] and [17]) if the following two conditions hold:

- (i)  $X$  is dimensionally homogeneous, i.e., every cell is contained in a closure of at least one  $n$ -cell, and
- (ii)  $X$  is locally  $(n - 1)$ -chainable, i.e., given any  $(n - 2)$ -cell  $v$ , every two  $n$ -cells  $\sigma$  and  $\sigma'$  incident to  $v$  can be joined by a sequence  $\sigma = \sigma_0, \sigma_1, \dots, \sigma_k = \sigma'$  where  $\sigma_i$  and  $\sigma_{i+1}$  are two adjacent  $n$ -cells incident to  $v$  for  $i = 0, 1, \dots, k - 1$ .

The boundary  $\partial X$  of  $X$  is the union of the closures of the  $(n - 1)$ -cells contained in the closure of exactly one  $n$ -cell. Using a regular barycentric subdivision we obtain that given any locally finite complex there is a locally finite simplicial complex such that any cell is a union of simplices.

**Definition 2.3.** Let  $U$  be a subset of a complex  $X$ . A function  $f : U \rightarrow \mathbb{R}$  is called *smooth* if for any  $n$ -cell  $\sigma$  of  $X$ , the restriction  $f|_{\sigma \cap U}$  can be extended to a smooth function on  $\mathbf{E}^i \cap U$

in the extended plane defined by  $\sigma$ . A map  $f : U \rightarrow Z \subset \mathbf{E}^M$  into a complex  $Z$  is called *smooth* if with respect to some affine coordinate system on  $\mathbf{E}^M$ ,  $f = (f^1, \dots, f^M)$  where  $f^j$  is smooth for every  $j = 1, \dots, M$ .

**Definition 2.4.** A Riemannian metric  $g_\sigma$  on a cell  $\sigma$  is the restriction to  $\sigma$  of a smooth Riemannian metric on its extended plane. A Riemannian metric  $g$  on  $X$  is a smooth Riemannian metric  $g_\sigma$  on each  $n$ -cell  $\sigma$  of  $X$  satisfying the additional property that if  $\tau$  is a face of  $\sigma$ , then  $g_\sigma|_\tau = g_\tau$ , where  $g_\sigma|_\tau$  denotes the restriction of the extension of  $g_\sigma$  to the extended plane of  $\sigma$ . In particular, the expressions of  $g_\sigma$  with respect to some affine coordinates in the extended plane are smooth functions in the sense of Definition 2.3.

**Definition 2.5.** A *smooth complex vector bundle of rank  $r$*  over a complex  $X$  is a topological space  $E$  and a continuous, surjective map  $\pi : E \rightarrow X$  such that:

- (1) for each  $x \in X$  the fibre  $\pi^{-1}(x)$  has the structure of a complex vector space, and
- (2) there exists an open cover  $\{U_\alpha\}_{\alpha \in I}$  of  $X$  such that for each  $\alpha \in I$  there exists a homeomorphism  $\varphi_\alpha : \pi^{-1}(U_\alpha) \rightarrow U_\alpha \times \mathbb{C}^r$  such that
  - (i)  $\varphi_\alpha$  restricts to a linear isomorphism  $\pi^{-1}(x) \cong \{x\} \times \mathbb{C}^r$  for each  $x \in U_\alpha$ , and
  - (ii) If  $U_\alpha \cap U_\beta \neq \emptyset$ , then the transition function  $g_{\alpha\beta} = \varphi_\alpha \circ \varphi_\beta^{-1} : U_\alpha \cap U_\beta \times \mathbb{C}^r \rightarrow U_\alpha \cap U_\beta \times \mathbb{C}^r$  induces a smooth map  $g_{\alpha\beta} : U_\alpha \cap U_\beta \rightarrow \text{GL}(r, \mathbb{C})$ .

A *section* of  $\pi : E \rightarrow X$  is a continuous map  $s : X \rightarrow E$  satisfying  $\pi \circ s = \text{id}_X$ . The section is *smooth* if on each local trivialisation  $\pi^{-1}(U_\alpha) \cong U_\alpha \times \mathbb{C}^r$  with projection onto the second factor denoted by  $p_2 : \pi^{-1}(U_\alpha) \rightarrow \mathbb{C}^r$ , the composition of  $U_\alpha \xrightarrow{s} \pi^{-1}(U_\alpha) \xrightarrow{p_2} \mathbb{C}^r$  is a smooth map as in Definition 2.3. Let  $\Omega^0(X, E)$  denote the vector space of all smooth sections of  $\pi : E \rightarrow X$ . If  $E$  is a smooth vector bundle, then so is any associated bundle formed by taking the dual, tensor product, etc. In particular, if  $E$  is smooth then  $\text{End}(E)$  is smooth.

**Definition 2.6.** A smooth complex  $p$ -form on a cell  $\sigma$  is the restriction to  $\sigma$  of a smooth complex  $p$  form on the extended plane of the cell. A *smooth  $p$ -form*  $\omega = \{\omega_\sigma\}_{\sigma \in \mathcal{F}}$  on a complex  $X$  with values in a smooth vector bundle  $E$  is a collection of smooth  $p$ -forms  $\omega_\sigma$  with values in  $E$  for each cell  $\sigma$  of  $X$ , with the additional property that if  $\tau$  is a face of  $\sigma$ , then  $\omega_\sigma|_\tau = \omega_\tau$ . In particular, the expressions of  $\omega_\sigma$  with respect to some affine coordinates in the extended plane are smooth functions in the sense of Definition 2.3. We denote  $\Omega^p(X, E)$  as the space of all smooth  $p$ -forms with values in  $E$ . If  $E$  is the trivial line bundle, then we write  $\Omega^p(X) = \Omega^p(X, E)$  and this is the space of smooth  $p$ -forms on  $X$ . Given a smooth  $p$ -form  $\omega = \{\omega_\sigma\}_{\sigma \in \mathcal{F}} \in \Omega^p(X)$ , we define  $d\omega = \{d\omega_\sigma\}_{\sigma \in \mathcal{F}}$  and note that this is a well-defined smooth  $(p+1)$ -form. Clearly,  $d^2 = 0$  and the complex  $(\Omega^*(X), d)$  denotes the smooth deRham complex. We denote by  $H_{dR}^p(X)$  the cohomology groups associated with this complex (cf. [19, Ch. VIII]).

**Definition 2.7.** A *smooth connection* on a smooth vector bundle  $\pi : E \rightarrow X$  is a  $\mathbb{C}$ -linear map  $D : \Omega^0(X, E) \rightarrow \Omega^1(X, E)$  that satisfies the Leibniz rule

$$D(fs) = (df)s + f(Ds), \quad f \in \Omega^0(X), s \in \Omega^0(X, E).$$

We denote the space of all smooth connections by  $\mathcal{A}^{\mathbb{C}}(E)$ .

The definition of  $D$  can be extended to bundle-valued forms in the usual way. More precisely, any element in  $\sigma \in \Omega^p(X, E)$  can be written as a linear combination of elements

of the form  $\sigma = s\omega$  with  $\omega \in \Omega^p(X)$  and  $s \in \Omega^0(X, E)$ , and define

$$(2.1) \quad D\sigma = s(d\omega) + (Ds) \wedge \omega.$$

*Remark 2.8.* Implicit in the definition of  $\Omega^1(X, E)$  is that 1-forms with values in  $E$  must agree on the interfaces between the cells in the complex  $X$ . Therefore, the definition above implies that a connection must map sections that agree on the interfaces to bundle-valued 1-forms that agree on the interfaces.

As for the case of a smooth vector bundle over a smooth manifold, with respect to a trivialization  $\varphi_\alpha : \pi^{-1}(U_\alpha) \rightarrow U_\alpha \times \mathbb{C}^r$ ,  $D = d + A_\alpha$  where  $(A_\alpha)_{ij}$  is a complex-valued smooth 1-form.  $A_\alpha$  is called the *connection form* of  $D$  with respect to the trivialization  $\varphi_\alpha$ . In different trivialization  $\varphi_\beta$  and with  $g_{\alpha\beta} = \varphi_\alpha \circ \varphi_\beta^{-1}$  we have,

$$(2.2) \quad A_\beta = g_{\alpha\beta}^{-1} d g_{\alpha\beta} + g_{\alpha\beta}^{-1} A_\alpha g_{\alpha\beta}$$

**Definition 2.9.** The *curvature* of a smooth connection  $D$  is the matrix valued 2-form  $F_D$  defined by

$$D^2 s = F_D s, \quad \text{for all } s \in \Omega^0(X, E).$$

Locally, we have  $(F_D)_\alpha = dA_\alpha + A_\alpha \wedge A_\alpha$ , where  $A_\alpha$  is the connection form of  $D$ . Furthermore,

$$(2.3) \quad (F_D)_\beta = g_{\alpha\beta}^{-1} (F_D)_\alpha g_{\alpha\beta},$$

and so the curvature form  $F_D$  is an element of  $\Omega^2(X, \text{End}(E))$ .

**Definition 2.10.** The *complex gauge group* is the group  $\mathcal{G}^{\mathbb{C}}(E)$  of all smooth automorphisms of  $E$ . If  $D$  is a smooth connection on  $E$  and  $g \in \mathcal{G}^{\mathbb{C}}(E)$ , then we define  $g(D) = g^{-1} \circ D \circ g$ . In local coordinates, the action of  $\mathcal{G}^{\mathbb{C}}(E)$  on  $\mathcal{A}^{\mathbb{C}}(E)$  is

$$(2.4) \quad g(d + A_\alpha) = d + g^{-1} dg + g^{-1} A_\alpha g.$$

**Definition 2.11.** A *smooth Hermitian metric*  $h = (h_\sigma)$  on a rank  $r$  complex vector bundle  $\pi : E \rightarrow X$  is a smooth section  $h$  of  $\text{End}(E)$  such that for each cell  $\sigma$  its restriction  $h_\sigma$  is a Hermitian metric and that if  $\tau$  is a face of  $\sigma$ , then  $h_\sigma|_\tau = h_\tau$ . A Hermitian metric in a trivialization  $\varphi_\alpha : \pi^{-1}(U_\alpha) \rightarrow U_\alpha \times \mathbb{C}^r$  is given locally by a smooth map  $\tilde{h}_\alpha$  from  $U_\alpha$  into the positive definite matrices in  $\text{GL}(r, \mathbb{C})$ , and the induced inner product on the fibres of  $E$  is

$$\langle s_1(x), s_2(x) \rangle = \overline{\varphi_\alpha(s_1(x))}^T \tilde{h}_\alpha(x) \varphi_\alpha(s_2(x)) \in \mathbb{C}.$$

**Definition 2.12.** A connection  $D$  on a vector bundle  $E$  with a Hermitian metric  $h$  is a *unitary connection* if the following equation is satisfied.

$$d \langle s_1, s_2 \rangle = \langle Ds_1, s_2 \rangle + \langle s_1, Ds_2 \rangle,$$

where  $\langle \cdot, \cdot \rangle$  is the pointwise inner product on the fibres of  $E$  induced by the metric  $h$ . The space of smooth unitary connections on  $E$  is denoted  $\mathcal{A}(E, h)$ . If  $D \in \mathcal{A}(E, h)$ , then the curvature  $F_D$  is a section of  $\Omega^2(\text{ad}(E))$ . In other words, with respect to a unitary frame field the curvature satisfies  $F_D^* = -F_D$ .

**Definition 2.13.** The *unitary gauge group*  $\mathcal{G}(E)$  is the subgroup of  $\mathcal{G}^{\mathbb{C}}(E)$  that preserves the Hermitian metric  $h$  on each fibre of  $E$ . The action on  $\mathcal{G}(E)$  on  $\mathcal{A}^{\mathbb{C}}(E)$  preserves the space  $\mathcal{A}(E, h)$ .

**Definition 2.14.** A connection  $D$  on a vector bundle  $E$  is *flat* if  $F_D = 0$ . Given a flat connection, we can define the twisted deRham complex  $(\Omega^*(X, E), D)$ . The cohomology groups will be denoted by  $H^p(X, E)$ .

**Definition 2.15.** A *flat structure* on a vector bundle  $\pi : E \rightarrow X$  is given by an open cover  $\{U_\alpha\}_{\alpha \in I}$  and trivialisations  $\{\varphi_\alpha\}_{\alpha \in I}$  for which the transition functions  $g_{\alpha\beta} = \varphi_\alpha \circ \varphi_\beta^{-1}$  are constant. A vector bundle with a flat structure is also called a *flat bundle*.

*Remark 2.16.* Equation (2.2) shows that the connection  $D = d$  (with zero connection form) is globally defined on a flat bundle. Thus a flat bundle clearly admits a connection of curvature zero. The converse is also true.

**Theorem 2.17.** *Let  $X$  be  $n$ -complex,  $U$  an open subset of  $X$  and  $E$  a smooth vector bundle with a smooth flat connection on  $U$ . Then  $E$  admits a flat structure.*

*Proof.* Given a flat connection  $D$  on  $E$ , fix a cell  $\sigma$ , a point  $x_0 \in \bar{\sigma} \cap U$  and consider a contractible neighbourhood  $V_\sigma$  of  $x_0$  in the extended plane of  $\sigma$ . Choose a local frame  $s_\sigma^0$  of  $E$  on  $V_\sigma$  and let  $A^\sigma$  be the corresponding connection form. We are assuming that the local frames  $s_\sigma^0$  patch together to define a piecewise smooth frame  $s_0$  in a neighbourhood of  $x_0$  in  $X$ . We are going to choose a different trivialisation  $s_\sigma$  for which the connection can be written as  $D = d$ . This can be done by solving the equation

$$(2.5) \quad \begin{aligned} g_\sigma^{-1} A^\sigma g_\sigma + g_\sigma^{-1} dg_\sigma &= 0 \\ \Leftrightarrow dg_\sigma &= -A^\sigma g_\sigma \end{aligned}$$

locally for a gauge transformation  $g_\sigma$ . By the result in the smooth case (this is an application of the Frobenius theorem) a solution  $g_\sigma$  exists and by multiplying by a constant matrix we may assume without loss of generality that  $g_\sigma(x_0) = id$ . This makes the solution unique and thus if a cell  $\tau$  is a face of a cell  $\sigma$  then, since  $A^\sigma|_\tau = A^\tau$ , it must be  $g_\sigma|_\tau = g_\tau$ . It follows that the new frames  $s_\sigma = g_\sigma \circ s_\sigma^0$  patch together to define a piecewise smooth frame  $s$  in a neighbourhood of  $x_0$  in  $X$ . The flat structure is now defined by the local frames  $\{s\}$ .  $\square$

**Definition 2.18.** A section  $s \in \Omega^0(X, E)$  is *parallel with respect to  $D$*  if  $Ds = 0$ . Given a smooth curve  $c : [a, b] \rightarrow X$ , a section  $s$  is *parallel along  $c$  with respect to  $D$*  if  $D_{c'(t)}s = 0$ . Given a curve  $c : [a, b] \rightarrow X$  and  $s_a \in \pi^{-1}(c(a))$  the *parallel transport of  $s$  along  $c$  with respect to  $D$*  is the section  $s : \pi^{-1}(c([a, b])) \rightarrow E$  which is given locally by the solution to the equation

$$\frac{ds(c(t))}{dt} + A_{c(t)}(c'(t))s(c(t)) = 0.$$

**Lemma 2.19.** *Let  $c_1, c_2 : [a, b] \rightarrow X$  be two closed smooth curves in  $X$  which are homotopy equivalent, and which satisfy  $x_0 = c_1(a) = c_1(b) = c_2(a) = c_2(b)$ . Let  $D$  be a smooth flat connection on a rank  $r$  bundle  $\pi : E \rightarrow X$ , and let  $s_1$  and  $s_2$  be the parallel transport with respect to  $D$  along  $c_1$  and  $c_2$  respectively, with initial condition  $s_0 \in \pi^{-1}(x_0)$ . If  $F_D = 0$  then  $s_1(c_1(b)) = s_2(c_2(b))$ .*

*Proof.* As usual note that it suffices to show that the holonomy is trivial around a homotopically trivial loop. If there is a homotopy equivalence between two loops that is constant except on a single cell, then standard theorems for smooth manifolds show that the holonomy around the two loops is the same. Given a homotopically trivial loop  $\gamma$ ,

there is a sequence of homotopy equivalences  $\gamma \sim \gamma_1, \gamma_1 \sim \gamma_2, \dots, \gamma_N \sim \text{id}$  between  $\gamma$  and the trivial loop (denoted  $\text{id}$ ), such that each homotopy equivalence is constant except on a single  $n$ -cell. For example, one can do this by identifying the fundamental group with the edge group of a simplicial complex (cf. [4], Section 6.4). Therefore, the holonomy of  $\gamma$  is the same as the holonomy of each  $\gamma_n$  along this sequence of homotopy equivalences, and so the holonomy of  $\gamma$  is trivial.  $\square$

**Definition 2.20.** A flat connection  $D$  on a rank  $r$  vector bundle  $\pi : E \rightarrow X$  defines a representation  $\rho : \pi_1(X) \rightarrow \text{GL}(r, \mathbb{C})$  called the *holonomy representation of  $D$* . A flat connection is called *irreducible* if its holonomy representation is irreducible. The space of irreducible, flat smooth connections is denoted  $\mathcal{A}^{\mathbb{C}, \text{irr}}(E)$ .

**Lemma 2.21.** A representation  $\rho : \pi_1(X) \rightarrow \text{GL}(r, \mathbb{C})$  defines a flat connection on a bundle  $\pi : E_\rho \rightarrow X$  with holonomy representation  $\rho$ . Moreover, the flat connection on  $E_\rho$  depends continuously on the representation  $\rho$ .

*Proof.* In the standard way, from a representation  $\rho : \pi_1(X) \rightarrow \text{GL}(r, \mathbb{C})$  we construct a flat vector bundle  $E_\rho \rightarrow X$ , with total space

$$(2.6) \quad E_\rho = \tilde{X} \times_\rho \mathbb{C}^r,$$

where  $\tilde{X}$  is the universal cover of  $X$ , and the equivalence is by deck transformations on the left factor  $\tilde{X}$ , and via the representation  $\rho$  on the right factor  $\mathbb{C}^r$ . On each trivialisation we have the trivial connection  $d$ , and since the transition functions of  $E$  are constant, then this connection is globally defined. Since the deck transformations depend continuously on the representation  $\rho$  then the flat connection on  $E_\rho$  depends continuously on  $\rho$ .  $\square$

**Corollary 2.22.** A flat connection on a vector bundle over a simply connected complex  $X$  is complex gauge equivalent to the trivial connection  $d$  on the trivial vector bundle.

**Definition 2.23.** The  $\text{SL}(r, \mathbb{C})$  character variety is the space of irreducible representations  $\rho : \pi_1(X) \rightarrow \text{SL}(r, \mathbb{C})$  modulo conjugation by  $\text{SL}(r, \mathbb{C})$

$$(2.7) \quad \mathcal{M}_{\text{char}} = \{\text{irreducible reps } \rho : \pi_1(X) \rightarrow \text{SL}(r, \mathbb{C})\} / \text{SL}(r, \mathbb{C})$$

The next Lemma is a trivial consequence of the path lifting property and is standard.

**Lemma 2.24.** If two characters defined by the representations  $\rho$  and  $\rho'$  belong to the same connected component of  $\mathcal{M}_{\text{char}}$  then the vector bundles  $E_\rho$  and  $E_{\rho'}$  are smoothly isomorphic.

In view of the above, let  $\mathcal{C}$  denote the set of connected components of  $\mathcal{M}_{\text{char}}$ . Then we can write

$$\mathcal{M}_{\text{char}} = \bigsqcup_{c \in \mathcal{C}} \mathcal{M}_{\text{char}}^c$$

and write  $E_c = E_\rho$  for any representative in the isomorphism class of bundles defined by  $\rho \in \mathcal{M}_{\text{char}}^c$ .

*Remark 2.25.* Since we are interested in the  $\text{SL}(r, \mathbb{C})$  character variety instead of the  $\text{GL}(r, \mathbb{C})$  character variety we need to fix determinants in our definitions of connections and gauge transformations. Henceforth we will impose the condition that all connection forms are traceless and all gauge transformations have determinant one. For the sake of notational simplicity we will keep the same notation as before for the various spaces of  $\text{SL}(r, \mathbb{C})$  connections and gauge groups.



**Proposition 2.26.**

$$\mathcal{A}_{flat}^{\mathbb{C},irr}(E_c)/\mathcal{G}^{\mathbb{C}}(E_c) \cong \mathcal{M}_{char}^c$$

*Proof.* The holonomy map applied to an irreducible flat connection  $D$  gives an irreducible representation  $\rho : \pi_1(X, x_0) \rightarrow \mathrm{GL}(r, \mathbb{C})$ . The action of a complex gauge transformation  $g \in \mathcal{G}^{\mathbb{C}}(E_c)$  on  $D$  induces the conjugate action of an element  $\xi = g(x_0) \in \mathrm{GL}(r, \mathbb{C})$  on  $\rho$ . Therefore we have a continuous map  $\tau : \mathcal{A}_{flat}^{\mathbb{C},irr}(E_c)/\mathcal{G}^{\mathbb{C}}(E_c) \rightarrow \mathcal{M}_{char}^c$ . Note that  $\tau([D_1]) = \tau([D_2])$  implies that the flat structures associated to  $D_1$  and  $D_2$  by Theorem 2.17 are complex gauge-equivalent, and so  $D_1$  and  $D_2$  are complex gauge-equivalent. Therefore  $\tau$  is injective.

Similarly, given a representation  $\rho : \pi_1(X, x_0) \rightarrow \mathrm{GL}(r, \mathbb{C})$  we construct a flat connection  $d$  on the flat bundle  $E_\rho$  as in the proof of Lemma 2.21. If we conjugate the representation by an element  $\xi \in \mathrm{GL}(r, \mathbb{C})$ , then the flat connection associated to this new representation is related to  $E_\rho$  by a global change of co-ordinates using the action of  $\xi$  on the fibres of  $E_\rho$ . Therefore the two flat bundles are complex gauge-equivalent, and so conjugate representations give  $\mathcal{G}^{\mathbb{C}}(E_c)$ -equivalent flat connections, which gives us a continuous map  $\zeta : \mathcal{M}_{char} \rightarrow \mathcal{A}_{flat}^{\mathbb{C},irr}(E_c)/\mathcal{G}^{\mathbb{C}}(E_c)$ . Lemma 2.21 shows that  $\tau \circ \zeta = \mathrm{id}$ . Since  $\tau$  is injective then this implies that  $\zeta \circ \tau = \mathrm{id}$  and so  $\tau$  is a homeomorphism  $\mathcal{A}_{flat}^{\mathbb{C},irr}(E_c)/\mathcal{G}^{\mathbb{C}}(E_c) \cong \mathcal{M}_{char}^c$ .  $\square$

**2.2. Relationship to Higgs bundles.** Given a complex  $X$  with universal cover  $\tilde{X}$ , fix an irreducible representation  $\rho : \pi_1(X) \rightarrow \mathrm{SL}(r, \mathbb{C})$ , and let  $E = \tilde{X} \times_\rho \mathbb{C}^r \rightarrow X$  be as before. We also fix a  $\rho$ -equivariant map  $u : \tilde{X} \rightarrow \mathrm{SL}(r, \mathbb{C})/\mathrm{SU}(r)$ , locally Lipschitz away from the 0-skeleton  $X^0$  of  $X$ . We now recall the basic construction from [8] and [16].

- (1) The complexified tangent space  $T_h^{\mathbb{C}}(\mathrm{SL}(r, \mathbb{C})/\mathrm{SU}(r))$  can be identified (independent of  $h$ ) with the space of traceless matrices and this gives a trivialization of the complexified tangent bundle  $T^{\mathbb{C}}(\mathrm{SL}(r, \mathbb{C})/\mathrm{SU}(r)) \cong \mathrm{SL}(r, \mathbb{C})/\mathrm{SU}(r) \times \mathfrak{sl}(r, \mathbb{C})$ .
- (2) In the trivialization given in (1) the Levi-Civita connection at a point  $h \in \mathrm{SL}(r, \mathbb{C})/\mathrm{SU}(r)$  has the form

$$\nabla_X Y = dY(X) - \frac{1}{2} (dh(X)h^{-1}Y + Yh^{-1}dh(X)),$$

where we use the notation  $h$  to indicate left translation by  $h$ .

- (3) The identification  $h^{-1}(T_h^{\mathbb{C}}(\mathrm{SL}(r, \mathbb{C})/\mathrm{SU}(r))) \cong T_{\mathrm{id}}^{\mathbb{C}}(\mathrm{SL}(r, \mathbb{C})/\mathrm{SU}(r)) \cong \mathfrak{sl}(r, \mathbb{C})$  gives another isomorphism  $\theta : T^{\mathbb{C}}(\mathrm{SL}(r, \mathbb{C})/\mathrm{SU}(r)) \rightarrow \mathrm{SL}(r, \mathbb{C})/\mathrm{SU}(r) \times \mathfrak{sl}(r, \mathbb{C})$ . It follows immediately from (2) that in the coordinates given by  $\theta$ , the Levi-Civita connection is given by

$$\begin{aligned} \nabla_X s &= h^{-1} \nabla_X (hs) \\ &= ds(X) + \frac{1}{2} [h^{-1}dh(X), s]. \end{aligned}$$

We thus conclude that in the above coordinates

$$(2.8) \quad \nabla = d + \frac{1}{2} [h^{-1}dh, \cdot].$$

- (4) The isomorphism  $\theta$  is equivariant with respect to the  $\mathrm{PSL}(r, \mathbb{C})$  action on the complexified tangent bundle  $T^{\mathbb{C}}(\mathrm{SL}(r, \mathbb{C})/\mathrm{SU}(r))$  and the adjoint representation on  $T_{\mathrm{id}}^{\mathbb{C}}(\mathrm{SL}(r, \mathbb{C})/\mathrm{SU}(r)) \cong \mathfrak{sl}(r, \mathbb{C})$ .

- (5) Given  $u$  as above, consider the pull-backs  $\mathcal{D} = u^*d$  and  $d_A = u^*\nabla$  on the trivial bundle  $\tilde{X} \times T_{id}^{\mathbb{C}}(\mathrm{SL}(r, \mathbb{C})/\mathrm{SU}(r)) \cong \tilde{X} \times \mathfrak{sl}(r, \mathbb{C})$ . First notice, that since  $u^*d$  is trivial and  $u$  is  $\rho$ -equivariant  $\mathcal{D}$  descends to a flat connection of holonomy  $\rho$  on  $E_\rho$ . Again, by the  $\rho$ -equivariance of  $u$  and (4) the connection  $d_A$  descends to a connection on  $ad(E_\rho)$  over  $X$ . Moreover, since its connection form acts by the adjoint representation it defines an  $\mathrm{SL}(r, \mathbb{C})$  connection on the bundle  $E_\rho$  over  $X$  and (2.8) implies

$$(2.9) \quad \mathcal{D} = d_A + \psi, \quad \psi = -\frac{1}{2}u^{-1}du.$$

Since  $\mathcal{D}$  is a flat connection, then

$$(2.10) \quad F_A + \psi \wedge \psi = 0$$

$$(2.11) \quad d_A\psi = 0.$$

### 2.3. The balancing condition.

**Definition 2.27.** A smooth 1-form  $\omega = \{\omega_\sigma\}_{\sigma \in \mathcal{F}} \in \Omega^1(X)$  satisfies the *balancing condition* if for every  $(n-1)$ -cell  $\tau$ , we have

$$(2.12) \quad \sum_{\sigma > \tau} \omega_\sigma(e_\sigma) = 0$$

where  $\sigma > \tau$  implies that  $\tau$  is a face of  $\sigma$ , and  $e_\sigma$  is an inward pointing normal vector field along  $\tau$  in  $\sigma$ . The set  $\Omega_{bal}^1(X)$  is the subset of  $\Omega^1(X)$  consisting of forms satisfying the balancing condition.

**Definition 2.28.** Let  $E$  be smooth vector bundle on  $X$  of rank  $r$  and let  $p : \tilde{X} \rightarrow X$  be the universal cover. We assume that the pullback bundle  $p^*(E)$  over  $\tilde{X}$  is trivial with a fixed trivialization  $p^*(E) \cong \tilde{X} \times \mathbb{C}^r$  (if the connection is flat then this is always valid by Corollary 2.22). A connection  $D \in \mathcal{A}^{\mathbb{C}, irr}(E)$  is called *balanced* if its pullback  $p^*(D)$  to  $p^*(E)$  can be written (in the given trivialization) as  $p^*(D) = d + A$  where all the components  $A_{ij} \in \Omega_{bal}^1(\tilde{X})$ . Let  $\mathcal{A}_{bal}^{\mathbb{C}, irr}(E)$  be the space of irreducible, smooth, balanced  $\mathrm{GL}(r, \mathbb{C})$  connections, and let  $\mathcal{A}_{bal}^{irr}(E)$  denote the space of irreducible, smooth, balanced connections compatible with the Hermitian metric  $h$  on  $E$ . In what follows, if the meaning is clear then the notation for the metric is suppressed.

**Definition 2.29.** Let  $E$  be as in the previous Definition. Given  $g \in \mathcal{G}^{\mathbb{C}}(E)$ , let  $\tilde{g}$  denote the induced gauge transformation of  $p^*(E)$ . We define  $\mathcal{G}_{bal}^{\mathbb{C}}(E)$  (resp.  $\mathcal{G}_{bal}(E)$ ) to be the group of complex (resp. unitary) gauge transformations such that  $g \in \mathcal{G}_{bal}^{\mathbb{C}}(E)$  (resp.  $g \in \mathcal{G}_{bal}(E)$ ) implies that  $d\tilde{g}_{ij} \in \Omega_{bal}^1(\tilde{X})$ .

*Remark 2.30.* Note that via (2.4), the group  $\mathcal{G}_{bal}^{\mathbb{C}}(E)$  acts on the space  $\mathcal{A}_{bal}^{\mathbb{C}, irr}(E)$ , and  $\mathcal{G}_{bal}(E)$  acts on  $\mathcal{A}_{bal}^{irr}(E)$ .

*Remark 2.31.* In this paper we are interested in flat bundles. Note that Corollary 2.22 implies that the pullback of a flat bundle to the universal cover is trivial. By choosing a trivialization it thus makes sense to talk about balanced connections and gauge transformations.

### 3. HARMONIC MAPS AND HIGGS BUNDLES

In this section we describe the relationship between Higgs bundles and harmonic maps from a complex  $X$  into the space  $\mathrm{SL}(n, \mathbb{C})/\mathrm{SU}(n)$ , a generalisation of the construction of [16] and [8]. From now on  $X$  will denote an admissible 2-dimensional simplicial complex without boundary. We will further assume that  $X$  is equipped with a Riemannian metric  $g$  such that for any 2-simplex  $\sigma$ ,  $(\sigma, g_\sigma)$  is isometric to an interior of an equilateral triangle in  $\mathbf{R}^2$  and for any 1-simplex  $\tau$ ,  $(\tau, g_\tau)$  is isometric with the open unit interval in  $\mathbf{R}$ . It is not hard to extend the results of this section to general Riemannian metrics and also general 2-dimensional complexes. We endow  $\mathrm{SL}(n, \mathbb{C})/\mathrm{SU}(n)$  with a Riemannian metric of non-positive sectional curvature such that  $\mathrm{SL}(n, \mathbb{C})$  acts by isometries.

#### 3.1. Estimates of harmonic maps.

**Theorem 3.1.** *Let  $X$  be a 2-complex as before with universal cover  $\tilde{X}$  and  $\rho : \pi_1(X) \rightarrow \mathrm{SL}(n, \mathbb{C})$  be an irreducible representation. Then there exists a unique  $\rho$ -equivariant harmonic map  $u : \tilde{X} \rightarrow Y := \mathrm{SL}(n, \mathbb{C})/\mathrm{SU}(n)$ .*

*Proof.* The existence is a special case of Theorem 4.5 of [11]. Uniqueness follows from [30].  $\square$

Let  $p$  be a vertex (i.e. 0-cell) of  $X$ . Given an 1-cell  $\tau$  of  $X$ , define  $\mathcal{S}_2(\tau)$  be the set of 2-cells of  $X$  containing  $\tau$  in its closure.

**Theorem 3.2.** *If  $u : X \rightarrow Y$  is a harmonic map, then for any 1-simplex  $\tau$  and 2-simplex  $\sigma \in \mathcal{S}_2(\tau)$  we have  $u \in C^\infty(\sigma \cup \tau)$ . (In other words, the restriction of  $u$  to  $\sigma$  is  $C^\infty$  up to  $\tau$  in the extended plane of  $\sigma$ ). Moreover, for every 1-simplex  $\tau$  and  $p \in \tau$  assume that  $u$  is given in a neighbourhood of  $u(p)$  in local coordinates by  $u = (u^1, \dots, u^M)$ .*

Then,

$$(3.1) \quad \sum_{\sigma > \tau} \frac{\partial u_j^m}{\partial e_\sigma} = 0$$

where  $\sigma > \tau$  implies that  $\tau$  is a face of  $\sigma$ , and  $e_\sigma$  is an inward pointing normal vector field along  $\tau$  in  $\sigma$ .

*Proof.* The fact that  $u^m \in C^\infty(\sigma \cup \tau)$  follows from Theorem 4 and Corollary 6 of [12]. Equation (3.1), follows from Corollary 5 of [12].  $\square$

For an edge  $\tau$  and  $\sigma \in \mathcal{S}_2(\tau)$ , we define polar coordinates  $(r, \theta)$  of  $\sigma \cup \tau$  centered at  $p$  by setting  $r$  to be the distance from  $p$  to a point  $q \in \sigma \cup \tau$  and  $\theta$  to be the angle between  $\tau$  and the line  $\overline{pq}$  connecting  $p$  and  $q$ . The next theorem is one of the main technical results of the paper and describes the singular behavior of harmonic maps near the lower dimensional strata.

**Theorem 3.3.** *Let  $u : X \rightarrow Y$  be a harmonic map. If  $(r, \theta)$  are the polar coordinates of  $\sigma \cup \tau$  centered at a 0-cell  $p$  and  $u$  is given in local coordinates  $(u^1, \dots, u^M)$  in a neighbourhood of  $u(p)$ , we have the following derivative bounds for  $u^m$  in a neighbourhood of  $p$ :*

$$\left| \frac{\partial u^m}{\partial r} \right| \leq Cr^{\alpha-1}, \quad \left| \frac{\partial u^m}{\partial \theta} \right| \leq Cr^\alpha$$

$$\begin{aligned} \left| \frac{\partial^2 u^m}{\partial r^2} \right| &\leq Cr^{\alpha-2}, \quad \left| \frac{\partial^2 u^m}{\partial r \partial \theta} \right| \leq Cr^{\alpha-1}, \quad \left| \frac{\partial^2 u^m}{\partial \theta^2} \right| \leq Cr^\alpha \\ \left| \frac{\partial^3 u^m}{\partial r^3} \right| &\leq Cr^{\alpha-3}, \quad \left| \frac{\partial^3 u^m}{\partial^2 r \partial \theta} \right| \leq Cr^{\alpha-2}, \quad \left| \frac{\partial^3 u^m}{\partial r \partial^2 \theta} \right| \leq Cr^{\alpha-1}, \quad \left| \frac{\partial^3 u^m}{\partial \theta^3} \right| \leq Cr^\alpha \end{aligned}$$

for some constants  $C > 0$  and  $\alpha > 0$  depending on the total energy of  $u$  and the geometry of the complex  $X$ . Furthermore,  $\alpha$  can be chosen independently of the choice of the 0-cell  $p$  of  $X$ .

*Proof.* Let  $\sigma = \sigma_1, \dots, \sigma_J$  be the 2-cells in  $\mathcal{S}_2(\tau)$ . For each  $j = 1, \dots, J$ , we let  $(x, y)$  be the Euclidean coordinates of  $\overline{\sigma_j \cup \tau}$  so that (i)  $p$  is given as  $(x, y) = (0, 0)$ , (ii) if  $(x, y) \in \tau$  then  $x > 0$  and  $y = 0$  and (iii) if  $(x, y) \in \sigma_j$  then  $x, y > 0$ . Let  $u_j^m = u^m|_{\sigma_j}$ .

We will now compute the first derivative bounds with respect to the polar coordinates  $r$  and  $\theta$ . By Theorem 6.2 of [11], we have the inequality

$$|\nabla u|^2(r, \theta) \leq Cr^{2\alpha-2}$$

for some  $\alpha > 0$ . More specifically,  $\alpha$  can be chosen to be the order of  $u$  at  $p$ , i.e.

$$\alpha = \lim_{r \rightarrow 0} \frac{r \int_{B_r(p)} |\nabla u|^2 d\mu}{\int_{\partial B_r(p)} d^2(u, u(p)) ds}.$$

Hence,

$$(3.2) \quad \left| \frac{\partial u_j^m}{\partial x} \right| \leq Cr^{\alpha-1} \quad \text{and} \quad \left| \frac{\partial u_j^m}{\partial y} \right| \leq Cr^{\alpha-1}.$$

Using the fact that  $x = r \cos \theta$  and  $y = r \sin \theta$ , we get

$$\frac{\partial u_j^m}{\partial r} = \frac{\partial u_j^m}{\partial x} \cos \theta + \frac{\partial u_j^m}{\partial y} \sin \theta \quad \text{and} \quad \frac{\partial u_j^m}{\partial \theta} = -\frac{\partial u_j^m}{\partial x} r \sin \theta + \frac{\partial u_j^m}{\partial y} r \cos \theta.$$

This immediately implies

$$\left| \frac{\partial u_j^m}{\partial r} \right| \leq Cr^{\alpha-1} \quad \text{and} \quad \left| \frac{\partial u_j^m}{\partial \theta} \right| \leq Cr^\alpha.$$

We will now establish the second derivative estimates of  $u_j^m$  for a points  $(r, \theta)$  on  $\sigma_j \cup \tau$  with  $\theta$  sufficiently small. We will need the following notations: for a function  $\varphi$  and a domain  $\Omega \subset \mathbb{R}^2$ , we set

$$\begin{aligned} |\varphi|_{0;\Omega} &= \sup_{p \in \Omega} |\varphi(p)| \\ |D\varphi|_{0;\Omega} &= \sup_{p \in \Omega} \max \left\{ \left| \frac{\partial \varphi}{\partial x}(p) \right|, \left| \frac{\partial \varphi}{\partial y}(p) \right| \right\} \\ |D^2\varphi|_{0;\Omega} &= \sup_{p \in \Omega} \max \left\{ \left| \frac{\partial^2 \varphi}{\partial x^2}(p) \right|, \left| \frac{\partial^2 \varphi}{\partial x \partial y}(p) \right|, \left| \frac{\partial^2 \varphi}{\partial y^2}(p) \right| \right\} \\ [\varphi]_{\beta;\Omega} &= \sup_{p, q \in \Omega, p \neq q} \frac{|\varphi(p) - \varphi(q)|}{|p - q|^\beta} \\ [D\varphi]_{\beta;\Omega} &= \sup_{p, q \in \Omega, p \neq q} \frac{1}{|p - q|^\beta} \max \left\{ \left| \frac{\partial \varphi}{\partial x}(p) - \frac{\partial \varphi}{\partial x}(q) \right|, \left| \frac{\partial \varphi}{\partial y}(p) - \frac{\partial \varphi}{\partial y}(q) \right| \right\} \end{aligned}$$

$$[D^2\varphi]_{\beta;\Omega} = \sup_{p,q \in \Omega, p \neq q} \frac{1}{|p-q|^\beta} \max \left\{ \left| \frac{\partial^2 \varphi}{\partial x^2}(p) - \frac{\partial^2 \varphi}{\partial x^2}(q) \right|, \left| \frac{\partial^2 \varphi}{\partial x \partial y}(p) - \frac{\partial^2 \varphi}{\partial x \partial y}(q) \right|, \left| \frac{\partial^2 \varphi}{\partial y^2}(p) - \frac{\partial^2 \varphi}{\partial y^2}(q) \right| \right\}.$$

Let  $T := \{(x, y) \in \mathbb{R}^2 : y \geq 0, y < \sqrt{3}x, y < -\sqrt{3}x + \sqrt{3}\}$ ,  $T^- = \{(x, -y) \in \mathbb{R}^2 : (x, y) \in T\}$  and  $\hat{T} = T \cup T^-$ . Fix  $m$  and  $j$  and define  $U : \hat{T} \rightarrow \mathbb{R}$  by setting

$$U(x, y) = \begin{cases} u_j^m(x, y) & \text{if } y \geq 0 \\ -u_j^m(x, -y) + \frac{2}{J} \sum_{j'=1}^J u_{j'}^m(x, -y) & \text{if } y < 0. \end{cases}$$

Let

$$(3.3) \quad \Gamma_j^m = \sum_{p,q=1}^M \Gamma_{pq}^m(u_j) \left( \frac{\partial u_j^p}{\partial x} \frac{\partial u_j^q}{\partial x} + \frac{\partial u_j^p}{\partial y} \frac{\partial u_j^q}{\partial y} \right)$$

where  $\Gamma_{pq}^m$  are the Christoffel symbols of  $Y$  with respect to the local coordinates  $(u^1, \dots, u^M)$ . Since the harmonic map equation

$$\Delta u_j^m = \Gamma_j^m$$

is satisfied in  $T$ , if we set

$$f(x, y) = \begin{cases} \Gamma_j^m(x, y) & \text{if } y \geq 0 \\ -\Gamma_j^m(x, -y) + \frac{2}{J} \sum_{j'=1}^J \Gamma_{j'}^m(x, -y) & \text{if } y < 0, \end{cases}$$

then  $U$  satisfies the Poisson equation

$$(3.4) \quad \Delta U = f$$

weakly in  $\hat{T}$ . Indeed, let  $\xi$  be a test function supported in a neighbourhood  $B_R(q)$  of a point  $q = (x_0, 0) \in \hat{T}$ . Since  $U$  is a  $C^1$  function we have by the divergence theorem,

$$\begin{aligned} \int_{\hat{T}} \operatorname{div}(\xi \nabla U) dx dy &= \int_T \operatorname{div}(\xi \nabla U) dx dy + \int_{T^-} \operatorname{div}(\xi \nabla U) dx dy \\ &= - \int_{x_0-R}^{x_0+R} \xi \frac{\partial u_j^m}{\partial y}(x, 0) dx \\ &\quad + \int_{x_0-R}^{x_0+R} \left( \xi \frac{\partial u_j^m}{\partial y}(x, 0) - \frac{2}{J} \sum_{j'=1}^J \xi \frac{\partial u_{j'}^m}{\partial y}(x, 0) \right) dx \\ &= 0 \end{aligned}$$

where the last equality is because of (3.1). On the other hand

$$\int_{\hat{T}} \operatorname{div}(\xi \nabla U) dx dy = \int_{\hat{T}} \nabla \xi \cdot \nabla U + \int_{\hat{T}} \xi f dx dy$$

which along with the previous equation implies (3.4). If  $B_{2R}(q) \subset \hat{T}$ , then elliptic regularity theory (cf. [18] or Lemma 3 on p13 of [31]) implies

$$R^{1+\beta} [DU]_{\beta; B_{\frac{3R}{2}}(q)} \leq C(|U|_{0; B_{2R}(q)} + R^2 |f|_{0; B_{2R}(q)}).$$

If we choose  $R$  to be the largest number so that  $B_{2R}(q) \subset \hat{T}$ , then  $R$  is proportional to  $r$  where  $r$  is the distance of  $q$  to the vertex  $p$ . Furthermore, the distance from  $p$  to any

point of  $B_{2R}(q)$  is bounded uniformly by some constant multiple of  $r$ . Hence, assuming  $U(0, 0) = 0$  without a loss of generality, we have

$$[DU]_{\beta; B_{\frac{3R}{2}}(p)} \leq C(r^{-1-\beta}|U|_{0; B_{2R}(p)} + r^{1-\beta}|f|_{0; B_{2R}(p)}) \leq C(r^{-1-\beta+\alpha} + r^{-\beta+2\alpha-1}) \leq Cr^{-\beta+\alpha-1}.$$

Here, we have used the Hölder continuity of  $u_j^m$  (hence of  $U$ ) near  $p$  with Hölder exponent  $\alpha$  (cf. Theorem 3.7 of [11]) and the inequalities of (3.2) along with the fact that  $f$  is quadratic in  $Du_j^m$  from (3.3). Thus, with  $B_{\frac{3R}{2}}^+(q) = B_{\frac{3R}{2}}(q) \cap \{y \geq 0\}$ , we obtain

$$[Du_j^m]_{\beta; B_{\frac{3R}{2}}^+(q)} \leq Cr^{-\beta+\alpha-1}.$$

This equation along with (3.2) and (3.3) implies that

$$(3.5) \quad [\Gamma_j^m]_{\beta; B_{\frac{3R}{2}}^+(q)} \leq C|Du_j^k|_{0; B_{\frac{3R}{2}}^+(q)}[Du_j^\ell]_{\beta; B_{\frac{3R}{2}}^+(q)} \leq Cr^{-\beta+2\alpha-2}.$$

We are now ready to prove the second derivative bounds of  $u_j^m$ . Note that we have the set of partial differential equations

$$(3.6) \quad \Delta u_j^m = \Gamma_j^m \quad j = 1, \dots, J \quad m = 1, \dots, M$$

in  $T$ , along with boundary conditions

$$(3.7) \quad u_j^m - u_1^m = 0 \quad j = 2, \dots, J \quad m = 1, \dots, M$$

$$(3.8) \quad \sum_{j=1}^J \frac{\partial u_j^m}{\partial y} = 0 \quad m = 1, \dots, M$$

in  $B = \{(x, y) \in \mathbb{R}^2 : y = 0, 0 < x < 1\}$ . This is a system of  $JM$  number of equations containing  $JM$  number of unknowns (i.e.  $u_j^m$ ) along with  $JM$  number of boundary conditions. If we assign weights  $s_j^m = 0$  to the equations, weights  $t_j^m = 2$  to the unknowns, weights  $r_j^m = -2$  for  $j = 2, \dots, M$  and  $r_1^m = -1$  to the boundary conditions, then this system is said to be elliptic with complementing boundary condition according to the elliptic regularity theory of [3] (or elliptic and coercive in [28]). Hence, we have the Schauder estimates (cf. Theorem 9.1 of [3]),

$$\begin{aligned} & R^2|D^2u_j^m|_{0; B_R^+(q)} + R^{2+\beta}[D^2u_j^m]_{\beta; B_R^+(q)} \\ & \leq C(|\Gamma_j^m|_{0; B_{\frac{3R}{2}}^+(q)} + R^{2+\beta}[\Gamma_j^m]_{\beta; B_{\frac{3R}{2}}^+(q)} + |u_j^m|_{0; B_{\frac{3R}{2}}^+(q)}). \end{aligned}$$

With the same choice of  $q$  and  $R$  as above, we obtain

$$|D^2u_j^m|_{0; B_R^+(q)} \leq C(|\Gamma_j^m|_{0; B_{\frac{3R}{2}}^+(q)} + r^\beta[\Gamma_j^m]_{\beta; B_{\frac{3R}{2}}^+(q)} + r^{-2}|u_j^m|_{0; B_{\frac{3R}{2}}^+(q)}).$$

The above inequality, along with (3.5), implies

$$|D^2u_j^m|_{0; B_R^+(q)} \leq C(r^{2\alpha-2} + r^{2\alpha-2} + r^{\alpha-2}) \leq Cr^{\alpha-2}.$$

Since

$$\frac{\partial^2 u_j^m}{\partial r^2} = \frac{\partial^2 u_j^m}{\partial x^2} \cos^2 \theta + 2 \frac{\partial^2 u_j^m}{\partial x \partial y} \sin \theta \cos \theta + \frac{\partial^2 u_j^m}{\partial y^2} \sin^2 \theta,$$

$$\begin{aligned} \frac{\partial^2 u_j^m}{\partial r \partial \theta} &= -\frac{\partial^2 u_j^m}{\partial x^2} r \sin \theta \cos \theta + \frac{\partial^2 u_j^m}{\partial x \partial y} r \cos^2 \theta - \frac{\partial u_j^m}{\partial x} \sin \theta \\ &\quad - \frac{\partial^2 u_j^m}{\partial x \partial y} r \sin^2 \theta + \frac{\partial^2 u_j^m}{\partial y^2} r \sin \theta \cos \theta + \frac{\partial u_j^m}{\partial y} \cos \theta, \\ \frac{\partial^2 u_j^m}{\partial \theta^2} &= \frac{\partial^2 u_j^m}{\partial x^2} r^2 \sin^2 \theta + 2 \frac{\partial^2 u_j^m}{\partial x \partial y} r^2 \sin^2 \theta + \frac{\partial^2 u_j^m}{\partial y^2} r^2 \cos \theta \\ &= -\frac{\partial u_j^m}{\partial x} r \cos \theta - \frac{\partial u_j^m}{\partial y} r \sin \theta, \end{aligned}$$

we immediately obtain

$$\left| \frac{\partial^2 u_j^m}{\partial r^2} \right| \leq C r^{\alpha-2}, \quad \left| \frac{\partial^2 u_j^m}{\partial r \partial \theta} \right| \leq C r^{\alpha-1} \quad \text{and} \quad \left| \frac{\partial^2 u_j^m}{\partial \theta^2} \right| \leq C r^\alpha$$

at  $(r, \theta)$  for  $\theta$  sufficiently small. This restriction on  $\theta$  is due to the choice of  $R$  and  $q$ . For  $(r, \theta)$  with  $\theta$  sufficiently large, we can use a similar argument using standard elliptic regularity theory (e.g. [18] or Lemma 3 on p13 of [31]) in the interior of  $\sigma$ . The third derivative estimates follow the same way from the first two by bootstrapping the elliptic equations (3.6) with boundary conditions (3.8) and (3.8).

Section 4 of [12] shows that the order of  $u$  at  $p$  can be bounded from below by  $2\lambda_v^{comb}$  where  $\lambda_v^{comb}$  is the combinatorial eigenvalue of the link of  $v$  which is always a positive quantity. Hence choosing  $\alpha$  to be the minimum of  $2\lambda_v^{comb}$  over all 0-cells of  $X$ , we have established the last assertion of the Theorem.  $\square$

**3.2. Weighted Sobolev spaces.** In this subsection we recall the important features of the weighted Sobolev spaces used in this paper. The main references are [2], [15], and [29]. In the following we fix a smooth vector bundle  $E$  of rank  $r$  over a 2-complex  $X$  with a Hermitian metric, and a fixed Riemannian metric on the base space  $X$ . Define the space  $C_0^\infty(E)$  to be the space of smooth sections  $s \in \Omega^0(X, E)$  that satisfy  $s(p) = 0$  whenever  $p$  is a vertex of  $X$ . In the local model  $\tilde{B}(r)$  around each vertex  $p$ , we define local co-ordinates  $(t, \theta) = (-\log r, \theta)$ , where  $(r, \theta)$  are the standard polar co-ordinates in a neighbourhood of the vertex  $p$ . To define a norm on  $C_0^\infty(E)$ , let  $\{x_i\}_{i=1, \dots, V}$  denote the vertices of  $X$  and choose disjoint open neighbourhoods  $U_{x_i}$  for each vertex  $x_i$ . Then cover the rest of  $X$  with open sets  $\{V_\alpha\}_{\alpha=1, \dots, K}$  that do not contain any of the vertices. For  $\delta \in \mathbb{R}$ , the space  $L_\delta^p$  is the completion of  $C_0^\infty(E)$  in the norm

$$(3.9) \quad \|s\|_{L_\delta^p} = \left( \sum_{i=1}^V \int_{U_{x_i}} e^{t\delta} |s|^p + \sum_{\alpha=1}^K \int_{V_\alpha} |s|^p \right)^{1/p}$$

where we use  $e^{t\delta}$  to denote the co-ordinates in a neighbourhood of a vertex. Away from all of the vertices,  $e^{t\delta}$  is bounded and  $s$  is continuous, and so the question of whether the norm  $\|\cdot\|_{L_\delta^p}$  is finite only depends on the choice of co-ordinates near each vertex. Different choices of  $V_\alpha$  will lead to equivalent norms.

Given a vertex  $p$ , and a trivialization of  $E$  near  $p$  we say that a connection is *trivial* in a neighbourhood of  $p$  if with respect to the above trivialization  $\nabla = d$ . Given a fixed

connection  $\nabla_0$  trivial near the vertices, and a positive integer  $k$ , we define the *weighted Sobolev space*  $L_{k,\delta}^q(E)$  as the completion of  $C_0^\infty(E)$  in the norm

$$(3.10) \quad \|s\|_{L_{k,\delta}^q} = \sum_{\ell=0}^k \|\nabla_0^\ell s\|_{L_\delta^q}$$

Note that in this paper we are considering bundles with a fixed trivialization on the universal cover (cf. Remark 2.31). Since the star of a vertex  $p$  in  $X$  is simply connected it follows that we have a fixed trivialization of  $E$  in a neighbourhood of  $p$ . It thus makes sense to talk about connections on  $E$  trivial near the vertices.

It is a standard fact that the spaces  $L_{k,\delta}^q$  do not change if we either: (a) change the connection  $\nabla_0$  outside a neighbourhood of the vertices of  $X$ , or (b) change the co-ordinates outside a neighbourhood of the vertices. The usual multiplication theorems for Sobolev spaces on compact manifolds carry over to the weighted Sobolev spaces studied here. To be more precise, we have that the multiplication map  $L_{s_1,\delta_1}^2 \times L_{s_2,\delta_2}^2 \rightarrow L_{s,\delta}^2$  is continuous if  $s_1, s_2 \geq s$ ,  $s < s_1 + s_2 - n/2$  and  $\delta < \delta_1 + \delta_2 + n/2$ , where  $n$  is the dimension of the complex  $X$ .

Following Section 3.1 of [15] we define the space of weighted connections  $\mathcal{A}_\delta^\mathbb{C}(E)$  to be the space of all connections whose connection form is an element of  $L_{1,\delta}^2$ , and the space  $\mathcal{A}_\delta(E) \subset \mathcal{A}_\delta^\mathbb{C}(E)$  to be the subset of all unitary connections. The weighted gauge group  $\mathcal{G}_\delta(E)$  is defined as follows. Let  $\nabla_0$  be a connection as above and define

$$(3.11) \quad \mathcal{R} = \left\{ v \in L_{2,loc}^2(\text{End}(E)) : \|\nabla_0 v\|_{L_{1,\delta}^2} < \infty \right\}$$

Then the *weighted gauge group* is defined as

$$(3.12) \quad \mathcal{G}_\delta(E) = \{v \in \mathcal{R} : vv^* = \text{id}, \det v = 1\}$$

and the complexified gauge group is

$$(3.13) \quad \mathcal{G}_\delta^\mathbb{C}(E) = \{v \in \mathcal{R} : \det v = 1\}.$$

The multiplication theorem for weighted Sobolev spaces shows that both  $\mathcal{G}_\delta(E)$  and  $\mathcal{G}_\delta^\mathbb{C}(E)$  have a group structure, and that there are well-defined actions of  $\mathcal{G}_\delta(E)$  on  $\mathcal{A}_\delta$  and  $\mathcal{G}_\delta^\mathbb{C}(E)$  on  $\mathcal{A}_\delta^\mathbb{C}(E)$  respectively.

Similarly we have balanced versions of these spaces  $\mathcal{G}_{bal,\delta}(E)$ ,  $\mathcal{A}_{bal,\delta}(E)$  and  $\Omega_{bal,\delta}^1(\text{ad}(E))$ . When a smooth pair  $(d_A, \psi) \in \mathcal{A}_{bal,\delta}(E) \times \Omega_{bal,\delta}^1(\text{ad}(E))$  solves the equations (2.10) and (2.11), then the *holonomy of the pair*  $(d_A, \psi)$  refers to the holonomy of the flat connection  $d_A + \psi \in \mathcal{A}_{bal,flat,\delta}^\mathbb{C}(E)$ .

**Proposition 3.4.** *If  $D_i \in \mathcal{A}_{bal,flat,\delta}^\mathbb{C}(E)$ ,  $i = 1, 2$  are smooth and  $\mathcal{G}_{bal,\delta}^\mathbb{C}(E)$ -gauge equivalent then they are  $\mathcal{G}_{bal}^\mathbb{C}(E)$ -gauge equivalent.*

*Proof.* Since the result is local, it follows by elliptic regularity.  $\square$

**Proposition 3.5.** *Let  $D \in \mathcal{A}_{bal,flat,\delta}^\mathbb{C}(E)$  be smooth. Then  $D$  has trivial holonomy around the vertices of  $X$ .*



*Proof.* For  $D = d + A$  write  $A(t, \theta) = B(t, \theta)dt + C(t, \theta)d\theta$ . Consider the family of loops  $c_t : [0, 2\pi] \rightarrow X$  given by  $c_t(\theta) = (t, \theta)$  and consider the holonomy equation from Definition 2.18 along  $c_t(\theta)$

$$(3.14) \quad \frac{ds_t(\theta)}{d\theta} + C(t, \theta)s_t(\theta) = 0 \quad \text{with } s_t(0) = id.$$

Lemma IV.4.1 on p54 of [22] implies

$$(3.15) \quad |s_t(\theta)| \leq |s_t(0)| \exp \left\{ \int_0^\theta |C(t, \theta)| d\theta \right\} \leq K \exp \left\{ \int_0^{2\pi} |C(t, \theta)| d\theta \right\}$$

where  $K$  is a dimensional constant. Since

$$\int_0^\infty e^{t\delta} \int_0^{2\pi} |C(t, \theta)|^2 d\theta dt < \infty$$

there exists a sequence  $t_i \rightarrow \infty$  such  $\int_0^{2\pi} |C(t_i, \theta)|^2 d\theta \rightarrow 0$ . By Cauchy-Schwarz we also have

$$(3.16) \quad \int_0^{2\pi} |C(t_i, \theta)| d\theta \rightarrow 0.$$

Combined with (3.15) this implies that  $|s_{t_i}(\theta)|$  is uniformly bounded. By integrating equation (3.14) with respect to  $\theta$ , we obtain from (3.16)

$$(3.17) \quad |s_{t_i}(2\pi) - s_{t_i}(0)| \leq \int_0^{2\pi} |s_{t_i}(\theta)| |C(t_i, \theta)| d\theta \rightarrow 0.$$

Since the holonomy is independent of  $t$  we obtain that  $s_{t_i}(2\pi) = s_{t_i}(0)$  and thus it must be trivial.  $\square$

Proposition 3.4 and Proposition 3.5 allow us to define the notion of conjugacy class of holonomy for a smooth flat connection  $D \in \mathcal{A}_{bal, flat, \delta}^{C, irr}(E)$  as follows.

**Definition 3.6.** Let  $D \in \mathcal{A}_{bal, flat, \delta}^{C, irr}(E)$  be a smooth flat connection and let  $\rho_* : \pi_1(X_*) \rightarrow \text{SL}(r, \mathbb{C})$  be the holonomy of  $D$ , where  $X_* = X \setminus X^0$  and  $X^0$  denotes the 0-skeleton of  $X$ . Since the star of a vertex is contractible, then Van Kampen's theorem implies that  $\pi_1(X) = \pi_1(X_*)/\pi$ , where  $\pi$  denotes the subgroup of  $\pi_1(X_*)$  generated by  $\cup_{p \in X^0} \pi_1(Lk(p))$ . By Proposition 3.5, the restriction of  $\rho_*$  to  $\pi$  is trivial hence it induces a homomorphism  $\rho : \pi_1(X) \rightarrow \text{SL}(r, \mathbb{C})$ . We say that the conjugacy class of holonomy of  $D$  is  $[\rho]$ . Notice that the map is well defined since gauge equivalent pairs yield conjugate holonomies. Furthermore,  $\rho$  is irreducible because  $D$  is irreducible.

## 4. EQUIVALENCE OF MODULI SPACES

**4.1. Higgs Moduli Space.** We fix a vector bundle  $E_c = E$  of rank  $r$  over a 2-complex  $X$  with a Hermitian metric, and a fixed Riemannian metric on the base space  $X$ .

**Definition 4.1.** The *Higgs moduli space* is the space  $\mathcal{M}_{Higgs}(E)$  of  $\mathcal{G}_{bal, \delta}(E)$  equivalence classes of pairs  $(d_A, \psi) \in \mathcal{A}_{bal, \delta}(E) \times \Omega_{bal, \delta}^1(\sqrt{-1} \text{ad}(E))$  that are *smooth, irreducible* and

solve the following equations

$$(4.1) \quad F_A + \psi \wedge \psi = 0$$

$$(4.2) \quad d_A \psi = 0$$

$$(4.3) \quad d_A^* \psi = 0.$$

We endow  $\mathcal{M}_{Higgs}(E)$  with the  $L^2_{1,\delta}$ -topology.

Given  $[(d_A, \psi)] \in \mathcal{M}_{Higgs}(E)$ , we can assign by Definition 3.6 the holonomy  $[\rho]$  of the flat connection  $d_A + \psi$  and set  $\alpha[(d_A, \psi)] := [\rho]$ . The map  $\alpha$  is well defined. The next proposition follows from continuous dependence of solutions of ODE upon the initial condition.

**Proposition 4.2.** *The map  $\alpha : \mathcal{M}_{Higgs}(E) \rightarrow \mathcal{M}_{char}^c$ , where  $\alpha[(d_A, \psi)] = [\rho]$ , is well defined and continuous.*

The following is the main theorem of this paper.

**Theorem 4.3.** *The map  $\alpha : \mathcal{M}_{Higgs}(E) \rightarrow \mathcal{M}_{char}^c$  is a homeomorphism.*

In the next section we will construct the inverse map. We end this section with a proposition that will be used later.

**Proposition 4.4.** *Let  $(d_{A_1}, \psi_1)$  and  $(d_{A_2}, \psi_2)$  be solutions to equations (4.1)-(4.3) and assume that they are  $\mathcal{G}_{bal,\delta}^{\mathbb{C}}(E)$ -gauge equivalent. Then they are  $\mathcal{G}_{bal,\delta}(E)$ -gauge equivalent.*

*Proof.* Assume that there exists  $g \in \mathcal{G}_{bal,\delta}^{\mathbb{C}}(E)$  such that  $(d_{A_1}, \psi_1) = g \cdot (d_{A_2}, \psi_2)$ , and we have to show that  $g$  is unitary. Let  $h = g^* g$  and we will show that  $h$  is constant. By [32] Lemma 3.1(d) we have the following point wise estimate away from the vertices (notice that the sign of our Laplacian is the opposite from Simpson's)

$$(4.4) \quad \Delta \operatorname{tr}(h) \leq 0.$$

Now since  $g$  is balanced, then so is  $\operatorname{tr} h$ , and therefore an application of Stokes' theorem on each face of  $X$  shows that

$$(4.5) \quad \begin{aligned} \int_X \Delta \operatorname{tr} h \, dx &= \lim_{r \rightarrow 0} \int_{X \setminus \bigcup_{0\text{-cells } v} B_r(v)} \Delta \operatorname{tr} h \, dx \\ &= \lim_{r \rightarrow 0} \sum_{2\text{-cells } \sigma} \int_{F \setminus \bigcup_{0\text{-cells } v} B_r(v)} \Delta \operatorname{tr} h \, dx \\ &= \lim_{r \rightarrow 0} \sum_{2\text{-cells } \sigma} \int_{\partial(F \setminus \bigcup_{0\text{-cells } v} B_r(v))} \frac{\partial \operatorname{tr} h}{\partial \nu} \, ds \end{aligned}$$

where  $\nu$  is the outwards pointing normal vector on  $\partial(\sigma \setminus \bigcup_{0\text{-cells } v} B_r(v))$ . The boundary  $\partial(\sigma \setminus \bigcup_{\text{vertices } v} B_r(v))$  consists of points on the 1-cells of  $\sigma$ , and points on  $\partial B_r(v) \cap \sigma$ . Breaking the integral into these two parts, we obtain

$$(4.6) \quad \sum_{2\text{-cells } \sigma} \int_{\partial(\sigma \setminus \bigcup_{0\text{-cells } v} B_r(v))} \frac{\partial \operatorname{tr} h}{\partial \nu} \, ds = \sum_{2\text{-cells } \sigma} \left( \sum_{1\text{-cells } \tau : \tau \cap \bar{\sigma} \neq \emptyset} \int_{\tau \setminus \bigcup_v B_r(v) \cap \tau} \frac{\partial \operatorname{tr} h}{\partial \nu} \, ds \right) + \sum_{2\text{-cells } \sigma} \int_{\bigcup_v \partial B_r(v) \cap \sigma} \frac{\partial \operatorname{tr} h}{\partial \nu} \, ds$$

The balancing condition shows that the first term is zero. Therefore we are left with

$$(4.7) \quad \int_X \Delta \operatorname{tr} h \, dx = \lim_{r \rightarrow 0} \sum_{2\text{-cells } \sigma} \int_{\cup_v \partial B_r(v) \cap \sigma} \frac{\partial \operatorname{tr} h}{\partial \nu} \, ds$$

In polar co-ordinates, each component of this integral becomes

$$(4.8) \quad \int_{\partial B_r(v) \cap F} \frac{\partial \operatorname{tr} h}{\partial \nu} \, ds = r \int_0^{\frac{\pi}{3}} \frac{\partial \operatorname{tr} h}{\partial r} \, d\theta$$

Since  $h \in \mathcal{G}(E)_{bal,\delta}^{\mathbb{C}}$  (and in particular, the integral of  $\frac{\partial^2 h}{\partial r^2}$  is bounded), then we have

$$(4.9) \quad \lim_{r \rightarrow 0} \left( \sigma \int_0^{\frac{\pi}{3}} \operatorname{tr} \left( \frac{\partial h}{\partial r} \right) \, d\theta \right) = 0$$

and so (4.7) becomes

$$(4.10) \quad \int_X \Delta \operatorname{tr} h \, dx = 0.$$

Combined with  $\Delta \operatorname{tr} h \leq 0$  from (4.4), we see that  $\Delta \operatorname{tr} h = 0$ . The second to the last formula in [32, p876] implies that  $D(h) = 0$  pointwise away from the vertices. This implies that the connection  $D$  splits according to the eigenspaces of  $h$ , and since the connection  $D$  is indecomposable, then  $h$  must be a constant multiple of the identity matrix, which concludes the proof.  $\square$

**4.2. The inverse map.** For an irreducible representation  $\rho : \pi_1(X) \rightarrow \operatorname{SL}(r, \mathbb{C})$ , with  $[\rho] \in \mathcal{M}_{char}^c$  and  $E = E_c$ , Theorem 3.3 then shows that there exists a unique  $\rho$ -equivariant harmonic map  $u : \tilde{X} \rightarrow \operatorname{SL}(r, \mathbb{C})/\operatorname{SU}(r)$ . As in subsection 2.2, let  $d_A$  and  $\psi$  be the associated unitary connection and Higgs field. Since  $u$  is harmonic,  $d_A$  is the pullback of the Levi-Civita connection on  $\operatorname{SL}(r, \mathbb{C})/\operatorname{SU}(r)$ , and  $\psi$  is the derivative of  $u$ , then we also have the equation

$$(4.11) \quad d_A^* \psi = 0$$

almost everywhere (in fact by Theorem 3.3 everywhere away from the 0-skeleton).

**Proposition 4.5.** *If  $u$  is harmonic,  $\alpha$  as in Theorem 3.3 and  $\delta < \alpha$ , then  $\mathcal{D} \in \mathcal{A}_{bal,flat,\delta}^{\mathbb{C}}(E)$ . The metric on the bundle  $E$  induces a decomposition of  $\mathcal{D}$  into skew-adjoint and self-adjoint parts,  $\mathcal{D} = d_A + \psi$ , where  $d_A \in \mathcal{A}_{bal,\delta}(E)$  and  $\psi \in \Omega_{bal,\delta}^1(i \operatorname{ad}(E))$ . Furthermore,  $\mathcal{D}$ ,  $d_A$  and  $\psi$  are smooth (over  $X_*$ ).*

*Proof.* The construction in Section 2.2 shows that the connection  $\mathcal{D}$  is induced from the trivial connection on the universal cover, hence it is clearly balanced, flat and  $L_{1,\delta}^2$ . Furthermore, since  $d_A = u^* \nabla$  and  $\psi = u^{-1} du$ , Theorem 3.3 and (2.9) imply that  $d_A$  and  $\psi$  are balanced. Therefore, since  $u : X \rightarrow \operatorname{SL}(r, \mathbb{C})/\operatorname{SU}(r)$  is a Lipschitz map over the compact space  $X$ , in order to show that  $d_A \in \mathcal{A}_{bal,\delta}(E)$  and  $\psi \in \Omega_{bal,\delta}^1(i \operatorname{ad}(E))_\delta$ , it suffices to show that to show that  $du \in L_{1,\delta}^2$ .

First we show that  $du \in L_\delta^2$ . Theorem 3.3 shows that  $|\frac{\partial u}{\partial r}| \leq Cr^{\alpha-1}$  and  $|\frac{\partial u}{\partial \theta}| \leq Cr^\alpha$  for some positive  $\alpha$ . Using the co-ordinate transformation  $r = e^{-t}$  we see that  $|\frac{\partial u}{\partial \theta}| \leq Ce^{-\alpha t}$ ,

and

$$\begin{aligned} \left| \frac{\partial u}{\partial t} \right| &= \left| \frac{\partial u}{\partial r} \frac{dr}{dt} \right| \\ &\leq Cr^{\alpha-1}r \\ &= Ce^{-\alpha t} \end{aligned}$$

Therefore,  $du \in L_\delta^2$  if  $\delta < \alpha$ . Similarly, we use the estimates on the second derivatives of  $u$  to show that  $du \in L_{1,\delta}^2$ . We have  $\left| \frac{\partial^2 u}{\partial \theta^2} \right| \leq Ce^{-\alpha t}$ , and we can compute

$$\begin{aligned} \left| \frac{\partial^2 u}{\partial t \partial \theta} \right| &= \left| \frac{\partial^2 u}{\partial r \partial \theta} \frac{dr}{dt} \right| \\ &\leq Cr^{\alpha-1}r \\ &= Ce^{-\alpha t} \end{aligned}$$

and similarly

$$\begin{aligned} \left| \frac{\partial^2 u}{\partial t^2} \right| &= \left| \frac{\partial}{\partial t} \left( \frac{\partial u}{\partial r} \frac{dr}{dt} \right) \right| \\ &= \left| \frac{\partial^2 u}{\partial r^2} \left( \frac{dr}{dt} \right)^2 + \frac{\partial u}{\partial t} \left( \frac{d}{dr} \frac{dr}{dt} \right) \frac{dr}{dt} \right| \\ &\leq Cr^{\alpha-2}r^2 + Cr^{\alpha-1}r^2 \\ &\leq Ce^{-\alpha t} + Ce^{-(\alpha+1)t} \\ &\leq Ce^{-\alpha t}, \end{aligned}$$

where in the last step we use the fact that  $t \geq 0$  near a vertex. Therefore,  $du \in L_{1,\delta}^2$  if  $\delta < \alpha$ .  $\square$

**Theorem 4.6.** *The map  $\beta : \mathcal{M}_{char}^c \rightarrow \mathcal{M}_{Higgs}(E)$  defined by  $\beta([\rho]) = [(d_A, \psi)]$  is a continuous inverse of  $\alpha$ .*

*Proof.* The first step is to show that the map  $\beta$  is well defined. Given  $\rho$ , Proposition 4.5 implies that  $d_A \in \mathcal{A}_{bal,\delta}(E)$  and  $\psi \in \Omega_{bal^1,\delta}(i \operatorname{ad}(E))$ . Moreover, we claim that the pair  $(d_A, \psi)$  is irreducible. If  $\rho_* : \pi_1(X_*) \rightarrow \operatorname{SL}(r, \mathbb{C})$  denotes the holonomy of the flat connection  $d_A + \psi$  then, as pointed out in Definition 3.6,  $\rho_* = \rho \circ p$ , where  $p : \pi_1(X_*) \rightarrow \pi_1(X) = \pi_1(X_*)/\pi$  is the natural quotient map. Since by assumption  $\rho$  is irreducible it follows that  $\rho_*$  is also irreducible proving our claim.

Now, let  $\rho$  and  $\rho' = \gamma\rho\gamma^{-1}$  be two representatives of  $[\rho]$  and let  $u$  and  $u'$  be the two corresponding equivariant harmonic maps. It follows that  $u' = \gamma \cdot u$  where  $\cdot$  denotes the action of  $\operatorname{SL}(r, \mathbb{C})$  on  $\operatorname{SL}(r, \mathbb{C})/\operatorname{SU}(r)$ . It follows that the induced decompositions  $\mathcal{D} = d_A + \psi$  on the universal cover agree, hence after taking the quotients by  $\rho$  and  $\rho' = \gamma\rho\gamma^{-1}$  respectively the corresponding pairs are complex gauge equivalent by  $\gamma$ . Proposition 4.4 then shows that they are  $\mathcal{G}_{bal,\delta}$  gauge equivalent which completes the proof that  $\beta$  is well defined.

Next we will first show that  $\alpha(\beta([\rho])) = [\rho]$ . Let  $\beta([\rho]) = [(d_A, \psi)]$ . According to (2.9), we have  $d_A + \psi = \mathcal{D}$ , where  $\mathcal{D}$  is the connection on  $\operatorname{ad}(E_\rho)$  induced by the trivial connection on the universal cover which has holonomy  $\rho$ . Hence,  $\alpha(\beta([\rho])) = [\rho]$ .

Conversely,  $\beta(\alpha([(d_A, \psi)])) = [(d_A, \psi)]$ . Indeed, let  $(d_B, \phi)$  be a smooth representative of  $\beta(\alpha([(d_A, \psi)]))$ . By applying  $\alpha$  on both sides and what we just proved,  $\alpha([(d_A, \psi)]) = \alpha([(d_B, \phi)])$ . In other words,  $(d_A, \psi)$  and  $(d_B, \phi)$  have conjugate holonomies. Since the holonomies of these pairs near the vertices are trivial by Proposition 3.5, then Proposition 2.26 implies that the corresponding flat connections (and hence also the pairs) are complex gauge equivalent. Thus Proposition 4.4 implies that  $(d_A, \psi)$  and  $(d_B, \phi)$  are  $\mathcal{G}_{bal, \delta}$  gauge equivalent, hence  $\beta(\alpha([(d_A, \psi)])) = [(d_A, \psi)]$ .

In order to prove continuity, let  $\rho_i \rightarrow \rho \in \mathcal{M}_{char}^c$  and let  $u_i, u$  be the associated equivariant harmonic maps. Fix a compact fundamental domain  $F \subset \tilde{X}$  for the action of  $\Gamma$  and define  $\rho_i$  equivariant maps  $\tilde{u}_i$  by setting  $\tilde{u}_i = u$  on  $F$  and extending  $\rho_i$  equivariantly on  $\tilde{X}$ . Since the  $u_i$  are harmonic, the energy  $E^{u_i}$  satisfies

$$E^{u_i} \leq E^{\tilde{u}_i} = E^u.$$

The global Hölder bound (cf. Theorem 3.12 of [11]) implies that there is a subsequence (we call it again by  $\{i\}$  by a slight abuse of notation) such that  $u_i \rightarrow u_\infty$  uniformly on  $F$ . Furthermore, the convergence of the representations  $\rho_i \rightarrow \rho$  implies that  $u_\infty$  is  $\rho$ -equivariant and Theorem 5.1 of [11] implies that  $u_\infty$  is harmonic. Finally, the uniqueness Theorem 4.6 of [11] implies that  $u_\infty = u$ . We have thus shown so far

$$u_i \rightarrow u \text{ locally uniformly.}$$

Let  $(d_{A_i}, \psi_i)$  denote the unitary connection and Higgs field associated with the harmonic map  $u_i$ . By Theorem 3.3 together with the proof of Proposition 4.5 (in this we use the third derivative estimates) we obtain that the  $L_{2, \delta}^2$ -norm of  $(A_i, \psi_i)$  is uniformly bounded, and thus there exists a subsequence (we call it again by  $\{i\}$  by a slight abuse of notation) such that  $(d_{A_i}, \psi_i) \rightarrow (d_A, \psi)$  weakly in  $L_{2, \delta}^2$  and hence strongly in  $L_{1, \delta}^2$ .  $\square$

## REFERENCES

- [1] D. Arapura, A. Dimca and R. Hain *On the fundamental groups of normal varieties*. Preprint (2014).
- [2] R. Adams. *Sobolev spaces*. Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1975. Pure and Applied Mathematics, Vol. 65.
- [3] S. Agmon, A. Douglis, and L. Nirenberg. *Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions II*. Comm. Pure Appl. Math., 17:35–92, 1964.
- [4] M. Armstrong. *Basic Topology*. Springer-Verlag 1983.
- [5] V. Balaji, P. Barik and D. Nagaraj. *A degeneration of moduli of Hitchin pairs*. Electron. Res. Announc. Math. Sci. 20 (2013), 103–108.
- [6] J. Carlson and D. Toledo. *Harmonic mappings of Kähler manifolds to locally symmetric spaces*. Publ. Math. IHES, tome 69, 1989, 173–201.
- [7] J. Chen. *On energy minimizing mappings between and into singular spaces*. Duke Math. J. 79 (1995) 77–99.
- [8] K. Corlette. *Flat G-bundles with canonical metrics*. J. Differential Geom. 28 (3): 361–382, 1988.
- [9] K. Corlette. *Archimedean superrigidity and hyperbolic geometry*. Ann. of Math. 135 (1992) 165–182.
- [10] K. Corlette and C. Simpson. *On the classification of rank-two representations of quasiprojective fundamental groups*. Compos. Math. 144(5):1271–1331, 2008.
- [11] G. Daskalopoulos and C. Mese. *Harmonic maps from 2-complexes*. Comm. Anal. Geom., 14(3):497–549, 2006.
- [12] G. Daskalopoulos and C. Mese. *Harmonic maps from a cell complex and geometric rigidity*. J. Differential Geom. 78(2):269–293, 2008.
- [13] G. Daskalopoulos and C. Mese. *Fixed point and rigidity theorems for harmonic maps from simplicial complexes into NPC spaces*. Geom. Dedicata (2009) 141 33–57.
- [14] G. Daskalopoulos and C. Mese. *Restrictions of fundamental groups of normal projective varieties*. In preparation.

- [15] G. Daskalopoulos and R. Wentworth. *Geometric quantization for the moduli space of vector bundles with parabolic structure*. In *Geometry, topology and physics (Campinas, 1996)*, pages 119–155. de Gruyter, Berlin, 1997.
- [16] S. K. Donaldson. *Twisted harmonic maps and the self-duality equations*. *Proc. London Math. Soc.* (3), 55(1):127–131, 1987.
- [17] J. Eells and B. Fuglede. *Harmonic maps between Riemannian polyhedra*. Cambridge Tracts in Mathematics 142, Cambridge University Press, Cambridge 2001.
- [18] D. Gilbarg and N. Trudinger. *Elliptic partial differential equations of second order*, volume 224 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 1983.
- [19] P. Griffiths and J. Morgan. *Rational homotopy theory and differential forms*, volume 16 of *Progress in Mathematics*. Birkhäuser Boston, Mass., 1981.
- [20] M. Gromov and R. Schoen. *Harmonic maps into singular spaces and  $p$ -adic superrigidity for lattices in groups of rank one*. *Publ. Math. IHES* 76 (1992) 165–246.
- [21] N. Hitchin. *The self duality equations on a Riemann surface*. *Proc. London Math. Soc.* 55 (1987), 59–126.
- [22] P. Hartman. *Ordinary differential equations*. John Wiley & Sons Inc., New York, 1964.
- [23] J. Jost. *Nonpositive curvature: geometric and analytic aspects*. *Lectures in Mathematics*. ETH Zürich, Birkhäuser Verlag 1997.
- [24] J. Jost, Y-H. Yang and K. Zuo. *The cohomology of a variation of polarized Hodge structures over a quasi-compact Kähler manifold*. *J. Algebraic Geom.* 16(3), 401–434, 2007.
- [25] J. Jost and K. Zuo. *Harmonic maps and  $Sl(r, \mathbb{C})$ -representations of fundamental groups of quasi-projective manifolds*. *J. Algebraic Geom.* 5(1) 77–106, 1996.
- [26] M. Kapovich and J. Kollar. *Fundamental groups of links of isolated singularities*. *J. Amer. Math. Soc.* 27 (2014), 929–952.
- [27] S. Lojasiewich. *Triangulations of semi-analytic sets*. *Ann. Scuola Norm. Sup. Pisa* (3) 18 (1964) 449–474.
- [28] D. Kinderlehrer, L. Nirenberg, and J. Spruck. *Regularity in elliptic free boundary problems*. *J. Analyse Math.* 34:86–119 (1979), 1978.
- [29] R. Lockhart and R. McOwen. *Elliptic differential operators on noncompact manifolds*. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4) 12, no. 3, 409–447, 1985.
- [30] C. Mese. *Uniqueness theorems for harmonic maps into metric spaces*. *Commun. Contemp. Math.* 4(4): 725–750, 2002.
- [31] L. Simon. *Theorems on regularity and singularity of energy minimizing maps*. *Lectures in Mathematics ETH Zürich*. Birkhäuser Verlag, Basel, 1996. Based on lecture notes by Norbert Hungerbühler.
- [32] C. Simpson. *Constructing variations of Hodge structure using Yang-Mills theory and applications to uniformization*. *Journal of the AMS*. Volume 1, 1988, 867–918.
- [33] C. Simpson. *Harmonic bundles on noncompact curves*. *Journal of the AMS*. Volume 3, 1990, 713–770.
- [34] C. Simpson. *Higgs bundles and local systems*. *Publ. Math. IHES*, tome 75, 1992, 5–95.
- [35] Y.T. Siu. *The complex analyticity of harmonic maps and the strong rigidity of compact Kähler manifolds*. *Ann. of Math.* 112 (1980) 73–111.

DEPARTMENT OF MATHEMATICS, BROWN UNIVERSITY, PROVIDENCE, RI 02912, USA  
*E-mail address:* daskal@math.brown.edu

DEPARTMENT OF MATHEMATICS, JOHNS HOPKINS UNIVERSITY, BALTIMORE, MD 21218, USA  
*E-mail address:* cmese@math.jhu.edu

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE 119076  
*E-mail address:* graeme@nus.edu.sg