

This is a repository copy of *HOMOTOPY GROUPS OF MODULI SPACES OF STABLE QUIVER REPRESENTATIONS*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/149821/>

Version: Accepted Version

Article:

Wilkin, Graeme Peter Desmond (2010) HOMOTOPY GROUPS OF MODULI SPACES OF STABLE QUIVER REPRESENTATIONS. *International Journal of Mathematics*. p. 1219.

<https://doi.org/10.1142/S0129167X1000646X>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

HOMOTOPY GROUPS OF MODULI SPACES OF STABLE QUIVER REPRESENTATIONS

GRAEME WILKIN

ABSTRACT. The purpose of this paper is to describe a method for computing homotopy groups of the space of α -stable representations of a quiver with fixed dimension vector and stability parameter α . The main result is that the homotopy groups of this space are trivial up to a certain dimension, which depends on the quiver, the choice of dimension vector, and the choice of parameter. As a corollary we also compute low-dimensional homotopy groups of the moduli space of α -stable representations of the quiver with fixed dimension vector, and apply the theory to the space of non-degenerate polygons in three-dimensional Euclidean space.

1. INTRODUCTION

Symplectic quotients are important examples of symplectic manifolds, which arise in many areas of mathematics. When the quotient is a smooth manifold there exists a well-developed research program to compute the cohomology of these quotients via equivariant Morse theory, begun by Atiyah and Bott in [2], and Kirwan in [19]. When the quotient is singular then in many cases of interest one would like to compute topological invariants of the smooth locus of this space, for which new techniques are needed. In the case of holomorphic bundles over a compact Riemann surface, an approach due to Daskalopoulos and Uhlenbeck is to use transversality theory to compute low-dimensional homotopy groups of the moduli space of stable holomorphic bundles. In this paper we extend the Daskalopoulos-Uhlenbeck techniques to the space of stable representations of a quiver, and compute homotopy groups of these moduli spaces.

Let $\text{Rep}(Q, \mathbf{v})$ denote the space of representations of a quiver with fixed dimension vector \mathbf{v} , and let $G_{\mathbf{v}}$ be the associated group acting on $\text{Rep}(Q, \mathbf{v})$ (see Section 2 for the precise definition). For a given stability parameter $\alpha \in \mathfrak{g}_{\mathbf{v}}^*$, denote the subset of α -stable points by $\text{Rep}(Q, \mathbf{v})^{\alpha-st}$, and the α -semistable points by $\text{Rep}(Q, \mathbf{v})^{\alpha-ss}$. In a typical situation, for almost all choices of α we have $\text{Rep}(Q, \mathbf{v})^{\alpha-st} = \text{Rep}(Q, \mathbf{v})^{\alpha-ss}$, and since the isotropy group of each α -stable point is the subgroup consisting of scalar multiples of the identity, then there is a smooth structure on the quotient $\text{Rep}(Q, \mathbf{v})^{\alpha-ss} //_{\alpha} G_{\mathbf{v}}^{\mathbb{C}} = \text{Rep}(Q, \mathbf{v})^{\alpha-st} / G_{\mathbf{v}}^{\mathbb{C}}$.

Various methods have already been employed to compute topological invariants of the quotient $\text{Rep}(Q, \mathbf{v})^{\alpha-st} / G_{\mathbf{v}}^{\mathbb{C}}$ in the case where the spaces $\text{Rep}(Q, \mathbf{v})^{\alpha-ss}$ and $\text{Rep}(Q, \mathbf{v})^{\alpha-st}$ co-incide. The major result in this direction is that of Reineke in [27], who used an argument involving counting

Date: August 31, 2010.

2000 Mathematics Subject Classification. Primary: 58D15; Secondary: 14D20, 32G13.

Key words and phrases. Representations of quivers, Transversality, Moduli spaces of polygons.

rational points (analogous to the work of Harder and Narasimhan in [9] for the moduli space of stable vector bundles) to compute Betti numbers of the quotient. When Q is the star-shaped polygon quiver studied in Section 4.2, Klyachko in [21] computed Betti numbers of $\text{Rep}(Q, \mathbf{v})^{\alpha-st}/G_{\mathbf{v}}^{\mathbb{C}}$ for the case where $\text{Rep}(Q, \mathbf{v})^{\alpha-ss} = \text{Rep}(Q, \mathbf{v})^{\alpha-st}$, and Kapovich and Millson in [13] studied the geometry of $\text{Rep}(Q, \mathbf{v})^{\alpha-st}/G_{\mathbf{v}}^{\mathbb{C}}$.

More recently, the results of [8] show that the Morse theory methods of Kirwan in [19] apply to the non-compact space $\text{Rep}(Q, \mathbf{v})$, and as a consequence there is a surjective map in equivariant cohomology

$$\kappa : H_{G_{\mathbf{v}}}^*(\text{Rep}(Q, \mathbf{v})) \rightarrow H_{G_{\mathbf{v}}}^*(\text{Rep}(Q, \mathbf{v})^{\alpha-ss}).$$

When the spaces $\text{Rep}(Q, \mathbf{v})^{\alpha-ss}$ and $\text{Rep}(Q, \mathbf{v})^{\alpha-st}$ co-incide then there is an additional isomorphism

$$H_{G_{\mathbf{v}}}^*(\text{Rep}(Q, \mathbf{v})^{\alpha-ss}) = H_{G_{\mathbf{v}}}^*(\text{Rep}(Q, \mathbf{v})^{\alpha-st}) \cong H^*(\mathcal{M}_{\alpha}(Q, \mathbf{v})^{st}) \otimes H^*(BU(1)),$$

and from the Morse theory one can recover the Betti number calculations of [27].

In general, when the spaces $\text{Rep}(Q, \mathbf{v})^{\alpha-ss}$ and $\text{Rep}(Q, \mathbf{v})^{\alpha-st}$ do not co-incide then there are three different quotient varieties of interest: the GIT quotient $\mathcal{M}_{\alpha}(Q, \mathbf{v}) := \text{Rep}(Q, \mathbf{v})//_{\alpha}G^{\mathbb{C}}$, the resolution of singularities $\tilde{\mathcal{M}}_{\alpha}(Q, \mathbf{v})$ of the GIT quotient (as studied by Kirwan in [20]), and the smooth locus $\mathcal{M}_{\alpha}(Q, \mathbf{v})^{st} := \text{Rep}(Q, \mathbf{v})^{\alpha-st}/G^{\mathbb{C}} \subset \mathcal{M}_{\alpha}(Q, \mathbf{v})$. The object of study in this paper is the space $\mathcal{M}_{\alpha}(Q, \mathbf{v})^{st}$, and the results derived here on the topology of this space cannot be reproduced by the methods of [27] or [8] when $\text{Rep}(Q, \mathbf{v})^{\alpha-ss} \neq \text{Rep}(Q, \mathbf{v})^{\alpha-st}$.

The moduli space of semistable holomorphic bundles of degree zero over a compact Riemann surface is an example of a singular Kähler quotient for which one would like to study the smooth locus: the moduli space of stable bundles. In the rank 2 case Kirwan applied her general theory for computing intersection Betti numbers of GIT quotients to obtain results about the low-dimensional Betti numbers (see [16] and Remark 3.4 in [17]), and later Daskalopoulos in [5] used infinite-dimensional methods in the spirit of Atiyah and Bott to obtain results about the homotopy groups of this space. Following this, Daskalopoulos and Uhlenbeck in [6] used an infinite-dimensional transversality argument to extend these results to the moduli space of stable bundles for any rank and degree, and obtained a simple formula for the homotopy groups of the moduli space of stable bundles up to a dimension bound depending on the rank of the bundle and the genus of the surface.

In this paper we extend the Daskalopoulos and Uhlenbeck approach to the moduli space of α -stable representations of a quiver with fixed dimension vector, as introduced by King in [14] (see Section 2 for more information). The difference between the approach of this paper and Kirwan's approach of [16] is twofold: (a) Kirwan's results do not apply *a priori* to the non-compact spaces $\text{Rep}(Q, \mathbf{v})$ under consideration here, one first needs to prove a gradient flow convergence result for the norm-square of the moment map on $\text{Rep}(Q, \mathbf{v})$ (which was done in [8]) as well as on the blow-ups of $\text{Rep}(Q, \mathbf{v})$ required to desingularise the quotient (the proof of which has not yet been carried out), and (b) in this paper the dimensional bound d_{min} can be computed directly by studying the

different possible destabilising subrepresentations, whereas in Kirwan's method one needs to also study the desingularised spaces, the calculations for which become much more complicated as the dimension vector \mathbf{v} increases in size. Of course, it should be mentioned that Kirwan's methods do apply to a general class of quotients of compact Kähler manifolds and appear as a corollary of her results on the intersection cohomology of Kähler quotients, while the scope of this paper is limited to computing homotopy groups of Kähler quotients of $\text{Rep}(Q, \mathbf{v})$.

The main result of this paper is a computation of the low-dimensional homotopy groups of the spaces $\text{Rep}(Q, \mathbf{v})^{\alpha-st}$ and $\mathcal{M}_\alpha(Q, \mathbf{v})^{st}$, up to a dimension that depends on the underlying quiver, the choice of dimension vector, and the stability parameter α . The following is the main theorem.

Theorem 1.1. *[Theorem 3.10] Let Q be a quiver, $\text{Rep}(Q, \mathbf{v})$ be the space of representations with fixed dimension vector \mathbf{v} , let $\alpha \in \mathfrak{g}_{\mathbf{v}}^*$ be a stability parameter, and let $d_{\min}(Q, \mathbf{v}, \alpha)$ be the associated minimal dimension as defined in Definition 3.1. Then $\pi_n(\text{Rep}(Q, \mathbf{v})^{\alpha-st}) = 0$ for all $n < d_{\min}(Q, \mathbf{v}, \alpha) - 1$.*

Corollary 1.2. *[Corollary 3.11] Let $PG_{\mathbf{v}} = G_{\mathbf{v}}/U(1)$ be the quotient of $G_{\mathbf{v}}$ by the subgroup consisting of scalar multiples of the identity. Then $\pi_n(\mathcal{M}_\alpha(Q, \mathbf{v})^{st}) \cong \pi_{n-1}(PG_{\mathbf{v}})$ for all $n < d_{\min}(Q, \mathbf{v}, \alpha) - 1$.*

As an application of the theory, some examples of interest are explicitly worked out in Section 4. The first example is a space that contains the moduli space of framed instantons on S^4 , via the ADHM construction of [3], and while the ADHM moduli space itself (which is a hyperkähler quotient, or alternately a Kähler quotient of a singular space) is beyond the scope of this paper, one can view these results as a first step towards studying the topology of this space. Again, it should be mentioned that in [18], Kirwan takes a different approach to the problem of studying the ADHM moduli space, by first desingularising the hyperkähler quotient of the space of semistable points, and then using general results (first derived in [15]) to describe the effect on low-dimensional homotopy groups of removing the strata corresponding to the singular points in the original moduli space.

The second example is the space of *non-degenerate* polygons in \mathbb{R}^3 (i.e. those polygons that do not lie in a line) modulo rigid motions of \mathbb{R}^3 . In Section 4.2 we compute the homotopy groups of the moduli space of non-degenerate polygons up to dimension 2 under certain conditions on the stability parameter.

Ultimately one would like to apply these methods to hyperkähler quotients, in particular the Nakajima quiver varieties from [26], and the moduli space of stable Higgs bundles of rank n and degree d over a compact Riemann surface from [12]. These can be viewed in two different ways: As a hyperkähler quotient $\mu_1^{-1}(0) \cap \mu_2^{-1}(0) \cap \mu_3^{-1}(0)/G$, or as a GIT quotient of a singular space $(\mu_2^{-1}(0) \cap \mu_3^{-1}(0)) // G^{\mathbb{C}}$. In the first case there is no stability condition that implies existence of common zeros of the three moment maps μ_1, μ_2, μ_3 , and in the second case the space $\mu_2^{-1}(0) \cap \mu_3^{-1}(0)$ is singular, and so *a priori* the methods of this paper do not apply to this situation. It would be a

very interesting problem to extend these techniques to stratified singular spaces in order to study hyperkähler quotients such as those described above.

This paper is organised as follows: Section 2 contains the background material and notational conventions used in this paper, Section 3 contains the proof of the main theorem, and the explicit examples are worked out in Section 4.

2. BACKGROUND AND NOTATION

A *quiver* Q is a directed graph, with vertices \mathcal{J} , edges \mathcal{E} and head/tail maps $h, t : \mathcal{J} \rightarrow \mathcal{E}$ that define the orientation of the edges. A *complex representation* of a quiver consists of a choice of complex vector spaces $\{V_j\}_{j \in \mathcal{J}}$ with a fixed Hermitian structure, and homomorphisms $\{A_a : V_{t(a)} \rightarrow V_{h(a)}\}_{a \in \mathcal{E}}$. The *dimension vector* of a representation is the vector $\mathbf{v} = (\dim V_j)_{j \in \mathcal{J}}$. The space of representations with fixed dimension vector is denoted

$$(2.1) \quad \text{Rep}(Q, \mathbf{v}) = \bigoplus_{a \in \mathcal{E}} \text{Hom}(V_{t(a)}, V_{h(a)}),$$

and the vector space consisting of the direct sum of all the vector spaces at each vertex is denoted

$$\text{Vect}(Q, \mathbf{v}) = \bigoplus_{j \in \mathcal{J}} V_j.$$

A *subrepresentation* of a representation $A \in \text{Rep}(Q, \mathbf{v})$ consists of vector spaces $\{V'_j \subseteq V_j\}_{j \in \mathcal{J}}$ such that $A_a(V'_{t(a)}) \subseteq V'_{h(a)}$ for all $a \in \mathcal{E}$, and homomorphisms $\{A'_a : V'_{t(a)} \rightarrow V'_{h(a)}\}_{a \in \mathcal{E}}$ such that each A'_a is the restriction of A_a to $V'_{t(a)}$.

Given such a subrepresentation, for each vertex $j \in \mathcal{J}$ there is an associated projection $\pi_j : V_j \rightarrow V'_j$ that is defined using orthogonal projection with respect to the Hermitian structure on V_j . If the dimension vector of the subrepresentation is $\mathbf{v}' = (\dim V'_j)_{j \in \mathcal{J}}$ then we write $\mathbf{v}' \leq \mathbf{v}$, the projection $\pi : \text{Vect}(Q, \mathbf{v}) \rightarrow \text{Vect}(Q, \mathbf{v}')$ is used to denote the vector spaces in the subrepresentation, and $A' \in \text{Rep}(Q, \mathbf{v}')$ is used to denote the induced homomorphisms $A'_a = \pi \circ A_a \circ \pi \in \text{Hom}(V'_{t(a)}, V'_{h(a)})$.

Given a fixed Hermitian structure on each vector space V_j , the group $G_{\mathbf{v}} = \times_{j \in \mathcal{J}} \mathbf{U}(V_j)$ acts on $\text{Rep}(Q, \mathbf{v})$ by

$$(2.2) \quad g \cdot \{A_a\}_{a \in \mathcal{E}} = \{g_{h(a)} A_a g_{t(a)}^{-1}\}_{a \in \mathcal{E}},$$

where $g = (g_j)_{j \in \mathcal{J}} \in \times_{j \in \mathcal{J}} \mathbf{U}(V_j)$. Let $\mathfrak{g}_{\mathbf{v}}$ denote the Lie algebra of $G_{\mathbf{v}}$. The Hermitian structure on each vector space V_j gives an identification

$$\mathfrak{g}_{\mathbf{v}} = \bigoplus_{j \in \mathcal{J}} \mathfrak{u}(V_j) \cong \bigoplus_{j \in \mathcal{J}} \mathfrak{u}(V_j)^* = \mathfrak{g}_{\mathbf{v}}^*,$$

and with respect to this Hermitian structure we define the adjoint A_a^* of each homomorphism $A_a : V_{t(a)} \rightarrow V_{h(a)}$.

The action of $G_{\mathbf{v}}$ preserves the symplectic structure (in fact the Kähler structure) on $\text{Rep}(Q, \mathbf{v})$ induced by the Hermitian structure on each V_j , and a moment map for this action is a map

$\mu : \text{Rep}(Q, \mathbf{v}) \rightarrow \mathfrak{g}_{\mathbf{v}}^* = \bigoplus_{j \in \mathcal{J}} \mathfrak{u}(V_j)^*$, where the component corresponding to $\mathfrak{u}(V_j)^*$ is

$$(2.3) \quad \mu_j(A) = \sqrt{-1} \sum_{\substack{a \in \mathcal{E} \\ h(a)=j}} A_a A_a^* - \sqrt{-1} \sum_{\substack{a \in \mathcal{E} \\ t(a)=j}} A_a^* A_a.$$

For convenience, in the rest of the paper the notation

$$(2.4) \quad \mu(A) = \sqrt{-1} \sum_{a \in \mathcal{E}} [A_a, A_a^*] \in \mathfrak{g}_{\mathbf{v}}^* \subset \text{End}(\text{Vect}(Q, \mathbf{v}))$$

is used, where each $A_a \in \text{Hom}(V_{t(a)}, V_{h(a)}) \subseteq \text{End}(\text{Vect}(Q, \mathbf{v}))$, and the commutator of two elements $X, Y \in \text{End}(\text{Vect}(Q, \mathbf{v}))$ is denoted $[X, Y]$.

A central element α of $\mathfrak{g}_{\mathbf{v}}^*$ consists of a choice of a real number α_j at each vertex $j \in \mathcal{J}$, denoted

$$\alpha = (\sqrt{-1} \alpha_j \text{id}_{V_j})_{j \in \mathcal{J}}.$$

Corollary 6.2 of [14] identifies the Kähler quotient

$$(2.5) \quad \mathcal{M}_{\alpha}(Q, \mathbf{v}) := \mu^{-1}(\alpha) / G_{\mathbf{v}}$$

with the *moduli space of α -semistable representations*, $\text{Rep}(Q, \mathbf{v}) //_{\alpha} G_{\mathbf{v}}^{\mathbb{C}}$, defined using GIT (see below for more details). This space depends on the quiver Q , the dimension vector \mathbf{v} , and the choice of $\alpha \in \mathfrak{g}_{\mathbf{v}}^*$, and note that since $\mu(A)$ consists of commutators then the quotient is empty if $\sum_{j \in \mathcal{J}} \alpha_j \dim_{\mathbb{C}} V_j \neq 0$.

The construction of King in [14] shows that when α is integral then this quotient can be described via Geometric Invariant Theory as follows. Let $\alpha = (\sqrt{-1} \alpha_j \text{id}_{V_j})_{j \in \mathcal{J}}$, with each $\alpha_j \in \mathbb{Z}$, and let $G_{\mathbf{v}}^{\mathbb{C}} = \times_{j \in \mathcal{J}} \text{GL}(V_j, \mathbb{C})$ denote the complexification of G . The action of the group $G_{\mathbf{v}}^{\mathbb{C}}$ lifts to an action on the trivial line bundle over $\text{Rep}(Q, \mathbf{v})$ by

$$g \cdot (\{A_a\}_{a \in \mathcal{E}}, \xi) = \left(\left\{ g_{h(a)} A_a g_{t(a)}^{-1} \right\}, \chi(g) \xi \right), \text{ for } \xi \in \mathbb{C},$$

where the character χ is given by

$$\chi(g) = \prod_{j \in \mathcal{J}} \det(g_j)^{-\alpha_j}.$$

A representation $A \in \text{Rep}(Q, \mathbf{v})$ is *α -semistable* if (given some nonzero $\xi \in \mathbb{C}$) the closure of the $G_{\mathbf{v}}^{\mathbb{C}}$ -orbit of (A, ξ) in $\text{Rep}(Q, \mathbf{v}) \times \mathbb{C}$ does not intersect the zero section, i.e.

$$\overline{G_{\mathbf{v}}^{\mathbb{C}} \cdot (A, \xi)} \cap (\text{Rep}(Q, \mathbf{v}) \times \{0\}) = \emptyset.$$

A representation $A \in \text{Rep}(Q, \mathbf{v})$ is *α -stable* if (given some nonzero $\xi \in \mathbb{C}$) the $G_{\mathbf{v}}^{\mathbb{C}}$ -orbit of (A, ξ) in $\text{Rep}(Q, \mathbf{v}) \times \mathbb{C}$ is closed, and, in addition, the isotropy group of A is the subgroup of scalar multiples of the identity in $G_{\mathbf{v}}^{\mathbb{C}}$.

Proposition 3.1 in [14] shows that the definitions given above are equivalent to the following slope-stability conditions (analogous to those for holomorphic bundles over a Kähler manifold).

Given a subrepresentation with dimension vector $\mathbf{v}' = (v_j)_{j \in \mathcal{J}}$, the *rank* and α -*degree* of the subrepresentation are given by

$$(2.6) \quad \text{rank}(Q, \mathbf{v}') = \sum_{j \in \mathcal{J}} v_j$$

$$(2.7) \quad \text{deg}_\alpha(Q, \mathbf{v}') = \sum_{j \in \mathcal{J}} -\alpha_j v_j$$

The α -*slope* is then defined to be the quotient $\text{slope}_\alpha(Q, \mathbf{v}') = \text{deg}_\alpha(Q, \mathbf{v}') / \text{rank}(Q, \mathbf{v}')$.

Lemma 2.1 (Proposition 3.1 of [14]). *Let Q be a quiver, \mathbf{v} a dimension vector and α a stability parameter satisfying $\text{deg}_\alpha(Q, \mathbf{v}) = 0$. A representation $A \in \text{Rep}(Q, \mathbf{v})$ is α -stable (resp. α -semistable) if and only if every subrepresentation satisfies*

$$(2.8) \quad \text{slope}_\alpha(Q, \mathbf{v}') < \text{slope}_\alpha(Q, \mathbf{v}) \quad (\text{resp. } \text{slope}_\alpha(Q, \mathbf{v}') \leq \text{slope}_\alpha(Q, \mathbf{v})).$$

The space of α -stable (resp. α -semistable) representations is denoted $\text{Rep}(Q, \mathbf{v})^{\alpha-st}$ (resp. $\text{Rep}(Q, \mathbf{v})^{\alpha-ss}$), and the GIT quotient is defined to be

$$(2.9) \quad \mathcal{M}_\alpha(Q, \mathbf{v}) = \text{Rep}(Q, \mathbf{v}) //_\alpha G_{\mathbf{v}}^{\mathbb{C}} := \text{Rep}(Q, \mathbf{v})^{\alpha-ss} // G_{\mathbf{v}}^{\mathbb{C}},$$

where the quotient $//$ identifies orbits whose closures intersect.

It follows from Theorem 6.1 of [14] that the Kähler quotient defined in (2.5) can be identified with the GIT quotient defined in (2.9) (see also [8] for an alternative proof using the gradient flow of $\|\mu - \alpha\|^2$).

Reineke in [27] computes Betti numbers of $\mathcal{M}_\alpha(Q, \mathbf{v})^{st}$ for choices of parameter and dimension vector for which the sets $\text{Rep}(Q, \mathbf{v})^{\alpha-st}$ and $\text{Rep}(Q, \mathbf{v})^{\alpha-ss}$ co-incide. Following on from this, the results of [8] show that there is a surjective map (known as the *Kirwan map*) $\kappa : H_{G_{\mathbf{v}}}^*(\text{Rep}(Q, \mathbf{v})) \rightarrow H_{G_{\mathbf{v}}}^*(\text{Rep}(Q, \mathbf{v})^{\alpha-ss})$ and that the Betti numbers can also be computed using the equivariant Morse theory of Atiyah and Bott [2] and Kirwan [19]. The Morse theory only gives information about the topology of $\text{Rep}(Q, \mathbf{v})^{\alpha-ss}$ and the purpose of this paper is to understand the low-dimensional topology of $\mathcal{M}_\alpha(Q, \mathbf{v})^{\alpha-st}$ regardless of whether $\text{Rep}(Q, \mathbf{v})^{\alpha-st}$ and $\text{Rep}(Q, \mathbf{v})^{\alpha-ss}$ co-incide.

If the sets $\text{Rep}(Q, \mathbf{v})^{\alpha-st}$ and $\text{Rep}(Q, \mathbf{v})^{\alpha-ss}$ do not co-incide then the quotient may have serious singularities (see for example [20]). In these cases there is a smooth structure on the quotient of the subset of stable representations

$$(2.10) \quad \mathcal{M}_\alpha(Q, \mathbf{v})^{st} = \text{Rep}(Q, \mathbf{v})^{\alpha-st} / G_{\mathbf{v}}^{\mathbb{C}}.$$

The situation in this paper is an extension to the case of quivers of the infinite-dimensional methods for studying the moduli space of stable holomorphic bundles over a compact Riemann surface used in [5] and [6]. For rank 2 bundles Daskalopoulos used an algebraic stratification of the space of semistable bundles to obtain results about the low-dimensional topology of the space of stable bundles (see Section 7 of [5] for more details). As noted on pp744-745 of [5], for higher rank bundles a stratification argument quickly becomes intractable, and in [6] a more general approach

using transversality is adopted. In this paper we follow the technique of [6], and use a transversality argument to obtain the results of Theorem 3.10.

Given a representation $A \in \text{Rep}(Q, \mathbf{v})$, the *infinitesimal action* of $G_{\mathbf{v}}^{\mathbb{C}}$ at A is given by

$$(2.11) \quad \begin{aligned} \rho_A : \mathfrak{g}_{\mathbf{v}}^{\mathbb{C}} &\rightarrow T_A \text{Rep}(Q, \mathbf{v}) \\ u &\mapsto \left. \frac{d}{dt} \right|_{t=0} e^{tu} \cdot A = ([u, A_a])_{a \in \mathcal{E}}. \end{aligned}$$

Given $A \in \text{Rep}(Q, \mathbf{v})^{\alpha-st}$, the infinitesimal deformations of the moduli space $\mathcal{M}_{\alpha}(Q, \mathbf{v})^{st}$ at the equivalence class containing A can be viewed in terms of the cohomology of a complex

$$(2.12) \quad \mathfrak{g}_{\mathbf{v}}^{\mathbb{C}} \xrightarrow{\rho_A} T_A \text{Rep}(Q, \mathbf{v}).$$

Remark 2.2. This complex can be interpreted in terms of the cohomology of the quiver representation V_A associated to A , via the following exact sequence, which first appeared in Section 2.1 of [28] (I thank the referee for this reference)

$$(2.13) \quad 0 \rightarrow \text{Hom}(V_A, V_A) \cong \ker \rho_A \rightarrow \mathfrak{g}_{\mathbb{C}} \xrightarrow{\rho_A} T_A \text{Rep}(Q, \mathbf{v}) \rightarrow \text{coker } \rho_A \cong \text{Ext}^1(V_A, V_A) \rightarrow 0.$$

The tangent space to the moduli space is isomorphic to $T_A \text{Rep}(Q, \mathbf{v}) / \text{im } \rho_A = \text{coker } \rho_A \cong \text{Ext}^1(V_A, V_A)$. Since an α -stable point has stabiliser consisting of scalar multiples of the identity in $G_{\mathbf{v}}^{\mathbb{C}}$, then $\dim_{\mathbb{C}} \ker \rho_A = 1$, and so

$$\begin{aligned} \dim_{\mathbb{C}} \text{coker } \rho_A - 1 &= \dim_{\mathbb{C}} \text{coker } \rho_A - \dim_{\mathbb{C}} \ker \rho_A \\ &= \dim_{\mathbb{C}} T_A \text{Rep}(Q, \mathbf{v}) - \dim_{\mathbb{C}} \mathfrak{g}_{\mathbf{v}}^{\mathbb{C}} \\ \Leftrightarrow \dim_{\mathbb{C}} \text{coker } \rho_A &= \dim_{\mathbb{C}} T_A \text{Rep}(Q, \mathbf{v}) - \dim_{\mathbb{C}} \mathfrak{g}_{\mathbf{v}}^{\mathbb{C}} + 1. \end{aligned}$$

When $A \in \text{Rep}(Q, \mathbf{v})$ is α -unstable, then one can still define the complex (2.12), however $\ker \rho_A$ may now be non-trivial. A particular case of interest is when A preserves the vector spaces $\text{Vect}(Q, \mathbf{v}')$ of a given subrepresentation with associated orthogonal projection π . In this case one can view π as an endomorphism $\pi \in \mathfrak{g}_{\mathbf{v}}^{\mathbb{C}} = \bigoplus_{j \in \mathcal{J}} \text{End}(V_j)$, and $\rho_A(\pi) = ([A_a, \pi])_{a \in \mathcal{E}}$. If $\pi : \text{Vect}(Q, \mathbf{v}) \rightarrow \text{Vect}(Q, \mathbf{v}')$ is a subrepresentation of A (i.e. $A_a(\pi(V_{t(a)})) \subseteq \pi(V_{h(a)})$ for each $a \in \mathcal{E}$), then $\pi \circ A_a \circ \pi = A_a \circ \pi = A_a \circ \pi \circ \pi$, or alternately $[A_a, \pi] \circ \pi = 0$ for each $a \in \mathcal{E}$. Conversely, if $[A_a, \pi] \circ \pi = 0$ for each $a \in \mathcal{E}$, then A_a preserves the vector spaces in the image of the projection π . Therefore we have proved the following lemma.

Lemma 2.3. $\pi : \text{Vect}(Q, \mathbf{v}) \rightarrow \text{Vect}(Q, \mathbf{v}')$ is a subrepresentation of $A \in \text{Rep}(Q, \mathbf{v})$ iff $\rho_A(\pi) \circ \pi = 0$.

Given a subrepresentation $\pi : \text{Vect}(Q, \mathbf{v}) \rightarrow \text{Vect}(Q, \mathbf{v}')$ and the orthogonal complement $\pi^{\perp} := (\text{id} - \pi) : \text{Vect}(Q, \mathbf{v}) \rightarrow \text{Vect}(Q, \mathbf{v}')^{\perp}$ we define the following subspaces of $\mathfrak{g}_{\mathbf{v}}^{\mathbb{C}}$ and $T_A \text{Rep}(Q, \mathbf{v})$.

$$\begin{aligned} \text{Hom}^0(\pi, \pi^{\perp}) &:= \bigoplus_{j \in \mathcal{J}} \text{Hom}(\pi(V_j), \pi^{\perp}(V_j)) \subseteq \mathfrak{g}_{\mathbf{v}}^{\mathbb{C}} \\ \text{Hom}^1(\pi, \pi^{\perp}) &:= \bigoplus_{a \in \mathcal{E}} \text{Hom}(\pi(V_{t(a)}), \pi^{\perp}(V_{h(a)})) \subseteq \text{Rep}(Q, \mathbf{v}) \end{aligned}$$

Moreover, if A preserves π then $\rho_A : \mathfrak{g}_{\mathbb{C}} \rightarrow T_A \text{Rep}(Q, \mathbf{v})$ induces a map

$$(2.14) \quad \text{Hom}^0(\pi, \pi^\perp) \xrightarrow{\rho_\pi} \text{Hom}^1(\pi, \pi^\perp),$$

where we identify $\text{Hom}^1(\pi, \pi^\perp) \subseteq \text{Rep}(Q, \mathbf{v})$ with a quotient of the tangent space $T_A \text{Rep}(Q, \mathbf{v})$. If A preserves both π and π^\perp , then we say that the representation *splits*. This viewpoint will become useful in the next section, where we use this to define a simple condition for a representation to be unstable.

Remark 2.4. The complex (2.14) also has a cohomological interpretation similar to that in Remark 2.2 via the following complex

$$(2.15) \quad 0 \rightarrow \text{Hom}(V_\pi, V/V_\pi) \rightarrow \text{Hom}^0(\pi, \pi^\perp) \xrightarrow{\rho_\pi} \text{Hom}^1(\pi, \pi^\perp) \rightarrow \text{Ext}(V_\pi, V/V_\pi) \rightarrow 0,$$

where $\text{Hom}(V_\pi, V/V_\pi) \cong \ker \rho_\pi$ and $\text{Ext}(V_\pi, V/V_\pi) \cong \text{coker } \rho_\pi$. The Euler characteristic of the subrepresentation (defined below) is then fixed independently of the choice of representations V_π and V/V_π

$$(2.16) \quad \begin{aligned} \chi(V_\pi, V/V_\pi) &:= \dim_{\mathbb{C}} \text{Hom}(V_\pi, V/V_\pi) - \dim_{\mathbb{C}} \text{Ext}^1(V_\pi, V/V_\pi) \\ &= \dim_{\mathbb{C}} \text{Hom}^0(\pi, \pi^\perp) - \dim_{\mathbb{C}} \text{Hom}^1(\pi, \pi^\perp). \end{aligned}$$

When the representation splits, then $V/V_\pi \cong V_{\pi^\perp}$, and so we denote

$$\chi(\pi, \pi^\perp) := \chi(V_\pi, V_{\pi^\perp}).$$

3. TRANSVERSALITY RESULTS

The purpose of this section is to prove Theorem 3.10, which shows that the homotopy groups of the space $\text{Rep}(Q, \mathbf{v})^{\alpha-st}$ are trivial up to a dimensional bound that depends on the quiver Q , the dimension vector \mathbf{v} and the parameter α . The key idea behind this is to show that for small values of n there exists a based homotopy equivalence F between a given map $s : S^n \rightarrow \text{Rep}(Q, \mathbf{v})^{\alpha-st}$ and the constant map, such that the image of F is contained in $\text{Rep}(Q, \mathbf{v})^{\alpha-st}$. Equivalently, we show that the image of F does not contain any representations that preserve a sub-representation violating the slope-stability condition of Lemma 2.1.

Given a dimension vector $\mathbf{v}' \leq \mathbf{v}$, the set of all possible subrepresentations $\pi : \text{Vect}(Q, \mathbf{v}) \rightarrow \text{Vect}(Q, \mathbf{v}')$ is given by

$$(3.1) \quad \text{Gr}(\mathbf{v}', \mathbf{v}) := \prod_{j \in \mathcal{J}} \text{Gr}(v'_j, v_j).$$

For $\pi \in \text{Gr}(\mathbf{v}', \mathbf{v})$, the value of $\dim_{\mathbb{R}} \text{Hom}^1(\pi, \pi^\perp) - \dim_{\mathbb{R}} \text{Hom}^0(\pi, \pi^\perp)$ depends only on the dimension vector \mathbf{v}' .

Definition 3.1. The *minimal dimension* of the triple (Q, \mathbf{v}, α) is

$$(3.2) \quad \begin{aligned} d_{\min}(Q, \mathbf{v}, \alpha) &:= \min_{\substack{\mathbf{v}' < \mathbf{v} \\ \text{slope}_\alpha(Q, \mathbf{v}') \geq \text{slope}_\alpha(Q, \mathbf{v})}} \dim_{\mathbb{R}} \text{Hom}^1(\pi, \pi^\perp) - \dim_{\mathbb{R}} \text{Hom}^0(\pi, \pi^\perp) \\ &= \min_{\substack{\mathbf{v}' < \mathbf{v} \\ \text{slope}_\alpha(Q, \mathbf{v}') \geq \text{slope}_\alpha(Q, \mathbf{v})}} (-2)\chi(\pi, \pi^\perp). \end{aligned}$$

Remark 3.2. Theorem 3.10 and Corollary 3.11 show that the value of $d_{\min}(Q, \mathbf{v}, \alpha) - 2$ gives an upper bound on the dimension of the homotopy groups of $\mathcal{M}(Q, \mathbf{v})^{\alpha-st}$ that can be computed using the methods of this paper. In Section 4 we compute $d_{\min}(Q, \mathbf{v}, \alpha)$ explicitly for each example studied, and it would be an extremely interesting problem to provide some general combinatorial bounds for $d_{\min}(Q, \mathbf{v}, \alpha)$ in terms of the quiver and dimension vector. Results of Schofield in [29] provide a starting point in this direction, as they give some conditions for which (in the notation of this paper) $\text{Ext}(V_\pi, V_{\pi^\perp})$ vanishes generally, which relates to d_{\min} via (2.16).

Let $T \rightarrow \text{Gr}(\mathbf{v}', \mathbf{v})$ denote the tautological bundle with fibre over $\pi \in \text{Gr}(\mathbf{v}', \mathbf{v})$ given by $\text{Vect}(Q, \mathbf{v}')$, a sub-bundle of the trivial $\text{Vect}(Q, \mathbf{v})$ bundle over $\text{Gr}(\mathbf{v}', \mathbf{v})$. Similarly, let $W^0 \rightarrow \text{Gr}(\mathbf{v}', \mathbf{v})$ be the tautological sub-bundle of the trivial bundle $\text{Gr}(\mathbf{v}', \mathbf{v}) \times \mathfrak{g}_{\mathbf{v}}^{\mathbb{C}}$, where the fibres of W^0 are the subspaces $\text{Hom}^0(\pi, \pi^\perp) \subseteq \mathfrak{g}_{\mathbf{v}}^{\mathbb{C}}$, and let $W^1 \rightarrow \text{Gr}(\mathbf{v}', \mathbf{v})$ be the tautological sub-bundle of the trivial bundle $\text{Gr}(\mathbf{v}', \mathbf{v}) \times \text{Rep}(Q, \mathbf{v})$, where the fibres of W^1 are the subspaces $\text{Hom}^1(\pi, \pi^\perp) \subseteq \text{Rep}(Q, \mathbf{v})$.

Given a compact manifold X , and basepoints $x_0 \in X$ and $A_0 \in \text{Rep}(Q, \mathbf{v})$, let $C^q(X, \text{Rep}(Q, \mathbf{v}))$ be the set of C^q maps from X to $\text{Rep}(Q, \mathbf{v})$, this has the structure of a C^q manifold. Let $\mathcal{C}(X) \subset C^q(X, \text{Rep}(Q, \mathbf{v}))$ be the subset of basepoint-preserving C^q maps from X to $\text{Rep}(Q, \mathbf{v})$.

Lemma 3.3. $\mathcal{C}(X)$ is a submanifold of $C^q(X, \text{Rep}(Q, \mathbf{v}))$, and the tangent space at a point $s \in \mathcal{C}(X)$ is

$$(3.3) \quad T_s \mathcal{C}(X) = \{\delta s : X \rightarrow \text{Rep}(Q, \mathbf{v}) : \delta s \in C^q, \delta s(x_0) = 0\}.$$

Proof. Define $L : C^q(X, \text{Rep}(Q, \mathbf{v})) \rightarrow \text{Rep}(Q, \mathbf{v})$ by $L(s) = s(x_0)$. Then $\mathcal{C}(X) = L^{-1}(A_0)$, and so by the implicit function theorem, the proof reduces to showing that A_0 is a regular value of L . The derivative of L at $s \in \mathcal{C}(X)$ is $\delta L_s(\delta s) = \delta s(x_0)$, and the Whitney extension theorem (cf. Appendix A in [1]) shows that given a prescribed derivative δs at x_0 there exists $s \in C^q(X, \text{Rep}(Q, \mathbf{v}))$ with $s(x_0) = A_0$ and the derivative of s at x_0 equal to δs . Therefore δL is surjective and A_0 is a regular value of L . \square

The space of basepoint-preserving homotopy equivalences between maps in $\mathcal{C}(X)$ is defined as follows. Let $\mathcal{S}(X) \subset C^q(X \times [0, 1], \text{Rep}(Q, \mathbf{v}))$ be the subset of maps $F : X \times [0, 1] \rightarrow \text{Rep}(Q, \mathbf{v})$ such that $F(x_0, t) = A_0$ for all $t \in [0, 1]$.

Lemma 3.4. The space $\mathcal{S}(X)$ has the structure of a C^q manifold, and the tangent space at $F \in \mathcal{S}(X)$ is

$$(3.4) \quad T_F \mathcal{S}(X) = \{\delta F : X \times [0, 1] \rightarrow \text{Rep}(Q, \mathbf{v}) : \delta F \in C^q, \delta F(x_0, t) = 0 \forall t \in [0, 1]\}.$$

The proof of this statement is similar to that of the previous lemma (which uses the implicit function theorem and Whitney extension theorem), and so it is omitted.

Given $s \in \mathcal{C}(X)$, the space of basepoint-preserving homotopy equivalences between s and the constant map is

$$(3.5) \quad \mathcal{S}(s) = \{F \in \mathcal{S}(X) : F(x, 0) = s(x), F(x, 1) = A_0, F(x_0, t) = A_0\}.$$

Since $\text{Rep}(Q, \mathbf{v})$ is a vector space then $\mathcal{S}(s)$ is clearly non-empty.

Lemma 3.5. $\mathcal{S}(s)$ is a C^q submanifold of $\mathcal{S}(X)$, and the tangent space of $\mathcal{S}(s)$ at F is

$$(3.6) \quad T_F \mathcal{S}(s) = \{\delta F : X \times [0, 1] \rightarrow \text{Rep}(Q, \mathbf{v}) : \\ \delta F \in C^q, \delta F(x, 0) = \delta F(x, 1) = \delta F(x_0, t) = 0, \forall t \in [0, 1], x \in X\}.$$

The proof of this lemma is similar to that of the previous lemma (see also the proof of Lemma 2.1 in [6]), and so it is omitted.

Now fix $X = S^n$ and $s : X \rightarrow \text{Rep}(Q, \mathbf{v})^{\alpha-st}$. For notation let $\mathcal{C} = \mathcal{C}(X)$, $\mathcal{S} = \mathcal{S}(s)$ and $Y = S^n \times [0, 1]$. Given a dimension vector $\mathbf{v}' \leq \mathbf{v}$, define the map

$$\mathcal{D} : \text{Gr}(\mathbf{v}', \mathbf{v}) \times \mathcal{S} \times Y \rightarrow W^1 \subseteq \text{Gr}(\mathbf{v}', \mathbf{v}) \times \text{Rep}(Q, \mathbf{v}) \\ \mathcal{D}(\pi, F, y) = \left(\pi, \pi^\perp \circ \rho_{F(y)}(\pi) \circ \pi \right)$$

Since the component of $\rho_{F(y)}(\pi)$ along each edge $a \in \mathcal{E}$ is given by $[F(y)_a, \pi]$, then for simplicity of notation we write $\rho_{F(y)}(\pi) = [F(y), \pi]$, where the meaning is that the component of $[F(y), \pi]$ corresponding to the edge $a \in \mathcal{E}$ is $[F(y)_a, \pi]$. For a given map $F \in \mathcal{S}(s)$, define $\mathcal{D}^F(\pi, y) = \mathcal{D}(\pi, F, y)$.

Remark 3.6. (i) Since $\pi^\perp \circ \pi = 0$ and $\pi \circ \pi = \pi$, then we can write

$$\mathcal{D}(\pi, F, y) = \left(\pi, \pi^\perp \circ [F(y), \pi] \circ \pi \right) = \left(\pi, \pi^\perp \circ F(y) \circ \pi \right).$$

(ii) Let \mathcal{O}_{W^1} denote the zero section of W^1 . Then $F(y) \in \text{Rep}(Q, \mathbf{v})$ preserves the subspaces defined by π iff $\pi^\perp \circ F(y) \circ \pi = 0$, which by the previous statement is equivalent to saying that $F(y)$ preserves π iff $(\pi, F, y) \in \mathcal{D}^{-1}(\mathcal{O}_{W^1})$.

(iii) The same computation as above shows that

$$F(y) \text{ preserves } \pi \iff \pi^\perp \circ [F(y), \pi] = 0 \iff [F(y), \pi] \circ \pi = 0.$$

Proposition 3.7. (i) $\mathcal{D} : \text{Gr}(\mathbf{v}', \mathbf{v}) \times \mathcal{S} \times Y \rightarrow W^1$ is a C^q map of fibre bundles over $\text{Gr}(\mathbf{v}', \mathbf{v})$.

(ii) \mathcal{D} is transverse to the zero section \mathcal{O}_{W^1} of W^1 .

(iii) Let W_π^1 be the fibre of W^1 over $\pi \in \text{Gr}(\mathbf{v}', \mathbf{v})$. Then

$$(3.7) \quad \text{codim}_{W^1} \mathcal{O}_{W^1} = \dim_{\mathbb{R}} W_\pi^1 = \dim_{\mathbb{R}} \text{Hom}^1(\pi, \pi^\perp)$$

$$(3.8) \quad \dim_{\mathbb{R}} \text{Gr}(\mathbf{v}', \mathbf{v}) \times Y = \dim_{\mathbb{R}} \text{Hom}^0(\pi, \pi^\perp) + n + 1$$

Proof. It is clear that \mathcal{D} maps fibres of $\mathrm{Gr}(\mathbf{v}', \mathbf{v}) \times \mathcal{S} \times Y$ to fibres of W^1 , and so the first statement follows from the fact that F is C^q .

To see the second statement, a calculation shows that the derivative of \mathcal{D} with respect to F at a point $(\pi, F, y) \in \mathcal{D}^{-1}(\mathcal{O}_{W^1})$ is given by

$$(3.9) \quad d_2 \mathcal{D}_{(\pi, F, y)}(\delta F) = \pi^\perp \circ \delta F \circ \pi.$$

Since $s : X \rightarrow \mathrm{Rep}(Q, \mathbf{v})^{\alpha-st}$, $A_0 \in \mathrm{Rep}(Q, \mathbf{v})^{\alpha-st}$, and $F \in \mathcal{S}(s)$, then $(\pi, F, y) \in \mathcal{D}^{-1}(\mathcal{O}_{W^1})$ implies that $y \notin X \times \{0\}$, $y \notin X \times \{1\}$ and $y \notin \{x_0\} \times [0, 1]$. Therefore, from (3.6), $\delta F(y)$ can take any value in $\mathrm{Rep}(Q, \mathbf{v})$, and so the above equation shows that $d_2 \mathcal{D}_{(\pi, F, y)}$ is onto $\mathrm{Hom}^1(\pi, \pi^\perp)$, the fibre of W^1 at π .

In the proof of the third statement, equation (3.7) follows by observing that $W_\pi^1 \cong \mathrm{Hom}^1(\pi, \pi^\perp)$. Equation (3.8) then follows from the definition of $\mathrm{Gr}(\mathbf{v}', \mathbf{v})$ in (3.1). \square

In the following Corollary, recall that a *residual subset* is the countable intersection of open dense subsets.

Corollary 3.8. *If $q \geq 1$ and $\dim_{\mathbb{R}} \mathrm{Hom}^0(\pi, \pi^\perp) + n + 1 < \dim_{\mathbb{R}} \mathrm{Hom}^1(\pi, \pi^\perp)$, then for F in a residual subset of \mathcal{S} , the image $\mathcal{D}^F(\mathrm{Gr}(\mathbf{v}', \mathbf{v}) \times S^n \times [0, 1])$ does not intersect \mathcal{O}_{W^1} .*

Proof. For $(\pi, F, y) \in \mathcal{D}^{-1}(\mathcal{O}_{W^1})$, let $p : T_{\mathcal{D}(\pi, F, y)} W^1 \rightarrow T_0 W_\pi^1$ be the natural projection map. Since the manifolds $\mathrm{Gr}(\mathbf{v}', \mathbf{v})$ and Y are finite-dimensional, as are the fibres W_π^1 of the bundle W^1 , then $p \circ (\delta \mathcal{D}^F)_{(\pi, y)} : T_\pi \mathrm{Gr}(\mathbf{v}', \mathbf{v}) \times T_y Y \rightarrow T_0 W_\pi^1$ is Fredholm of index

$$\dim_{\mathbb{R}} \mathrm{Gr}(\mathbf{v}', \mathbf{v}) + \dim_{\mathbb{R}} Y - \dim_{\mathbb{R}} W_\pi^1 = \dim_{\mathbb{R}} \mathrm{Hom}^0(\pi, \pi^\perp) + n + 1 - \dim \mathrm{Hom}^1(\pi, \pi^\perp).$$

Therefore a combination of Proposition 3.7 together with the Daskalopoulos-Uhlenbeck version of the transversal density theorem (Theorems A2 and A4 in [6]) shows that the set

$$\mathcal{S}_{\mathcal{O}_{W^1}} = \{F \in \mathcal{S} : \mathcal{D}^F \text{ is transversal to } \mathcal{O}_{W^1}\}$$

is residual in \mathcal{S} .

Since the dimension of the fibres of W^1 (or alternately, the codimension of \mathcal{O}_{W^1} in W^1) is given by $\dim \mathrm{Hom}^1(\pi, \pi^\perp)$, and by assumption we have $\dim \mathrm{Gr}(\mathbf{v}', \mathbf{v}) + n + 1 < \dim \mathrm{Hom}^1(\pi, \pi^\perp)$, then the map $\mathcal{D}^F : \mathrm{Gr}(\mathbf{v}', \mathbf{v}) \times S^n \times [0, 1] \rightarrow W^1$ is transverse to the zero section \mathcal{O}_{W^1} iff the image $\mathcal{D}^F(\mathrm{Gr}(\mathbf{v}', \mathbf{v}) \times S^n \times [0, 1])$ does not intersect \mathcal{O}_{W^1} . \square

Remark 3.9. This corollary shows that there is a residual set of C^1 basepoint-preserving homotopy equivalences F between $s : S^n \rightarrow \mathrm{Rep}(Q, \mathbf{v})^{\alpha-st}$ and the constant map, such that no representation in the image of F preserves a sub-representation in $\mathrm{Gr}(\mathbf{v}', \mathbf{v})$.

Theorem 3.10. *If $n + 1 < d_{\min}(Q, \mathbf{v}, \alpha)$ then $\pi_n(\mathrm{Rep}(Q, \mathbf{v})^{\alpha-st}) = 0$.*

Proof. Fix a continuous map $s : S^n \rightarrow \mathrm{Rep}(Q, \mathbf{v})^{\alpha-st}$. We aim to show that there is a basepoint-preserving homotopy equivalence in $\mathrm{Rep}(Q, \mathbf{v})^{\alpha-st}$ between s and the constant map. Firstly note

that since the set of stable points $\text{Rep}(Q, \mathbf{v})^{\alpha-st}$ is open in the vector space $\text{Rep}(Q, \mathbf{v})$, then we can perturb s (while remaining in $\text{Rep}(Q, \mathbf{v})^{\alpha-st}$) so that $s \in C^1$ (cf. Theorem 10.21 of [25]).

Corollary 3.8 then shows that for each $\mathbf{v}' < \mathbf{v}$ there is a residual set $\mathcal{S}(s)_{\mathbf{v}'}^0$ of homotopy equivalences F from s to the constant map such that no representation in the image of F preserves a subrepresentation in $\text{Gr}(\mathbf{v}', \mathbf{v})$.

There are a finite number of choices of $\mathbf{v}' < \mathbf{v}$ such that $\text{slope}_\alpha(Q, \mathbf{v}') \geq \text{slope}_\alpha(Q, \mathbf{v})$, and so the set

$$\mathcal{S}(s)_\alpha^0 = \bigcap_{\substack{\mathbf{v}' < \mathbf{v} \\ \text{slope}_\alpha(Q, \mathbf{v}') \geq \text{slope}_\alpha(Q, \mathbf{v})}} \mathcal{S}(s)_{\mathbf{v}'}^0$$

is residual in $\mathcal{S}(s)$. In other words, there is a residual set $\mathcal{S}(s)_\alpha^0$ of C^1 homotopy equivalences between s and the constant map, such that for every $F \in \mathcal{S}(s)_\alpha^0$ the image $F(S^n \times [0, 1])$ is contained in $\text{Rep}(Q, \mathbf{v})^{\alpha-st}$.

Since residual sets are non-empty, then there is at least one basepoint-preserving homotopy equivalence in $\text{Rep}(Q, \mathbf{v})^{\alpha-st}$ between s and the constant map, and so $\pi_n(\text{Rep}(Q, \mathbf{v})^{\alpha-st}) = 0$. \square

Corollary 3.11. *If $n + 1 < d_{\min}(Q, \mathbf{v}, \alpha)$ then $\pi_n(\mathcal{M}_\alpha(Q, \mathbf{v})^{st}) \cong \pi_{n-1}(PG_{\mathbf{v}})$.*

Proof. The group $PG_{\mathbf{v}}$ acts freely on $\text{Rep}(Q, \mathbf{v})^{\alpha-st}$, and the quotient is

$$\mathcal{M}_\alpha(Q, \mathbf{v})^{st} = \text{Rep}(Q, \mathbf{v})^{\alpha-st} / PG_{\mathbf{v}}.$$

The fibre bundle

$$PG_{\mathbf{v}} \rightarrow \text{Rep}(Q, \mathbf{v})^{\alpha-st} \rightarrow \mathcal{M}_\alpha(Q, \mathbf{v})^{st}$$

induces a long exact sequence of homotopy groups

$$\cdots \rightarrow \pi_n(\text{Rep}(Q, \mathbf{v})^{\alpha-st}) \rightarrow \pi_n(\mathcal{M}_\alpha(Q, \mathbf{v})^{st}) \rightarrow \pi_{n-1}(PG_{\mathbf{v}}) \rightarrow \pi_{n-1}(\text{Rep}(Q, \mathbf{v})^{\alpha-st}) \rightarrow \cdots,$$

and therefore an application of Theorem 3.10 shows that

$$\pi_n(\text{Rep}(Q, \mathbf{v})^{\alpha-st}) = \pi_{n-1}(\text{Rep}(Q, \mathbf{v})^{\alpha-st}) = 0$$

for $n < d_{\min}(Q, \mathbf{v}, \alpha) - 1$. This shows that from the long exact sequence above, we have the isomorphism

$$\pi_n(\mathcal{M}_\alpha(Q, \mathbf{v})^{st}) \cong \pi_{n-1}(PG_{\mathbf{v}}),$$

which completes the proof. \square

4. EXAMPLES

This section contains two applications of Theorem 3.10 and Corollary 3.11. The first is to study the low-dimensional topology of moduli spaces that contain the stable loci of Nakajima quiver varieties, an explicit calculation is performed for the case of the ADHM quiver with the results in Theorem 4.2. The second application of the main theorem is to the moduli space of polygons in \mathbb{R}^3 , which parametrises polygons in \mathbb{R}^3 with a fixed number of sides (determined by the quiver), and fixed sidelengths (determined by the choice of parameter), up to equivalence by rigid motions

in \mathbb{R}^3 . For almost all choices of parameter (we call these *generic* parameters) there do not exist any polygons that lie in a line, and the moduli space is smooth. For non-generic parameters, *degenerate polygons* (those that lie in a line) correspond to singularities in the moduli space $\mathcal{M}_\alpha(Q, \mathbf{v})$, since there are rotations of \mathbb{R}^3 that leave such polygons invariant. In Theorem 4.5 we compute the low-dimensional homotopy groups of the space of *non-degenerate polygons* (those that are not contained in a line), in the case where the parameter α is non-generic.

4.1. The “Doubled” ADHM quiver. It is well-known that the moduli space of framed instantons on S^4 has a description in terms of a finite-dimensional hyperkähler quotient construction, known as the *ADHM construction* (see [3], [7] for more details).

This finite-dimensional hyperkähler quotient construction is a special case of what is now known as a *Nakajima quiver variety*, where a new quiver \tilde{Q} is constructed by adding a “dual” edge to each edge on the original quiver Q , and the moduli space $\mathcal{M}_{(\alpha,\beta)}^{hk}(Q, \mathbf{v})$ is defined to be the hyperkähler quotient of $\text{Rep}(\tilde{Q}, \mathbf{v}) = T^* \text{Rep}(Q, \mathbf{v})$ with parameters α and β (see [26] for a complete description and [23], [24] for more examples).

There is an inclusion of each of these hyperkähler quotients $\mathcal{M}_{(\alpha,\beta)}^{hk}(Q, \mathbf{v}) \hookrightarrow \mathcal{M}_\alpha(\tilde{Q}, \mathbf{v})$ into the moduli space associated to the quiver \tilde{Q} , with the same dimension vector and stability parameter. The purpose of this section is to use Theorem 3.10 to study the topology of the space $\mathcal{M}_0(\tilde{Q}, \mathbf{v})^{st}$, which contains $\mathcal{M}_{(0,0)}^{hk}(Q, \mathbf{v})$, in the case of the quiver corresponding to the ADHM construction of framed instantons on S^4 . It is worth mentioning also that the method used is more general than just this one specific example, one can also prove analogous results for any Nakajima quiver variety by computing d_{min} for each example.

Fix a vector bundle E over S^4 with rank n and instanton number k . Then the quiver \tilde{Q} under consideration is given as follows.


$$(4.1) \quad \begin{array}{c} \begin{array}{c} \curvearrowright \quad \curvearrowleft \\ \bullet \mathbb{C}^k \\ \uparrow \quad \downarrow \\ \dots \quad \dots \\ \uparrow \quad \downarrow \\ \bullet \mathbb{C} \end{array} \end{array}$$

(note that this differs slightly from Donaldson’s description in [7], here we use the construction of Crawley-Boevey in [4] which identifies the moduli space associated to the quiver described above with the solutions to the equations from [7]).

Let $\alpha = 0$ be the stability parameter. There are n arrows from the vertex \mathbb{C}^k to the vertex \mathbb{C} , another n arrows from the vertex \mathbb{C} to the vertex \mathbb{C}^k , and two arrows from \mathbb{C}^k to itself. In this case $G_{\mathbf{v}} = \text{U}(k) \times \text{U}(1)$, and $PG_{\mathbf{v}} \cong \text{U}(k)$.

Remark 4.1. When $\alpha = 0$ then every representation is α -semistable, and a representation is α -stable if and only if there are no non-trivial subrepresentations.


A subrepresentation π can have one of the following forms (the diagram below also includes the quiver for π^\perp). Firstly we have the case where the vertex \mathbb{C} is contained in the subrepresentation π . Since π cannot have dimension vector \mathbf{v} , then $k_1 < k$ in the diagram below.



The elements of $\text{Hom}^1(\pi, \pi^\perp)$ correspond to maps from the vertex \mathbb{C}^{k_1} to \mathbb{C}^{k-k_1} (there are two edges with this property) and maps from the vertex \mathbb{C} to \mathbb{C}^{k-k_1} (there are n edges with this property). Therefore we see that $\dim_{\mathbb{R}} \text{Hom}^1(\pi, \pi^\perp) = 4k_1(k - k_1) + 2n(k - k_1)$. A similar computation shows that $\dim_{\mathbb{R}} \text{Hom}^0(\pi, \pi^\perp) = 2k_1(k - k_1)$. Therefore in this case

$$(4.2) \quad \dim_{\mathbb{R}} \text{Hom}^1(\pi, \pi^\perp) - \dim_{\mathbb{R}} \text{Hom}^0(\pi, \pi^\perp) = 2(k_1 + n)(k - k_1).$$

The second case is when the vertex \mathbb{C} is contained in π^\perp . Since π cannot have dimension vector $\mathbf{0}$, then $k_1 > 0$ in the diagram below.



A calculation shows that

$$\dim_{\mathbb{R}} \text{Hom}^1(\pi, \pi^\perp) = 4k_1(k - k_1) + 2nk_1,$$

and $\dim_{\mathbb{R}} \text{Hom}^0(\pi, \pi^\perp) = 2k_1(k - k_1)$. Therefore in this case

$$(4.3) \quad \dim_{\mathbb{R}} \text{Hom}^1(\pi, \pi^\perp) - \dim_{\mathbb{R}} \text{Hom}^0(\pi, \pi^\perp) = 2k_1(k - k_1 + n).$$

In both cases we see that $\dim_{\mathbb{R}} \text{Hom}^1(\pi, \pi^\perp) - \dim_{\mathbb{R}} \text{Hom}^0(\pi, \pi^\perp) \geq 2(k - 1 + n)$, and so $d_{\min} \geq 2(k - 1 + n)$. Therefore Theorem 3.10 gives the following result.

Theorem 4.2. *Let \tilde{Q} be the quiver in (4.1), and let $\alpha = 0$. Then*

$$\pi_j \left(\text{Rep}(\tilde{Q}, \mathbf{v})^{\alpha-st} \right) = 0$$


for all $j < 2(k - 1 + n) - 1$, and

$$\pi_j \left(\mathcal{M}_\alpha(\tilde{Q}, \mathbf{v})^{st} \right) = \pi_{j-1}(\text{U}(k))$$

for all $j < 2(k - 1 + n) - 1$.


4.2. The moduli space of non-degenerate polygons in \mathbb{R}^3 . The topology and geometry of the moduli space of polygons in \mathbb{R}^3 with fixed sidelengths has been well-studied, see for example [21], [13], [11], [10]. When the sidelengths are chosen so that some polygons may lie in a line (we call these *degenerate* polygons) then the moduli space has singularities, corresponding to the fact that a degenerate polygon is invariant under a circle subgroup of the rotation group. The purpose of this section is to prove new results about the low-dimensional topology of the moduli space of *non-degenerate* polygons.

An n -sided polygon in \mathbb{R}^3 corresponds to a choice of basepoint $b \in \mathbb{R}^3$, as well as n vectors e_1, \dots, e_n such that $e_1 + \dots + e_n = 0$, where the vertices of the polygon are then given by $v_k = b + e_1 + \dots + e_k$. The *moduli space of polygons in \mathbb{R}^3 with fixed sidelengths* is the space of all polygons in \mathbb{R}^3 with basepoint 0 and given sidelengths s_1, \dots, s_n , modulo rotations by $\text{SO}(3)$. It is well-known (see for example [22]) that this moduli space corresponds to the representation variety of a quiver, given as follows. Let Q be the “star-shaped” quiver below, with one central vertex and n outer vertices (one for each side of the polygon). Fix a dimension vector \mathbf{v} , with dimension 2 at the central vertex, and dimension 1 at each of the outer vertices. In this case $G_{\mathbf{v}} = \text{U}(2) \times \text{U}(1)^n$, and $PG_{\mathbf{v}} \cong \text{U}(2) \times \text{U}(1)^{n-1}$.



With the conventions of Section 2, the parameter α_j at the j^{th} vertex is chosen so that $\alpha_j = -s_j$ (where s_j is the sidelength of the j^{th} edge of the polygon in \mathbb{R}^3), and the parameter α_0 is then given by $\alpha_0 = -\frac{1}{2} \sum_{j=1}^n \alpha_j$, i.e. the associated central element $\alpha \in \mathfrak{g}_{\mathbf{v}}^*$ is trace-free.


Since $\alpha_j < 0$ for each $j = 1, \dots, n$ then by the definition of slope stability in (2.8) we see that a representation which is not α -stable must correspond to one of the following two cases. The first case consists of those subrepresentations π that do not contain any subspaces of the central vertex (as for the examples in the previous section, the diagram below includes the subrepresentation π^\perp). The outer vertices in the following diagram have been re-labelled, so that α_{j_i} for $1 \leq i \leq k$ are in the subrepresentation π , and α_{j_i} for $k + 1 \leq i \leq n$ are in the subrepresentation π^\perp .



Note that since the subrepresentation π is non-trivial, then $k \geq 1$. The elements of $\text{Hom}^1(\pi, \pi^\perp)$ correspond to maps from the vertices in the quiver on the left (corresponding to the subrepresentation π) to the central vertex in the quiver on the right (corresponding to π^\perp). A calculation shows that $\dim_{\mathbb{R}} \text{Hom}^1(\pi, \pi^\perp) = 4k$, and that $\dim_{\mathbb{R}} \text{Hom}^0(\pi, \pi^\perp) = 0$. Therefore, in this case

$$(4.4) \quad \dim_{\mathbb{R}} \text{Hom}^1(\pi, \pi^\perp) - \dim_{\mathbb{R}} \text{Hom}^0(\pi, \pi^\perp) = 4k \geq 4.$$

The other possible type of subrepresentation occurs when the subrepresentation π contains a complex dimension one subspace of the central vertex, described in the diagram below.



Unlike the previous case (where any value of k between 1 and n could occur), the range of possible values of k is limited by the choice of α_j at each vertex (i.e. the sidelengths of the polygon). To see this, let \mathbf{v}' be the dimension vector associated to the subrepresentation π , and note that for the condition $\text{slope}_\alpha(Q, \mathbf{v}') \geq \text{slope}_\alpha(Q, \mathbf{v})$ to hold we must have

$$\deg(Q, \mathbf{v}') = -\alpha_0 - \sum_{i=1}^k \alpha_{j_i} \geq 0,$$

which places a restriction on the possible configurations that can occur in this case (recall that $\alpha_0 > 0$ and that $\alpha_j < 0$ for $j = 1, \dots, k$).

A calculation shows that $\dim_{\mathbb{R}} \text{Hom}^1(\pi, \pi^\perp) = 2k$, and that $\dim_{\mathbb{R}} \text{Hom}^0(\pi, \pi^\perp) = 2$. Therefore, in this case

$$(4.6) \quad \dim_{\mathbb{R}} \text{Hom}^1(\pi, \pi^\perp) - \dim_{\mathbb{R}} \text{Hom}^0(\pi, \pi^\perp) = 2k - 2.$$

In analogy with [10], define a subset $J \subset \{1, \dots, n\}$ to be *strictly short* if

$$(4.7) \quad -\sum_{j \in J} \alpha_j < -\sum_{j \notin J} \alpha_j.$$

Equivalently, J is strictly short if $-\alpha_0 - \sum_{j \in J} \alpha_j < 0$.

Definition 4.3. For a positive integer ℓ , the parameter α is ℓ -*strictly short* if every subset $J \subset \{1, \dots, n\}$ with ℓ elements is strictly short. If there is no positive integer ℓ for which α is ℓ -strictly short then α is called *0-strictly short*.

Remark 4.4. If α is ℓ -strictly short, then there are no subrepresentations of the type in diagram (4.5) with $k \leq \ell$. Therefore by (4.6) we have

$$(4.8) \quad \dim_{\mathbb{R}} \text{Hom}^1(\pi, \pi^\perp) - \dim_{\mathbb{R}} \text{Hom}^0(\pi, \pi^\perp) \geq 2\ell.$$

Equations (4.4) and (4.8) show that d_{min} takes the following values when α is ℓ -strictly short.

d_{min}	ℓ
0	0
2	1
4	2 or greater

Together with Theorem 3.10 and Corollary 3.11, the above calculations lead to the following theorem.

Theorem 4.5. (i) *If α is 2-strictly short then*

- $\pi_j(\text{Rep}(Q, \mathbf{v})^{\alpha-st}) = 0$ for $j \leq 2$.
- $\pi_j(\mathcal{M}_\alpha(Q, \mathbf{v})^{st}) = \pi_{j-1}(\mathbf{U}(2) \times \mathbf{U}(1)^{n-1})$ for $j \leq 2$.

(ii) *If α is 1-strictly short then $\mathcal{M}_\alpha(Q, \mathbf{v})^{st}$ and $\text{Rep}(Q, \mathbf{v})^{\alpha-st}$ are both connected.*

ACKNOWLEDGMENTS

The author would like to thank Georgios Daskalopoulos for his advice and encouragement, as well as the referee for some useful observations and background references.

REFERENCES

- [1] Ralph Abraham and Joel Robbin. *Transversal mappings and flows*. An appendix by Al Kelley. W. A. Benjamin, Inc., New York-Amsterdam, 1967.
- [2] M. F. Atiyah and R. Bott. The Yang-Mills equations over Riemann surfaces. *Philos. Trans. Roy. Soc. London Ser. A*, 308(1505):523–615, 1983.
- [3] M. F. Atiyah, N. J. Hitchin, V. G. Drinfel'd, and Yu. I. Manin. Construction of instantons. *Phys. Lett. A*, 65(3):185–187, 1978.
- [4] William Crawley-Boevey. Geometry of the moment map for representations of quivers. *Compositio Math.*, 126(3):257–293, 2001.
- [5] Georgios D. Daskalopoulos. The topology of the space of stable bundles on a compact Riemann surface. *J. Differential Geom.*, 36(3):699–746, 1992.
- [6] Georgios D. Daskalopoulos and Karen K. Uhlenbeck. An application of transversality to the topology of the moduli space of stable bundles. *Topology*, 34(1):203–215, 1995.
- [7] S. K. Donaldson. Instantons and geometric invariant theory. *Comm. Math. Phys.*, 93(4):453–460, 1984.
- [8] Megumi Harada and Graeme Wilkin. Morse theory of the moment map for representations of quivers. Available at arxiv:0807.4734.
- [9] G. Harder and M. S. Narasimhan. On the cohomology groups of moduli spaces of vector bundles on curves. *Math. Ann.*, 212:215–248, 1974/75.
- [10] J.-C. Hausmann and A. Knutson. The cohomology ring of polygon spaces. *Ann. Inst. Fourier (Grenoble)*, 48(1):281–321, 1998.
- [11] Jean-Claude Hausmann and Allen Knutson. Polygon spaces and Grassmannians. *Enseign. Math. (2)*, 43(1-2):173–198, 1997.
- [12] N. J. Hitchin. The self-duality equations on a Riemann surface. *Proc. London Math. Soc. (3)*, 55(1):59–126, 1987.
- [13] Michael Kapovich and John J. Millson. The symplectic geometry of polygons in Euclidean space. *J. Differential Geom.*, 44(3):479–513, 1996.
- [14] A. D. King. Moduli of representations of finite-dimensional algebras. *Quart. J. Math. Oxford Ser. (2)*, 45(180):515–530, 1994.
- [15] Frances Kirwan. On spaces of maps from Riemann surfaces to Grassmannians and applications to the cohomology of moduli of vector bundles. *Ark. Mat.*, 24(2):221–275, 1986.
- [16] Frances Kirwan. On the homology of compactifications of moduli spaces of vector bundles over a Riemann surface. *Proc. London Math. Soc. (3)*, 53(2):237–266, 1986.
- [17] Frances Kirwan. Rational intersection cohomology of quotient varieties. *Invent. Math.*, 86(3):471–505, 1986.

- [18] Frances Kirwan. Geometric invariant theory and the Atiyah-Jones conjecture. In *The Sophus Lie Memorial Conference (Oslo, 1992)*, pages 161–186. Scand. Univ. Press, Oslo, 1994.
- [19] Frances Clare Kirwan. *Cohomology of quotients in symplectic and algebraic geometry*, volume 31 of *Mathematical Notes*. Princeton University Press, Princeton, NJ, 1984.
- [20] Frances Clare Kirwan. Partial desingularisations of quotients of nonsingular varieties and their Betti numbers. *Ann. of Math. (2)*, 122(1):41–85, 1985.
- [21] Alexander A. Klyachko. Spatial polygons and stable configurations of points in the projective line. In *Algebraic geometry and its applications (Yaroslavl, 1992)*, Aspects Math., E25, pages 67–84. Vieweg, Braunschweig, 1994.
- [22] Hiroshi Konno. On the cohomology ring of the hyperkähler analogue of the polygon spaces. *Contemporary Mathematics*, 309:129–149, 2002.
- [23] P. B. Kronheimer. The construction of ALE spaces as hyper-Kähler quotients. *J. Differential Geom.*, 29(3):665–683, 1989.
- [24] Peter B. Kronheimer and Hiraku Nakajima. Yang-Mills instantons on ALE gravitational instantons. *Math. Ann.*, 288(2):263–307, 1990.
- [25] John M. Lee. *Introduction to smooth manifolds*, volume 218 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2003.
- [26] Hiraku Nakajima. Instantons on ALE spaces, quiver varieties, and Kac-Moody algebras. *Duke Math. J.*, 76(2):365–416, 1994.
- [27] Markus Reineke. The Harder-Narasimhan system in quantum groups and cohomology of quiver moduli. *Invent. Math.*, 152(2):349–368, 2003.
- [28] Claus Michael Ringel. Representations of K -species and bimodules. *J. Algebra*, 41(2):269–302, 1976.
- [29] Aidan Schofield. General representations of quivers. *Proc. London Math. Soc. (3)*, 65(1):46–64, 1992.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COLORADO, BOULDER, CO 80303
E-mail address: graeme.wilkin@colorado.edu