



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/149941/>

Version: Accepted Version

Article:

Kirkham, E.J. and Levita, L. (2020) Early life stress is associated with reduced avoidance of threatening facial expressions. *Development and Psychopathology*, 32 (3). pp. 1059-1067. ISSN: 0954-5794

<https://doi.org/10.1017/s095457941900110x>

This article has been published in a revised form in *Development and Psychopathology*, <http://doi.org/10.1017/s095457941900110x>. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works. © Cambridge University Press 2019.

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



CAMBRIDGE
UNIVERSITY PRESS

Early life stress is associated with reduced avoidance of threatening facial expressions

Journal:	<i>Development and Psychopathology</i>
Manuscript ID	DPP-2018-00027.R3
Manuscript Type:	Regular Article
Keyword:	early life stress, emotional facial expressions, anger, avoidance, approach
Abstract:	<p>Early neglect or maltreatment has been associated with changes in children's processing of emotional facial expressions, including a hypersensitivity to the emotion of anger. This may facilitate the avoidance of danger in a maltreating environment. However, few studies have examined whether experiences of early life stress (ELS) are associated with atypical avoidance responses towards emotional facial expressions, nor whether effects of ELS can be observed in adult participants. The present study therefore examined the effect of ELS on adults' approach-avoidance tendencies towards angry, happy and neutral facial expressions. Surprisingly, higher levels of ELS were associated with reduced avoidance of angry facial expressions amongst individuals with no evidence of mental illness. In contrast, there was no evidence of a relationship between ELS and avoidance of angry facial expressions amongst individuals with experience of mental illness. These novel findings suggest that ELS-related changes in social cognition can be observed years after the ELS itself occurred.</p>

SCHOLARONE™
Manuscripts

Abstract

Early neglect or maltreatment has been associated with changes in children's processing of emotional facial expressions, including a hypersensitivity to the emotion of anger. This may facilitate the avoidance of danger in a maltreating environment. However, few studies have examined whether experiences of early life stress (ELS) are associated with atypical avoidance responses towards emotional facial expressions, nor whether effects of ELS can be observed in adult participants. The present study therefore examined the effect of ELS on adults' approach-avoidance tendencies towards angry, happy and neutral facial expressions. Surprisingly, higher levels of ELS were associated with reduced avoidance of angry facial expressions amongst individuals with no evidence of mental illness. In contrast, there was no evidence of a relationship between ELS and avoidance of angry facial expressions amongst individuals with experience of mental illness. These novel findings suggest that ELS-related changes in social cognition can be observed years after the ELS itself occurred.

Keywords: early life stress, emotional facial expressions, anger, avoidance, approach

1
2
3 Early life stress (ELS) such as abuse or neglect has been associated with altered
4 processing of emotional facial expressions (Gibb, Schofield, & Coles, 2009; McCrory et al.,
5 2013; Pollak, Messner, Kistler, & Cohn, 2009). Specifically, children who experienced ELS
6 show hyper-sensitivity to the emotion of anger (Maheu et al., 2010; McCrory et al., 2011;
7 Pollak, 2008). Compared to non-maltreated children, these individuals can identify anger
8 with fewer perceptual cues (Pollak & Sinha, 2002) and misattribute the emotion of anger to
9 neutral facial expressions (Pollak & Kistler, 2002). Pollak et al. (2009) presented children
10 with a continuum of facial expressions which transitioned gradually from a neutral expression
11 to a prototypical expression of anger. Children who had experienced ELS identified anger
12 earlier in the continuum than those who had not been maltreated. These findings are
13 supported by fMRI studies which have found heightened amygdala activation to angry facial
14 expressions amongst children and adults who experienced ELS (Dannowski et al., 2012; Di
15 Iorio et al., 2017; Maheu et al., 2010; McCrory et al., 2011; Redlich et al., 2015).

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33 Furthermore, previous research suggests that ELS-related changes in facial emotion
34 processing are specific to the emotion of anger; responses to happy facial expressions do not
35 appear to be related to experiences of ELS (Gibb et al., 2009; Pollak & Kistler, 2002; Pollak
36 et al., 2009; Pollak & Sinha, 2002). The observation that ELS-related changes in facial
37 emotion processing appear to be specific to anger supports the theory that these changes
38 represent an adaptation to an abusive environment (McCrory & Viding, 2015; Pollak et al.,
39 2009). In a maltreating context, the earlier a child can identify a threat (as indexed by the
40 presence of anger), the better their chances of avoiding a harmful outcome. However, despite
41 the intuitive value of this explanation, there has been very little investigation into whether
42 experiences of ELS are indeed associated with atypical avoidance of threat as depicted by
43 angry facial expressions.

1
2
3 In addition, the majority of behavioural research into the relationship between ELS
4 and atypical responding to facial expressions has been carried out with children rather than
5 with adults. One study which did focus on adults, conducted by Gibb et al. (2009), found that
6 those who experienced high levels of ELS were better able to identify subtle expressions of
7 anger when compared to adults who did not experience ELS. This suggests that ELS-related
8 alterations in responsivity to facial expressions of anger are not specific to childhood but can
9 also be observed during adulthood (Doretto & Scivoletto, 2018). However, it remains unclear
10 whether these alterations in face processing extend to atypical avoidance of such expressions.
11
12
13
14
15
16
17
18
19
20

21 Atypical avoidance of emotional facial expressions has been found amongst adults
22 with depression (Derntl et al., 2011), a condition which is positively associated with
23 experiences of ELS (Kessler et al., 2010). Seidel et al. (2010) found that individuals with
24 depression were more avoidant in their responses to emotional facial expressions than healthy
25 controls, and that these effects were most pronounced amongst female participants. This latter
26 finding is supported by work with healthy participants which found that female participants
27 showed greater levels of self-reported avoidance of emotional facial expressions than male
28 participants (Miller, Chabriac, & Molet, 2013). Given that experiences of ELS intersect with
29 depression, anxiety and gender (Cooke & Weathington, 2014; Heim & Nemeroff, 2001), it is
30 essential that any investigation of atypical avoidance of emotional facial expressions
31 following ELS controls for participants' gender and levels of depression and anxiety.
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 Emerging evidence suggests that the relationships between ELS and cognitive
47 processes may present differently in adults who have gone on to develop a mental illness than
48 in those who have remained mentally well (Suzuki, Poon, Humari & Cleare, 2015). For
49 example, Saleh et al. (2017) found that, amongst people with depression, higher levels of
50 ELS were associated with reduced processing speed, whereas amongst people without
51 depression, higher levels of ELS were associated with increased processing speed. Their
52
53
54
55
56
57
58
59
60

1
2
3 findings support the theory of stress inoculation (Meichenbaum & Cameron, 1989), which
4
5 states that whilst chronic, uncontrollable early stress increases vulnerability to mental illness,
6
7 moderate, manageable early stress may entrain resilience to future sources of stress
8
9
10 (Southwick & Charney, 2012; Ashokan, Sivasubramanian & Mitra, 2016). In light of this, the
11
12 present study included additional exploratory analysis to examine whether the interaction
13
14 between ELS and the presence of mental illness was associated with adults' approach-
15
16 avoidance of emotional facial expressions.
17
18

19
20 The present study therefore examined the relationship between adults' experiences of
21
22 ELS and their approach-avoidance responses towards emotional facial expressions. Using a
23
24 similar approach to Seidel, Habel, Kirschner, Gur, and Derntl (2010), participants were
25
26 required to indicate their tendency to approach or avoid images of male and female angry,
27
28 happy and neutral facial expressions. It was hypothesised that higher levels of ELS would be
29
30 associated with increased avoidance responses to angry facial expressions. Furthermore, it
31
32 was tentatively predicted that the relationship between ELS and approach-avoidance
33
34 responses to angry facial expressions may vary according to the presence or absence of
35
36 mental illness. In addition, it was hypothesised that ELS would not be associated with
37
38 approach-avoidance responses to happy facial expressions.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Method

Participants

Three hundred and ninety-eight individuals took part in the study. As the study included investigation of gender effects, two participants who described their gender as “other” were excluded. An additional participant was excluded due to incorrect completion of a questionnaire. The final sample therefore consisted of 395 participants (235 female) with a mean age of 23.25 years ($SD = 8.92$). All participants were from the United Kingdom. Ninety participants reported that they had been diagnosed with a mental illness at some time in their lives. Participants gave informed consent and were made aware that they could withdraw from the study at any time. The study was approved by the Department of Psychology Ethics Committee, University of Sheffield. Forty-eight participants (recruited via www.prolific.ac) received £2.10 for completing the study; the remainder were offered entry to a £20 prize draw.

Materials

Face stimuli. Colour photographs of angry, happy and neutral facial expressions were taken from the “FACES” database, a validated database of facial expressions (Ebner, Riediger, & Lindenberger, 2010). All of the colour photographs depict frontal views of the head, neck and upper shoulders of young (aged 19 – 31) Caucasian individuals. Pilot testing indicated that use of the full stimulus set (48 images depicting different individuals) would lead to attrition. Consequently, the full set was split into two sets of 24 images, and each participant viewed one of these sets. The stimulus sets were each comprised of 12 male faces and 12 female faces, with four faces of each gender depicting each facial expression (angry, happy and neutral).

Questionnaires. ELS was measured with the Child Abuse and Trauma Scale (CATS; Sanders & Becker-Lausen, 1995), a 38 item questionnaire concerning the respondent’s home

environment during their childhood and adolescence. Example items include “Were you expected to follow a strict code of behavior in your home?”, “Did your parents insult you or call you names?” and “How often were you left at home alone as a child?”. Responses to each question are measured using a five point Likert scale ranging from 0 (Never) to 4 (Always), resulting in possible total scores between 0 and 152. The scale showed excellent internal consistency within the current sample ($n = 395$, $\alpha = 0.93$). Participants’ psychological wellbeing was measured using the Hospital Anxiety and Depression Scale (HADS; Zigmond & Snaith, 1983), a 14-item questionnaire which provides two scores indicating levels of anxiety and depression. Respondents indicate the extent to which they experience each symptom using a scale ranging from 0 to 3. Thus, each subscale provides a total score between 0 and 21. Within the present sample the HADS showed good internal consistency for both the anxiety subscale ($n = 395$, $\alpha = 0.86$) and the depression subscale ($n = 395$, $\alpha = 0.81$).

Procedure

The study was carried out online via Qualtrics (www.qualtrics.com). After providing consent and demographic information, participants were shown one set of 24 images of male and female models depicting angry, happy and neutral facial expressions. Participants used four “sliders” located below each image (Figure 1) to indicate, on a 100 point scale, the extent to which they would approach the person in the image, the extent to which they would avoid the person in the image, how angry they perceived the person to be, and how happy they perceived them to be. The order of the rating sliders was counterbalanced across participants. The task was presented across 24 web pages, with each page displaying one image positioned above the four sliders (Figure 1). The order of image presentation was randomised across participants. After completing this task, participants completed the questionnaires measuring ELS, depression and anxiety levels.

[FIGURE 1 HERE]

Statistical analyses

Statistical analyses were performed using IBM SPSS Statistics for Windows, Version 22.0. Several measures were non-normally distributed; thus, where appropriate, original p values are reported alongside bias-corrected and accelerated 95% bootstrap confidence intervals based on 1000 samples. Prior to analysis, a single measure of approach-avoidance responsiveness was created by subtracting scores on the avoid scale from scores on the approach scale. Scores above zero on this approach-avoidance measure indicate a tendency towards approach, whilst scores below zero indicate a tendency towards avoidance. Three separate measures of approach-avoidance were calculated, one for each facial expression (angry, happy and neutral). To examine behavioural performance in the task, the approach-avoidance scores for each of the facial expressions were entered into a repeated-measures analysis of variance (ANOVA) and the mean scores for each expression were examined.

The aim of this research was to examine the effect of ELS on approach-avoidance of emotional facial expressions. To this end, a regression model was constructed in which the control variables of participant gender, age, mental illness (as indicated by a self-reported diagnosis and/or a score above the suggested clinical cut-off for one or both of the HADS scales), anxiety scores (HADS-A) and depression scores (HADS-D) were entered in the first step and the variable of interest, ELS score (CATS), was entered in the second step. In light of the widely reported relationship between ELS and mental illness, an interaction term was also created by multiplying standardised CATS scores by the dichotomous mental illness variable. This interaction term was entered in the third step of the model. The regression model was used to predict each of the three outcome variables: approach-avoidance of angry

1
2
3 facial expressions, approach-avoidance of happy expressions and approach-avoidance of
4 neutral expressions. After fitting the regression model, the residuals were examined, and any
5 cases with residuals $> \pm 3$ standard deviations were removed (Field, 2005). The analysis
6 was then re-run, and the results of the final analyses were reported. The number of cases
7 removed from each analysis is detailed in the notes below the table containing the analysis.
8
9
10
11
12
13
14
15 The highest number of cases removed from any individual analysis was five.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

Results

Demographics

Mean scores for age, ELS (CATS), depression (HADS-D) and anxiety (HADS-A) are presented in Table 1 and Table 2. Given the widely-reported relationship between ELS and depression and anxiety scores, Pearson's correlations were used to examine whether the present sample also showed this expected relationship. Due to the significant difference between male and female participants' depression and anxiety scores (Table 1), correlations were performed separately for each gender. ELS was significantly positively associated with depression scores in both male ($r(158) = .49, p < .001, \text{BCa } 95\% \text{ CI } [.35, .61]$) and female participants ($r(233) = .42, p < .001, \text{BCa } 95\% \text{ CI } [.27, .56]$). Similarly, ELS scores were significantly positively correlated with anxiety scores in both male ($r(158) = .41, p < .001, \text{BCa } 95\% \text{ CI } [.25, .54]$) and female participants ($r(233) = .46, p < .001, \text{BCa } 95\% \text{ CI } [.35, .57]$).

[TABLES 1 AND 2 HERE]

Task reliability

As the present study used a new task to measure participants' responses towards emotional facial expressions, internal consistency for each rating scale was examined using Cronbach's alpha. Cronbach's alpha was calculated for each scale in response to each facial expression (Supplementary Table 1). These alpha values were above 0.8 for all measures, indicating good reliability across the task.

Task performance

Mean approach-avoidance responses to the three facial expressions (angry, happy and neutral) are presented in Figure 2. As the approach-avoidance score was created by subtracting the avoidance score from the approach score, values above zero indicate a tendency to approach and values below zero indicate a tendency to avoid. A repeated measures ANOVA was used to examine whether participants' approach-avoidance responses differed between the three facial expressions. There was a significant effect of facial expression on approach-avoidance responses ($F(1.70, 670.81) = 1122.05, p < .001$). Examination of the mean approach-avoidance scores revealed that participants showed avoidance of angry facial expressions ($M = -36.41, SD = 32.15$) and approach of happy expressions ($M = 54.05, SD = 31.25$), whilst the mean approach-avoidance score for neutral expressions was close to zero ($M = -0.16, SD = 31.06$). As would be expected, pairwise comparisons using the Bonferroni correction demonstrated that participants were significantly more likely to avoid angry facial expressions than happy or neutral facial expressions, and significantly more likely to approach happy facial expressions than neutral expressions (all p values $< .001$).

[FIGURE 2 HERE]

Early life stress and approach-avoidance of angry facial expressions

Hierarchical regression analysis was used to examine the effect of early life stress on approach-avoidance responses to angry facial expressions (Table 3). The control variables of participants' gender, age, mental illness, anxiety scores (HADS-A) and depression scores (HADS-D) were entered in the first step of the regression model, whilst scores on the

1
2
3 measure of ELS (CATS) were entered in the second step of the model, and the interaction
4
5 term reflecting ELS multiplied by mental illness was entered in the third step.
6
7

8 The final regression model predicted a significant amount of the variance in approach-
9
10 avoidance responses towards angry facial expressions ($R^2 = .09$, $F(7, 385) = 5.64$, $p < .001$).
11
12 The addition of ELS scores in the second step of the model was unable to explain a
13
14 significant additional amount of variance in approach-avoidance responses towards angry
15
16 facial expressions ($\Delta R^2 = .00$, $F(1,386) = 1.58$, $p = .21$). However, the addition of the
17
18 interaction term (standardised ELS scores multiplied by mental illness) in the third step of the
19
20 model was able to explain a significant amount of additional variance in approach-avoidance
21
22 responses towards angry facial expressions ($\Delta R^2 = .01$, $F(1,385) = 5.40$, $p = .02$). Due to the
23
24 observed inter-relationships between ELS, depression and anxiety in the present sample,
25
26 variance inflation factor (VIF) values were used to examine multicollinearity between the
27
28 predictor variables. All predictors showed VIF values below 10, indicating that the extent of
29
30 multicollinearity was acceptable (Field, 2005).
31
32
33
34

35 Examination of the beta weights (Table 3) revealed that the interaction between ELS
36
37 and mental illness predicted a significant additional amount of the variance in approach-
38
39 avoidance of angry facial expressions. To examine this interaction, the correlation between
40
41 ELS and approach-avoidance of angry facial expressions was examined in participants with
42
43 no evidence of mental illness, and in participants with evidence of mental illness. It was
44
45 found that, amongst participants with no evidence of mental illness, higher ELS scores were
46
47 significantly associated with *reduced* avoidance of angry facial expressions ($r(159) = .20$, $p =$
48
49 $.01$, $[0.02, 0.35]$; Figure 3). In contrast, there was no significant relationship between ELS
50
51 scores and approach-avoidance of angry facial expressions in those with evidence of mental
52
53 illness ($r(230) = 0.01$, $p = .92$, $[-0.14, 0.15]$; Figure 3). In addition, examination of the beta
54
55 weights showed that female participants were significantly more avoidant than male
56
57
58
59
60

1
2
3 participants in their responses to angry facial expressions, whilst younger participants were
4
5 significantly more avoidant of these expressions than older participants. In light of the
6
7 significant gender differences in depression and anxiety scores (Table 1), and the significant
8
9 interaction of ELS and mental illness on approach-avoidance of angry facial expressions, an
10
11 interaction term for ELS, mental illness and participant gender was created and entered as an
12
13 additional step in the regression model. There was no significant effect of this interaction
14
15 term on approach-avoidance of angry facial expressions (Supplementary Table 2).
16
17
18
19
20

21
22 *[FIGURE 3 HERE]*
23
24

25 **Early life stress and approach-avoidance of happy and neutral facial expressions**

26
27 The hierarchical regression model was then used to examine the effect of early life
28
29 stress on approach-avoidance responses to happy facial expressions, and approach-avoidance
30
31 responses to neutral facial expressions (Table 3). The control variables of participants'
32
33 gender, age, mental illness, anxiety scores (HADS-A) and depression scores (HADS-D) were
34
35 entered in the first step of the regression model, whilst scores on the measure of ELS (CATS)
36
37 were entered in the second step of the model, and the interaction term of ELS by mental
38
39 illness was entered in the third step.
40
41
42

43
44 The final regression model predicted a significant amount of the variance in approach-
45
46 avoidance responses towards happy facial expressions ($R^2 = .13$, $F(7, 382) = 7.93$, $p < .001$)
47
48 and towards neutral facial expressions ($R^2 = .12$, $F(7, 384) = 7.40$, $p < .001$). The addition of
49
50 ELS scores in the second step of the model was unable to explain a significant additional
51
52 amount of variance in approach-avoidance responses towards happy ($\Delta R^2 = .00$, $F(1,383) =$
53
54 0.33 , $p = .57$) or neutral ($\Delta R^2 = .00$, $F(1,385) = 0.03$, $p = .88$) facial expressions. Similarly,
55
56 the addition of the interaction term (standardised ELS scores multiplied by mental illness) in
57
58 the third step of the model was also unable to explain a significant amount of additional
59
60

1
2
3 variance in approach-avoidance responses towards happy facial expressions ($\Delta R^2 = .00$,
4
5 $F(1,382) = 1.16, p = .28$) or neutral facial expressions ($\Delta R^2 = .00, F(1,384) = 0.04, p = .83$).
6
7

8 Examination of the beta weights (Table 3) revealed that, compared to male
9
10 participants, female participants were less avoidant of happy facial expressions and more
11
12 avoidant of neutral facial expressions. Older age was associated with reduced avoidance of
13
14 both happy and neutral facial expressions. In line with previous research, higher depression
15
16 scores were significantly associated with increased avoidance of happy facial expressions.
17
18 Neither ELS scores nor the interaction between ELS scores and mental illness were
19
20 significantly related to approach-avoidance responses to happy or neutral facial expressions.
21
22
23
24
25
26
27
28
29
30
31

32 [TABLE 3 HERE]
33
34

35 **Summary of findings**

36 Taking the analyses together, there was evidence for an effect of participant gender on
37
38 responses to emotional facial expressions, such that female participants were more avoidant
39
40 than male participants when responding to angry and neutral facial expressions, but less
41
42 avoidant when responding to happy facial expressions. In addition, older age was associated
43
44 with reduced avoidance of all emotional facial expressions (angry, happy and neutral). There
45
46 was an emotion-specific effect of depression scores, such that higher depression scores were
47
48 associated with greater avoidance of happy facial expressions, but were not associated with
49
50 avoidance responses to angry or neutral expressions. Crucially, in mentally healthy
51
52 participants, higher levels of ELS predicted *reduced* avoidance of angry facial expressions.
53
54 This relationship was not present amongst participants with experience of mental illness.
55
56
57
58
59
60

Discussion

This study investigated the effects of ELS on adults' self-reported approach-avoidance tendencies towards photographs of angry, happy and neutral facial expressions. Contrary to predictions, higher ELS scores were associated with reduced self-reported avoidance of angry facial expressions. This effect was only present amongst participants with no evidence of mental illness; amongst participants with experience of mental illness, there was no significant relationship between ELS scores and approach-avoidance of angry facial expressions. Experiences of ELS were not associated with approach-avoidance tendencies towards happy or neutral facial expressions. It was also found that participant gender and age were associated with approach-avoidance tendencies towards emotional facial expressions; female participants were more avoidant of angry and neutral facial expressions, and less avoidant of happy facial expressions, than male participants, whilst older participant age was associated with reduced avoidance of angry, happy and neutral facial expressions. Higher depression scores on the HADS-D were associated with increased avoidance of happy facial expressions, but were unrelated to approach-avoidance responses to angry or neutral facial expressions.

In this study with adult participants, higher levels of ELS did not predict increased avoidance of angry facial expressions. This is in contrast to previous studies with children in which ELS was associated with hyper-sensitivity to the emotion of anger (da Silva Ferreira, Crippa, & Osorio, 2014), as demonstrated by maltreated children's ability to detect anger with fewer perceptual and facial cues (Pollak et al., 2009; Pollak & Sinha, 2002). Furthermore, the finding that higher levels of ELS were not associated with increases in self-reported avoidance appears to contradict the suggestion that ELS-related changes in responsivity to anger serve the purpose of facilitating increased avoidance of danger (McCrory & Viding, 2015; Pollak et al., 2009).

1
2
3 It is interesting to note that the present study found some evidence for the opposite
4 pattern of responsivity; higher levels of ELS were associated with *reduced* self-reported
5 avoidance of angry facial expressions amongst individuals with no evidence of mental illness.
6
7 Given the novelty of these findings, a thorough understanding of their significance requires
8 additional investigation. However, one highly speculative explanation for the effects is the
9
10 notion that reduced avoidance in response to angry facial expressions could represent a
11
12 marker of resilience or stress inoculation, following manageable levels of ELS (Ashokan et
13
14 al., 2016; Ellis, Bianchi, Griskevicius & Frankenhuis, 2017). This explanation could suggest
15
16 that those who had experienced moderate levels of ELS would be best equipped to deal with
17
18 psychosocial challenge (such as how to respond to an angry facial expression). Unlike those
19
20 with very low levels of ELS, these individuals have encountered and learned to manage
21
22 stressful situations, yet they have not been exposed to the more severe levels of stress
23
24 experienced by those who go on to develop mental health conditions.
25
26
27
28
29
30
31
32

33 Indeed, on a conceptual level, the findings align with work by Yamamoto et al.
34
35 (2017), who examined the effects of ELS on mentally healthy participants. They found that
36
37 experience of ELS was associated with increased amygdala activation during a sad mood
38
39 induction, and that this increased amygdala activation was in turn associated with lower
40
41 scores on a measure of depression symptomology. Yamamoto et al. (2017) suggested that the
42
43 ELS-related activity they observed in the amygdala represented a marker of resilience, rather
44
45 than of vulnerability, to future mental illness. Similarly, the present finding of a pattern of
46
47 responsivity which varies according to mental health status accords with work by Saleh et al.
48
49 (2017), who found that the relationship between ELS and processing speed varied according
50
51 to levels of depression symptomology.
52
53
54
55

56 Despite the potential value of a resilience narrative in explaining the present findings,
57
58 it is acknowledged that there was no evidence of a relationship between responses to angry
59
60

1
2
3 facial expressions and measures of depression and anxiety. This limits the proposed
4
5 explanation for the results; if anxiety and depression are not in themselves associated with
6
7 avoidance, it is unclear why alterations in avoidance responsivity should influence an
8
9 individual's propensity for anxiety and depression. Further work should investigate this
10
11 apparent discrepancy in greater detail.
12
13

14
15 It is interesting to acknowledge the patterns of emotion specificity in the present set of
16
17 findings. The effects of ELS on approach-avoidance responses were specific to angry facial
18
19 expressions, with no evidence for any relationships between ELS scores and responses to
20
21 happy or neutral facial expressions. This specificity is broadly in accordance with previous
22
23 work, which has highlighted a specific role for the emotion of anger in studies of face
24
25 processing following ELS (Maheu et al., 2010; McCrory et al., 2011; Pollak, 2008).
26
27 Similarly, the present study found that the effect of depression scores on approach-avoidance
28
29 was specific to happy facial expressions (with higher depression scores associated with
30
31 increased avoidance of happy expressions); depression scores were unrelated to approach-
32
33 avoidance of angry or neutral facial expressions. This finding contributes to literature
34
35 highlighting a bias away from happy emotional cues amongst people with depression
36
37 (Bourke, Douglas & Porter, 2010; Leppanen, 2006), and extends work which found that those
38
39 with depression are more likely than people without depression to perceive happy expressions
40
41 as neutral rather than happy (Gur et al., 1992; Surguladze et al., 2004).
42
43
44
45
46

47 The present work also found significant effects of participant gender and age on
48
49 approach-avoidance tendencies. Specifically, female participants were more likely to avoid
50
51 angry and neutral facial expressions than male participants, an effect which was independent
52
53 of levels of ELS. This is in partial accordance with Miller et al.'s (2013) finding that female
54
55 participants kept a greater simulated distance between themselves and others than male
56
57 participants, irrespective of the others' emotional expressions. The present finding of a
58
59
60

1
2
3 relationship between older age and reduced avoidance of emotional facial expressions was
4 unexpected; to the authors' knowledge, age has not previously been associated with
5 avoidance of emotional facial expressions. Further work is required in order to clarify the
6 relationships between gender, age, and processing of specific emotional expressions.
7
8
9

10
11
12 One of the strengths of this work is its demonstration that early experiences do not
13 have to be extreme to have a long-lasting impact on cognitive processing. However, it is
14 acknowledged that the relatively low levels of ELS in the present sample ($M = 28.42$, out of a
15 maximum score of 152 on the CATS) do give rise to difficulties when attempting to compare
16 the findings to other work in the field, which has typically focused on severe ELS such as
17 abuse that has been documented by authorities (e.g. McCrory et al., 2011; Pollak et al., 2009).
18 Furthermore, whilst steps were taken to identify those likely to be experiencing symptoms of
19 mental illness at the time of the study, clinical interviews were not used, and as such it is
20 possible that some of the participants identified as mentally healthy were experiencing
21 symptoms of mental illness that were not detected by the self-report questionnaires.
22
23
24
25
26
27
28
29
30
31
32
33
34

35
36 The present study was limited by the use of a retrospective self-report measure of ELS
37 and by its cross-sectional design. However, a review of the validity of adult retrospective
38 reports by Hardt and Rutter (2004) concluded that, whilst not all adults will disclose abuse or
39 neglect when asked, cases where abuse or neglect are disclosed are generally reliable.
40 Furthermore, both retrospective reports and cross-sectional designs play an important role in
41 advancing knowledge about the pervasive long-term effects of ELS (e.g. Dannlowski et al.,
42 2012; Du et al., 2016; Edwards, Holden, Felitti, & Anda, 2003). An additional limitation of
43 the present study concerns the task itself. The task was carried out online, and participants
44 were given as much time as they required to respond to each facial expression. As a result, it
45 is possible that some participants were not paying attention to the task. However, the fact that
46 the sample as a whole performed the task as expected suggests that participants did pay
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 attention when responding. It is also acknowledged that the task lacked ecological validity; it
4
5 is not known whether participants' approach and avoidance tendencies towards photographs
6
7 reflect the responses they would make in real life situations. In hindsight, the use of separate
8
9 scales for approach and avoidance was an oversight, as it reduced the ability to compare the
10
11 findings as clearly with other studies which have used one scale ranging from approach to
12
13 avoidance (e.g. Seidel et al., 2010; Derntl et al., 2011). The creation of one scale from the two
14
15 scales was an attempt to remedy this, though it is recognised that this approach has not been
16
17 validated by previous studies.
18
19
20

21
22 In spite of these limitations, this study represents, to the authors' knowledge, the first
23
24 investigation of the relationship between ELS and adults' self-reported approach-avoidance
25
26 of emotional facial expressions. It was found that, in participants with no evidence of mental
27
28 illness, higher ELS scores were associated with *reduced* avoidance of angry facial
29
30 expressions. This finding represents a novel and interesting contribution to the field,
31
32 demonstrating that ELS-related alterations in social cognition can be observed many years
33
34 after the occurrence of ELS, and that these alterations may vary according to the mental
35
36 health of the individual.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- Ashokan, A., Sivasubramanian, M. & Mitra, R. (2016). Seeding stress resilience through inoculation. *Neural Plasticity*, Article ID 4928081, doi: <https://doi.org/10.1155/2016/4928081>
- Bjelland, I., Dahl, A. A., Haug, T. T., & Neckelmann, D. (2002). The validity of the Hospital Anxiety and Depression Scale - An updated literature review. [Review]. *Journal of Psychosomatic Research*, 52(2), 69-77. doi: 10.1016/s0022-3999(01)00296-3
- Bourke, C., Douglas, K. & Porter, R. (2010). Processing of facial emotion expression in major depression: A review. *Australian and New Zealand Journal of Psychiatry*, 44(8), 681-696. doi: 10.3109/00048674.2010.496359
- Cooke, B. M., & Weathington, J. M. (2014). Human and animal research into sex-specific effects of child abuse. *Hormones and Behavior*, 65(4), 416-426. doi: 10.1016/j.yhbeh.2014.03.004
- da Silva Ferreira, G. C., Crippa, J. A. S., & Osorio, F. D. (2014). Facial emotion processing and recognition among maltreated children: a systematic literature review. *Frontiers in Psychology*, 5. doi: 10.3389/fpsyg.2014.01460
- Dannlowski, U., Stuhrmann, A., Beutelmann, V., Zwanzger, P., Lenzen, T., Grotegerd, D., . . . Kugel, H. (2012). Limbic scars: long-term consequences of childhood maltreatment revealed by functional and structural magnetic resonance imaging. *Biol Psychiatry*, 71(4), 286-293. doi: 10.1016/j.biopsych.2011.10.021
- Derntl, B., Seidel, E. M., Eickhoff, S. B., Kellermann, T., Gur, R. C., Schneider, F., & Habel, U. (2011). Neural correlates of social approach and withdrawal in patients with major depression. *Social Neuroscience*, 6(5-6), 482-501. doi: 10.1080/17470919.2011.579800

Running head: EARLY LIFE STRESS AND REDUCED AVOIDANCE OF THREAT 20

- 1
2
3 Di Iorio, C. R., Carey, C. E., Michalski, L. J., Corral-Frias, N. S., Conley, E. D., Hariri, A. R.,
4
5 & Bogdan, R. (2017). Hypothalamic-pituitary-adrenal axis genetic variation and early
6
7 stress moderates amygdala function. *Psychoneuroendocrinology*, *80*, 170-178. doi:
8
9 10.1016/j.psyneuen.2017.03.016
10
11
12 Doretto, V., & Scivoletto, S. (2018). Effects of Early Neglect Experience on Recognition and
13
14 Processing of Facial Expressions: A Systematic Review. *Brain Sciences*, *8*(1), 10.
15
16
17 Du, L., Wang, J. J., Meng, B., Yong, N., Yang, X. Y., Huang, Q. L., . . . Hu, H. (2016). Early
18
19 life stress affects limited regional brain activity in depression. *Scientific Reports*, *6*.
20
21 doi: 10.1038/srep25338
22
23
24 Ebner, N. C., Riediger, M., & Lindenberger, U. (2010). FACES-A database of facial
25
26 expressions in young, middle-aged, and older women and men: Development and
27
28 validation. *Behavior Research Methods*, *42*(1), 351-362. doi: 10.3758/brm.42.1.351
29
30
31 Edwards, V. J., Holden, G. W., Felitti, V. J., & Anda, R. F. (2003). Relationship between
32
33 multiple forms of childhood maltreatment and adult mental health in community
34
35 respondents: Results from the adverse childhood experiences study. *American Journal*
36
37 *of Psychiatry*, *160*(8), 1453-1460. doi: 10.1176/appi.ajp.160.8.1453
38
39
40 Ellis, B.J., Bianchi, J., Griskevicius, V. & Frankenhuis, W.E. (2017). Beyond risk and
41
42 protective factors: An adaptation-based approach to resilience. *Perspectives on*
43
44 *Psychological Science*, *12*(4), 561-587. doi: 10.1177/1745691617693054
45
46
47 Field, A. (2005). *Discovering statistics using SPSS* (2 ed.). London: Sage Publications Ltd.
48
49
50 Gibb, B. E., Schofield, C. A., & Coles, M. E. (2009). Reported History of Childhood Abuse
51
52 and Young Adults' Information-Processing Biases for Facial Displays of Emotion.
53
54 *Child Maltreatment*, *14*(2), 148-156. doi: 10.1177/1077559508326358
55
56
57
58
59
60

- 1
2
3 Gur, R. C., Erwin, R. J., Gur, R. E., Zvil, A. S., Heimberg, C., & Kraemer, H. C. (1992).
4
5 Facial Emotion Discrimination .2. Behavioral Findings in Depression. *Psychiatry*
6
7 *Research*, 42(3), 241-251. doi:Doi 10.1016/0165-1781(92)90116-K
8
9
10 Hardt, J., & Rutter, M. (2004). Validity of adult retrospective reports of adverse childhood
11
12 experiences: review of the evidence. *Journal of Child Psychology and Psychiatry*,
13
14 45(2), 260-273. doi: 10.1111/j.1469-7610.2004.00218.x
15
16
17 Hayes, A. F. (2013). *Introduction to mediation, moderation, and conditional process*
18
19 *analysis: A regression-based approach*. New York, NY, US: Guilford Press.
20
21
22 Heim, C., & Nemeroff, C. B. (2001). The role of childhood trauma in the neurobiology of
23
24 mood and anxiety disorders: Preclinical and clinical studies. *Biol Psychiatry*, 49(12),
25
26 1023-1039. doi: Doi 10.1016/S0006-3223(01)01157-X
27
28
29 Kessler, R. C., Angermeyer, M., Anthony, J. C., De Graaf, R., Demyttenaere, K., Gasquet, I.,
30
31 ... Ustün, T. B. (2007). Lifetime prevalence and age-of-onset distributions of mental
32
33 disorders in the World Health Organization's World Mental Health Survey
34
35 Initiative. *World psychiatry : official journal of the World Psychiatric Association*
36
37 *(WPA)*, 6(3), 168-76. doi: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2174588>
38
39
40 Kessler, R. C., McLaughlin, K. A., Green, J. G., Gruber, M. J., Sampson, N. A., Zaslavsky,
41
42 A. M., . . . Williams, D. R. (2010). Childhood adversities and adult psychopathology
43
44 in the WHO World Mental Health Surveys. [Article]. *British Journal of Psychiatry*,
45
46 197(5), 378-385. doi: 10.1192/bjp.bp.110.080499
47
48
49 Leppanen, J. M. (2006). Emotional information processing in mood disorders: a review of
50
51 behavioral and neuroimaging findings. *Current Opinion in Psychiatry*, 19(1), 34-39.
52
53
54 Maheu, F. S., Dozier, M., Guyer, A. E., Mandell, D., Peloso, E., Poeth, K., . . . Ernst, M.
55
56 (2010). A preliminary study of medial temporal lobe function in youths with a history
57
58
59
60

1
2
3 of caregiver deprivation and emotional neglect. *Cognitive Affective & Behavioral*
4
5 *Neuroscience*, 10(1), 34-49. doi: 10.3758/Cabn.10.1.34
6
7

8 Malinosky-Rummell, R., & Hansen, D. J. (1993). Long-term consequences of childhood
9
10 physical abuse. [Article]. *Psychological Bulletin*, 114(1), 68-79. doi: 10.1037//0033-
11
12 2909.114.1.68
13

14 McCrory, E. J., De Brito, S. A., Kelly, P. A., Bird, G., Sebastian, C. L., Mechelli, A., . . .
15
16 Viding, E. (2013). Amygdala activation in maltreated children during pre-attentive
17
18 emotional processing. *British Journal of Psychiatry*, 202(4), 269-276. doi:
19
20 10.1192/bjp.bp.112.116624
21
22

23
24 McCrory, E. J., De Brito, S. A., Sebastian, C. L., Mechelli, A., Bird, G., Kelly, P. A., &
25
26 Viding, E. (2011). Heightened neural reactivity to threat in child victims of family
27
28 violence. *Current Biology*, 21(23), R947-R948.
29

30
31 McCrory, E. J., & Viding, E. (2015). The theory of latent vulnerability: Reconceptualizing
32
33 the link between childhood maltreatment and psychiatric disorder. *Development and*
34
35 *Psychopathology*, 27(2), 493-505. doi: 10.1017/s0954579415000115
36
37

38 McLaughlin, K. A., Sheridan, M. A., & Lambert, H. K. (2014). Childhood adversity and
39
40 neural development: Deprivation and threat as distinct dimensions of early
41
42 experience. *Neuroscience and Biobehavioral Reviews*, 47, 578-591. doi:
43
44 10.1016/j.neubiorev.2014.10.012
45
46

47 Meichenbaum, D. & Cameron, R. (1989). Stress inoculation training. In *Stress Reduction*
48
49 *and Prevention*, D. Meichenbaum & M. Jaremko, Eds., pp. 115-154, New York,
50
51 USA: Springer.
52

53
54 Miller, H. C., Chabriac, A. S., & Molet, M. (2013). The Impact of Facial Emotional
55
56 Expressions and Sex on Interpersonal Distancing as Evaluated in a Computerized
57
58
59
60

- 1
2
3 Stop-Distance Task. *Canadian Journal of Experimental Psychology-Revue*
4
5 *Canadienne De Psychologie Experimentale*, 67(3), 188-194. doi: 10.1037/a0030663
6
7
8 Perry, R., & Sullivan, R. M. (2014). Neurobiology of Attachment to an Abusive Caregiver:
9
10 Short-Term Benefits and Long-Term Costs. *Developmental Psychobiology*, 56(8),
11
12 1626-1634. doi: 10.1002/dev.21219
13
14
15 Pollak, S. D. (2008). Mechanisms Linking Early Experience and the Emergence of Emotions:
16
17 Illustrations From the Study of Maltreated Children. *Current Directions in*
18
19 *Psychological Science*, 17(6), 370-375. doi: 10.1111/j.1467-8721.2008.00608.x
20
21
22 Pollak, S. D., & Kistler, D. J. (2002). Early experience is associated with the development of
23
24 categorical representations for facial expressions of emotion. *Proceedings of the*
25
26 *National Academy of Sciences of the United States of America*, 99(13), 9072-9076.
27
28 doi: 10.1073/pnas.142165999
29
30
31 Pollak, S. D., Messner, M., Kistler, D. J., & Cohn, J. F. (2009). Development of perceptual
32
33 expertise in emotion recognition. *Cognition*, 110(2), 242-247. doi:
34
35 10.1016/j.cognition.2008.10.010
36
37
38 Pollak, S. D., & Sinha, P. (2002). Effects of early experience on children's recognition of
39
40 facial displays of emotion. *Developmental Psychology*, 38(5), 784-791. doi:
41
42 10.1037//0012-1649.38.5.784
43
44
45 Rainecki, C., Sarro, E., Rincon-Cortes, M., Perry, R., Boggs, J., Holman, C. J., . . . Sullivan, R.
46
47 M. (2015). Paradoxical Neurobehavioral Rescue by Memories of Early-Life Abuse:
48
49 The Safety Signal Value of Odors Learned during Abusive Attachment.
50
51 *Neuropsychopharmacology*, 40(4), 906-914. doi: 10.1038/npp.2014.266
52
53
54 Redlich, R., Stacey, D., Opel, N., Grotegerd, D., Dohm, K., Kugel, H., . . . Dannlowski, U.
55
56 (2015). Evidence of an IFN-gamma by early life stress interaction in the regulation of
57
58
59
60

1
2
3 amygdala reactivity to emotional stimuli. *Psychoneuroendocrinology*, 62, 166-173.

4
5 doi: 10.1016/j.psyneuen.2015.08.008

6
7 Saleh, A., Potter, G., McQuoid, D., Boyd, B., Turner, R., MacFall, J., & Taylor, W. (2017).

8
9 Effects of early life stress on depression, cognitive performance and brain

10
11 morphology. *Psychological Medicine*, 47(1), 171-181.

12
13 doi:10.1017/S0033291716002403

14
15 Sanders, B., & Becker-Lausen, E. (1995). The Measurement of Psychological Maltreatment -

16
17 Early Data on the Child-Abuse and Trauma Scale. *Child Abuse & Neglect*, 19(3), 315-

18
19 323. doi: Doi 10.1016/S0145-2134(94)00131-6

20
21 Seidel, E. M., Habel, U., Finkelmeyer, A., Schneider, F., Gur, R. C., & Derntl, B. (2010).

22
23 Implicit and explicit behavioral tendencies in male and female depression. *Psychiatry*

24
25 *Research*, 177(1-2), 124-130. doi: 10.1016/j.psychres.2010.02.001

26
27 Seidel, E. M., Habel, U., Kirschner, M., Gur, R. C., & Derntl, B. (2010). The Impact of Facial

28
29 Emotional Expressions on Behavioral Tendencies in Women and Men. *Journal of*

30
31 *Experimental Psychology-Human Perception and Performance*, 36(2), 500-507. doi:

32
33 10.1037/a0018169

34
35 Southwick, S.M. & Charney, D.S. (2012). The science of resilience: Implications for the

36
37 prevention and treatment of depression. *Science*, 338(6103), 79-82, doi:

38
39 10.1126/science.1222942

40
41 Surguladze, S. A., Young, A. W., Senior, C., Brebion, G., Travis, M. J., & Phillips, M. L.

42
43 (2004). Recognition accuracy and response bias to happy and sad facial expressions in

44
45 patients with major depression. *Neuropsychology*, 18(2), 212-218. doi:10.1037/0894-

46
47 4105.18.2.212

- 1
2
3 Suzuki, A., Poon, L., Kumari, V., & Cleare, A. J. (2015). Fear Biases in Emotional Face
4
5 Processing Following Childhood Trauma as a Marker of Resilience and Vulnerability
6
7 to Depression. *Child Maltreatment*, 20(4), 240-250. doi:10.1177/1077559515600781
8
9
10 Yamamoto, T., Toki, S., Siegle, G. J., Takamura, M., Takaishi, Y., Yoshimura, S., . . .
11
12 Yamawaki, S. (2017). Increased amygdala reactivity following early life stress: a
13
14 potential resilience enhancer role. *Bmc Psychiatry*, 17. doi:ARTN 2710.1186/s12888-
15
16 017-1201-x
17
18
19
20 Zigmond, A. S., & Snaith, R. P. (1983). The Hospital Anxiety and Depression Scale. *Acta*
21
22 *Psychiatr Scand*, 67(6), 361-370. doi: DOI 10.1111/j.1600-0447.1983.tb09716.x
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Tables

Table 1

Scores on demographic variables for male and female participants

Variable	Participants			<i>p</i>	95% CI
	All (<i>n</i> = 395)	Male (<i>n</i> = 160)	Female (<i>n</i> = 235)		
Age	23.25 (8.92)	22.45 (8.48)	23.80 (9.18)	.14	[-3.08, 0.53]
ELS	28.42 (18.22)	27.41 (15.39)	29.11 (19.93)	.34	[-5.25, 1.67]
Anxiety	8.23 (4.36)	7.54 (4.16)	8.69 (4.44)	.009**	[-2.01, -0.37]
Depression	4.11 (3.49)	4.68 (3.42)	3.73 (3.49)	.007**	[0.29, 1.64]

Note. Values inside brackets indicate standard deviation of the mean. Independent *t* tests used for comparison of means between male and female participants.

ELS measured with the Child Abuse and Trauma Scale (Sanders & Becker-Lausen, 1995), anxiety and depression measured with the Hospital Anxiety and Depression Scale (Zigmond & Snaith, 1983). ** = $p < .01$.

Table 2

Scores on demographic variables for participants with and without evidence of mental illness

Variable	Participants		<i>p</i>	95% CI
	With mental illness (<i>n</i> = 234)	Without mental illness (<i>n</i> = 161)		
Age	23.38 (8.89)	23.07 (8.99)	0.77	[-2.14, 1.62]
ELS	34.48 (19.62)	19.62 (11.21)	0.001**	[-17.82, -11.60]
Anxiety	10.82 (3.59)	4.46 (1.98)	0.001**	[-6.92, -5.80]
Depression	5.41 (3.68)	2.23 (2.04)	0.001**	[-3.76, -2.54]

Note. Values inside brackets indicate standard deviation of the mean. Independent *t* tests used for comparison of means between participants with and without evidence of mental illness (measured as self-reported diagnosis and/or clinical score on one or both HADS subscales). ELS measured with the Child Abuse and Trauma Scale (Sanders & Becker-Lausen, 1995), anxiety and depression measured with the Hospital Anxiety and Depression Scale (Zigmond & Snaith, 1983).

** = $p < .01$.

Running head: EARLY LIFE STRESS AND REDUCED AVOIDANCE OF THREAT

Table 3

Predictors of approach-avoidance responses to angry, happy and neutral facial expressions

Expression	Predictor	Step 1				Step 2				Step 3			
		<i>B</i>	<i>SE B</i>	95% CI	β	<i>B</i>	<i>SE B</i>	95% CI	β	<i>B</i>	<i>SE B</i>	95% CI	β
Angry	Gender	-13.88	3.39	[-20.31, -7.69]	-0.22**	-14.04	3.37	[-20.40, -7.99]	-0.22**	-13.83	3.37	[-20.34, -7.53]	-0.22**
	Age	0.64	0.20	[0.25, 1.00]	0.18**	0.63	0.20	[0.23, 1.00]	0.18**	0.61	0.20	[0.21, 0.99]	0.18**
	Anxiety	0.25	0.57	[-0.77, 1.45]	0.04	0.14	0.59	[-0.94, 1.34]	0.02	0.21	0.58	[-0.91, 1.45]	0.03
	Depression	0.20	0.58	[-0.89, 1.31]	0.02	0.03	0.59	[-1.07, 1.09]	0.00	0.04	0.59	[-1.05, 1.11]	0.00
	Mental illness	-3.93	4.43	[-12.84, 5.59]	-0.06	-4.52	4.45	[-13.25, 4.77]	-0.07	-8.23	4.85	[-17.47, 1.44]	-0.13
	ELS					2.22	1.81	[-1.54, 5.79]	0.07	10.26	4.34	[1.13, 18.49]	0.33*
	ELS*MI [†]									-9.93	4.67	[-18.75, -0.20]	-0.27*
	<i>R</i> ²	.08***				.08***				.09***			
Happy	Gender	5.94	2.85	[0.09, 11.88]	0.10*	6.02	2.84	[0.29, 12.02]	0.10*	6.12	2.84	[0.17, 12.35]	0.11*
	Age	0.49	0.15	[0.22, 0.78]	0.15**	0.49	0.15	[0.22, 0.79]	0.15**	0.49	0.15	[0.21, 0.77]	0.15**
	Anxiety	0.63	0.53	[-0.29, 1.51]	0.10	0.68	0.53	[-0.26, 1.64]	0.10	0.71	0.53	[-0.26, 1.63]	0.11
	Depression	-2.22	0.52	[-3.35, -1.17]	-0.27**	-2.15	0.52	[-3.31, -1.06]	-0.26**	-2.15	0.52	[-3.31, -1.11]	-0.26**
	Mental illness	-6.90	4.11	[-15.51, 1.72]	-0.12	-6.67	4.16	[-15.01, 1.58]	-0.12	-8.22	4.43	[-17.28, 0.71]	-0.14
	ELS					-0.90	1.65	[-4.26, 2.28]	-0.03	2.46	3.77	[-5.01, 10.82]	0.09
	ELS*MI [†]									-4.14	4.16	[-12.23, 2.94]	-0.12
	<i>R</i> ²	.12***				.12***				.13***			
Neutral	Gender	-6.79	3.18	[-13.22, -0.87]	-0.11*	-6.77	3.18	[-13.35, -0.74]	-0.11*	-6.75	3.18	[-13.25, -0.76]	-0.11*
	Age	1.02	0.19	[0.63, 1.43]	0.30**	1.02	0.19	[0.65, 1.42]	0.30**	1.02	0.19	[0.64, 1.42]	0.30**
	Anxiety	-0.39	0.52	[-1.41, 0.67]	-0.06	-0.38	0.53	[-1.41, 0.73]	-0.06	-0.37	0.53	[-1.41, 0.74]	-0.05
	Depression	-0.46	0.51	[-1.50, 0.51]	-0.05	-0.44	0.52	[-1.49, 0.58]	-0.05	-0.44	0.52	[-1.47, 0.58]	-0.05
	Mental illness	-1.62	4.04	[-9.78, 6.17]	-0.03	-1.55	4.08	[-9.65, 6.12]	-0.03	-1.88	4.22	[-9.92, 6.23]	-0.03
	ELS					-0.26	1.71	[-3.51, 2.80]	-0.01	0.44	3.58	[-7.00, 7.03]	0.01
	ELS*MI [†]									-0.87	4.02	[-8.39, 6.79]	-0.02
	<i>R</i> ²	.12***				.12***				.12***			


Note. Positive scores indicate increased approach, negative scores indicate increased avoidance. Each analysis had a slightly different *n* due to the removal of

cases with residuals > +/- 3 standard deviations, angry: *n* = 393 (2 cases removed), happy *n* = 390 (5 cases removed), neutral *n* = 392 (3 cases removed).

Running head: EARLY LIFE STRESS AND REDUCED AVOIDANCE OF THREAT

1 Gender refers to participant's gender and is coded as 0 = male, 1 = female. ELS = early life stress, measured using the Child Abuse and Trauma Scale (Sanders
2 & Becker-Lausen, 1995), note that scores on this variable were standardised prior to inclusion in the model. Anxiety and depression measured using the
3
4 Hospital Anxiety and Depression Scale (HADS; Zigmond & Snaith, 1983). * = $p < .05$, ** = $p < .01$, *** = $p < .001$, †MI = Mental illness, measured as self-
5
6 reported diagnosis and/or a score above the clinical cut-off for one or both of the HADS scales, coded as mental illness = 1, no mental illness = 0.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46


For Review Only



18 Task stimuli. Example of a web page viewed by participants. Note in the actual task the image of the facial
19 expression was positioned above the "sliders".
20

21 227x81mm (120 x 120 DPI)
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Mean approach-avoidance scores across all participants (n = 395) in response to angry, happy and neutral facial expressions. Positive scores indicate a tendency to approach whilst negative scores indicate a tendency to avoid. Error bars display the standard error of the mean.

128x94mm (150 x 150 DPI)

A. Participants with mental illness



B. Participants without mental illness



Scatter plots depicting the correlation between early life stress and approach-avoidance responses to angry facial expressions in (A) participants with evidence of mental illness, and (B) participants without evidence of mental illness. Positive scores on the y axis indicate greater approach, whilst negative scores indicate greater avoidance.