



This is a repository copy of *Deformations of Saito-Kurokawa type and the Paramodular Conjecture*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/150254/>

Version: Accepted Version

Article:

Berger, T. orcid.org/0000-0002-5207-6221 and Klosin, K. (2020) Deformations of Saito-Kurokawa type and the Paramodular Conjecture. *American Journal of Mathematics*, 142 (6). pp. 1821-1875. ISSN 0002-9327

<https://doi.org/10.1353/ajm.2020.0052>

© 2020 Johns Hopkins University Press. This is an author-produced version of a paper subsequently published in *American Journal of Mathematics*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

DEFORMATIONS OF SAITO-KUROKAWA TYPE AND THE PARAMODULAR CONJECTURE

TOBIAS BERGER¹ AND KRZYSZTOF KLOSIN²
WITH AN APPENDIX BY CRIS POOR³, JERRY SHURMAN⁴, AND DAVID S.
YUEN⁵

ABSTRACT. We study short crystalline, minimal, essentially self-dual deformations of a mod p non-semisimple Galois representation $\bar{\sigma}$ with $\bar{\sigma}^{\text{ss}} = \chi^{k-2} \oplus \rho \oplus \chi^{k-1}$, where χ is the mod p cyclotomic character and ρ is an absolutely irreducible reduction of the Galois representation ρ_f attached to a cusp form f of weight $2k - 2$. We show that if the Bloch-Kato Selmer groups $H_f^1(\mathbf{Q}, \rho_f(1 - k) \otimes \mathbf{Q}_p/\mathbf{Z}_p)$ and $H_f^1(\mathbf{Q}, \rho(2 - k))$ have order p , and there exists a characteristic zero absolutely irreducible deformation of $\bar{\sigma}$ then the universal deformation ring is a dvr. When $k = 2$ this allows us to establish the modularity part of the Paramodular Conjecture in cases when one can find a suitable congruence of Siegel modular forms. As an example we prove the modularity of an abelian surface of conductor 731. When $k > 2$, we obtain an $R^{\text{red}} = T$ theorem showing modularity of all such deformations of $\bar{\sigma}$.

1. INTRODUCTION

In analogy with the Taniyama-Shimura Conjecture (proved by Wiles et al.), in the 1980s Yoshida proposed a conjecture postulating that abelian surfaces over \mathbf{Q} should correspond to Siegel modular forms. The most important progress on this conjecture is due to Tilouine and Pilloni ([Til06, Pil12]). Their results show p -adic modularity of abelian surfaces A (see also forthcoming work of Boxer, Calegari, Gee, and Pilloni on modularity, see [BCGP18]) under the assumption that the residual Galois representation $A(\overline{\mathbf{Q}})[p]$ has image containing $\text{PSp}_4(\mathbf{F}_p)$ and, more significantly, is itself modular. The latter assumption essentially calls for an analogue of Serre's conjecture for $\bar{\sigma}_A$, which at present appears out of reach.

Yoshida himself proved some cases of his conjecture in [Yos80] when $\text{End}_{\mathbf{Q}}(A) \neq \mathbf{Z}$. In 2014, Brumer and Kramer proposed a refinement of Yoshida's Conjecture by specifying the level of the Siegel modular form [BK14]. This is sometimes referred to as the *Paramodular Conjecture*.

Conjecture 1.1 (Brumer - Kramer). *For every isogeny class of abelian surfaces A/\mathbf{Q} of conductor N with $\text{End}_{\mathbf{Q}} A = \mathbf{Z}$ there exists a weight 2 Siegel modular form F , which is not in the space spanned by the Saito-Kurokawa lifts, has level $K(N)$ and rational eigenvalues, where $K(N)$ is the*

The first author's research was supported by an LMS Research in Pairs Grant and the EPSRC Grant EP/R006563/1. The second author was supported by a Collaboration for Mathematicians Grant #578231 from the Simons Foundation and by a PSC-CUNY award jointly funded by the Professional Staff Congress and the City University of New York.

paramodular group of level N defined by $K(N) = \gamma M_4(\mathbf{Z}) \gamma^{-1} \cap \mathrm{Sp}_4(\mathbf{Q})$ with $\gamma = \mathrm{diag}[1, 1, 1, N]$. The L -series of A and F should agree and the p -adic representation of $T_p(A) \otimes \mathbf{Q}_p$ should be isomorphic to the one associated to F for any p prime to N where $T_p(A)$ is the p -adic Tate module.

Brumer-Kramer [BK14] and Poor-Yuen [PY15] verified Conjecture 1.1 in several cases by proving the non-existence of an abelian surface for certain (small) conductors N for which there are no non-lift Siegel modular forms of level $K(N)$. When A acquires extra endomorphisms over a quadratic field the paramodularity of A has in some cases been proven using theta lifts, see [JLR12] and [BDPS15].

At the time of writing this article the only genuinely symplectic case (i.e., concerning surfaces whose associated Galois representations are absolutely irreducible and not induced from 2-dimensional representations) of Conjecture 1.1 in which the abelian surface exists that had been fully verified was the case of the abelian surface of conductor 277, see [BPP⁺ar], which now also contains a proof for abelian surfaces of conductors 353 and 587.

In this article we study Galois representations arising from abelian surfaces (as in Conjecture 1.1) with rational torsion. We propose a method to prove modularity of such representations which we carry out here under certain assumptions. This case is not covered by the work of Tilouine and Pilloni et al. It has the advantage of not requiring modularity of $\bar{\sigma}_A$ and replacing this assumption with a construction of congruences between paramodular Saito-Kurokawa lifts and stable Siegel modular forms. In other words we do not have a need for an analogue of Serre's Conjecture. While potentially more accessible, proving the existence of such congruences is still a considerable problem. In this article however we mostly focus on the Galois representation side (deformation theory) of the modularity problem, and make use of results where the required congruence has been constructed by others (mostly by Poor and Yuen). For the purposes of this article Poor, Yuen and Shurman have kindly agreed to prove the existence of such a congruence in the case of forms of paramodular level 731 (see Appendix). See also [PY15] for other examples. A full modularity result would require the existence of such congruences in a more general context, a problem which is a subject of joint work in progress with J. Brown. As a consequence of the results proved in this article we are able to establish the modularity part of the Paramodular Conjecture in new (genuinely symplectic) cases (in particular the first composite level case) and provide a way to verify more, when the appropriate ingredients are supplied.

Let us explain the results and the methods of this paper in more detail. Let $p > 3$ be a prime such that $p \nmid N$. Let A be as in Conjecture 1.1 and suppose that A has a rational p -torsion point, semi-abelian reduction at $\ell \mid N$, and a polarization of degree prime to p . This implies that the semisimplification $\bar{\sigma}_A^{\mathrm{ss}}$ of the residual Galois representation $\bar{\sigma}_A : G_{\mathbf{Q}} \rightarrow \mathrm{GL}_4(\mathbf{F}_p)$ attached to A (i.e., afforded by the p -torsion Galois module $A(\overline{\mathbf{Q}})[p]$) has the form

$$\bar{\sigma}_A^{\mathrm{ss}} \cong 1 \oplus \chi \oplus \rho^{\mathrm{ss}},$$

where χ is the mod p cyclotomic character and ρ is a two-dimensional representation. Assume that ρ is absolutely irreducible (it is automatically odd). Serre's Conjecture (now a theorem of Khare and Wintenberger) implies that ρ arises as a mod p reduction of the Galois representation ρ_f attached to a modular form f of weight 2 and level $\Gamma_0(N)$. If the sign of the functional equation for f is -1 , then $\bar{\sigma}_A^{\text{ss}}$ is the mod p reduction of the Galois representation attached to a weight 2, level $K(N)$ Siegel Hecke eigenform $\text{SK}(f)$ which is the *paramodular Saito-Kurokawa lift* of f . In fact, regardless of the sign of f , $\bar{\sigma}^{\text{ss}}$ arises from a congruence level Saito-Kurokawa lift of f , see section 9.2 for examples.

Our first result shows that one can construct a Galois-invariant lattice L in the space \mathbf{Q}_p^4 with $G_{\mathbf{Q}}$ -action via σ_A with respect to which the residual Galois representation is non-semisimple. More precisely, we can ensure that it is block-upper-triangular with a specific order of the Jordan-Hölder factors on the diagonal $(1, \rho, \chi)$ - this order plays an important role in controlling the deformations (see below). For the modularity argument it is crucial to know that there is only one isomorphism class of such residual representations as long as we require that they be short crystalline, minimally ramified and give rise to desired extensions of the Jordan-Hölder factors of $\bar{\sigma}_A$. The existence of such a lattice can be proved by essentially following the standard method of Ribet (cf. Proposition 2.1 in [Rib76]) adapted to a higher-dimensional setting. The uniqueness of the residual representation is trickier and requires us to study iterated Fontaine-Laffaille extensions as well as to control ramification at the primes dividing N .

We then study short crystalline, minimal and essentially self-dual deformations of the residual representation $\bar{\sigma}_A$. Let R denote the reduced universal deformation ring. Its structure relevant for our purposes can be controlled by two sets of data: its ideals of reducibility $\{I_{\mathcal{P}}\}_{\mathcal{P}}$ corresponding to all possible partitions \mathcal{P} of the set of Jordan-Hölder factors of $\bar{\sigma}_A$ and the quotients $\{R/I_{\mathcal{P}}\}_{\mathcal{P}}$. Roughly speaking, the former control trace-irreducible deformations, while the latter the trace-reducible ones. In the case when $I_{\mathcal{P}}$ is the total ideal of reducibility (i.e., corresponds to the most refined partition), the quotient $R/I_{\mathcal{P}}$ can be shown to be related to the Bloch-Kato Selmer group H_{-1} , where $H_i := H_f^1(\mathbf{Q}, \rho_f(i) \otimes \mathbf{Q}_p/\mathbf{Z}_p)$, by a generalization of the approach used in [BK13]. On the other hand understanding of the ideals of reducibility $I_{\mathcal{P}}$ requires an entirely new approach as the methods used in [BK13] cannot be extended to our current situation.

To overcome this problem we draw on ideas of Bellaïche and Chenevier contained in sections 8 and 9 of their indispensable book [BC09] which has served us as an inspiration on many occasions. There a problem concerning representations with multiple Jordan-Hölder factors is studied in characteristic zero (as opposed to our characteristic p situation). We, too, are able to prove that in our situation (assuming that the divisible Bloch-Kato Selmer group H_0 is of corank at most 1) all the (a priori different) ideals of reducibility in fact coincide and as a consequence we are able to deduce that they are principal. This approach poses a considerable amount of technical

difficulties because, among other things, we are not able to control ramification in the same way as Bellaïche and Chenevier do, one reason being that their proof of the splitting of extensions at ramified primes (see proof of Proposition 8.2.10 in [BC09]) does not extend to characteristic p .

So, while following the general strategy of [BC09] we need a different way of manufacturing the necessary ingredients, and in particular are led to working with a specific minimality condition. Happily this turns out to also be the correct condition for yielding deformations corresponding to Siegel modular forms of squarefree paramodular level (see Proposition 8.5). In some sense section 6, where the ideal of reducibility is studied, comprises the technical heart of the paper.

We show that if H_0 is of corank ≤ 1 and $\#H_{-1} = p$ (that $\#H_{-1} \geq p$ is automatic thanks to the existence of the lattice L), then R is a discrete valuation ring (see Theorem 7.8). It means that the deformation σ_A of $\bar{\sigma}_A$ is the unique characteristic zero deformation. To conclude modularity of A we now need a non-lift Siegel modular form F of weight 2 and level $K(N)$ whose Galois representation σ_F reduces (after semisimplification) to $\bar{\sigma}_A^{\text{ss}}$. This is achieved by exhibiting the existence of a congruence between a candidate F and the paramodular Saito-Kurokawa lift $\text{SK}(f)$ (see section 9 and the Appendix). Irreducibility of σ_F (which we prove) and uniqueness of the non-semisimple residual representation (discussed above) now guarantee that σ_F is a characteristic zero deformation of $\bar{\sigma}_A$. So, we must have $\sigma_A \cong \sigma_F$, proving the modularity of A . This can be viewed as the main application of our method in this paper, stated as Theorem 8.6. If a matching bound on \mathbf{T}/J were available (see next paragraph) our method would in fact be enough to prove a full $R = T$ theorem. We plan to address this problem in a future paper.

Let us note that we prove our results for a residual representation $\bar{\sigma}$ such that $\bar{\sigma}^{\text{ss}} \cong \chi^{k-2} \oplus \chi^{k-1} \oplus \rho$ for any integer $k \geq 2$. The case $k = 2$ (discussed above) is the most interesting because it corresponds to abelian surfaces and thus to Conjecture 1.1. In fact for even $k > 9$ and $N = 1$ we prove a full modularity result $R = \mathbf{T}$ using a congruence result of Brown [Bro11], which provides us with the desired matching bound on \mathbf{T}/J in this case.

Let us now comment on the restrictiveness of our results. The most serious assumptions are the ones on the Selmer groups. It is worth noting that while for H_{-1} we require a specific bound on its order, for H_0 we only require that is of corank 1. In fact, it is important that for our method the corank assumption of the latter group is sufficient since H_0 is infinite if the central L -value of f is zero. It is here that the order of the Jordan-Hölder factors on the diagonal of the residual representation is crucial - a different order could swap the conditions on H_0 and H_{-1} making our theorem empty in this case.

On the other hand the Selmer group H_{-1} is non-critical which poses certain difficulty in computing its order. We overcome this by using Kato's result [Kat04] towards the Main Conjecture of Iwasawa Theory and slightly adapting a control theorem of [SU14] to relate the order to a special value of a p -adic (rather than classical) L -function of f (cf. section 2.3).

The assumptions on the Selmer groups are central to our method and in fact without them R cannot be a dvr.

We also require that p is not a congruence prime for f and that each $\ell \mid N$ satisfies $p \nmid 1 + w_{f,\ell}$, where $w_{f,\ell}$ is the local (at ℓ) Atkin-Lehner sign of f . The first condition allows us to relate the order of $R/I_{\mathcal{P}}$ to the order of H_{-1} , while the latter allows us to prove that the classes in $H^1(\mathbf{Q}, \rho_f \otimes \mathbf{Q}_p/\mathbf{Z}_p)$ are automatically unramified at $\ell \mid N$ (see Proposition 2.7). It is conceivable that both of these conditions could be relaxed, but we do not know of a way to do it. What is important is that these conditions are often satisfied for small primes which has been our goal since it is often the case that abelian surfaces possess rational p -torsion for such primes. Unfortunately, the method to prove principality of the ideal of reducibility requires that p does not divide $d!$, where d is the dimension of the Galois representations (cf. section 1.2 of [BC09]), which forces us to exclude $p = 2, 3$. As for elliptic curves the size of the torsion subgroup of rational points on abelian surfaces is conjectured to be bounded, but so far this has only been proven in special cases. Importantly for us it is known though by work e.g. of Flynn [Fly90] that there are infinitely many abelian surfaces with rational torsion points for certain primes p .

While we hope that this article is only the first half of a larger undertaking of proving a full $R = \mathbf{T}$ theorem for abelian surfaces in Conjecture 1.1 with rational torsion (the second half being the mentioned work in progress on constructing congruences), it is worth noting that even without the matching work on the Hecke side, it allows us to prove modularity of several new examples. In section 9 we work out the details for an abelian surface of conductor 731, and discuss other examples.

The paper is organized as follows. We begin by assembling necessary results concerning Selmer groups in section 2 and define the relevant deformation problem in section 3. The lattice L is constructed in section 4 and the uniqueness of the residual representation is proved in section 5. Section 6 is devoted to the study of the ideals of reducibility $I_{\mathcal{P}}$, while in Section 7 we study the quotients $R/I_{\mathcal{P}}$ and conclude the proof that R is a dvr. In section 8 we prepare the ground for applications to examples of abelian surfaces satisfying our conditions and state the application to the Paramodular Conjecture (Theorem 8.6). Finally, in section 9 we prove the modularity of an abelian surface of conductor $N = 731$ and discuss other examples. We conclude the paper by proving an $R = \mathbf{T}$ theorem in the case of $k > 2$ and $N = 1$ in section 10.

We would like to thank David Savitt for helping us with the proof of Proposition 5.1 for the prime p and Jim Brown, Armand Brumer, Kenneth Kramer, Chris Skinner and Eric Urban for helpful conversations related to the topics of this article. We would also like to thank the anonymous referee for some helpful comments on the manuscript.

2. SELMER GROUPS

For each prime ℓ of \mathbf{Q} we fix embeddings $\overline{\mathbf{Q}} \hookrightarrow \overline{\mathbf{Q}}_{\ell} \hookrightarrow \mathbf{C}$. Let $p > 3$ be a prime. Throughout this paper E will denote a sufficiently large finite

extension of \mathbf{Q}_p , \mathcal{O} its valuation ring with uniformizer ϖ and residue field \mathbf{F} . Let ϵ be the p -adic cyclotomic character. We will write χ for its mod ϖ reduction.

2.1. Definitions and first properties. In this section we define (local and global) Selmer groups which will be in use throughout the paper and recall some of their basic properties - for a more detailed treatment we refer the reader to section 5 of [BK13]. For Σ a finite set of finite places of \mathbf{Q} containing p we write G_Σ for the Galois group of the maximal extension of \mathbf{Q} unramified outside of Σ and infinity. We write $G_{\mathbf{Q}_\ell}$ for the absolute Galois group of \mathbf{Q}_ℓ . Let M be an \mathcal{O} -module with an \mathcal{O} -linear action of $G = G_K$ or G_Σ . We call M a p -adic G -module over \mathcal{O} if one of the following holds:

- (1) M is *finitely generated*, i.e. a finitely generated \mathbf{Z}_p -module and the G -action is continuous for the p -adic topology on M ;
- (2) M is *discrete*, i.e. a torsion \mathbf{Z}_p -module of finite corank (i.e. M is isomorphic as a \mathbf{Z}_p -module to $(\mathbf{Q}_p/\mathbf{Z}_p)^r \oplus M'$ for some $r \geq 0$ and some \mathbf{Z}_p -module M' of finite order) and the G -action on M is continuous for the discrete topology on M ;
- (3) M is a finite-dimensional \mathbf{Q}_p -vector space and the G -action is continuous for the p -adic topology on M .

M is both finitely generated and discrete if and only if it is of finite cardinality.

Given a p -adic G_Σ -module M we assume that we have a finite-singular structure \mathcal{S} on M in the sense of [Wes00], i.e. for each prime $\ell \in \Sigma$ a choice of \mathcal{O} -submodule $H_{f,\mathcal{S}}^1(\mathbf{Q}_\ell, M) \subseteq H^1(\mathbf{Q}_\ell, M)$. We then define two global Selmer groups for M :

$$(2.1) \quad H_{f,\mathcal{S}}^1(\mathbf{Q}, M) = \ker(H^1(G_\Sigma, M) \rightarrow \prod_{\ell \in \Sigma} H^1(\mathbf{Q}_\ell, M)/H_{f,\mathcal{S}}^1(\mathbf{Q}_\ell, M))$$

and a “relaxed” Selmer group (no conditions at primes $\ell \in \Sigma \setminus \{p\}$)

$$H_{\Sigma,\mathcal{S}}^1(\mathbf{Q}, M) := \ker\left(H^1(G_\Sigma, M) \rightarrow H^1(\mathbf{Q}_p, M)/H_{f,\mathcal{S}}^1(\mathbf{Q}_p, M)\right).$$

We consider the following local finite-singular structures $H_{f,\mathcal{S}}^1(\mathbf{Q}_\ell, M)$ (dropping the subscript \mathcal{S} for the place at infinity and at p , where we fix the choice of finite-singular structure):

We always take $H_f^1(\mathbf{Q}_\ell, M) = 0$ for $\ell \mid \infty$.

For $\ell = p$ we define the *crystalline* local finite-singular structure as follows. Let $T \subseteq V$ be a G_K -stable \mathbf{Z}_p -lattice and put $W = V/T$. For $n \geq 1$, put

$$W_n = \{x \in W : \varpi^n x = 0\} \cong T/\varpi^n T.$$

Following Bloch and Kato [BK90] we define $H_f^1(\mathbf{Q}_\ell, V) = \ker(H^1(\mathbf{Q}_\ell, V) \rightarrow H^1(\mathbf{Q}_\ell, B_{\text{crys}} \otimes V))$, denote by $H_f^1(\mathbf{Q}_\ell, T)$ its pullback via the natural map $T \hookrightarrow V$ and let $H_f^1(\mathbf{Q}_\ell, W) = \text{im}(H_f^1(\mathbf{Q}_\ell, V) \rightarrow H^1(\mathbf{Q}_\ell, W))$. Finally, we set $H_f^1(\mathbf{Q}_\ell, W_n)$ to be the inverse image of $H_f^1(\mathbf{Q}_\ell, W)$ under the map $H^1(\mathbf{Q}_\ell, W_n) \rightarrow H_f^1(\mathbf{Q}_\ell, W)$.

For finitely generated p -adic $G_{\mathbf{Q}_p}$ -modules we recall the theory of Fontaine-Laffaille [FL82], following the exposition in [CHT08] Section 2.4.1. Let

$\mathcal{MF}_{\mathcal{O}}$ (“Dieudonné modules”) denote the category of finitely generated \mathcal{O} -modules M together with a decreasing filtration $\mathrm{Fil}^i M$ by \mathcal{O} -submodules which are \mathcal{O} -direct summands with $\mathrm{Fil}^0 M = M$ and $\mathrm{Fil}^{p-1} M = (0)$ and Frobenius linear maps $\Phi^i : \mathrm{Fil}^i M \rightarrow M$ with $\Phi^i|_{\mathrm{Fil}^{i+1} M} = p\Phi^{i+1}$ and $\sum \Phi^i \mathrm{Fil}^i M = M$. They define an exact, fully faithful covariant functor \mathbf{G} of \mathcal{O} -linear categories from $\mathcal{MF}_{\mathcal{O}}$ (in their notation $\mathbb{G}_{\bar{v}}$ and $\mathcal{MF}_{\mathcal{O}, \bar{v}}$) to the category of finitely generated \mathcal{O} -modules with continuous action by $G_{\mathbf{Q}_p}$. Its essential image is closed under taking subquotients, finite direct sums and contains quotients of lattices in short crystalline representations defined as follows: We call V a continuous finite-dimensional $G_{\mathbf{Q}_p}$ -representation over \mathbf{Q}_p *short crystalline* if, for all places $v \mid p$, $\mathrm{Fil}^0 D = D$ and $\mathrm{Fil}^{p-1} D = (0)$ for the filtered vector space $D = (B_{\mathrm{crys}} \otimes_{\mathbf{Q}_p} V)^{G_v}$ defined by Fontaine.

For any p -adic $G_{\mathbf{Q}_p}$ -module M of finite cardinality in the essential image of \mathbf{G} (we will call such modules short crystalline) we define $H_f^1(\mathbf{Q}_p, M)$ as the image of $\mathrm{Ext}_{\mathcal{MF}_{\mathcal{O}}}^1(1_{\mathrm{FD}}, D)$ in $H^1(\mathbf{Q}_p, M) \cong \mathrm{Ext}_{\mathcal{O}[G_{\mathbf{Q}_p}]}^1(1, M)$, where $\mathbf{G}(D) = M$ and 1_{FD} is the unit filtered Dieudonné module defined in Lemma 4.4 of [BK90].

For any complete Noetherian \mathbf{Z}_p -algebra we define a representation $\rho : G_{\mathbf{Q}_p} \rightarrow \mathrm{GL}_n(A)$ to be crystalline if for each Artinian quotient A' of A , $\rho \otimes A'$ lies in the essential image of \mathbf{G} .

Remark 2.1. Note that for short crystalline representations V we gave two different definitions for $H_f^1(\mathbf{Q}_p, W_n)$ (via pullback of $H_f^1(\mathbf{Q}_p, W)$ and via the \mathbf{G} -functor). That these agree follows from the proof of Proposition 2.2 in [DFG04] or [BK13] Lemma 5.3.

For primes $\ell \neq p$ we define the *unramified* local finite-singular structure on any p -adic $G_{\mathbf{Q}_\ell}$ -module M over \mathcal{O} as

$$H_{f, \mathrm{ur}}^1(\mathbf{Q}_\ell, M) = H_{\mathrm{ur}}^1(\mathbf{Q}_\ell, M) = \ker(H^1(\mathbf{Q}_\ell, M) \rightarrow H^1(\mathbf{Q}_{\ell, \mathrm{ur}}, M)),$$

where $\mathbf{Q}_{\ell, \mathrm{ur}}$ is the maximal unramified extension of \mathbf{Q}_ℓ .

Let V be a continuous finite-dimensional $G_{\mathbf{Q}_\ell}$ -representation over \mathbf{Q}_p and $T \subseteq V$ be a $G_{\mathbf{Q}_\ell}$ -stable \mathbf{Z}_p -lattice and put $W = V/T$. Bloch-Kato then define the following *minimal* finite-singular structures on V , T and W :

$$H_{f, \mathrm{min}}^1(\mathbf{Q}_\ell, V) = H_{\mathrm{ur}}^1(\mathbf{Q}_\ell, V),$$

$$H_{f, \mathrm{min}}^1(\mathbf{Q}_\ell, T) = i^{-1} H_f^1(\mathbf{Q}_\ell, V) \text{ for } T \xrightarrow{i} V$$

and

$$H_{f, \mathrm{min}}^1(\mathbf{Q}_\ell, W) = \mathrm{im}(H_f^1(\mathbf{Q}_\ell, V) \rightarrow H^1(\mathbf{Q}_\ell, W)).$$

Remark 2.2. Following [Rub00] Definition 1.3.4 we define $H_{f, \mathrm{min}}^1(\mathbf{Q}_\ell, W_n)$ as the inverse image of $H_{f, \mathrm{min}}^1(\mathbf{Q}_\ell, W)$ under the map $H^1(\mathbf{Q}_\ell, W_n) \rightarrow H^1(\mathbf{Q}_\ell, W)$.

By [Rub00] Lemma 1.3.5 we have $H_{f, \mathrm{min}}^1(\mathbf{Q}_\ell, W) = H_{\mathrm{ur}}^1(\mathbf{Q}_\ell, W)_{\mathrm{div}}$ and if W^{I_ℓ} is divisible we further have $H_{f, \mathrm{min}}^1(\mathbf{Q}_\ell, W) = H_{\mathrm{ur}}^1(\mathbf{Q}_\ell, W)$, $H_{f, \mathrm{min}}^1(\mathbf{Q}_\ell, T) = H_{\mathrm{ur}}^1(\mathbf{Q}_\ell, T)$ and $H_{f, \mathrm{min}}^1(\mathbf{Q}_\ell, W_n) = H_{\mathrm{ur}}^1(\mathbf{Q}_\ell, W_n)$ (for the latter using also [Rub00] Lemma 1.3.8 and its proof).

Lemma 2.3 ([Wes00] Lemma II.3.1). *Let $0 \rightarrow M' \xrightarrow{i} M \xrightarrow{j} M'' \rightarrow 0$ be an exact sequence of p -adic G_Σ -modules over \mathcal{O} . Assume that the modules have been given the induced finite-singular structures, i.e. such that $H_{f,S}^1(\mathbf{Q}_\ell, M'') = j_* H_{f,S}^1(\mathbf{Q}_\ell, M)$ and $H_{f,S}^1(\mathbf{Q}_\ell, M') = i_*^{-1} H_{f,S}^1(\mathbf{Q}_\ell, M)$ for all ℓ .*

Then there is an exact sequence

$$0 \rightarrow H^0(\mathbf{Q}, M') \rightarrow H^0(\mathbf{Q}, M) \rightarrow H^0(\mathbf{Q}, M'') \rightarrow H_{f,S}^1(\mathbf{Q}, M') \rightarrow H_{f,S}^1(\mathbf{Q}, M) \rightarrow H_{f,S}^1(\mathbf{Q}, M'').$$

The following proposition summarizes the facts we will need about the Selmer groups we have defined:

Proposition 2.4. (1) *Suppose $W^{G_\Sigma} = 0$. Then $H_\Sigma^1(\mathbf{Q}, W_n) \cong H_\Sigma^1(\mathbf{Q}, W)[\varpi^n]$ and $H_{f,\min}^1(\mathbf{Q}, W_n) \cong H_{f,\min}^1(\mathbf{Q}, W)[\varpi^n]$.*
 (2) *If the module W^{I_ℓ} is divisible for every $\ell \in \Sigma \setminus \{p\}$, then $H_{f,\text{ur}}^1(\mathbf{Q}, W_n) = H_{f,\min}^1(\mathbf{Q}, W_n)$ and $H_{f,\text{ur}}^1(\mathbf{Q}, W) = H_{f,\min}^1(\mathbf{Q}, W)$.*
 (3) *If W is short crystalline at p then one can identify $H_f^1(\mathbf{Q}_p, W_n)$ as the image of $\text{Ext}_{\mathcal{MF}_\mathcal{O}}^1(1_{\text{FD}}, D)$, where D maps to W_n by the Fontaine-Laffaille functor.*

Proof. The first statement follows from Lemma 2.3 applied to the sequence $0 \rightarrow W[\varpi^n] \rightarrow W \xrightarrow{\times \varpi^n} W \rightarrow 0$ since the minimal and crystalline finite-singular structures on W_n are induced from the one on W by definition. (This statement is also stated as [Rub00], Lemma 1.5.4 (where this Selmer group is denoted $\mathcal{S}^\emptyset(\mathbf{Q}, W)$) upon noting that our assumption forces injectivity of the map $H^1(G_\Sigma, W_n) \rightarrow H^1(G_\Sigma, W)[\varpi^n]$ by [Rub00], Lemma 1.2.2(i)).

The second statement follows immediately from Remark 2.2, and the final one from Remark 2.1. \square

In the following we will write $H_f^1(\mathbf{Q}, M)$ for $H_{f,\text{ur}}^1(\mathbf{Q}, M)$ and $H_\Sigma^1(\mathbf{Q}, M)$ for $H_{\Sigma,\text{ur}}^1(\mathbf{Q}, M)$ for any p -adic G_Σ -module M .

2.2. More properties of Selmer groups. Assume that $k \geq 2$ is an even integer such that $p > 2k - 2$. Note that under this assumption the p -adic Galois representation ρ_f associated to an eigenform of weight $2k - 2$ of level not divisible by p is short crystalline. This assumption will be in force throughout the paper.

Proposition 2.5. *Let $\rho_f : G_\Sigma \rightarrow \text{GL}_2(E)$ be the Galois representation attached to a newform $f \in S_{2k-2}(N)$ for N a square-free integer with $p \nmid N$ and $\Sigma = \{\ell \mid N\} \cup \{p\}$. Suppose also that the residual representation $\bar{\rho}_f$ is absolutely irreducible and ramified at every prime $\ell \mid N$. Then for $i = 1, 2$*

- (i) $(\rho_f(i - k) \otimes E/\mathcal{O})^{I_\ell}$ is divisible for all $\ell \neq p$
- (ii) $(\rho_f(i - k) \otimes E/\mathcal{O})^{G_\Sigma} = 0$.

Proof. For the convenience of the reader we include the proof, however one can also consult the proof of Lemma 3.4 in [PW11].

(i) Set $\varrho := \rho_f(i - k)$ ($i = 1, 2$). We choose a basis $\{e_1, e_2\}$ for the space $V = V_\varrho$ so that ϱ with respect to that basis is valued in $\mathrm{GL}_2(\mathcal{O})$. We set $T = \mathcal{O}e_1 \oplus \mathcal{O}e_2$ (a G_Σ -stable \mathcal{O} -lattice in V) and set $W = V/T$.

One has by our assumption and by Theorem 3.26(3)(b) of [Hid00] that

$$(2.2) \quad \rho_f|_{G_{\mathbf{Q}_\ell}} \cong_E \begin{bmatrix} \psi\epsilon & * \\ & \psi \end{bmatrix}, \quad \rho_f|_{I_\ell} \neq 1,$$

where $\psi : G_{\mathbf{Q}_\ell} \rightarrow E^\times$ denotes the unramified character sending Frob_ℓ to $a_\ell(f)$. So, there exists $A \in \mathrm{GL}_2(E)$ such that

$$A\varrho|_{G_\ell}A^{-1} = \begin{bmatrix} \psi\epsilon^{i-k+1} & * \\ & \psi\epsilon^{i-k} \end{bmatrix}.$$

First let's note that ϱ is only tamely ramified at ℓ and so the image of I_ℓ is pro-cyclic. Suppose that $g \in I_\ell$ is such that $\varrho(g)$ topologically generates the image of I_ℓ . Conjugating $\varrho(g)$ by a matrix of the form $\mathrm{diag}(u\varpi^n, 1)$ for a suitable n and unit u we may assume that $A\varrho(g)A^{-1} = \begin{bmatrix} 1 & 1 \\ & 1 \end{bmatrix} \in \mathrm{GL}_2(\mathcal{O})$.

Write $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. By enlarging E (so that $\sqrt{\det A} \in E$) we may also assume that $\det A = 1$. So,

$$\varrho(g) = A^{-1} \begin{bmatrix} 1 & 1 \\ & 1 \end{bmatrix} A = \begin{bmatrix} 1 + cd & d^2 \\ -c^2 & 1 - cd \end{bmatrix}.$$

Since ϱ is valued in $\mathrm{GL}_2(\mathcal{O})$ we must have $c^2 \in \mathcal{O}$ and $d^2 \in \mathcal{O}$. Since \mathcal{O} is integrally closed in E this implies $c, d \in \mathcal{O}$.

Since $\bar{\rho}_f$ is ramified at ℓ so is $\bar{\varrho}$, i.e., we must have that either $c \in \mathcal{O}^\times$ or $d \in \mathcal{O}^\times$. Let $v \in V$. Then $v + T \in W^{I_\ell}$ if and only if $g \cdot v - v \in T$.

Writing v in the basis $\{e_1, e_2\}$ as $v = \begin{bmatrix} x \\ y \end{bmatrix}$ the above relation assumes the form: $\begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} \mathcal{O} \\ \mathcal{O} \end{bmatrix} \in W^{I_\ell}$ if and only if

$$\begin{bmatrix} cdx + d^2y \\ -c^2x - cdy \end{bmatrix} = \varrho(g) \begin{bmatrix} x \\ y \end{bmatrix} - \begin{bmatrix} x \\ y \end{bmatrix} \in \begin{bmatrix} \mathcal{O} \\ \mathcal{O} \end{bmatrix}$$

which happens if and only if

$$d(cx + dy) \in \mathcal{O} \quad \text{and} \quad -c(cx + dy) \in \mathcal{O}.$$

Now suppose $c \in \mathcal{O}^\times$. Then from the second relation we get $cx + dy \in \mathcal{O}$. Since this condition also implies that the first expression is in \mathcal{O} , we get that in this case $\begin{bmatrix} x \\ y \end{bmatrix} + T \in W^{I_\ell}$ if and only if $cx + dy \in \mathcal{O}$. Similarly if $d \in \mathcal{O}^\times$, then we again get the exact same condition. Hence we conclude that

$$W^{I_\ell} = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} \mathcal{O} \\ \mathcal{O} \end{bmatrix} \mid x, y \in E, cx + dy \in \mathcal{O} \right\}.$$

Suppose $c \in \mathcal{O}^\times$. Then

$$W^{I_\ell} = \left\{ \begin{bmatrix} -c^{-1}dy \\ y \end{bmatrix} + \begin{bmatrix} \mathcal{O} \\ \mathcal{O} \end{bmatrix} \mid y \in E \right\}.$$

Consider the map from the module on the right to E/\mathcal{O} given by

$$\begin{bmatrix} -c^{-1}dy \\ y \end{bmatrix} + \begin{bmatrix} \mathcal{O} \\ \mathcal{O} \end{bmatrix} \mapsto y + \mathcal{O}.$$

It is clearly a map of \mathcal{O} -modules and clearly surjective. Suppose that $\begin{bmatrix} -c^{-1}dy \\ y \end{bmatrix} + \begin{bmatrix} \mathcal{O} \\ \mathcal{O} \end{bmatrix} \in \ker$. Then $y \in \mathcal{O}$, which implies that $c^{-1}dy \in \mathcal{O}$, so the map is also injective, hence an isomorphism. Analogous map works in the case when $d \in \mathcal{O}^\times$. Since E/\mathcal{O} is divisible this proves that W^{I_ℓ} is divisible.

(ii) Let T, W be as in the proof of (i). Clearly if $v \in W^{G_\Sigma}$, then some \mathcal{O} -multiple of v is a non-zero element in $W[\varpi]$. Hence it is enough to show that $W[\varpi]^{G_\Sigma} = 0$. However, we have (see e.g. [Rub00], Lemma 1.2.2(iii)) $W[\varpi] \cong T/\varpi T \cong \bar{\rho}$, so the claim follows from the irreducibility of $\bar{\rho}_f$. \square

Corollary 2.6. *Let ρ_f be as in Proposition 2.5. For $i = 1, 2$ one has*

$$H_{f,\min}^1(\mathbf{Q}, \bar{\rho}_f(i-k)) = H_{f,\text{ur}}^1(\mathbf{Q}, \bar{\rho}_f(i-k)).$$

Proof. This follows from Propositions 2.5 and 2.4(1)+(2). \square

Proposition 2.7. *Let $\rho_f : G_\Sigma \rightarrow \text{GL}_2(E)$ be the Galois representation attached to a newform $f \in S_{2k-2}(N)$ with $p \nmid N$. Suppose also that the residual representation $\bar{\rho}_f$ is absolutely irreducible and ramified at every prime $\ell \mid N$. One has $H_\Sigma^1(\mathbf{Q}, \rho_f(2-k) \otimes E/\mathcal{O}) = H_f^1(\mathbf{Q}, \rho_f(2-k) \otimes E/\mathcal{O})$ if N is square-free, $\Sigma = \{\ell \mid N\} \cup \{p\}$, and $p \nmid 1 + w_{f,\ell}$ for all primes $\ell \mid N$. Here $w_{f,\ell} \in \{1, -1\}$ is the eigenvalue of the Atkin-Lehner involution at ℓ .*

Remark 2.8. Note that when N is prime $w_{f,N} = (-1)^{k-1}W_f$, where $W_f \in \{1, -1\}$ is the sign of the functional equation of f (see e.g. [Shi71] Theorem 3.66).

Proof. We use Lemma 5.6 in [BK13] with $V = \rho_f(2-k)$. Then $V^* = V^\vee(1) = V$.

By (2.2) we have that $(V^*)^{I_\ell} = \psi(3-k)$ and hence we get (using the notation of [BK13])

$$P_\ell(V^*, 1) = \det(1 - \text{Frob}_\ell |_{(V^*)^{I_\ell}}) = 1 - a_\ell(f)\ell^{3-k}.$$

By Theorem 3 in [AL70] (but with Hecke action normalized as in [Miy89], see [Miy89] Theorem 4.6.17(2) for the correct power of ℓ in our normalization) we know that $a_\ell(f) = -\ell^{k-2}w_{f,\ell}$. This shows that under the conditions of the Proposition we have $\text{val}_p(P_\ell(V^*, 1)) = 0$ for all $\ell \nmid N$. To be able to use Lemma 5.6 in [BK13], we also need to know that W^{I_ℓ} is divisible, where $W = V/T$ with V as above (i.e, the two dimensional vector space over E on which G_Σ acts via $\rho_f(2-k)$) and $T \subset V$ a G_Σ -stable \mathcal{O} -lattice. However, this follows from Proposition 2.5(i). We note here that the order of the Selmer group does not depend on the choice of T since ρ is absolutely irreducible (cf. e.g., Lemma 9.7 in [Klo09]). \square

Proposition 2.9. *One has*

$$H_f^1(\mathbf{Q}, \mathbf{F}(1)) = H_f^1(\mathbf{Q}, \mathbf{F}(-1)) = 0.$$

Proof. For the latter Selmer group this is essentially Proposition 6.16 along with Theorem 6.17 in [Was97] which tell us that the p -part of the class group of the splitting field of χ^{-1} is trivial (as the numerator of B_2 (the second Bernoulli number) is 1). This implies that $H_{f,\min}^1(\mathbf{Q}, W_1) = 0$ for $W = \mathbf{Q}_p/\mathbf{Z}_p(-1)$. By Proposition 2.4 we also have that $H_{f,\min}^1(\mathbf{Q}, W_1) = H_{f,\text{ur}}^1(\mathbf{Q}, W_1) = H_f^1(\mathbf{Q}, W_1)$ since $W^{\ell} = 0$ for all $\ell \neq p$.

On the other hand [Rub00] Proposition 1.6.4(ii) tells us that $H_f^1(\mathbf{Q}, \mathbf{Q}_p/\mathbf{Z}_p(1))$ fits into a short exact sequence

$$0 \rightarrow \mathbf{Z}^\times \otimes \mathbf{Q}_p/\mathbf{Z}_p \rightarrow H_f^1(\mathbf{Q}, \mathbf{Q}_p/\mathbf{Z}_p(1)) \rightarrow \text{Cl}(\mathbf{Q}) \rightarrow 0.$$

This implies that $H_f^1(\mathbf{Q}, \mathbf{Q}_p/\mathbf{Z}_p(1)) = 0$, and therefore $H_f^1(\mathbf{Q}, \mathbf{F}(1)) = H_f^1(\mathbf{Q}, \mathbf{Q}_p/\mathbf{Z}_p(1))[p] = 0$. \square

2.3. Computing the non-critical Selmer group. We will be interested in bounding from above the order of $H_f^1(\mathbf{Q}, \rho_f(1-k) \otimes E/\mathcal{O})$. In the case that $k = 2$ the motif $\rho_f(1-k)$ is not critical in the sense of Deligne. However, we will show how it can still be bounded by a certain p -adic L -value using Iwasawa theory for $p \neq 3$.

Let ω be the Teichmüller lift of χ . Suppose $f \in S_2(\Gamma_0(N))$ (with $p \nmid N$) is a newform ordinary at p and that $\bar{\rho}_f$ is absolutely irreducible and ramified at all primes $\ell \mid N$. We will denote by $L_p^{\text{an}}(f, \omega^{-1}) \in \mathcal{O}[[T]]$ the Manin-Vishik p -adic L -function for $f \otimes \omega^{-1}$ (see e.g. [Kat04] Theorem 16.2 where the p -adic L -function is defined for the p -stabilisation of f as an element of $\tilde{\Lambda} = \mathcal{O}[\Delta][[T]]$; we take its projection onto the component of $\tilde{\Lambda}$ where Δ acts by ω^{-1}).

Proposition 2.10. *For $p \neq 3$ we have*

$$\text{val}_p(\#H_f^1(\mathbf{Q}, \rho_f(-1) \otimes E/\mathcal{O})) \leq \text{val}_p(\#\mathcal{O}/L_p^{\text{an}}(f, \omega^{-1}, T = p)).$$

Proof. Write T_f for a choice of a G_Σ -invariant lattice inside the Galois representation $V_f = \rho_f : G_\Sigma \rightarrow \text{GL}_2(E)$. Since f is ordinary there is a unique D_p -stable line V_f^+ of V_f on which I_p acts via ϵ . Set $T_f^+ := T_f \cap V_f^+$. Following [SU14] (p. 34) for any continuous character $\xi : G_{F,\Sigma} \rightarrow \mathcal{O}^\times$ we define the Greenberg Selmer group

$$\text{Sel}_{F,E}^\Sigma(f, \xi) := \ker \left\{ H^1(G_{F,\Sigma}, T_f \otimes \xi \otimes E/\mathcal{O}) \rightarrow H^1(I_p, (T_f/T_f^+) \otimes \xi \otimes E/\mathcal{O}) \right\},$$

where $F = \mathbf{Q}$ or $F = \mathbf{Q}_\infty$, the cyclotomic \mathbf{Z}_p -extension of \mathbf{Q} . We are interested in the case where $\xi = \omega^{m-1}\epsilon^{-m}$ for $m = 1$, while Skinner and Urban are mainly interested in the case where $\xi = \omega^m\epsilon^{-m}$. We set $X_{F,E}^\Sigma(f, \xi) := \text{Hom}_{\mathcal{O}}(\text{Sel}_{F,E}^\Sigma(f, \xi), E/\mathcal{O})$ to be the Pontryagin dual. By a theorem due to Kato ([SU14], Theorem 3.15) we know that $X_{\mathbf{Q}_\infty,E}^\Sigma(f, \xi)$ is a torsion Λ -module, where $\Lambda = \mathcal{O}[[\text{Gal}(\mathbf{Q}_\infty/\mathbf{Q})]] = \mathcal{O}[[T]]$. Set $Ch_{\mathbf{Q}_\infty,E}^\Sigma(f, \omega^{-1})$ to be the characteristic ideal of $X_{\mathbf{Q}_\infty,E}^\Sigma(f, \omega^{-1})$. The Main Conjecture of Iwasawa Theory states that $Ch_{\mathbf{Q}_\infty,E}^\Sigma(f, \omega^{-1})$ is a principal ideal generated by $L_p^{\text{an}}(f, \omega^{-1}, T)$ (cf. Conjecture 3.24 in [SU14]). By Theorem 17.14 in [Kat04] (but see also Theorem 3.25 in [SU14], whose notation is consistent

with ours) the ideal $Ch_{\mathbf{Q}_{\infty}, E}^{\Sigma}(f, \omega^{-1})$ divides $L_p^{\text{an}}(f, \omega^{-1}, T)$ in Λ . Note that the assumption in Theorem 17.14 in [Kat04] that the image of ρ_f contains $\text{SL}_2(\mathbf{Z}_p)$ can be replaced by $\bar{\rho}_f$ being ramified at some $\ell \parallel N$, as explained in [Ski16] pages 187/8.

To relate this to our $H_f^1(\mathbf{Q}, \rho_f(-1) \otimes E/\mathcal{O})$ we need a control theorem which was proved by Skinner and Urban in the case $\xi = \omega^m \epsilon^{-m}$ (Corollary 3.21(i) in [SU14] which should say $m \neq 0$ instead of $m \neq 1$). Let us only indicate here what changes are needed to adapt it to the case $\xi = \epsilon^{-1} = \omega^{-1} \cdot (\omega/\epsilon)^m$ with $m = 1$.

Theorem 2.11 (Control Theorem). *One has*

$$\#X_{\mathbf{Q}, E}^{\Sigma}(f, \epsilon^{-1}) \mid \#\Lambda/(T - p, Ch_{\mathbf{Q}_{\infty}, E}^{\Sigma}(f, \omega^{-1})).$$

Proof. Skinner and Urban prove the corresponding statement (for $\xi = \omega^m \epsilon^{-m}$) by a series of results culminating in Proposition 3.20 which is then used to prove Corollary 3.21 (here we only care about one divisibility as stated above instead of equality). The assumption that f be p -stabilized in [SU06] Section 3.3.13 is not necessary. The first result is Proposition 3.10. The proof of the only part relevant for us, the injectivity of the map $\text{Sel}_{\mathbf{Q}, E}^{\Sigma}(f, \epsilon^{-1}) \rightarrow \text{Sel}_{\mathbf{Q}_{\infty}, E}^{\Sigma}(f, \epsilon^{-1})^{G_{\Sigma}}$, carries over unchanged to our situation. As the next step we need an analogue of Proposition 3.20 (stating that $X_{\mathbf{Q}_{\infty}, E}^{\Sigma}(f, \omega^{-1})$ has no non-zero pseudo-null Λ -submodules) for which it is sufficient to prove corresponding analogues of Lemmas 3.18 and 3.19. Let us explain what changes are needed to the proof of Lemma 3.18 which for us only holds for $p \neq 3$. While in [SU14] the module denoted there by $M^{-}[x]|_{I_p}$ is isomorphic to $E/\mathcal{O}(\omega^m \epsilon^{-m})$ we have $M^{-}[x]|_{I_p} \cong E/\mathcal{O}(\omega^{m-1} \epsilon^{-m})$ (and $m = 1$). The assumption that $p \neq 3$ is needed in our case to ensure that $M^{-}[\mathfrak{m}]^*(1) = \mathbf{F}(\omega^2)$ is ramified at p (note $\epsilon = \omega \bmod \varpi$ and $\mathfrak{m} = (x, \varpi)$), where $*$ denotes the Pontryagin dual. Also in our case $M^{-}[x]^{I_p} = 0$, hence in particular finite. Lemma 3.19 follows the same way as in [SU14]. Having all this, Corollary 3.21 is also proved the same way as in [SU14]. \square

We continue with the proof of Proposition 2.10. As a corollary of the control theorem we obtain the following inequality:

$$\#X_{\mathbf{Q}, E}^{\Sigma}(f, \epsilon^{-1}) \leq \#\mathcal{O}/L_p^{\text{an}}(f, \omega^{-1}, T = p)$$

(cf. Theorem 3.36 in [SU14]). Note that Skinner and Urban work within the critical range of the relevant L -function, while we are outside of this range, so in particular we do not relate the p -adic L -value to a classical one.

Let us now show that $\#X_{\mathbf{Q}, E}^{\Sigma}(f, \epsilon^{-1}) \geq \#H_f^1(\mathbf{Q}, \rho_f(-1) \otimes E/\mathcal{O})$. First note that $\#X_{\mathbf{Q}, E}^{\Sigma}(f, \epsilon^{-1}) = \#\text{Sel}_{\mathbf{Q}, E}^{\Sigma}(f, \epsilon^{-1})$ since the group on the left is finite. We will show that $\text{Sel}_{\mathbf{Q}, E}^{\Sigma}(f, \epsilon^{-1}) \supset H_f^1(\mathbf{Q}, \rho_f(-1) \otimes E/\mathcal{O})$. The conditions at $\ell \neq p$ defining both groups coincide by combining Propositions 2.5(i) and 2.4(2), so it is enough to compare conditions at p . There we have $H_f^1(\mathbf{Q}_p, \rho_f(-1) \otimes E/\mathcal{O}) \subset H_g^1(\mathbf{Q}_p, \rho_f(-1) \otimes E/\mathcal{O})$ which follows from (3.7) in [BK90] by taking images in the cohomology with divisible coefficients. On the other hand the diagram at the bottom of p. 87 in [Och00] tells us

that $H_g^1(\mathbf{Q}_p, \rho_f(-1) \otimes E/\mathcal{O})$ injects into

$$\begin{aligned} H_{\text{Gr}}^1(\mathbf{Q}_p, \rho_f(-1) \otimes E/\mathcal{O}) &:= \ker(H^1(\mathbf{Q}_p, T_f(-1) \otimes E/\mathcal{O}) \rightarrow H^1(I_p, T_f(-1) \otimes E/\mathcal{O})) \\ &\hookrightarrow \ker(H^1(\mathbf{Q}_p, T_f(-1) \otimes E/\mathcal{O}) \rightarrow H^1(I_p, (T_f/T_f^+)(-1) \otimes E/\mathcal{O})). \end{aligned}$$

Indeed, $H_g^1(\mathbf{Q}_p, \rho_f(-1) \otimes E/\mathcal{O})$ is defined as the image of $H_g^1(\mathbf{Q}_p, V_f(-1))$ in $H^1(\mathbf{Q}_p, T_f(-1) \otimes E/\mathcal{O})$ and according to the diagram the defining map factors through $H_{\text{Gr}}^1(\mathbf{Q}_p, \rho_f(-1) \otimes E/\mathcal{O})$. The corresponding inclusions of global Selmer groups follow. \square

3. DEFORMATIONS

Let $\rho : G_\Sigma \rightarrow \text{GL}_2(\mathbf{F})$ be a continuous absolutely irreducible odd Galois representation of determinant χ^{2k-3} , short crystalline at p and such that for every prime $\ell \in \Sigma \setminus \{p\}$ one has $\rho|_{I_\ell} \cong \begin{bmatrix} 1 & * \\ 0 & 1 \end{bmatrix} \neq I_2$. Consider a residual (short crystalline) representation

$$\bar{\sigma} := \begin{bmatrix} \chi^{k-2} & a & b \\ & \rho & c \\ & & \chi^{k-1} \end{bmatrix} : G_\Sigma \rightarrow \text{GL}_4(\mathbf{F}),$$

where a, c are assumed to be non-trivial classes in $H^1(\mathbf{Q}, \rho(1-k))$ (no assumption on b).

Remark 3.1. We have

$$\text{Hom}(\rho, \chi^{k-2}) = \rho^\vee \otimes \chi^{k-2} = \rho \otimes \chi^{3-2k} \otimes \chi^{k-2} = \rho(1-k) = \text{Hom}(\chi^{k-1}, \rho),$$

so a and c are indeed both classes in $H^1(\mathbf{Q}, \rho(1-k))$.

$$\text{Set } N_1 = \begin{bmatrix} 0 & & & \\ & 0 & 1 & \\ & & 0 & \\ & & & 0 \end{bmatrix} \text{ and for any prime } \ell \neq p \text{ let } t_p : I_\ell \rightarrow \mathbf{Z}_p(1) \text{ be}$$

the tame character. We will always assume that $\bar{\sigma}$ as above is *semi-abelian* by which we mean that for each $\ell \in \Sigma \setminus \{p\}$ one has

$$(3.1) \quad \bar{\sigma}|_{I_\ell} \cong \exp(t_p N_1).$$

Lemma 3.2. *Let $x \in G_\Sigma$ be such that $\bar{\sigma}(x)$ generates the image of I_ℓ . Then $\bar{\sigma}$ is isomorphic to a representation of the form*

$$\bar{\sigma}' = \begin{bmatrix} \chi^{k-2} & *_1 & *_2 \\ & \rho & *_3 \\ & & \chi^{k-1} \end{bmatrix}$$

and such that

$$\bar{\sigma}'(x) - I_4 = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & * & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \neq 0,$$

i.e., the matrix conjugating $\bar{\sigma}|_{I_\ell}$ to $\exp(t_\ell N_1)$ does not change the matrix form of $\bar{\sigma}$.

Proof. Without loss of generality assume that $\rho(x) = \begin{bmatrix} 1 & \alpha \\ 0 & 1 \end{bmatrix}$ with $\alpha \neq 0$.

Write $\alpha_{i,j}$ for the (i,j) -entry of $\bar{\sigma}$. Then

$$\bar{\sigma}(x) - I_4 = \begin{bmatrix} 0 & \alpha_{1,2}(x) & \alpha_{1,3}(x) & \alpha_{1,4}(x) \\ 0 & 0 & \alpha & \alpha_{2,4}(x) \\ 0 & 0 & 0 & \alpha_{3,4}(x) \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Since $\text{rank}(\bar{\sigma}(x) - I_4) = \text{rank}(N_1) = 1$, we must also have that $M := \begin{bmatrix} \alpha_{1,3}(x) & \alpha_{1,4}(x) \\ \alpha & \alpha_{2,4}(x) \end{bmatrix}$ has rank 1. Thus there exists $\beta \in \mathbf{F}$ such that $\alpha_{1,3}(x) = \beta\alpha$ and $\alpha_{1,4}(x) = \beta\alpha_{2,4}(x)$. Hence we have

$$AMD^{-1} = \begin{bmatrix} 0 & 0 \\ \alpha & 0 \end{bmatrix}, \quad \text{where } A = \begin{bmatrix} 1 & -\beta \\ 0 & 1 \end{bmatrix}, D^{-1} = \begin{bmatrix} 1 & -\alpha_{2,4}(x)/\alpha \\ 0 & 1 \end{bmatrix} \in \text{GL}_2(\mathbf{F}).$$

Thus

$$\begin{bmatrix} A & \\ & D \end{bmatrix} (\bar{\sigma}(x) - I_4) \begin{bmatrix} A & \\ & D \end{bmatrix}^{-1} = \begin{bmatrix} 0 & \alpha'_{1,2}(x) & 0 & 0 \\ 0 & 0 & \alpha & 0 \\ 0 & 0 & 0 & \alpha'_{3,4}(x) \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

However, since the last matrix has rank one and $\alpha \neq 0$, we must have $\alpha'_{1,2}(x) = \alpha'_{3,4}(x) = 0$. Furthermore note that

$$(3.2) \quad \bar{\sigma}' := \begin{bmatrix} A & \\ & D \end{bmatrix} \bar{\sigma} \begin{bmatrix} A & \\ & D \end{bmatrix}^{-1} \\ = \begin{bmatrix} A & \\ & D \end{bmatrix} \begin{bmatrix} \chi^{k-2} & * & * \\ & \rho & * \\ & & \chi^{k-1} \end{bmatrix} \begin{bmatrix} A & \\ & D \end{bmatrix}^{-1} = \begin{bmatrix} A & \\ & D \end{bmatrix} \begin{bmatrix} \chi^{k-2} & *' & *' \\ & \rho & *' \\ & & \chi^{k-1} \end{bmatrix} \begin{bmatrix} A & \\ & D \end{bmatrix}^{-1}.$$

□

Corollary 3.3. *For $\ell \in \Sigma \setminus \{p\}$ we have $\bar{\sigma}|_{I_\ell} \cong 1 \oplus \rho|_{I_\ell} \oplus 1$. In particular, the extensions given by the entries a and c of $\bar{\sigma}$ are both split when restricted to I_ℓ .*

Let $\text{LCN}(E)$ be the category of local complete Noetherian \mathcal{O} -algebras with residue field \mathbf{F} .

Definition 3.4. For an object A of $\text{LCN}(E)$ we will say that $\sigma : G_\Sigma \rightarrow \text{GL}_4(A)$ is *minimal* if $\sigma|_{I_\ell} \cong_{\text{Tot}(A)} \exp(t_p N_1)$ for every prime $\ell \in \Sigma \setminus \{p\}$. Here $\text{Tot}(A)$ denotes the total ring of fractions of A .

Note that any minimal $\sigma : G_\Sigma \rightarrow \text{GL}_4(A)$ is tamely ramified at all $\ell \in \Sigma \setminus \{p\}$, so $\sigma(I_\ell)$ is a pro-cyclic group for all such ℓ .

We consider the deformation problem for $\bar{\sigma}$ where to every object A of $\text{LCN}(E)$ we assign the set of strict equivalence classes of representations $\sigma : G_\Sigma \rightarrow \text{GL}_4(A)$ which are short crystalline, have determinant ϵ^{4k-6} , are minimal and reduce to $\bar{\sigma}$ modulo the maximal ideal \mathfrak{m}_A of A .

Theorem 3.5. *The above deformation problem is representable.*

Proof. It is easy to check that $\bar{\sigma}$ has scalar centralizer, so by e.g. [CHT08] Proposition 2.2.9 it is enough to verify that short crystallinity and minimality are deformation conditions. The former is due to Ramakrishna (cf. Theorem 1.1 of [Ram93]). The latter can be proven similar to [Maz97] Section 29. \square

We will write R' for the universal deformation ring and $(\sigma')^{\text{univ}} : G_\Sigma \rightarrow \text{GL}_4(R')$ for the universal deformation.

Consider the following antiinvolution $\tau : R'[G_\Sigma] \rightarrow R'[G_\Sigma]$ given by $\tau(g) = \epsilon^{2k-3}(g)g^{-1}$. Clearly τ induces the following permutation on the set \mathcal{I} of Jordan-Hölder factors of $\bar{\sigma}$:

$$\chi^{k-1} \mapsto \chi^{k-2}, \quad \rho \mapsto \rho, \quad \chi^{k-2} \mapsto \chi^{k-1}.$$

For $A \in \text{LCN}(E)$ we will say that a short crystalline deformation $\sigma : G_\Sigma \rightarrow \text{GL}_4(A)$ is τ -self-dual if $\text{tr } \sigma = \text{tr } \sigma \circ \tau$. Note that for semisimple σ and fields A this is equivalent to $\sigma^\vee \cong \sigma(3-2k)$.

Proposition 3.6. *The functor assigning to an object $A \in \text{LCN}(E)$ the set of strict equivalence classes of τ -self-dual short crystalline (at p), minimal deformations to $\text{GL}_4(A)$ is representable by the quotient of R' by the ideal generated by $\{\text{tr } (\sigma')^{\text{univ}}(g) - \text{tr } (\sigma')^{\text{univ}}(\tau(g)) \mid g \in G_\Sigma\}$. We will denote this quotient by R and will write σ^{univ} for the corresponding universal deformation.*

We write R^{red} for the quotient of R by its nilradical and σ^{red} for the corresponding (universal) deformation, i.e., the composite of σ^{univ} with $R \twoheadrightarrow R^{\text{red}}$.

4. LATTICE

Theorem 4.1. *Let E be a finite extension of \mathbf{Q}_p with valuation ring \mathcal{O} , uniformizer ϖ and $\mathbf{F} = \mathcal{O}/\varpi\mathcal{O}$. Let $\sigma : G_\Sigma \rightarrow \text{GL}_n(E)$ be an absolutely irreducible Galois representation. Suppose there exists a G_Σ -stable \mathcal{O} -lattice Λ in the space of σ such that*

$$\bar{\sigma}_\Lambda \cong \begin{bmatrix} \rho_1 & * \\ 0 & \rho_2 \end{bmatrix}$$

with $\rho_i : G_\Sigma \rightarrow \text{GL}_{n_i}(\mathbf{F})$ (for $i = 1, 2$) having scalar centralizer, ρ_1 semisimple and such that none of the irreducible constituents of ρ_1^{ss} is isomorphic to any of the irreducible constituents of ρ_2^{ss} . Then there exists a G_Σ -stable lattice Λ' in the space of σ such that

$$\bar{\sigma}_{\Lambda'} \cong \begin{bmatrix} \rho_1 & * \\ 0 & \rho_2 \end{bmatrix} \not\cong \rho_1 \oplus \rho_2.$$

Let us note that if ρ_2 is semisimple, then Theorem 4.1 follows from a generalization of Theorem in [Urb99] - cf. Remark (a) in [Urb99].

We begin the proof of Theorem 4.1 with the following lemma.

Lemma 4.2. *Let ρ_1, ρ_2 be as in Theorem 4.1. Assume that $\bar{\sigma} : G_\Sigma \rightarrow \mathrm{GL}_n(\mathbf{F})$ given by*

$$\bar{\sigma} = \begin{bmatrix} \rho_1 & f \\ & \rho_2 \end{bmatrix}$$

is isomorphic to $\rho_1 \oplus \rho_2$. Then there exists a matrix

$$M = \begin{bmatrix} I_{n_1} & B \\ & I_{n_2} \end{bmatrix} \in \mathrm{GL}_n(\mathbf{F})$$

such that $M\bar{\sigma}M^{-1} = \begin{bmatrix} \rho_1 & \\ & \rho_2 \end{bmatrix}$.

Proof. By assumption there exists $h = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathrm{GL}_n(\mathbf{F})$ (with correct dimensions) such that

$$(4.1) \quad h \begin{bmatrix} \rho_1 & f \\ & \rho_2 \end{bmatrix} h^{-1} = \begin{bmatrix} \rho_1 & 0 \\ & \rho_2 \end{bmatrix}$$

which implies that $\rho_2 C = C \rho_1$, i.e., that $C \in \mathrm{Hom}_{G_\Sigma}(\rho_1, \rho_2)$. The image of C is a quotient of ρ_1 which is a submodule of ρ_2 . However, by our assumption on the irreducible constituents of ρ_1^{ss} and ρ_2^{ss} , this image must be zero, so $C = 0$. Then (4.1) implies that A centralizes ρ_1 and D centralizes ρ_2 , hence $A = \alpha I_{n_1}$ and $D = \delta I_{n_2}$ with $\alpha, \delta \in \mathbf{F}^\times$. By simply scaling h by α^{-1} we may assume without loss of generality that $\alpha = 1$. Finally (with $C = 0$), the B -entry of $h \begin{bmatrix} \rho_1 & f \\ & \rho_2 \end{bmatrix} h^{-1}$ equals

$$(4.2) \quad -\alpha \rho_1 \alpha^{-1} B \delta^{-1} + \alpha f \delta^{-1} + B \rho_2 \delta^{-1} = \rho_1 B \delta^{-1} + f \delta^{-1} + B \rho_2 \delta^{-1} = 0.$$

From this we see that the expression in (4.2) equals zero regardless of the value of δ , so we may assume that $\delta = 1$. \square

Proof of Theorem 4.1. In this we follow Ribet [Rib76], the proof of Proposition 2.1. Let the notation be as in the statement of the Theorem. Suppose no Λ' exists. Then by Lemma 4.2 there exists $h = \begin{bmatrix} I_{n_1} & B \\ & I_{n_2} \end{bmatrix}$ (where B here is any lift to $M_{n_1 \times n_2}(\mathcal{O})$ of B in lemma) such that for all $g \in G_\Sigma$ one has

$$h \sigma_\Lambda(g) h^{-1} = \begin{bmatrix} \alpha & \varpi \beta \\ \varpi \gamma & \delta \end{bmatrix} \quad \text{with } \alpha \equiv \rho_1(g) \pmod{\varpi}, \delta \equiv \rho_2(g) \pmod{\varpi}$$

and $\alpha, \beta, \gamma, \delta$ with entries in \mathcal{O} . Then

$$\tau_1(g) := \begin{bmatrix} I_{n_1} & \\ & \varpi I_{n_2} \end{bmatrix} h \sigma_\Lambda(g) h^{-1} \begin{bmatrix} I_{n_1} & \\ & \varpi I_{n_2} \end{bmatrix}^{-1} = \begin{bmatrix} \alpha & \beta \\ \varpi^2 \gamma & \delta \end{bmatrix}.$$

The reduction of τ_1 still satisfies the assumptions of Lemma 4.2, hence by that lemma there exists $h_1 = \begin{bmatrix} I_{n_1} & B_1 \\ & I_{n_2} \end{bmatrix} \in \mathrm{GL}_n(\mathcal{O})$ such that $h_1 \tau_1 h_1^{-1} = \begin{bmatrix} \alpha_1 & \varpi \beta' \\ \varpi^2 \gamma' & \delta' \end{bmatrix}$, where $\alpha', \beta', \gamma' = \gamma, \delta$ have entries in \mathcal{O} . We continue this way as Ribet does to conclude in the end that σ itself is reducible, which leads to a contradiction. This finishes the proof of Theorem 4.1. \square

Corollary 4.3. *Let $\sigma : G_\Sigma \rightarrow \mathrm{GL}_4(E)$ be an absolutely irreducible short crystalline Galois representation. Suppose that $\bar{\sigma}^{\mathrm{ss}} = \chi^{k-2} \oplus \rho \oplus \chi^{k-1}$ with ρ absolutely irreducible and that $\sigma|_{I_\ell} \cong_E \exp(t_p N_1)$ for all $\ell \in \Sigma \setminus \{p\}$. Then there exists a G_Σ -stable lattice Λ in the space of σ such that*

$$\bar{\sigma}_\Lambda = \begin{bmatrix} \chi^{k-2} & *_1 & *_2 \\ & \rho & *_3 \\ & & \chi^{k-1} \end{bmatrix}$$

*is semi-abelian with $*_3$ a non-split extension of χ^{k-1} by ρ and $*_1$ a non-trivial extension of ρ by χ^{k-2} .*

Proof. Applying Theorem 6.1 in [Bro11] with $\mathcal{R} = \mathcal{O}$ and $I = \varpi\mathcal{O}$ (note that $\mathcal{T} \cong \mathcal{O}$ because any lattice in E is isomorphic to \mathcal{O}) we get that there exists a lattice Λ' in the space of σ such that

$$\bar{\sigma}_{\Lambda'} = \begin{bmatrix} \chi^{k-2} & 0 & *_2 \\ & \rho & *_3 \\ & & \chi^{k-1} \end{bmatrix} \not\cong \chi^{k-2} \oplus \rho \oplus \chi^{k-1}.$$

We now claim that $\bar{\sigma}_{\Lambda'}$ cannot be equivalent to a representation of the same shape where $*_3 = 0$. Indeed, suppose it were, then $\bar{\sigma}_{\Lambda'} \cong \rho \oplus \begin{bmatrix} \chi^{k-2} & *_2 \\ & \chi^{k-1} \end{bmatrix}$.

Let $\ell \in \Sigma \setminus \{p\}$. If $x \in G_\Sigma$ is such that $\sigma(x)$ generates the image of I_ℓ , then the rank of $\sigma(x) - I_4$ must be one. Since $\sigma_{\Lambda'} \cong_E \sigma$, we also must have $\mathrm{rank}(\sigma_{\Lambda'}(x) - I_4) = 1$, and so also $\mathrm{rank}(\bar{\sigma}_{\Lambda'}(x) - I_4) = 1$. Let us prove this last implication. Since $\rho|_{I_\ell} \neq 1$, we must have $\mathrm{rank}(\bar{\sigma}_{\Lambda'}(x) - I_4) \geq 1$. So, we need to prove a rank one matrix cannot reduce mod ϖ to a matrix of a higher rank. Let A be a rank one matrix with entries in \mathcal{O} . Then every row is a scalar multiple of the first non-zero row. These scalars are of the form $u\varpi^n$, where $n \in \mathbf{Z}$. Pick a row for which n is minimal and by making this row first (permutation matrix has entries in \mathcal{O}), we may now assume that all $n \geq 0$, i.e, that all the scalars are in \mathcal{O} . Thus every row of the reduction of A mod ϖ is a scalar multiple of the first row. This establishes that $\mathrm{rank}(\bar{\sigma}_{\Lambda'}(x) - I_4) \leq 1$.

However, if $\bar{\sigma}_{\Lambda'} \cong \begin{bmatrix} \rho & & \\ & \chi^{k-2} & *_2 \\ & & \chi^{k-1} \end{bmatrix}$, then this rank condition forces $*_2$ to

be unramified at ℓ . Thus in this case, $\sigma_{\Lambda'}$ has a direct summand isomorphic to $\begin{bmatrix} \chi^{k-2} & * \\ & \chi^{k-1} \end{bmatrix}$ with $*$ unramified away from p . This direct summand is short crystalline at p since $\sigma_{\Lambda'}$ is, and so gives rise to an element in $H_f^1(\mathbf{Q}, \mathbf{F}(-1))$. Since this group is trivial (by Proposition 2.9), this implies that $\bar{\sigma}_{\Lambda'}$ is semisimple, contradicting our assumption.

So we must have that $*_3$ gives a non-trivial extension of χ^{k-1} by ρ . Now, apply Theorem 4.1 with $\rho_1 = \chi^{k-2}$ and $\rho_2 = \begin{bmatrix} \rho & *_3 \\ & \chi^{k-1} \end{bmatrix}$ (note that ρ_2 has

scalar centralizer). This gives us

$$(4.3) \quad \bar{\sigma}_\Lambda = \begin{bmatrix} \chi^{k-2} & *1 & *2 \\ & \rho & *3 \\ & & \chi^{k-1} \end{bmatrix} \not\cong \chi^{k-2} \oplus \begin{bmatrix} \rho & *3 \\ & \chi^{k-1} \end{bmatrix}$$

with $*3$ a non-split extension of χ^{k-1} by ρ . It remains to show that $*1$ gives rise to a non-trivial extension of ρ by χ^{k-2} . Suppose that one has $\begin{bmatrix} \chi^{k-2} & *1 \\ & \rho \end{bmatrix} \cong \chi^{k-2} \oplus \rho$. Then $*1$ corresponds to a coboundary in $H^1(\mathbf{Q}, \text{Hom}(\rho, \chi^{k-2}))$.

More precisely, there exists a matrix $f \in M_{n_1 \times n_2}(\mathbf{F})$ (i.e., a map in $\text{Hom}_{\mathbf{F}}(\rho, \chi^{k-2})$) such that

$$\begin{bmatrix} \chi^{k-2} & *1 \\ & \rho \end{bmatrix} \cong \begin{bmatrix} \chi^{k-2} & \chi^{k-2}f - f\rho \\ 0 & \rho \end{bmatrix}.$$

However, note that

$$\begin{bmatrix} I_{n_1} & f & \\ & I_{n_2} & \\ & & I_{n_3} \end{bmatrix} \begin{bmatrix} \chi^{k-2} & a & b \\ & \rho & c \\ & & \chi^{k-1} \end{bmatrix} \begin{bmatrix} I_{n_1} & f & \\ & I_{n_2} & \\ & & I_{n_3} \end{bmatrix}^{-1} = \begin{bmatrix} \chi^{k-2} & 0 & b + fc \\ & \rho & c \\ & & \chi^{k-1} \end{bmatrix}.$$

So, it remains to show that

$$\begin{bmatrix} \chi^{k-2} & 0 & b \\ & \rho & c \\ & & \chi^{k-1} \end{bmatrix} \cong \chi^{k-2} \oplus \begin{bmatrix} \rho & * \\ & \chi^{k-1} \end{bmatrix}.$$

Note that we have

$$\begin{bmatrix} 0_{n_2 \times n_1} & I_{n_2} & \\ & 0_{n_1 \times n_2} & \\ & & I_{n_3} \end{bmatrix} \begin{bmatrix} \chi^{k-2} & 0 & b \\ & \rho & c \\ & & \chi^{k-1} \end{bmatrix} \begin{bmatrix} 0_{n_2 \times n_1} & I_{n_2} & \\ I_{n_1} & 0_{n_1 \times n_2} & \\ & & I_{n_3} \end{bmatrix}^{-1} = \begin{bmatrix} \rho & \chi^{k-2} & c \\ & \chi^{k-2} & b \\ & & \chi^{k-1} \end{bmatrix}.$$

By the same argument as before $b \in H^1(G_\Sigma, \text{Hom}_{\mathbf{F}}(\chi^{k-1}, \chi^{k-2}))$ must again be a coboundary, i.e., $b(g) = \chi^{k-2}(g)f - f\rho(g)$ for some matrix f . But then

$$\text{again the matrix } \begin{bmatrix} I_{n_1} & & \\ & I_{n_2} & f \\ & & I_{n_3} \end{bmatrix} \text{ conjugates } \begin{bmatrix} \rho & & c \\ & \chi^{k-2} & b \\ & & \chi^{k-1} \end{bmatrix} \text{ to } \begin{bmatrix} \rho & & c \\ & \chi^{k-2} & 0 \\ & & \chi^{k-1} \end{bmatrix} \cong$$

$\chi^{k-2} \oplus \begin{bmatrix} \rho & *3 \\ & \chi^{k-1} \end{bmatrix}$, which leads to a contradiction to (4.3). The fact that $\bar{\sigma}_\Lambda$ is semi-abelian follows again from the minimality of σ . \square

Corollary 4.4. *Let $\sigma, \tau : G_\Sigma \rightarrow \text{GL}_4(E)$ be two absolutely irreducible short crystalline at p Galois representations such that their restriction to I_ℓ is isomorphic to $\exp(t_p N_1)$ for all $\ell \in \Sigma \setminus \{p\}$. Suppose that $\bar{\sigma}^{\text{ss}} = \bar{\tau}^{\text{ss}} = \chi^{k-2} \oplus \rho \oplus \chi^{k-1}$ with ρ absolutely irreducible. Suppose that $\dim_{\mathbf{F}} H_f^1(\mathbf{Q}, \rho(1-k)) \leq 1$. Then there exists a G_Σ -stable lattice Λ in the space of σ and a G_Σ -stable lattice Λ' in the space of τ such that*

$$\bar{\sigma}_\Lambda = \begin{bmatrix} \chi^{k-2} & a & b \\ & \rho & c \\ & & \chi^{k-1} \end{bmatrix} \quad \text{and} \quad \bar{\tau}_{\Lambda'} = \begin{bmatrix} \chi^{k-2} & a & b' \\ & \rho & c \\ & & \chi^{k-1} \end{bmatrix},$$

both semi-abelian, with a, c both non-trivial elements of $H_f^1(\mathbf{Q}, \rho(1-k))$.

Remark 4.5. Note that in Corollary 4.4 we only assume an upper bound on the dimension of $H_f^1(\mathbf{Q}, \rho(1-k))$, but in fact as argued at the beginning of the proof of the corollary our assumptions imply that we always have $\dim_{\mathbf{F}} H_f^1(\mathbf{Q}, \rho(1-k)) \geq 1$.

Proof. By Corollary 4.3 we can find lattices, so that both reductions have the above shape, are short crystalline at p and semi-abelian. By Corollary 3.3 we know that the extensions induced by the entries a and c lie in $H_f^1(\mathbf{Q}, \rho(1-k))$. Since the latter group is one-dimensional we can conjugate one of them to ensure that both representations have the same a -entries and c -entries. \square

5. UNIQUENESS OF ITERATED RESIDUAL EXTENSIONS

In this section we assume that ρ is the mod ϖ reduction of $\rho_f : G_{\Sigma} \rightarrow \mathrm{GL}_2(E)$, the Galois representation attached to a newform $f \in S_{2k-2}(N)$, where N squarefree, $p \nmid N$, k is even and $\Sigma = \{\ell \mid N\} \cup \{p\}$. We also assume that ρ is absolutely irreducible and ramified at every $\ell \mid N$. By Proposition 2.5 (see Corollary 2.6) we then have that $H_{f,\min}^1(\mathbf{Q}, \rho(1-k)) = H_{f,\mathrm{ur}}^1(\mathbf{Q}, \rho(1-k))$ which we will simply abbreviate to $H_f^1(\mathbf{Q}, \rho(1-k))$. We assume in this section that

$$\dim_{\mathbf{F}} H_f^1(\mathbf{Q}, \rho(1-k)) = 1.$$

The goal of this section is to prove the following proposition.

Proposition 5.1. *Let $\bar{\sigma}, \bar{\tau} : G_{\Sigma} \rightarrow \mathrm{GL}_4(\mathbf{F})$ be two Galois representations of the form*

$$\bar{\sigma} = \begin{bmatrix} \chi^{k-2} & a & b \\ & \rho & c \\ & & \chi^{k-1} \end{bmatrix}, \quad \bar{\tau} = \begin{bmatrix} \chi^{k-2} & a' & b' \\ & \rho & c' \\ & & \chi^{k-1} \end{bmatrix}$$

with the extensions $\begin{bmatrix} \chi^{k-2} & a \\ & \rho \end{bmatrix}$, $\begin{bmatrix} \chi^{k-2} & a' \\ & \rho \end{bmatrix}$, $\begin{bmatrix} \rho & c \\ & \chi^{k-1} \end{bmatrix}$ and $\begin{bmatrix} \rho & c' \\ & \chi^{k-1} \end{bmatrix}$ all giving rise to non-zero elements of $H_f^1(\mathbf{Q}, \rho(1-k)) = \mathbf{F}$. Suppose also that both $\bar{\sigma}$ and $\bar{\tau}$ are short crystalline and semi-abelian. Then $\bar{\sigma} \cong \bar{\tau}$.

Proof. Denote the cohomology classes corresponding to the extensions in the statement of the Proposition by ϕ, ϕ', ψ, ψ' respectively. Note that they indeed give rise to elements in H_f^1 by Corollary 3.3. Then there exist $\alpha, \gamma \in \mathbf{F}^{\times}$ such that $\phi' = \alpha\phi$ and $\psi' = \gamma\psi$. The functions a, a', c, c' are given by $a(g) = \phi(g)\rho(g)$, $a'(g) = \phi'(g)\rho(g)$, $c(g) = \psi(g)\chi^{k-1}(g)$ and $c'(g) = \psi'(g)\chi^{k-1}(g)$, so in particular $a' = \alpha a$ and $c' = \gamma c$. Hence

$$\begin{bmatrix} 1 & & \\ & \alpha I_2 & \\ & & \alpha\gamma \end{bmatrix} \bar{\tau} \begin{bmatrix} 1 & & \\ & \alpha I_2 & \\ & & \alpha\gamma \end{bmatrix}^{-1} = \begin{bmatrix} \chi^{k-2} & a & \alpha^1\gamma^{-1}b \\ & \rho & c \\ & & \chi^{k-1} \end{bmatrix}.$$

Thus we may assume without loss of generality that $a' = a$ and $c' = c$. We will express $\bar{\sigma}$ as an iterated extension and apply Lemma 2.3 to show that such extensions are a torsor under $H_f^1(\mathbf{Q}, \mathbf{F}(-1)) = 0$ from which the uniqueness of such $\bar{\sigma}$ will follow.

Consider $\bar{\sigma}$ as in the statement of the Proposition. Its subrepresentation corresponding to the 3×3 upper-left block gives rise to the following exact sequence of G_Σ -representations:

$$(5.1) \quad 0 \rightarrow \mathrm{Hom}_{\mathbf{F}}(\chi^{k-1}, \chi^{k-2}) \rightarrow \mathrm{Hom}_{\mathbf{F}}(\chi^{k-1}, \begin{bmatrix} \chi^{k-2} & a \\ & \rho \end{bmatrix}) \rightarrow \mathrm{Hom}_{\mathbf{F}}(\chi^{k-1}, \rho) \rightarrow 0,$$

and after restricting to the decomposition group at v , also to a corresponding short exact sequence of $G_{\mathbf{Q}_v}$ -representations.

We first consider the case $v = p$: Write \mathcal{M} for the category of filtered Dieudonne modules as defined in [CHT08] Section 2.4.1. Since \mathcal{M} is closed under subquotients, there exist objects M_1, M_2, M_3, M of \mathcal{M} corresponding to the restrictions to $G_{\mathbf{Q}_p}$ of χ^{k-2} , ρ , χ^{k-1} and the subrepresentation $\begin{bmatrix} \chi^{k-2} & a \\ & \rho \end{bmatrix} : G_\Sigma \rightarrow \mathrm{GL}_3(\mathbf{F})$ respectively. Since $\mathrm{Hom}(\chi^{k-1}, M) = M(1-k)$ we see that all the objects in (5.1) are in the essential image of \mathbf{G} . Hence we can apply Lemma 5.3 in [BK13] to conclude that it gives rise to an exact sequence of Selmer groups

$$(5.2) \quad \begin{aligned} H^0(\mathbf{Q}_p, \mathrm{Hom}_{\mathbf{F}}(\chi^{k-1}, \rho)) &\rightarrow H_f^1(\mathbf{Q}_p, \mathrm{Hom}_{\mathbf{F}}(\chi^{k-1}, \chi^{k-2})) \rightarrow H_f^1(\mathbf{Q}_p, \mathrm{Hom}_{\mathbf{F}}(\chi^{k-1}, \begin{bmatrix} \chi^{k-2} & a \\ 0 & \rho \end{bmatrix})) \\ &\rightarrow H_f^1(\mathbf{Q}_p, \mathrm{Hom}_{\mathbf{F}}(\chi^{k-1}, \rho)) \rightarrow 0. \end{aligned}$$

It follows from (5.2) that (in the terminology of [Wes00], p.4) the finite-singular structures on the first and last non-zero term in the $G_{\mathbf{Q}_p}$ sequence (5.1) are induced from the middle term.

Let us now show that this is also true for all primes $v \neq p$ when we take H_f^1 to be the unramified structure defined in section 2. So, suppose that $v \mid N$. It suffices to show that the sequence

$$(5.3) \quad \begin{aligned} H^0(\mathbf{Q}_v, \mathrm{Hom}_{\mathbf{F}}(\chi^{k-1}, \rho)) &\rightarrow H_{\mathrm{ur}}^1(\mathbf{Q}_v, \mathrm{Hom}_{\mathbf{F}}(\chi^{k-1}, \chi^{k-2})) \rightarrow H_{\mathrm{ur}}^1(\mathbf{Q}_v, \mathrm{Hom}_{\mathbf{F}}(\chi^{k-1}, \begin{bmatrix} \chi^{k-2} & a \\ 0 & \rho \end{bmatrix})) \\ &\rightarrow H_{\mathrm{ur}}^1(\mathbf{Q}_v, \mathrm{Hom}_{\mathbf{F}}(\chi^{k-1}, \rho)) \rightarrow 0 \end{aligned}$$

with $H_{\mathrm{ur}}^1(\mathbf{Q}_v, M) := \ker(H^1(\mathbf{Q}_v, M) \rightarrow H^1(I_v, M))$ is exact. Let us rewrite (5.1) as

$$(5.4) \quad 0 \rightarrow \mathbf{F}(-1) \rightarrow \begin{bmatrix} \chi^{-1} & a' \\ & \rho(1-k) \end{bmatrix} \rightarrow \rho(1-k) \rightarrow 0$$

and call the extension in the middle \mathcal{E} .

By Corollary 3.3 we have $\mathcal{E}^{I_v} = \mathbf{F}(-1)^{I_v} \oplus \rho(1-k)^{I_v}$, so in particular the sequence

$$0 \rightarrow \mathbf{F}(-1)^{I_v} \rightarrow \mathcal{E}^{I_v} \rightarrow \rho(1-k)^{I_v} \rightarrow 0$$

is exact. Since every module in that sequence has an action of $G := \mathrm{Gal}(\mathbf{Q}_v^{\mathrm{ur}}/\mathbf{Q}_v)$, we get a long exact sequence in cohomology of that group

(and note that $(M^{I_v})^G = M^{G_v}$, where $G_v = G_{\mathbf{Q}_v}$):

$$(5.5) \quad 0 \rightarrow \mathbf{F}(-1)^{G_v} \rightarrow \mathcal{E}^{G_v} \rightarrow \rho(1-k)^{G_v} \rightarrow H^1(\mathbf{Q}_v^{\text{ur}}/\mathbf{Q}_v, \mathbf{F}(-1)^{I_v}) \rightarrow \\ H^1(\mathbf{Q}_v^{\text{ur}}/\mathbf{Q}_v, \mathcal{E}^{I_v}) \rightarrow H^1(\mathbf{Q}_v^{\text{ur}}/\mathbf{Q}_v, \rho(1-k)^{I_v}) \rightarrow H^2(\mathbf{Q}_v^{\text{ur}}/\mathbf{Q}_v, \mathbf{F}(-1)^{I_v})$$

Note that the last group is zero since G has cohomological dimension one. For all three modules M one has $H^1(\mathbf{Q}_v^{\text{ur}}/\mathbf{Q}_v, M^{I_v}) = H_{\text{ur}}^1(\mathbf{Q}_v, M)$ by Lemma 3.2 (i) in [Rub00].

Since we now showed that the finite-singular structures on the first and last non-zero term in the sequence 5.1 are induced from the middle term for all $G_{\mathbf{Q}_v}$, Lemma 2.3 tells us that we have an exact sequence

$$(5.6) \quad H_f^1(\mathbf{Q}, \text{Hom}_{\mathbf{F}}(\chi^{k-1}, \chi^{k-2})) \rightarrow H_f^1(\mathbf{Q}, \text{Hom}_{\mathbf{F}}(\chi^{k-1}, \begin{bmatrix} \chi^{k-2} & a \\ 0 & \rho \end{bmatrix})) \rightarrow H_f^1(\mathbf{Q}, \text{Hom}_{\mathbf{F}}(\chi^{k-1}, \rho)).$$

Since $H_f^1(\mathbf{Q}, \text{Hom}_{\mathbf{F}}(\chi^{k-1}, \chi^{k-2})) = H_f^1(\mathbf{Q}, \mathbf{F}(-1)) = 0$ by Proposition 2.9, we see that the middle Selmer group in (5.6) injects into the last one. Hence the cohomology class in $H_f^1(\mathbf{Q}, \text{Hom}_{\mathbf{F}}(\chi^{k-1}, \rho))$ corresponding to the quotient representation $\begin{bmatrix} \rho & c \\ 0 & \chi^{k-1} \end{bmatrix}$ of $\bar{\sigma}$ determines a short crystalline extension of χ^{k-1} by $\begin{bmatrix} \chi^{k-2} & a \\ 0 & \rho \end{bmatrix}$ uniquely. Since both $\bar{\sigma}$ and $\bar{\tau}$ are such extensions, we get $\bar{\sigma} \cong \bar{\tau}$. \square

6. THE IDEALS OF REDUCIBILITY

Let $\rho_f : G_{\Sigma} \rightarrow \text{GL}_2(E)$ be the Galois representation attached to a newform $f \in S_{2k-2}(N)$ for N a square-free integer with $p \nmid N$ and $\Sigma = \{\ell \mid N\} \cup \{p\}$. Suppose also that the residual representation $\rho := \bar{\rho}_f$ is absolutely irreducible and ramified at every prime $\ell \mid N$. Then $\rho|_{I_{\ell}} \cong \begin{bmatrix} 1 & * \\ & 1 \end{bmatrix}$ (see (2.2)). Let

$$\bar{\sigma} = \begin{bmatrix} \chi^{k-2} & a & b \\ & \rho & c \\ & & \chi^{k-1} \end{bmatrix} : G_{\Sigma} \rightarrow \text{GL}_4(\mathbf{F})$$

be short crystalline at p and semi-abelian with a, c non-trivial classes in $H_f^1(\mathbf{Q}, \rho(1-k))$.

The universal deformation σ^{red} (resp. its trace) gives rise to an R^{red} -algebra morphism (which we denote by the same letter) $\sigma^{\text{red}} : R^{\text{red}}[G_{\Sigma}] \rightarrow M_4(R^{\text{red}})$ (resp. $T = \text{tr } \sigma^{\text{red}} : R^{\text{red}}[G_{\Sigma}] \rightarrow R^{\text{red}}$). We define $\ker \sigma^{\text{red}}$ (resp. $\ker T$) as in [BC09], section 1.2.4.

Since R , and hence also R^{red} , is Noetherian, both rings have a finite number of minimal primes. Then the total ring of fractions $\text{Tot}(R^{\text{red}})$ of R^{red} is a finite product of fields. In fact we have an injection

$$R^{\text{red}} \hookrightarrow \text{Tot}(R^{\text{red}}) = \prod_{\mathcal{P}} R_{\mathcal{P}}^{\text{red}},$$

where \mathcal{P} runs over minimal primes of R^{red} and the localization $R_{\mathcal{P}}^{\text{red}}$ is a field (cf. [BC09], Proposition 1.3.11).

To ease notation in this section we will write G for G_{Σ} . Let us note that both $R^{\text{red}}[G]/\ker T$ and $R^{\text{red}}[G]/\ker \sigma^{\text{red}}$ are Cayley-Hamilton quotients of $R^{\text{red}}[G]$ in the sense of [BC09], section 1 and we have a canonical R^{red} -algebra map $\varphi : R^{\text{red}}[G]/\ker \sigma^{\text{red}} \rightarrow R^{\text{red}}[G]/\ker T$.

Theorem 6.1. *One has the following:*

- (i) *Let $? \in \{T, \sigma\}$ and $i, j \in \{k-1, \rho, k-2\}$. Let $S^?$ be $R^{\text{red}}[G]/\ker T$ (when $? = T$) or $R^{\text{red}}[G]/\ker \sigma^{\text{red}}$ (when $? = \sigma$). Then there are data \mathcal{E} of idempotents for which $S^?$ is a GMA (Generalized Matrix Algebra) and there exist R^{red} -submodules $\mathcal{A}_{i,j}^?$ of $S^?$ which satisfy*

$$\mathcal{A}_{i,j}^? \mathcal{A}_{j,k}^? \subset \mathcal{A}_{i,k}^?, \quad T : \mathcal{A}_{i,i}^? \xrightarrow{\sim} R^{\text{red}}, \quad T(\mathcal{A}_{i,j}^? \mathcal{A}_{j,i}^?) \subset \mathfrak{m}_{R^{\text{red}}},$$

and

$$S^? \cong \text{GMA}(\mathcal{A}^?) := \begin{bmatrix} \mathcal{A}_{k-2,k-2}^? & M_{1,2}(\mathcal{A}_{k-2,\rho}^?) & \mathcal{A}_{k-2,k-1}^? \\ M_{2,1}(\mathcal{A}_{\rho,k-2}^?) & M_{2,2}(\mathcal{A}_{\rho,\rho}^?) & M_{2,1}(\mathcal{A}_{\rho,k-1}^?) \\ \mathcal{A}_{k-1,k-2}^? & M_{1,2}(\mathcal{A}_{k-1,\rho}^?) & \mathcal{A}_{k-1,k-1}^? \end{bmatrix}.$$

The isomorphism $S^? \cong \text{GMA}(\mathcal{A}^?)$ (and also $S^? \cong \text{GMA}(A^?)$ below) is an R^{red} -algebra morphism.

- (ii) *For \mathcal{E} as in (i) one has*

$$S^? \cong \text{GMA}(A^?) := \begin{bmatrix} R^{\text{red}} & M_{1,2}(A_{k-2,\rho}^?) & A_{k-2,k-1}^? \\ M_{2,1}(A_{\rho,k-2}^?) & M_{2,2}(R^{\text{red}}) & M_{2,1}(A_{\rho,k-1}^?) \\ A_{k-1,k-2}^? & M_{1,2}(A_{k-1,\rho}^?) & R^{\text{red}} \end{bmatrix} \subset M_4(\text{Tot}(R^{\text{red}})),$$

where $A_{i,j}^?$ are fractional ideals of R^{red} (and $A_{i,j}^{\sigma}$ are ideals of R^{red}) that satisfy the Chasles relations

$$A_{i,j}^? \mathcal{A}_{j,i}^? \subset \mathcal{A}_{i,k}^?, \quad A_{i,i}^? = R^{\text{red}}, \quad A_{i,j}^? \mathcal{A}_{j,i}^? \subset \mathfrak{m}_{R^{\text{red}}}.$$

- (iii) *The data \mathcal{E} can be adapted so that the R^{red} -algebra map $\varphi' : \text{GMA}(\mathcal{A}^{\sigma}) \rightarrow \text{GMA}(\mathcal{A}^T)$ induced from φ has the property that $\varphi'(\mathcal{A}_{i,j}^{\sigma}) = \mathcal{A}_{i,j}^T$.*

Proof. The statements (i), (ii) for $? = T$ and (i) for $? = \sigma$ are a direct consequence of Theorem 1.4.4 in [BC09]. To show that we can also obtain (ii) for $? = \sigma$ we argue as in [BK13], Proposition 2.8 which uses Lemma 1.3.7 of [BC09]. Finally, we proceed as in the proof of Lemma 2.5 in [BK13] to show that \mathcal{E} can be adapted to satisfy (iii). \square

Remark 6.2. The data $\mathcal{E} = \{e_i, \psi_i \mid i \in \{\chi^{k-2}, \rho, \chi^{k-1}\}\}$ in Theorem 6.1 can be (and will be) chosen so that for $i \in \{\chi^{k-2}, \rho, \chi^{k-1}\}$ the maps $\psi_i : e_i S^{\sigma} e_i \xrightarrow{\sim} M_{d_i, d_i}(R^{\text{red}})$ satisfy $\psi_i \otimes \mathbf{F} \cong i$. Here $d_i = \dim i$. From now on fix \mathcal{E} .

Using Theorem 6.1 we get the following commutative diagram of R^{red} -algebras:

$$\begin{array}{ccc} \text{GMA}(\mathcal{A}^\sigma) & \xleftarrow[\iota_1]{\sim} R^{\text{red}}[G_\Sigma]/\ker \sigma^{\text{red}} & \xrightarrow[\sim]{\iota_2} \text{GMA}(A^\sigma) \\ \downarrow \varphi' & & \downarrow \varphi \\ \text{GMA}(\mathcal{A}^T) & \xleftarrow[\iota_3]{\sim} R^{\text{red}}[G_\Sigma]/\ker T & \xrightarrow[\sim]{\iota_4} \text{GMA}(A^T) \end{array}$$

where the maps ι_i are the ones given by Theorem 6.1. Using Lemma 1.3.8 (resp. Proposition 1.3.12) of [BC09] we get that the composite $\iota_2 \circ \iota_1^{-1}$ (resp. $\iota_4 \circ \iota_3^{-1}$) give rise to isomorphisms of R^{red} -modules $f_{i,j}^\sigma : \mathcal{A}_{i,j}^\sigma \xrightarrow{\sim} A_{i,j}^\sigma$ (resp. $f_{i,j}^T : \mathcal{A}_{i,j}^T \xrightarrow{\sim} A_{i,j}^T$). We define R^{red} -module surjections $\phi_{i,j}$ to make the following diagram commute

$$(6.1) \quad \begin{array}{ccc} \mathcal{A}_{i,j}^\sigma & \xrightarrow{f_{i,j}^\sigma} & A_{i,j}^\sigma \\ \varphi' \downarrow & & \downarrow \phi_{i,j} \\ \mathcal{A}_{i,j}^T & \xrightarrow{f_{i,j}^T} & A_{i,j}^T \end{array}$$

Definition 6.3 ([BC09] Definition 1.5.2). Let $\mathcal{P} = (\mathcal{P}_1, \dots, \mathcal{P}_s)$ be a partition of the set $\mathcal{I} = \{\chi^{k-2}, \rho, \chi^{k-1}\}$. The ideal of reducibility $I^{\mathcal{P}}$ (associated with partition \mathcal{P}) is the smallest ideal I of R^{red} with the property that there exist pseudocharacters $T_1, \dots, T_s : R^{\text{red}}[G]/IR^{\text{red}}[G] \rightarrow R^{\text{red}}/I$ such that

- (i) $T \otimes R^{\text{red}}/I = \sum_{l=1}^s T_l$,
- (ii) for each $l \in \{1, \dots, s\}$, $T_l \otimes \mathbf{F} = \sum_{\rho' \in \mathcal{P}_l} \text{tr } \rho'$.

To shorten notation we will sometimes write $k-1, k-2, \rho$ for the elements of \mathcal{I} instead of $\chi^{k-1}, \chi^{k-2}, \rho$.

Proposition 6.4 (Proposition 1.5.1 in [BC09]). *For every partition \mathcal{P} the corresponding ideal of reducibility $I^{\mathcal{P}}$ exists. Furthermore, let S be any Cayley-Hamilton quotient of (R^{red}, T) (we will only use (R^{red}, T) and $(R^{\text{red}}, \sigma^{\text{red}})$) and choose data of idempotents \mathcal{E} as in Theorem 6.1 so that $S \cong \text{GMA}(\mathcal{A}^?)$. Then $I^{\mathcal{P}}$ is given by the following formula (whose sides do not depend on the choice of S or \mathcal{E})*

$$I^{\mathcal{P}} = \sum_{\substack{(i,j) \\ i,j \text{ not in the same } \mathcal{P}_l}} T(\mathcal{A}_{i,j}^? \mathcal{A}_{j,i}^?).$$

Corollary 6.5. *For $? \in \{T, \sigma^{\text{red}}\}$ let $A_{i,j}^?$ be defined as in Theorem 6.1 (ii). Then one has*

$$I^{\mathcal{P}} = \sum_{\substack{(i,j) \\ i,j \text{ not in the same } \mathcal{P}_l}} A_{i,j}^? A_{j,i}^?.$$

Proof. Multiplication between elements of $\alpha \in \mathcal{A}_{i,j}^?$ and $\beta \in \mathcal{A}_{j,k}^?$ corresponds to correct matrix multiplication, i.e., one puts α in the (i, j) th spot (which may be a block) of a matrix and β in the (j, k) th spot and completes

both matrices by putting zeros elsewhere. Then $\alpha\beta$ is the (i, k) th spot in the matrix obtained as a product of the matrices above. The corollary follows from the commutativity of the following diagram

$$\begin{array}{ccc} \mathcal{A}_{i,j}^? \mathcal{A}_{j,i}^? & \hookrightarrow & \mathcal{A}_{i,i} \\ f_{i,j}^? \otimes f_{j,i}^? \downarrow & & \sim \downarrow T \\ \mathcal{A}_{i,j}^? \mathcal{A}_{j,i}^? & \hookrightarrow & R^{\text{red}} \end{array}$$

since $f_{i,i}^? = T : \mathcal{A}_{i,i} \rightarrow R^{\text{red}}$. \square

In our situation we have 4 possible partitions of \mathcal{P} and the following 4 corresponding ideals of reducibility.

(6.2)

$$\begin{aligned} \mathcal{P} = \{\chi^{k-2}, \rho\} \cup \{\chi^{k-1}\} &\implies I^{\mathcal{P}} = I^{k-1} = A_{k-1,k-2}A_{k-2,k-1} + A_{\rho,k-1}A_{k-1,\rho} \\ \mathcal{P} = \{\chi^{k-1}, \rho\} \cup \{\chi^{k-2}\} &\implies I^{\mathcal{P}} = I^{k-2} = A_{k-1,k-2}A_{k-2,k-1} + A_{\rho,k-2}A_{k-2,\rho} \\ \mathcal{P} = \{\chi^{k-2}, \chi^{k-1}\} \cup \{\rho\} &\implies I^{\mathcal{P}} = I^{\rho} = A_{k-2,\rho}A_{\rho,k-2} + A_{k-1,\rho}A_{\rho,k-1} \\ \mathcal{P} = \{\chi^{k-2}\} \cup \{\rho\} \cup \{\chi^{k-1}\} &\implies I^{\mathcal{P}} = I^{\text{tot}} = A_{k-1,k-2}A_{k-2,k-1} + A_{\rho,k-1}A_{k-1,\rho} \\ &\quad + A_{\rho,k-2}A_{k-2,\rho} \end{aligned}$$

We have the following analogue of [BC09], Lemma 9.3.1:

Theorem 6.6. *One has*

- (i) *All the ideals of reducibility coincide with I^{tot} ,*
- (ii) $I^{\text{tot}} = A_{k-2,\rho}^T A_{\rho,k-2}^T$,
- (iii) *If, in addition, $\dim_{\mathbf{F}} H_f^1(\mathbf{Q}, \rho(2-k)) = 1$ and $p \nmid 1 + w_{f,\ell}$ for all $\ell \mid N$, then I^{tot} is a principal ideal of R^{red} .*

Proof. We will prove Theorem 6.6 by a sequence of Lemmas some pertaining to $A_{i,j}^T$ and some to $A_{i,j}^{\sigma}$.

Lemma 6.7. *One has*

$$(6.3) \quad A_{k-2,\rho}^T A_{\rho,k-2}^T = A_{k-1,\rho}^T A_{\rho,k-1}^T.$$

Proof. This is proved exactly as Lemma 8.2.16 of [BC09] using the antiinvolution τ . \square

We will need the following analogue of [BC09], Lemma 8.3.1:

Lemma 6.8. *We have*

$$A_{k-1,k-2}^{\sigma} = A_{k-1,\rho}^{\sigma} A_{\rho,k-2}^{\sigma}.$$

The strategy to prove Lemma 6.8 follows that of [BC09], Lemma 8.3.1, but instead of using Ext_T we use Lemma 6.9 below. For $\rho_i, \rho_j \in \{\chi^{k-1}, \chi^{k-2}, \rho\}$, $\rho_i \not\cong \rho_j$, set $A'_{i,j} := A_{i,l}^{\sigma} A_{l,j}^{\sigma}$, where $\rho_l \in \{\chi^{k-1}, \chi^{k-2}, \rho\}$, $\rho_i \not\cong \rho_l \not\cong \rho_j$.

Lemma 6.9. *There is an injection*

$$\text{Hom}_{R^{\text{red}}}(A_{i,j}^{\sigma}/A'_{i,j}, \mathbf{F}) \hookrightarrow H_{\Sigma}^1(\mathbf{Q}, \text{Hom}(\rho_j, \rho_i)).$$

If either (i) or (ii) hold, where

- (i) $\rho_i = \chi^{k-2}$ and $\rho_j = \chi^{k-1}$,
- (ii) $\rho_i = \chi^{k-1}$ and $\rho_j = \chi^{k-2}$,

then the image of the injection is contained in H_f^1 .

Proof. For simplicity in this proof only we write R for R^{red} . By [BC09], Theorem 1.5.5, taking $J = \mathfrak{m}$ there is an injection

$$\iota_{i,j} : \text{Hom}_R(A_{i,j}^\sigma/A'_{i,j}, \mathbf{F}) \hookrightarrow \text{Ext}_{R[G]/\mathfrak{m}R[G]}^1(\rho_j, \rho_i).$$

We have

$$\mathfrak{m}R[G] = \ker(R[G] \rightarrow \mathbf{F}[G]) = (\mathfrak{m}R)[G],$$

hence $R[G]/\mathfrak{m}R[G] = (R/\mathfrak{m}R)[G] = \mathbf{F}[G]$. Thus we get an injection

$$\iota_{i,j} : \text{Hom}_R(A_{i,j}^\sigma/A'_{i,j}, \mathbf{F}) \hookrightarrow H^1(\mathbf{Q}, \text{Hom}(\rho_j, \rho_i)).$$

It remains to prove that the image is contained in the correct Selmer group. By [BC09], Theorem 1.5.6(1) the image consists precisely of the $S/\mathfrak{m}S$ -extensions of ρ_i by ρ_j , where

$$S := (R[G]) / (\ker \sigma)(R[G]).$$

By [BC09], Theorem 1.5.6(2) any $S/\mathfrak{m}S$ -extension is a quotient of $M_j/\mathfrak{m}M_j \oplus \rho_i$, where $M_j = SE_j$ and E_j are defined as in [BC09], p. 21. Since $p > 2k-2$, the representations χ^{k-2} , χ^{k-1} and ρ are short crystalline. Since the category of such representations is closed under taking subobjects, quotients and finite direct sums it suffices therefore to prove that M_j is short crystalline. By [BC09], section 1.5.4, one has $S = M_j \oplus S(1 - E_j)$, hence in particular M_j is an S -submodule, and hence also an $\mathcal{O}[G]$ -submodule of S . One has $M_j \subset M_j \otimes \text{Tot}(R^{\text{red}})$ and the latter Galois module is isomorphic to the representation $(\text{Tot}(R^{\text{red}})^4, \sigma^{\text{red}})$, which is short crystalline. Thus M_j is short crystalline.

Let us now check that if $\rho_i = \chi^{k-2}$ and $\rho_j = \chi^{k-1}$ (or vice versa), then the extensions in the image of $\iota_{i,j}$ are unramified away from p . Let $\ell \mid N$ be a prime. By Remark 6.2 we can conjugate σ^{red} so that it is adapted to the data of idempotents \mathcal{E} . Abusing notation we will in this proof denote this conjugate still by σ^{red} . This implies in particular that $e_\rho(\sigma \otimes \mathbf{F})e_\rho \cong \rho$ and that

$$(6.4) \quad \sigma^{\text{red}}(S^\sigma) = \begin{bmatrix} R & M_{1,2}(A_{k-2,\rho}^\sigma) & A_{k-2,k-1}^\sigma \\ M_{2,1}(A_{\rho,k-2}^\sigma) & M_{2,2}(R) & M_{2,1}(A_{\rho,k-1}^\sigma) \\ A_{k-1,k-2}^\sigma & M_{1,2}(A_{k-1,\rho}^\sigma) & R \end{bmatrix}.$$

Let $X \in I_\ell$ be such that $\sigma^{\text{red}}(X)$ topologically generates $\sigma^{\text{red}}(I_\ell)$. Write $\alpha_{k,l}$ for the (k,l) -entry of $Y := \sigma^{\text{red}}(X) - I_4$. Since ρ is ramified at ℓ at least one of the entries $\alpha_{2,2}$, $\alpha_{2,3}$, $\alpha_{3,2}$, $\alpha_{3,3}$ lies in R^\times . To fix attention let us assume that $\alpha_{2,3} \in R^\times$. The proof in the other three cases is identical.

The construction of the extensions in the image of $\iota_{i,j}$ is given in [BC09], p. 37. Note that while there are in general three choices for i (corresponding to $\rho_i = \chi^{k-2}$, χ^{k-1} or ρ), and the same holds for j , there are four choices for k and for l .

If $\rho_i = \rho_j$, let $a_{ii} : S/\mathfrak{m}S \rightarrow e_{\rho_i}(S/\mathfrak{m}S)e_{\rho_i}$ be the canonical projection defined in Lemma 1.5.4 of [BC09], where e_{ρ_i} is the idempotent corresponding

to ρ_i . (For notation and terminology see [BC09].) We note that a_{ii} composed with the isomorphism $e_{\rho_i}(S/\mathfrak{m}S)e_{\rho_i} \cong \mathbf{F}$ gives a representation isomorphic to ρ_i . If $\rho_i \neq \rho_j$, then the projection $a_{ij} : S/\mathfrak{m}S \rightarrow e_{\rho_i}(S/\mathfrak{m}S)e_{\rho_j}$ is defined in an analogous way and is the mod \mathfrak{m} -reduction of the canonical projection $\tilde{a}_{ij} : S \rightarrow e_{\rho_i}Se_{\rho_j}$ defined again in the same way. The codomain of \tilde{a}_{ij} is isomorphic to \mathcal{A}_{ij}^σ (via, say, a map ϕ) which then is isomorphic to A_{ij}^σ via the map $f_{i,j}^\sigma$. The composite $f_{i,j}^\sigma \circ \phi \circ \tilde{a}_{ij}$ equals $\epsilon_i \sigma^{\text{red}} \epsilon_j$, where

$$\epsilon_i = \begin{cases} \text{diag}(1, 0, 0, 0) & \text{if } \rho_i = \chi^{k-2} \\ \text{diag}(0, 1, 1, 0) & \text{if } \rho_i = \rho \\ \text{diag}(0, 0, 0, 1) & \text{if } \rho_i = \chi^{k-1}. \end{cases}$$

Here we are only interested in the case when $\rho_i = \chi^{k-1}$ and $\rho_j = \chi^{k-2}$ or vice versa (but the proof in the second case is identical to the first, so we omit it).

Let $f \in \text{Hom}(A_{\chi,1}^\sigma/A'_{\chi,1}, \mathbf{F})$. Recall that we have $A'_{\chi,1} = A_{\chi,\rho}^\sigma A_{\rho,1}^\sigma$. Since χ and 1 are unramified to show that the extension

$$\begin{bmatrix} a_{\chi\chi} & f \circ a_{\chi 1} \\ & a_{11} \end{bmatrix}$$

is unramified at ℓ it suffices to show that $a_{\chi 1}(I_\ell) \subset A'_{\chi,1}$. Note that $a_{\chi 1}(X) =$

$$\alpha_{4,1}. \text{ Since } Y \text{ is a conjugate of } \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ and } A_{k-1,\rho}^\sigma \subset R \text{ (by Theorem}$$

6.1 (ii)) while $\alpha_{2,3} \in R^\times$ we see that the fourth row of Y is a scalar multiple of the second row and that this scalar $\alpha \in \text{Tot}(R)$ in fact lies in R . If $\alpha = 0$, then $a_{\chi 1}(X) = \alpha_{4,1} = 0$ and we are done, otherwise let s be the largest integer such that $\alpha \in \mathfrak{m}^s$. Then $\alpha_{4,1} = \alpha \alpha_{2,1}$. If $\alpha_{2,1} = 0$, we are done, otherwise let r be the largest integer such that $\alpha_{2,1} \in \mathfrak{m}^r$. Then $\alpha_{4,1} = \alpha \alpha_{2,1} \in \mathfrak{m}^{r+s}$. Finally $\alpha_{4,3} = \alpha \alpha_{2,3} \in \mathfrak{m}^s - \mathfrak{m}^{s+1}$ since $\alpha_{2,3}$ is a unit. Comparing the matrix

$$Y = \begin{bmatrix} * & * & * & * \\ \alpha_{2,1} & * & \alpha_{2,3} & * \\ * & * & * & * \\ \alpha \alpha_{2,1} & * & \alpha \alpha_{2,3} & * \end{bmatrix}$$

with (6.4) we see that by the definition of r we get that $A_{\rho,1}^\sigma \supset \mathfrak{m}^r$ (since $A_{\rho,1}^\sigma$ is the ideal of R generated by the entries of σ falling in the corresponding spots and $\alpha_{2,1} \in \mathfrak{m}^r - \mathfrak{m}^{r+1}$). Similarly by the definition of s and the fact that $\alpha_{4,3} \in \mathfrak{m}^s - \mathfrak{m}^{s+1}$ we get that $A_{\chi,\rho}^\sigma \supset \mathfrak{m}^s$ and so, $A'_{\chi,1} = A_{\chi,\rho}^\sigma A_{\rho,1}^\sigma \supset \mathfrak{m}^{r+s}$. Thus, since $\alpha_{4,1} \in \mathfrak{m}^{r+s}$ we get $\alpha_{4,1} \in A'_{\chi,1}$. This completes the proof of the lemma. \square

Proof of Lemma 6.8. We have by Proposition 2.9 that $H_f^1(\mathbf{Q}, \text{Hom}(\chi^{k-2}, \chi^{k-1})) = 0$. Thus by Lemma 6.9 and Nakayama's Lemma we get that $A_{k-1,k-2}^\sigma = A'_{k-1,k-2}$. \square

Lemma 6.10. *We have*

$$A_{k-1,k-2}^T = A_{k-1,\rho}^T A_{\rho,k-2}^T.$$

Proof. This follows by applying $\phi_{k-1,k-2}$ to both sides of the equality in Lemma 6.8 and noting that $\phi_{k-1,k-2}$ preserves multiplication after identifying the elements $\alpha \in A_{k-1,\rho}^\sigma$ and $\beta \in A_{\rho,k-2}^\sigma$ as matrices with non-zero entries in the correct spots, i.e., that one has $\phi_{k-1,k-2}(\alpha\beta) = \phi_{k-1,\rho}(\alpha)\phi_{\rho,k-2}(\beta)$. \square

We can now finish the proof of parts (i) and (ii) of Theorem 6.6. First we apply the Chasles relation to see that one has $A_{k-2,k-1}^T A_{k-1,\rho}^T \subset A_{k-2,\rho}^T$ (cf. Theorem 6.1). So, using Lemma 6.8 one gets $A_{k-2,k-1}^T A_{k-1,k-2}^T \subset A_{k-2,\rho}^T A_{\rho,k-2}^T$. This gives

$$I^{\text{tot}} = A_{k-2,k-1}^T A_{k-1,k-2}^T + A_{k-2,\rho}^T A_{\rho,k-2}^T + A_{k-1,\rho}^T A_{\rho,k-1}^T = A_{k-2,\rho}^T A_{\rho,k-2}^T.$$

This proves (i) and (ii). To prove (iii) we begin with the following lemma.

Lemma 6.11. *One has $A_{k-2,\rho}^\sigma = R^{\text{red}}$ and $A_{\rho,k-2}^\sigma$ is a principal ideal of R^{red} .*

Proof. We have

$$\sigma^{\text{red}}(R^{\text{red}}[G_\Sigma]) \cong \begin{bmatrix} R^{\text{red}} & M_{1,2}(A_{k-2,\rho}^\sigma) & A_{k-2,k-1}^\sigma \\ M_{2,1}(A_{\rho,k-2}^\sigma) & M_{2,2}(R^{\text{red}}) & M_{2,1}(A_{\rho,k-1}^\sigma) \\ A_{k-1,k-2}^\sigma & M_{1,2}(A_{k-1,\rho}^\sigma) & R^{\text{red}} \end{bmatrix}.$$

Recall that

$$\bar{\sigma} = \begin{bmatrix} \chi^{k-2} & a & b \\ & \rho & c \\ & & \chi^{k-1} \end{bmatrix},$$

with a and c both non-trivial classes. Thus a conjugate of σ^{red} which is adapted to the data of idempotents \mathcal{E} will have the corresponding properties. More precisely, $\sigma^{\text{red}} \otimes \mathbf{F}$ will have a subquotient which is a non-split extension of ρ by χ^{k-2} and one which is a non-split extension of χ^{k-1} by ρ . Thus we get that $A_{k-2,\rho}^\sigma = A_{\rho,k-1}^\sigma = R^{\text{red}}$. Then by Chasles relations we get that also $A_{k-2,k-1}^\sigma = R^{\text{red}}$. Finally, by Theorem 6.1(ii) we have that $A_{ij}^\sigma A_{ji}^\sigma \subset \mathfrak{m}$ for each pair $i \neq j$. Thus we get that each of $A_{k-1,k-2}^\sigma$, $A_{k-1,\rho}^\sigma$ and $A_{\rho,k-2}^\sigma$ is contained in \mathfrak{m} .

By Lemma 6.8 we have $A_{k-1,k-2}^\sigma = A_{k-1,\rho}^\sigma A_{\rho,k-2}^\sigma \subset \mathfrak{m} A_{\rho,k-2}^\sigma$. This gives us R^{red} -module maps (note that $A_{\rho,k-1}^\sigma = R^{\text{red}}$)

$$\frac{A_{\rho,k-2}^\sigma}{A_{\rho,k-1}^\sigma A_{k-1,k-2}^\sigma} \twoheadrightarrow \frac{A_{\rho,k-2}^\sigma}{\mathfrak{m} A_{\rho,k-2}^\sigma} \implies \text{Hom}_{R^{\text{red}}} \left(\frac{A_{\rho,k-2}^\sigma}{\mathfrak{m} A_{\rho,k-2}^\sigma}, \mathbf{F} \right) \hookrightarrow \text{Hom}_{R^{\text{red}}} \left(\frac{A_{\rho,k-2}^\sigma}{A_{\rho,k-1}^\sigma A_{k-1,k-2}^\sigma}, \mathbf{F} \right).$$

We have (the first isomorphism following from our assumptions in part (iii) of Theorem 6.6 and from Propositions 2.4(1) and 2.7 while the first

injection following from Lemma 6.9)

$$(6.5) \quad \mathbf{F} \cong H_{\Sigma}^1(\mathbf{Q}, \mathrm{Hom}(\chi^{k-2}, \rho)) \hookrightarrow \mathrm{Hom}_{R^{\mathrm{red}}}(A_{\rho, k-2}^{\sigma}/A'_{\rho, k-2}, \mathbf{F}) = \\ = \mathrm{Hom}_{R^{\mathrm{red}}}\left(\frac{A_{\rho, k-2}^{\sigma}}{A_{\rho, k-1}^{\sigma}A_{k-1, k-2}^{\sigma}}, \mathbf{F}\right) \hookrightarrow \mathrm{Hom}_{R^{\mathrm{red}}}\left(\frac{A_{\rho, k-2}^{\sigma}}{\mathfrak{m}A_{\rho, k-2}^{\sigma}}, \mathbf{F}\right),$$

so, the (clearly non-trivial) \mathbf{F} -vector space $\frac{A_{\rho, k-2}^{\sigma}}{\mathfrak{m}A_{\rho, k-2}^{\sigma}}$ is one-dimensional. Thus, $A_{\rho, k-2}^{\sigma}$ is generated over R by one element by Nakayama's lemma. \square

Lemma 6.12. *One has $A_{k-2, \rho}^T = R^{\mathrm{red}}$ and $A_{\rho, k-2}^T$ is a principal ideal of R^{red} .*

Proof. As explained in the proof of Lemma 6.10 this follows by applying $\phi_{k-2, \rho}$ (resp. $\phi_{\rho, k-2}$) to $A_{k-2, \rho}^{\sigma}$ (resp. $A_{\rho, k-2}^{\sigma}$) and using Lemma 6.11 along with the fact that both ϕ maps are R -linear. \square

Lemma 6.12 along with part (ii) of the Theorem imply part (iii). \square

7. SUFFICIENT CONDITIONS FOR R^{red} TO BE A DISCRETE VALUATION RING

Let $\rho : G_{\Sigma} \rightarrow \mathrm{GL}_2(\mathbf{F})$ be a continuous absolutely irreducible and odd Galois representation of determinant χ^{2k-3} , short crystalline at p and such that for every prime $\ell \in \Sigma \setminus \{p\}$ one has $\rho|_{I_{\ell}} \cong \begin{bmatrix} 1 & * \\ 0 & 1 \end{bmatrix} \neq I_2$. Let R_{ρ} be the universal deformation ring for deformations $\rho' : G_{\Sigma} \rightarrow \mathrm{GL}_2(A)$ (here A is an object of $\mathrm{LCN}(E)$) of ρ which have determinant equal to ϵ^{2k-3} , are short crystalline at p and minimal in the sense that they satisfy

$$(7.1) \quad \rho'|_{I_{\ell}} \cong \begin{bmatrix} 1 & * \\ 0 & 1 \end{bmatrix}.$$

We will assume that $R_{\rho} \cong \mathcal{O}$, i.e., is a discrete valuation ring (cf. [BK13], section 6.1 for discussion of this assumption as well as section 9 in the current paper). This implies that the set of strict equivalence classes of short crystalline, minimal deformations of ρ to $\mathrm{GL}_2(\mathcal{O})$ with determinant equal to ϵ^{2k-3} contains a single element. We will write $\tilde{\rho}$ for a fixed representative of this strict equivalence class.

Consider short crystalline, semi-abelian $\bar{\sigma} : G_{\Sigma} \rightarrow \mathrm{GL}_4(\mathbf{F})$ of the form

$$\bar{\sigma} = \begin{bmatrix} \chi^{k-2} & a & b & \\ & \rho & c & \\ & & & \chi^{k-1} \end{bmatrix}$$

with a, c giving rise to non-zero elements of $H_f^1(\mathbf{Q}, \rho(1-k))$ (cf. Corollary 3.3).

We assume in this section that

$$\dim_{\mathbf{F}} H_f^1(\mathbf{Q}, \rho(1-k)) = 1.$$

For A an object of $\text{LCN}(E)$ we will call a matrix $M \in \text{GL}_4(A)$ upper-triangular if

$$M = \begin{bmatrix} * & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \\ 0 & 0 & 0 & * \end{bmatrix}.$$

We will say that a homomorphism $\sigma' : G_\Sigma \rightarrow \text{GL}_4(A)$ is upper-triangular if for all $g \in G_\Sigma$, the matrix $\sigma'(g) \in \text{GL}_4(A)$ is upper-triangular. Finally, we say that a deformation $\sigma : G_\Sigma \rightarrow \text{GL}_4(A)$ of $\bar{\sigma}$ is upper-triangular if there exists a member σ' of the strict equivalence class of σ which is upper-triangular.

Lemma 7.1. *Every short crystalline minimal deformation σ of $\bar{\sigma}$ to $\text{GL}_4(A)$ such that $\text{tr } \sigma = T_1 + T_2 + T_3$ with T_1, T_2, T_3 pseudocharacters such that $T_1 \equiv \chi^{k-2}$, $T_2 \equiv \text{tr } \rho$, $T_3 \equiv \chi^{k-1} \pmod{\mathfrak{m}_A}$ is upper-triangular.*

Proof. This is Theorem 1.1 in [Bro08]. Note that the Artinian assumption in [Bro08] is unnecessary. See also remarks to Theorem 1 in [Urb99]. \square

Lemma 7.2. *Consider an upper-triangular deformation of $\bar{\sigma}$ to $\text{GL}_4(\mathbf{F}[X]/X^2)$. Then every upper-triangular member σ' of its strict equivalence class has the form*

$$\sigma' = \begin{bmatrix} \chi^{k-2} & * & * & \\ & \rho & * & \\ & & \chi^{k-1} & \end{bmatrix}.$$

Proof. We need to show that the only deformations of χ^{k-2} , ρ , χ^{k-1} are the trivial ones. Suppose that $\chi^{k-2} + \alpha X$ is a deformation of χ^{k-2} to $(\mathbf{F}[X]/X^2)^\times$. Then since χ^{2-k} is also crystalline, we get that $1 + \chi^{2-k}\alpha X$ is crystalline. Then $\alpha' := \chi^{2-k}\alpha$ is a homomorphism from G_Σ to the additive group \mathbf{F} . Let $\ell \mid N$ be a prime. Note that the total ring of fractions of $\mathbf{F}[X]/X^2$ is $\mathbf{F}[X]/X^2$. If $x \in G_\Sigma$ is such that $\sigma'(I_\ell)$ is generated by $\sigma'(x)$ then by minimality we know that $\sigma'(x) - I_4$ is conjugate (over

$\mathbf{F}[X]/X^2$) to $\begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & f & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ for $f \in \mathbf{F}^\times$. Using the upper-triangular form

of σ' this implies that α must be unramified at ℓ . Hence the character $1 + \alpha'X : G_\Sigma \rightarrow (\mathbf{F}[X]/X^2)^\times$ can only be ramified at p . However, crystallinity forces the character to be trivial (by a simple modification of the proof of Lemma 9.6 in [BK13]), so $\alpha = 0$. Similarly one proves the only deformation of χ^{k-1} is the trivial one. Finally, the claim for ρ follows from the assumption that R_ρ is a dvr. \square

Proposition 7.3. *There do not exist any non-trivial upper-triangular deformations of $\bar{\sigma}$ to $\mathbf{F}[X]/X^2$.*

Proof. First note that $\begin{bmatrix} \chi^{k-2} & a \\ & \rho \end{bmatrix}$ is a subrepresentation of $\bar{\sigma}$ and that $\begin{bmatrix} \rho & c \\ & \chi^{k-1} \end{bmatrix}$ is a quotient of $\bar{\sigma}$ hence they are both short crystalline since $\bar{\sigma}$ is. Thus (by

Corollary 3.3) a gives rise to a (non-zero) element of $H_f^1(\mathbf{Q}, \text{Hom}(\rho, \chi^{k-2})) = H_f^1(\mathbf{Q}, \rho(1-k))$ and c gives rise to a (non-zero) element of $H_f^1(\mathbf{Q}, \text{Hom}(\chi^{k-1}, \rho)) = H_f^1(\mathbf{Q}, \rho(1-k))$ (cf. Remark 3.1). Since this last Selmer group is assumed to be one-dimensional, a and c give rise to linearly dependent cohomology classes.

By Lemma 7.2 any upper-triangular deformation ϱ of $\bar{\sigma}$ to $\text{GL}_4(\mathbf{F}[X]/X^2)$ has the form

$$\varrho = \begin{bmatrix} \chi^{k-2} & a + Xa' & b + Xb' \\ & \rho & c + Xc' \\ & & \chi^{k-1} \end{bmatrix}$$

for a', b', c' valued in \mathbf{F} .

Let us now show that all the off-block-diagonal entries are unramified outside p . Suppose $\ell \mid N$. By Corollary 3.3 we see that a, b, c are all unramified at ℓ . Write $a' = [a_1 \ a_2]$, $c' = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$ with $a_i, c_i : G_\Sigma \rightarrow \mathbf{F}$. Let $x \in G_\Sigma$ be such that $\varrho(x)$ generates $\varrho(I_\ell)$. After conjugating ρ (if necessary), we may assume that $\rho(x) = \begin{bmatrix} 1 & 1 \\ & 1 \end{bmatrix}$. As the (2,3)-entry of $\varrho(x) - I_4$ lies in \mathbf{F}^\times minimality of ϱ forces the first row of $\varrho(x) - I_4$ to be a scalar multiple of the second row and the fourth column of $\varrho(x) - I_4$ to be a scalar multiple of the third column. This gives $a_1(x) = c_2(x) = 0$. Set $E_0 := \begin{bmatrix} A & \\ & D \end{bmatrix}$ with $A = \begin{bmatrix} 1 & -a_2(x)X \\ & 1 \end{bmatrix}$ and $D = \begin{bmatrix} 1 & c_1(x)X \\ & 1 \end{bmatrix}$. One has $E_0(\varrho(x) - I_4)E_0^{-1} = \begin{bmatrix} 0 & 0 & 0 & b'(x)X \\ 0 & 0 & * & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$. Note that $E_0 \equiv I_4 \pmod{\varpi}$, so the conjugate $E_0\varrho E_0^{-1}$ is strictly equivalent to ϱ . Hence we may assume that $a_2(x) = c_1(x) = 0$. Since again the first row must be a scalar multiple of the second row we also get $b'(x) = 0$.

Note that $\begin{bmatrix} \chi^{k-2} & a + Xa' \\ & \rho \end{bmatrix}$ is a subrepresentation of ϱ (with $a + Xa'$ unramified away from p). Then using that $\dim_{\mathbf{F}} H_f^1(\mathbf{Q}, \rho(1-k)) = 1$ we obtain by Proposition 7.2 of [BK13] (where the assumption 6.1(ii) is satisfied for ρ and for χ^{k-2} it can be replaced by the lack of a non-trivial deformation of χ^{k-2} to $(\mathbf{F}[X]/X^2)^\times$ - cf. Proof of Lemma 7.2 above) that $\begin{bmatrix} \chi^{k-2} & a + Xa' \\ & \rho \end{bmatrix}$ can be conjugated (over $\mathbf{F}[X]/X^2$) to the representation $\begin{bmatrix} \chi^{k-2} & a \\ & \rho \end{bmatrix}$, i.e., there exists $A \in I_3 + XM_3(\mathbf{F}[X]/X^2)$ such that $A \begin{bmatrix} \chi^{k-2} & a + Xa' \\ & \rho \end{bmatrix} A^{-1} = \begin{bmatrix} \chi^{k-2} & a \\ & \rho \end{bmatrix}$. Then we have

$$\begin{bmatrix} A & 0 \\ 0 & 1 \end{bmatrix} \varrho \begin{bmatrix} A & 0 \\ 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} \chi^{k-2} & a & b'' + Xb''' \\ & \rho & c'' + Xc''' \\ & & \chi^{k-1} \end{bmatrix}.$$

Since the image of A in $M_3(\mathbf{F})$ is the identity, we see that $b'' = b$ and $c'' = c$, so in particular this is still a deformation of $\bar{\sigma}$. Hence we conclude that ϱ is

equivalent to $\begin{bmatrix} \chi^{k-2} & a & b + Xb' \\ & \rho & c + Xc' \\ & & \chi^{k-1} \end{bmatrix}$ which we will still denote by ϱ .

Let V be the representation space of ϱ (i.e., the free $\mathbf{F}[X]/X^2$ -module of rank 4 on which G_Σ acts via ϱ). Let W be the G_Σ -invariant free $\mathbf{F}[X]/X^2$ -submodule of V spanned by the vector ${}^t[1, 0, 0, 0]$. Then G_Σ acts on the quotient V/W via

$$\varrho' := \begin{bmatrix} \rho & c + Xc' \\ & \chi^{k-1} \end{bmatrix}.$$

Hence we can again conclude by Proposition 7.2 in [BK13] that there exists $A \in I_3 + XM_3(\mathbf{F}[X]/X^2)$ such that $A\varrho'A^{-1} = \begin{bmatrix} \rho & c \\ & \chi^{k-1} \end{bmatrix}$. Then we have

$$\begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} \varrho \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix}^{-1} = \begin{bmatrix} \chi^{k-2} & a + Xa'' & b + Xb'' \\ & \rho & c \\ & & \chi^{k-1} \end{bmatrix}.$$

Denote the right-hand side of the above equation by ϱ' . Now, let us re-write ϱ' in the \mathbf{F} -basis of $(\mathbf{F}[X]/X^2)^4$ given by

$$\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} X \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ X \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ X \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ X \end{bmatrix} \right\}.$$

We get that ϱ' in the basis \mathcal{B} has the form

$$\begin{bmatrix} \chi^{k-2} & a & b \\ & \rho & c \\ & & \chi^{k-1} \\ & a'' & b'' & \chi^{k-2} & a & b \\ & & & & \rho & c \\ & & & & & & \chi^{k-1} \end{bmatrix}.$$

Taking a sequence of submodules and quotients we get

$$\begin{bmatrix} \chi^{k-2} & a & b \\ & \rho & c \\ & & \chi^{k-1} \\ & a'' & b'' & \chi^{k-2} \end{bmatrix} \rightarrow \begin{bmatrix} \rho & c \\ a'' & b'' & \chi^{k-2} \end{bmatrix},$$

which if we conjugate it by $\begin{bmatrix} I_2 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$ will give us

$$(7.2) \quad \begin{bmatrix} \rho & & c \\ a'' & \chi^{k-2} & b'' \\ & & \chi^{k-1} \end{bmatrix},$$

from which we can extract a short crystalline 2-dimensional subrepresentation of the form

$$\begin{bmatrix} \rho & \\ a'' & \chi^{k-2} \end{bmatrix},$$

and again (by crystallinity of the above representation) we see that a'' (which is unramified away from p by an argument as before) gives rise to an element of $H_f^1(\mathbf{Q}, \text{Hom}(\rho, \chi^{k-2})) = H_f^1(\mathbf{Q}, \rho(1-k))$. So, by one-dimensionality of the latter there exists a constant $\alpha \in \mathbf{F}$ such that $a'' = \alpha a$.

Hence we get that

$$(7.3) \quad \varrho = \begin{bmatrix} \chi^{k-2} & a & b + Xb' \\ & \rho & c + Xc' \\ & & \chi^{k-1} \end{bmatrix} \cong \begin{bmatrix} \chi^{k-2} & a + X\alpha a & b + Xb'' \\ & \rho & c \\ & & \chi^{k-1} \end{bmatrix} = \varrho',$$

and the isomorphism is by conjugation by an element of $I_4 + XM_4(\mathbf{F})$. Furthermore we note that

$$M_0 \varrho' M_0^{-1} = \begin{bmatrix} \chi^{k-2} & a & b + Xb'' \\ & \rho & c \\ & & \chi^{k-1} \end{bmatrix},$$

where

$$M_0 = \begin{bmatrix} 1 & & \\ & (1 + \alpha X)I_2 & \\ & & 1 + \alpha X \end{bmatrix} \in 1 + XM_4(\mathbf{F}).$$

Hence we conclude that we can take $a'' = 0$.

Then we see that $\begin{bmatrix} \chi^{k-2} & b'' \\ & \chi^{k-1} \end{bmatrix}$ is a quotient of the representation in (7.2). Thus, b'' gives rise to an element of $H_{\Sigma}^1(\mathbf{Q}, \mathbf{F}(-1))$. By the argument from the beginning of the proof we see that b'' is unramified at all primes $\ell \mid N$, so it in fact gives rise to an element of $H_f^1(\mathbf{Q}, \mathbf{F}(-1))$ which is zero by Proposition 2.9. \square

Let $I_{R'} \subset R'$ denote the total ideal of reducibility corresponding to the universal deformation $(\sigma')^{\text{univ}}$ and $I_R \subset R$ the total ideal of reducibility corresponding to the universal deformation σ^{univ} .

Corollary 7.4. *The structure maps $\mathcal{O} \rightarrow R'/I_{R'}$, $\mathcal{O} \rightarrow R/I_R$ and $\mathcal{O} \rightarrow R^{\text{red}}/I^{\text{tot}}$ are all surjective.*

Proof. By Lemma 7.11 of [BK13] it is enough to show that the structure map $\mathcal{O} \rightarrow R'/I_{R'}$ is surjective. We need the following lemma.

Lemma 7.5. *Let $I \subset R'$ be an ideal such that $R'/I \in \text{LCN}(E)$. Then $I \supset I'_R$ if and only if $(\sigma')^{\text{univ}} \bmod I$ is an upper-triangular deformation of $\bar{\sigma}$ to $\text{GL}_4(R'/I)$.*

Proof. The proof of Corollary 7.8 in [BK13] carries over to three Jordan-Hölder factors with the application of Theorem 7.7 in [BK13] replaced by an application of Lemma 7.1 above. \square

The rest of the proof of Corollary 7.4 is the same as the proof of Proposition 7.10 in [BK13] with the application of Corollary 7.8 in [BK13] now replaced with an application of Lemma 7.5. \square

Theorem 7.6. *Let $\tilde{\rho}$ be a representative of the unique deformation of ρ to characteristic zero. Suppose that $\#H_f^1(\mathbf{Q}, \tilde{\rho}(1-k) \otimes E/\mathcal{O}) \leq \#\mathcal{O}/L\mathcal{O}$ for some $L \in \mathcal{O}$. Then $\#R^{\text{red}}/I^{\text{tot}} \leq \#\mathcal{O}/L\mathcal{O}$.*

Proof. Again by Lemma 7.11 of [BK13] it is enough to show that $\#R'/I_{R'} \leq \#\mathcal{O}/L\mathcal{O}$. First let us note that it follows from the proof of Lemma 7.2 that there do not exist any non-trivial short crystalline deformations of χ^{k-2} and of χ^{k-1} to $(\mathbf{F}[X]/X^2)^\times$. Let R_{k-2} (resp. R_{k-1}) denote the universal short crystalline deformation ring of χ^{k-2} (resp. χ^{k-1}). Then we can adapt the proof of Proposition 7.10 in [BK13] with S there replaced by R_{k-2} to show that $R_{k-2}/\varpi R_{k-2} = \mathbf{F}$ and hence by Nakayama's lemma the structure map $\mathcal{O} \rightarrow R_{k-2}$ is onto. However, since ϵ^{k-2} is a deformation of χ^{k-2} to \mathcal{O}^\times , we must in fact have $R_{k-2} = \mathcal{O}$ and so ϵ^{k-2} is a unique such deformation. The same conclusion holds for R_{k-1} . Assume that $\#R'/I_{R'} = \#\mathcal{O}/\varpi^s\mathcal{O} > \#\mathcal{O}/L\mathcal{O}$ (we allow $s = \infty$ here). Then by Lemma 7.5 there exists an upper-triangular deformation $\sigma : G_\Sigma \rightarrow \text{GL}_4(\mathcal{O}/\varpi^s\mathcal{O})$ of $\bar{\sigma}$. By the above argument and using the fact that $\tilde{\rho}$ is the unique deformation of ρ to $\text{GL}_2(\mathcal{O})$ the deformation σ must have the form

$$\begin{bmatrix} \epsilon^{k-2} & *_1 & *_2 \\ & \tilde{\rho} & *_3 \\ & & \epsilon^{k-1} \end{bmatrix},$$

where the diagonal pieces are understood to be taken mod ϖ^s .

Arguing as in the proof of Proposition 7.3 we see that minimality guarantees that the classes of $*_1$ and $*_3$ are unramified away from p and thus give rise to classes in $H_f^1(\mathbf{Q}, \tilde{\rho}(1-k) \otimes E/\mathcal{O}[\varpi^s]) = H_f^1(\mathbf{Q}, \tilde{\rho}(1-k) \otimes E/\mathcal{O})[\varpi^s]$ (cf. Proposition 2.4(1)). Since the reduction $\bar{\sigma}$ of σ has the property that its entry corresponding to $*_3$ above gives rise to a non-trivial element in $H_f^1(\mathbf{Q}, \rho(1-k))$, we conclude that the image of $*_3$ is not contained in $\varpi\mathcal{O}/\varpi^s\mathcal{O}$, hence gives rise to an element of $H_f^1(\mathbf{Q}, \tilde{\rho}(1-k) \otimes E/\mathcal{O})$ of order $\#\mathcal{O}/\varpi^s$, which is a contradiction. \square

Corollary 7.7. *The rings R' , R and R^{red} are all topologically generated as an \mathcal{O} -algebra by the set $\{\text{tr } \sigma(\text{Frob}_l) \mid l \nmid Np\}$, where σ stands for $(\sigma')^{\text{univ}}$, σ^{univ} and σ^{red} respectively.*

Proof. It is enough to prove this for R' as the other rings are quotients of it. For this one can use the same proof as the one of Proposition 7.13 in [BK13] replacing again the application of Corollary 7.8 with Lemma 7.5. \square

The upshot of this section is the following theorem identifying conditions when the universal deformation ring is a discrete valuation ring. To make the statement self-contained we will include all the assumptions made so far.

Theorem 7.8. *Let $\rho_f : G_\Sigma \rightarrow \text{GL}_2(E)$ be the Galois representation attached to a newform $f \in S_{2k-2}(N)$ for N a square-free integer with $p \nmid N$ and*

$\Sigma = \{\ell \mid N\} \cup \{p\}$ such that $p \nmid 1 + w_{f,\ell}$ for all primes $\ell \mid N$. Suppose also that the residual representation $\rho := \bar{\rho}_f$ is absolutely irreducible and ramified at every prime $\ell \mid N$. Assume that $\dim_{\mathbf{F}} H_f^1(\mathbf{Q}, \rho(2-k)) = 1$ and that $\#H_f^1(\mathbf{Q}, \rho_f(1-k) \otimes E/\mathcal{O}) \leq \#\mathbf{F}$. Furthermore assume that R_ρ is a dvr. Let $\bar{\sigma} : G_\Sigma \rightarrow \mathrm{GL}_4(\mathbf{F})$ be a continuous representation short crystalline at p and semi-abelian of the form

$$\bar{\sigma} = \begin{bmatrix} \chi^{k-2} & a & b \\ & \rho & c \\ & & \chi^{k-1} \end{bmatrix}$$

with a, c giving rise to non-zero elements of $H_f^1(\mathbf{Q}, \rho(1-k))$. Suppose $\bar{\sigma}$ admits a deformation σ to $\mathrm{GL}_4(\mathcal{O})$ which is τ -self-dual.

Then σ gives rise to an \mathcal{O} -algebra isomorphism $R_\sigma^{\mathrm{red}} = R^{\mathrm{red}} \xrightarrow{\sigma} \mathcal{O}$.

Proof. The \mathcal{O} -algebra map $R \rightarrow \mathcal{O}$ induced by σ factors through R^{red} since \mathcal{O} is torsion-free. Theorem 6.6 combined with Theorem 7.6 imply that the maximal ideal of R^{red} is principal. Because of the surjection to \mathcal{O} its generator is not nilpotent. As in [Cal06] Lemma 3.4 we can therefore deduce that R^{red} is a discrete valuation ring and the surjection an isomorphism $R^{\mathrm{red}} \cong \mathcal{O}$. \square

8. APPLICATION TO THE PARAMODULAR CONJECTURE

The following conjecture is due to Brumer and Kramer (cf. Conjecture 1.1.1 in [BPP⁺ar]) and is often referred to as the Paramodular Conjecture.

Conjecture 8.1 (Brumer - Kramer). *For every isogeny class of abelian surfaces A/\mathbf{Q} of conductor N with $\mathrm{End}_{\mathbf{Q}} A = \mathbf{Z}$ there exists a weight 2 Siegel modular form F , which is not in the space spanned by the Saito-Kurokawa lifts, has level $K(N)$ and rational eigenvalues, where $K(N)$ is the paramodular group of level N defined by $K(N) = \gamma M_4(\mathbf{Z}) \gamma^{-1} \cap \mathrm{Sp}_4(\mathbf{Q})$ with $\gamma = \mathrm{diag}[1, 1, 1, N]$. The L -series of A and F should agree and the p -adic representation of $T_p(A) \otimes \mathbf{Q}_p$ should be isomorphic to the one associated to F for any p prime to N where $T_p(A)$ is the p -adic Tate module.*

In this section we will show how our results can be used to verify this conjecture in some cases when A has rational p -torsion. More precisely, our method will only allow for the verification of the second claim, i.e., the existence of the isomorphism of the Tate module with the representation associated to F .

Let $p > 2$ be a prime and let A be an abelian surface of square-free conductor N as in Conjecture 8.1. Suppose that $p \nmid N$ and that A has a polarization of degree prime to p and a \mathbf{Q} -rational point of order p . Then the p -adic Tate module gives rise to a Galois representation

$$\sigma_A := V_p(A) : G_\Sigma \rightarrow \mathrm{GL}_4(\mathbf{Q}_p),$$

where $\Sigma = \{p\} \cup \{\ell \mid N\}$, short crystalline at p (by [ST68] and [Fon82]), which is absolutely irreducible since it is semisimple and $\mathrm{End}_{\mathbf{Q}_p[G_\Sigma]}(V_p(A)) =$

$\text{End}_{\mathbf{Q}}(A) \otimes \mathbf{Q}_p = \mathbf{Q}_p$ by [Fal83]. Furthermore, under our assumptions the semisimple residual representation $\bar{\sigma}_A^{\text{ss}} : G_{\Sigma} \rightarrow \text{GL}_4(\mathbf{F}_p)$ has the form

$$\bar{\sigma}_A^{\text{ss}} = 1 \oplus \rho^{\text{ss}} \oplus \chi,$$

where, as before, χ is the mod p cyclotomic character and ρ is a two-dimensional representation. The representation σ_A has determinant ϵ^2 and hence $\det \rho = \chi$, so in particular, ρ is odd.

We will assume that ρ is absolutely irreducible, ramified at all $\ell \mid N$. It follows from Serre's conjecture [KW09] (see Proposition 4 of [Ser87] for the determination of the Serre weight) that $\rho \cong \bar{\rho}_f$ where $\rho_f : G_{\Sigma} \rightarrow \text{GL}_2(\mathbf{Q}_p)$ is the Galois representation attached to a newform $f \in S_2(N)$ and $\bar{\rho}_f$ denotes its mod p reduction.

We now assume that A has semi-abelian reduction at all $\ell \mid N$, i.e. the reduction is an extension of an elliptic curve by a torus and that p does not divide the Tamagawa number at ℓ , i.e. the number of connected components of the special fiber of a Neron model of A at ℓ . Under these assumptions [LD98] Proposition 1.3 proves that $\bar{\sigma}_A|_{I_{\ell}} \cong \exp(t_{\ell}N_1)$. The proof generalizes to $A[p^n]$ as [ST68] Lemma 1 and 2 hold in this generality, so we deduce that $\sigma_A|_{I_{\ell}} \cong \exp(t_{\ell}N_1)$.

For a positive integer N and an integer $k \geq 2$ we will denote by $\mathcal{S}_k(N)^{\text{para}}$ the space of Siegel modular forms of genus 2, (parallel) weight k and paramodular level N . For a prime $\ell \nmid N$, write $T_{\ell,2}$ for the Hecke operator given by the double coset $K(N) \text{diag}(1, 1, \ell, \ell)K(N)$, $T_{\ell,1}$ for the Hecke operator given by the double coset $K(N) \text{diag}(1, \ell, \ell^2, \ell)K(N)$, and $T_{\ell,0}$ for the Hecke operator given by the double coset $\ell K(N)$ (see e.g. p. 18 of [Pil17]). We will say that $F \in \mathcal{S}_k(N)^{\text{para}}$ is an eigenform if F is an eigenform for $T_{\ell,i}$ for all $\ell \nmid N$ and $i = 0, 1, 2$. If F is an eigenform we will write $\lambda_{\ell,i}(F)$ for the eigenvalue of $T_{\ell,i}$ corresponding to F .

Theorem 8.2 (Taylor, Weissauer, Laumon). *For $k \geq 2$ let $F \in S_k(N)^{\text{para}}$ be an eigenform. There exists a finite extension E of \mathbf{Q}_p and a continuous semisimple representation*

$$\sigma_F : G_{\Sigma} \rightarrow \text{GL}_4(E)$$

with $\sigma_F^{\vee} \cong \sigma_F \otimes \epsilon^{3-2k}$ such that for all primes $\ell \nmid Np$ one has $\det(I_4 - X\sigma_F(\text{Frob}_{\ell}))^{-1} = Q_{\ell}(X)$ for the Hecke polynomial

$$Q_{\ell}(X) = 1 - \lambda_{\ell,2}(F)X + \ell(\lambda_{\ell,1}(F) + (1 + \ell^2)\lambda_{\ell,0}(F))X^2 - \ell^3\lambda_{\ell,2}(F)\lambda_{\ell,0}(F)X^3 + \ell^6\lambda_{\ell,0}(F)^2X^4,$$

where $\lambda_{\ell,0}(F) = \ell^{2k-6}$.

Remark 8.3. This result is well-known, but for the benefit of the reader we explain the relationship to results in the literature. The eigenform F generates an automorphic representation π that decomposes into a direct sum $\pi_1 \oplus \dots \oplus \pi_n$ of irreducible cuspidal automorphic representations of GSp_4 whose local representations agree away from N . Laumon [Lau05] and Weissauer [Wei05] constructed the Galois representations for irreducible automorphic representations π with π_{∞} holomorphic discrete series (and hence for eigenforms F as above for weight $k > 2$ as all π_i give rise to the same semi-simple Galois representation). The result for $k = 2$ can be

deduced from this using Taylor's argument in Example 1 of section 1.3 in [Tay91]. [Jor10] and [Mok14] Proposition 4.14 provide an alternative proof (for [Mok14] under the restriction that F be of type (G) in the sense of [Sch17] section 2.1, i.e. such that all π_i have a cuspidal transfer to GL_4 , but see Remark [Mok14] 4.15).

Theorem 8.4 ([FC90], [Jor12] Theorem 3.1). *Let F be as in Theorem 8.2, $k \geq 2$ and $p \nmid N$. If $k = 2$ assume that F has pairwise distinct roots for the Hecke polynomial $Q_p(F)$. Then σ_F is crystalline at p and short crystalline if $p > 2k - 2$.*

Proposition 8.5. *For $k \geq 2$ and N squarefree let $F \in S_k(N)^{\mathrm{para}}$ be an eigenform of type (G) in the sense of [Sch17] section 2.1. If $k = 2$ we further assume that σ_F is absolutely irreducible and that $\bar{\sigma}_F^{\mathrm{ss}}$ is ramified at ℓ for all $\ell \mid N$. Then we have*

$$\sigma_F|_{I_\ell} \cong \exp(t_p N_1) \text{ for all } \ell \mid N.$$

Proof. For $k > 2$ this is proved by [Mok14] Theorem 3.5 (local-global compatibility up to Frobenius semisimplification) and [Sor10] Corollary 1 (monodromy rank 1). (We know by [Sch17] Proposition 1.2.1 that type (G) paramodular eigenforms give rise to generic local representations, so that Sorenson's result applies.)

As for $k = 2$ [Mok14] Theorem 4.14 only proves local-global compatibility up to semisimplification we argue as follows: Theorem 1.1 of [MT15] allows us to p -adically approximate the Hecke eigenvalues of a weight 2 form F by those of cohomological Hecke eigenforms F_n . This means that the corresponding Galois representations σ_{F_n} are trace convergent to σ_F in the sense of [BCKL05], i.e. $\mathrm{tr}(\sigma_{F_n})(g)$ converges to $\mathrm{tr}(\sigma_F)(g)$ in \mathbf{C}_p for all $g \in G_{\mathbf{Q}}$. As we assume that σ_F is absolutely irreducible we can invoke Theorem 1.2 of [BCKL05] to deduce that σ_{F_n} are physically convergent to σ_F , i.e. such that there exist \mathbf{C}_p -bases for σ_{F_n} and ρ such that the matrix entries of σ_{F_n} converge to the corresponding entries of σ_F . This result also tells us that the σ_{F_n} are absolutely irreducible for $n \gg 0$ so we know that the F_n are eventually of type (G). As these are of weight $k > 2$ we know that $\sigma_{F_n}|_{I_\ell} \cong \exp(t_p N_1)$ and the rank of the monodromy remains 1 in the limit by our assumption that $\bar{\sigma}_F^{\mathrm{ss}}$ is ramified. \square

Theorem 8.6. *Let A and $\rho = \bar{\rho}_f$ be as above and assume that $p \nmid 1 + w_{f,\ell}$ for all $\ell \mid N$, $R_\rho = \mathcal{O}$, $\#H_f^1(\mathbf{Q}, \bar{\rho}_f) = \#\mathbf{F}$ and $\#H_f^1(\mathbf{Q}, \rho_f(-1) \otimes E/\mathcal{O}) \leq \#\mathbf{F}$.*

Furthermore suppose that there exists $F \in \mathcal{S}_2(N)^{\mathrm{para}}$ such that $\lambda_{\ell,2}(F) \equiv 1 + \ell + a_\ell(f) \pmod{p}$ for all primes $\ell \nmid Np$ and such that its Hecke polynomial at some prime $\ell \nmid Np$ vanishes neither at 1 nor at $1/\ell$. We also assume that F has pairwise distinct roots for the Hecke polynomial $Q_p(F)$. Then $\sigma_A \cong \sigma_F$. In particular, A is paramodular of level N .

Proof. The congruence, Tchebotarev's density Theorem and the Brauer-Nesbitt Theorem imply that the semisimplification $\bar{\sigma}_F^{\mathrm{ss}}$ of the mod p reduction of σ_F has the form $\bar{\sigma}_F^{\mathrm{ss}} = 1 \oplus \bar{\rho}_f \oplus \chi = \bar{\sigma}_A^{\mathrm{ss}}$.

We will show below that σ_F is absolutely irreducible. Using Corollary 4.4 (and the fact that σ_F is short crystalline at p - cf. Theorem 8.4) we can then

choose G_Σ -stable lattices in σ_A and in the representation space of σ_F so that with respect to these lattices $\bar{\sigma}_A$ and $\bar{\sigma}_F$ have the form as in Corollary 4.4. Then by Proposition 5.1 we get that in fact $\bar{\sigma}_A \cong \bar{\sigma}_F$. By adjusting the basis of σ_F if necessary we can assume that $\bar{\sigma}_A = \bar{\sigma}_F$, hence we obtain that σ_F is a deformation of $\bar{\sigma}_A$. Let R be the quotient of the universal deformation ring of $\bar{\sigma}_A$ as in Proposition 3.6. Then by Theorem 7.8 we see that σ_A and σ_F both give rise to an isomorphism $R^{\text{red}} \cong \mathcal{O}$ of \mathcal{O} -algebras. Hence we get a commutative diagram

$$\begin{array}{ccc}
 \mathcal{O} & & \\
 \searrow^{\text{id}} & & \\
 & R^{\text{red}} & \xrightarrow[\sigma_A]{\sim} \mathcal{O} \\
 \searrow^{\text{id}} & \downarrow \sigma_F & \\
 & \mathcal{O} &
 \end{array}$$

from which we see that $\sigma_A = \sigma_F$ as maps from R^{red} to \mathcal{O} , which implies that the representations σ_F and σ_A are isomorphic since they are both composites of the universal deformation with the map $R^{\text{red}} \rightarrow \mathcal{O}$.

Let us now show that σ_F is absolutely irreducible. Indeed, note that σ_F cannot be the sum of 4 characters because its reduction has an absolutely irreducible two-dimensional component $\bar{\rho}_f$. Suppose that σ_F^{ss} splits as a direct sum of two irreducible 2-dimensional representations $\sigma_1 \oplus \sigma_2$. Note that $\sigma_1^\vee \not\cong \sigma_2 \otimes \epsilon^{-1}$ since without loss of generality $\bar{\sigma}_1 = \rho \neq \bar{\sigma}_2^{\text{ss}} = 1 \oplus \chi$. We therefore have $\sigma_1^\vee \cong \sigma_1 \otimes \epsilon^{-1}$, which implies $\det(\sigma_1) = \det(\sigma_2) = \epsilon$. Indeed we must have $\det(\sigma_1) = \epsilon\phi$ for some quadratic character ϕ reducing to the identity mod p since $\det(\bar{\sigma}_1) = \chi$. Since $p > 2$ the character ϕ has to be trivial.

We can therefore argue as in case (v) on p. 46 of [SU06]. Note that σ_1 has to be odd since ρ is, so σ_2 must be odd as well. Since the automorphic representation π corresponding to F can be transferred to an isobaric automorphic representation on GL_4 the assumptions in Theorem C of [Ram13] are satisfied, which tells us that $L^S(\sigma_1^\vee \otimes \sigma_2, 1) \neq 0$ for some finite set of places S . This means that the standard L -function $L^S(s, \pi, \text{std}) = \zeta^S(s) L^S(\sigma_1^\vee \otimes \sigma_2, s)$ has a pole at $s = 1$, which by [KRS92] implies that π is endoscopic, i.e. corresponds to type (Y) in [Sch17]. Since F is paramodular this contradicts [Sch17] Lemma 2.2.1. This excludes the case $\sigma_A = \sigma_1 \oplus \sigma_2$.

It remains to show that σ_A cannot split as $\sigma_1 \oplus \chi_1$ with σ_1 a 3-dimensional representation and χ_1 a character. If it did, then the spin L -function of A would have a linear factor. Since the only short crystalline, minimal deformations of 1 (resp. χ) are 1 (resp. ϵ) - cf. the beginning of the proof of Lemma 7.2 - we must have that χ_1 is either 1 or ϵ . Thus the linear factor of the local spin L -function must be $1 - \ell^{-s}$ (if $\chi_1 = 1$) or $1 - \epsilon(\ell)\ell^{-s} = 1 - \ell\ell^{-s}$. But the local spin L -factor is the local Hecke polynomial at ℓ with X replaced by ℓ^{-s} , so we're done by our assumption. \square

9. EXAMPLES

In this section we work out in detail how our result proves paramodularity of an abelian surface of conductor 731. We also discuss other cases where the result may be applicable without going into details.

9.1. Conductor $N = 731$. We will show that for an abelian surface of this conductor (using a congruence result proved in the Appendix) the conditions for Theorem 8.6 are satisfied. This establishes a new case of the Paramodular Conjecture.

For $N = 731 = 17 \cdot 43$ there exists an abelian surface A of conductor N , which arises as the Jacobian of the hyperelliptic curve with equation

$$y^2 + (x^3 + x^2)y = x^5 + 2x^4 - x - 3.$$

According to LMFDB [LMF17b] it has semi-abelian reduction at 17 and 43 and has a rational point of order $p = 5$. We first note that $5 \nmid 1 \pm \ell$ for $\ell = 17, 43$.

Since A has rational 5-torsion and is principally polarised we know that $\bar{\sigma}_{A,p}^{\text{ss}} = 1 \oplus \rho \oplus \chi$. Furthermore A has semi-abelian reduction and has Tamagawa numbers at 17 and 43 not divisible by 5, which tells us that if ρ were reducible over any finite extension of \mathbf{Q}_p its semisimplification would be unramified away from 5, i.e. of the form $\chi^i \oplus \chi^{1-i} \pmod{5}$ for $i = 1$ or 2 . We check on specific Frobenius elements that the trace of ρ is not equal to $\chi^i + \chi^{1-i}$ for $i = 1, 2$ so that ρ has to be absolutely irreducible. Serre's conjecture implies that ρ is modular by a cuspidal weight 2 eigenform of level 17, 43 or 731 (see section 8). Using MAGMA [BCP97] we check (by comparing the linear term of the Euler factor of A at ℓ with $1 + \ell + a_\ell(f)$) that the only newform giving rise to ρ is the modular form corresponding to the elliptic curve E (Cremona label 731a1) of conductor 731 given by

$$y^2 + xy + y = x^3 - 539x + 4765,$$

which has rank 1 and trivial rational torsion. This also shows that 5 is not a congruence prime for the corresponding weight 2 modular form, i.e. that $R_\rho = \mathbf{Z}_p$. By consulting LMFDB [LMF17a] we know that $\bar{\rho}_{E,p}$ is ramified at 17 and 43 (since E has non-split reduction at both primes). We also note that E has good ordinary reduction at 5.

We can check the Selmer group conditions as follows: We have $H_f^1(\mathbf{Q}, \bar{\rho}_{E,p}) = H_f^1(\mathbf{Q}, \rho_{E,p} \otimes \mathbf{Q}_p/\mathbf{Z}_p)[p] = \text{Sel}_p(E)[p]$ where the first equality follows from Proposition 2.4 and for the second one see e.g. [Rub00] Proposition 1.6.8. Here $\text{Sel}_p(E)$ is defined by

$$(9.1) \quad 0 \rightarrow E(\mathbf{Q}) \otimes \mathbf{Q}_p/\mathbf{Z}_p \rightarrow \text{Sel}_p(E) \rightarrow \text{III}(E)[p^\infty] \rightarrow 0.$$

Using snake lemma on the diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & E(\mathbf{Q}) \otimes \mathbf{Q}_p/\mathbf{Z}_p & \longrightarrow & \text{Sel}_p(E) & \longrightarrow & \text{III}(E)[p^\infty] \longrightarrow 0 \\ & & \downarrow \cdot p & & \downarrow \cdot p & & \downarrow \cdot p \\ 0 & \longrightarrow & E(\mathbf{Q}) \otimes \mathbf{Q}_p/\mathbf{Z}_p & \longrightarrow & \text{Sel}_p(E) & \longrightarrow & \text{III}(E)[p^\infty] \longrightarrow 0 \end{array}$$

we get an exact sequence

$$0 \rightarrow E(\mathbf{Q}) \otimes \mathbf{Q}_p/\mathbf{Z}_p[p] \rightarrow \mathrm{Sel}_p(E)[p] \rightarrow \mathrm{III}(E)[p] \rightarrow 0.$$

Since $\bar{\rho}_{E,p}$ is absolutely irreducible, we conclude that E has no rational p -torsion and hence $(E(\mathbf{Q}) \otimes \mathbf{Q}_p/\mathbf{Z}_p)[p] = E(\mathbf{Q})/pE(\mathbf{Q})$. This implies that (9.1) gives rise to a short exact sequence

$$0 \rightarrow E(\mathbf{Q})/pE(\mathbf{Q}) \rightarrow H_f^1(\mathbf{Q}, \bar{\rho}_{E,p}) \rightarrow \mathrm{III}(E)[p] \rightarrow 0,$$

Since the rank of E is 1 we therefore deduce that $\#H_f^1(\mathbf{Q}, \bar{\rho}_{E,p}) = 5$ if $\mathrm{III}(E)[5] = 0$. According to LMFDB the analytic order of $\mathrm{III}(E)$ is 1. By Skinner et al. [JSW17] the p -part of the BSD formula is satisfied, so $\mathrm{III}(E)[5] = 0$ (alternatively this was checked by Grigorov, Jorza, Patrikis, Stein, and Tarnita, see Theorem 3.27 in [GJP⁺09]).

It now remains to bound the order of $\#H_f^1(\mathbf{Q}, \rho_{E,p}(-1) \otimes \mathbf{Q}_p/\mathbf{Z}_p)$ (which we know to be non-trivial) from above by 5. Setting $f \in S_2(\Gamma_0(731))$ in Proposition 2.10 to be the newform corresponding to E it is enough to calculate p -valuation of the p -adic L -series $L(E, \omega^{-1}) \in \mathbf{Z}_p[[T]]$ specialized at $T = p$. Using SAGE we confirm that this p -valuation is indeed 1.

As proved in the Appendix there is a unique non-lift paramodular eigenform F new at level 731 (non-vanishing modulo 5 and with Hecke eigenvalue $\lambda_{5,2}(F) = 0$) that is congruent modulo 5 to a Gritsenko lift (paramodular Saito-Kurokawa lift) of a modular form of level dividing 731. This congruence of Fourier expansions implies a congruence of Hecke eigenvalues modulo 5.

Using MAGMA we can rule out that F is congruent to all but the Saito-Kurokawa lift of the rational modular form f corresponding to E . Since $\lambda_{5,2}(F) = 0$ its Hecke polynomial at 5 is of the form $1 + cT^2 + 25T^4$ for some $c \in \mathbf{Z}$, which either has distinct roots or more than one repeated root. As $a_5(f) \equiv -1 \pmod{5}$ the Hecke polynomial of the Saito-Kurokawa lift of f (given by $(1 - \alpha T)(1 - \beta T)(1 - T)(1 - 5T)$ with $\alpha + \beta = a_5(f)$) is congruent to $1 - T^2$ modulo 5, so we deduce that the roots of the Hecke polynomial of F are distinct modulo 5.

Theorem A.1(4) in the appendix also shows that the Hecke polynomial at 2 is given by $4T^4 + 2T^3 + 2T^2 + T + 1 = (1 - T + 2T^2)(1 + 2T + 2T^2)$. Its complex roots do not include 1, 2 or $1/2$.

9.2. Other examples. Poor and Yuen have found candidate paramodular forms for some other examples of abelian surfaces. For the seven conductors in Table 5 of [PY15] one can check that only the abelian surface of conductor $N = 277$ involves a congruence (for the torsion prime $p = 3$, but not for $p = 5$) with the paramodular Saito-Kurokawa lift of a modular form corresponding to an elliptic curve. In this case the p -valuation of the p -adic L -series gives us a bound of 9 on $\#H_f^1(\mathbf{Q}, \rho_f(-1) \otimes E/\mathcal{O})$. It may still be possible that the Bloch-Kato Selmer group has order 3, but we were not able to confirm this (and this would fall outside of the range of applicability of our results as we exclude $p = 3$). However, this abelian surface has been proved to be paramodular by Brumer et al. using the Faltings-Serre method, but a proof has not yet appeared in print.

When the conjectured congruence is with the Saito-Kurokawa lift of a modular form with non-rational Fourier coefficients it is sometimes possible to check that $R_{\bar{\rho}_f}$ is a discrete valuation ring. This is the case for the examples of conductor $N = 349, 353$ and 389 on the list in [PY15]. These surfaces have rational p -torsion for $p = 13, 11$ and 5 , respectively. We have, however, not tried to check the Selmer group assumptions for these cases.

Another example we want to highlight is that of conductor $N = 997$, where there exists an abelian surface with 3-torsion. In this case the expected congruence is between a paramodular non-lift and a Saito-Kurokawa lift of a modular form corresponding to a rank 2 elliptic curve. As the root number of the modular form is $+1$ this Saito-Kurokawa lift has to be of congruence level, rather than paramodular as the other cases. It would be interesting to confirm the existence of a matching paramodular non-lift Siegel modular form in this case, but Theorem 8.6 does not apply because $p = 3$ and the elliptic curve rank > 1 should imply $\#H_f^1(\mathbf{Q}, \bar{\rho}_{E,p}) > 3$.

10. MODULARITY THEOREM FOR COHOMOLOGICAL WEIGHTS k

In this section we will prove a modularity theorem in cases $k > 2$ and $N = 1$. The sole reason for these two restrictions is that at present only in this case we have a result (which is due to Brown) providing us with enough congruences among Siegel modular forms [Bro11]. This result also allows us to replace the assumption that $\#H_f^1(\mathbf{Q}, \rho_f(1-k) \otimes E/\mathcal{O}) \leq \#\mathbf{F}$ with the weaker assumption that $\dim_{\mathbf{F}} H_f^1(\mathbf{Q}, \rho_f(1-k)) = 1$. In particular, the proof does not proceed via proving that R^{red} is a discrete valuation ring - in fact the latter property is not implied by our $R = T$ theorem. We again include all the assumptions to make the statement self-contained, however to make their use more transparent we will separate the assumptions that are necessary for Brown's congruence result (collecting them below as Assumption 10.1 - cf. Theorem 5.4 and Corollary 5.6 in [Bro11]) from the assumptions required on the deformation side (which will be spelled out in the statement of Theorem 10.2).

Let k be a positive integer and p a prime such that $p > 2k - 2$.

Assumption 10.1. Suppose $k > 9$ is even. Assume also that there exists $N \in \mathbf{Z}_{>1}$ and a fundamental discriminant $D < 0$ such that $\chi_D(-1) = -1$ and $p \nmid ND[\text{Sp}_4(\mathbf{Z}) : \Gamma_0^{(2)}(N)]$. Here $\Gamma_0^{(2)}(N)$ is the congruence subgroup of $\text{Sp}_4(\mathbf{Z})$ of level N . Let $f \in S_{2k-2}(1)$ be a p -ordinary newform. Also assume that there exists a Dirichlet character ψ of conductor N such that

$$\text{val}_{\varpi}(L^{\Sigma}(3-k, \psi)L^{\text{alg}}(k-1, f, \chi_D)L^{\text{alg}}(1, f, \psi)L^{\text{alg}}(2, f, \psi)) = 0.$$

For precise definitions of the L -factors, cf. [Bro11], section 5.

Theorem 10.2. *Assume Assumption 10.1. Let $\rho_f : G_{\Sigma} \rightarrow \text{GL}_2(E)$ be the Galois representation attached to the newform f and write ρ for the residual representation $\bar{\rho}_f$, which we assume to be absolutely irreducible. Here $\Sigma = \{p\}$. Assume that*

$$\dim_{\mathbf{F}} H_f^1(\mathbf{Q}, \rho(1-k)) = \dim_{\mathbf{F}} H_f^1(\mathbf{Q}, \rho(2-k)) = 1.$$

Furthermore assume that R_ρ is a dvr. Let $\sigma : G_\Sigma \rightarrow \mathrm{GL}_4(E)$ be an absolutely irreducible short crystalline Galois representation with $\det \sigma = \epsilon^{4k-6}$, satisfying $\sigma^\vee \cong \sigma(3-2k)$ and such that its residual representation $\bar{\sigma}$ (defined only up to semisimplification) satisfies

$$\bar{\sigma}^{\mathrm{ss}} \cong \chi^{k-2} \oplus \rho \oplus \chi^{k-1}.$$

Then σ is modular, i.e., there exists a cuspidal non-lift Siegel modular eigenform F of weight k , level one such that $L_{\mathrm{spin}}(s, F) = L(s, \sigma)$.

Proof. Let σ be as above. Then by Corollary 4.3, there is a G_Σ -stable lattice Λ in the space of σ such that

$$\bar{\sigma} = \begin{bmatrix} \chi^{k-2} & a & b \\ & \rho & c \\ & & \chi^{k-1} \end{bmatrix}$$

with a and c non-trivial classes in $H_f^1(\mathbf{Q}, \rho(1-k))$. Let R^{red} denote, as before, the (reduced, self-dual) universal deformation ring of $\bar{\sigma}$.

Let Φ' denote the set of linearly independent Siegel modular eigenforms of weight k , level 1 which are orthogonal to the subspace spanned by Saito-Kurokawa lifts. By Theorems 8.2 and 8.4 we get that for every $F \in \Phi'$ there exists a Galois representation

$$\sigma_F : G_\Sigma \rightarrow \mathrm{GL}_4(E)$$

which is short crystalline, satisfies $\det \sigma_F = \epsilon^{4k-6}$ and is τ -self-dual. Let $\Phi \subset \Phi'$ be the subset consisting of forms F such that $\bar{\sigma}_F^{\mathrm{ss}} \cong \chi^{k-1} \oplus \rho \oplus \chi^{k-2}$. For $F \in \Phi$ the representation σ_F is absolutely irreducible which can be proven as in [Bro07] (cf. the arguments following the proof of Proposition 8.3). Write \mathbf{T} for the quotient of \mathbf{T}'_m acting on the space spanned by Φ . Here \mathbf{T}' denotes the Hecke algebra generated over \mathcal{O} by the operators $T_{\ell,0}$, $T_{\ell,1}$ and $T_{\ell,2}$ for all primes $\ell \neq p$ and \mathbf{T}'_m denotes the completion of \mathbf{T}' at the maximal ideal corresponding to Φ .

Thanks to Kato's result (Theorem 17.14 in [Kat04]) we can use an argument similar to that in the proof of Proposition 2.10 to deduce that

$$(10.1) \quad \#H_f^1(\mathbf{Q}, \rho_f(1-k) \otimes E/\mathcal{O}) \leq \#\mathcal{O}/L^{\mathrm{alg}}(k, f).$$

Here $L^{\mathrm{alg}}(k, f)$ is the algebraic part of the L -function of f (for precise normalization see [Bro11]). For this we apply an analogue of the control theorem (Theorem 2.11) for $\xi = \epsilon^{-m}$ with $m = k-1$ and the fact that the p -adic L -function interpolates the classical L -value at critical points.

We will now demonstrate that Φ is non-empty by showing that $\mathbf{T} \neq 0$. Since the Selmer group on the left of (10.1) is non-trivial, we get that $\mathrm{val}_\varpi(L^{\mathrm{alg}}(k, f)) > 0$. Let $\mathrm{SK}(f)$ denote the Saito-Kurokawa lift of f . It is a Siegel modular eigenform of weight k and level 1. Let J be the ideal of \mathbf{T} generated by the image of the annihilator $\mathrm{Ann}_{\mathbf{T}'_m}(\mathrm{SK}(f))$ of $\mathrm{SK}(f)$ under the map $\mathbf{T}'_m \twoheadrightarrow \mathbf{T}$. Then it follows from Corollary 5.6 in [Bro11] that $\#\mathbf{T}/J \geq \#\mathcal{O}/L^{\mathrm{alg}}(k, f)$. In particular $\Phi \neq \emptyset$.

For every $F \in \Phi$, we get by Corollary 4.3 that we may assume that $\bar{\sigma}_F = \begin{bmatrix} \chi^{k-2} & *_1 & *_2 \\ & \rho & *_3 \\ & & \chi^{k-1} \end{bmatrix}$. Since σ_F are short crystalline (by our assumption that $p > 2k - 2$) Proposition 5.1 then shows that we have $\bar{\sigma}_F \cong \bar{\sigma}$. We further get an \mathcal{O} -algebra map

$$\phi : R^{\text{red}} \rightarrow \prod_{F \in \Phi} \mathcal{O} \supset \mathbf{T}.$$

Clearly $\phi(R^{\text{red}}) \supset \mathbf{T}$, but we in fact get that $\phi(R^{\text{red}}) = \mathbf{T}$ by Corollary 7.7. The map ϕ descends to a map $\bar{\phi} : R^{\text{red}}/I^{\text{tot}} \rightarrow \mathbf{T}/J$. Now combining Corollary 7.4 with Theorem 7.6, inequality (10.1) and the fact $\#\mathbf{T}/J \geq \#\mathcal{O}/L^{\text{alg}}(k, f)$, we conclude that $\bar{\phi}$ is an isomorphism. Since I^{tot} is principal by Theorem 6.6, ϕ is also an isomorphism by the commutative algebra criterion [BK13], Theorem 4.1. \square

REFERENCES

- AL70. A. O. L. Atkin and J. Lehner, *Hecke operators on $\Gamma_0(m)$* , Math. Ann. **185** (1970), 134–160.
- BC09. J. Bellaïche and G. Chenevier, *p -adic families of Galois representations and higher rank Selmer groups*, Astérisque (2009), no. 324.
- BCGP18. G. Boxer, F. Calegari, T. Gee, and V. Pilloni, *Abelian surfaces over totally real fields are potentially modular*, Preprint (2018), arXiv: 1812.09269.
- BCKL05. J. Bellaïche, G. Chenevier, C. Khare, and M. Larsen, *Converging sequences of p -adic Galois representations and density theorems*, Int. Math. Res. Not. (2005), no. 59, 3691–3720.
- BCP97. W. Bosma, J. Cannon, and C. Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), no. 3-4, 235–265, Computational algebra and number theory (London, 1993).
- BDPS15. T. Berger, L. Dembélé, A. Pacetti, and M. H. Sengün, *Theta lifts of Bianchi modular forms and applications to paramodularity*, J. Lond. Math. Soc. (2) **92** (2015), no. 2, 353–370.
- BK90. S. Bloch and K. Kato, *L -functions and Tamagawa numbers of motives*, The Grothendieck Festschrift, Vol. I, Progr. Math., vol. 86, Birkhäuser Boston, Boston, MA, 1990, pp. 333–400.
- BK13. T. Berger and K. Klosin, *On deformation rings of residually reducible Galois representations and $R = T$ theorems*, Math. Ann. **355** (2013), no. 2, 481–518.
- BK14. A. Brumer and K. Kramer, *Paramodular abelian varieties of odd conductor*, Trans. Amer. Math. Soc. **366** (2014), no. 5, 2463–2516.
- BPP⁺ar. A. Brumer, A. Pacetti, C. Poor, G. Tornaria, J. Voight, and D. S. Yuen, *On the paramodularity of typical abelian surfaces*, Algebra and Number Theory (to appear).
- Bro07. J. Brown, *Saito-Kurokawa lifts and applications to the Bloch-Kato conjecture*, Compos. Math. **143** (2007), no. 2, 290–322.
- Bro08. ———, *Residually reducible representations of algebras over local Artinian rings*, Proc. Amer. Math. Soc. **136** (2008), no. 10, 3409–3414.
- Bro11. ———, *On the cuspidality of pullbacks of Siegel Eisenstein series and applications*, Int. Math. Res. Not. IMRN **7** (2011), 1706–1756.
- Cal06. F. Calegari, *Eisenstein deformation rings*, Compos. Math. **142** (2006), no. 1, 63–83.
- CHT08. L. Clozel, M. Harris, and R. Taylor, *Automorphy for some l -adic lifts of automorphic mod l Galois representations*, Publ. Math. Inst. Hautes Études Sci.

- (2008), no. 108, 1–181, With Appendix A, summarizing unpublished work of Russ Mann, and Appendix B by Marie-France Vignéras.
- DFG04. F. Diamond, M. Flach, and L. Guo, *The Tamagawa number conjecture of adjoint motives of modular forms*, Ann. Sci. École Norm. Sup. (4) **37** (2004).
- Fal83. G. Faltings, *Endlichkeitssätze für abelsche Varietäten über Zahlkörpern*, Invent. Math. **73** (1983), no. 3, 349–366.
- FC90. G. Faltings and C.-L. Chai, *Degeneration of abelian varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 22, Springer-Verlag, Berlin, 1990, With an appendix by David Mumford.
- FL82. J.-M. Fontaine and G. Laffaille, *Construction de représentations p -adiques*, Ann. Sci. École Norm. Sup. (4) **15** (1982), no. 4, 547–608 (1983).
- Fly90. E. V. Flynn, *Large rational torsion on abelian varieties*, J. Number Theory **36** (1990), no. 3, 257–265.
- Fon82. J.-M. Fontaine, *Sur certains types de représentations p -adiques du groupe de Galois d'un corps local; construction d'un anneau de Barsotti-Tate*, Ann. of Math. (2) **115** (1982), no. 3, 529–577.
- GJP⁺09. G. Grigorov, A. Jorza, S. Patrikis, W. A. Stein, and C. Tarnita, *Computational verification of the Birch and Swinnerton-Dyer conjecture for individual elliptic curves*, Math. Comp. **78** (2009), no. 268, 2397–2425.
- Hid00. H. Hida, *Modular forms and Galois cohomology*, Cambridge Studies in Advanced Mathematics, vol. 69, Cambridge University Press, Cambridge, 2000.
- JLR12. J. Johnson-Leung and B. Roberts, *Siegel modular forms of degree two attached to Hilbert modular forms*, J. Number Theory **132** (2012), no. 4, 543–564.
- Jor10. A. Jorza, *Crystalline representations for GL_2 over quadratic imaginary fields*, Ph.D. thesis, Princeton University, 2010.
- Jor12. ———, *p -adic families and Galois representations for $GS_p(4)$ and $GL(2)$* , Math. Res. Lett. **19** (2012), no. 5, 987–996.
- JSW17. D. Jetchev, C. Skinner, and X. Wan, *The Birch and Swinnerton-Dyer formula for elliptic curves of analytic rank one*, Camb. J. Math. **5** (2017), no. 3, 369–434.
- Kat04. K. Kato, *p -adic Hodge theory and values of zeta functions of modular forms*, Astérisque (2004), no. 295, ix, 117–290, Cohomologies p -adiques et applications arithmétiques. III.
- Klo09. K. Klosin, *Congruences among automorphic forms on $U(2, 2)$ and the Bloch-Kato conjecture*, Annales de l'institut Fourier **59** (2009), no. 1, 81–166.
- KRS92. S. S. Kudla, S. Rallis, and D. Soudry, *On the degree 5 L -function for $Sp(2)$* , Invent. Math. **107** (1992), no. 3, 483–541.
- KW09. C. Khare and J.-P. Wintenberger, *Serre's modularity conjecture. I*, Invent. Math. **178** (2009), no. 3, 485–504.
- Lau05. G. Laumon, *Fonctions zêtas des variétés de Siegel de dimension trois*, Astérisque (2005), no. 302, 1–66, Formes automorphes. II. Le cas du groupe $GSp(4)$.
- LD98. P. Le Duff, *Représentations galoisiennes associées aux points d'ordre l des jacobiniennes de certaines courbes de genre 2*, Bull. Soc. Math. France **126** (1998), no. 4, 507–524.
- LMF17a. The LMFDB Collaboration, *The L -functions and Modular Forms Database*, Home page of the Elliptic Curve Isogeny Class 731.a.1, <http://www.lmfdb.org/EllipticCurve/Q/731/a/1>, 2017, [Online; accessed 12 October 2017].
- LMF17b. ———, *The L -functions and Modular Forms Database*, Home page of the Genus 2 Curve Isogeny Class 731.a.12427.1, <http://www.lmfdb.org/Genus2Curve/Q/731/a/12427/1>, 2017, [Online; accessed 12 October 2017].
- Maz97. B. Mazur, *An introduction to the deformation theory of Galois representations*, Modular forms and Fermat's last theorem (Boston, MA, 1995), Springer, New York, 1997, pp. 243–311.

- Miy89. T. Miyake, *Modular forms*, Springer-Verlag, Berlin, 1989, Translated from the Japanese by Yoshitaka Maeda.
- Mok14. C. P. Mok, *Galois representations attached to automorphic forms on GL_2 over CM fields*, Compos. Math. **150** (2014), no. 4, 523–567.
- MT15. C. P. Mok and F. Tan, *Overconvergent families of Siegel-Hilbert modular forms*, Canad. J. Math. **67** (2015), no. 4, 893–922.
- Och00. T. Ochiai, *Control theorem for Bloch-Kato’s Selmer groups of p -adic representations*, J. Number Theory **82** (2000), no. 1, 69–90.
- Pil12. V. Pilloni, *Modularité, formes de Siegel et surfaces abéliennes*, J. Reine Angew. Math. **666** (2012), 35–82.
- Pil17. ———, *Higher coherent cohomology and p -adic modular forms of singular weight*, preprint available at <http://perso.ens-lyon.fr/vincent.pilloni/complexhidatheorygsp4.pdf>.
- PW11. R. Pollack and T. Weston, *On anticyclotomic μ -invariants of modular forms*, Compos. Math. **147** (2011), no. 5, 1353–1381.
- PY15. C. Poor and D. S. Yuen, *Paramodular cusp forms*, Math. Comp. **84** (2015), no. 293, 1401–1438.
- Ram93. R. Ramakrishna, *On a variation of Mazur’s deformation functor*, Compositio Math. **87** (1993), no. 3, 269–286.
- Ram13. D. Ramakrishnan, *Decomposition and parity of Galois representations attached to $GL(4)$* , Automorphic representations and L -functions, Tata Inst. Fundam. Res. Stud. Math., vol. 22, Tata Inst. Fund. Res., Mumbai, 2013, pp. 427–454.
- Rib76. K. A. Ribet, *A modular construction of unramified p -extensions of $Q(\mu_p)$* , Invent. Math. **34** (1976), no. 3, 151–162.
- RS07. B. Roberts and R. Schmidt, *Local newforms for $GSp(4)$* , Lecture Notes in Mathematics, vol. 1918, Springer, Berlin, 2007.
- Rub00. K. Rubin, *Euler systems*, Annals of Mathematics Studies, vol. 147, Princeton University Press, Princeton, NJ, 2000, Hermann Weyl Lectures. The Institute for Advanced Study.
- Sch17. R. Schmidt, *Packet structure and paramodular forms*, Transactions of the AMS (2017).
- Ser87. J.-P. Serre, *Sur les représentations modulaires de degré 2 de $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$* , Duke Math. J. **54** (1987), no. 1, 179–230.
- Shi71. G. Shimura, *Introduction to the arithmetic theory of automorphic functions*, Publications of the Mathematical Society of Japan, No. 11. Iwanami Shoten, Publishers, Tokyo, 1971, Kanô Memorial Lectures, No. 1.
- Ski16. C. Skinner, *Multiplicative reduction and the cyclotomic main conjecture for GL_2* , Pacific J. Math. **283** (2016), no. 1, 171–200.
- Sor10. C. M. Sorensen, *Galois representations attached to Hilbert-Siegel modular forms*, Doc. Math. **15** (2010), 623–670.
- ST68. J.-P. Serre and J. Tate, *Good reduction of abelian varieties*, Ann. of Math. (2) **88** (1968), 492–517.
- SU06. C. Skinner and E. Urban, *Sur les déformations p -adiques de certaines représentations automorphes*, J. Inst. Math. Jussieu **5** (2006), no. 4, 629–698.
- SU14. ———, *The Iwasawa Main Conjectures for GL_2* , Invent. Math. **195** (2014), no. 1, 1–277.
- Tay91. R. Taylor, *Galois representations associated to Siegel modular forms of low weight*, Duke Math. J. **63** (1991), no. 2, 281–332.
- Til06. J. Tilouine, *Nearly ordinary rank four Galois representations and p -adic Siegel modular forms*, Compos. Math. **142** (2006), no. 5, 1122–1156, With an appendix by Don Blasius.
- Urb99. E. Urban, *On residually reducible representations on local rings*, J. Algebra **212** (1999), no. 2, 738–742.
- Was97. L. C. Washington, *Introduction to cyclotomic fields*, second ed., Graduate Texts in Mathematics, vol. 83, Springer-Verlag, New York, 1997.

- Wei05. R. Weissauer, *Four dimensional Galois representations*, Astérisque (2005), no. 302, 67–150, Formes automorphes. II. Le cas du groupe $GS(4)$.
- Wes00. T. Weston, *On Selmer groups of geometric Galois representations*, Thesis, Harvard University, Cambridge, 2000.
- Yos80. H. Yoshida, *Siegel's modular forms and the arithmetic of quadratic forms*, Invent. Math. **60** (1980), no. 3, 193–248.

APPENDIX A. BY CRIS POOR³, JERRY SHURMAN⁴, AND DAVID S. YUEN⁵

We prove the following theorem.

Theorem A.1. *The space $\mathcal{S}_2(\mathbb{K}(731))$ has dimension 19, and its subspace of Gritsenko lifts has dimension 18. There exists a new Hecke eigenform $f_{731} \in \mathcal{S}_2(\mathbb{K}(731))$ such that:*

- (1) f_{731} has integer Fourier coefficients with content 1.
- (2) f_{731} is not a Gritsenko lift.
- (3) f_{731} is congruent modulo 5 to a Gritsenko lift Hecke eigenform that has integer Fourier coefficients.
- (4) the first few Hecke eigenvalues of f_{731} are

$$\lambda_2 = -1, \lambda_3 = 0, \lambda_4 = -2, \lambda_5 = 0,$$

and its spin 2-Euler factor is

$$1 + x + 2x^2 + 2x^3 + 4x^4.$$

The proof will take several steps. The existence argument is motivated by techniques in [8, 6, 11], while the congruence is shown using a new technique.

A.1. Notation and Jacobi Restriction method. Let $\mathcal{S}_k(\mathbb{K}(N))$ denote the space of weight k Siegel cusp forms for the level N paramodular group, or, more briefly, the space of weight k , level N paramodular cusp forms. We use the notation “[d]” for the image under projection onto the first d Jacobi coefficients, and “(d)” for the subspace where the first $d - 1$ Jacobi coefficients vanish; because all Jacobi coefficients have index divisible by N , this condition is that the Jacobi coefficients of index $N, 2N, \dots, (d - 1)N$ are 0. Thus

$\mathcal{S}_k(\mathbb{K}(N))[d] = \mathcal{S}_k(\mathbb{K}(N))$ elements truncated to the first d Jacobi coefficients,
 $\mathcal{S}_k(\mathbb{K}(N))(d) = \mathcal{S}_k(\mathbb{K}(N))$ elements with vanishing first $d - 1$ Jacobi coefficients,

and there is an exact sequence

$$0 \longrightarrow \mathcal{S}_k(\mathbb{K}(N))(d + 1) \longrightarrow \mathcal{S}_k(\mathbb{K}(N)) \longrightarrow \mathcal{S}_k(\mathbb{K}(N))[d] \longrightarrow 0.$$

Analogous notation is understood to hold with either Fricke eigenspace $\mathcal{S}_k(\mathbb{K}(N))^\pm$ of $\mathcal{S}_k(\mathbb{K}(N))$, or any Atkin–Lehner eigenspace, and the exact sequence persists.

Let $J_{k,m}^{\text{cusp}}$ denote the space of weight k , index m Jacobi cusp forms. Using the Fourier–Jacobi expansions of paramodular cusp forms, we view two maps as containments for simplicity,

$$\mathcal{S}_k(\mathbb{K}(N)) \subset \bigoplus_{j=1}^{\infty} J_{k,jN}^{\text{cusp}} \subset \mathbb{C}^\infty.$$

The latter containment entails some chosen ordering of the index sets for Fourier expansions of Jacobi forms. For any prime p , let a subscript p denote the map from subsets of \mathbb{C}^∞ to subsets of \mathbb{F}_p^∞ that reduces integral elements modulo p ,

$$V_p = (V \cap \mathbb{Z}^\infty) \bmod p, \quad V \subset \mathbb{C}^\infty.$$

Thus

$$\mathcal{S}_k(\mathbb{K}(N))_p \subset \bigoplus_{j=1}^{\infty} (\mathcal{J}_{k,jN}^{\text{cusp}})_p \subset \mathbb{F}_p^\infty.$$

The spaces $\mathcal{S}_k(\mathbb{K}(N))$ and $\mathcal{J}_{k,m}^{\text{cusp}}$ have bases in \mathbb{Z}^∞ by results of [16] and [3], respectively. Thus $\dim_{\mathbb{F}_p} \mathcal{S}_k(\mathbb{K}(N))_p = \dim_{\mathbb{C}} \mathcal{S}_k(\mathbb{K}(N))$ and $\dim_{\mathbb{F}_p} (\mathcal{J}_{k,m}^{\text{cusp}})_p = \dim_{\mathbb{C}} \mathcal{J}_{k,m}^{\text{cusp}}$. Extending the notation from above,

$\mathcal{S}_k(\mathbb{K}(N))[d]_p =$ reduction modulo p of elements of $\mathcal{S}_k(\mathbb{K}(N))[d] \cap \mathbb{Z}^\infty$,
 $\mathcal{S}_k(\mathbb{K}(N))_p =$ reduction modulo p of elements in $\mathcal{S}_k(\mathbb{K}(N)) \cap \mathbb{Z}^\infty$,
 $\mathcal{S}_k(\mathbb{K}(N))_p(d) =$ elements of $\mathcal{S}_k(\mathbb{K}(N))_p$ with first $d - 1$ Jacobi coefficients 0,
and in consequence of an integral basis of $\mathcal{S}_k(\mathbb{K}(N))$,

$$\begin{aligned} \dim \mathcal{S}_k(\mathbb{K}(N))[d] &= \dim \mathcal{S}_k(\mathbb{K}(N))[d]_p, \\ \dim \mathcal{S}_k(\mathbb{K}(N))(d) &\leq \dim \mathcal{S}_k(\mathbb{K}(N))_p(d). \end{aligned}$$

Again, analogous notation is understood to hold with either Fricke eigenspace or any Atkin–Lehner eigenspace of $\mathcal{S}_k(\mathbb{K}(N))$, and the dimension relations persist.

For either Fricke eigenvalue $\epsilon = \pm 1$, the Jacobi Restriction method [2, 5] runs with d as a parameter, returning a basis of a finite-dimensional complex vector space \mathcal{J}_d^ϵ of truncated formal Fourier–Jacobi expansions such that

$$\mathcal{S}_k(\mathbb{K}(N))^\epsilon[d] \subset \mathcal{J}_d^\epsilon \subset \bigoplus_{j=1}^d \mathcal{J}_{k,jN}^{\text{cusp}}.$$

An additional parameter, \det_{\max} , bounds the Fourier coefficient index determinants in the calculation, thus making the method an algorithm; this parameter must be chosen so that each space $\mathcal{J}_{k,jN}^{\text{cusp}}$ with $j \leq d$ is determined by the Fourier coefficients whose indices satisfy the determinant bound. For simplicity we suppress \det_{\max} from the notation \mathcal{J}_d^ϵ , along with the weight k and the level N . We can also run the Jacobi Restriction method modulo a prime p . In this case, the algorithm returns a basis of a finite dimensional vector space $\mathcal{J}_{d,p}^\epsilon$ over \mathbb{F}_p such that

$$\mathcal{S}_k(\mathbb{K}(N))^\epsilon[d]_p \subset \mathcal{J}_{d,p}^\epsilon \subset \bigoplus_{j=1}^d (\mathcal{J}_{k,jN}^{\text{cusp}})_p.$$

The Jacobi Restriction method can also be run on any Atkin–Lehner eigenspace. For a vector of signs describing an Atkin–Lehner eigenspace, the Jacobi Restriction method returns a basis of a space $\mathcal{J}_d^{\text{signs}}$ or $\mathcal{J}_{d,p}^{\text{signs}}$ that contains $\mathcal{S}_k(\mathbb{K}(N))^{\text{signs}}[d]$ or $\mathcal{S}_k(\mathbb{K}(N))^{\text{signs}}[d]_p$.

The methods of [11] let us try to show that $\mathcal{S}_2(\mathbb{K}(N))(d)$ or $\mathcal{S}_2(\mathbb{K}(N))_p(d)$ is 0 for a certain d . The method we will use here requires us to span enough

of $\mathcal{S}_4(\mathbb{K}(N))$, namely a subspace of codimension less than $\dim \mathbb{J}_{2,N}^{\text{cusp}}$. A formula from [4] gives $\dim \mathcal{S}_4(\mathbb{K}(731)) = 972$, and we also have $\dim \mathbb{J}_{2,731}^{\text{cusp}} = 18$ from [3].

A.2. Spanning $\mathcal{S}_4(\mathbb{K}(731))$. For $\epsilon = (-1)^k$, let $\text{Grit} : \mathbb{J}_{k,N}^{\text{cusp}} \rightarrow \mathcal{S}_k(\mathbb{K}(N))^\epsilon$ be the linear map sending a Jacobi cusp form to its Gritsenko lift. Let g_1, \dots, g_{18} be a basis of the space of Gritsenko lifts in the Fricke plus space $\mathcal{S}_2(\mathbb{K}(731))^+$, and let $\tilde{g}_1, \dots, \tilde{g}_{138}$ be a basis of the space of Gritsenko lifts in the Fricke plus space $\mathcal{S}_4(\mathbb{K}(731))^+$. The set

$$S = \{T_\ell(g_i g_j), \tilde{g}_k : 1 \leq i \leq j \leq 18, 1 \leq \ell \leq 4, 1 \leq k \leq 138\}$$

turns out to span at least a 719-dimensional space in $\mathcal{S}_4(\mathbb{K}(731))^+$, as shown by its Fourier coefficients for indices out to determinant 206. Establishing this required expanding the g_i initially out to indices of determinant 3296. Thus $\dim \mathcal{S}_4(\mathbb{K}(731))^+ \geq 719$, and we may select a basis of a 719-dimensional subspace. For $\ell \nmid N$, the Hecke operator T_ℓ employed here is the $T_{\ell,2}$ of the main article.

Note that $731 = 17 \cdot 43$. Let $\mathcal{S}_4(\mathbb{K}(731))^{+,-}$ denote the subspace of $\mathcal{S}_4(\mathbb{K}(731))$ where the Atkin–Lehner operators AL_{17} and AL_{43} have respective eigenvalues ± 1 , and similarly for $\mathcal{S}_4(\mathbb{K}(731))^{-,+}$. Jacobi Restriction with parameters $d = 6$, $p = 5$, and $\det_{\max} = 18275$ on these Atkin–Lehner subspaces returns spaces $\mathcal{J}_{6,5}^{+,-}$ and $\mathcal{J}_{6,5}^{-,+}$ such that

$$\begin{aligned} \mathcal{S}_4(\mathbb{K}(731))^{+,-}[6]_5 &\subset \mathcal{J}_{6,5}^{+,-}, & \dim_{\mathbb{F}_5} \mathcal{J}_{6,5}^{+,-} &= 125, \\ \mathcal{S}_4(\mathbb{K}(731))^{-,+}[6]_5 &\subset \mathcal{J}_{6,5}^{-,+}, & \dim_{\mathbb{F}_5} \mathcal{J}_{6,5}^{-,+} &= 128. \end{aligned}$$

Even though $719 + 125 + 128 = 972$, it is not mathematically certain that 719, 125, and 128 are the dimensions of the Fricke plus space and the two Atkin–Lehner subspaces of the Fricke minus space in $\mathcal{S}_4(\mathbb{K}(731))$; however, in practice we are confident that they are, and this guided our search for spanning elements.

We construct a Borcherds product $f_{4,731} \in \mathcal{S}_4(\mathbb{K}(731))^-$. The website [17] gives the details of the construction, but here we note that the Borcherds product’s leading theta block was located among more than 354,000 candidate theta blocks in $\mathbb{J}_{4,731}^{\text{cusp}}$ of q -vanishing order 2. Its symmetrizations

$$h^{+,-}, h^{-,+} = f_{4,731} \pm \text{AL}_{17}(f_{4,731}) \in \mathcal{S}_4(\mathbb{K}(731))^{+,-}, \mathcal{S}_4(\mathbb{K}(731))^{-,+}$$

are both nonzero. To generate many more elements of these Atkin–Lehner subspaces, we use a technique called *bootstrapping* [10], created to overcome the obstacle that, computationally, a Hecke operator returns shorter truncations of paramodular Fourier expansions than it receives. This shortening allows only a few iterations of Hecke operators, even starting with very long vectors that are expensive to compute. A brief summary of bootstrapping in the present context is as follows.

- (1) We have a sufficiently long expansion of $h^{+,-}$ to identify it modulo 5 in $\mathcal{J}_{6,5}^{+,-}$. The vector that we identify it as modulo 5 is longer.
- (2) Using the longer vector in $\mathcal{J}_{6,5}^{+,-}$, we have sufficiently many Fourier coefficients to apply T_2 , obtaining a shorter expansion of $T_2 h^{+,-}$

modulo 5. But this vector is sufficiently long to identify in $\mathcal{J}_{6,5}^{+,-}$, again giving us a long enough expansion to apply T_2 again, and so on.

This technique shows that $T_2^i h^{+,-}$ for $i = 0, 1, \dots, 124$ are linearly independent modulo 5, and so $\dim_{\mathbb{F}_5} \mathcal{S}_4(\mathbb{K}(731))^{+,-}[6]_5 \geq 125$. Similarly, bootstrapping with $h^{-,+}$ gives $\dim_{\mathbb{F}_5} \mathcal{S}_4(\mathbb{K}(731))^{-,+}[6]_5 \geq 124$. Thus altogether we have shown that $\dim_{\mathbb{F}_5} \mathcal{S}_4(\mathbb{K}(731))^{-}[6]_5 \geq 125 + 124 = 249$, and we may select a basis modulo 5 of a 249-dimensional subspace of $\mathcal{S}_4(\mathbb{K}(731))_5^-$. So, because $719 + 249 = 968$, altogether we have reductions modulo 5 of linearly independent elements that span a codimension 4 subspace of $\mathcal{S}_4(\mathbb{K}(731))$. This is comfortably beyond the needed codimension $\dim J_{2,731}^{\text{cusp}} - 1 = 17$.

A.3. Determining $\mathcal{S}_2(\mathbb{K}(731))^-$. The $H_4(N, 2, 1)^-$ method of Proposition 4.6 in [11] uses a low-codimension subspace in $\mathcal{S}_4(\mathbb{K}(N))^-$ to show that $\mathcal{S}_2(\mathbb{K}(N))_p^-(2) = 0$. Briefly, the idea is as follows. Suppose that some h in $\mathcal{S}_2(\mathbb{K}(731))_5^-(2)$ is nonzero. Then the space $W = h \text{ Grit}(J_{2,731}^{\text{cusp}})_5 \subseteq \mathcal{S}_4(\mathbb{K}(731))_5^-$ has dimension 18. But certain Fourier coefficients of every element w of W must be zero modulo 5, because such w are a particular kind of product. We make a $249 \times r$ matrix M of coefficients from a basis of the known 249-dimensional subspace of $\mathcal{S}_4(\mathbb{K}(731))_5^-$ where the r coefficients of every $w \in W$ must be zero, and we compute the rank of M modulo 5 to be 249. Because the 249-dimensional subspace has codimension at most 4 in $\mathcal{S}_4(\mathbb{K}(731))_5^-$, we conclude that $\dim W \leq 4$, contradicting $\dim W = 18$. This proves that $\mathcal{S}_2(\mathbb{K}(731))_5^-(2) = 0$, which in turn proves that $\mathcal{S}_2(\mathbb{K}(731))_5^-(2) = 0$. Further, Jacobi Restriction with parameters $d = 1$, $p = 5$, and $\det_{\max} = 1462$ gives $\mathcal{J}_{1,5}^- = 0$, and so $\mathcal{S}_2(\mathbb{K}(731))_5^-[1] = 0$, and the exact sequence gives $\mathcal{S}_2(\mathbb{K}(731))^- = 0$.

A.4. Determining $\mathcal{S}_2(\mathbb{K}(731))^+$. The $H_4(N, 6, 1)^+$ method of Proposition 4.6 in [11] shows that $\mathcal{S}_2(\mathbb{K}(731))_5^+(6) = 0$, using very similar reasoning. This implies that $\mathcal{S}_2(\mathbb{K}(731))_5^+(6) = 0$. These critical vanishing results are ultimately what allow us to prove equalities and congruences in $\mathcal{S}_2(\mathbb{K}(731))$. The exact sequence says that the projection map $\mathcal{S}_2(\mathbb{K}(731))^+ \rightarrow \mathcal{S}_2(\mathbb{K}(731))^+[5]$ is an isomorphism. Further, Jacobi Restriction with parameters $d = 5$, $p = 5$, and $\det_{\max} = 1462$ gives $\dim \mathcal{J}_{5,5}^+ = 19$, so $\dim \mathcal{S}_2(\mathbb{K}(731))^+[5] \leq 19$. Thus $\dim \mathcal{S}_2(\mathbb{K}(731))^+ \leq 19$, which is to say that there is at most one dimension of nonlift eigenforms.

The existence of a nonlift, given in section A.6, will complete the proof that $\dim \mathcal{S}_2(\mathbb{K}(731))^+ = 19$.

A.5. Borchers products. Our current best method for finding a paramodular form that is not a Gritsenko lift in the low weight 2 and the fairly high level 731 is to try constructing Borchers products (see [8, 9, 11]).

However, the algorithm from [9] shows that there are no nonlift Borchers products in $\mathcal{S}_2(\mathbb{K}(731))$. The method from [6, 8] is to go to twice the level and search for a Borchers product in the minus Fricke subspace $\mathcal{S}_2(\mathbb{K}(1462))^-$.

Indeed, we find two Borcherds products

$$f_{2,1462,a}, f_{2,1462,b} \in \mathcal{S}_2(\mathbb{K}(1462))^-.$$

See the website [17] for their constructions.

A.6. Tracing down from $\mathcal{S}_2(\mathbb{K}(1462))^-$ to $\mathcal{S}_2(\mathbb{K}(731))^+$. We prove the existence of a nonlift in $\mathcal{S}_2(\mathbb{K}(731))^+$ by using the trace down map from [11],

$$\mathrm{TrDn} : \mathcal{S}_2(\mathbb{K}(1462)) \rightarrow \mathcal{S}_2(\mathbb{K}(731))^+.$$

Here the codomain is the Fricke plus space because $\mathcal{S}_2(\mathbb{K}(731))^- = 0$, as shown above. Tracing down the Borcherds products $f_{2,1462,a}$ and $f_{2,1462,b}$ requires many more Fourier coefficients than we can directly compute. So we run Jacobi Restriction on $\mathcal{S}_2(\mathbb{K}(1462))^-$ with parameters $d = 4$ and $\det_{\max} = 5848$, and get $\dim \mathcal{J}_4^- = 2$. After identifying the projections of the Borcherds products in \mathcal{J}_4^- , we can compute two Fourier coefficients of $\mathrm{TrDn}(f_{2,1462,a})$ and $\mathrm{TrDn}(f_{2,1462,b})$, but this is insufficient for our purposes. To remedy this, we use the fact that the Borcherds products are Fricke minus forms to compute further Fourier coefficients for indices that are beyond the fourth Jacobi coefficient but still within the \det_{\max} bound. This permits the computation of 46 Fourier coefficients of $\mathrm{TrDn}(f_{2,1462,a})$ and $\mathrm{TrDn}(f_{2,1462,b})$. The computed coefficients of $\mathrm{TrDn}(f_{2,1462,a})$ show that it is linearly independent of the Gritsenko lifts in $\mathcal{S}_2(\mathbb{K}(731))^+$. This proves that $\dim \mathcal{S}_2(\mathbb{K}(731))^+ = 19$.

Because there is only one nonlift eigenform in $\mathcal{S}_2(\mathbb{K}(731))^+$, it must scale to have rational coefficients, and then rescale to a nonlift eigenform f_{731} that has integer Fourier coefficients with content 1. Jacobi Restriction with parameters $d = 5$ and $\det_{\max} = 1462$ returns $\dim \mathcal{J}_5^+ = 19$. Hence $\mathcal{S}_2(\mathbb{K}(731))^+[5] = \mathcal{J}_5^+$ and we have enough coefficients to compute eigenvalues on the Hecke operators through T_5 . Especially, the nonlift eigenform f_{731} has eigenvalues

$$\lambda_2 = -1, \lambda_3 = 0, \lambda_4 = -2, \lambda_5 = 0.$$

We refer to [13, RS07] for the theory of global paramodular newforms, and note here that the nonlift eigenform f_{731} must be a newform because by [12] the spaces $\mathcal{S}_2(\mathbb{K}(1))$, $\mathcal{S}_2(\mathbb{K}(17))$, and $\mathcal{S}_2(\mathbb{K}(43))$ have no nonlifts.

A.7. Proving the mod 5 congruence. First we compute the action of T_2 on the 18-dimensional space of Gritsenko lifts in $\mathcal{S}_2(\mathbb{K}(731))^+$. The characteristic polynomial of T_2 on this space, factored into irreducibles over \mathbb{Q} , is

$$\begin{aligned} & (x-4)(x-2)^2(x-1) \\ & \cdot (x^6 - 17x^5 + 112x^4 - 358x^3 + 566x^2 - 400x + 101) \\ & \cdot (x^8 - 25x^7 + 264x^6 - 1530x^5 + 5286x^4 - 11046x^3 + 13465x^2 - 8612x + 2174). \end{aligned}$$

Thus there is one Gritsenko lift eigenform with $\lambda_2 = 4$, which we denote g and scale to have integer coefficients with content 1. The first few eigenvalues

of g can be computed,

$$\lambda_2 = 4, \lambda_3 = 5, \lambda_4 = 8, \lambda_5 = 5.$$

We now prove that the Fourier expansion of f_{731} is congruent to an integer multiple of the Fourier expansion of g modulo 5. For any $h \in \mathcal{S}_2(\mathbb{K}(731))^+$ with integer coefficients, introduce correspondingly

$$h[5] \in \mathcal{S}_2(\mathbb{K}(731))^+[5], \quad h_5 \in \mathcal{S}_2(\mathbb{K}(731))_5^+, \quad h[5]_5 \in \mathcal{S}_2(\mathbb{K}(731))^+[5]_5.$$

For any $v \in \mathcal{S}_2(\mathbb{K}(731))^+[5]_5$, there are enough computed Fourier coefficients of T_2v to determine T_2v as an element of $\mathcal{S}_2(\mathbb{K}(731))^+[5]_5$, and so we can compute a matrix representative of the map

$$T_2[5]_5 : \mathcal{S}_2(\mathbb{K}(731))^+[5]_5 \longrightarrow \mathcal{S}_2(\mathbb{K}(731))^+[5]_5.$$

Basic linear algebra on the matrix representative shows that the $T_2[5]_5$ -eigenspace with eigenvalue 4,

$$V = \{h \in \mathcal{S}_2(\mathbb{K}(731))^+[5]_5 : T_2[5]_5 h = 4h\},$$

is 2-dimensional. Although we did not have enough Fourier coefficients to compute the matrix representative of $T_4[5]_5$ on $\mathcal{S}_2(\mathbb{K}(731))^+[5]_5$, we do have enough to compute its matrix representative on the space V , which has a smaller set of determining Fourier coefficients, and then basic linear algebra shows that the $T_4[5]_5$ -eigenspace with eigenvalue 3 in V is 1-dimensional. Both $f_{731}[5]_5$ and $g[5]_5$ lie in V , and both have $T_4[5]_5$ -eigenvalue 3. They cannot be linearly independent, because then they would span V , giving the contradiction that $T_4[5]_5|_V = 3\text{Id}_V$. Thus $f_{731}[5]_5$ and $g[5]_5$ are linearly dependent, and so there exist an integer $\beta \neq 0 \pmod{5}$ such that $f_{731}[5]_5 + \beta g[5]_5 = 0 \pmod{5}$. This implies that

$$(f_{731} + \beta g)_5 \in \mathcal{S}_2(\mathbb{K}(731))_5^+(6).$$

We earlier proved the critical result that $\mathcal{S}_2(\mathbb{K}(731))_5^+(6) = 0$, and this implies $(f_{731} + \beta g)_5 = 0$. This shows that f_{731} is congruent to a multiple of g modulo 5. The multiple of g is still a Gritsenko lift, so f_{731} is congruent modulo 5 to a Gritsenko lift. This completes the proof of Theorem A.1.

We thank Ralf Schmidt for helpful discussions.

APPENDIX REFERENCES

1. J. Arthur. *The endoscopic classification of representations*, volume 61 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2013. Orthogonal and symplectic groups.
2. J. Breeding, II, C. Poor, and D. S. Yuen. Computations of spaces of paramodular forms of general level. *J. Korean Math. Soc.*, 53(3):645–689, 2016.
3. M. Eichler and D. Zagier. *The theory of Jacobi forms*, volume 55 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 1985.
4. T. Ibukiyama and H. Kitayama. Dimension formulas of paramodular forms of square-free level and comparison with inner twist. To appear in *Journal of the Mathematical Society of Japan*.
5. T. Ibukiyama, C. Poor, and D. S. Yuen. Jacobi forms that characterize paramodular forms. *Abh. Math. Semin. Univ. Hambg.*, 83(1):111–128, 2013.
6. C. Poor, R. Schmidt, and D. S. Yuen. Paramodular forms of level 16 and supercuspidal representations. In preparation.

7. C. Poor, R. Schmidt, and D. S. Yuen. Paramodular forms of level 8 and weights 10 and 12. To appear in *Int. J. Num. Th.*
8. C. Poor, J. Shurman, and D. S. Yuen. Nonlift weight two paramodular eigenform constructions. <https://arxiv.org/abs/1805.04137>
9. C. Poor, J. Shurman, and D. S. Yuen. Finding all Borcherds lift paramodular cusp forms of a given weight and level. <https://arxiv.org/abs/1803.11092>
10. C. Poor, J. Shurman, and D. S. Yuen. Weight three paramodular forms and applications. In preparation.
11. C. Poor, J. Shurman, and D. S. Yuen. Siegel paramodular forms of weight 2 and squarefree level. *Int. J. Number Theory*, 13(10):2627–2652, 2017.
12. C. Poor and D. S. Yuen. Paramodular cusp forms. *Math. Comp.*, 84(293):1401–1438, 2015.
13. B. Roberts and R. Schmidt. On modular forms for the paramodular groups. In *Automorphic forms and zeta functions*, pages 334–364. World Sci. Publ., Hackensack, NJ, 2006.
14. B. Roberts and R. Schmidt. *Local Newforms for GSp(4)*, volume 1918 of *Lecture Notes in Mathematics*. Springer, Berlin, 2007.
15. R. Schmidt. Packet structure and paramodular forms. *Transactions of the AMS*, (17), 2017.
16. G. Shimura. On the Fourier coefficients of modular forms of several variables. *Nachr. Akad. Wiss. Göttingen Math.-Phys. Kl. II*, (17):261–268, 1975.
17. D. S. Yuen. Degree 2 Siegel paramodular forms of weight 2 and level 731, 2017. <http://siegelmodularforms.org/pages/degree2/paramodular-wt2-lev731/>.

¹SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SHEFFIELD, HICKS BUILDING, HOUNSFIELD ROAD, SHEFFIELD S3 7RH, UK.

Email address: tberger@cantab.net

²DEPARTMENT OF MATHEMATICS, QUEENS COLLEGE OF THE CITY UNIVERSITY OF NEW YORK, 65-30 KISSENA BLVD, QUEENS, NY 11367, USA

Email address: krzysztof.klosin@qc.cuny.edu

³DEPARTMENT OF MATHEMATICS, FORDHAM UNIVERSITY, BRONX, NY 10458, USA

Email address: poor@fordham.edu

⁴DEPARTMENT OF MATHEMATICS, REED COLLEGE, PORTLAND, OR 97202 USA

Email address: jerry@reed.edu

⁵DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAWAII, 2565 MCCARTHY MALL, HONOLULU, HI, 96822 USA

Email address: yuen888@hawaii.edu