

This is a repository copy of *Contrasting nitrogen fertilisation rates alter mycorrhizal contribution to barley nutrition in a field trial*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/151531/>

Version: Accepted Version

Article:

Thirkell, Thomas James, Cameron, Duncan D. and Hodge, Angela orcid.org/0000-0002-5181-697X (2019) Contrasting nitrogen fertilisation rates alter mycorrhizal contribution to barley nutrition in a field trial. *Frontiers in Plant Science*. 1312. ISSN: 1664-462X

<https://doi.org/10.3389/fpls.2019.01312>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Contrasting nitrogen fertilisation rates alter mycorrhizal contribution to barley nutrition in a field trial

Authors: Tom Thirkell ^{*1,2,a}, Duncan Cameron ², Angela Hodge ¹

* corresponding author.

Author affiliations and contact information:

Tom Thirkell: ¹ Department of Biology, University of York, York, UK; ² Department of Animal and Plant Sciences, University of Sheffield, UK; ^a *Present address:* Centre for Plant Sciences, University of Leeds, UK. *Email:* t.j.thirkell@leeds.ac.uk.

Duncan Cameron: ² Department of Animal and Plant Sciences, University of Sheffield, UK.
Email: d.cameron@sheffield.ac.uk.

Angela Hodge: ¹ Department of Biology, University of York, York, UK. *Email:* angela.hodge@york.ac.uk.

Word count: 4352

Running title: Fungal-mediated barley nitrogen nutrition.

Language style: British English editing preferred.

Abstract

Controlled environment studies show that arbuscular mycorrhizal fungi (AMF) may contribute to plant nitrogen (N) uptake, but the role of these near-ubiquitous symbionts in crop plant N nutrition under natural field conditions remains largely unknown. In a field trial, we tested the effects of N fertilisation and barley (*Hordeum vulgare* L.) cultivar identity on the contribution of AMF to barley N uptake using ^{15}N tracers added to rhizosphere soil compartments. AMF were shown capable of significantly increasing plant ^{15}N acquisition from root exclusion zones, and this was influenced by nitrogen addition type, N fertiliser application rate and barley cultivar identity. Our data demonstrate a previously overlooked potential route of crop plant N uptake which may be influenced substantially and rapidly in response to shifting agricultural management practices.

Key words

Arbuscular mycorrhiza, nitrogen, barley, field trial, plant ecophysiology.

34 Introduction

35 Nitrogen (N) is usually the most limiting mineral nutrient to plant growth (Agren et al., 2012)
36 and maintaining modern agricultural production requires frequent and substantial application
37 of fertiliser to farm soils. In various forms an estimated 50 MT year⁻¹ fertiliser N is applied to
38 agricultural land worldwide (Ladha et al., 2016). Assimilation of applied N by crops may be
39 under 50 % (Ladha et al., 2005, Masclaux-Daubresse et al., 2010); a significant fraction of
40 this applied N is wasted – lost through processes including volatilisation, microbial
41 immobilisation, runoff and leaching (Ladha et al., 2016, Cameron et al., 2013). There is
42 economic and ecological pressure on farmers to optimise the N uptake efficiency of crop
43 plants (Hawkesford, 2014) and by reducing the reliance on non-renewable inputs, improve
44 the sustainability of agriculture (Pretty, 2008). This progress will require the integration of
45 biological and ecological processes into agriculture, and better understanding of soil
46 microbial communities and their roles in nutrient cycling (Rillig et al., 2016, Pretty, 2018).

47 As near-ubiquitous symbionts of cereal crops, arbuscular mycorrhizal fungi (AMF) are prime
48 targets to investigate the role of soil biota in improving agricultural sustainability (Gosling et
49 al., 2006, Thirkell et al., 2017, Rillig et al., 2019). The majority of land plant species engage
50 in symbiosis with these fungi, which may aid plants' mineral nutrient uptake from soils, in
51 exchange for photosynthetic carbon (C) from their plant hosts (Smith and Read, 2008). The
52 influence that AMF mycelia may exert over nutrient dynamics in agricultural systems is not
53 limited to direct effects on plant nutrient acquisition however; the presence of AMF has been
54 shown to reduce mineral fertiliser leaching (Cavagnaro et al., 2015) and to influence
55 greenhouse gas emissions (Storer et al., 2018). While the role of AMF in biogeochemical
56 cycles is undoubtedly complex, of pressing need is to determine the extent to which plants
57 rely on these symbionts for mineral nutrient acquisition.

58 It is well established that AMF can contribute to plant N uptake (Ames et al., 1983, Hodge et
59 al., 2001, Leigh et al., 2009, Thirkell et al., 2016), but the extent to which this takes place,
60 and whether it is ecologically or agriculturally relevant is unclear (Smith and Smith, 2011a).
61 This is in part due to relatively little experimental attention. There remains in the literature a
62 focus on the role of AMF in plant phosphorus (P) uptake (Smith and Smith, 2011a,
63 Karasawa et al., 2012, Ezawa and Saito, 2018), and consideration of symbiotic N uptake is
64 often restricted to diazotrophic bacteria while AMF are often overlooked (Garcia et al., 2016).

65 Improved access to poorly-mobile soil P is, in most instances, the primary benefit of AMF to
66 their plant hosts (Smith and Read, 2008). The relative immobility of inorganic P (Pi) in soil
67 means that plant uptake of Pi from the rhizosphere can outpace Pi diffusion from the
68 surrounding bulk soil and the subsequent P-depletion zones that form around the root are
69 narrow and sharply defined. By engaging in symbiosis with AMF, with a mycelium spreading
70 several centimetres beyond the rhizosphere, the plant effectively increases the volume of
71 soil from which it can acquire nutrients, particularly poorly mobile ions such as Pi (Sanders
72 and Tinker, 1973, Hodge, 2017). Nitrate (NO₃⁻) and ammonium (NH₄⁺), the predominant
73 forms in which plants and fungi acquire N (Marschner, 2011), are more mobile in soil than
74 orthophosphate (Tinker and Nye, 2000). Despite this, a zone of N-depletion may still form
75 around the root (Brackin et al., 2017), in which case AMF may facilitate improved N capture
76 for their plant hosts. With smaller diameters than plant roots, AMF hyphae may also
77 penetrate soil micropores more effectively than a plant root, and thereby be present when
78 inorganic N forms are released through microbial decomposition processes and effectively
79 scavenge for this released inorganic N (Hodge, 2014).

80 Results from microcosm studies are conflicting as to the importance of AMF in plant N
81 uptake (Hodge and Storer, 2015). While a number of studies have shown no improvement

of N uptake by AM plants versus non-mycorrhizal counterparts (Cui and Caldwell, 1996a, Cui and Caldwell, 1996b, Reynolds et al., 2005 Kahkola et al., 2012), it is possible that AMF make an invisible contribution to nutrient acquisition which cannot easily be identified without the use of isotope tracing techniques. Mycorrhizal downregulation of plant root phosphate transporters has been identified in a number of studies (Smith et al., 2003, Smith et al., 2004). In this situation, AMF may be responsible for the majority of a plant's P acquisition, but root transporter downregulation may result in reduced plant P uptake compared to non-mycorrhizal control plants (Smith et al., 2003, Smith et al., 2004). Whether a similar phenomenon occurs in mycorrhizal root N uptake remains unclear. Isotope tracing data does, however, show that AMF can transfer substantial amount of N to a host plant (Leigh et al., 2009, Thirkell et al., 2016), while the contribution of AMF to field-grown plant N uptake is unknown.

AMF are capable of acquiring N from decomposing organic sources (Leigh et al., 2009, Hodge and Fitter, 2010, Barrett et al., 2014, Thirkell et al., 2016) and even to acquire some organic N directly from the hyphosphere, notably as amino acids (Hawkins et al., 2000, Breuninger et al., 2004, Whiteside et al., 2012a, Whiteside et al., 2012b, Tisserant et al., 2012) and perhaps as dipeptides (Belmondo et al., 2014). As in plants however, the vast majority of N acquired by AMF is thought to be as NO_3^- or NH_4^+ (Govindarajulu et al., 2005, Bucking and Kafle, 2015). Greater N uptake as NO_3^- might be expected as it is usually more abundant than NH_4^+ because of rapid nitrification (Marschner, 2011). However, because N acquired as NO_3^- must be reduced to NH_4^+ before further assimilation, it should be energetically favourable for AMF to acquire N as NH_4^+ (Hodge et al., 2010, Courty et al., 2015). Corroborative data remains equivocal as to AMF 'preference' for N types (Johansen et al., 1993, Hawkins and George, 2001). As NO_3^- and NH_4^+ are the most commonly-used forms of fertiliser in Western agriculture, the need to understand mycorrhizal plant acquisition of these N sources is pressing.

Nutrient trade between partners in AM symbioses shows considerable variation in response to biotic factors such as plant and fungal genotype (Smith et al., 2004), in addition to abiotic factors including soil nutrient status (Johnson et al., 2015). Despite substantial experimental data, predictability of the extent to which plants benefit from AMF colonisation remains poor. For example, no universally beneficial fungal isolate has been identified and comparatively few plants are obligate symbionts with AMF.

Despite the widespread distribution of AMF (Smith and Read, 2008, Davison et al., 2015) and the readiness with which they colonise most staple crop plant roots (Smith and Smith, 2011a), little is understood about the function of AMF in the field (Lekberg and Helgason, 2018, Ryan and Graham, 2018). Most published material on the function of AMF is derived from studies conducted under controlled conditions, often comparing AM plants with non-AM controls. While such experiments have provided much valuable data and insight, their findings cannot directly be extrapolated to the field scale, as the occurrence of non-AM cereals in most arable soils is unlikely (Smith and Smith, 2011a). Despite disruptive practices such as tilling and the application of fungicides, there remains a substantial AMF spore bank (and therefore inoculum potential) in agricultural soils (Sosa-Hernandez et al., 2018) and it is very likely that plants in arable field soil will be colonised by AMF (Smith and Smith, 2011a). Further research is needed to begin to understand how AMF might affect crop plant nutrient uptake *in situ*.

Adding ^{15}N isotope tracers to mesh-walled soil compartments in a field trial, we examined the role of AMF in the N acquisition by barley (*Hordeum vulgare* L.) cultivars 'Meridian' and 'Maris Otter'. Isotopic ^{15}N labelling was carried out in plots receiving contrasting N application rates to test the impact of N availability on nutrient transfer in the symbiosis,

testing the hypothesis that increased N fertilisation would result in more AMF transfer of N to host plants. N tracers were added as NH_4^+ or NO_3^- to investigate the relative uptake and transfer of different N sources by AMF.

Materials and Methods

Field trial design

Data were gathered from a larger field trial, designed and implemented at Sancton, East Riding of Yorkshire (co-ordinates 53°51'10.2"N 0°35'29.1"W), by ADAS (Pendeford, Wolverhampton, UK). The ADAS trial was set up to test how barley yield compares among 6 application rates of ammonium nitrate (NH_4NO_3) fertiliser (Nitram, CF Fertiliser, Ince, Cheshire, UK) ranging from 0 – 300 kg ha⁻¹. The soil at the trial site comprises a silty rendzina, with a significant proportion of chalk fragments (UKSO, 2016). Soil mineral N, quantified shortly before sowing, was 29.9 kg N Ha⁻¹, of which 28 kg was nitrate-N and 1.9 kg ammonium-N. The field site on which the trial was based is a commercial arable farm, with barley (*Hordeum vulgare* L.), oilseed rape (*Brassica napus* L.) and wheat (*Triticum aestivum* L.) grown in a rotation.

The ADAS trial used plots measuring 12 m x 1.5 m, clustered in groups of 6 by N application rate, with each variety represented once per cluster. Each N application rate was applied to 3 replicate clusters, of 6 varieties, meaning 18 clusters in total, with a combined area of 1944 m². Experimental clusters of N application rates were separated to each side by buffer zones 6 m wide, and at each end by buffer zones 3 m long (Fig.1). Owing to the logistical challenges of sampling the entire trial, the experimental work presented here is gathered from two of the N application rates (60 kg ha⁻¹ (N rate 2 in Fig. 1), and 280 kg ha⁻¹ (N rate 5 in Fig. 1)), and two of the barley cultivars: KWS Meridian (KWS UK Ltd, Thriplow, Hertfordshire, UK), a 6-row feedstock barley; and Maris Otter (Robin Appel, Waltham Chase, Hampshire, UK), a 2-row malting barley, giving 4 treatment groups, with 3 replicate plots per treatment. Meridian and Maris Otter were chosen from the panel of 6 cultivars available in the trial as they represent contrasting ages of barley varieties, developed in the 1960s and 2000s respectively. Further, Maris Otter is a malting barley, characterised by a low grain protein content, while Meridian was developed as a feedstock barley, with a higher grain protein (and therefore N) content. Experimental sampling and isotope labelling were carried out during the post-anthesis, grain filling period - approximate growth stages 70-80 (Zadoks, 1985).

Intraradical and extraradical AMF quantification

AMF colonisation of both barley varieties was confirmed and then quantified by staining of roots collected from the trial plots. Roots were collected from between 5 and 15 cm below the surface. After clearing in 10 % (w/v) KOH for 20 minutes at 70 °C, roots were rinsed in de-ionised water, acidified in 1 % (v/v) HCl at 25 °C for 10 minutes and then stained in Trypan Blue at 25 °C for 20 minutes. Roots were then rinsed again in de-ionised water before being left in a 50 % (v/v) glycerol solution for 24 hours, before being mounted onto microscope slides to allow quantification of root length colonisation (RLC) using the gridline intersect method (McGonigle et al., 1990).

Soil samples were collected from between 5 and 15 cm below the soil surface. As AMF hyphal turnover can be rapid, (Staddon et al., 2003), hyphal extraction took place within 6 hours of collection to minimise loss due to decomposition. Extraradical hyphal quantity in the plots was determined using an adapted method from Staddon et al. (1999). Briefly, samples of known mass (5-10 g) were suspended in 500 mL of de-ionised water and agitated with a magnetic stirrer plate in order to free the hyphae from soil particles. From this, 200 mL was decanted to a smaller beaker on a magnetic stirrer. Aliquots (10 mL) were removed and

vacuum filtered through 0.45 µm nylon mesh (Anachem, Bedfordshire, UK) and hyphal length density (HLD) was quantified using the gridline intersect method (Hodge, 2001).

¹⁵N stable isotope labelling

The AMF contribution to barley N uptake was investigated by adding a solution of ¹⁵N (as either (¹⁵NH₄)₂SO₄ or K¹⁵NO₃), into mesh-walled cores, into which AMF hyphae could access but plant roots could not, or (as controls for diffusion and mass flow of the added N) cores into which neither AMF hyphae or roots could access. Isotopic ¹⁵N was added in the form of Long Ashton nutrient solution (LAS) (Smith et al., 1983), which can be prepared variously to provide ¹⁵N as ¹⁵NH₄⁺ or ¹⁵NO₃⁻ in equimolar concentrations. The LAS was made to the standard protocols except N being 300% the original concentrations. Each core received 5 mL of LAS, containing 0.683 mg ¹⁵N. (Long Ashton nutrient solution protocol is included in Supplementary Information document 1)

Hyphal access cores were constructed following an adapted method from Johnson et al. (2001). Lengths of PVC tubing (length 85 mm, internal diameter 13 mm, external diameter 16 mm; internal volume 9.9 cm³) with 2 windows cut in the sides of the lower 2/3 of the tube so that 50 % of the side area was open, were wrapped in a 20 µm nylon mesh (John Stanier and Co., Whitefield, Manchester, UK), fixed with Tensol adhesive cement (Bostik Inc., Wauwatosa, Wisconsin, USA). The open bottom end of each tube was covered with the same size mesh. Control cores, which allowed diffusion and mass flow of solutes but prevent hyphal ingrowth, were covered with 0.45 µm nitrocellulose membrane mesh to prevent root and hyphal ingrowth. Cores were filled with a 1/1 (v/v) mixture of silica sand and TerraGreen® (calcinated attapulgite clay, Oil-Dri, Cambridgeshire, UK), which had been sterilised by autoclaving (121 °C for 44 minutes), providing a uniform substrate into which the ¹⁵N solutions could be added.

Each of these cores was then placed inside another, slightly larger core, constructed in the same manner (length 75 mm internal diameter 18, external diameter 21). These cores were also covered in a 20 µm nylon mesh. Such a 'core in a core' design allows the placement of zones of defined and uniform size into the soil, to which ¹⁵N label solutions could be added. A small (approx. 1 mm) air gap is made between the external mesh wall of one core and the internal mesh wall of the other, which should reduce the rapid diffusion of N from the site of addition, which has been a problem in studies where ¹⁵N has been added (Smith and Smith, 2011b). Diffusion and mass flow are unlikely to be prevented entirely, as the pressure of soil on the sides of the core may push the mesh together so that the two layers of mesh make contact. However, the system provides a more stable labelling zone than using a single core, where one mesh layer may be easily damaged (Johnson et al., 2001).

Each of the 12 experimental plots received four cores (1. No AMF Access + ¹⁵NH₄⁺; 2. AMF Access + ¹⁵NH₄⁺; 3. No AMF Access + ¹⁵NO₃⁻; 4. AMF Access + ¹⁵NO₃⁻), spaced 3 m apart to avoid contamination of ¹⁵N from neighbouring cores (Fig. 2). Placement of cores took place 8 weeks before label addition, to allow hyphal ingrowth from the bulk soil. A piece of tape was placed over the top of cores to minimise contamination. This tape was removed for ¹⁵N addition and then replaced.

Sample collection and preparation

After 7 days, the nearest plant to each labelling core was cut at ground level and removed, dried at 70 °C for 48 hours and homogenised in a kitchen blender (Morphy Richards, Mexborough, South Yorkshire, UK) then in a ball mill (MM400 Ball Mill, Retsch GmbH, Haan, Germany). Homogenised shoot samples of known mass (3 mg ± 0.5 mg) were used to quantify ¹⁵N and N content, performed by isotope ratio mass spectrometry (IRMS) (PDZ 2020, Sercon Ltd, Crewe, UK).

Statistical analysis

For all data, statistical analysis was performed using the “R 3.1.0” statistical package, through the “RStudio” integrated development environment (R foundation for Statistical Computing, Vienna, Austria). Data were tested for normality using Shapiro-Wilk and Kolmogorov-Smirnov tests, and Levene’s test was used to confirm homogeneity of variance. Where these tests suggested data did not match test assumptions, data were square-root or log-transformed prior to analysis. Data for root length colonisation, hyphal length density, barley N concentration and biomass were tested by two-way ANOVA, using N addition rate and barley variety as explanatory variables. As two additional explanatory variables were added in the trial for ^{15}N uptake (N addition type, ammonium / nitrate; AM treatment, access / no access), and the small number of replicates in the ADAS field trial, it was not possible to test these factors and the N addition rate and barley cultivar at once. As such, data were split into barley cultivar and N application rate for the ^{15}N data and tested by two-way ANOVA. Here, ^{15}N enrichment was the response variable, while N type and AMF access treatment were the explanatory variables.

Results

Shoot acquisition of ^{15}N added to mesh cores was significantly improved by allowing AMF access into cores, but only when added as $^{15}\text{NO}_3^-$, and only in the High-N plots of Meridian barley (Fig. 3). T-tests indicate that only in High-N Meridian plots receiving $^{15}\text{NO}_3^-$ were ^{15}N enrichment levels greater in AM access treatment than in no access controls ($T_2 = 4.48$, $p = 0.023$) (Supplementary Information, Figure 1). Two-way ANOVA showed that in High-N Meridian, there was a significant effect of N source ($F_{1,8} = 12.73$, $p = 0.007$) and AMF access to cores ($F_{1,8} = 27.86$, $p = 0.007$). There was also a significant interaction between N source and AMF access ($F_{1,8} = 14.25$, $p = 0.005$) (Fig. 3). In High-N Meridian with AMF access, the harvested plants, i.e. those individuals closest to the core to which the isotope label was added, acquired on average 1.62% of the ^{15}N supplied. Other treatment groups saw no greater plant uptake of ^{15}N where AMF could access the isotope label than in no-access controls. Excepting High-N Meridian plots, mean shoot ^{15}N content did not differ among treatments and controls, indicating similar plant acquisition of N following diffusion/mass flow out and into the soil, but minimal fungal-mediated uptake.

All plant roots studied were found to be colonised by AMF, indicating a substantial inoculum potential of the soil at the trial site, although no differences were found between cultivar or N-rate treatments ($p > 0.05$). Mean colonisation was 33.7 % (± 3.52 % SEM) across all treatment groups (Fig. 4). Extraradical mycelium (ERM) hyphal densities, measured in the zones to which ^{15}N was added, were not different among treatment groups ($p > 0.05$). Mean ERM hyphal density across all treatments was 2.49 m g $^{-1}$ DW soil (± 0.31 m g $^{-1}$ SEM). In both cultivars, High-N plots supported ~ 60 % higher shoot N content than Low-N plots ($F_{1,8} = 74.55$, $p < 0.001$), and shoot N concentration was significantly higher in High-N than Low-N plots ($F_{1,8} = 84.28$, $p < 0.001$). Mean shoot N concentration was 9.30 mg g $^{-1}$ DW in Low-N blocks of Maris Otter, and 14.75 mg g $^{-1}$ DW in the High-N. Meridian showed a very similar trend, as N concentration increased from 9.57 mg g $^{-1}$ DW in Low-N plots to 14.38 mg g $^{-1}$ DW in the High N. Shoot N concentration and content did not differ between the two cultivars tested. Shoot DW did not differ between the varieties or the N addition rates.

Discussion

The enrichment of ^{15}N in barley shoots suggests a role for AM-facilitated N acquisition by crop plants, a phenomenon not previously observed in a field setting. Moreover, our data suggest this route of N uptake is dependent upon barley cultivar identity, the N form added and the rate at which N has previously been applied to the plots. AMF have been shown to

transfer substantial quantities of N to plants in root organ culture experiments (Jin et al., 2005) although caution must be exercised before extrapolating these values to crop plant systems as they are far-removed from realistic mycorrhizal physiology. Whole-plant microcosm studies conducted under greenhouse conditions have given mixed results as to whether AMs may contribute to plant N nutrition (Hodge and Storer, 2015). Our data provide the first suggestion that AMF may have a role in cereal crop N uptake in the field. Our data also suggest that short-term changes in N fertilisation regimes can elicit shifts in AM functioning.

While our data suggest a preference for AMF to transfer N to plants when provided to this system as NO_3^- rather than NH_4^+ , previous experimental evidence as to inorganic N source preference by AMF is equivocal (Johansen et al., 1993, Hawkins and George, 2001). Higher uptake of NO_3^- than NH_4^+ is contrary to models which suggest NH_4^+ acquisition should be less energetically expensive (Govindarajulu et al., 2005). Hyphal NH_4^+ uptake may be retarded by problems of charge balancing that are perhaps not encountered when N is acquired as NO_3^- . Simultaneous uptake of NO_3^- and cations such as K^+ , Ca^{2+} or Mg^{2+} from the soil may avoid changes in electrochemical potential across exchange surfaces, allowing N acquisition. Meanwhile, NH_4^+ uptake would require proton secretion (or anion uptake), which may shift soil pH making further NH_4^+ uptake more difficult. Nitrate-N comprised over 90 % of the available N in the soil before the trial was planted, a trend which is not unusual, as NO_3^- often dominates inorganic N pools in arable soils (Marschner, 2011). These relative abundances of N sources may have led to AMF hyphal physiology being acclimated to nitrate uptake (Garraway and Evans, 1984), meaning suddenly-available NH_4^+ could not be acquired effectively. Although the movement and cycling of nitrate and ammonium are known to be influenced by soil moisture (Homyak et al., 2017), precipitation data for the site (Supplementary Information, Table 1) indicates no extraordinary rainfall in the weeks over which the experiment took place, suggesting this was of minor importance here.

While recovery of only 1.6% of the ^{15}N label seems low, total ^{15}N recovery is likely to have been greater than the data suggests. Our data are derived from the aboveground tissue of one plant proximal to the mesh-walled core into which isotopes were added, and it is probable that the roots of numerous plants would have been in close proximity to the core. As such, further ^{15}N is likely to have been acquired by multiple plants. Furthermore, greater ^{15}N uptake into plant shoots may have been recorded if the shoot tissue samples had been taken longer after ^{15}N addition to the mesh-walled cores.

Mesh-walled exclusion cores have been used to quantify AMF-plant nutrient dynamics in a number of studies (Johnson et al., 2001, Field et al., 2012, Field et al., 2016), and are of particular utility where the establishment of truly non-mycorrhizal control plants is not feasible, as in this study. The use of a 0.45 μm nitrocellulose membrane to exclude AMF in-growth to soil compartments is a well-established methodology in the literature (Hodge et al., 2001, Leigh et al., 2009, Thirkell et al., 2016, Storer et al., 2018), although some concerns arise in relation to the effects of such small pore sizes on solute movement, although in the case of studies investigating mycorrhizal P uptake, such effects have been determined to be insignificant (see Zhang et al., 2016, Svenningsen et al., 2018). Our data show increased plant ^{15}N uptake in plots only where N was supplied as nitrate, to Meridian barley, and in plots which had received high rates of N fertiliser (Figure 3). Were the movement of N through these systems determined by the porosity of the membranes used in 'no access' treatments, we might expect ^{15}N enrichment in all plots which received $^{15}\text{NO}_3^-$, which is not the case. Alternative control treatments to disentangle the effects of AMF on plant nutrition might be tested further in future studies to determine the relative merits of each method. Non-mycorrhiza-forming mutants of a number of cereals have been developed (Paszkowski et al., 2006, Watts-Williams and Cavagnaro, 2015) but to date no mycorrhiza-defective

barley mutants are available against which data from hyphal exclusion experiments can be compared. Furthermore, an AMF-colonised plant is morphologically (Gutjahr et al., 2009) and physiologically (Luginbuehl and Oldroyd, 2017) distinct from one which remains uncolonised, and comparisons between AM and mycorrhiza-defective mutants may erroneously conflate these differences and ascribe all contrasts to the lack of mycorrhizas. Combinations of experimental approaches may be employed here to improve the rigour of field experimentation, although the logistics of such trials may prove represent a significant challenge.

Identifying the mechanisms responsible for differential nitrogen transfer from fungus to plant are beyond the scope of this study, but a number of possibilities may be considered. Numerous studies have demonstrated shifts in AMF community composition or structure following N fertilisation, in grassland (Egerton-Warburton and Allen, 2000, Egerton-Warburton et al., 2007, Antoninka et al., 2011, Jiang et al., 2018) and arable systems (Verbruggen et al., 2010, Avio et al., 2013, Liu et al., 2014, Williams et al., 2017). As AMF isolates are known to be functionally different (Avio et al., 2006, Mensah et al., 2015) any N-driven shifts in AMF community have the potential to influence the N cycling in the system. Future experimental testing of the AMF community composition within cereal roots, combined with isotopic tracer studies may elucidate any link between the structure and function of AMF communities in agronomic systems.

Conclusions

Our data show that AMF transfer of N to plant hosts is influenced by agricultural management decisions, here the cultivar of barley and the rate at which inorganic N fertiliser is supplied. The extent to which symbiotic soil microbes might enhance total nutrient uptake in the field remains to be tested; despite demonstrating a mechanism by which plants acquire N, our data cannot indicate whether non-AMF plants in the same field conditions might show enhanced nutrition. Further experimental investigation is required for a wider perspective on the influence of these fungi on their crop plant hosts, and therefore their importance in agricultural systems.

Acknowledgements

This work was supported by a BBSRC White Rose DTP grant: BB/J014443/1. We thank Kate Storer and Pete Berry from ADAS for permission to use their field trial, and for discussion about the experiment. We thank Andrew Manfield, on whose land the trial was sited. We thank Heather Walker for help with the IRMS. This manuscript is an adapted and revised version of a thesis chapter submitted by Tom Thirkell for the degree of Doctor of Philosophy at the University of York, UK, in 2017. Experimental data is presented in the supplementary material.

Author contributions

TT, DC and AH designed the study. TT carried out experimental work, data analysis and wrote the initial draft of the manuscript. All authors contributed to revisions of the manuscript, and read and approved the final submitted version.

Declaration of interest

The authors declare that the submitted work was carried out in the absence of any personal, professional or financial conflict of interest.

369

370

Figure 1. ADAS experiment established at Sancton, East Riding of Yorkshire, UK. Six barley (*Hordeum vulgare* L.) cultivars were planted at the trial site, and received one of 6 N addition rates, ranging from 0 to 300 kg ha⁻¹. Each combination of barley cultivar and N rate was replicated 3 times. Each plot has 3 numbers, denoting: plot identity, N addition rate and barley cultivar, reading top to bottom. Nitrogen addition rate '2' represents 60 kg ha⁻¹ and '5' is 280 kg ha⁻¹. Plot colours also represent N addition rate. Meridian barley is denoted by '4' and Maris Otter by '5'. Asterisks (*) represent plots from which root samples were taken for analysis of root length colonisation and to which ¹⁵N tracer was added. Reproduced with permission by Kate Storer, ADAS.

Figure 2. Diagram of ¹⁵N addition experiment. PVC cores were inserted adjacent to barley (*Hordeum vulgare* L.) plants, four cores per plot, spaced 3 m apart. Cores were organised as follows A1 – AMF Access + Ammonium (NH₄⁺); A2 - No AMF Access + Nitrate (NO₃⁻); A3 - AMF Access + NO₃⁻; A4 - No AMF Access + NH₄⁺. Each core received 0.683 mg ¹⁵N added as Long Ashtons nutrient solution. Plant shoots closest to the core (B1-4) were removed, dried and homogenised for N analysis. Blue boxes (B1-4) represent shoot samples taken.

Figure 3. Excess ¹⁵N content in Maris Otter and Meridian shoots (calculated by subtracting shoot ¹⁵N content in each 'Access' unit from the mean of the corresponding values in the 'No Access' units). Shoot ¹⁵N enrichment was significantly higher than 'no access' controls when supplied as nitrate to Meridian barley in High-N plots. Circles represent individual data points, boxplot centre bars represent the median values. High-N + ammonium groups are represented by white bars, High-N + nitrate by light blue bars, Low-N + ammonium by dark grey bars and Low-N + nitrate by dark blue bars. Data shown are means ± SEM, n = 3. Bars sharing the same letter are not significantly different.

Figure 4. Percentage root length colonisation, as determined by Trypan Blue staining, was not significantly different between treatments. All inspected plants were colonised by arbuscular mycorrhizal fungi (AMF), confirmed by presence of characteristic structures, arbuscules and vesicles. Mean colonisation ranged from 28.5 % in Maris Otter in Low N, to 38.0 % in Meridian Low-N, but no groups were significantly different. Circles represent individual data points. High-N groups are denoted by green bars, Low-N groups are denoted by yellow bars. "N.S.D." indicates that there were no significant differences among treatment means. Data shown are means ± SEM, n = 3.

References

- Agren, G.I., Wetterstedt, J.a.M. , Billberger, M.F.K. (2012). Nutrient limitation on terrestrial plant growth - modeling the interaction between nitrogen and phosphorus. *New Phytologist*, 194, 953-960.10.1111/j.1469-8137.2012.04116.x.
- Ames, R.N., Reid, C.P.P., Porter, L.K. , Cambardella, C. (1983). Hyphal uptake and transport of nitrogen from 2 ¹⁵N-labeled sources by *Glomus mosseae*, a vesicular arbuscular mycorrhizal fungus. *New Phytologist*, 95, 381-396.10.1111/j.1469-8137.1983.tb03506.x.
- Antoninka, A., Reich, P.B. , Johnson, N.C. (2011). Seven years of carbon dioxide enrichment, nitrogen fertilization and plant diversity influence arbuscular mycorrhizal fungi in a grassland ecosystem. *New Phytologist*, 192, 200-214.10.1111/j.1469-8137.2011.03776.x.

414 Avio, L., Castaldini, M., Fabiani, A., Bedini, S., Sbrana, C., Turrini, A. , Giovannetti, M.
415 (2013). Impact of nitrogen fertilization and soil tillage on arbuscular mycorrhizal
416 fungal communities in a mediterranean agroecosystem. *Soil Biology & Biochemistry*,
417 67, 285-294.10.1016/j.soilbio.2013.09.005.

418 Avio, L., Pellegrino, E., Bonari, E. , Giovannetti, M. (2006). Functional diversity of arbuscular
419 mycorrhizal fungal isolates in relation to extraradical mycelial networks. *New*
420 *Phytologist*, 172, 347-357.10.1111/j.1469-8137.2006.01839.x.

421 Barrett, G., Campbell, C.D. , Hodge, A. (2014). The direct response of the external mycelium
422 of arbuscular mycorrhizal fungi to temperature and the implications for nutrient
423 transfer. *Soil Biology & Biochemistry*, 78, 109-117

424 Belmondo, S., Fiorilli, V., Perez-Tienda, J., Ferrol, N., Marmeisse, R. , Lanfranco, L. (2014).
425 A dipeptide transporter from the arbuscular mycorrhizal fungus *Rhizophagus*
426 *irregularis* is upregulated in the intraradical phase. *Frontiers in Plant Science*,
427 5.10.3389/fpls.2014.00436.

428 Brackin, R., Atkinson, B.S., Sturrock, C.J. , Rasmussen, A. (2017). Roots-eye view: Using
429 microdialysis and microct to non-destructively map root nutrient depletion and
430 accumulation zones. *Plant Cell and Environment*, 40, 3135-3142.10.1111/pce.13072.

431 Breuninger, M., Trujillo, C.G., Serrano, E., Fischer, R. , Requena, N. (2004). Different
432 nitrogen sources modulate activity but not expression of glutamine sythetase in
433 arbuscular mycorrhizal fungi. *Fungal Genetics and Biology*, 542-552

434 Bucking, H. , Kafle, A. (2015). Role of arbuscular mycorrhizal fungi in the nitrogen uptake of
435 plants: Current knowledge and research gaps. *Agronomy-Basel*, 5, 587-
436 612.10.3390/agronomy5040587.

437 Cameron, K.C., Di, H.J. , Moir, J.L. (2013). Nitrogen losses from the soil/plant system: A
438 review. *Annals of Applied Biology*, 162, 145-173.10.1111/aab.12014.

439 Cavagnaro, T.R., Bender, S.F., Asghari, H.R. , Van Der Heijden, M.G.A. (2015). The role of
440 arbuscular mycorrhizas in reducing soil nutrient loss. *Trends in Plant Science*, 20,
441 283-290.10.1016/j.tplants.2015.03.004.

442 Courty, P.E., Smith, P., Koegel, S., Redecker, D. , Wipf, D. (2015). Inorganic nitrogen uptake
443 and transport in beneficial plant root-microbe interactions. *Critical Reviews in Plant*
444 *Sciences*, 34, 4-16.10.1080/07352689.2014.897897.

445 Cui, M. , Caldwell, M.M. (1996a). Facilitation of plant phosphate acquisition by arbuscular
446 mycorrhizas from enriched soil patches .1. Roots and hyphae exploiting the same
447 soil volume. *New Phytologist*, 133, 453-460.10.1111/j.1469-8137.1996.tb01912.x.

448 Cui, M.Y. , Caldwell, M.M. (1996b). Facilitation of plant phosphate acquisition by arbuscular
449 mycorrhizas from enriched soil patches .2. Hyphae exploiting root-free soil. *New*
450 *Phytologist*, 133, 461-467.10.1111/j.1469-8137.1996.tb01913.x.

451 Davison, J., Moora, M., Opik, M., Adholeya, A., Ainsaar, L., Ba, A., Burla, S., Diedhiou, A.G.,
 452 Hiiesalu, I., Jairus, T., Johnson, N.C., Kane, A., Koorem, K., Kochar, M., Ndiaye, C.,
 453 Partel, M., Reier, U., Saks, U., Singh, R., Vasar, M., Zobel, M. (2015). Global
 454 assessment of arbuscular mycorrhizal fungus diversity reveals very low endemism.
 455 *Science*, 349, 970-973.10.1126/science.aab1161.

456 Egerton-Warburton, L.M., Allen, E.B. (2000). Shifts in arbuscular mycorrhizal communities
 457 along an anthropogenic nitrogen deposition gradient. *Ecological Applications*, 10,
 458 484-496.10.1890/1051-0761(2000)010[0484:siamca]2.0.co;2.

459 Egerton-Warburton, L.M., Johnson, N.C., Allen, E.B. (2007). Mycorrhizal community
 460 dynamics following nitrogen fertilization: A cross-site test in five grasslands.
 461 *Ecological Monographs*, 77, 527-544.10.1890/06-1772.1.

462 Ezawa, T., Saito, K. (2018). How do arbuscular mycorrhizal fungi handle phosphate? New
 463 insight into fine-tuning of phosphate metabolism. *New Phytologist*, 220, 1116-
 464 1121.10.1111/nph.15187.

465 Field, K.J., Cameron, D.D., Leake, J.R., Tille, S., Bidartondo, M.I., Beerling, D.J. (2012).
 466 Contrasting arbuscular mycorrhizal responses of vascular and non-vascular plants to
 467 a simulated palaeozoic CO₂ decline. *Nature Communications*,
 468 3.10.1038/ncomms1831.

469 Field, K.J., Rimington, W.R., Bidartondo, M.I., Allinson, K.E., Beerling, D.J., Cameron, D.D.,
 470 Duckett, J.G., Leake, J.R., Pressel, S. (2016). Functional analysis of liverworts in
 471 dual symbiosis with Glomeromycota and Mucoromycotina fungi under a simulated
 472 palaeozoic CO₂ decline. *ISME Journal*, 10, 1514-1526.10.1038/ismej.2015.204.

473 Garcia, K., Doidy, J., Zimmermann, S.D., Wipf, D., Courty, P.E. (2016). Take a trip through
 474 the plant and fungal transportome of mycorrhiza. *Trends in Plant Science*, 21, 937-
 475 950.10.1016/j.tplants.2016.07.010.

476 Garraway, M.O., Evans, R.C. 1984. *Fungal nutrition and physiology*, New York, USA, John
 477 Wiley & Sons.

478 Gosling, P., Hodge, A., Goodlass, G., Bending, G.D. (2006). Arbuscular mycorrhizal fungi
 479 and organic farming. *Agriculture Ecosystems & Environment*, 113, 17-
 480 35.10.1016/j.agee.2005.09.009.

481 Govindarajulu, M., Pfeffer, P.E., Jin, H.R., Abubaker, J., Douds, D.D., Allen, J.W., Bucking,
 482 H., Lammers, P.J., Shachar-Hill, Y. (2005). Nitrogen transfer in the arbuscular
 483 mycorrhizal symbiosis. *Nature*, 435, 819-823.10.1038/nature03610.

484 Gutjahr, C., Casieri, L., Paszkowski, U. (2009). *Glomus intraradices* induces changes in root
 485 system architecture of rice independently of common symbiosis signaling. *New*
 486 *Phytologist*, 182, 829-837.10.1111/j.1469-8137.2009.02839.x.

487 Hawkesford, M.J. (2014). Reducing the reliance on nitrogen fertilizer for wheat production.
 488 *Journal of Cereal Science*, 59, 276-283.10.1016/j.jcs.2013.12.001.

489 Hawkins, H.J. , George, E. (2001). Reduced N-15-nitrogen transport through arbuscular
490 mycorrhizal hyphae to *Triticum aestivum* L. Supplied with ammonium vs. nitrate
491 nutrition. *Annals of Botany*, 87, 303-311.10.1006/anbo.2000.1305.

492 Hawkins, H.J., Johansen, A. , George, E. (2000). Uptake and transport of organic and
493 inorganic nitrogen by arbuscular mycorrhizal fungi. *Plant and Soil*, 226, 275-
494 285.10.1023/a:1026500810385.

495 Herman, D.J., Firestone, M.K., Nuccio, E. , Hodge, A. (2012). Interactions between an
496 arbuscular mycorrhizal fungus and a soil microbial community mediating litter
497 decomposition. *FEMS Microbiology Ecology*, 80, 236-247.10.1111/j.1574-
498 6941.2011.01292.x.

499 Hodge, A. (2001). Arbuscular mycorrhizal fungi influence decomposition of, but not plant
500 nutrient capture from, glycine patches in soil. *New Phytologist*, 151, 725-
501 734.10.1046/j.0028-646x.2001.00200.x.

502 Hodge, 2014 (2014). Interactions between arbuscular mycorrhizal fungi and organic material
503 substrates. *Advances in Applied Microbiology* 89, 47-99. doi:10.1016/b978-0-12-800259-
504 9.00002-0.

505 Hodge, A. (2017). "Accessibility of inorganic and organic nutrients for mycorrhizas," in
506 Mycorrhizal mediation of soil fertility, structure and carbon storage, ed N. Johnson, C.
507 Gehring, J. Jansa (Amsterdam, Netherlands: Elsevier, 129-148.

508 Hodge, A., Campbell, C.D. , Fitter, A.H. (2001). An arbuscular mycorrhizal fungus
509 accelerates decomposition and acquires nitrogen directly from organic material.
510 *Nature*, 413, 297-299.10.1038/35095041.

511 Hodge, A. , Fitter, A.H. (2010). Substantial nitrogen acquisition by arbuscular mycorrhizal
512 fungi from organic material has implications for N cycling. *Proceedings of the*
513 *National Academy of Sciences of the United States of America*, 107, 13754-
514 13759.10.1073/pnas.1005874107.

515 Hodge, A., Helgason, T. , Fitter, A.H. (2010). Nutritional ecology of arbuscular mycorrhizal
516 fungi. *Fungal Ecology*, 3, 267-273.10.1016/j.funeco.2010.02.002.

517 Hodge, A. , Storer, K. (2015). Arbuscular mycorrhiza and nitrogen: Implications for individual
518 plants through to ecosystems. *Plant and Soil*, 386, 1-19.10.1007/s11104-014-2162-1.

519 Homyak, P.M., Allison, S.D., Huxman, T.E., Goulden, M.L. , Treseder, K.K. (2017). Effects of
520 drought manipulation on soil nitrogen cycling: A meta-analysis. *Journal of*
521 *Geophysical Research-Biogeosciences*, 122, 3260-3272.10.1002/2017jg004146.

522 Jiang, S.J., Liu, Y.J., Luo, J.J., Qin, M.S., Johnson, N.C., Opik, M., Vasar, M., Chai, Y.X.,
523 Zhou, X.L., Mao, L., Du, G.Z., An, L.Z. , Feng, H.Y. (2018). Dynamics of arbuscular
524 mycorrhizal fungal community structure and functioning along a nitrogen enrichment
525 gradient in an alpine meadow ecosystem. *New Phytologist*, 220, 1222-
526 1235.10.1111/nph.15112.

527 Jin, H., Pfeffer, P.E., Douds, D.D., Piotrowski, E., Lammers, P.J. , Shachar-Hill, Y. (2005).
528 The uptake, metabolism, transport and transfer of nitrogen in an arbuscular
529 mycorrhizal symbiosis. *New Phytologist*, 168, 687-696.10.1111/j.1469-
530 8137.2005.01536.x.

531 Johansen, A., Jakobsen, I. , Jensen, E.S. (1993). Hyphal transport by a vesicular-arbuscular
532 mycorrhizal fungus of N applied to the soil as ammonium or nitrate. *Biology and*
533 *Fertility of Soils*, 16, 66-70.10.1007/bf00336518.

534 Johnson, D., Leake, J.R. , Read, D.J. (2001). Novel in-growth core system enables
535 functional studies of grassland mycorrhizal mycelial networks. *New Phytologist*, 152,
536 555-562.10.1046/j.0028-646X.2001.00273.x.

537 Johnson, N.C. (2010). Resource stoichiometry elucidates the structure and function of
538 arbuscular mycorrhizas across scales. *New Phytologist*, 185, 631-
539 647.10.1111/j.1469-8137.2009.03110.x.

540 Johnson, N.C., Wilson, G.W.T., Wilson, J.A., Miller, R.M. , Bowker, M.A. (2015). Mycorrhizal
541 phenotypes and the law of the minimum. *New Phytologist*, 205, 1473-
542 1484.10.1111/nph.13172.

543 Kahkola, A.K., Nygren, P., Leblanc, H.A., Pennanen, T. , Pietikainen, J. (2012). Leaf and
544 root litter of a legume tree as nitrogen sources for cacaos with different root
545 colonisation by arbuscular mycorrhizae. *Nutrient Cycling in Agroecosystems*, 92, 51-
546 65.10.1007/s10705-011-9471-z.

547 Karasawa, T., Hodge, A. , Fitter, A.H. (2012). Growth, respiration and nutrient acquisition by
548 the arbuscular mycorrhizal fungus *glomus mosseae* and its host plant *Plantago*
549 *lanceolata* in cooled soil. *Plant Cell and Environment*, 35, 819-828.10.1111/j.1365-
550 3040.2011.02455.x.

551 Ladha, J.K., Pathak, H., Krupnik, T.J., Six, J. , Van Kessel, C. (2005). Efficiency of fertilizer
552 nitrogen in cereal production: Retrospects and prospects. *Advances in Agronomy*,
553 Vol 87, 87, 85-156.10.1016/s0065-2113(05)87003-8.

554 Ladha, J.K., Tirol-Padre, A., Reddy, C.K., Cassman, K.G., Verma, S., Powlson, D.S., Van
555 Kessel, C., Richter, D.D., Chakraborty, D. , Pathak, H. (2016). Global nitrogen
556 budgets in cereals: A 50-year assessment for maize, rice, and wheat production
557 systems. *Scientific Reports*, 6.10.1038/srep19355.

558 Leigh, J., Hodge, A. , Fitter, A.H. (2009). Arbuscular mycorrhizal fungi can transfer
559 substantial amounts of nitrogen to their host plant from organic material. *New*
560 *Phytologist*, 181, 199-207.10.1111/j.1469-8137.2008.02630.x.

561 Lekberg, Y. , Helgason, T. (2018). In situ mycorrhizal function - knowledge gaps and future
562 directions. *New Phytologist*, 220, 957-962.10.1111/nph.15064.

563 Liu, W., Jiang, S.S., Zhang, Y.L., Yue, S.C., Christie, P., Murray, P.J., Li, X.L. , Zhang, J.L.
564 (2014). Spatiotemporal changes in arbuscular mycorrhizal fungal communities under
565 different nitrogen inputs over a 5-year period in intensive agricultural ecosystems on

566 the north china plain. *FEMS Microbiology Ecology*, 90, 436-453.10.1111/1574-
567 6941.12405.

568 Luginbuehl, L.H. , Oldroyd, G.E.D. (2017). Understanding the arbuscule at the heart of
569 endomycorrhizal symbioses in plants. *Current Biology*, 27, R952-
570 R963.10.1016/j.cub.2017.06.042.

571 Marschner, H. 2011. *Mineral nutrition of higher plants*, London, Academic Press.

572 Masclaux-Daubresse, C., Daniel-Vedele, F., Dechorgnat, J., Chardon, F., Gaufichon, L. ,
573 Suzuki, A. (2010). Nitrogen uptake, assimilation and remobilization in plants:
574 Challenges for sustainable and productive agriculture. *Annals of Botany*, 105, 1141-
575 1157.10.1093/aob/mcq028.

576 McGonigle, T.P., Miller, M.H., Evans, D.G., Fairchild, G.L. , Swan, J.A. (1990). A new
577 method which gives an objective-measure of colonization of roots by vesicular
578 arbuscular mycorrhizal fungi. *New Phytologist*, 115, 495-501.10.1111/j.1469-
579 8137.1990.tb00476.x.

580 Mensah, J.A., Koch, A.M., Antunes, P.M., Kiers, E.T., Hart, M. , Bucking, H. (2015). High
581 functional diversity within species of arbuscular mycorrhizal fungi is associated with
582 differences in phosphate and nitrogen uptake and fungal phosphate metabolism.
583 *Mycorrhiza*, 25, 533-546.10.1007/s00572-015-0631-x.

584 Paszkowski, U., Jakovleva, L. , Boller, T. (2006). Maize mutants affected at distinct stages of
585 the arbuscular mycorrhizal symbiosis. *Plant Journal*, 47, 165-173.10.1111/j.1365-
586 313X.2006.02785.x.

587 Pretty, J. (2008). Agricultural sustainability: Concepts, principles and evidence. *Philosophical*
588 *Transactions of the Royal Society B-Biological Sciences*, 363, 447-
589 465.10.1098/rstb.2007.2163.

590 Pretty, J. (2018). Intensification for redesigned and sustainable agricultural systems.
591 *Science*, 362, 908-+.10.1126/science.aav0294.

592 Reynolds, H.L., Hartley, A.E., Vogelsang, K.M., Bever, J.D. , Schultz, P.A. (2005).
593 Arbuscular mycorrhizal fungi do not enhance nitrogen acquisition and growth of old-
594 field perennials under low nitrogen supply in glasshouse culture. *New Phytologist*,
595 167, 869-880.10.1111/j.1469-8137.2005.01455.x.

596 Rillig, M.C., Aguilar-Trigueros, C.A., Camenzind, T., Cavagnaro, T.R., Degrune, F.,
597 Hohmann, P., Lammel, D.R., Mansour, I., Roy, J., Van Der Heijden, M.G.A. , Yang,
598 G.W. (2019). Why farmers should manage the arbuscular mycorrhizal symbiosis.
599 *New Phytologist*, 222, 1171-1175.10.1111/nph.15602.

600 Rillig, M.C., Sosa-Hernandez, M.A., Roy, J., Aguilar-Trigueros, C.A., Valyi, K. , Lehmann, A.
601 (2016). Towards an integrated mycorrhizal technology: Harnessing mycorrhiza for
602 sustainable intensification in agriculture. *Frontiers in Plant Science*,
603 7.10.3389/fpls.2016.01625.

604 Ryan, M.H. , Graham, J.H. (2018). Little evidence that farmers should consider abundance
605 or diversity of arbuscular mycorrhizal fungi when managing crops. *New Phytologist*,
606 220, 1092-1107.10.1111/nph.15308.

607 Sanders, F. E. and Tinker, P.B. (1973). Phosphate inflow into mycorrhizal roots. *Pesticide*
608 *Science* 4, 385-395. doi:10.1002/ps.2780040316.

609 Smith, F.A. , Smith, S.E. (2011a). What is the significance of the arbuscular mycorrhizal
610 colonisation of many economically important crop plants? *Plant and Soil*, 348, 63-
611 79.10.1007/s11104-011-0865-0.

612 Smith, G.S., Johnston, C.M. , Cornforth, I.S. (1983). Comparison of nutrient solutions for
613 growth of plants in sand culture. *New Phytologist*, 94, 537-548.10.1111/j.1469-
614 8137.1983.tb04863.x.

615 Smith, S.E. , Read, D.J. 2008. *Mycorrhizal symbiosis*, London, Academic Press.

616 Smith, S.E. , Smith, F.A. (2011b). Roles of arbuscular mycorrhizas in plant nutrition and
617 growth: New paradigms from cellular to ecosystem scales. *Annual Review of Plant*
618 *Biology*, Vol 62, 62, 227-250.10.1146/annurev-arplant-042110-103846.

619 Smith, S.E., Smith, F.A. , Jakobsen, I. (2003). Mycorrhizal fungi can dominate phosphate
620 supply to plants irrespective of growth responses. *Plant Physiology*, 133, 16-
621 20.10.1104/pp.103.024380.

622 Smith, S.E., Smith, F.A. , Jakobsen, I. (2004). Functional diversity in arbuscular mycorrhizal
623 (AM) symbioses: The contribution of the mycorrhizal P uptake pathway is not
624 correlated with mycorrhizal responses in growth or total P uptake. *New Phytologist*,
625 162, 511-524.10.1111/j.1469-8137.2004.01039.x.

626 Sosa-Hernandez, M.A., Roy, J., Hempel, S., Kautz, T., Kopke, U., Uksa, M., Schlöter, M.,
627 Caruso, T. , Rillig, M.C. (2018). Subsoil arbuscular mycorrhizal fungal communities in
628 arable soil differ from those in topsoil. *Soil Biology & Biochemistry*, 117, 83-
629 86.10.1016/j.soilbio.2017.11.009.

630 Staddon, P.L., Fitter, A.H. , Graves, J.D. (1999). Effect of elevated atmospheric co₂ on
631 mycorrhizal colonization, external mycorrhizal hyphal production and phosphorus
632 inflow in *Plantago lanceolata* and *Trifolium repens* in association with the arbuscular
633 mycorrhizal fungus *Glomus mosseae*. *Global Change Biology*, 5, 347-
634 358.10.1046/j.1365-2486.1999.00230.x.

635 Storer, K., Coggan, A., Ineson, P. , Hodge, A. (2018). Arbuscular mycorrhizal fungi reduce
636 nitrous oxide emissions from N₂O hotspots. *New Phytologist*, 220, 1285-
637 1295.10.1111/nph.14931.

638 Svenningsen, N.B., Watts-Williams, S.J., Joner, E.J., Battini, F., Efthymiou, A., Cruz-
639 Paredes, C., Nybroe, O. , Jakobsen, I. (2018). Suppression of the activity of
640 arbuscular mycorrhizal fungi by the soil microbiota. *ISME Journal*, 12, 1296-
641 1307.10.1038/s41396-018-0059-3.

642 Thirkell, T.J., Cameron, D.D. , Hodge, A. (2016). Resolving the 'nitrogen paradox' of
643 arbuscular mycorrhizas: Fertilization with organic matter brings considerable benefits
644 for plant nutrition and growth. *Plant Cell and Environment*, 39, 1683-
645 1690.10.1111/pce.12667.

646 Thirkell, T.J., Charters, M.D., Elliott, A.J., Sait, S.M. , Field, K.J. (2017). Are mycorrhizal
647 fungi our sustainable saviours? Considerations for achieving food security. *Journal of*
648 *Ecology*, 105, 921-929.10.1111/1365-2745.12788.

649 Tinker, P.B. , Nye, P.H. 2000. *Solute movement in the rhizosphere*, Oxford, Oxford
650 University Press.

651 Tisserant, E., Kohler, A., Dozolme-Seddas, P., Balestrini, R., Benabdellah, K., Colard, A.,
652 Croll, D., Da Silva, C., Gomez, S.K., Koul, R., Ferrol, N., Fiorilli, V., Formey, D.,
653 Franken, P., Helber, N., Hijri, M., Lanfranco, L., Lindquist, E., Liu, Y., Malbreil, M.,
654 Morin, E., Poulain, J., Shapiro, H., Van Tuinen, D., Waschke, A., Azcon-Aguilar, C.,
655 Becard, G., Bonfante, P., Harrison, M.J., Kuster, H., Lammers, P., Paszkowski, U.,
656 Requena, N., Rensing, S.A., Roux, C., Sanders, I.R., Shachar-Hill, Y., Tuskan, G.,
657 Young, J.P.W., Gianinazzi-Pearson, V. , Martin, F. (2012). The transcriptome of the
658 arbuscular mycorrhizal fungus *Glomus intraradices* (DAOM 197198) reveals
659 functional tradeoffs in an obligate symbiont. *New Phytologist*, 193, 755-
660 769.10.1111/j.1469-8137.2011.03948.x.

661 Ukso. 2016. *Soilscapes for England and Wales* [Online]. Available:
662 <http://www.ukso.org/SoilsOfEngWales/home.html> [Accessed 20/08 2016].

663 Verbruggen, E., Roling, W.F.M., Gamper, H.A., Kowalchuk, G.A., Verhoef, H.A. , Van Der
664 Heijden, M.G.A. (2010). Positive effects of organic farming on below-ground
665 mutualists: Large-scale comparison of mycorrhizal fungal communities in agricultural
666 soils. *New Phytologist*, 186, 968-979.10.1111/j.1469-8137.2010.03230.x.

667 Watts-Williams, S.J. , Cavagnaro, T.R. (2015). Using mycorrhiza-defective mutant
668 genotypes of non-legume plant species to study the formation and functioning of
669 arbuscular mycorrhiza: A review. *Mycorrhiza*, 25, 587-597.10.1007/s00572-015-
670 0639-2.

671 Whiteside, M.D., Digman, M.A., Gratton, E. , Treseder, K.K. (2012a). Organic nitrogen
672 uptake by arbuscular mycorrhizal fungi in a boreal forest. *Soil Biology &*
673 *Biochemistry*, 55, 7-13.10.1016/j.soilbio.2012.06.001.

674 Whiteside, M.D., Garcia, M.O. , Treseder, K.K. (2012b). Amino acid uptake in arbuscular
675 mycorrhizal plants. *Plos One*, 7.10.1371/journal.pone.0047643.

676 Williams, A., Manoharan, L., Rosenstock, N.P., Olsson, P.A. , Hedlund, K. (2017). Long-term
677 agricultural fertilization alters arbuscular mycorrhizal fungal community composition
678 and barley (*Hordeum vulgare*) mycorrhizal carbon and phosphorus exchange. *New*
679 *Phytologist*, 213, 874-885.10.1111/nph.14196.

680 Zadoks, J.C. (1985). A decimal code for the growth-stages of cereals. *Agriculture Biology &*
681 *Environmental Sciences*, 14-14

682 Zhang, L., Xu, M.G., Liu, Y., Zhang, F.S., Hodge, A. , Feng, G. (2016). Carbon and
683 phosphorus exchange may enable cooperation between an arbuscular mycorrhizal
684 fungus and a phosphate-solubilizing bacterium. *New Phytologist*, 210, 1022-
685 1032.10.1111/nph.13838.