



This is a repository copy of *Infection/inflammation-associated preterm delivery within 14 days of presentation with symptoms of preterm labour : A multivariate predictive model.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/151656/>

Version: Published Version

Article:

Amabebe, E. orcid.org/0000-0002-3924-5270, Reynolds, S. orcid.org/0000-0002-6463-8471, He, X. et al. (3 more authors) (2019) Infection/inflammation-associated preterm delivery within 14 days of presentation with symptoms of preterm labour : A multivariate predictive model. PLOS ONE, 14 (9). e0222455.

<https://doi.org/10.1371/journal.pone.0222455>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>


RESEARCH ARTICLE

Infection/inflammation-associated preterm delivery within 14 days of presentation with symptoms of preterm labour: A multivariate predictive model

Emmanuel Amabebe¹ , Steven Reynolds², Xiaoya He¹, Robyn Wood¹, Victoria Stern¹, Dilly O. C. Anumba^{1*}

1 Academic Unit of Reproductive and Developmental Medicine, University of Sheffield, Sheffield, England, United Kingdom, **2** Academic Unit of Radiology, University of Sheffield, Sheffield, England, United Kingdom

* d.o.c.anumba@sheffield.ac.uk



OPEN ACCESS

Citation: Amabebe E, Reynolds S, He X, Wood R, Stern V, Anumba DOC (2019) Infection/inflammation-associated preterm delivery within 14 days of presentation with symptoms of preterm labour: A multivariate predictive model. PLoS ONE 14(9): e0222455. <https://doi.org/10.1371/journal.pone.0222455>

Editor: Luca Giannella, Azienda Ospedaliero Universitaria Ospedali Riuniti di Ancona Umberto I G M Lancisi G Salesi, ITALY

Received: April 15, 2019

Accepted: August 29, 2019

Published: September 12, 2019

Copyright: © 2019 Amabebe et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Funding: This studies were supported by the Medical Research Council, UK (grant number: MR/J014788/1) to DA. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. <https://mrc.ukri.org>.

Abstract

Multi-marker tests hold promise for identifying symptomatic women at risk of imminent preterm delivery (PTD, <37 week's gestation). This study sought to determine the relationship of inflammatory mediators and metabolites in cervicovaginal fluid (CVF) with spontaneous PTD (sPTD) and delivery within 14 days of presentation with symptoms of preterm labour (PTL). CVF samples from 94 (preterm = 19, term = 75) singleton women with symptoms of PTL studied between 19⁺⁰–36⁺⁶ weeks' gestation were analysed for cytokines/chemokines by multiplexed bead-based immunoassay, while metabolites were quantified by enzyme-based spectrophotometry in a subset of 61 women (preterm = 16, term = 45). Prevalence of targeted vaginal bacterial species was determined for 70 women (preterm = 14, term = 66) by PCR. Overall, 10 women delivered within 14 days of sampling. Predictive capacities of individual biomarkers and cytokine-metabolite combinations for sPTD and delivery within 14 days of sampling were analysed by logistic regression models and area under the receiver operating characteristic curve. *Fusobacterium* sp., *Mobiluncus mulieris* and *Mycoplasma hominis* were detected in more preterm-delivered than term women ($P < 0.0001$), while, *M. curtisii* was found in more term-delivered than preterm women ($P < 0.0001$). RANTES (0.91, 0.65–1.0), IL-6 (0.79, 0.67–0.88), and Acetate/Glutamate ratio (0.74, 0.61–0.85) were associated with delivery within 14 days of sampling (AUC, 95% CI). There were significant correlations between cytokines and metabolites, and several cytokine-metabolite combinations were associated with sPTD or delivery within 14 days of sampling (e.g. L/D-lactate ratio +Acetate/Glutamate ratio+IL-6: 0.84, 0.67–0.94). Symptomatic women destined to deliver preterm and within 14 days of sampling express significantly higher pro-inflammatory mediators at mid to late gestation. In this cohort, IL-6, Acetate/Glutamate ratio and RANTES were associated with delivery within 14 days of sampling, consistent with their roles in modulating infection-inflammation-associated preterm labour in women presenting with symptoms of preterm birth. Replication of these observations in larger cohorts of women could show potential clinical utility.

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: Ace/Glx, Acetate/Glutamate ratio; *a. u.*, arbitrary unit; AUC, Area under the Receiver Operating Characteristics (ROC) curve; BMI, Body mass index; BV, Bacterial vaginosis; CCL5, chemokine (C-C motif) ligand 5; CI, Confidence interval; conc, absolute concentration; CVF, Cervicovaginal fluid; EBS, Enzyme-based spectrophotometry; IFN- γ , interferon gamma; *IL-*, interleukin; NF- κ B, nuclear factor- κ B; *n.i.*, ¹H-NMR-derived normalised integral; NMR, Nuclear Magnetic Resonance Spectroscopy; MAPK, mitogen-activated protein kinase; NPV, negative predictive value; *qfFN*, quantitative Fetal fibronectin; PGE₂, prostaglandin E2; PPRM, Preterm premature rupture of membranes; PPV, positive predictive value; PTD, preterm delivery; PTL, preterm labour; RANTES, regulated on activation, normal T cell expressed and secreted; Sen, sensitivity; sIL-6R α , soluble form of IL-6 receptor α ; Spec, specificity; sPTD, spontaneous preterm delivery; TNF-r1, tumor necrosis factor receptor 1; +LR, positive likelihood ratio; -LR, negative likelihood ratio.

Introduction

Preterm delivery (PTD, before 37 weeks of gestation) remains the dominant global cause of perinatal morbidity and mortality. Preterm labour (PTL) is suspected when a woman presents with frequent uterine contractions occurring at least once every 5–8 minutes, cervical dilation >2 cm and cervical effacement ($\geq 50\%$) before 37 weeks' gestation [1]. Approximately 45% of PTDs are preceded by spontaneous PTL, while ~30% follow preterm premature rupture of membranes (PPROM) [2]. Majority of women presenting with symptoms of PTL do not eventually deliver preterm and would benefit from better prognostication of those most likely to imminently deliver preterm [3]. Accurate identification of these pregnant women facilitates prompt clinical decision-making, maternal treatment with steroids to aid fetal lung maturation, minimises unnecessary hospitalisations, and improves triaging of patients to centres with optimal neonatal care facilities.

The most commonly employed clinical test for predicting imminent PTD are quantitative fetal fibronectin (qfFN) and insulin-like growth factor binding protein-1 (Actim Partus) [4–7], however, due to the heterogeneity of the pathophysiology of PTD, several studies have explored other maternal clinical, inflammation and biochemical markers to predict PTD within 7–14 days in women presenting with symptoms of PTL [3, 8]. Previously we showed that decreased cervicovaginal fluid (CVF) acetate, probably produced by vaginal mixed anaerobes in the absence of *Lactobacillus* dominance, appears a suitable “rule-out” marker for delivery within 2 weeks in symptomatic women [9, 10]. Low CVF glutamate is also associated with reduced vaginal fluid acidity and bacterial vaginosis (BV) [11], an established risk factor of PTD [12–14]. We have also recently demonstrated the predictive value of several CVF pro-inflammatory mediators, either singly or in combination with qfFN for spontaneous PTD (sPTD) in high risk women without symptoms of PTL [15]. Combining multiple potential markers implicated in the pathogenesis of sPTD may provide a more accurate clinical screening approach [7, 16, 17].

Therefore, to determine whether CVF metabolites and cytokines/chemokines assessment enhanced the prediction of sPTD and delivery within 14 days of presentation with symptoms suggestive of PTL, we investigated the relationship between these markers of infection/inflammatory and sPTD and delivery within 14 days of presentation. We hypothesised that symptomatic pregnant women who eventually deliver prematurely, within 14 days of presentation/sampling, will show higher CVF pro-inflammatory mediators and acetate/glutamate ratio compared to those that delivered afterwards. Furthermore, a multivariable approach may provide a more comprehensive analysis of the patho-mechanism of sPTD.

Methods

In this study, CVF samples were obtained from 94 women presenting with symptoms of PTL (19⁺⁰–36⁺⁶ weeks' gestation) at the Labour Ward Assessment Suite of the Jessop Wing Maternity Hospital, Sheffield, UK,—regular uterine contractions (at least once every 10 mins) and cervical dilatation <3 cm, intact fetal membranes and no clinical evidence of genitourinary tract infection, abnormal cervical cytology or vaginal bleeding. CVF samples were obtained prior to any vaginal examination or clinical intervention such as administration of antibiotics, tocolytics, steroids, or any vaginal pessary.

Study participants were enrolled between January 2014 and March 2017 and closely monitored until delivery outcomes were ascertained. Immediately after the swab samples were obtained, quantitative fetal fibronectin (qfFN), vaginal sonographic determination of cervical length (CL) as well as vaginal pH measurement were also performed as previously reported [9, 10, 18, 19].

These studies were approved by the Yorkshire and Humber (Sheffield) Committee of the UK National Research Ethics Service (REC Number 13/YH/0167).

CVF sample collection and preparation

Through a sterile Cusco's vaginal speculum, 2 high vaginal swabs were obtained from the posterior vaginal fornix of each pregnant woman with sterile Dacron swab (Deltalab Eurotubo 300263, Fisher Scientific, UK) at presentation. The swabs were processed as previously described [9, 10, 18, 19]. Briefly, immediately after collection, swabs were stored at -20°C and processed within 1–3 days. The swabs were processed by placing in a 1.5 μl microfuge tube containing 400 μl (bacterial DNA extraction) and 600 μl (metabolite analysis) isotonic Phosphate Buffered Saline (PBS). The microfuge tube containing the cut end of the swab in PBS was vortexed for 5 minutes to wash the CVF into solution. The swab tip was safely discarded, and the remaining solution was centrifuged for 3 minutes at 13,000 rpm after which the supernatant was aspirated into a fresh tube and preserved at -80°C until further analysis.

CVF cytokine measurement

From the 400 μl CVF sample reserved for bacterial DNA extraction, 50 μl was aspirated and transferred into a new 1.5 μl microfuge tube and analysed for IL-1 α , IL-1 β , IL-2, IL-4, IL-6, IL-8, IL-10, IL-12p20, RANTES, TNF-r1, and IFN- γ by multiplexed bead-based immunoassay (BDTM Cytometric Bead Array, BD Biosciences, CA, USA). This was performed according to the BD CBA Human Soluble Protein Master Buffer Kit instruction (See text in [S1 File](#)). Briefly, standard solutions of each cytokine (for calibration curve), and 25 μl CVF samples were mixed with a 25 μl solution of capture beads-antibodies conjugate (left for 1 hour) and then 25 μl Phycoerythrin detection reagent (left for additional 2 hours) at room temperature. The samples were then washed, centrifuged individually, left for resuspension, and then transferred to the plate. Mean Fluorescence Intensity (MFI) were generated by a Life Technologies Attune Acoustic Focusing Cytometry and Attune NxT Cytometric Software v.2.1. Actual cytokine concentrations in each sample was extrapolated from the generated standard curve. Cytokine analyses were performed by staff blinded from patient's clinical data and eventual delivery outcome.

$^1\text{H-NMR}$ Spectroscopy

From the CVF samples dissolved in 600 μl PBS, 400 μl containing 20 μl D_2O was analysed by ^1H -Nuclear Magnetic Resonance (NMR) spectroscopy using a 400 MHz (9.4T) Bruker Avance III NMR spectrometer (Bruker BioSpin GmbH, Karlsruhe, DE), with 5mm BBO probe. Lactate, acetate, glutamate and other metabolites implicated in vaginal host-microbial activities, infection and PTD were identified in the $^1\text{H-NMR}$ spectrum ([S1 Fig](#)). Further analyses, integration and normalisation to obtain normalised integrals (n.i.) of identified metabolites were also performed as described previously [9, 10, 18, 19].

CVF metabolite measurement by spectrophotometry

The NMR-guided metabolites were chosen as the basis for a cheaper clinically applicable test [20]. We measured absolute concentrations of metabolites by enzyme-based spectrophotometric assays (Megazyme, IE): acetate (K-ACETGK 08/14), glutamate (K-GLUT 07/12), L- and D-lactate (K-LATE 07/14 and K-DATE 04/14), glucose (K-GLUHK-110A/K-GLUHK-220A 07/14), pyruvate (K-PYRUV 04/14), urea (K-URAMR 07/17), and succinate (K-SUCC 01/14), from a randomly selected subgroup of 61 women (preterm = 16, term = 45, i.e. all women

whose delivery outcomes were known at the time of analysis). All measurements except pyruvate and urea were performed on the ChemWell[®] 2910 auto-analyser (Awareness Technology, USA). Measurement of urea was performed on the ChemWell[®]-T auto-analyser (Awareness Technology, USA), while pyruvate measurement was on the MegaQuant wave spectrophotometer (Megazyme, IE).

Polymerase chain reaction (PCR)

To identify the potential microbial composition/stimulus inducing the expression of metabolites and cytokines, the prevalence of targeted vaginal bacterial species was also determined for a random subgroup of 70 women (preterm = 14, term = 66) by PCR (i.e. all women whose delivery outcomes were known at the time of analysis). Bacterial 16S rDNA were extracted using the QIAamp DNA mini kit (Qiagen, UK) and amplified by genus/species-specific primers (Sigma-Aldrich, UK, [S1 Table](#)) [21–24], as previously described [15, 18]. Amplification was performed on an Applied Biosystems 2720 Thermal cycler (Life Technologies, UK) using a 25 µl reaction mix containing 12.5 µl AmpliTaq Gold DNA polymerase (Applied Biosystems, UK), 5 ng genomic DNA template, 1 µl each of 10 µM forward and reverse primers. The cycling criteria included 95°C (5 mins)–denaturation, followed by 35 cycles of 95°C (1 min)—denaturing, 50–62°C depending on the primer sets (1 min)—annealing, 72°C (1 min)—elongation, with a final extension at 72°C (7 mins). The amplicons were visualized/confirmed on a UV-transilluminator by 1% agarose gel electrophoresis stained with ethidium bromide. Positive results were assigned according to the presence of bands of appropriate size relative to the DNA marker. Bacterial 16S rDNA standards were also used as positive controls.

Statistical analysis

Data were subjected to Shapiro-Wilk normality tests prior to analyses. Differences in maternal clinical data, CVF cytokine and metabolite concentrations between term- and preterm-delivered women as well as between women that delivered within and after 2 weeks of sampling were determined by Mann-Whitney *U* test. The relationships between cytokine, metabolite expression levels and maternal clinical and demographic variables were determined by non-parametric Spearman's (rho) correlation coefficients. *P*-values < 0.05 were considered statistically significant. Bonferroni corrections were applied for multiple measurements and correlations. The differences in the prevalence of vaginal bacterial species between preterm- and term-delivered women were determined by Fisher's exact test. Predictive capacities of CVF cytokine-metabolite combinations for sPTD (< 37 weeks of gestation) and delivery within 2 weeks of sampling were analysed by logistic regression models and area under the receiver operating characteristic curve (AUC). All analyses were performed using SPSS 24 (SPSS Inc., IL, USA), GraphPad Prism 7.03 (GraphPad Software, Inc. USA), and MedCalc 18.9 (MedCalc Software bvba, Ostend, BE; <http://www.medcalc.org>; 2018) statistical software packages.

Results

Maternal demographic and clinical details

Ninety-four women (preterm = 19, term = 75) with singleton pregnancies presenting with symptoms of PTL were enrolled in the study. Nineteen (20.2%) women delivered preterm, of which 10 (10.6%) delivered within 2 weeks of sampling. All women included in this study experienced spontaneous onset of labour, while those that underwent elective caesarean section without prior labour were excluded. Details of maternal demographic and clinical data

Table 1. Maternal clinical and demographic details according to delivery outcomes.

Characteristic	Term (N = 75)	Preterm (N = 19)	P-value	>2 weeks (N = 84)	<2 weeks (N = 10)	P-value
Age, years	26.8±5.7 (n = 70)	29.9±8.1 (n = 17)	0.2	26.9±5.8 (n = 74)	32.7±8.4 (n = 9)	0.03
BMI, kg/m ²	25.4±5.0 (n = 60)	28.6±5.7 (n = 14)	0.02	25.8±5.4 (n = 65)	26.2±3.6 (n = 7)	0.4
GAAP, weeks	30±3.8 (n = 75)	28.8±3.2 (n = 19)	0.2	29.8±3.9 (n = 80)	29.5±2.8 (n = 10)	0.6
GAAD, weeks	39.0±1.2 (n = 71)	31.6±3.1 (n = 19)	<0.0001	38.3±2.3 (n = 80)	30.2±2.9 (n = 10)	<0.0001
qfFN, ng/ml	19.2±31.6 (n = 45)	215.5±227.8 (n = 10)	0.005	44.0±105.0 (n = 47)	187.6±233.1 (n = 5)	0.1
CL, mm	30.9±10.1 (n = 42)	18.4±13.4 (n = 12)	0.002	28.7±10.8 (n = 45)	18.7±17.4 (n = 6)	0.1
Vaginal pH	4.2±0.6 (n = 41)	4.3±0.8 (n = 11)	0.9	4.2±0.6 (n = 43)	4.6±0.9 (n = 6)	0.3

Data are presented as Median ± standard deviation (SD). Nine of the preterm women delivered after 2 weeks of sampling. *BMI*, body mass index; *qfFN*, quantitative fetal fibronectin; *CL*, cervical length; *GAAP*, gestational age at presentation; *GAAD*, gestational age at delivery. n = reduced study population where participants' consent and/or data are absent.

<https://doi.org/10.1371/journal.pone.0222455.t001>

are presented in [Table 1](#). The preterm-delivered women had significantly higher BMI ($P = 0.02$) and qfFN ($P = 0.005$) and shorter CL ($P = 0.002$) compared to the term-delivered women. Most of the demographic and clinical characteristics were similar in women who delivered within and after 2 weeks of sampling, except that the women who delivered within 2 weeks of sampling were older ($P = 0.03$) than those that delivered later.

Prevalence of vaginal anaerobic bacterial species

The individual prevalence of *Fusobacterium* sp. (21%), *Mobiluncus mulieris* (18.4%) and *Mycoplasma hominis* (19.5%) was found to be higher in the preterm women compared to their term counterparts ($P < 0.0001$), while, *M. curtisii* was found in 11.6% more term-delivered than preterm women ($P < 0.0001$) ([Fig 1](#)). Furthermore, *Fusobacterium* sp. and *M. hominis* were more prevalent in women who delivered within 2 weeks of sampling, whereas *M. curtisii* and *M. mulieris* were more prevalent in women that delivered later than 2 weeks of sampling ($P < 0.0001$) ([Fig 1](#)). The prevalence of *Gardnerella vaginalis*, *Bacteroides* sp., and *Lactobacillus* sp., which were present in at least 77%, 48% and 98% of the women respectively regardless of the sub-grouping, did not differ significantly between the groups. Group B *Streptococcus* was only identified in one woman who delivered at term.

CVF cytokine concentrations

CVF cytokine concentrations in relation to delivery outcomes are presented in [Table 2](#). TNF- α ($P = 0.006$), RANTES ($P = 0.02$), IL-6 ($P = 0.03$) were significantly higher in preterm-delivered women compared to the term women. Additionally, RANTES ($P = 0.02$) and IL-6 ($P = 0.03$) were significantly higher in women who delivered within 2 weeks of sampling compared to those who delivered afterward. None of the other cytokines i.e. IL-1 α and β , and IL-8 showed significant differences in relation to delivery outcomes between the cohorts, while IL-2, IL-4, IL-10, IL-12p70 and IFN- γ were below the detectable limit of the assay kit in all the samples and were omitted in subsequent analyses.

Metabolite concentration measured by spectrophotometry

We have previously shown good correlation between ¹H-NMR and biochemical assay [[10](#)]. [Table 3](#) shows differences in metabolite concentrations measured by spectrophotometry between study cohorts. Similar to ¹H-NMR-derived normalised integrals, the absolute concentration of acetate was higher in the women who delivered within 2 weeks of sampling




Fig 1. Prevalence of vaginal mixed anaerobic bacterial species in relation to delivery outcome. (A) Preterm vs. term. The preterm-delivered women had higher prevalence of *Fusobacterium* sp. (Fuso), *Mobiluncus mulieris* (Mm), *Mycoplasma hominis* (Mh), whereas *Mobiluncus curtisii* (Mc) was more prevalent in term-delivered women though to a lesser proportion. (B) Delivery within and after 2 weeks. *Fusobacterium* sp. and *M. hominis* were more prevalent in women who delivered within 2 weeks of sampling, whereas *M. curtisii* and *M. mulieris* were more prevalent in women that delivered later than 2 weeks of sampling.

<https://doi.org/10.1371/journal.pone.0222455.g001>

compared to those that delivered later ($P = 0.047$), as reported previously in a smaller population [10]. Although the glutamate concentration was not significantly different, the ratio (Ace/Glx_{conc.}) was more than 4-fold higher in the women who delivered within 2 weeks ($P = 0.03$) and was associated with delivery within 2 weeks of presentation with PTL (AUC = 0.74) (Table 3 and Fig 2). Predictive utility of spectrophotometry by ROC analysis was similar to the ¹H-NMR-derived ratio (Ace/Glx_{n.i.}).

There were no significant differences in the concentration of the L- and D- lactate, glucose, succinate, pyruvate, and urea between any of the cohorts, and none of these metabolites was predictive of PTD nor delivery within 2 weeks of sampling individually.

Association of CVF cytokines, metabolites and maternal clinical data

The correlations between CVF pro-inflammatory mediators, metabolites (measured by enzyme-based spectrophotometry) and maternal clinical and demographic details are presented in Fig 3. After Bonferroni correction to minimise Type 1 error, TNF- α positively correlated with IL-1 α , IL-1 β , IL-6 and negatively with CL. IL-1 β also correlated positively with L/D-lactate ratio which correlated positively with vaginal pH. Furthermore, D-lactate correlated positively with glutamate and inversely with vaginal pH. CL and qfFN were also negatively correlated.

Table 2. Comparison of cervicovaginal fluid cytokine/chemokine concentrations (pg/ml) in relation to delivery outcomes.

Cytokine/chemokine	Term (N = 75)	Preterm (N = 19)	P-value
TNF-r1	781.4 (283.3–1814.5) (n = 75)	2350.7 (645.7–3795.8) (n = 19)	0.006
IL-1β	369.5 (69.4–760.8) (n = 61)	785.6 (133.5–1653.4) (n = 9)	0.2
RANTES	40.8 (12.8–67.3) (n = 9)	101.4 (70.7–541.3) (n = 6)	0.02
IL-6	38.3 (6.1–147.2) (n = 47)	207.6 (47.1–524.5) (n = 13)	0.01
IL-8	771.6 (299.3–1385.3) (n = 20)	490.3 (402.5–1827.8) (n = 4)	0.8
IL-1α	546.2 (218.7–935.5) (n = 62)	1062.4 (397.6–1816.5) (n = 17)	0.3
Delivery within 2 weeks of sampling			
	> 2 weeks (N = 84)	< 2 weeks (N = 10)	P-value
TNF-r1	922.5 (323.2–1936.5) (n = 80)	2083.8 (454.6–3187.0) (n = 10)	0.2
IL-1β	362.1 (75.2–1018.3) (n = 62)	628.8 (125.9–1883.5) (n = 4)	0.6
RANTES	40.8 (17.9–79.2) (n = 11)	289.1 (89.7–660.2) (n = 4)	0.02
IL-6	47.8 (6.1–147.2) (n = 51)	385.8 (56.8–592.9) (n = 7)	0.03
IL-8	490.3 (273.3–0.0) (n = 22)	490.3 (473.3–0.0) (n = 2)	0.4
IL-1α	583.6 (231.8–1138.7) (n = 67)	1062.4 (298.5–1940.9) (n = 9)	0.3

Data presented as median (25th - 75th percentile). Samples without a particular cytokine or cytokine concentration below the detectable limit of the assay kit (indicated as a value of zero) were omitted in subsequent analyses, hence the varied *n* numbers. Significant *P*-values are marked in bold.

<https://doi.org/10.1371/journal.pone.0222455.t002>

Receiver operating characteristic analysis of CVF biomarkers and clinical details for spontaneous PTB

As shown in [S2 Table](#), RANTES, TNF-r1, IL-6, Ace/Glx_{conc}, BMI, qfFN, which were significantly increased and CL, which was significantly decreased in the preterm women were able to determine risk of sPTD individually, but only RANTES, IL-6, Ace/Glx_{conc} and maternal age were associated with delivery within 2 weeks of sampling.

Furthermore, certain cytokine-metabolite combinations containing lactate, acetate, Ace/Glx_{conc}, IL-6, and TNF-r1 were associated with sPTD, while others were associated with delivery within 2 weeks of sampling ([S2 Table](#)).

RANTES, which had the highest individual predictive values for sPTD (AUC = 0.87) and delivery within 2 weeks of sampling (AUC = 0.91) ([Fig 4](#) and [S2 Table](#)) could not be included in the combination (multibiomarker) models by logistic regression due to reduced number of samples with detectable RANTES concentrations.

Table 3. Cervicovaginal fluid metabolites concentrations in relation to delivery outcome.

Metabolites	Term	Preterm	AUC (95% CI)	P-value
Acetate	0.02 (0.01–0.03) n = 41	0.04 (0.02–0.08) n = 15	0.79(0.66–0.89)	0.0007
Glutamate	0.07 (0.04–0.10) n = 42	0.07(0.04–0.13) n = 15	0.53(0.40–0.67)	0.7
Acetate/Glutamate ratio	0.27 (0.14–0.50) n = 42	0.83 (0.30–1.89) n = 15	0.74 (0.60–0.85)	0.006
Delivery within 2 weeks				
	> 2 weeks	< 2 weeks	AUC (95% CI)	P-value
Acetate	0.021 (0.01–0.03) (n = 47)	0.054 (0.02–0.08) (n = 8)	0.73 (0.59–0.84)	0.047
Glutamate	0.068 (0.04–0.10) (n = 49)	0.063 (0.05–0.08) (n = 8)	0.54 (0.41–0.68)	0.7
Acetate/Glutamate ratio	0.282 (0.16–0.58) n = 47	1.153 (0.40–1.77) n = 8	0.74 (0.61–0.85)	0.03

Concentrations of metabolites (g/l) derived by spectrophotometry are presented as median (25th and 75th percentiles). AUC, area under the ROC curve; CI, confidence interval.

<https://doi.org/10.1371/journal.pone.0222455.t003>




Fig 2. Receiver operating characteristic curve analysis of cervicovaginal fluid metabolites in predicting delivery within 2 weeks of presentation with symptoms of preterm labour. (A) ¹H-NMR-derived acetate/glutamate ratio (Ace/Glx, AUC = 0.86, 0.76–0.93), Acetate (AUC = 0.78, 0.68–0.87) and Glutamate (AUC = 0.68, 0.57–0.78); (B) Metabolite concentrations measured by enzyme-based spectrophotometry: Ace/Glx (AUC = 0.74, 0.61–0.85), Acetate (AUC = 0.73, 0.59–0.84) and Glutamate (AUC = 0.54, 0.41–0.68). *n.i.*, normalised integral.

<https://doi.org/10.1371/journal.pone.0222455.g002>

Discussion

This study investigated the relationship of CVF inflammatory mediators, microbiota metabolites and risk of imminent sPTD in women presenting with PTL. We observed that women who delivered preterm or within 2 weeks of sampling demonstrated higher prevalence of potentially pathogenic vaginal anaerobes in the CVF, and showed significantly higher markers of vaginal dysbiosis (Ace/Glx ratio), inflammation (RANTES, TNFr-1 and IL-6), shorter CL




Fig 3. Correlation of cervicovaginal fluid pro-inflammatory mediators, metabolite concentrations and maternal clinical and demographic details. Spearman's correlation coefficients (*r*, colour) range from -1 (dark blue) to +1 (dark red); and significant *P*-values (< 0.05) stated in the coloured boxes. Due to the high number (28) of compared measures, a Bonferroni corrected value of 0.05/364 = *p* < 0.000137 was applied to minimise Type 1 error. *Ace/Glx*, acetate/glutamate ratio; *CL*, cervical length; *BMI*, body mass Index; *FFN*, quantitative fetal fibronectin; *Gest.*, gestational; *g.w*, gestational week; *g.d*, gestational day; *labour*, latency i.e. interval between presentation and delivery.

<https://doi.org/10.1371/journal.pone.0222455.g003>




Fig 4. Receiver operating characteristic curve analysis of cervicovaginal fluid RANTES (A-B) and IL-6 (C-D) in predicting preterm delivery and delivery within 2 weeks of presentation with symptoms of preterm labour. AUC, area under the ROC curve.

<https://doi.org/10.1371/journal.pone.0222455.g004>

and increased qfFN. Cytokine-metabolite combinations of total lactate, L/D-lactate ratio, acetate, acetate/glutamate ratio, IL-6, TNF- α were also associated with sPTD and delivery within 2 weeks of presentation.

The observed cytokine-metabolite alterations were accompanied by high prevalence of anaerobes such as *Fusobacterium* sp., *M. hominis* and *M. mulieris* in women destined to deliver preterm; while those that delivered within 2 weeks of sampling showed higher prevalence of *Fusobacterium* sp. and *M. hominis* only. This is consistent with reports of elevated CVF IL-6 and positive cultures of *Ureaplasma urealyticum* and *M. hominis* in the chorioamniotic membranes in women with symptoms of PTL [25]. Our observations are also consistent with reports that vaginal communities dominated by anaerobes are potentially associated with greater pro-inflammatory cytokine responses than *Lactobacilli*-dominated niches [26, 27].

Increased vaginal concentrations of acetate and glutamate have contrasting implications for reproductive health. Acetate in the vaginal milieu is produced majorly by anaerobes, especially when *Lactobacilli* are deficient [28–30], and high amounts are associated with infection and PTD [9, 10, 18, 30]. Contrastingly, high vaginal glutamate level, similar to D-lactate, is associated with healthy microbiota dominated by *Lactobacilli* [11, 31] and decreased prevalence of anaerobes that are known to metabolise amino acids to their catabolic by-products [11] including acetate [18, 30]. For instance, *Fusobacterium* sp. which was more prevalent in the women who delivered preterm and within 2 weeks of sampling, ferments glutamate to acetate via the hydroxyglutarate pathway [32, 33]. Such metabolic processes explain our observation that the Ace/Glx ratio was associated with sPTD and delivery within 2 weeks of sampling. It is noteworthy that glutamate alone was not associated with delivery within 2 weeks, while acetate alone was to a lesser extent associated with imminent sPTD [9, 10]. Therefore, the Ace/Glx ratio was more strongly associated with delivery within 14 days of sampling than either acetate

or glutamate individually. Furthermore, decreased glutamate, D-lactate, and increased L/D-lactate ratio were associated with high vaginal pH.

Despite our limited sample size, an association of CVF RANTES, IL-6 and TNF- α with sPTD within 14 days was also demonstrated. Additionally, the combination of metabolite indices (acetate, glutamate, L- and D-lactate) and cytokine concentrations were similarly associated with imminent sPTD, observations which are reflective of vaginal host-microbial interactions [34], being downstream products of gene expression and protein synthesis [35–37]. Our study gives additional insight into the role played by the vaginal ecosystem and host immune responses in the propagation of sPTD [38], and whether specific cytokine-metabolite combinations can be used to screen for women at high risk of imminent sPTD.

Elevated blood, serum and CVF RANTES at mid to late gestation has been associated with sPTD in both symptomatic [3, 37] and asymptomatic women [15, 39], as well as murine models [40]. We recently reported decreased CVF RANTES from mid to late second trimester in asymptomatic women who delivered at term compared to their preterm counterparts in whom levels remained unchanged [15]. RANTES is a potent chemoattractant and inducer of inflammatory mediators involved in parturition and is produced by immune, endothelial and gestational tissues [37, 41, 42].

Apart from RANTES, TNF- α (that binds TNF- α) and IL-6 were also increased in women who delivered preterm. IL-6 was additionally associated with delivery within 2 weeks. These pro-inflammatory mediators are implicated in the NF- κ B-p38MAPK-orchestrated inflammatory cascade implicated in sPTL leading to myometrial contraction, cervical remodelling, and degradation/rupture of fetal membranes [31, 43–45]. TNF- α is detectable early in the amniotic fluid following bacterial colonisation and stimulates the production of IL-6, IL-8, matrix metalloproteinases and PGE₂ during infection-associated PTL [42]. IL-6 activates the acute phase response [46] and, along with TNF- α , which were correlated in our study, correlates with qfFN [47] and is associated with PTD [48–50].

An altered vaginal microbiota with reduced *Lactobacillus* sp. dominance may increase vaginal pH creating a permissive environment for the proliferation of pathogenic anaerobes [51]. There is also a direct link between vaginal dysbiosis, short cervix [45, 52], and disrupted fetal membranes with consequent leakage of fFN into the cervicovaginal space [7, 45, 53]. Therefore, a high CVF Ace/Glx ratio could be suggestive of a dysbiosis with associated reduced D-lactate, glutamate and vaginal acidity, increased expression of pro-inflammatory mediators, shortened cervix and elevated qfFN as shown in this study and other reports [3, 15, 25, 30, 31, 45, 50].

Hitherto, no multivariable model of inflammatory mediators and cervical length associated with imminent sPTD in symptomatic [3, 54] and asymptomatic women [15, 55] has achieved clinical utility. Interestingly, we observed amongst others that combination of CVF: 1) acetate, lactate, IL-6 and TNF- α ; and 2) Ace/Glx ratio, L/D-lactate ratio and IL-6 were associated with sPTD and delivery within 14 days of sampling, respectively.

Two recent extensive systematic reviews of the accuracy of qfFN for the prediction of PTD within 7 to 10 days of testing in symptomatic women reported pooled sensitivities of 75% and 77%; and pooled specificities of 79% and 83% respectively [56, 57]. Both our single and multivariable cytokine-metabolite models on a similar cohort showed greater sensitivities and specificities for determining sPTD within 2 weeks of sampling respectively. However, there is the need to test these findings in larger and more diverse study populations.

Conclusion

Symptomatic women destined to deliver preterm within 2 weeks express significantly higher pro-inflammatory mediators at mid to late gestation. In this cohort, RANTES, IL-6, and

acetate/glutamate ratio; were more associated with imminent preterm delivery compared to other biomarkers. The combination of CVF metabolites and pro-inflammatory mediators appears able to distinguish symptomatic women at risk of imminent preterm birth. This approach may improve the triaging of such women, allowing patients to be discharged to out-patient care whilst sign-posting neonatal high-dependency care to those at highest risk. A larger study is required to confirm these observations and determine the cost utility of this approach in clinical care.

Supporting information

S1 Fig. Cervicovaginal fluid metabolite peaks identified on ¹H-NMR spectrum at 400 MHz and 294K. BCAA, branched chain amino acids; *ppm*, parts per million.
(TIF)

S1 Table. Primer sequences used for bacterial 16S rDNA amplification. GBS, group B *Streptococcus*.
(PDF)

S2 Table. Predictive performance of biomarkers for imminent preterm delivery. AUC, area under the ROC curve; CI, confidence interval; Sens, sensitivity; Spec, specificity; PPV, positive predictive value; NPV, negative predictive value; +LR, positive likelihood ratio; -LR, negative likelihood ratio; BMI, body mass index; CL, cervical length; qfFN, quantitative fetal fibronectin; Ace/Glx_{conc}, ratio of acetate to glutamate concentrations. All metabolite concentrations were measured by enzyme-based spectrophotometry.
(PDF)

S1 File. CVF cytokine measurement.
(PDF)

S1 Dataset. CVF cytokines, metabolites and PCR data.
(ZIP)

Acknowledgments

We are grateful to the women who consented to participate in these studies, Dr Graham Stafford who co-supervised EA during his PhD studies, and the management and staff of Megazyme, Ireland particularly Dr Vincent A. McKie and Dr Orlaith Dowling for providing infrastructural and technical support toward the spectrophotometric experiments.

Parts of this work have been presented at the 64th Annual Scientific Meeting of the Society for Reproductive Investigation, Orlando 2017, Rep Sci 24(Suppl 1):175A-175A (F-018) (Poster); the 19th annual conference of the British Maternal and Fetal Medicine Society, Amsterdam, BJOG 2017, 124 (S2):127, Oral (O.PO.7) and Poster (P.PO. 17); 66th annual scientific meeting of the Society for Reproductive Investigation, Paris 2019, Rep Sci 26(Suppl 1):301A-301A (S-008) (Poster) and 301A-302A (S-009) (Poster); and 21st annual scientific meeting of the British Maternal and Fetal Medicine Society, Edinburgh, BJOG 2019, 126(S1): 132–133 (EP. 323, 327) (Posters).

Author Contributions

Conceptualization: Emmanuel Amabebe, Dilly O. C. Anumba.

Data curation: Emmanuel Amabebe, Steven Reynolds, Xiaoya He, Robyn Wood, Victoria Stern, Dilly O. C. Anumba.

Formal analysis: Emmanuel Amabebe, Steven Reynolds, Xiaoya He, Robyn Wood, Victoria Stern, Dilly O. C. Anumba.

Funding acquisition: Dilly O. C. Anumba.

Investigation: Emmanuel Amabebe, Xiaoya He, Robyn Wood, Victoria Stern, Dilly O. C. Anumba.

Methodology: Emmanuel Amabebe, Steven Reynolds, Xiaoya He, Robyn Wood, Victoria Stern, Dilly O. C. Anumba.

Project administration: Emmanuel Amabebe, Dilly O. C. Anumba.

Resources: Victoria Stern, Dilly O. C. Anumba.

Supervision: Emmanuel Amabebe, Dilly O. C. Anumba.

Writing – original draft: Emmanuel Amabebe, Steven Reynolds, Xiaoya He, Robyn Wood, Victoria Stern, Dilly O. C. Anumba.

Writing – review & editing: Emmanuel Amabebe, Steven Reynolds, Dilly O. C. Anumba.

References

1. Bianchi-Jassir F, Seale AC, Kohli-Lynch M, Lawn JE, Baker CJ, Bartlett L, et al. Preterm Birth Associated With Group B Streptococcus Maternal Colonization Worldwide: Systematic Review and Meta-analyses. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America*. 2017; 65(Suppl 2):S133–S42. <https://doi.org/10.1093/cid/cix661> PMC5850429. PMID: [29117329](https://pubmed.ncbi.nlm.nih.gov/29117329/)
2. Goldenberg RL, Culhane JF, Iams JD, Romero R. Preterm birth 1—Epidemiology and causes of preterm birth. *Lancet*. 2008; 371(9606):75–84. [https://doi.org/10.1016/S0140-6736\(08\)60074-4](https://doi.org/10.1016/S0140-6736(08)60074-4) WOS:000252192600033. PMID: [18177778](https://pubmed.ncbi.nlm.nih.gov/18177778/)
3. Tsiartas P, Holst R, Wennerholm U, Hagberg H, Hougaard D, Skogstrand K, et al. Prediction of spontaneous preterm delivery in women with threatened preterm labour: a prospective cohort study of multiple proteins in maternal serum. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2012; 119(7):866–73.
4. Goldenberg RL. The management of preterm labor. *Obstetrics & Gynecology*. 2002; 100(5, Part 1):1020–37.
5. Honest H, Bachmann LM, Gupta JK, Kleijnen J, Khan KS. Accuracy of cervicovaginal fetal fibronectin test in predicting risk of spontaneous preterm birth: systematic review. *British Medical Journal*. 2002; 325(7359):301–4C. <https://doi.org/10.1136/bmj.325.7359.301> WOS:000177482500015. PMID: [12169504](https://pubmed.ncbi.nlm.nih.gov/12169504/)
6. Ramsey PS, Andrews WW. Biochemical predictors of preterm labor: fetal fibronectin and salivary estriol. *Clinics in perinatology*. 2003; 30(4):701–33. PMID: [14714920](https://pubmed.ncbi.nlm.nih.gov/14714920/)
7. Heng YJ, Liong S, Permezel M, Rice GE, Di Quinzio MK, Georgiou HM. Human cervicovaginal fluid biomarkers to predict term and preterm labor. *Frontiers in physiology*. 2015;6.
8. Liong S, Di Quinzio M, Fleming G, Permezel M, Rice G, Georgiou H. New biomarkers for the prediction of spontaneous preterm labour in symptomatic pregnant women: a comparison with fetal fibronectin. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2015; 122(3):370–9.
9. Amabebe E, Reynolds S, Stern VL, Parker JL, Stafford GP, Paley MN, et al. Identifying metabolite markers for preterm birth in cervicovaginal fluid by magnetic resonance spectroscopy. *Metabolomics*. 2016; 12(4):1–11.
10. Amabebe E, Reynolds S, Stern V, Stafford G, Paley M, Anumba DOC. Cervicovaginal Fluid Acetate: A Metabolite Marker of Preterm Birth in Symptomatic Pregnant Women. *Frontiers in Medicine*. 2016; 3(48). <https://doi.org/10.3389/fmed.2016.00048> PMID: [27777928](https://pubmed.ncbi.nlm.nih.gov/27777928/)
11. Srinivasan S, Morgan MT, Fiedler TL, Djukovic D, Hoffman NG, Raftery D, et al. Metabolic Signatures of Bacterial Vaginosis. *mBio*. 2015; 6(2):e00204–15. <https://doi.org/10.1128/mBio.00204-15> PMID: [25873373](https://pubmed.ncbi.nlm.nih.gov/25873373/)
12. Lamont RF. Advances in the Prevention of Infection-Related Preterm Birth. *Frontiers in immunology*. 2015; 6(566). <https://doi.org/10.3389/fimmu.2015.00566> PMID: [26635788](https://pubmed.ncbi.nlm.nih.gov/26635788/)
13. Nelson DB, Hanlon A, Nachamkin I, Haggerty C, Mastrogiannis DS, Liu C, et al. Early Pregnancy Changes in Bacterial Vaginosis-Associated Bacteria and Preterm Delivery. *Paediatric and perinatal epidemiology*. 2014; 28(2):88–96. <https://doi.org/10.1111/ppe.12106> PMID: [24405280](https://pubmed.ncbi.nlm.nih.gov/24405280/)

14. Foxman B, Wen A, Srinivasan U, Goldberg D, Marrs CF, Owen J, et al. Mycoplasma, bacterial vaginosis-associated bacteria BVAB3, race, and risk of preterm birth in a high-risk cohort. *American journal of obstetrics and gynecology*. 2014; 210(3):226.e1–e2267. Epub 2013/10/04. <https://doi.org/10.1016/j.ajog.2013.10.003> PMID: 24096128.
15. Amabebe E, Chapman DR, Stern VL, Stafford G, Anumba DOC. Mid-gestational changes in cervicovaginal fluid cytokine levels in asymptomatic pregnant women are predictive markers of inflammation-associated spontaneous preterm birth. *Journal of Reproductive Immunology*. 2018; 126:1–10. <https://doi.org/10.1016/j.jri.2018.01.001> PMID: 29367099
16. Chan RL. Biochemical markers of spontaneous preterm birth in asymptomatic women. *BioMed research international*. 2014;2014.
17. Georgiou HM, Di Quinzio MK, Permezel M, Brennecke SP. Predicting Preterm Labour: Current Status and Future Prospects. *Disease markers*. 2015; 2015:435014. Epub 2015/07/15. <https://doi.org/10.1155/2015/435014> PMID: 26160993; PubMed Central PMCID: PMC4486247.
18. Amabebe E. Analysis of cervicovaginal fluid metabolome and microbiome in relation to preterm birth [PhD thesis]. White Rose eTheses Online: University of Sheffield; 2016.
19. Stafford GP, Parker JL, Amabebe E, Kistler J, Reynolds S, Stern V, et al. Spontaneous Preterm Birth Is Associated with Differential Expression of Vaginal Metabolites by Lactobacilli-Dominated Microflora. *Frontiers in Physiology*. 2017; 8(615). <https://doi.org/10.3389/fphys.2017.00615> PMID: 28878691
20. Amabebe E, Reynolds S, Stern V, Stafford G, Paley M, Anumba D. Prognostic Capacity of Cervicovaginal Fluid Acetate-Glutamate Ratio for Risk of Preterm Delivery within Two Weeks of Presentation with Symptoms of Preterm Labor. *Reproductive Sciences*. 2017; 24:175A-A. WOS:000399043900390.
21. Bernhard AE, Field KG. A PCR assay To discriminate human and ruminant feces on the basis of host differences in Bacteroides-Prevotella genes encoding 16S rRNA. *Applied and environmental microbiology*. 2000; 66(10):4571–4. Epub 2000/09/30. <https://doi.org/10.1128/aem.66.10.4571-4574.2000> PMID: 11010920; PubMed Central PMCID: PMC92346.
22. Ling Z, Kong J, Liu F, Zhu H, Chen X, Wang Y, et al. Molecular analysis of the diversity of vaginal microbiota associated with bacterial vaginosis. *BMC genomics*. 2010; 11. <https://doi.org/10.1186/1471-2164-11-488> WOS:000282790600001. PMID: 20819230
23. Walter J, Margosch D, Hammes WP, Hertel C. Detection of Fusobacterium species in human feces using genus-specific PCR primers and denaturing gradient gel electrophoresis. *Microbial Ecology in Health and Disease*. 2002; 14(3):129–32. <https://doi.org/10.1080/089106002320644294> BCI: BCI200300038741.
24. Zariffard MR, Saifuddin M, Sha BE, Spear GT. Detection of bacterial vaginosis-related organisms by real-time PCR for Lactobacilli, Gardnerella vaginalis and Mycoplasma hominis. *FEMS immunology and medical microbiology*. 2002; 34(4):277–81. Epub 2002/11/22. <https://doi.org/10.1111/j.1574-695X.2002.tb00634.x> PMID: 12443827.
25. Jacobsson B, Mattsby-Baltzer I, Hagberg H. Interleukin-6 and interleukin-8 in cervical and amniotic fluid: relationship to microbial invasion of the chorioamniotic membranes. 2005; 112(6):719–24. <https://doi.org/10.1111/j.1471-0528.2005.00536.x> PMID: 15924526
26. Witkin S, Linhares I. Why do lactobacilli dominate the human vaginal microbiota? 2017; 124(4):606–11. <https://doi.org/10.1111/1471-0528.14390> PMID: 28224747
27. Smith SB, Ravel J. The vaginal microbiota, host defence and reproductive physiology. 2017; 595(2):451–63. <https://doi.org/10.1113/JP271694> PMID: 27373840
28. Vitali B, Cruciani F, Picone G, Parolin C, Donders G, Laghi L. Vaginal microbiome and metabolome highlight specific signatures of bacterial vaginosis. *Eur J Clin Microbiol*. 2015; 34(12):2367–76.
29. Laghi L, Picone G, Cruciani F, Brigidi P, Calanni F, Donders G, et al. Rifaximin modulates the vaginal microbiome and metabolome in women affected by bacterial vaginosis. *Antimicrobial agents and chemotherapy*. 2014; 58(6):3411–20. <https://doi.org/10.1128/AAC.02469-14> PMID: 24709255
30. Aldunate M, Srbinovski D, Hearps AC, Latham CF, Ramsland PA, Gugasyan R, et al. Antimicrobial and immune modulatory effects of lactic acid and short chain fatty acids produced by vaginal microbiota associated with eubiosis and bacterial vaginosis. *Frontiers in physiology*. 2015; 6:164. <https://doi.org/10.3389/fphys.2015.00164> PMID: 26082720
31. Amabebe E, Anumba DOC. The Vaginal Microenvironment: The Physiologic Role of Lactobacilli. 2018; 5(181). <https://doi.org/10.3389/fmed.2018.00181> PMID: 29951482
32. Wolfgang Buckel HAB. Two Pathways of Glutamate Fermentation by Anaerobic Bacteria. *Journal of Bacteriology*. 1974; 117(3):1248–60. PMID: 4813895
33. Ramezani M, Resmer KL, White RL. Glutamate racemization and catabolism in *Fusobacterium varium*. *The FEBS Journal*. 2011; 278(14):2540–51. <https://doi.org/10.1111/j.1742-4658.2011.08179.x> PMID: 21575137

34. Gajer P, Brotman RM, Bai GY, Sakamoto J, Schuette UME, Zhong X, et al. Temporal Dynamics of the Human Vaginal Microbiota. *Sci Transl Med*. 2012; 4(132). ARTN 132ra52 <https://doi.org/10.1126/scitranslmed.3003605> ISI:000303596400004. PMID: [22553250](https://pubmed.ncbi.nlm.nih.gov/22553250/)
35. Fanos V, Atzori L., Makarenko K., Melis G.B., Ferrazi E. Metabolomics application in maternal-fetal medicine. *BioMed research international*. 2013; 2013(720514):<http://dx.doi.org/10.1155/2013/720514>.
36. Romero R, Mazaki-Tovi S., Vaisbuch E., Kusanovic J.P., Chaiworapongsa T., Gomez R., Nien J.K., Yoon B.H., Mazor M., Luo J., Banks D., Ryals J., Beecher C. Metabolomics in premature labor: a novel approach to identify patients at risk for preterm delivery. *J Matern Fetal Neonatal Med*. 2010; 23(12):1344–59. <https://doi.org/10.3109/14767058.2010.482618> PMID: [20504069](https://pubmed.ncbi.nlm.nih.gov/20504069/)
37. Hamilton SA, Tower CL, Jones RL. Identification of Chemokines Associated with the Recruitment of Decidual Leukocytes in Human Labour: Potential Novel Targets for Preterm Labour. *PLOS ONE*. 2013; 8(2):e56946. <https://doi.org/10.1371/journal.pone.0056946> PMID: [23451115](https://pubmed.ncbi.nlm.nih.gov/23451115/)
38. Ghartey J, Bastek JA, Brown AG, Anglim L, Elovitz MA. Women with preterm birth have a distinct cervicovaginal metabolome. *American Journal of Obstetrics and Gynecology*. 2015; 212(6):776. e1–e12.
39. Chow SS, Craig ME, Jones CA, Hall B, Cateau J, Lloyd AR, et al. Differences in amniotic fluid and maternal serum cytokine levels in early midtrimester women without evidence of infection. *Cytokine*. 2008; 44(1):78–84. Epub 2008/08/16. <https://doi.org/10.1016/j.cyto.2008.06.009> PMID: [18703348](https://pubmed.ncbi.nlm.nih.gov/18703348/)
40. Yang Q, El-Sayed Y, Rosenberg-Hasson Y, Hirschberg DL, Nayak NR, Schilling J, et al. Multiple cytokine profile in plasma and amniotic fluid in a mouse model of pre-term labor. *Am J Reprod Immunol*. 2009; 62(5):339–47. Epub 2009/10/09. <https://doi.org/10.1111/j.1600-0897.2009.00743.x> PMID: [19811468](https://pubmed.ncbi.nlm.nih.gov/19811468/)
41. Athayde N, Romero R, Maymon E, Gomez R, Pacora P, Araneda H, et al. A role for the novel cytokine RANTES in pregnancy and parturition. *American Journal of Obstetrics and Gynecology*. 1999; 181(4):989–94. [https://doi.org/10.1016/s0002-9378\(99\)70337-6](https://doi.org/10.1016/s0002-9378(99)70337-6) PMID: [10521766](https://pubmed.ncbi.nlm.nih.gov/10521766/)
42. Vrachnis N, Karavolos S, Iliodromiti Z, Sifakis S, Siristatidis C, Mastorakos G, et al. Impact of mediators present in amniotic fluid on preterm labour. *In Vivo*. 2012; 26(5):799–812. PMID: [22949593](https://pubmed.ncbi.nlm.nih.gov/22949593/)
43. Agrawal V, Hirsch E. Intrauterine infection and preterm labor. *Seminars in Fetal and Neonatal Medicine*. 2012; 17(1):12–9. <https://doi.org/10.1016/j.siny.2011.09.001> PMID: [21944863](https://pubmed.ncbi.nlm.nih.gov/21944863/)
44. Keelan JA. Pharmacological inhibition of inflammatory pathways for the prevention of preterm birth. *Journal of Reproductive Immunology*. 2011; 88(2):176–84. <https://doi.org/10.1016/j.jri.2010.11.003> PMID: [21236496](https://pubmed.ncbi.nlm.nih.gov/21236496/)
45. Keelan JA. Intrauterine inflammatory activation, functional progesterone withdrawal, and the timing of term and preterm birth. *Journal of Reproductive Immunology*. 2018; 125:89–99. <https://doi.org/10.1016/j.jri.2017.12.004> PMID: [29329080](https://pubmed.ncbi.nlm.nih.gov/29329080/)
46. Kalinka J, Sobala W, Wasiela M, Brzezińska-Błaszczak E. Decreased Proinflammatory Cytokines in Cervicovaginal Fluid, as Measured in Midgestation, are Associated with Preterm Delivery. *American Journal of Reproductive Immunology*. 2005; 54(2):70–6. <https://doi.org/10.1111/j.1600-0897.2005.00289.x> PMID: [16105098](https://pubmed.ncbi.nlm.nih.gov/16105098/)
47. Inglis SR, Jeremias J, Kuno K, Lescale K, Peeper Q, Chervenak FA, et al. Detection of tumor necrosis factor-alpha, interleukin-6, and fetal fibronectin in the lower genital tract during pregnancy: relation to outcome. *Am J Obstet Gynecol*. 1994; 171(1):5–10. Epub 1994/07/01. [https://doi.org/10.1016/s0002-9378\(94\)70069-9](https://doi.org/10.1016/s0002-9378(94)70069-9) PMID: [8030732](https://pubmed.ncbi.nlm.nih.gov/8030732/)
48. Paternoster DM, Stella A, Gerace P, Manganelli F, Plebani M, Snijders D, et al. Biochemical markers for the prediction of spontaneous pre-term birth. 2002; 79(2):123–9. [https://doi.org/10.1016/S0020-7292\(02\)00243-6](https://doi.org/10.1016/S0020-7292(02)00243-6)
49. Lockwood CJ, Ghidini I, Wein R, Lapinski R, Casal D, Berkowitz RL. Increased interleukin-6 concentrations in cervical secretions are associated with preterm delivery. *American Journal of Obstetrics and Gynecology*. 1994; 171(4):1097–102. [https://doi.org/10.1016/0002-9378\(94\)90043-4](https://doi.org/10.1016/0002-9378(94)90043-4) PMID: [7943078](https://pubmed.ncbi.nlm.nih.gov/7943078/)
50. Kalan AM, Simhan HN. Mid-trimester cervical inflammatory milieu and sonographic cervical length. *American Journal of Obstetrics and Gynecology*. 2010; 203(2):126.e1–e5. <https://doi.org/10.1016/j.ajog.2010.03.013>
51. Smith SB, Ravel J. The vaginal microbiota, host defence and reproductive physiology. *The Journal of Physiology*. 2016:n/a-n/a. <https://doi.org/10.1113/JP271694> PMID: [27373840](https://pubmed.ncbi.nlm.nih.gov/27373840/)
52. Donders GG, Van Calsteren C, Bellen G, Reybrouck R, Van den Bosch T, Riphagen I, et al. Association between abnormal vaginal flora and cervical length as risk factors for preterm birth. *Ultrasound in Obstetrics & Gynecology*. 2010.
53. Agrawal V, Hirsch E, editors. *Intrauterine infection and preterm labor*. *Seminars in Fetal and Neonatal Medicine*; 2012: Elsevier.

54. Holst RM, Hagberg H, Wennerholm UB, Skogstrand K, Thorsen P, Jacobsson B. Prediction of spontaneous preterm delivery in women with preterm labor: analysis of multiple proteins in amniotic and cervical fluids. *Obstet Gynecol.* 2009; 114(2 Pt 1):268–77. Epub 2009/07/23. <https://doi.org/10.1097/AOG.0b013e3181ae6a08> PMID: 19622987.
55. Vogel I, Goepfert AR, Thorsen P, Skogstrand K, Hougaard DM, Curry AH, et al. Early second-trimester inflammatory markers and short cervical length and the risk of recurrent preterm birth. *J Reprod Immunol.* 2007; 75(2):133–40. Epub 2007/04/20. <https://doi.org/10.1016/j.jri.2007.02.008> PMID: 17442403.
56. Deshpande S, van Asselt A, Tomini F, Armstrong N, Allen A, Noake C, et al. Rapid fetal fibronectin testing to predict preterm birth in women with symptoms of premature labour: a systematic review and cost analysis. 2013.
57. Boots AB, Sanchez-Ramos L, Bowers DM, Kaunitz AM, Zamora J, Schlattmann P. The short-term prediction of preterm birth: a systematic review and diagnostic metaanalysis. *American Journal of Obstetrics and Gynecology.* 2014; 210(1):54.e1–.e10. <http://dx.doi.org/10.1016/j.ajog.2013.09.004>.