



UNIVERSITY OF LEEDS

This is a repository copy of *The relationship between physical examination and ultrasonography for large entheses is best for the Achilles tendon and patellar tendon origin*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/151885/>

Version: Accepted Version

Article:

Aydin, SZ, Bakirci, S, Kasapoglu, E et al. (8 more authors) (2020) The relationship between physical examination and ultrasonography for large entheses is best for the Achilles tendon and patellar tendon origin. *Journal of Rheumatology*, 47 (7). pp. 1026-1030. ISSN 0315-162X

<https://doi.org/10.3899/jrheum.190169>

© 2019 The Journal of Rheumatology. This is an author produced version of an article published in *The Journal of Rheumatology*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

The relationship between physical examination and ultrasonography for large entheses is best for the Achilles tendon and patellar tendon origin

Sibel Zehra Aydin, Sibel Bakirci, Esen Kasapoglu, Concepcion Castillo-Gallego, Fatıma Arslan Alhussain, Zoe R Ash, Esra Kurum, Dennis McGonagle, Helena Marzo-Ortega, Dafna Gladman, Lihi Eder

Key Indexing Terms: Ultrasound, physical examination, enthesopathy

This research is supported by the National Institute for Health Research (NIHR) Leeds Biomedical Research Centre. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

Funding: None

Conflict of interests: None relevant for the manuscript

Sibel Zehra Aydin, MD, Assoc Prof, University of Ottawa Faculty of Medicine, Ottawa Hospital Research Institute, Department of Internal Medicine, Division of Rheumatology, Ottawa, ON, CANADA.

Sibel Bakirci, MD, University of Ottawa Faculty of Medicine, Department of Internal Medicine, Division of Rheumatology, Ottawa, ON, CANADA.

Esen Kasapoglu, MD, Assoc. Prof, Istanbul Medeniyet University, Faculty of Medicine, Department of Internal Medicine, Division of Rheumatology, Istanbul, TURKEY.

Concepcion Castillo-Gallego, MD, Complejo Hospitalario Torrecárdenas, Department of Internal Medicine, Division of Rheumatology, Almería, SPAIN.

Fatıma Arslan Alhussain, MD, Istanbul Medeniyet University, Department of Internal Medicine, Istanbul, TURKEY.

Zoe Ash, MD, NIHR Leeds Biomedical Research Centre, Leeds Teaching Hospitals Trust and Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds. Leeds, UK.

Esra Kurum, Assistant Professor, University of California, Department of Statistics, Riverside, California, USA

Dennis McGonagle, MD, Prof, NIHR Leeds Biomedical Research Centre, Leeds Teaching Hospitals Trust and Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds. Leeds, UK.

Helena Marzo-Ortega, MD, NIHR Leeds Biomedical Research Centre, Leeds Teaching Hospitals Trust and Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds. Leeds, UK.

Dafna Gladman, MD, Professor of Medicine, University of Toronto, Department of Internal Medicine, Division of Rheumatology, Krembil research Institute, Toronto Western Hospital, Toronto, CANADA.

Lihi Eder, MD, PhD, Assistant Professor, University of Toronto, Women's College Research Institute, Women's College Hospital, Rheumatology, Toronto, CANADA.

Corresponding author: Sibel Zehra Aydin

1967 Riverside Drive, Ottawa, ON, K1H 7W9, Canada

Telephone number:6137388400

Fax number:6137388371

saydin@toh.ca

Running head: Accuracy of enthesal ultrasound

Abstract

Background: To investigate the relationship between physical examination (PE) and sonographic features of enthesitis, based on anatomical sites.

Methods: The analysis was done using merged raw data of 3 studies on 2298 entheses.

Results: Patients with clinical Achilles enthesitis had more abnormalities on ultrasound (hypoechoogenicity ($p < 0.001$), thickening ($p = 0.01$), Doppler ($p = 0.002$) and erosions ($p = 0.02$). The patellar tendon origin also correlated with PE but distal patellar tendon insertion and plantar aponeurosis were uncoupled from the ultrasound.

Conclusion: The relationship between clinical and sonographic findings for large entheses is dependent on the anatomical site and is best for the Achilles tendon and patellar tendon origin.

Introduction:

Enthesitis is a characteristic sign and hallmark of spondyloarthritis (SpA) and is clinically defined as pain or tenderness at the attachment site of a tendon/ligament to the bone with, or without, swelling. However, physical examination (PE) is neither sensitive nor specific for the evaluation of enthesitis (1). Ultrasonography (US) has been increasingly used for the assessment of enthesitis as it has the advantage of visualizing both soft tissue and bony changes (2,3). Comparison between US with PE has mostly been reported in relationship to the summation of total US scores with overall PE scores from multiple enthesal sites, rather than elementary lesions of enthesitis on US and per site (4-6). US is usually considered to be more sensitive than PE to detect enthesitis. However, it is also true whereby patients with clinical enthesal tenderness sometimes have no US feature of enthesitis (7,8). Unlike synovitis, it is not feasible to evaluate and validate sonographic or clinical enthesitis against the “gold standard” of tissue biopsy, so the relevance of clinical and imaging findings for enthesitis is difficult to disentangle.

Our hypothesis is that PE may be overrating enthesitis at certain sites whereas the link between US and PE can be better in others. There is limited information on the relationship between clinical and imaging findings from individual US lesions on multiple enthesal sites.

Materials and methods

The raw data of three previous studies were used for this analysis (8-10). All 3 studies were approved by 3 different ethic boards (Marmara University Ethics Board, No: 09.2014.0143, Leeds (East) REC 09/H1306/105; The University Health Network REB# is 08-0126-AE). The Marmara University Ethics Board was contacted and was declared that additional approval was not required for additional analysis by combining the raw data. The first study by Aydin et al comparing the enthesal differences in PsA, psoriasis and healthy controls (8), had one sonographer (SZA) and PE was performed by one investigator (ZRA) on the same day, blinded to each other's assessment. A Logiq-E9 (General Electric, Wauwatosa, Wisconsin, USA) was used to scan 12 entheses: quadriceps insertion, patellar tendon origin and insertion, Achilles tendon and plantar aponeurosis insertions and lateral epicondyle for the common extensor tendon origin, bilaterally. The elementary lesions defined by the Outcome Measures in Rheumatology (OMERACT) group have been used: hypoechogenicity, thickening, enthesal PD signals, erosions, enthesophytes and calcifications (11). All findings were graded between 0-3, quantitatively for thickening and erosions and semi-quantitatively for the other lesions to determine lesion severity on US (8).

For the second study by Arslan et al, comparing the differences between PsA and ankylosing spondylitis (AS) (9) the sonographer was the same (SZA) using the same methodology as the aforementioned study, except the US machine being an MyLab70-XVG (Esaote, Genoa, Italy), with a 6-18 Mhz linear transducer. One clinician performed the PE of the entheses on the same day as the US (FA), blinded to each other's assessment. For these 2 studies, only psoriasis or PsA data were extracted.

The third study by Polachek et al examined the association between sonographic enthesitis and the severity of radiographic features of damage in the peripheral and axial joints in patients with PsA (10). The US scans were done by one sonographer (LE) using a MyLab70-XVG (Esaote, Genoa, Italy) equipped with a 6–18 MHz linear transducer. Clinical assessment of the entheses was performed at the same day by the rheumatologist evaluating the patient. MADrid Sonographic Enthesitis Index (MASEI) scoring system was used in this study (12). Therefore, in addition, the triceps tendon insertion was also scanned, however this site was not analyzed as not being included in the previous studies. The same elementary lesions were investigated. There were some differences for the scoring of the severity of the lesions: Doppler signals and erosions were scored as 0 or 3 whereas hypoechogenicity and thickening were scored as 0 or 1.

Statistical analysis

All analysis was done per enthesal site. As there were some differences between the scoring methods, two types of analysis were performed, using the appropriate data:

- 1) The presence and absence of each sonographic elementary lesion was compared with findings on PE at the same enthesal site, by using all 3 data sources as this was captured by all.
- 2) The weighted analysis including the scoring of the findings were only performed by using the first 2 databases as scoring were done between 0-3 for all lesions.

The frequency of each elementary lesion on US was explored and presented as frequencies (percentages). The dependence between PE and US scores was assessed using chi-square or Fisher's exact tests, as appropriate. SPSS V-21 was used for analysis (SPSS Inc., Chicago, IL, USA).

Results:

A total of 2298 entheses from 377 patients (341 with psoriatic arthritis, 36 with psoriasis) were compared using US and PE.

The presence of elementary lesions:

Patients with clinical Achilles enthesitis had more frequent abnormalities on US (hypoechoogenicity: $p < 0.001$, thickening: $p = 0.01$, Doppler positivity: $p = 0.002$ and erosions: $p = 0.02$) (Table). Similarly, hypoechoogenicity ($p = 0.001$) and enthesophytes/calcifications ($p = 0.028$) at the patellar tendon origin were more common in patients with clinical enthesitis and there was a tendency for more erosions but it did not reach statistical significance ($p = 0.065$). The clinical quadriceps enthesitis was related to hypoechoogenicity on US ($p = 0.001$) and patients with clinical enthesitis on the lateral elbow had more frequent Doppler signals ($p = 0.007$). The rest of the enthesial sites were uncoupled from the US features, especially distal patellar tendon insertion and plantar aponeurosis (Table).

We have repeated the analysis by removing the psoriasis patients. Only focusing on PsA patients, the results were very similar with the whole group, with the exception of patellar tendon origin for calcifications (data not given).

The severity of elementary lesions on US:

Quadriceps tendon insertion: Clinical enthesitis was linked to severity of hypoechoogenicity ($p = 0.026$) and calcifications ($p = 0.020$) on US (Supplementary Table, Figure).

Patellar tendon origin: Patients with clinical enthesitis had more severe hypoechoogenicity ($p < 0.001$), thickening ($p < 0.001$), enthesophytes ($p < 0.001$) and calcifications ($p = 0.003$) on US.

Achilles enthesitis: Clinical enthesitis was associated with the severity of hypoechoogenicity ($p = 0.008$) and power Doppler ($p = 0.048$) on US.

Common extensor tendon origin: The severity of hypoechoogenicity ($p = 0.018$) and power Doppler ($p = 0.017$) was associated with clinical enthesitis (Supplementary Table, Figure).

Discussion:

The findings from this study confirm that the link between PE and US for enthesitis is dependent on the anatomical site. Patellar tendon origin and Achilles entheses are the 2 sites where PE is significantly linked to US findings, supporting the construct validity of US to visualize enthesitis.

The discrepancies between PE and US across different enthesial sites can be due to the difficulties to identify the entheses accurately by PE and different pain thresholds at different enthesial sites. For sites where PE is not linked to US such as the plantar aponeurosis or the patellar tendon insertion, US may improve the assessment by eliminating the false positives and negatives of PE. However, there may also be technical difficulties to detect some of the enthesial changes by US, such as the very low prevalence of Doppler signals in the plantar aponeurosis or US not being capable of detecting osteitis. Our group has previously reported the relationship between each elementary lesion with PE findings directly at the enthesial insertions using US on 21 patients with SpA, for the entheses around the knee

only (13). That study suggested that clinical enthesitis was associated with more hypoechogenicity and thickening on US. To the best of our knowledge, this is the first study on a large number of entheses and at multiple sites to explore the agreement between various sonographic features and PE to understand the clinical significance of individual lesions on US.

The enthesis is a very important structure in SpA, not only since it is frequently involved but also due to its significant impact on patients' pain, global assessment and quality of life. The recognition of enthesitis is important both at diagnosis and at follow up to decide the most appropriate treatment. However, the enthesis is probably the most difficult musculoskeletal structure to assess as the same sites are commonly affected by mechanical tendinopathies/enthesopathies and due to the proximity of fibromyalgia tender points (14). It is important to accurately assess the cause of pain at the enthesal insertions not to over or under-treat the patients.

The major strengths of this study are the large number of entheses and the representation of 2 experienced sonographers' in 3 different settings. As the same scoring method was not applied, it was not possible to include all patients to link the severity of the US features with PE but using the same definitions of elementary lesions, the presence/absence data were comparable. There were multiple clinical assessors for the 3rd study which may be considered as a limitation, however this perfectly reflects real-life experience and the assessors have been trained by the same individual.

In summary, the relationship between US and PE for enthesitis assessment depends on the enthesal site. US may be used to prove the presence of enthesal inflammation when their diagnostic uncertainty or when disease activity is not clear and/or therapies are considered. In the absence of a gold standard histological method, we believe that these findings provide a platform for the assessment of clinically relevant enthesitis. Future research should aim to confirm these findings and further validate the currently existing clinical scoring systems for enthesitis in SpA.

References

1. McGonagle D, Khan MA, Marzo-Ortega H, O'Connor P, Gibbon W, Emery P. Enthesitis in spondyloarthritis. *Curr Opin Rheumatol* 1999;11:244-50.
2. Kaeley GS, Eder L, Aydin SZ, Gutierrez M, Bakewell C. Enthesitis: A hallmark of psoriatic arthritis. *Semin Arthritis Rheum* 2018;48:35-43.
3. Gandjbakhch F, Terslev L, Joshua F, Wakefield RJ, Naredo E, D'Agostino MA. Ultrasound in the evaluation of enthesitis: status and perspectives. *Arthritis Res Ther* 2011;13:188.
4. Husic R, Gretler J, Felber A, Graninger WB, Duftner C, Hermann J, et al. Disparity between ultrasound and clinical findings in psoriatic arthritis. *Ann Rheum Dis* 2014;73:1529-36.
5. Bandinelli F, Prignano F, Bonciani D, Bartoli F, Collaku L, Candelieri A, et al. Ultrasound detects occult enthesal involvement in early psoriatic arthritis independently of clinical features and psoriasis severity. *Clin Exp Rheumatol* 2013;31:219-24.
6. Borman P, Koparal S, Babaoglu S, Bodur H. Ultrasound detection of enthesal insertions in the foot of patients with spondyloarthritis. *Clin Rheumatol* 2006;25:373-7.
7. Balint PV, Kane D, Wilson H, McInnes IB, Sturrock RD. Ultrasonography of enthesal insertions in the lower limb in spondyloarthritis. *Ann Rheum Dis* 2002;61:905-10.
8. Aydin SZ, Castillo-Gallego C, Ash ZR, Marzo-Ortega H, Emery P, Wakefield RJ, et al. Ultrasonographic assessment of nail in psoriatic disease shows a link between onychopathy and distal interphalangeal joint extensor tendon enthesopathy. *Dermatology* 2012;225:231-5.
9. Alhussain FA, Gunal EK, Kurum E, Bakirci Ureyen S, Ozturk AB, McGonagle D, et al. Greater magnitude of enthesal micro-damage and repair in Psoriatic Arthritis compared to Ankylosing Spondylitis on Ultrasound. *Rheumatology (Oxford)* 2019;58:299-303.
10. Polachek A, Cook R, Chandran V, Gladman DD, Eder L. The association between sonographic enthesitis and radiographic damage in psoriatic arthritis. *Arthritis Res Ther* 2017;19:189.
11. Terslev L, Naredo E, Iagnocco A, Balint PV, Wakefield RJ, Aegerter P, et al. Defining enthesitis in spondyloarthritis by ultrasound: results of a Delphi process and of a reliability reading exercise. *Arthritis Care Res (Hoboken)* 2014;66:741-8.
12. de Miguel E, Cobo T, Munoz-Fernandez S, Naredo E, Uson J, Acebes JC, et al. Validity of enthesal ultrasound assessment in spondyloarthritis. *Ann Rheum Dis* 2009;68:169-74.

13. Aydin SZ, Tan AL, Hodsgon R, Grainger A, Emery P, Wakefield RJ, et al. Comparison of ultrasonography and magnetic resonance imaging for the assessment of clinically defined knee enthesitis in spondyloarthritis. *Clin Exp Rheumatol* 2013; 31:933-6.
14. Roussou E, Ciurtin C. Clinical overlap between fibromyalgia tender points and enthesitis sites in patients with spondyloarthritis who present with inflammatory back pain. *Clin Exp Rheumatol* 2012;30:24-30.