

This is a repository copy of Ventilation CT : voxel-level comparison with HP gas MRI.

White Rose Research Online URL for this paper: http://eprints.whiterose.ac.uk/152330/

Version: Accepted Version

## **Proceedings Paper:**

Tahir, B. orcid.org/0000-0003-0531-3519, Marshall, H., Hughes, P. et al. (9 more authors) (2016) Ventilation CT : voxel-level comparison with HP gas MRI. In: European Respiratory Journal. ERS Lung Science Conference 2016, 10-13 Mar 2016, Estoril, Portugal. European Respiratory Society .

https://doi.org/10.1183/13993003.congress-2016.pa744

© 2016 The Authors. This is an author-produced version of a paper subsequently published in ERS International Congress 2016 abstracts. Uploaded in accordance with the publisher's self-archiving policy.

## Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

## Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/

## Ventilation CT: Voxel-level comparison with hyperpolarized Helium-3 & Xenon-129 MRI

Tahir BA<sup>1,2</sup>, Marshall H<sup>2</sup>, Hughes PJ<sup>2</sup>, Stewart NJ<sup>2</sup>, Horn FC<sup>2</sup>, Collier GJ<sup>2</sup>, Norquay G<sup>2</sup>, Hart KA<sup>1</sup>, Swinscoe JA<sup>1</sup>, Hatton MQ<sup>1</sup>, Wild JM<sup>2</sup> and Ireland RH<sup>1,2</sup>

<sup>1</sup>Academic Unit of Clinical Oncology and <sup>2</sup>Academic Radiology, University of Sheffield, UK

**Purpose:** To compare the spatial correlation of ventilation surrogates computed from inspiratory and expiratory breath-hold CT with hyperpolarized Helium-3 & Xenon-129 MRI in a cohort of lung cancer patients.

**Methods:** 5 patients underwent expiration & inspiration breath-hold CT. Xenon-129 & <sup>1</sup>H MRI were also acquired at the same inflation state as inspiratory CT. This was followed immediately by acquisition of Helium-3 & <sup>1</sup>H MRI in the same breath and at the same inflation state as inspiratory CT. Expiration CT was deformably registered to inspiration CT for calculation of ventilation CT from voxel-wise differences in Hounsfield units. Inspiration CT and the Xenon-129's corresponding anatomical <sup>1</sup>H MRI were registered to Helium-3 MRI via the same-breath anatomical <sup>1</sup>H MRI. This enabled direct comparison of CT ventilation with Helium-3 MRI & Xenon-129 MRI for the median values in corresponding regions of interest, ranging from finer to coarser in-plane dimensions of 10 by 10, 20 by 20, 30 by 30 and 40 by 40, located within the lungs as defined by the same-breath <sup>1</sup>H MRI lung mask. Spearman coefficients were used to assess voxel-level correlation.

**Results:** The median Spearman's coefficients of ventilation CT with Helium-3 & Xenon-129 MRI for ROIs of 10 by 10, 20 by 20, 30 by 30 and 40 by 40 were 0.52, 0.56, 0.60 and 0.68 and 0.40, 0.42, 0.52 and 0.70, respectively.

**Conclusion**: This work demonstrates a method of acquiring CT & hyperpolarized gas MRI (Helium-3 & Xenon-129 MRI) in similar breath-holds to enable direct spatial comparison of ventilation maps. Initial results show moderate correlation between ventilation CT & hyperpolarized gas MRI, improving for coarser regions which could be attributable to the inherent noise in CT intensity, non-ventilatory effects and registration errors at the voxel-level. Thus, it may be more beneficial to quantify ventilation at a more regional level.