



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/152334/>

Version: Accepted Version

Proceedings Paper:

Chan, H.-F., Parra-Robles, J., Collier, G. et al. (2017) Comparison of in-vivo lung morphometry models from multiple b-value diffusion-weighted MRI in healthy controls, IPF and COPD patients. In: European Respiratory Journal. ERS International Congress 2017, 09-13 Sep 2017, Milan, Italy. European Respiratory Society. ISSN: 0903-1936. EISSN: 1399-3003.

<https://doi.org/10.1183/1393003.congress-2017.oa4644>

This is an author-submitted, peer-reviewed version of an abstract that has been accepted for publication in the European Respiratory Journal, prior to copy-editing, formatting and typesetting. This version of the manuscript may not be duplicated or reproduced without prior permission from the copyright owner, the European Respiratory Society. The publisher is not responsible or liable for any errors or omissions in this version of the manuscript or in any version derived from it by any other parties. The final, copy-edited, published article, which is the version of record, is available without a subscription 18 months after the date of issue publication.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

3518

Comparison of in-vivo lung morphometry models from multiple b-value diffusion-weighted MRI in healthy controls, IPF and COPD patients

Morphology, Imaging

H. Chan^{1*}, **J. Parra-Robles**², **G. Collier**¹, **J. Wild**¹

¹

Academic Unit of Radiology, University of Sheffield - Sheffield

²

(United Kingdom), Department of Bioengineering, Universidad Carlos III de Madrid - Madrid (Spain)

Introduction

Two models of gas diffusion MRI, namely the cylinder model (CM) [Yablonskiy, D. A., et al. (2002). PNAS 99(5)] and the stretched exponential model (SEM) [Parra-Robles, J., et al. (2014). Proc ISMRM:3529], have been proposed for quantitative measurement of alveolar microstructural length scales (L_m for CM and L_{mD} for SEM) from multiple b-value hyperpolarised gas diffusion-weighted (DW) MRI.

Aims

Evaluate and compare estimates of lung morphometry from two models of

³

He gas diffusion in healthy controls, idiopathic pulmonary fibrosis (IPF) and chronic obstructive pulmonary disease (COPD) patients.


Methods

46 subjects underwent multiple b-value ³He DW-MRI: 19 healthy controls, 13 IPF and 14 COPD patients. Maps of L_m and L_{mD} values were generated

using the CM and SEM respectively, and mean values for each subject were derived.

Results

A statistically significant difference ($p < 0.001$) was obtained between healthy controls and patient groups for both mean L_m and L_{mD} values. L_{mD} and L_m were significantly correlated ($p < 0.001$) in a power relationship regression fit model ($R^2 = 0.957$) (Figure 1).



Conclusion

The non-linear relationship observed results from inherent differences in model assumptions. E.g. the CM is optimised for a limited range of alveolar diameters (< 400 Mm), whereas SEM does not make geometrical assumptions and is potentially valid over a wider range of alveolar sizes.