



This is a repository copy of *Thromboprophylaxis in lower limb immobilisation after injury (TiLLI)*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/153445/>

Version: Published Version

---

**Article:**

Horner, D., Goodacre, S. [orcid.org/0000-0003-0803-8444](https://orcid.org/0000-0003-0803-8444), Pandor, A. [orcid.org/0000-0003-2552-5260](https://orcid.org/0000-0003-2552-5260) et al. (6 more authors) (2019) Thromboprophylaxis in lower limb immobilisation after injury (TiLLI). *Emergency Medicine Journal*. 208944. pp. 1-6. ISSN 1472-0205

<https://doi.org/10.1136/emered-2019-208944>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC) licence. This licence allows you to remix, tweak, and build upon this work non-commercially, and any new works must also acknowledge the authors and be non-commercial. You don't have to license any derivative works on the same terms. More information and the full terms of the licence here:  
<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>



OPEN ACCESS

# Thromboprophylaxis in lower limb immobilisation after injury (TiLLI)

Daniel Horner <sup>1,2</sup>, Steve Goodacre <sup>2</sup>, Abdullah Pandor,<sup>2</sup> Timothy Nokes,<sup>3</sup> Jonathan Keenan,<sup>3</sup> Beverley Hunt,<sup>4</sup> Sarah Davis <sup>2</sup>, John W Stevens,<sup>5</sup> Kerstin Hogg<sup>6</sup>

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/emmermed-2019-208944>).

<sup>1</sup>Emergency Department, Salford Royal Hospitals NHS Trust, Salford, UK

<sup>2</sup>Centre for Urgent and Emergency Care Research (CURE), University of Sheffield, Sheffield, UK

<sup>3</sup>Departments of Haematology and Trauma/Orthopaedics, Plymouth Hospitals NHS Trust, Plymouth, UK

<sup>4</sup>Departments of Haematology and Rheumatology, Guy's & St Thomas's NHS Foundation Trust, London, UK

<sup>5</sup>Department of Health Economics and Decision Science, SCHARR, University of Sheffield, Sheffield, UK

<sup>6</sup>Department of Medicine, McMaster University, Hamilton, Ontario, Canada

## Correspondence to

Dr Daniel Horner, Salford Royal Hospitals NHS Trust, Salford M6 8HD, UK; [danielhorner@nhs.net](mailto:danielhorner@nhs.net)

Received 17 July 2019

Revised 10 October 2019

Accepted 17 October 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Horner D, Goodacre S, Pandor A, *et al.* *Emerg Med J* Epub ahead of print: [please include Day Month Year]. doi:10.1136/emmermed-2019-208944

## ABSTRACT

Venous thromboembolic disease is a major global cause of morbidity and mortality. An estimated 10 million episodes are diagnosed yearly; over half of these episodes are provoked by hospital admission/procedures and result in significant loss of disability adjusted life years. Temporary lower limb immobilisation after injury is a significant contributor to the overall burden of venous thromboembolism (VTE). Existing evidence suggests that pharmacological prophylaxis could reduce overall VTE event rates in these patients, but the proportional reduction of *symptomatic* events remains unclear. Recent studies have used different pharmacological agents, dosing regimens and outcome measures. Consequently, there is wide variation in thromboprophylaxis strategies, and international guidelines continue to offer conflicting advice for clinicians. In this review, we provide a summary of recent evidence assessing both the clinical and cost effectiveness of thromboprophylaxis in patients with temporary immobilisation after injury. We also examine the evidence supporting stratified thromboprophylaxis and the validity of widely used risk assessment methods.

## CASE 1

A 60-year-old woman attends your department after a fall downstairs, injuring her right ankle. X-rays confirm a Weber B fracture to the lateral malleolus without talar shift. There is no pain to the medial aspect. The orthopaedic plan is for below-knee plaster immobilisation in a non-weight-bearing cast, conservative management and outpatient review in 7–10 days' time. On further assessment, a body mass index (BMI) of 35 is apparent, along with a history of chronic obstructive pulmonary disease.

## CASE 2

A 25-year-old man attends your department after a crush injury to his left foot sustained at work. X-rays reveal undisplaced fourth and fifth metatarsal neck fractures. The orthopaedic team advise a walking boot and to allow partial weight bearing as tolerated until outpatient review later that week. On further assessment, the patient has no other medical history and no family history of venous thromboembolism (VTE). He takes no medications. Despite analgesia in the department, he remains unable to put any weight through the affected limb.

## QUESTION

Should these patients be offered thromboprophylaxis to reduce their short-term risk of VTE?

To answer that question, we will provide a short summary of the problem, followed by a review of the evidence and a practical guide to decision-making and intervention in these clinical scenarios.

## BACKGROUND

VTE is a significant global health burden, with incident events alone costing the UK an estimated £640 million and the USA an estimated \$7–10 billion each year.<sup>1,2</sup> Within the last decade, VTE has resulted in more deaths than prostate cancer, breast cancer, road traffic accidents and AIDS combined.<sup>3</sup>

Temporary immobilisation after injury accounts for approximately 2% of all VTE cases in registry data.<sup>4</sup> These cases are potentially preventable with early pharmacological thromboprophylaxis. However, it is unclear which patients will benefit from such intervention, which will be harmed, whether clinical outcomes are improved as a result of treatment and whether prophylaxis is cost-effective. As a result, international guidelines continue to recommend different management options; the UK advises routine assessment and individualised prescribing, for example, whereas US guidelines advise against thromboprophylaxis in this population.<sup>5,6</sup>

This issue was recently highlighted as a research priority for emergency medicine, through the James Lind Alliance Priority Setting Partnership in the UK.<sup>7</sup> While the debate continues, case report data and media coverage continue to highlight recurrent and tragic outcomes.<sup>8,9</sup> It remains unclear whether these tragedies were avoidable. In many regions, legal rulings and recommendations by the coroner/medical examiner delivered with the aim of avoiding future deaths have forced the clinical agenda without recourse to scientific evidence.<sup>10</sup> Practice remains variable with differing levels of engagement and awareness, despite the relative frequency of the problem. It is imperative that a cross-specialty consensus is reached on management of VTE risk, to ensure that clinicians and patients can be adequately informed and counselled on the merits of intervention.

This article summarises recent work on this topic and provides pragmatic guidance and decision support for clinicians.

## What is the risk of VTE following lower limb injury and temporary immobilisation?

A recent Cochrane review on this topic suggests wide variation in the incidence of VTE following lower

limb immobilisation without thromboprophylaxis, reporting a range between 4.3% and 40.0% within the literature.<sup>11</sup> These figures are based on the outcome of 'any VTE', including routine screening and detection of isolated distal deep vein thrombosis (DVT). There is ongoing debate regarding the clinical relevance of such a composite endpoint and the sensitivity of the contributing diagnostics.<sup>12-14</sup> Of more importance to clinicians and patients is the incidence of morbidity after immobilisation. This is more challenging to define but suggested by many to be the incidence of only clinically relevant or symptomatic VTE, warranting repeat hospital attendance and/or intervention.<sup>15</sup> Accurate estimates of this endpoint suffer from limitations in the injured population regarding subjectivity. When should a clinician or patient suspect VTE in a leg that is painful and swollen at baseline from injury? However, recent studies with transparent reporting of clinically relevant outcomes demonstrate a high level of consistency. The recent Cochrane review extracted data from six studies<sup>16-21</sup> involving just under 3000 randomised patients; symptomatic VTE occurred in 1.8%–5.5% of conservatively managed patients. These results suggest that without thromboprophylaxis, approximately 1 in every 50 patients will suffer a symptomatic VTE event (including symptomatic distal and proximal DVT and pulmonary embolism (PE)) following temporary immobilisation after injury. This is likely to be an underestimate given the range previously mentioned, the exclusion of high-risk patients from trial design and the general health equity of patients willing to engage with research.

### Is thromboprophylaxis clinically effective?

Can pharmacological thromboprophylaxis reduce this morbidity burden and, if so, by what margin? We performed a recently updated systematic review and network meta-analysis to compare any pharmacological intervention against control for this defined population.<sup>22,23</sup> This review included the Prevention of Thrombosis after Lower Leg Plaster Cast (POT-CAST) trial,<sup>21</sup> which has reignited debate on this important topic, and other recent randomised studies assessing the novel use of fondaparinux for this indication. We identified 13 trials with 6857 patients suitable for inclusion.<sup>16-21,24-30</sup> Interventions included prophylactic dose low molecular weight heparin (LMWH) or fondaparinux only. We examined four additional trials that had been excluded from the previous Cochrane review and one trial published subsequently. We found no trial evidence examining the use of direct oral anticoagulants (DOACs) in this situation. Risk of bias was present in all included studies. We found that both LMWH and fondaparinux significantly reduced the odds of clinically detected DVT, PE and any VTE. Intervention appeared to halve the risk of VTE across all classifications.<sup>23</sup>

It follows that the absolute 2% risk of clinically significant VTE following temporary immobilisation after injury can be

effectively halved to 1% with prophylactic dose anticoagulation. Although this may represent a potentially high number needed to treat, this value would depend primarily on baseline risk (which is likely to vary within the population) and also would be considered in the context of prevalence. Approximately 70 000 patients are immobilised and discharged from EDs in the UK alone every year.<sup>23</sup> Even a 1% absolute risk reduction in the VTE event rate in this cohort would therefore represent prevention of 700 clinically relevant VTE events. Assuming these events would be detected and treated as provoked VTE in the absence of thromboprophylaxis, they would incur 63 000 patient days of therapeutic dose anticoagulation, with accompanying costs and harms.

### What are the harms of thromboprophylaxis?

Clear definitions are now in place for the identification and classification of bleeding events in trials of anticoagulation and thromboprophylaxis.<sup>31-33</sup> Using these, our above systematic review found limited evidence of harm from prophylactic dose anticoagulation in this population. Only four major bleeding events were identified in total, with insufficient evidence of an increase within the LMWH group (OR 1.45, 95% CrI 0.08 to 32.17). Although these results are inconclusive given the low frequency of events, only four events within a trial sample of just under 7000 patients provide some reassurance on safety.

Minor bleeding as a secondary outcome varied across the population, being found in 0%–10.5% of patients treated with LMWH, in 0%–1.5% of patients treated with fondaparinux and in 0%–6.8% of the control groups. No cases of death attributable to VTE or intervention were identified within the study, and in those trials monitoring for the incidence of heparin-induced thrombocytopenia (HIT), no cases were detected. The most common adverse event of infection appeared equal between control and intervention groups when strictly defined.<sup>21</sup>

### Do the benefits of thromboprophylaxis justify the risks?

Thromboprophylaxis reduces the risk of thromboembolism but may increase the risk of bleeding. Our data were inconclusive on this latter aspect, but it remains one of the primary clinical concerns regarding the use of thromboprophylaxis. We undertook decision-analytic modelling to determine how these risks might compare in people undergoing lower limb immobilisation due to injury.<sup>34</sup> Table 1 summarises the results. The risk of fatal PE (which often drives decisions and recommendations regarding thromboprophylaxis) is very small, and the reduction in this risk associated with thromboprophylaxis is roughly matched by an increase in the risk of fatal bleeding or non-fatal intracranial haemorrhage. Overall, the risk of death or non-fatal intracranial haemorrhage is about 1 in 4000, whether thromboprophylaxis is

**Table 1** Predicted clinical outcomes per 100 000 patients with lower limb immobilisation due to injury

	Outcomes at 6 months per 100 000 patients							Outcomes at 5 years per 100 000 patients				
	Fatal PE	Fatal bleed	Non-fatal ICH	Other major bleed*	Non-fatal PE	Symptomatic DVT	Asymptomatic DVT	PTS	PE survivor with CTEPH	PE survivor without CTEPH	ICH survivor	Dead (any cause)
No prophylaxis	12	9	5	26	415	907	7052	1859	11	397	5	1133
Prophylaxis	7	12	8	35	225	492	3820	1007	6	215	7	1129

\*Patients having other major bleeds could also have a DVT or non-fatal PE.

CTEPH, chronic thromboembolic pulmonary hypertension; DVT, deep vein thrombosis; ICH, intracranial haemorrhage; PE, pulmonary embolism; PTS, post-thrombotic syndrome.

given or not. These are tragic events and, given the large population undergoing immobilisation, will occur with some frequency. However, our analysis suggests that thromboprophylaxis will not markedly change their incidence.

The main benefits of thromboprophylaxis lie in preventing non-fatal PE, symptomatic DVT and, in particular, asymptomatic DVT. Benefits for patients are assumed to accrue in the form of reduced long-term complications, particularly post-thrombotic syndrome. We estimated that, taking all effects into account, thromboprophylaxis results in 0.015 additional quality-adjusted life years (QALYs) per patient treated (95% credible interval (CrI) 0.004 to 0.029), indicating an overall benefit. However, it is uncertain whether a reduced risk of asymptomatic DVT leads to a reduced risk of post-thrombotic syndrome, so this finding should be treated with some caution.<sup>35</sup>

### Is thromboprophylaxis cost-effective?

Given the cost implications of prophylaxis, the low absolute event rate and the potential implications of any recommendations it is necessary to examine the cost effectiveness of this intervention. Our recent work for the Health Technology Assessment Programme examined this in detail.<sup>23</sup> We examined the wider literature to generate estimates of incidence for VTE, post-thrombotic syndrome, chronic thromboembolic pulmonary hypertension, intracranial haemorrhage, major bleeding and death among other potential outcomes. We subsequently assigned costs to these outcomes then designed a decision tree and used a Markov model to generate estimates of overall cost effectiveness.

Based on the mean costs and QALYs gained, we found an incremental cost-effectiveness ratio (ICER) for thromboprophylaxis compared with no thromboprophylaxis of £13 524. Based on standard thresholds used by the National Institute for Health and Care Excellence (NICE) in England, this value would fall below the ICER threshold of £20 000 and, as such, the intervention would broadly be considered as cost-effective.

We went on to examine the potential cost effectiveness of using a risk assessment method (RAM) to recommend stratified thromboprophylaxis. Several published RAMs were available with threshold prognostic accuracy data, to inform the modelling.<sup>36–37</sup> We found that a RAM could potentially improve the cost effectiveness of thromboprophylaxis, depending on the threshold score at which thromboprophylaxis would be given. The optimal threshold, assuming the RAM operated with diagnostic characteristics similar to the Leiden Thrombosis Risk in Plaster (cast) (L-TRiP(cast)) score, would result in a sensitivity of 84%–89% and a specificity of 46%–55% for predicting VTE.

### Can we safely risk stratify patients to maximise clinical and cost effectiveness?

Using risk factors to tailor thromboprophylaxis recommendations is a common strategy in hospitals.<sup>38</sup> There is face validity to the concept of risk prediction and tailored therapy in temporary lower limb immobilisation, which is strengthened by the low absolute risk and costs/inconvenience of treatment.<sup>15</sup>

We examined the literature to try and identify whether established individual VTE risk factors can predict the likelihood of subsequent disease in this ambulatory cohort of patients. A further systematic review was conducted, identifying 15 studies and 80 678 patients for inclusion. Meta-analysis of data was not possible due to significant variation between studies regarding data collection methods. All studies were deemed to be at moderate to severe risk of bias. We found advancing age and injury pattern to be consistently associated with increased VTE risk. BMI was the third most consistent individual risk highlighted, although overall results were conflicting. A total of 12 other risk factors analysed did not demonstrate any consistency in association with VTE across the dataset.<sup>39</sup> A subsequent publication supports our findings, reporting injury pattern, family history and BMI to be the individual risk factors most associated with VTE risk, among the POT CAST trial cohort.<sup>40</sup>

We conducted a further systematic review and engaged topic experts to try and identify available RAMs for use in this population. This work identified seven RAMs, three of which have undergone attempted external validation.<sup>36–41–46</sup> Variation and essential characteristics of the identified RAMs are presented in table 2. All studies looking to derive or externally validate a RAM were deemed to be at high risk of bias and did not report discrimination or calibration within an appropriate external population.

Prognostic accuracy measures for the three scores evaluated in two validation studies are presented in table 3. RAM sensitivity varies from 57.1% to 92.6% and specificity varies from 4.76% to 60.8%. As an ordinal score with additional validation data, The L-TRiP(cast) RAM is displayed in this table using thresholds denoting optimal performance and to allow direct comparison with other validated scores. The area under the receiver operating characteristic curve for the L-TRiP(cast) score ranged from 0.77 (95% CI 0.66 to 0.87) in the derivation cohort to 0.77 (95% CI 0.58 to 0.96) and 0.95 (95% CI 0.91 to 0.99) in the two subsequent validation cohorts. In addition, subsequent retrospective evaluation of the L-TRiP(cast) score within the POT-CAST cohort reports a lower area under the curve value of 0.69 (95% CI 0.58, 0.80).<sup>40</sup> As such, although the concept of a RAM is intuitive in this situation, the available supporting

**Table 2** Summary of design characteristics and threshold levels of identified RAMs

Risk assessment model	Acronym/descriptor	Derivation	Design	Incorporation of bleeding risk?	Number of variables	Threshold (suggested cut-point)	Attempted validation?
Roberts <i>et al</i> <sup>42</sup>	The GEMNet guideline	EC	Dichotomous	Yes	11	N/A	Yes
Keenan <i>et al</i> <sup>49</sup>	The Plymouth Rule	EC	Ordinal	No	14	>2	Yes
Nemeth <i>et al</i> <sup>36</sup>	The L-TRiP(cast) score	Regression	Ordinal	No	14	>8	Yes
Saragas <i>et al</i> <sup>41</sup>	The modified Caprini score	EC	Ordinal	No	36	>1	No
Eingartner <i>et al</i> <sup>45</sup>	N/A	EC	Ordinal	No	9	>1	No
Haque <i>et al</i> <sup>46</sup>	N/A	EC	Ordinal	No	14	>2	No
Giannadakis <i>et al</i> <sup>44</sup>	N/A	EC	Dichotomous	No	12	N/A	No

EC, Expert Consensus; GEMNet, Guidelines in Emergency Medicine Network; L-TRIP(cast), Leiden Thrombosis Risk in Plaster (cast); N/A, not applicable; RAM, risk assessment method.

**Table 3** Diagnostic performance of the L-TRiP(cast), GEMNet and Plymouth risk assessment models

Author	Roberts <i>et al</i>	Keenan <i>et al</i>	Nemeth <i>et al</i>	
Risk assessment model	GEMNet	Plymouth	L-TRiP(cast) with a cut-point of 8 or above	L-TRiP(cast) with a cut-point of 9 or above
Sensitivity (95% CI)	85.7% (62.6% to 96.2%)	57.1% (33.4% to 77.4%)	92.6%	80.8%
Specificity (95% CI)	4.76% (0.2% to 25.9%)	52.4% (30.3% to 73.6%)	39.7%	60.8%
Positive predictive value (95% CI)	47.4% (31.3% to 64.0%)	54.5% (32.7% to 74.9%)	3.8%	5.0%
Negative predictive value (95% CI)	25.0% (1.3% to 78.1%)	55.0% (32.0% to 76.2%)	99.5%	99.2%
Likelihood ratio positive (95% CI)	0.90 (0.73 to 1.10)	1.20 (0.67 to 2.15)	1.5	2.1
Likelihood ratio negative (95% CI)	3.00 (0.16 to 55.31)	0.81 (0.46 to 1.46)	0.2	0.3
Proportion receiving thromboprophylaxis (95% CI)	90.5% (76.5% to 96.9%)	52.4% (36.6% to 67.7%)	87.8%	74.7%

evidence is weak. Further RAMs (such as the Trauma, Immobilisation and Patients' Characteristics (TIP) score) continue to be published using Delphi consensus methodology and other derivation techniques, reporting improved performance metrics on internal validation.<sup>47</sup> All proposed RAMs remain in need of robust external validation.

### What are the key uncertainties and challenges moving forward?

Can I use a DOAC for this indication?

There is no trial evidence examining the use of DOAC medications in this clinical scenario. As such, there are no available data on the clinical or cost effectiveness of these drugs. Any such use would be off licence and approved as part of a trust protocol using local pharmacy and thrombosis committee governance structures.

Many trusts have taken this option and are starting to publish prospective datasets recording their experience.<sup>10 48</sup> Evaluation of this strategy with organisational support and careful oversight is to be encouraged and will provide estimates of effectiveness and safety that may support the case for future research.

Does this evidence apply to temporary splints as well as plaster casts?

The majority of research in this area focusses on immobilisation in plaster. As such, all estimates of effectiveness are not necessarily generalisable to other forms of immobilisation, which may allow partial weight bearing or less restriction in mobility. However, a caution should be noted here; often patients are treated with advice to partially weight bear when they clearly cannot do so. These patients remain at risk of VTE through immobility. NICE guideline NG89 currently defines lower limb immobilisation as 'any clinical decision taken to manage the affected limb in a way that would prevent normal weight-bearing status, or use of that limb, or both'.<sup>5</sup> Many patients who are managed with crutches and a splint would meet this definition and therefore justify risk assessment. This does not mandate prophylaxis but does recommend consideration of risk and shared decision-making (SDM).

Which RAM is best?

The evidence provided earlier highlights the lack of external validation data for any RAM. The Guidelines in Emergency Medicine Network (GEMNet), Plymouth rule, L-TRiP(cast) and

TIP score all appear to have limited supporting/validation data and variable strengths and weaknesses.<sup>36 42 47 49</sup> If using a RAM, clinicians should focus on consistency and serial audit to ensure appropriate use of the RAM, proportional use of thromboprophylaxis, patient outcomes and national benchmarking. For transparency and comparison, an example score sheet for each of the four RAMs previously mentioned can be found in online supplementary figures S1-S4.

Do these patients need blood tests prior to commencement of LMWH?

National guidelines from the British Committee for Standards in Haematology (BCSH) are clear that any patient receiving any form of heparin should have an estimate of baseline renal function and platelet count recorded.<sup>50</sup> These tests have a clear rationale: ensuring appropriate dosing and agent choice, absence of coagulopathy and recording baseline platelet count to clarify the degree of any future drop. These tests are equally necessary prior to commencement on any DOAC agent. The same BCSH guidance clarifies that adverse events such as HIT are rare enough with *prophylactic* dose LMWH that further routine platelet counts are unnecessary, and repeat testing should be based only on clinical concern.

How long should these patients remain on prophylactic anticoagulation and who should follow them up?

The vast majority of the trial data pertaining to these patients continued prophylactic anticoagulation for the duration of immobilisation in plaster and until return to baseline mobility. This often constitutes a period of 4–6 weeks. However, with the advent of newer immobilisation strategies and virtual fracture clinics, it is becoming clear that many patients are being encouraged to mobilise early after injury and often return to weight-bearing status and full mobility earlier than initially predicted. As such, it is vital that orthopaedic teams have an interest in this issue; decisions to remove a plaster cast and encourage mobilisation may facilitate earlier cessation of thromboprophylaxis. In addition, bleeding problems early in management may adjust the risk/benefit profile.

How do I use SDM in this context?

SDM involves the clinician offering options and describing their risks and benefits, the patient expressing his or her preferences and values, and then both jointly agreeing on a treatment decision. Emergency physicians appear supportive of this approach, and a number of tools have been developed, or are being developed, to support SDM in emergency care.<sup>51 52</sup> RAMs such as the Plymouth rule have been designed to allow potential completion by patients and to guide subsequent informed discussion. However, we are not aware of any published decision tools specifically incorporating SDM relevant to thromboprophylaxis in lower limb immobilisation. Data presented here could be used to support SDM and to develop such a tool.

### Cases: outcome

#### Case 1

This patient has both temporary and permanent risk factors for VTE. Her age and BMI are also of concern. She has a Plymouth score of 4 and an L-TRiP(cast) score of 10 and would meet GEMNet criteria for recommendation of thromboprophylaxis. As such, she should be counselled regarding her acutely increased risk of VTE. She should be informed that her absolute risk is likely to be higher than 2%.

Guidance regarding the clinical signs and symptoms of VTE should be provided in verbal and written form, and measures advised to mitigate risk (such as adequate hydration and mobilisation of the unaffected limb). The first and second authors would offer and prescribe prophylactic dose LMWH or a DOAC, respectively, up to the point of next orthopaedic review. The patient would be counselled on the minor risk of bleeding and informed that any treatment can be revisited at further orthopaedic follow-up, depending on clinical progress. A focal point of contact for immediate concern would be provided.

## Case 2

This patient has temporary but no clear permanent risk factors for VTE. He is also young and the immobilisation method chosen is less restrictive. He has a Plymouth score of 0 and an L-TRIP(cast) score of 3 and would not meet GEMNet criteria for recommendation of thromboprophylaxis. He should still be informed of a potential VTE risk, but this is likely to be an absolute risk lower than 2%.

Guidance regarding the clinical signs and symptoms of VTE should be provided in verbal and written form, and measures advised to mitigate risk (such as adequate hydration and mobilisation of the affected and unaffected limb). The first and second author would not routinely offer prophylactic anticoagulation for this clinical scenario, based on the low VTE risk and the limited application of the available data to non-rigid immobilisation. The patient should be informed that any signs or symptoms of VTE should be disclosed at further orthopaedic follow-up or earlier as required. A point of focal contact for immediate concern should be provided.

## SUMMARY

In patients with temporary lower limb immobilisation after trauma, the absolute risk of symptomatic VTE is low, at approximately 2%. Current evidence suggests that pharmacological prophylaxis can significantly reduce this risk. The benefits of thromboprophylaxis are achieved mainly through reduction of morbidity rather than lives saved. Pharmacological prophylaxis appears to be cost-effective.

Risk assessment can help inform SDM and individually tailor thromboprophylaxis, but there is limited evidence of external validation for any specific method at present. A key aspect of the risk assessment process is the sharing of information; clinicians must inform patients that there is an increased risk of VTE with temporary immobilisation and what the common presenting features are, even if the absolute risk is low.

**Twitter** Daniel Horner @rcemprof

**Acknowledgements** The authors thank all additional members of the core project group for HTA 15/187/06 for input and commentary throughout the work, including Shan Bennet and Robin Pierce-Williams for providing patient and public involvement. The authors also thank Mark Clowes for his assistance with the initial literature searches and article sourcing. We are also indebted to Joanne Hinde for assistance with logistics and administration.

**Contributors** DH and SG conceived the idea for this review and also acted as co-chief investigators for HTA 15/187/06. With regard to the wider project, AP was responsible for systematic review work; JWS was responsible for the statistical aspects; SD was responsible for the decision modelling and cost-effectiveness work; and TN, BH, KH and JK provided clinical expertise throughout the project. DH and SG were responsible for the drafting of this paper, although all authors provided comments on the drafts and read and approved the final version. DH is the guarantor for the paper.

**Funding** Much of the work cited in this review was funded by the UK National Institute for Health Research (NIHR) Health Technology Assessment Programme (project number 15/187/06). The views expressed in this review are those of the

authors and not necessarily those of the NIHR HTA Programme. Any errors are the responsibility of the authors. The funders had no role in the study design; collection, analysis and interpretation of data; writing of the manuscript; and decision to submit the manuscript for publication.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iDs

Daniel Horner <http://orcid.org/0000-0002-0400-2017>

Steve Goodacre <http://orcid.org/0000-0003-0803-8444>

Sarah Davis <http://orcid.org/0000-0002-6609-4287>

## REFERENCES

- House, Of Cet *et al.* *The prevention of venous thromboembolism in hospitalised patients - second report of session 2004-2005*. London: The Stationary Office, 2005.
- Grosse SD, Nelson RE, Nyarko KA, *et al.* The economic burden of incident venous thromboembolism in the United States: a review of estimated attributable healthcare costs. *Thromb Res* 2016;137:3–10.
- ISTH, Steering Cet *et al.* Thrombosis: a major contributor to the global disease burden. *J Thromb Haemost* 2014;12:1580–90.
- Bertoletti L, Righini M, Bounameaux H, *et al.* Acute venous thromboembolism after non-major orthopaedic surgery or post-traumatic limb immobilisation. *Thromb Haemost* 2011;105:739–41.
- NICE. Venous thromboembolism in over 16s: reducing the risk of hospital-acquired deep vein thrombosis or pulmonary embolism (NG89), 2018. Available: <https://www.nice.org.uk/guidance/ng89/chapter/Recommendations#terms-used-in-this-guideline> [Accessed 11 Jun 2019].
- Falck-Ytter Y, Francis CW, Johanson NA, *et al.* Prevention of VTE in orthopedic surgery patients: antithrombotic therapy and prevention of thrombosis, 9th ED: American College of chest physicians evidence-based clinical practice guidelines. *Chest* 2012;141:e278S–325.
- Smith J, Keating L, Flowerdew L, *et al.* An emergency medicine research priority setting partnership to establish the top 10 research priorities in emergency medicine. *Emerg Med J* 2017;34:454–6.
- Chen L, Soares D. Fatal pulmonary embolism following ankle fracture in a 17-year-old girl. *J Bone Joint Surg Br* 2006;88-B:400–1.
- Irish, Times. Coroner to highlight blood clot risks following sudden death of CO Tyrone church Minister Irishnews.com, 2019. Available: <https://www.irishnews.com/news/2018/08/08/news/coroner-to-highlight-blood-clot-risks-following-sudden-death-of-co-tyrone-church-minister-1402184/> [Accessed 11 Jun 2019].
- Macleod R, Roberts C. Personal communication. personal email regarding the use of DOAC medications as thromboprophylaxis for patients in lower limb plaster, as well as description of recent section 28 recommendation by coroner to avoid future deaths that risk assessment and thromboprophylaxis for this indication become mandatory 2017.
- Zee AAG, van Lieshout K, van der Heide M, *et al.* Low molecular weight heparin for prevention of venous thromboembolism in patients with lower-limb immobilization. *Cochrane Database Syst Rev* 2017;376.
- Horner D, Hogg K, Body R. Should we be looking for and treating isolated calf vein thrombosis? *Emerg Med J* 2016;33:431–7.
- Horner D, Hogg K, Body R, *et al.* The anticoagulation of calf thrombosis (act) project: results from the randomized controlled external pilot trial. *Chest* 2014;146:1468–77.
- Jones T, Horner D. Bet 2: striking the right balance: anticoagulation and isolated distal deep vein thrombosis. *Emerg Med J* 2017;34:334–6.
- Nemeth B, Cannegieter SC. Venous thrombosis following lower-leg cast immobilization and knee arthroscopy: from a population-based approach to individualized therapy. *Thromb Res* 2019;174:62–75.
- Bruntink MM, Groutars YME, Schipper IB, *et al.* Nadroparin or fondaparinux versus no thromboprophylaxis in patients immobilised in a below-knee plaster cast (protect): a randomised controlled trial. *Injury* 2017;48:936–40.
- Jørgensen PS, Warming T, Hansen K, *et al.* Low molecular weight heparin (Innohep) as thromboprophylaxis in outpatients with a plaster cast. *Thromb Res* 2002;105:477–80.
- Kock H-J, Schmit-Neuerburg KP, Hanke J, *et al.* Thromboprophylaxis with low-molecular-weight heparin in outpatients with plaster-cast immobilisation of the leg. *The Lancet* 1995;346:459–61.
- Lapidus LJ, Ponzer S, Elvin A, *et al.* Prolonged thromboprophylaxis with dalteparin during immobilization after ankle fracture surgery: a randomized placebo-controlled, double-blind study. *Acta Orthop* 2007;78:528–35.

- 20 Lassen MR, Borris LC, Nakov RL. Use of the low-molecular-weight heparin reviparin to prevent deep-vein thrombosis after leg injury requiring immobilization. *N Engl J Med* 2002;347:726–30.
- 21 van Adrichem RA, Nemeth B, Algra A, *et al.* Thromboprophylaxis after knee arthroscopy and Lower-Leg casting. *N Engl J Med* 2017;376:515–25.
- 22 Horner D, Stevens J, Pandor A, *et al.* Pharmacological thromboprophylaxis to prevent venous thromboembolism in patients with temporary lower limb immobilization after injury: systematic review and network meta-analysis. *J Thromb Haemost* 2019.
- 23 Pandor A, Horner D, Davis S, *et al.* Pharmacological thromboprophylaxis for lower limb immobilisation after injury: systematic review and economic evaluation. *Health Technology Assessment*. 2018.
- 24 Gehling H, Giannadakis K, Lefering R, *et al.* [Prospective randomized pilot study of ambulatory prevention of thromboembolism. 2 times 500 mg aspirin (ASS) vs. clivarin 1750 (NMH)]. *Unfallchirurg* 1998;101:42–9.
- 25 Goel D, Buckley R, deVries G, *et al.* Prophylaxis of deep-vein thrombosis in fractures below the knee: a prospective randomised controlled trial. *The Journal of Bone and Joint Surgery* 2009;91.
- 26 Kujath P, Spannagel U, Habscheid W. Incidence and prophylaxis of deep venous thrombosis in outpatients with injury of the lower limb. *Haemostasis* 1993;23 Suppl 1:20–6.
- 27 Lapidus LJ, Rosfors S, Ponzer S, *et al.* Prolonged thromboprophylaxis with dalteparin after surgical treatment of Achilles tendon rupture: a randomized, placebo-controlled study. *J Orthop Trauma* 2007;21:52–7.
- 28 Samama CM, Lecoules N, Kierzek G, *et al.* Comparison of fondaparinux with low-molecular-weight heparin for venous thromboembolism prevention in patients requiring rigid or semi-rigid immobilization for isolated non-surgical below-knee injury. *Annales Francaises de Medecine d'Urgence* 2014;4:153–66.
- 29 Selby R, Geerts WH, Kreder HJ, *et al.* A double-blind, randomized controlled trial of the prevention of clinically important venous thromboembolism after isolated lower leg fractures. *J Orthop Trauma* 2015;29:224–30.
- 30 Zheng X, Li D-Y, Wangyang Y, *et al.* Effect of chemical thromboprophylaxis on the rate of venous thromboembolism after treatment of foot and ankle fractures. *Foot Ankle Int* 2016;37:1218–24.
- 31 Kaatz S, Ahmad D, Spyropoulos AC, *et al.* Definition of clinically relevant non-major bleeding in studies of anticoagulants in atrial fibrillation and venous thromboembolic disease in non-surgical patients: communication from the SSC of the ISTH. *J Thromb Haemost* 2015;13:2119–26.
- 32 Schulman S, Angeras U, Bergqvist D, *et al.* Definition of major bleeding in clinical investigations of antithrombotic medicinal products in surgical patients. *J Thromb Haemost* 2010;8:202–4.
- 33 Schulman S, Kearon C, *et al.* Subcommittee on Control of Anticoagulation of the Scientific and Standardization Committee of the International Society on Thrombosis and Haemostasis. Definition of major bleeding in clinical investigations of antithrombotic medicinal products in non-surgical patients. *J Thromb Haemost* 2005;3:692–4.
- 34 Davis S, Goodacre S, Pandor A, *et al.* Decision-analysis modelling of the effects of thromboprophylaxis for people with lower limb immobilisation for injury. *Br J Haematol* 2019.
- 35 Kahn SR, Ginsberg JS. Relationship between deep venous thrombosis and the postthrombotic syndrome. *Arch Intern Med* 2004;164:17–26.
- 36 Nemeth B, Adrichem R, Aván, Hylckama Vlieg Aván, *et al.* Venous Thrombosis Risk after Cast Immobilization of the Lower Extremity: Derivation and Validation of a Clinical Prediction Score, L-TRIP(cast), in Three Population-Based Case–Control Studies. *PLoS Med* 2015;12:e1001899–e99.
- 37 Watson U, Hickey BA, Jones HM, *et al.* A critical evaluation of venous thromboembolism risk assessment models used in patients with lower limb cast immobilisation. *Foot and Ankle Surgery* 2016;22:191–5.
- 38 Caprini JA, Arcelus JI, Reyna J. Effective risk stratification of surgical and nonsurgical patients for venous thromboembolic disease. *Semin Hematol* 2001;38:12–19.
- 39 Horner D, Pandor A, Goodacre S, *et al.* Individual risk factors predictive of venous thromboembolism in patients with temporary lower limb immobilization due to injury: a systematic review. *J Thromb Haemost* 2019;17:329–44.
- 40 Nemeth B, van Adrichem R, Nelissen R, *et al.* Individualized thromboprophylaxis in patients with Lower-Leg cast Immobilization—A validation and subgroup analysis in the POT-CAST trial. *Thromb Haemost* 2019;119:1508–16.
- 41 Saragas NP, Ferrao PNF, Jacobson BF, *et al.* The benefit of pharmacological venous thromboprophylaxis in foot and ankle surgery. *S Afr Med J* 2017;107:327–30.
- 42 Roberts C, Horner D, Coleman G, *et al.* Guidelines in emergency medicine network (GEMNet): guideline for the use of thromboprophylaxis in ambulatory trauma patients requiring temporary limb immobilisation. *Emerg Med J* 2013;30:968–82.
- 43 Nokes TJCak J. Venous thromboembolism risk assessment V2. plymouth (Derriford Hospital) 2010.
- 44 Giannadakis K, Gehling H, Sitter H, *et al.* Is medicamentous thrombosis prophylaxis necessary in conservative outpatient therapy with lower limb injuries? *Unfallchirurg* 2000;103:475–8.
- 45 Eingartner C, Hontzsch D, Lang E, *et al.* Individualized prophylaxis for thromboembolism in outpatients – A practical way out between ‘ever’ and never? *Aktuelle Traumatologie* 1995;25:1–5.
- 46 Haque S, Bishnoi A, Khairandish H, *et al.* Thromboprophylaxis in ambulatory trauma patients with foot and ankle fractures: prospective study using a risk scoring system. *Foot Ankle Spec* 2016;9:388–93.
- 47 Douillet D, Nemeth B, Penalzoza A, *et al.* Venous thromboembolism risk stratification for patients with lower limb trauma and cast or brace immobilization. *PLoS One* 2019;14:e0217748.
- 48 Haque S, Davies MB. Oral thromboprophylaxis in patients with ankle fractures immobilized in a below the knee cast. *Foot and Ankle Surgery* 2015;21:266–8.
- 49 Keenan J, Hall MJ, Nokes TJC. A practical VTE risk assessment score tool for patients treated with lower limb cast immobilization. *Injury Extra* 2009;40:199–200.
- 50 Baglin T, Barrowcliffe TW, Cohen A, *et al.* Guidelines on the use and monitoring of heparin. *Br J Haematol* 2006;133:19–34.
- 51 Probst MA, Kanzaria HK, Frosch DL, *et al.* Perceived appropriateness of shared decision-making in the emergency department: a survey study. *Acad Emerg Med* 2016;23:375–81.
- 52 Melnick ER, Probst MA, Schoenfeld E, *et al.* Development and testing of shared decision making interventions for use in emergency care: a research agenda. *Acad Emerg Med* 2016;23:1346–53.