

This is a repository copy of *Integrated processing of sugarcane bagasse:Arabinoxylan extraction integrated with ethanol production*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/153613/>

Version: Accepted Version

---

**Article:**

Campbell, G. M., Čukelj Mustač, N., Alyassin, M. et al. (6 more authors) (2019) Integrated processing of sugarcane bagasse:Arabinoxylan extraction integrated with ethanol production. *Biochemical Engineering Journal*. pp. 31-40. ISSN 1873-295X

<https://doi.org/10.1016/j.bej.2019.03.001>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

Manuscript Number:

Title: Integrated processing of sugarcane bagasse: Arabinoxylan extraction integrated with ethanol production

Article Type: VSI: Bioprocess Development

Keywords: sugarcane bagasse, wheat bran, arabinoxylans, bioethanol, biorefinery integration

Corresponding Author: Professor Grant Murray Campbell, PhD

Corresponding Author's Institution: University of Huddersfield

First Author: Grant Murray Campbell, PhD

Order of Authors: Grant Murray Campbell, PhD; Nikolina Cukelj; Mohammad Alyassin; Leo Gomez; Rachael Simister; James Flint; Darren Phillips; Mark Gronnow; Nicholas Westwood

Abstract: A proof-of-concept project compared extraction of arabinoxylans (AX) from sugarcane bagasse and wheat bran via alkaline hydrogen peroxide followed by enzyme-assisted extraction with combinations of feruloyl esterases and a xylanase. Bagasse contains comparable amounts of AX to wheat bran, but with a much lower arabinoxylan substitution on the xylan backbone (A:X ratio of around 0.2 compared with 0.6 for wheat bran), hence offering AX products with distinctive functionality and potential end uses. In the current work, bagasse released its AX more readily than wheat bran, and released a wider range of molecular weights. Use of feruloyl esterase and xylanase enzymes on their own or following alkaline peroxide extraction did not enhance AX release substantially; however, the xylanase appeared to be effective at reducing the size of AX molecules, and there is scope to optimise the effects of enzymes to produce specific AX product fractions. As bagasse frequently arises within the context of bioethanol production, integration of AX extraction with ethanol production could allow economic production of a portfolio of AX products, as has been demonstrated in principle for AX co-production in a wheat ethanol plant.

Dr James Winterburn  
Guest Editor *Biochemical Engineering Journal*  
– “Bioprocesses Development” Special Issue

17<sup>th</sup> November 2018

Dear Dr Winterburn,

We are pleased to submit to the *Biochemical Engineering Journal* a manuscript entitled:

“Integrated processing of sugarcane bagasse:  
Arabinoxylan extraction integrated with ethanol production.”

for consideration for the upcoming Special Issue entitled “Bioprocess Development”. The work arose from a proof-of-concept project funded by the Lignocellulosic Biorefinery Network (LBNet). We are aware that the small scope of the project limits the findings; however, the findings are sufficiently robust to support the objectives and conclusions, while the paper contains additional value in terms of some specific technical outcomes as well the broader perspectives it illustrates that are well aligned with the Bioresources and Biorefinery Engineering theme and with the momentum of research in this area. We therefore hope that the manuscript will be considered well suited to publication in this Special Issue of the *Biochemical Engineering Journal*.

Declaration of interests: None.

Yours sincerely,



**Professor Grant Campbell**  
*Professor of Chemical Engineering*

Professor Grant Campbell PhD, CEng, CSci, FICHEM, FIFST, SFHEA  
Professor of Chemical Engineering, School of Applied Sciences. Tel: +44(0)1484 472 134; Email: grant.campbell@hud.ac.uk

Queensgate, Huddersfield, HD1 3DH, UK

+44 (0) 1484 422288 +44 (0) 1484 516151

Patron: HRH The Duke of York, KG

Vice-Chancellor: Professor Bob Cryan DL MBA DSc CEng FIET FHEA



**Integrated processing of sugarcane bagasse:  
Arabinoxylan extraction integrated with ethanol production.**

***Highlights***

- Bagasse released arabinoxylan more readily than bran, with a wider range of sizes
- Feruloyl esterase and xylanase enzymes could help to produce specific AX fractions
- Integration of AX extraction with ethanol production may allow economic production

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Integrated processing of sugarcane bagasse:  
Arabinoxylan extraction integrated with ethanol production.**

Campbell GM<sup>a\*</sup>, Čukelj N<sup>a1</sup>, Alyassin, M<sup>a</sup>, Gomez LD<sup>b</sup>, Simister R<sup>b</sup>, Flint J<sup>c</sup>, Philips DJ<sup>d</sup>, Gronnow MJ<sup>d</sup>  
and Westwood NJ<sup>e</sup>

a. School of Applied Sciences, University of Huddersfield, Queensgate, Huddersfield

b. Centre for Novel Agricultural Products, Department of Biology, University of York, Heslington,  
York, UK

c. College of Science, Joseph Banks Laboratories, University of Lincoln

d. Biorenewables Development Centre, 1 Hassacarr Close, Chessingham Park, Dunnington, York, UK

e. School of Chemistry and Biomedical Sciences, University of St Andrews

1. Currently at: Faculty of Food Technology and Biotechnology, University of Zagreb, Croatia

\* Author for correspondence, g.campbell@hud.ac.uk

**Abstract**

A proof-of-concept project compared extraction of arabinoxylans (AX) from sugarcane bagasse and wheat bran via alkaline hydrogen peroxide followed by enzyme-assisted extraction with combinations of feruloyl esterases and a xylanase. Bagasse contains comparable amounts of AX to wheat bran, but with a much lower arabinoxylan substitution on the xylan backbone (A:X ratio of around 0.2 compared with 0.6 for wheat bran), hence offering AX products with distinctive functionality and potential end uses. In the current work, bagasse released its AX more readily than wheat bran, and released a wider range of molecular weights. Use of feruloyl esterase and xylanase enzymes on their own or following alkaline peroxide extraction did not enhance AX release substantially; however, the xylanase appeared to be effective at reducing the size of AX molecules, and there is scope to optimise the effects of enzymes to produce specific AX product fractions. As bagasse frequently arises within the context of bioethanol production, integration of AX extraction with ethanol production could allow economic production of a portfolio of AX products, as has been demonstrated in principle for AX co-production in a wheat ethanol plant.

**Keywords:** sugarcane bagasse, wheat bran, arabinoxylans, bioethanol, biorefinery integration.

**Abbreviations**

AX	Arabinoxylan
UoH	University of Huddersfield
UoL	University of Lincoln
UoY	University of York
UoStA	University of St Andrews
BDC	Biorenewables Development Centre
LBNet	Lignocellulosic Biorefinery Network
P2PNet	Plants to Products Network
CE-High	High cut-off fraction (from ultrafiltration over 10 kDa) following chemical extraction

48	CE-Low	Low cut-off fraction (<10 kDa) following chemical extraction
49	EE-High	High cut-off fraction following enzyme-assisted extraction
50	EE-Low	Low cut-off fraction following enzyme-assisted extraction
51	XYL	$\beta$ -Xylanase ( <i>C. mixtus</i> , PRO-E0051, Prozomix UK)
52	FE-E0355	Feruloyl esterase ( <i>A. cellulolyticus</i> , PRO-E0355, Prozomix UK)
53	FE-E0356	Feruloyl esterase ( <i>A. cellulolyticus</i> , PRO-E0356, Prozomix UK)

54  
55 **Declarations of interest:** None.

## 56 57 58 **Introduction**

59  
60 For a reaction to take place, entities must come together and interact under the right conditions.  
61 The interactions leading to the current work arose from a workshop organized in September 2014 by  
62 the Lignocellulosic Biorefinery Network (LBNNet, <https://lb-net.net>), a Biotechnology and Biological  
63 Sciences Research Council Network in Industrial Biotechnology and Bioenergy (BBSRC NIBB). At this  
64 workshop, one of those who eventually formed a consortium asked of another, “If I want to use a  
65 source of lignin for a chemical conversion process I have developed, what lignin would make a good  
66 source?”, to which the other advised “One that already arises naturally within existing biorefineries,  
67 such that transportation costs are negated and the infrastructure and integration opportunities are  
68 in place to enable the lignin processing to be undertaken economically, and in which the raw  
69 material might already be partially degraded through earlier processing, to give more ready access  
70 to the lignin.” At a later point in the workshop, the participants were asked to write on pieces of  
71 paper what we would like to research, and place them on the floor. One of us wrote  
72 “Arabinoxylans”, based on a long-standing interest in this subject as a promising co-product of  
73 ethanol biorefineries, as the ethanol is used to precipitate the arabinoxylans, making the production  
74 of AX potentially economically viable in that context [1]. Meanwhile a representative of AB Sugar  
75 wrote “Something valuable from sugarcane bagasse (not furfural, we already do that)”. Seeing  
76 these notes together prompted the idea that arabinoxylans might similarly be extracted from  
77 bagasse in an integrated biorefinery producing ethanol from sugarcane, and that the residual  
78 bagasse following AX extraction might reveal a suitable source of lignin for further processing. A  
79 proof-of-concept project was constructed, to demonstrate the feasibility of AX extraction from  
80 sugarcane bagasse, and to examine the residual lignin, following partial deconstruction during the  
81 AX extraction process, for its suitability as a feedstock for conversion into a phenolic monomer using  
82 an established reaction sequence [2]. A parallel activity studying wheat bran was included, to  
83 extend the scope and make the findings more immediately applicable to the UK context (where  
84 bioethanol production is largely from wheat). Enzyme enhancement of the AX release, a scale-up  
85 component and a techno-economic analysis of AX production were also included in the project, to  
86 give a consortium comprising the Universities of Huddersfield, St Andrews, York, Lincoln and  
87 Nottingham along with the Biorenewables Development Centre in York  
88 (<http://www.biorenewables.org/>). Later a student project supported by the Plants to Products  
89 Network (another BBSRC NIBB, <http://www.nibbp2p.org>) extended the project by undertaking a  
90 bioethanol pinch analysis, following the approaches of Martinez et al. [3], to minimise ethanol usage  
91 while producing a range of AX products including arabinoxylan-oligosaccharides (AXOS) [4].

92  
93 This paper describes the proof-of-concept work undertaken to demonstrate AX extraction from  
94 sugarcane bagasse and to compare it with extraction from wheat bran in terms of yield, composition

95 and responsiveness of the extraction process to enzyme enhancement. A future paper will describe  
1 96 the investigation of the residual bagasse following AX extraction, in terms of the nature of its lignin  
2 97 and its suitability for further processing, including further integration opportunities with ethanol and  
3 98 butanol.  
4 99

5 100 Global sugarcane production is around 1.9 billion tonnes per annum  
6 101 (<https://www.statista.com/statistics/249604/sugar-cane-production-worldwide/>), resulting in  
7 102 around 570 million tonnes wet bagasse or half this amount if dried. Sugarcane bagasse thus  
8 103 represents a major waste stream arising from sugar and alcohol industries, typically containing  
9 104 around 40-50% cellulose, 25-35% hemicelluloses (predominantly xylans) and 20-30% lignin [5,6]. A  
10 105 focus of previous work has been to deploy feruloyl esterases and xylanases to assist the  
11 106 saccharification of bagasse to increase recovery of fermentable sugars, by removing the  
12 107 hemicelluloses that (to put it simply) link lignin and cellulose, thus increasing accessibility of the  
13 108 latter to cellulases and hence the release of glucose [7]. A difference in the current work is that the  
14 109 intention was not to hydrolyse arabinoxylan hemicelluloses to their constituent sugars, but rather to  
15 110 release and recover them as intact large AX molecules, in which form they have potential as  
16 111 functional food ingredients and non-food products [8,9]. The context of the ethanol biorefinery  
17 112 gives scope for AX co-production to be economic as a result of integration with ethanol production  
18 113 (used for precipitating the AX), as has been shown previously for AX production from wheat bran in  
19 114 a wheat ethanol biorefinery [1]. Wheat bran typically contains 20-30% AX [10,11], similar to  
20 115 bagasse, but with a much higher ratio of arabinose to xylose units; in wheat bran the A:X ratio is  
21 116 typically in the range 0.5-0.6 [12,13]), while in sugarcane bagasse it is much lower at around 0.2  
22 117 typically [14]. This “cleaner” xylan backbone with fewer arabinose substitutions is likely to exhibit  
23 118 different functional properties compared with wheat bran AX, including reduced solubility and  
24 119 greater susceptibility to enzyme action, as well as effects on viscosity and gel formation, and  
25 120 performance in food products or animal feed.  
26 121  
27 122

## 28 123 **Materials and Methods**

29 124  
30 125 Sugarcane bagasse (25 kg) and wheat bran (50 kg) were sourced commercially by AB Sugar and  
31 126 provided to the project. The bagasse and bran were milled at the Biorenewables Development  
32 127 Centre (BDC) using a Retsch Cutting Mill SM 300 (Retsch GmbH, Germany) with a 2.00 mm screen,  
33 128 and AX extraction studies undertaken at the University of Huddersfield (UoH). Proximate analysis  
34 129 and arabinoxylan (AX) content measurements were performed at the University of York (UoY).  
35 130 Lignin studies, to be presented in a future paper, were undertaken at the University of St Andrews  
36 131 (UoStA).  
37 132

38 133 Figure 1 describes the chemical extraction process using alkaline hydrogen peroxide applied to the  
39 134 milled wheat bran and sugarcane bagasse, based on the work of Hollmann and Lindhauer [15] and  
40 135 Du et al. [16]. The residue material was then subjected to enzyme treatment to see if further  
41 136 release of AX resulted, using selected combinations of  $\beta$ -Xylanase (*C. mixtus*, PRO-E0051, referred to  
42 137 here as XYL) and two types of Feruloyl esterase (*A. cellulolyticus*, PRO-E0355 and PRO-E0356,  
43 138 referred to here as FE-E0355 and FE-E0356) from Prozomix UK. Enzyme-assisted extraction on its  
44 139 own was also investigated, along with chemical extraction followed by further extraction with buffer  
45 140 solution, and buffer extraction on its own.  
46 141  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

142 In total eight extractions were performed for each feedstock:

- 1 143 1. Direct buffer extraction (control for all trials, particularly Trials 7-8);
- 2 144 2. Chemical extraction followed by buffer extraction (control for Trials 3-6);
- 3 145 3. Chemical extraction followed by FE-E0355 and XYL;
- 4 146 4. Chemical extraction followed by FE-E0356 and XYL;
- 5 147 5. Chemical extraction followed by FE-E0355;
- 6 148 6. Chemical extraction followed by FE-E0356;
- 7 149 7. Direct enzyme extraction with FE-E0355 and XYL;
- 8 150 8. Direct enzyme extraction with FE-E0356 and XYL.

9 151  
10 152 The supernatant following treatment, centrifugation and filtering was separated by ultrafiltration  
11 153 over a 10 kDa membrane (see below for details). The chemical extraction thus yielded High cut-off  
12 154 (CE-High) and Low cut-off (CE-Low) fractions and a pellet of residual material; enzyme extractions  
13 155 directly on the raw material or on the pellet following chemical extraction similarly yielded High cut-  
14 156 off (EE-High) and Low cut-off (EE-Low) fractions and a pellet. A single chemical extraction was  
15 157 performed to produce residual material for subsequent enzyme extraction in Trials 3-6; a second  
16 158 chemical extraction was performed for subsequent buffer extraction (Trial 2).

17 159  
18 160 For the chemical extraction, 50 g bran or 11 g bagasse (because of the lower bulk density of the  
19 161 latter restricting the amount that could be processed in a bottle) was weighed into a 1 L Duran  
20 162 bottle. Foaming is a problem during AX extraction, so 15 drops of anti-foaming agent (Dimeticon  
21 163 SILFAR® SE 4, Wacker Chemie AG, Germany) were added, then 400 mL of 2% hydrogen peroxide  
22 164 (Fisher Scientific UK Limited, analytical grade) was added very slowly, with stirring with a magnetic  
23 165 stirrer. 5 more drops of anti-foaming agent were added, followed by the final 100 mL of hydrogen  
24 166 peroxide solution. The pH was adjusted to 11.5 with 50% NaOH. The bottle was placed in a 50°C  
25 167 water bath and the solution stirred for 4 h. The pH was controlled every hour and readjusted to 11.5  
26 168 if needed. After cooling to room temperature the pH of the mixture was adjusted to 7 with  
27 169 concentrated sulphuric acid. The solution was centrifuged for 15 minutes at 4000 rpm using a  
28 170 Beckman GS-6S centrifuge (Beckman Coulter Life Sciences, USA). The supernatant was filtered and  
29 171 the remaining solids washed with 150 mL of water, then centrifuged, filtered, washed with 150 mL  
30 172 water and centrifuged again. The final supernatant was filtered and pooled with the previous two.  
31 173 The residual solid (the pellet) was placed in the oven to dry overnight at 50°C.

32 174  
33 175 The high molecular weight material in the supernatant was concentrated by ultrafiltration using a  
34 176 Vivaflow™ 200 system (Sartorius Stedim Biotech GmbH, Germany) with polyethersulfone  
35 177 membranes with a molecular weight cut-off of 10 kDa. (Ultrafiltration would be done as part of a  
36 178 commercial process to reduce the amount of ethanol needed subsequently to precipitate the AX  
37 179 [1,15]; in the current work the fractions were not precipitated. The ultrafiltration served to separate  
38 180 the released AX into larger MW and smaller MW fractions.) Retentate was recycled until the volume  
39 181 was reduced to one fifth of the original. The retentate (High cut-off) and permeate (Low cut-off)  
40 182 fractions were freeze-dried using a Christ Freeze Dryer Alpha 1-4 LDplus, (Martin Christ  
41 183 Gefriertrocknungsanlagen GmbH, Germany) at -47.8°C and 0.35 mbar. Samples of freeze-dried High  
42 184 cut-off and Low cut-off material and oven-dried pellets were sent to UoY for AX analysis (see below).  
43 185 Samples of the dried pellet material were also sent to the University of St Andrews (UoStA) for  
44 186 assessment of its lignin, to be described in a future paper.

45 187

46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



188 For enzyme-assisted extraction trials on raw wheat bran or sugarcane bagasse or on pellets after  
1 189 chemical extraction, material (30 g for the wheat bran and 6 g for the sugarcane bagasse) was  
2 190 weighed into a 1 L Duran bottle with a magnetic stirrer. 300 mL of a buffer (composed of 107.4 mL  
3 191 of 0.2 M disodium phosphate, 42.6 mL of citric acid and 150 mL of water) at pH 6.5 was added, the  
4 192 bottles placed in a 37°C water bath and the pH of the solution readjusted to 6.5 with citric acid when  
5 193 the temperature reached 37°C. The different enzymes were added at a level of 10 µg/10 g for the  
6 194 XYL and FE-E0355 and 5µg/10 g for the FE-E0356, and the solution was stirred for 2 h. As above,  
7 195 samples were centrifuged, filtered and washed, in this case using 100 mL of water for each washing,  
8 196 and the residual solid pellet oven-dried overnight at 50°C. Again the supernatant was passed  
9 197 through ultrafiltration over 10 kDa and the retentate (High cut-off, EE-H) and permeate (Low cut-off,  
10 198 EE-L) freeze dried and sent to UoY for analysis along with the oven-dried pellet, with pellet samples  
11 199 also sent to UoStA for lignin analysis.

16 200  
17 201 Due to the limited scope of this small proof-of-concept project, replicate enzyme extractions were  
18 202 not performed, the aim being to demonstrate broad effects in relation to AX yields from wheat bran  
19 203 and bagasse and the potential effects of enzyme-assisted extraction. Five replicate chemical  
20 204 extractions were performed, as enzyme extraction Trials 3-6 and buffer Trial 2 each required a  
21 205 chemical extraction first, although High and Low cut-off fractions were subsequently produced for  
22 206 only two of these (2 and 3), and only these two pellets were sent for lignin analysis. Subsequent  
23 207 similar work with replication has confirmed the broad trends reported here and shown sufficient  
24 208 reproducibility to have confidence in the trends, which are reported and discussed here within the  
25 209 limits of the acknowledged lack of replication.

26 210  
27 211 Proximate analysis of the wheat bran and sugar cane bagasse was undertaken as follows:

#### 28 212 *Lignin determination: acetyl bromide method*

29 213 Biomass powder was weighed out (4 mg) into 2 mL tubes. The biomass was heated at 50°C for 3  
30 214 hours after adding 250 µL of acetyl bromide solution (25% acetyl bromide and 75% glacial acetic acid  
31 215 by volume) and vortexing every 15 minutes. After the samples were cooled to room temperature,  
32 216 the contents were transferred into 5 mL volumetric flasks. A further 1 mL of NaOH (2 mol L<sup>-1</sup>) was  
33 217 used to rinse the tubes pouring the NaOH into the 5 mL flasks. 175 µL of hydroxylamine HCl (0.5 mol  
34 218 L<sup>-1</sup>) was added to the volumetric flasks and, after vortexing, the latter were filled up to 5 mL with  
35 219 glacial acetic acid and mixed several times. Finally, in order to measure the 280 nm UV adsorption  
36 220 by spectrophotometer, 100 µL of each sample was diluted in 900 µL of glacial acetic acid. The  
37 221 amount of lignin was calculated using the following formula: [absorbance/(coefficient pathlength)] ·  
38 222 [(total volume · 100%)/biomass weight], where coefficient = 15.69, pathlength = 1, total volume = 5,  
39 223 biomass weight = 4.

#### 40 224 *Non-cellulosic monosaccharide determination*

41 225 Following the method of Fry [17], biomass dry powder (4 mg) was partially hydrolyzed by adding 0.5  
42 226 mL of trifluoroacetic acid (TFA, 2 mol L<sup>-1</sup>). Then, the vials were flushed with dry argon, mixed and  
43 227 heated at 100°C for 4 hours, mixing periodically. The vials were then cooled to room temperature  
44 228 and dried in centrifugal evaporator with fume extraction overnight. The pellets were washed twice  
45 229 with 500 µL of 2-propanol and vacuum dried. Finally, the samples were resuspended in 200 µL of  
46 230 deionised water, filtered with 0.45 µm PTFE filters, and analyzed by HPAEC (see below).

47 231  
48 232  
49 233  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

234 It became evident that these hydrolysis conditions were inadequate to release all the AX in the  
1 235 original wheat bran (see below, where the mass balance indicates more AX in the extracted fractions  
2 236 than appeared to be present initially in the bran). The wheat bran was therefore hydrolysed under a  
3 237 range of conditions, to investigate the effects on AX measurement and to draw conclusions about  
4 238 the most appropriate conditions for AX analysis in wheat bran. The bran samples were hydrolysed in  
5 239 4M TFA (2 mL per 4 mg bran) for 1, 2 and 4 hours, and in 2M (2 mL per 4 mg bran) for 4 and 6 hours  
6 240 at 120°C. In addition, samples were pretreated in 98% TFA for 1, 2, 4 and 6 hours at room  
7 241 temperature, then diluted to 4M and boiled for 1 hour at 120°C.  
8 242

#### 12 243 *Crystalline cellulose*

13 244 Biomass dry pellets after TFA hydrolysis were washed once with 1.5 mL of water, and twice using 1.5  
14 245 mL of acetone. The dried pellets were left to air dry overnight before complete hydrolysis by adding  
15 246 90 µL of 72% w/w sulphuric acid, incubating at room temperature for 4 hours. 1.89 mL of water was  
16 247 subsequently added and the sample was heated for 4 hours at 120°C. The glucose content of the  
17 248 supernatant was assessed using the colorimetric Anthrone assay, using a glucose standard curve.  
18 249

19 250 Analysis of the sugar compositions of the wheat bran and bagasse samples and of fractions and  
20 251 residues following the various chemical and enzyme extractions was undertaken as follows:  
21 252

22 253 Monosaccharide analysis was performed by high performance anion-exchange chromatography  
23 254 (HPAEC) (Dionex IC 2500) on a Dionex CarboPac PA-10 column with integrated amperometry  
24 255 detection [18]. The separated monosaccharides were quantified using external calibration with an  
25 256 equimolar mixture of nine monosaccharide standards (arabinose, fucose, galactose, galacturonic  
26 257 acid, glucose, glucuronic acid, mannose, rhamnose, and xylose), which were subjected to TFA  
27 258 hydrolysis in parallel with the samples.  
28 259

## 36 260 **Results and Discussion**

37 261  
38 262  
39 263 Table 1 shows the compositions of the wheat bran and sugarcane bagasse used in the current work.  
40 264 The wheat bran had a significant starch content; therefore for the scale-up work undertaken at BDC,  
41 265 the wheat bran was washed with water to remove starch prior to extraction, in line with the  
42 266 recommendation of Du et al. [16]. The bagasse had 28% lignin and nearly 20% AX, suggesting it was  
43 267 a promising candidate for recovery of both materials. The analytical procedure used indicated 10%  
44 268 lignin in the wheat bran, although subsequent NMR work suggested a much lower lignin content, in  
45 269 line with other recent reports that wheat bran contains less lignin than previously thought. These  
46 270 results also suggest an AX content of only 8.64% in the wheat bran; this figure is lower than the 20-  
47 271 30% generally expected for wheat bran, and later proved to be incompatible with the mass balance  
48 272 for AX recovered in the various fractions and residues, which suggested an AX content in the original  
49 273 material of around 24% (see below). The A:X ratio for the wheat bran was 0.57, and for the bagasse  
50 274 0.21, in line with typical values expected from the literature, and showing the much “cleaner” xylan  
51 275 backbone for the bagasse AX, with fewer arabinose substitutions compared with the wheat AX.  
52 276

53 277 Tables 2 and 3 report the crude yields, AX concentrations and hence AX yields from 50 g wheat bran  
54 278 and 11 g sugarcane bagasse, respectively, in the starting materials and in the fractions following the  
55 279 various chemical, enzyme and buffer treatments. The second column in each table is for chemical  
56 280 extraction only, showing the data from two replicates, with good agreement. Considering the wheat  
57  
58  
59  
60  
61  
62  
63  
64  
65

281 bran results first, chemical extraction of around 50.6 g of bran (moisture content 9.5%) yielded a  
1 282 residual dry pellet weighing 29.94 g (averaged from Trials 2 and 3), a High cut-off (CE-High) of 8.21 g,  
2 283 and a Low cut-off (CE-Low) of 4.40 g (total 42.55 g, c.f. about 45 g solids in the original material).  
3 284 The AX concentration in the pellet was 35.81% and in the High and Low cut-off fractions was 16.32%  
4 285 and 1.25%, respectively. This implies a total amount of AX in the pellet and two fractions of  
5 286  $29.94 \times 35.81\% + 8.21 \times 16.32\% + 4.40 \times 1.25\% = 10.72 + 1.34 + 0.06 = 12.12$  g. The mass balance  
6 287 therefore implies an AX concentration of  $12.12/50.6 = 24\%$  in the original wheat bran, higher than  
7 288 the 8.64% reported in Table 1, and more in line with the expected AX content of wheat bran,  
8 289 suggesting that the 8.64% figure is erroneous.

290  
13 291 It is well known that the appropriate hydrolysis conditions for this sort of analysis depend on the  
14 292 nature of the sample, with different samples requiring different combinations of time, temperature  
15 293 and acidity to get an optimum balance between release and degradation of monosaccharides  
16 294 [19,20]. The bran sample was therefore reanalysed under a range of hydrolysis conditions as  
17 295 described above. Figure 2 shows the AX contents calculated for each of the hydrolysis regimes.  
18 296 Under similar conditions to the original analysis (2M for 4 hours, but at a higher temperature,  
19 297 120°C), the measurement was similar at 8.79%, while 2M for 6 hours released more AX to give  
20 298 14.04%. 4M for 1 or 2 hours released even more (18.58-19.71%), but 4M for 4 hours appears to give  
21 299 substantial degradation and a final measurement of only 5.50%. Pretreatment in concentrated acid  
22 300 appeared effective at releasing more AX for effective hydrolysis, giving measurements of around  
23 301 26.7% after 1 or 2 hours of pretreatment, decreasing to 22.8 and 20.6% after 4 and 6 hours,  
24 302 suggesting degradation at the high acid concentration despite the low temperature. Overall, these  
25 303 results demonstrate that an AX content of around 24%, as implied by the mass balance, is plausible,  
26 304 but that measuring the AX content of raw wheat bran requires different hydrolysis conditions  
27 305 compared to extracts or the residual pellet following extraction.

28 306  
29 307 Thus, of a total of around 12 g AX in the original 50 sample, 1.4 g or 11.5% was released by the  
30 308 alkaline hydrogen peroxide extraction process. This is much lower than the recoveries of 46-50%  
31 309 reported by Hollmann and Lindhauer [15] and Du et al. [16] for alkali-extracted AX from wheat bran  
32 310 using similar conditions. In those studies the wheat bran was boiled in 70% ethanol at 80°C for 4  
33 311 hours prior to alkaline H<sub>2</sub>O<sub>2</sub> treatment, which Hollmann and Lindhauer [15] advised was necessary to  
34 312 achieve high yields; this step was omitted in the current work (for safety and cost reasons in relation  
35 313 to the planned scale-up work, and because related unpublished work from our labs on extraction  
36 314 from maize meal had found that this step was not needed, as also confirmed by work from Doner  
37 315 and Hicks [21] on AX extraction from maize fibre). The omission of this ethanol boiling step is  
38 316 possibly the reason for the lower yields than in this previously reported work.

39 317  
40 318 Considering the sugarcane bagasse results, chemical extraction of around 11.5 g of bagasse  
41 319 (moisture content 9.3%) yielded a residual dry pellet weighing 5.43 g (averaged from Trials 2 and 3),  
42 320 a High cut-off (CE-High) of 3.11 g, and a Low cut-off (CE-Low) of 8.70 g. The total appears to be  
43 321 17.34 g, c.f. about 10.5 g solids in the original material; the mass balance does not give good  
44 322 agreement in this case. This is probably because the Low cut-off was extrapolated from the solids  
45 323 left after freeze-drying dilute samples (and the freeze-dried samples may not have been completely  
46 324 dry), such that the 8.70 g figure is not accurate, while overall the mass balance from just 11 g of  
47 325 bagasse is inherently less accurate than that from 50 g wheat bran, and the contribution from salts  
48 326 formed on neutralisation relatively greater.

49 327

50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

328 The AX concentration in the pellet was 14.02% (averaged from Trials 2 and 3, although higher in  
1 329 Trials 4-6) and in the High and Low cut-off fractions was 23.64% and 7.23%, respectively. This  
2 330 implies a total amount of AX in the pellet and two fractions of  $5.53 \times 14.02\% + 3.11 \times 23.64\% +$   
3 331  $8.70 \times 7.23\% = 0.775 + 0.735 + 0.629 = 2.139$  g, compared with  $11.54 \times 19.58\% = 2.260$  g of AX in the  
4 332 original sample. This mass balance appears reasonable, although the uncertainty over the Low cut-  
5 333 off contribution is acknowledged. In this case it appears that the AX left in the pellet (assuming this  
6 334 to be a more accurate figure) is about one-third of the AX in the original material ( $0.775/2.25 = 34\%$ ),  
7 335 implying nearly two-thirds has been released, compared with only 11.5% for the wheat bran. This  
8 336 suggests sugarcane bagasse may be very amenable to AX extraction, as it appears to yield its AX  
9 337 more readily than wheat bran. The readiness of the bagasse to release its AX compared with wheat  
10 338 bran suggests the expensive and hazardous ethanol boiling step advised by Hollmann and Lindhauer  
11 339 [15] for wheat bran may not be needed in a bagasse-based AX extraction process. As noted above,  
12 340 Doner and Hicks [21] advised that dewaxing with toluene-ethanol was unnecessary for AX extraction  
13 341 from maize fibre.  
14 342

15 343 Figure 3 shows the crude yields of High and Low cut-off material following the various wheat bran  
16 344 extractions, the AX concentrations in the fractions and hence the absolute yields of AX in each  
17 345 fraction. Trial 1 shows the recovery of High and Low fractions from just extraction with Buffer.  
18 346 Clearly, while quite a lot of small molecular weight material (<10 kDa) was recovered in the Low cut-  
19 347 off fraction, it contained very little AX, while some high MW AX was extracted just with the use of  
20 348 Buffer. Chemical extraction is shown in Trials 2 and 3, with good agreement, showing roughly twice  
21 349 as much High cut-off material as Low was recovered, but that the latter contained very little AX, such  
22 350 that the majority of the recovered AX was in the High cut-off fraction, giving a yield of around 2.6%  
23 351 compared with only 0.62% with Buffer. Thus chemical extraction using alkaline hydrogen peroxide  
24 352 was somewhat successful at releasing high MW AX from wheat bran. Further extraction with Buffer  
25 353 (Trial 2) recovered a little more high MW AX.  
26 354

27 355 Turning to Trials 7 and 8, use of FEA enzymes in combination with XYL appeared to enhance release  
28 356 of AX a little compared with just Buffer, but not substantially. Trials 3-7 taken together indicate that  
29 357 enzyme treatment following chemical extraction was able to release a little more AX, with the  
30 358 balance changing towards low MW material when the xylanase was included. This makes sense,  
31 359 although the absence of any detectable AX in the fractions from Trial 6 is unexpected. However, in  
32 360 general the enzymes did not dramatically enhance the further extraction of AX, and it is not possible  
33 361 to infer any meaningful differences in the performance of the different enzyme combinations.  
34 362

35 363 Figure 4 shows the equivalent results for sugarcane bagasse. Clearly the patterns are overall quite  
36 364 contrasting to those for the wheat bran. Most obviously, from Figure 4(c), the recovery of AX by  
37 365 chemical extraction was much greater than for wheat bran for both High and Low cut-off material,  
38 366 with yields of around 6-7% for high MW AX and 5.5% for low MW AX, compared with 2.6% and 0.1%,  
39 367 respectively, for wheat bran. Thus the bagasse released its AX much more readily following chemical  
40 368 extraction, and released a more balanced profile of AX between large and small MW molecules; this  
41 369 is also apparent in Figure 4(a) which shows much greater crude yields of Low cut-off material than  
42 370 High, in contrast to the yields from wheat bran. A consequence of the ready release of AX under  
43 371 chemical extraction is that there is therefore less material for the enzymes subsequently to work on,  
44 372 hence the subsequent enzyme treatments yield very little extra AX, although again there is evidence  
45 373 that the presence of the xylanase shifts the balance towards smaller molecules, as expected; this is  
46 374 clearer in Figure 5 which presents an expanded view of the Absolute yields, to allow the effects of  
47 375

375 the enzyme treatments, such as they are, to be seen more clearly. Even on their own, however,  
1 376 without prior chemical extraction, the enzymes release little more than Buffer alone (Trials 7-8 *c.f.*  
2 377 Trial 1), with the extra being entirely small MW material. It is recognised, however, that this small  
3 378 study did not explore a wider range of enzyme dosages and incubation conditions; it is likely that the  
4 379 effects of the enzymes could be enhanced under optimised conditions.  
5 380

8 381 Figure 6 shows the A:X ratios in the various wheat bran and bagasse extracts. In line with the  
9 382 starting material and as expected from literature reports, the wheat bran extracts have much higher  
10 383 A:X ratios than those from bagasse. Given the limitations of the work, not much more can be read  
11 384 into the fine detail of Figure 6, beyond noting that for the bagasse extracts, the smaller molecular  
12 385 weight material (Low cut-off fractions) consistently had lower A:X ratios than the corresponding  
13 386 larger MW fractions, whereas the wheat-derived AX presents a more mixed picture. It is well  
14 387 established that in general different parts of the biomass structure contain AX with different  
15 388 molecular weights and A:X ratios, reflective of different botanical functions of AX in different parts of  
16 389 the plant [22]. For the bagasse extracts, it appears to be consistently the case that material initially  
17 390 released by whichever means (chemical extraction, enzymes or buffer) has higher A:X ratios than  
18 391 material released subsequently via further extraction with buffer or enzymes, again reflecting  
19 392 differences in the nature of AX material given up easily compared with that released on further  
20 393 processing. The picture is less consistent for the wheat bran extracts, reflecting that the initial  
21 394 release was less extensive from the wheat bran than from the bagasse, such that comparisons are  
22 395 less dominated by that initial release; for the bagasse, so much was released initially that the  
23 396 remaining AX material is understandably quite different, whereas for the wheat bran, so little was  
24 397 released at all that what was released at any point was similar in structure. The A:X ratios greater  
25 398 than 1 for some of the Low cut-off wheat bran fractions (Trials 1 and 2) are probably erroneous,  
26 399 arising from errors in measuring very low concentrations of A and X in these samples (see Figure  
27 400 2(b)), although some components of wheat outer layers (cross-cells and pericarp) can have A:X  
28 401 ratios great than 1 [22].  
29 402

38 403 As noted already, the cleaner xylan chains of bagasse AX would offer somewhat different properties  
39 404 compared to those of wheat AX. This is an important consideration in developing commercial  
40 405 products; the challenge is to understand the functional performance and potential uses of AX  
41 406 fractions as affected by molecular weight and A:X ratio, and hence to understand which feedstocks  
42 407 and extraction processes are suitable for producing specific fractions. Even then, the likely scenario  
43 408 is not that specific fractions would be targeted for exclusive production, but rather that processing  
44 409 would co-produce a range of AX fractions, each suitable for different end-use applications, including  
45 410 small AXOS fractions with prebiotic functionality in food and in animal feed, alongside mid-range and  
46 411 large molecular weight fractions offering gradations of product functionality in relation to viscosity,  
47 412 gel formation and interaction with other food components [4]. In this respect, commercialisation of  
48 413 AX-based products is likely to follow the fractionation paradigm of crude oil cracking, to produce a  
49 414 range of products and to find markets for each. The use of enzymes would form part of the  
50 415 approach for creating specific fractions with targeted end-uses.  
51 416

57 417 The above observations and comments regarding the results from the current work are made in full  
58 418 recognition of the limits of replication and accuracy of the study; nevertheless, the overall patterns  
59 419 are clear, relative to the objectives of the work and the wider commercial context, and lead  
60 420 confidently to the following conclusions:  
61 421

62 421

63  
64  
65

- 422 1. Arabinoxylans can be extracted from sugarcane bagasse via similar protocols previously used for  
423 wheat bran, yielding AX with a lower A:X ratio than for wheat bran.
- 424 2. Bagasse released its AX more readily than wheat bran, and released a wider range of AX  
425 molecules, with a greater proportion of small MW (<10 kDa) molecules. In the current work,  
426 around two-thirds of the AX in the bagasse was released by chemical extraction, more or less  
427 equally divided between High and Low cut-off material, compared with just 11% of mostly large  
428 MW AX initially in the wheat bran.
- 429 3. Within the conditions used, feruloyl esterase and xylanase enzymes had small effects on  
430 releasing AX from raw wheat bran or sugarcane bagasse or from residues following chemical  
431 extraction, with the xylanase tending to shift the balance from large to small MW molecules.

432  
433 Following these results, larger scale extractions of AX from bran and bagasse were performed at the  
434 Biorenewables Development Centre (BDC) using alkaline hydrogen peroxide extraction. Bran was  
435 washed with water prior to extraction to remove the starch. 25 kg of bran and bagasse, in 5 kg  
436 batches, were subject to chemical extraction, centrifugation, ultrafiltration, ethanol precipitation,  
437 recovery and drying. A total of 1250 g of wheat bran extract (5% yield) of 54% purity and 848 g of  
438 bagasse extract (17% yield) at 52% purity were produced, with much of the rest being analysed as  
439 glucose, either from residual starch or from cellulose.

## 440 441 442 **Conclusions**

443  
444 The hypotheses that arabinoxylans could be extracted from sugarcane bagasse using similar  
445 protocols used for wheat bran, and that enzyme treatment might enhance the extraction, were  
446 investigated in a small proof-of-concept project. Bagasse was shown to be a promising source of AX  
447 in terms of its content (around 20%) and structure (with a low A:X ratio) and the readiness with  
448 which it yielded its AX to give a balanced release of both large (>10 kDa) and small (<10 kDa)  
449 molecules. Use of feruloyl esterase and xylanase enzymes on their own or following alkaline  
450 peroxide extraction was not particularly effective at enhancing AX release; however, there was  
451 evidence that the xylanase was effective at reducing the size of AX molecules, and there is scope to  
452 optimise the action of the enzymes through a more comprehensive study of dosage and incubation  
453 effects.

454  
455 Thus, the metaphorical reactions that arose from the LBNet workshop were successful in  
456 demonstrating the proof of concept, but the literal reactions in relation to optimising AX extraction  
457 from wheat bran and sugarcane bagasse retain some scope for further enhancement.

458  
459 As bagasse frequently arises within the context of bioethanol production, integration of AX  
460 extraction with ethanol production could allow economic production of AX products, as has been  
461 demonstrated in principle for AX co-production in a wheat ethanol plant [1]. Further processing of  
462 the now lignin-rich residue could give even further opportunities within the biorefinery, both for co-  
463 production of additional products and for further integration (particularly if the lignin processing  
464 also involves ethanol). The nature and additional processing of the lignin in the residual fractions  
465 from the current work were therefore studied further, and will be the topic of a future paper.

466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565

468 **Acknowledgements**

1 469  
2  
3 470 The authors gratefully acknowledge the funding for “Project POC02\_NOV 14 Campbell” from the  
4 471 Lignocellulosic Biorefinery Network (LBNNet), funded by the Biotechnology and Biological Sciences  
5 472 Research Council (BBSRC). Dr Matthew White and Prof Julian Cooper from AB Sugar are gratefully  
6  
7 473 acknowledged for supporting the project through provision of wheat bran and sugarcane bagasse  
8 474 and ongoing guidance throughout the project. Dr Gavin Thomas from the University of York and Dr  
9 475 Maggie Smallwood from BDC are thanked for facilitation of the project during its inception and  
10  
11 476 implementation. Alice Tibessart is gratefully acknowledged for performing some of the extraction  
12 477 work during her placement visit from Chimie ParisTech.

13 478

14  
15 47916 480 **References**

- 17 481  
18  
19 482 [1] N. Misailidis, G.M. Campbell, C. Du, J. Sadhukhan, M. Mustafa, F. Mateos-Salvador, R.M.  
20 483 Weightman, Evaluating the feasibility of commercial arabinoxylan production in the context  
21 484 of a wheat biorefinery principally producing ethanol. Part 2. Process simulation and economic  
22 485 analysis, *Chem. Eng. Res. Des.* 87 (2009). doi:10.1016/j.cherd.2008.12.028.  
23  
24 486 [2] C.S. Lancefield, O.S. Ojo, F. Tran, N.J. Westwood, Isolation of functionalized phenolic  
25 487 monomers through selective oxidation and CO bond cleavage of the  $\beta$ -O-4 linkages in Lignin,  
26 488 *Angew. Chemie - Int. Ed.* (2015). doi:10.1002/anie.201409408.  
27 489 [3] E. Martinez-Hernandez, J. Sadhukhan, G.M. Campbell, Integration of bioethanol as an in-  
28 490 process material in biorefineries using mass pinch analysis, *Appl. Energy*. 104 (2013).  
29 491 doi:10.1016/j.apenergy.2012.11.054.  
30  
31 492 [4] E. Martinez-Hernandez, A. Tibessart, G.M. Campbell, Conceptual design of integrated  
32 493 production of arabinoxylan products using bioethanol pinch analysis, *Food Bioprod. Process.*  
33 494 (2018). doi:10.1016/j.fbp.2018.08.005.  
34 495 [5] Z.C. Geng, J.X. Sun, S.F. Liang, F.Y. Zhang, Y.Y. Zhang, F. Xu, R.C. Sun, Characterization of  
35 496 water- and alkali-soluble hemicellulosic polymers from sugarcane bagasse, *Int. J. Polym. Anal.*  
36 497 *Charact.* (2006). doi:10.1080/10236660600649095.  
37  
38 498 [6] P. Feng, R. Jun-Li, X. Feng, B. Jing, P. Pai, S. Run-Cang, Comparative study of hemicelluloses  
39 499 obtained by graded ethanol precipitation from sugarcane bagasse, *J. Agric. Food Chem.* 57  
40 500 6305-6317. (2009). doi:10.1021/jf900986b.  
41 501 [7] C.M.P. Braga, P. da S. Delabona, D.J. da S. Lima, D.A.A. Paixão, J.G. da C. Pradella, C.S. Farinas,  
42 502 Addition of feruloyl esterase and xylanase produced on-site improves sugarcane bagasse  
43 503 hydrolysis, *Bioresour. Technol.* (2014). doi:10.1016/j.biortech.2014.07.115.  
44 504 [8] M. Aguedo, H.A. Ruiz, A. Richel, Non-alkaline solubilization of arabinoxylans from destarched  
45 505 wheat bran using hydrothermal microwave processing and comparison with the hydrolysis by  
46 506 an endoxylanase, *Chem. Eng. Process. Process Intensif.* (2015).  
47 507 doi:10.1016/j.cep.2015.07.020.  
48  
49 508 [9] L. Jacquemin, A. Mogni, R. Zeitoun, C. Guinot, C. Sablayrolles, L. Saulnier, P.Y. Pontalier,  
50 509 Performance evaluation of a semi-industrial production process of arabinoxylans from wheat  
51 510 bran, *Process Biochem.* (2015). doi:10.1016/j.procbio.2015.01.015.  
52  
53 511 [10] C. Maes, J.A. Delcour, Alkaline hydrogen peroxide extraction of wheat bran non-starch  
54 512 polysaccharides, *J. Cereal Sci.* (2001). doi:10.1006/jcfs.2001.0377.  
55 513 [11] C. Maes, J.A. Delcour, Structural characterisation of water-extractable and water-  
56 514 unextractable arabinoxylans in wheat bran, *J. Cereal Sci.* 35 (2002) 315–326.  
57 515 doi:10.1006/jcfs.2001.0439.  
58  
59 516 [12] C.M. Courtin, J.A. Delcour, Arabinoxylans and endoxylanases in wheat flour bread-making, *J.*  
60 517 *Cereal Sci.* (2002). doi:10.1006/jcfs.2001.0433.  
61 518 [13] J.J. Ordaz-Ortiz, L. Saulnier, Structural variability of arabinoxylans from wheat flour.  
62 519 Comparison of water-extractable and xylanase-extractable arabinoxylans, *J. Cereal Sci.* 42

63

64

65

- 520 (2005) 119–125. doi:10.1016/j.jcs.2005.02.004.
- 1 521 [14] A.P. de Souza, D.C.C. Leite, S. Pattathil, M.G. Hahn, M.S. Buckeridge, Composition and  
2 522 Structure of Sugarcane Cell Wall Polysaccharides: Implications for Second-Generation  
3 523 Bioethanol Production, *Bioenergy Res.* 6 (2013) 564–579. doi:10.1007/s12155-012-9268-1.
- 4 524 [15] J. Hollmann, M.G. Lindhauer, Pilot-scale isolation of glucuronoarabinoxylans from wheat  
5 525 bran, *Carbohydr. Polym.* (2005). doi:10.1016/j.carbpol.2004.09.015.
- 6 526 [16] C. Du, G.M. Campbell, N. Misailidis, F. Mateos-Salvador, J. Sadhukhan, M. Mustafa, R.M.  
7 527 Weightman, Evaluating the feasibility of commercial arabinoxylan production in the context  
8 528 of a wheat biorefinery principally producing ethanol. Part 1. Experimental studies of  
9 529 arabinoxylan extraction from wheat bran, *Chem. Eng. Res. Des.* 87 (2009) 1232–1238.  
10 530 doi:10.1016/j.cherd.2008.12.027.
- 11 531 [17] S.C. Fry, *The growing plant cell wall: chemical and metabolic analysis*, The Blackburn Press,  
12 532 New Jersey, USA, 1988. doi:10.1111/j.1469-8137.2003.00980.x.
- 13 533 [18] L. Jones, J.L. Milne, D. Ashford, S.J. McQueen-Mason, Cell wall arabinan is essential for guard  
14 534 cell function, *Proc. Natl. Acad. Sci.* (2003). doi:10.1073/pnas.1832434100.
- 15 535 [19] S. Ranganathan, D.G. Macdonald, N.N. Bakhshi, Kinetic studies of wheat straw hydrolysis  
16 536 using sulphuric acid, *Can. J. Chem. Eng.* 63 (1985) 840–844. doi:10.1002/cjce.5450630521.
- 17 537 [20] S. Willför, A. Pranovich, T. Tamminen, J. Puls, C. Laine, A. Suurnäkki, B. Saake, K. Uotila, H.  
18 538 Simolin, J. Hemming, B. Holmbom, Carbohydrate analysis of plant materials with uronic acid-  
19 539 containing polysaccharides-A comparison between different hydrolysis and subsequent  
20 540 chromatographic analytical techniques, *Ind. Crops Prod.* 29 (2009) 571–580.  
21 541 doi:10.1016/j.indcrop.2008.11.003.
- 22 542 [21] L.W. Doner, K.B. Hicks, Isolation of hemicellulose from corn fiber by alkaline hydrogen  
23 543 peroxide extraction, *Cereal Chem.* (1997). doi:10.1094/CCHEM.1997.74.2.176.
- 24 544 [22] L. Saulnier, P.E. Sado, G. Branlard, G. Charmet, F. Guillon, Wheat arabinoxylans: Exploiting  
25 545 variation in amount and composition to develop enhanced varieties, *J. Cereal Sci.* 46 (2007)  
26 546 261–281. doi:10.1016/j.jcs.2007.06.014.
- 27 547
- 28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



Table

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

Table 1. Compositions of wheat bran and sugarcane bagasse (%w/w dry basis).

Component	Wheat bran	Sugarcane bagasse
Starch	29.8	ND
Hemicellulose	16.6	22.1
Cellulose	21.8	28.3
Lignin	10.4	22.0
Other	21.2	27.6.8
AX (=0.88×(A+X))	8.64	19.58
A:X ratio	0.57	0.21

Table 2. Crude yields, AX concentrations and AX yields from wheat bran and its fractions following extraction under different treatments.

	1. Buffer (Control)	2. Chemical + Buffer (Control)	3. Chemical + Enzymes FEA E0355 + XYL E0051	4. Chemical + Enzymes FEA E0356 + XYL 0051	5. Chemical + Enzyme FEA E0355	6. Chemical + Enzyme FEA E0356	7. Enzymes FEA E0355 + XYL E0051	8. Enzymes FEA E0356 + XYL E0051
Initial bran (g)	30.34	50.73	50.20	50.47 <sup>c</sup>	50.47 <sup>c</sup>	50.47 <sup>c</sup>	30.61	30.47
AX concentration (%) <sup>a</sup>	8.64	8.64	8.64	8.64	8.64	8.64	8.64	8.64
AX amount (g) <sup>a</sup>	2.62	4.38	4.34	4.36	4.36	4.36	2.64	2.63
CE Pellet (g)		30.29	29.58	29.21	29.54	30.27		
AX concentration (%)		34.05	38.50	33.67	36.98	35.87		
AX amount in CE pellet (g)		10.31	11.39	9.84	10.92	10.86		
CE High (g)		8.11	8.31					
AX concentration (%)		16.43	16.21					
AX amount in CE High (g)		1.33	1.35					
Absolute yield (%)		2.63	2.68					
CE Low (g)		3.84	4.97					
AX concentration (%)		1.75	0.74					
AX amount in CE Low (g)		0.067	0.037					
Absolute yield (%)		0.13	0.07					
EE Pellet (g)	14.63 <sup>b</sup>	16.02	15.28	15.04	14.10	15.06	15.88	15.71
AX concentration (%)	31.00	38.19	35.82	31.21	35.74	22.61	36.07	31.85
AX amount in EE pellet (g)	4.54	6.12	5.47	4.69	5.04	3.41	5.73	5.00
Absolute yield (%)	14.95	12.06	10.90	9.30	9.99	6.75	18.71	16.42
EE High (g)	1.81	2.00	1.21	1.32	1.62	2.23	2.69	3.36
AX concentration (%)	10.42	12.2	21.25	19.46	23.23	0.00	8.43	5.22
AX amount in EE High (g)	0.19	0.24	0.26	0.26	0.38	0.00	0.23	0.18
Absolute yield (%)	0.62	0.48	0.51	0.51	0.75	0.00	0.74	0.58
EE Low (g)	4.78	0.97	3.20	2.64	3.20	1.95	4.93	4.97
AX concentration (%)	1.15	0.29	13.47	11.83	0.47	0.21	3.72	2.70
AX amount in EE Low (g)	0.05	0.00	0.43	0.31	0.02	0.00	0.18	0.13
Absolute yield (%)	0.18	0.01	0.86	0.62	0.03	0.01	0.60	0.44

a. The data reported for AX concentration and amount in the bran appear to be erroneous, as they indicate less AX in the raw material than in the residual pellet; a starting concentration of 24% is more in line with the mass balance and with the expected AX content of wheat bran.

b. The results for the pellet and high- and low-cutoff fractions following just buffer extraction are reported in these EE rows, but these are not to be understood as having undergone enzyme treatment.

c. Initial weights averaged from Trials 2 and 3.

Table 3. Crude yields, AX concentrations and AX yields from sugarcane bagasse and its fractions following extraction under different treatments.

	1. Buffer (Control)	2. Chemical + Buffer (Control)	3. Chemical + Enzymes FEA E0355 + XYL E0051	4. Chemical + Enzymes FEA E0356 + XYL 0051	5. Chemical + Enzyme FEA E0355	6. Chemical + Enzyme FEA E0356	7. Enzymes FEA E0355 + XYL E0051	8. Enzymes FEA E0356 + XYL E0051
Initial bran (g)	6.34	11.31	11.76	11.54 <sup>b</sup>	11.54 <sup>b</sup>	11.54 <sup>b</sup>	6.29	6.15
AX concentration (%)	19.54	19.54	19.54	19.54	19.54	19.54	19.54	19.54
AX amount (g)	1.24	2.21	2.30	2.25	2.25	2.25	1.23	1.20
CE Pellet (g)		5.37	5.48	5.68	5.36	5.39		
AX concentration (%)		14.56	13.47	16.06	20.52	17.75		
AX amount in CE pellet (g)		0.78	0.74	0.91	1.10	0.96		
CE High (g)		2.91	3.30					
AX concentration (%)		23.23	24.05					
AX amount in CE High (g)		0.68	0.79					
Absolute yield (%)		5.98	6.75					
CE Low (g)		7.97	9.43					
AX concentration (%)		7.49	6.97					
AX amount in CE Low (g)		0.60	0.66					
Absolute yield (%)		5.28	5.59					
EE Pellet (g)	3.36 <sup>a</sup>	4.12	3.94	4.25	4.23	4.18	3.54	3.93
AX concentration (%)	29.81	17.37	18.42	15.95	17.07	21.25	23.84	25.95
AX amount in EE pellet (g)	1.00	0.72	0.73	0.68	0.72	0.89	0.84	1.02
Absolute yield (%)	15.80	6.33	6.17	5.88	6.26	7.70	13.42	16.58
EE High (g)	1.00	1.17	0.96	1.00	1.12	1.02	0.92	0.97
AX concentration (%)	3.86	5.05	4.56	2.82	3.88	5.26	4.17	3.72
AX amount in EE High (g)	0.039	0.06	0.04	0.03	0.04	0.05	0.04	0.04
Absolute yield (%)	0.61	0.52	0.37	0.24	0.38	0.46	0.61	0.59
EE Low (g)	3.01	3.04	3.13	3.06	2.97	3.36	2.78	2.94
AX concentration (%)	1.44	0.44	0.85	1.10	0.46	0.66	2.25	1.86
AX amount in EE Low (g)	0.043	0.01	0.03	0.03	0.01	0.02	0.06	0.05
Absolute yield (%)	0.68	0.12	0.23	0.29	0.12	0.19	1.00	0.89

a. The results for the pellet and high- and low-cutoff fractions following just buffer extraction are reported in these EE rows, but these are not to be understood as having undergone enzyme treatment.

b. Initial weights averaged from Trials 2 and 3.

**Figure**

[Click here to download Figure: Figures, 171118.pdf](#)

Figure 1. Procedure for chemical extraction of AX from wheat bran or sugarcane bagasse, yielding High cut-off (CE-High) and Low cut-off (CE-Low) fractions and a residual pellet; and procedure for further enzyme treatment of the pellet to yield further High (EE-High) and Low (EE-Low) cut-off fractions.




Figure 2. Measurement of arabinoxylan content in wheat bran following different hydrolysis conditions.




Figure 3. (a) Yield, (b) Concentration and (c) Absolute Yield in fractions following arabinoxylan extraction from wheat bran.






Figure 4. (a) Yield, (b) Concentration and (c) Absolute Yield in fractions following arabinoxylan extraction from sugarcane bagasse.






Figure 5. Expanded view of Absolute Yields in fractions following arabinoxylan extraction from sugarcane bagasse under various chemical and enzyme treatments.






Figure 6. Arabinose:Xylose ratios in fractions following arabinoxylan extraction from (a) wheat bran and (b) sugarcane bagasse.

