

This is a repository copy of *Comparing the impact of future cropland expansion on global biodiversity and carbon storage across models and scenarios*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/153734/>

Version: Accepted Version

Article:

Molotoks, Amy, Henry, Roslyn C, Stehfest, Elke et al. (6 more authors) (2020) Comparing the impact of future cropland expansion on global biodiversity and carbon storage across models and scenarios. Philosophical Transactions Of The Royal Society Of London Series B - Biological Sciences. ISSN: 1471-2970

<https://doi.org/10.1098/rstb.2019.0189>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

PHILOSOPHICAL TRANSACTIONS OF THE ROYAL SOCIETY B

BIOLOGICAL SCIENCES

Comparing the impact of future cropland expansion on global biodiversity and carbon storage across models and scenarios

Journal:	<i>Philosophical Transactions B</i>
Manuscript ID	RSTB-2019-0189.R2
Article Type:	Research
Date Submitted by the Author:	n/a
Complete List of Authors:	Molotoks, Amy; University of Aberdeen, Institute of Biological & Environmental Sciences, School of Biological Sciences, 23 St Machar Drive, AB24 3UU; University of York, SEI York, Environment and Geography Henry, Roslyn; School of Geosciences, University of Edinburgh, Edinburgh, UK Stehfest, Elke; Netherlands Environmental Assessment Agency (PBL), Doelman, Jonathan; Netherlands Environmental Assessment Agency (PBL) Petr, Havlík; International Institute for Applied Systems Analysis (IIASA), Krisztin, Tamas; International Institute for Applied Systems Analysis (IIASA) Alexander, Peter; University of Edinburgh School of GeoSciences, ; SRUC, Dawson, Terry; King's College London Smith, Pete; University of Aberdeen, Institute of Biological & Environmental Sciences
Issue Code (this should have already been entered but please contact the Editorial Office if it is not present):	SACKLER
Subject:	Environmental Science < BIOLOGY
Keywords:	Land use change, biodiversity, carbon storage, integrated models

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Author-supplied statements

Relevant information will appear here if provided.

Ethics

Does your article include research that required ethical approval or permits?:
This article does not present research with ethical considerations

Statement (if applicable):
CUST_IF_YES_ETHICS :No data available.

Data

It is a condition of publication that data, code and materials supporting your paper are made publicly available. Does your paper present new data?:
Yes

Statement (if applicable):
Datasets and code will be deposited in a general repository e.g. Dyrad before publication

Conflict of interest

I/We declare we have no competing interests

Statement (if applicable):
CUST_STATE_CONFLICT :No data available.

Authors' contributions

This paper has multiple authors and our individual contributions were as below

Statement (if applicable):
AM coordinated contributions and led analysis on AZE sites and carbon storage. RH led analysis on hotspots and statistical analysis. All authors contributed to the design, drafting and revision of the article and all authors have given final approval of the version to be published.

Comparing the impact of future cropland expansion on global biodiversity and carbon storage across models and scenarios

Amy Molotoks^{1, 2*}, Roslyn Henry³, Elke Stehfest⁵, Jonathan Doelman⁵, Petr Havlik⁶, Tamás Krisztin⁶, Peter Alexander^{3, 4}, Terence P. Dawson⁷, Pete Smith¹

¹ Institute of Biological and Environmental Sciences, University of Aberdeen, 23 St Machar Drive, Aberdeen, AB24 3UU, UK

² Stockholm Environment Institute York, Department of Environment and Geography, University of York, York YO10 5NG, UK

³ School of Geosciences, University of Edinburgh, Edinburgh, UK

⁴ Global Academy of Agriculture and Food Security, The Royal (Dick) School of Veterinary Studies, University of Edinburgh, Edinburgh, UK

⁵ PBL Netherlands Environmental Assessment Agency, Bezuidenhoutseweg 30, 2594 AV The Hague, The Netherlands

⁶ IIASA, Schlossplatz 1, A-2361 Laxenburg, Austria

⁷ Department of Geography, King's College London, The Strand, London WC2R 2LS, UK.

Keywords: Land-use change, biodiversity, carbon storage, integrated models

Summary

Land-use change is a direct driver of biodiversity and carbon storage loss. Projections of future land-use often include notable expansion of cropland areas in response to changes in climate and food demand, although there are large uncertainties in results between models and scenarios. This study examines these uncertainties by comparing three different socio-economic scenarios (SSP1-3) across three models (IMAGE, GLOBIOM and PLUMv2). It assesses the impacts on biodiversity metrics and direct carbon loss from biomass and soil as a direct consequence of cropland expansion. Results show substantial variation between models and scenarios, with little overlap across all nine projections. Although SSP1 projects the least impact, there are still significant impacts projected. IMAGE and GLOBIOM project the greatest impact across carbon storage and biodiversity metrics due to both extent and location of cropland expansion. Furthermore, for all the biodiversity and carbon metrics used, there is a greater proportion of variance explained by model used. This demonstrates the importance of improving the accuracy of land-based models. Incorporating effects of land-use change in biodiversity impact assessments would also help better prioritise future protection of biodiverse and carbon-rich areas.

Introduction

Land-use change is a key direct driver of biodiversity loss (IPCC 2001; Parmesan and Yohe, 2003) and is one of the main drivers of species extinctions (Pimm et al., 2014). It is also expected to be exacerbated by climate change, which can also impact indirectly on biodiversity in a number of ways (Smith et al., 2018). For example, there is a negative global impact on crop production, which is projected to be high in the coming decades. For each degree-Celsius increase in global mean temperature, a 3.1-7.4% reduction in global yields of major crops is estimated (Zhao et al., 2017). This means cropland area will likely need to expand to meet increasing demand for food (Godfray et al., 2010; Delzeit et al., 2017), particularly in countries with growing food needs and limited access to technology which would allow sustainable intensification (Alexandratos and Bruinsma, 2012).

Cropland expansion is known to have severe adverse effects on natural biodiversity (Pimm and Raven, 2000; Purvis, Jones and Mace, 2000; Baillie, Hilton-Taylor and Stuart, 2004), through loss and fragmentation of habitats (Foley et al., 2005). Conversely, land-use and land-cover change (LULCC) also impacts climate change, and has accounted for an estimated 12.5% of anthropogenic carbon emissions from 1990-2010 (Houghton et al., 2012). Clearing natural ecosystems for crop

*Author for correspondence (amy.molotoks@york.ac.uk).

production also releases carbon dioxide into the atmosphere as stored carbon is released from biomass and soil (West et al., 2010). Human and natural responses to climate change are interconnected, with the majority of future model simulations of global cropland expansion exceeding the 15% planetary boundary in order to meet future food-supply targets (Henry et al., 2018). Therefore, research on food production systems and ecosystem impacts should be prioritised (Hannah et al., 2013).

Future land-use change has been explored through the application of modelling based upon the narratives for the shared socioeconomic pathways (SSPs, Popp et al., 2017; Riahi et al., 2017). Model results indicate a range of potential future land-use outcomes and have typically focused on consequences for greenhouse gas emissions, food provisions and prices. However, there has been less focus on potential consequences for biodiversity (Chaudhary and Mooers, 2018). Furthermore, in a recent review of biodiversity scenarios, Titeux et al. (2016) highlighted that biodiversity scenario analysis typically neglects consequences related to land-use change, but rather focuses on direct impacts of climate change. Thus, exploration of biodiversity impacts of future land-use change scenarios, which are partially driven by climate change, warrants further research.

There are, however, large uncertainties associated with model-based projections of future global land-use change (Schmitz et al., 2014). Existing studies have highlighted that both the total global quantity (Alexander et al., 2017) and regional specific land-use changes (Prestele et al., 2016) vary greatly according to the model. Similarly, other aspects such as climate-change responses and bioenergy impacts (Popp et al., 2014; Von Lampe et al., 2014) vary between models. While model inter-comparisons have considered differences in land-use change and associated climate impacts between models, no previous comparison has examined variation in biodiversity and carbon storage. On a global scale, studies have shown a high correlation between species richness and carbon storage, with a strong association between carbon stocks and mammal, bird and amphibian distributions (Strassburg et al., 2010). Although plot level studies observe weaker correlations, a strong association has also been observed at a national level, with a high proportion of threatened species relying on carbon-rich habitats in tropical regions (Sheil et al., 2016). Cropland expansion threatens both carbon storage and biodiversity, with consequences for ecosystem functioning (Tscharntke et al., 2005; Don, Schumacher and Freibauer, 2011; Delzeit et al., 2017). This global study therefore aims to compare the impact of cropland expansion projections on biodiversity and carbon storage across three different models and three different SSPs. This process allows for the quantification of variability in biodiversity and carbon outcomes associated with model and scenario which is important for more holistic assessments of the impact of land-use change. Differentiating the effect of extent and location can also be used to determine the relative importance of improving the accuracy of land-based models or socio-economic scenarios, for the purposes of prioritising areas for biodiversity conservation and carbon storage in the future.

Methods

Land-use models

Model outputs from the modelling teams GLOBIOM (Havlik et al., 2011), IMAGE (Stehfest et al., 2014) and PLUMv2 (Alexander et al., 2018) were collected, each looking at the time period 2010-2050. GLOBIOM is a global recursively dynamic partial equilibrium model which integrates the agricultural, bioenergy and forestry sectors (Havlik et al., 2011), with its main drivers being population, GDP, input prices, bioenergy demand, taxes and yields (Dumollard et al. 2012). It requires geographic information and land profitability of crop production for its land allocation (Havlik et al., 2011), basing its cropland expansion on a land rent approach (Schmitz et al., 2013). In comparison, the land component of IMAGE (Stehfest et al., 2014) uses a computable general equilibrium model, MAGNET (Woltjer and Kuiper, 2014), to calculate agricultural demand, trade and production. There are six key drivers for IMAGE: demography, economic growth, policy and governance, technological development, culture and lifestyle and natural resource availability, with a regression-based suitability assessment allocating land-use change (Stehfest et al., 2014). PLUMv2 is a global land-use and food-system model that combines spatially-explicit, biophysically-derived yield responses with socio-economic scenario data to project future demand, land use, and management inputs (Alexander et al., 2018). For each country and time-step, the agricultural land use and level of imports or exports is determined through a least-cost optimisation that meets the demand for food and bioenergy commodities in each country. GLOBIOM uses the crop model EPIC (Dumollard et al., 2012) whilst IMAGE uses the dynamic global vegetation model LPJmL (Müller et al., 2016) to determine cropland yields, both producing a spatially explicit output at 0.5°* 0.5° gridded resolution. Similarly, PLUMv2 (Alexander et al., 2018) uses a dynamic global vegetation model, LPJ-GUESS, to provide crop yield responses on a 0.5° * 0.5° grid (IPCC, 2014).

Scenarios

The models described can be used to simulate the effects of different shared socio-economic pathways (SSPs) (Fricko et al., 2017; van Vuuren et al., 2017; Alexander et al., 2018; Doelman et al., 2018) which are defined as ‘reference pathways describing plausible alternative trends in the evolution of society and ecosystems over a century timescale’ (O’Neill et al. 2014). SSP1 represents low challenges for mitigation and adaptation to climate change, SSP2 is moderate and SSP3 is high. SSP1 is the ‘greenest’ with sustainable development proceeding at a high pace, lessening global inequalities. There is rapid

technological change towards low carbon energy sources and high productivity of land whilst SSP3 has slow technological change, a rapidly growing population with unmitigated emissions. Investments in human capital are also low, with high inequality, reduced trade flows and large numbers of people being left vulnerable to climate change with low adaptive capacity. SSP2 is an intermediate case between SSP1 and SSP3 and represents a future where development trends are neither extreme, but follow a middle-of-the-road pathway consistent with typical patterns observed over the past century (O'Neill et al., 2017).

Biodiversity metrics

AZE sites

The Alliance for Zero Extinction (AZE) sites are identified by three criteria: it must contain at least one individual species which has evaluated as Endangered or Critically Endangered under the International Union for Conservation of Nature (IUCN) criteria; it is the sole area where this species occurs, containing over 95% of the known resident population; and it has a definable boundary (Alliance for Zero Extinction, 2010). These species often have little official protection, making them extremely vulnerable to external threats such as habitat destruction (Ricketts et al., 2005). Currently, 587 sites for 920 species of mammals, birds, amphibians, reptiles, conifers, and reef-building corals have been identified with 81% AZE sites being found within a biodiversity hotspot. These sites are therefore an important indicator of biological significance and the impact of future cropland expansion could threaten them further. So, the AZE dataset was used in a spatial overlay, as in Molotoks et al. (2018), to examine infringement of cropland expansion on AZE sites. The sum of AZE sites per region was then calculated per model and per scenario to estimate the total number of sites impacted.

Conservation International (CI) hotspots

Cropland expansion projections within CI hotspots was also explored, the criteria for which account for vascular plant species richness. CI hotspots identify regions of importance for biodiversity and to qualify, a region must be threatened - i.e. contain at most 30% of its original natural vegetation - yet contain at least 1,500 different species of endemic vascular plants. The 35 CI hotspots cover 2.3% of the land surface but support 50% of the world's endemic plant species and 43% of vertebrate endemic species (Myers et al., 2000; Mittermeier et al., 2004). CI hotspot shapefile data were converted to 0.5° raster maps. Any 0.5° cell containing CI hotspot polygon data is classified as a CI hotspot irrespective of hotspot size. The CI map is therefore binary and cells are classified as either a CI hotspot or not.

Vertebrate species rich regions

As another biodiversity metric, maps of vertebrate species richness, small-range vertebrate species richness, and threatened species richness were considered (Jenkins, Pimm and Joppa, 2013; Pimm et al., 2014). The resolution of the vertebrate species richness maps was decreased from 0.1° to 0.5° resolution to match the resolution of the three models involved in our analysis; the mean species richness was calculated for each grid cell. For all taxa, the distribution of species richness across grid cells is right-skewed: most cells contain a few species, while there are a few cells with a large number of species. For each taxon therefore the mean species richness values of grid cells was converted into percentile values and 'species-rich regions' assumed to be cells in the 90th percentile of grid cells.

Cropland expansion projected by PLUMv2, IMAGE and GLOBIOM in vertebrate species-rich regions was explored across the three SSP scenarios. Furthermore, for each model and SSP combination, regions of threat: regions with high biodiversity (either CI hotspot or vertebrate species rich region) under pressure from cropland expansion, were identified. An overall threat index for all species per grid cell was then calculated. This is the percentage of cropland expansion projected by 2050 from the models multiplied by the summed richness index of amphibians, birds and mammals. For the threat index, it was assumed that each species is equally important regardless of taxon. Calculating the threat index allowed comparisons of the location of threatened areas between the models and SSPs.

Carbon storage

Biomass

To examine storage loss in vegetation, cropland expansion projections for each model and scenario were overlaid with a combined dataset of C storage in fourteen forest types (Molotoks et al., 2018). Vegetation C stocks presented by Ruesch and Gibbs (2008) for land covers represented in the Global Land Cover 2000 map (Arino et al., 2010) were used to calculate carbon loss at 1km resolution in tonnes per hectare. This represents the total biomass C stored in both above and below ground vegetation. Where cropland expansion projections overlapped with forests, it is assumed the C stored is lost as a result of vegetation being cleared. Building on the methodology used in Molotoks et al. (2018), the mean value of carbon

present in tonnes per hectare, and the area and the percentage of cropland expansion for each individual grid cell were used to calculate an estimated total C loss.

Soil

Soil carbon stocks represented in the Harmonized World Soil Database (FAO/IIASA/ISRIC/ISS-CAS/JRC v.1.1, 2009) were also examined. 30-arc second resolution grids for each land use represented in the Global Land Cover 2009 map were used (Arino et al. 2010), using the total organic soil C stock density to a depth of 1 m reported by Hiederer and Kochy (2011). The mean value of carbon present for each grid cell, majority land cover, and figures from a global meta-analysis of land-use change impacts on soil organic carbon (SOC) (Guo and Gifford, 2002) were used to calculate estimates of soil carbon loss. For example, there is an estimated 42% and 59% loss of SOC when converting to cropland from forest and grassland, respectively (Guo and Gifford, 2002).

Statistical analysis

A similar approach to the statistical analysis as Prestele et al. (2016) and Alexander et al. (2017) was taken, identifying the sources of variance in the results for each of the different biodiversity metrics considered, by fitting multiple linear regressions with model and SSP as variables. Interaction terms were not considered, and the variance associated with such interactions is incorporated within the residuals. An analysis of variance (ANOVA) was then performed on the regression models to extract the type II sum of squares values for each variable to partition the relative importance of model and scenario. The statistical analysis here is not used to draw inferential conclusions with regards to whether the models or SSP scenarios have statistically significant effects on cropland expansion and, consequently, biodiverse regions. Rather, the variance of the results is partitioned to indicate the level of variance that can be associated with model choice or SSP scenario.

Results

To summarise, across all metrics, SSP1 typically has the lowest impacts on biodiversity and carbon storage. PLUMv2 in general shows the least impact on carbon storage, whilst IMAGE has the highest impact across biodiversity metrics. The highest impact on carbon storage is also seen in IMAGE, but there is variation between carbon loss from biomass and from soil. For all metrics used, both for biodiversity and carbon storage, the majority of variance is explained by the model used (Table 1).

[Table 1 here]

Biodiversity metrics

AZE sites

[Figure 1 here]

For all three models, cropland expansion infringing on AZE sites is lowest under SSP1 (Figure 1d). In the SSP1 scenarios, IMAGE projections show the greatest impact on AZE sites globally while in the SSP2 and SSP3 scenarios, GLOBIOM projections show the greatest impact (Figure 1d). For example in SSP2, 102 sites are projected to be impacted by cropland expansion in South America alone (Figure 1b). PLUMv2 projections show the smallest impact across all scenarios at a global level and across most regions (Figure 1). However, while there is variation in the number of AZE sites that cropland is projected to expand into across the SSPs, SSP accounted for only 21.4% of the variation in model results (Table 1). A larger fraction of the variation (63.5%) in the AZE results is explained by the model (Table 1).

Europe is almost consistently the region with the fewest sites impacted across all models and SSPs, whilst the Americas are the most highly impacted. South America has the highest numbers of AZE sites impacted by cropland expansion across all SSPs (Figure 1a-c). There is, however, variation within the models. For example, IMAGE projections show higher numbers of AZE sites impacted in North America than South America for SSP1 and 3 (Figure 1 a, c). Similarly, PLUMv2 projects a slightly higher number of AZE sites impacted in North America than South America in SSP2 (Figure 1b). There is also variation across other regions between model projections. IMAGE consistently projects the highest numbers of AZE sites impacted in Africa and Oceania across all three scenarios, whilst GLOBIOM projections show higher impacts for Europe and South America (Figure 1a-c).

Vertebrate species rich regions and CI hotspots

[Figure 2 here]

As with AZE sites, the smallest areas of cropland expansion in vertebrate species rich and CI hotspots regions are found in the SSP1 scenarios (Figure 2). SSP3 has the largest impacts, projecting the greatest area (Figures S1-4) with a high threat index in all three models (Figure 3). Yet the majority of variation is explained by the model (Table 1).

[Figure 3 here]

Figure 3 shows this variation between the models. South East Asia is the most heavily affected in PLUMv2 projections, whilst West Africa and the Cerrado region in Brazil show the most cropland expansion in GLOBIOM projections (Figure 3). GLOBIOM also projects the greatest levels of total cropland expansion in all species rich regions under SSP2 and SSP3 (Figure 2). For IMAGE projections, a wide range of areas in the tropics are shown to be affected, including South East Asia, Central Africa and the fringes of the Amazon rainforest in South America (Figure 3).

Carbon storage

For all three models, SSP1 has the lowest estimated carbon losses, for both the total estimates and individual estimates from biomass and soil, with the lowest estimates consistently shown in PLUMv2 projections (Figure 4d). Across all scenarios, IMAGE projections show the highest total losses of carbon with the greatest total estimate from SSP2 at over 46 Gigatonnes of carbon (GtC) lost from soil and biomass combined (Figure 4d). However, GLOBIOM generally has larger projected losses for soil carbon (Figure 4d), with higher carbon loss from temperate regions including North America (Figure 4 a-c).

[Figure 4 here]

As with biodiversity metrics, the model used also explains the greatest proportion of variance for carbon loss (Table 1). Africa and Oceania consistently have the largest impacts from IMAGE projections, whereas Europe and North America have the highest losses from GLOBIOM, and PLUMv2 shows higher losses in Asia (Figure 4a-c).

Discussion

SSPs scenarios are intended to have different environmental implications and therefore modelled differences between the SSPs are not unsurprising. While global land-use models differ by design, they all aim to model the same global system, capturing the same dynamics, and therefore ideally generate similar results under single scenarios. While all three models demonstrate some commonality in overall results, the models still vary considerably in their estimates of cropland expansion within SSPs. Our results are therefore in agreement with previous studies investigating uncertainties in land-use projections. For example, Alexander et al., (2017) and Prestele et al., (2016) both found large differences in land-cover projections between models with the highest variability occurring in future cropland areas. Our study is the first, however, to consider the implications of similarities and differences in land-cover projections arising under different models for biodiversity and carbon. Highlighting uncertainties between modelling approaches in terms of biodiversity and carbon impacts is important for conservation goals and climate change mitigation. When informed by model outcomes, conservation or mitigation measures could be misled when uncertainty is not considered. Conversely, identifying similarities between models across different metrics will help to identify key regions for prioritisation to ensure conservation and mitigation targets are met.

Biodiversity perspectives

The biodiversity results demonstrate similar broad patterns across models. For example, SSP1 consistently has the lowest levels of cropland expansion in AZE sites, vertebrate species rich regions and CI hotspots across all models (Figure 1, 2). Our results therefore agree with the findings from Chaudhary and Mooers (2018) who used land use model projections from the land-use harmonization dataset (LUH2) and found SSP1 resulted in the lowest land-use change driven global biodiversity loss. SSP1 is characterised by slow population growth, global sustainability and low vulnerability to climate change (van Vuuren and Carter, 2014). The world's growing population, coupled with increased affluence, are major drivers of food demand (Bajželj et al., 2014), so slow population growth will greatly influence the amount of cropland expansion. There is also strong land-use change regulation in SSP1 to avoid environmental trade-offs, and large assumed improvements in agricultural productivity (Popp et al., 2016) which would limit cropland expansion and subsequent encroachment into biodiverse regions (Foley et al., 2011). In contrast, across most models, the greatest levels of cropland expansion in biodiversity metrics examined are projected under SSP3, with the exception of IMAGE (Figure 2). SSP3 is characterised by limited land-use regulation and continued deforestation and, therefore, increased cropland expansion and subsequent changes in biodiverse regions are expected (Popp et al., 2017). Given the agreement of the models for SSP1, it is important for conservation that policy decisions strive for a global future characterised by land-use change regulation and 'green' choices, protecting biodiverse regions from cropland expansion.

There are certain areas that the models agree will experience cropland expansion (Figure 3, Figure S1-4) within the species rich regions and AZE sites. This agreement highlights them as areas of particular conservation concern. For example, under

SSP3, areas of Central Africa that contain high numbers of mammal species would be at risk (Figure S3). In terms of AZE sites, the models all project the greatest number affected by cropland expansion will be those in the Americas (Figure 1). Mexico (classified as North America in this study) is almost always the country with the highest numbers of AZE sites affected across all models, followed by Peru and Columbia (Appendix A). Other studies have also identified Mexico as a country expected to experience large habitat declines for a number of species by 2050 due to food production and consumption increases (Visconti et al., 2011). High levels of species richness and a large number of AZE sites cluster in the tropics. High levels of cropland expansion are projected in these areas as well, therefore the tropics and sub-tropics are where the threat index is found to be highest across the models (Figure 3). The tropics are also likely to see the greatest benefit to biodiversity in terms of the most species preserved, if global warming is constrained from 1.5 to 2°C (Smith et al., 2018). Hence, indirect impacts of climate change via land-use change could affect similar areas to those experiencing direct impacts of climate change. In particular, previous studies have highlighted areas in Central and South America as global priorities for adaptation of both agriculture and biodiversity in the face of climate change (Hannah et al., 2013; Warren et al., 2013; Smith et al., 2018). The three models here all agree that cropland expansion is expected in the tropics with notable impacts on AZE sites in the Americas. Thus, our results similarly suggest areas in Central and South America as conservation priorities, regardless of the SSP considered.

Despite similar general patterns across SSPs and some local spatial agreement in projected land change, there is considerable variability in the overall estimates of cropland expansion between models and, consequently, the effects on biodiverse regions and AZE sites (Figure 1, 2). Within SSPs, PLUMv2 consistently displays the lowest rates of cropland expansion, followed by IMAGE and GLOBIOM. Consequently the impact of cropland expansion in AZE sites, CI hotspots and species rich regions is lowest in projections produced by PLUMv2 and highest in projections produced by GLOBIOM. Furthermore, the larger cropland expansion with GLOBIOM results in larger areas of the temperate zones, such as North America, arising as regions of threat (Figure 3) compared to PLUMv2 and IMAGE. The lower cropland expansion observed in PLUMv2 likely result from the inclusion of crop and location specific fertiliser, irrigation intensification and the modelling of adaptation. GLOBIOM determines the amount of land that will be converted to agriculture through the use of a land supply curve (Eitelberg, van Vliet and Verburg, 2015). It has a high estimate of cropland availability as it is based on estimates of land productivity, relying mainly on biophysical production constraints (Havlik et al. 2011); hence, it has the largest extent of cropland expansion estimates of the three models (Figure 2).

Carbon perspectives

Similar to the biodiversity metrics, across models, SSP1 has the lowest estimated carbon losses. However estimates of carbon losses differ considerably between models. PLUMv2 consistently projects the lowest levels of carbon loss while, despite greater global cropland expansion with GLOBIOM, IMAGE projects the highest estimates of total carbon loss across SSPs at a global level. This global-level effect is largely driven by the location of cropland expansion in IMAGE compared to the other models. IMAGE projects high rates of cropland expansion in Central Africa, where some of the largest intact areas of tropical forest cover are located (Hansen, Stehman and Potapov, 2010; Potapov et al., 2012). Tropical vegetation currently stores ~340 billion tonnes of carbon and therefore higher rates of cropland expansion in Central Africa, as projected using IMAGE, results in higher levels of total carbon loss compared to the other models (West et al., 2010). This finding corroborates previous work and demonstrates the importance of considering not only uncertainty surrounding the magnitude of global cropland expansion but also the spatial location (Prestele et al., 2016). Our results serve to highlight that the location of cropland expansion has implications for carbon storage and, hence, the prioritisation of land conservation to mitigate carbon losses should consider the influence of models used to generate projections and the potential uncertainty involved.

Despite model differences, this study demonstrates that future cropland expansion has a significant negative impact on carbon storage. As much as 46 GtC is projected to be lost before 2050 (Figure 4), which is 3.4 times greater than the current annual global anthropogenic GHG emissions (IPCC, 2014), at a time when it is essential to minimise such emissions (Smith et al., 2016). Although models vary in their global estimates of potentially available cropland (Eitelberg, van Vliet and Verburg, 2015), large areas of remaining potentially cultivatable land are currently beneath tropical forests (Smith, 2013). Deforestation of the tropics for cropland expansion could lead to large-scale biodiversity and carbon losses. Although the feedback is not captured within all the models examined here, carbon loss contributes directly to climate change that, in turn, results in negative impacts on crop yield and increases the need for further cropland expansion. Consequently, future assessments of the impact of climate change on biodiversity and carbon storage should also consider the indirect effects of climate through land-use and land-management change (Smith et al., 2018).

Dealing with uncertainty in land based modelling studies

Our aim is to demonstrate the similarities and differences between models and scenarios concerning the impact of cropland expansion on carbon storage and biodiversity metrics. Given the apparent agreement between models and different metrics, we have highlighted SSP1 as the most desirable scenario for both biodiversity and carbon storage, although this scenario still projects high future impacts on metrics examined. For example, between 14-30 Gt C are projected to be lost in this

scenario (Figure 4d), 5-10,000,000 square kilometres of CI hotspots converted to cropland (Figure 2d) and up to 241 AZE sites impacted by this land use change (Figure 1d). This emphasises the need for a redoubling of efforts in SSP1 to avoid severe environmental impacts of future cropland expansion. Furthermore, we have identified regions that could be considered as priorities for both biodiversity and carbon storage loss (e.g. the Americas). However, there remains considerable variability in the estimates of cropland expansion between models within individual SSPs (Figure 2). Our results therefore demonstrate that intrinsic model characteristics can over or under-estimate cropland expansion irrespective of the scenario of interest. Model characteristics, parameterizations and institutional assumptions often lead to divergent land-use outcomes. Differences between the models here likely arise because of assumptions regarding cropland intensification, adaptation and estimates of cropland productivity. Furthermore, previous land-use model inter-comparisons have highlighted uncertainty arising from differences between initial land use input data, bioenergy production assumptions and yield responses to climate change associated with the underlying crop models (Nelson et al., 2014; Popp et al., 2014; Schmitz et al., 2014; Alexander et al., 2016; Prestele et al., 2016). For example Alexander et al. (2016) found substantial differences in starting cropland areas across 17 different models. Models often allocate land-use change based on land use in adjacent grid cells in former time steps (e.g. cropland expansion at the edge of existing agricultural area). Therefore, starting conditions can have a large influence on the dynamics of cropland expansion in future time steps (Alexander et al., 2016). The models used here and in other comparison studies also have different underlying crop yield models. Hence crop yield responses to inputs such as fertiliser and climate change can differ and ultimately affect the area of cropland required to meet projected demand for crop production (Nelson et al., 2014).

Conclusions

Here, we highlight firstly that even in the most environmentally sustainable pathway, there are significant impacts on biodiversity and carbon storage. Hence the importance of going beyond measures taken in the SSP1 scenario is emphasised. Secondly, the existence of uncertainty in land-use change projections needs to be acknowledged when designing conservation or mitigation strategies. Models are frequently selected for biodiversity or carbon studies based on user familiarity and accessibility and rarely are the results from more than a single model considered. Our intention is not to identify model results that are more plausible or the most accurate model. However, we show that it would be beneficial to include a range of models and scenarios when studying land use effects on biodiversity and carbon such that model uncertainty can be explored and areas for prioritisation identified. This is particularly important for prioritising AZE sites as the vast majority are unprotected, yet host small, restricted populations (Ricketts et al., 2005) of endemic, rare and threatened species (McDonald, Kareiva, and Forman, 2008). They are particularly vulnerable to external threats, as 95% of each individual species are found within the boundaries of their site (AZE, 2010). Hence, increased accuracy of land-based modelling studies could help prioritise sites to protect, thereby reducing potential future species extinctions. Recent studies have urged caution when using a single model for estimates of land-use change for environmental assessments (Prestele et al., 2016); here, we would urge the same from a biodiversity and carbon storage perspective. Previous efforts to model scenario outcomes, RCPs or SSPs, on biodiversity may also benefit from reassessment within the context of other land-use models to generate uncertainty. Focusing conservation efforts and climate mitigation in regions where models agree there will be substantial impacts could be an effective approach to conservation. Furthermore, considering results across different types of metrics (e.g. species rich regions, AZE sites and carbon stocks), could provide a comprehensive picture of biodiversity and carbon change, allowing for a holistic and cost-effective approach to prioritisation.

Additional Information

Acknowledgements

Conservation International, The Alliance for Zero Extinction and the Food and Agricultural Organisation (FAO) are acknowledged for provision of data.

Funding statement

This work contributes to the Belmont Forum/FACCE-JPI DEVIL project (NE/M021327/1) and the NERC funded Soils-R-GREAT project (NE/P019455/1). AM is supported by a Biotechnology and Biological Sciences Research Council (BBSRC) EastBio Studentship (<http://www.eastscotbiobdp.ac.uk/>) (grant number BB/M010996/1) and the Global Challenges Research Fund Trade, Development and the Environment Hub project (ES/S008160/1). PA and RH are supported by the UK's Global Food Security Programme project 'Resilience of the UK food system to Global Shocks' (RUGS, BB/N020707/1).

Competing Interests

The authors have no competing interests.

Authors' Contributions

AM coordinated contributions and led analysis on AZE sites and carbon storage. **RH** led analysis on hotspots and statistical analysis. All authors contributed to the design, drafting and revision of the article and all authors have given final approval of the version to be published.

References

- Alexander, P., Prestele, R., Verburg, P. H., Arneth, A., Baranzelli, C., Batista e Silva, F., Brown, C., Butler, A., Calvin, K., Dendoncker, N., Doelman, J. C., Dunford, R., Engström, K., Eitelberg, D., Fujimori, S., Harrison, P. A., Hasegawa, T., Havlik, P., Holzhauser, S., Humpenöder, F., Jacobs-Crisioni, C., Jain, A. K., Krisztin, T., Kyle, P., Laval, C., Lenton, T., Liu, J., Meiyappan, P., Popp, A., Powell, T., Sands, R. D., Schaldach, R., Stehfest, E., Steinbuks, J., Tabeau, A., van Meijl, H., Wise, M. A. and Rounsevell, M. D. A. (2017) 'Assessing uncertainties in land cover projections', *Global Change Biology*, 23(2), pp. 767–781. doi: 10.1111/gcb.13447.
- Alexander, P., Rabin, S., Anthoni, P., Henry, R., Pugh, T., Rounsevell, M. and Arneth, A. (2018) 'Adaptation of global land use and management intensity to changes in climate and atmospheric carbon dioxide', *Global Change Biology*.
- Alexandratos, N. and Bruinsma, J. (2012) 'World agriculture towards 2030/2050: the 2012 revision', FAO, Rome: ESA Working paper, 12(3). doi: 10.1016/S0264-8377(03)00047-4.
- Alliance for Zero Extinction (2010) 'Available at: www.zeroextinction.org. Accessed 2/6/2018'.
- Arino, O., Ramos, J., Kalogirou, V., Defourny, P. and Achard, F. (2010) 'GlobCover 2009. ESA Living Planet Symposium, 27 June - 2 July 2010, Bergen, Norway'.
- Baillie, J., Hilton-Taylor, C. and Stuart, S. N. (2004) '2004 IUCN red list of threatened species: a global species assessment', IUCN. doi: 10.2305/IUCN.CH.2005.3.en.
- Bajželj, B., Richards, K. S., Allwood, J. M., Smith, P., Dennis, J. S., Curmi, E. and Gilligan, C. A. (2014) 'Importance of food-demand management for climate mitigation', *Nature Climate Change*, 4(10), pp. 924–929. doi: 10.1038/nclimate2353.
- Chaudhary, A. and Mooers, A. O. (2018) 'Terrestrial vertebrate biodiversity loss under future global land use change scenarios', *Sustainability*, 10(8), p. 2768. doi: 10.3390/su10082764.
- Delzeit, R., Zabel, F., Meyer, C. and Václavík, T. (2017) 'Addressing future trade-offs between biodiversity and cropland expansion to improve food security', *Regional Environmental Change*, 17(5), pp. 1429–1441. doi: 10.1007/s10113-016-0927-1.
- Doelman, J. C., Stehfest, E., Tabeau, A., van Meijl, H., Lassaletta, L., Gernaat, D. E. H. J., Hermans, K., Harmsen, M., Daioglou, V., Biemans, H., van der Sluis, S. and van Vuuren, D. P. (2018) 'Exploring SSP land-use dynamics using the IMAGE model: Regional and gridded scenarios of land-use change and land-based climate change mitigation', *Global Environmental Change*. Elsevier Ltd, 48, pp. 119–135. doi: 10.1016/j.gloenvcha.2017.11.014.
- Don, A., Schumacher, J. and Freibauer, A. (2011) 'Impact of tropical land-use change on soil organic carbon stocks - a meta-analysis', *Global Change Biology*, 17(4), pp. 1658–1670. doi: 10.1111/j.1365-2486.2010.02336.x.
- Dumollard, G., Herrero, M. and Havlik, P. (2012) 'Climate change, agriculture and food security: a comparative review of global modelling approaches. CCAFS Working Paper no. 34.', CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS). Copenhagen, Denmark. Available online at: www.ccafs.cgiar.org.
- Eitelberg, D. A., van Vliet, J. and Verburg, P. H. (2015) 'A review of global potentially available cropland estimates and their consequences for model-based assessments', *Global Change Biology*, 21(3), pp. 1236–1248. doi: 10.1111/gcb.12733.
- FAO/IIASA/ISRIC/ISS-CAS/JRC (2009) 'Harmonized World Soil Database (version 1.1). FAO, Rome, Italy and IIASA, Laxenburg, Austria.'
- Foley, J. A., DeFries, R., Asner, G. P., Barford, C., Bonan, G., Carpenter, S. R., Chapin, F. S., Coe, M. T., Daily, G. C., Gibbs, H. K., Helkowski, J. H., Holloway, T., Howard, E. A., Kucharik, C. J., Monfreda, C., Patz, J. A., Prentice, I.

- C., Ramankutty, N., Snyder, P. K. and Land (2005) 'Global Consequences of Land Use', *Science*, 309(5734), pp. 570–574. doi: 10.1126/science.1111772.
- Foley, J. A., Ramankutty, N., Brauman, K. a., Cassidy, E. S., Gerber, J. S., Johnston, M., Mueller, N. D., O'Connell, C., Ray, D. K., West, P. C., Balzer, C., Bennett, E. M., Carpenter, S. R., Hill, J., Monfreda, C., Polasky, S., Rockström, J., Sheehan, J., Siebert, S., Tilman, D. and Zaks, D. P. M. (2011) 'Solutions for a cultivated planet', *Nature*, 478(7369), pp. 337–342. doi: 10.1038/nature10452.
- Fricko, O., Havlik, P., Rogelj, J., Klimont, Z., Gusti, M., Johnson, N., Kolp, P., Strubegger, M., Valin, H., Amann, M., Ermolieva, T., Forsell, N., Herrero, M., Heyes, C., Kindermann, G., Krey, V., McCollum, D. L., Obersteiner, M., Pachauri, S., Rao, S., Schmid, E., Schoepp, W. and Riahi, K. (2017) 'The marker quantification of the Shared Socioeconomic Pathway 2: A middle-of-the-road scenario for the 21st century', *Global Environmental Change*. Elsevier Ltd, 42, pp. 251–267. doi: 10.1016/j.gloenvcha.2016.06.004.
- Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F., Pretty, J., Robinson, S., Thomas, S. M. and Toulmin, C. (2010) 'Food Security: The Challenge of Feeding 9 Billion People', *Science*, 327(5967), pp. 812–818. doi: 10.1126/science.1185383.
- Guo, L. B. and Gifford, R. M. (2002) 'Soil carbon stocks and land use change: A meta analysis', *Global Change Biology*, 8(4), pp. 345–360. doi: 10.1046/j.1354-1013.2002.00486.x.
- Hannah, L., Ikegami, M., Hole, D. G., Seo, C., Butchart, S. H. M., Peterson, A. T. and Roehrdanz, P. R. (2013) 'Global climate change adaptation priorities for biodiversity and food security.', *PloS one*, 8(8), p. e72590. doi: 10.1371/journal.pone.0072590.
- Hansen, M. C., Stehman, S. V. and Potapov, P. V. (2010) 'Quantification of global gross forest cover loss', *Proceedings of the National Academy of Sciences*, 107(19), pp. 8650–8655. doi: 10.1073/pnas.0912668107.
- Havlik, P., Schneider, U. A., Schmid, E., Bottcher, H., Fritz, S., Skalsky, R., Aoki, K., De Cara, S., Kindermann, G., Kraxner, F., Leduc, S., Mccallum, I., Mosnier, A., Sauer, T. and Obersteiner, M. (2011) 'Global land-use implications of first and second generation biofuel targets', *Energy Policy*, 39, pp. 5690–5702. doi: 10.1016/j.enpol.2010.03.030.
- Henry, R. C., Engström, K., Olin, S., Alexander, P., Arneth, A. and Rounsevell, M. D. A. (2018) 'Food supply and bioenergy production within the global cropland planetary boundary', *PLoS ONE*, 13(3), pp. 1–17. doi: 10.1371/journal.pone.0194695.
- Hiederer, R. and Köchy, M. (2011) 'Global soil organic carbon estimates and the harmonized world soil database: European Commission Joint Research Centre Scientific and Technical Research series.', EUR 25225EN, Publications Office of the European Union, p. p.79. doi: 10.2788/13267.
- Houghton, R. A., House, J. I., Pongratz, J., van der Werf, G. R., DeFries, R. S., Hansen, M. C., Le Quéré, C. and Ramankutty, N. (2012) 'Carbon emissions from land use and land-cover change', *Biogeosciences*, 9(12), pp. 5125–5142. doi: 10.5194/bg-9-5125-2012.
- IPCC (2001) 'Climate Change 2001: Impacts, Adaptation, and Vulnerability, Contribution of Working Group II to the Third Assessment Report of the Intergovernmental Panel on Climate Change (eds McCarthy, J. J., Canziani, O. F., Leary, N. A., Dokken, D. J. & White, K. S.)', Cambridge Univ. Press, Cambridge, pp. 105–143. doi: 10.1021/bc700444t.
- IPCC (2014) 'Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Edenhofer, O., R. Pichs-Madruga, Y. Sokona, E. Farahani, S. Kadner, K. Seyboth, A. Adler', Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. doi: 10.1017/CBO9781107415416.
- Jenkins, C. N., Pimm, S. L. and Joppa, L. N. (2013) 'Global patterns of terrestrial vertebrate diversity and conservation', *Proceedings of the National Academy of Sciences*, 110(28), pp. E2602–E2610. doi: 10.1073/pnas.1302251110.
- Von Lampe, M., Willenbockel, D., Ahammad, H., Blanc, E., Cai, Y., Calvin, K., Fujimori, S., Hasegawa, T., Havlik, P., Heyhoe, E., Kyle, P., Lotze-Campen, H., Mason d'Croz, D., Nelson, G. C., Sands, R. D., Schmitz, C., Tabeau, A., Valin, H., van der Mensbrugghe, D. and van Meijl, H. (2014) 'Why do global long-term scenarios for agriculture

- differ? An overview of the AgMIP global economic model intercomparison', *Agricultural Economics*, 45(1), pp. 3–20. doi: 10.1111/agec.12086.
- McDonald, R. I., Kareiva, P. and Forman, R. T. T. (2008) 'The implications of current and future urbanization for global protected areas and biodiversity conservation', *Biological Conservation*, 141(6), pp. 1695–1703. doi: 10.1016/j.biocon.2008.04.025.
- Mittermeier, R., Robles Gil, P., Hoffmann, M., Pilgrim, J., Brooks, T., Goettsch Mittermeier, C., Lamoreux, J. and Fonseca, G. (2004) *Hotspots revisited. Earth's biologically richest and most endangered terrestrial ecoregions.*, Conservation International.
- Molotoks, A., Stehfest, E., Doelman, J., Albanito, F., Fitton, N., Dawson, T. P. and Smith, P. (2018) 'Global projections of future cropland expansion to 2050 and direct impacts on biodiversity and carbon storage', *Global Change Biology*, 24(12), pp. 5895–5908. doi: 10.1111/gcb.14459.
- Müller, C., Stehfest, E., Van Minnen, J. G., Strengers, B., Von Bloh, W., Beusen, A. H. W., Schaphoff, S., Kram, T. and Lucht, W. (2016) 'Drivers and patterns of land biosphere carbon balance reversal', *Environmental Research Letters*. IOP Publishing, 11(4). doi: 10.1088/1748-9326/11/4/044002.
- Myers, N., Mittermeier, R. A., Mittermeier, C. G., Da Fonseca, G. A. B. and Kent, J. (2000) 'Biodiversity hotspots for conservation priorities', *Nature*. Nature Publishing Group, 403(6772), p. 853.
- Nelson, G.C., Van Der Mensbrugghe, D., Ahammad, H., Blanc, E., Calvin, K., Hasegawa, T., Havlik, P., Heyhoe, E., Kyle, P., Lotze-Campen, H. and von Lampe, M. (2014) 'Agriculture and climate change in global scenarios: why don't the models agree', *Agricultural Economics*, 45(1), pp.85-101. doi:10.1111/agec.12091
- O'Neill, B. C., Kriegler, E., Ebi, K. L., Kemp-Benedict, E., Riahi, K., Rothman, D. S., van Ruijven, B. J., van Vuuren, D. P., Birkmann, J., Kok, K., Levy, M. and Solecki, W. (2017) 'The roads ahead: Narratives for shared socioeconomic pathways describing world futures in the 21st century', *Global Environmental Change*. Elsevier Ltd, 42, pp. 169–180. doi: 10.1016/j.gloenvcha.2015.01.004.
- O'Neill, B. C., Kriegler, E., Riahi, K., Ebi, K. L., Hallegatte, S., Carter, T. R., Mathur, R. and van Vuuren, D. P. (2014) 'A new scenario framework for climate change research: The concept of shared socioeconomic pathways', *Climatic Change*, 122(3), pp. 387–400. doi: 10.1007/s10584-013-0905-2.
- Parmesan, C. and Yohe, G. (2003) 'A globally coherent fingerprint of climate change impacts across natural systems', *Nature*, 421, pp. 37–42.
- Pimm, S. L., Jenkins, C. N., Abell, R., Brooks, T. M., Gittleman, J. L., Joppa, L. N., Raven, P. H., Roberts, C. M. and Sexton, J. O. (2014) 'The biodiversity of species and their rates of extinction, distribution, and protection', *Science*, 344(6187). doi: 10.1126/science.1246752.
- Pimm, S. L. and Raven, P. (2000) 'Extinction by numbers', *Nature*, 403(6772), pp. 843–845. doi: 10.1038/35002708.
- Popp, A., Calvin, K., Fujimori, S., Havlik, P., Humpenöder, F., Stehfest, E., Bodirsky, B. L., Dietrich, J. P., Doelmann, J. C., Gusti, M., Hasegawa, T., Kyle, P., Obersteiner, M., Tabeau, A., Takahashi, K., Valin, H., Waldhoff, S., Weindl, I., Wise, M., Kriegler, E., Lotze-Campen, H., Fricko, O., Riahi, K. and Vuuren, D. P. va. (2017) 'Land-use futures in the shared socio-economic pathways', *Global Environmental Change*. Elsevier Ltd, 42, pp. 331–345. doi: 10.1016/j.gloenvcha.2016.10.002.
- Popp, A., Rose, S. K., Calvin, K., Van Vuuren, D. P., Dietrich, J. P., Wise, M., Stehfest, E., Humpenöder, F., Kyle, P., Van Vliet, J., Bauer, N., Lotze-Campen, H., Klein, D. and Kriegler, E. (2014) 'Land-use transition for bioenergy and climate stabilization: Model comparison of drivers, impacts and interactions with other land use based mitigation options', *Climatic Change*, 123(3–4), pp. 495–509. doi: 10.1007/s10584-013-0926-x.
- Potapov, P. V., Turubanova, S. A., Hansen, M. C., Adusei, B., Broich, M., Altstatt, A., Mane, L. and Justice, C. O. (2012) 'Quantifying forest cover loss in Democratic Republic of the Congo, 2000-2010, with Landsat ETM+ data', *Remote Sensing of Environment*. Elsevier Inc., 122, pp. 106–116. doi: 10.1016/j.rse.2011.08.027.
- Prestele, R., Alexander, P., Rounsevell, M. D. A., Arneth, A., Calvin, K., Doelman, J., Eitelberg, D. A., Engström, K., Fujimori, S., Hasegawa, T., Havlik, P., Humpenöder, F., Jain, A. K., Krisztin, T., Kyle, P., Meiyappan, P., Popp, A., Sands, R. D., Schaldach, R., Schüngel, J., Stehfest, E., Tabeau, A., Van Meijl, H., Van Vliet, J. and Verburg, P. H.

- (2016) 'Hotspots of uncertainty in land-use and land-cover change projections: a global-scale model comparison', *Global Change Biology*, 22(12), pp. 3967–3983. doi: 10.1111/gcb.13337.
- Purvis, A., Jones, K. E. and Mace, G. M. (2000) 'Extinction', *BioEssays*, 22(12), pp. 1123–1133.
- Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaserna, J. C., KC, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T., Havlik, P., Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J., Strefler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A. and Tavoni, M. (2017) 'The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview', *Global Environmental Change*, 42, pp. 153–168. doi: 10.1016/j.gloenvcha.2016.05.009.
- Ricketts, T. H., Dinerstein, E., Boucher, T., Brooks, T. M., Butchart, S. H. M., Hoffmann, M., Lamoreux, J. F., Morrison, J., Parr, M., Pilgrim, J. D., Rodrigues, A. S. L., Sechrest, W., Wallace, G. E., Berlin, K., Bielby, J., Burgess, N. D., Church, D. R., Cox, N., Knox, D., Loucks, C., Luck, G. W., Master, L. L., Moore, R., Naidoo, R., Ridgely, R., Schatz, G. E., Shire, G., Strand, H., Wettengel, W. and Wikramanayake, E. (2005) 'Pinpointing and preventing imminent extinctions', *Proceedings of the National Academy of Sciences of the United States of America*, 102(51), pp. 18497–18501.
- Ruesch, A. and Gibbs, H. K. (2008) 'New IPCC Tier-1 Global Biomass Carbon Map for the Year 2000.' Oak Ridge National Laboratory, Carbon Dioxide Information Analysis Center, Oak Ridge, TN. Available at: <http://cdiac.ornl.gov>.
- Schmitz, C., van Meijl, H., Kyle, P., Nelson, G. C., Fujimori, S., Gurgel, A., Havlik, P., Heyhoe, E., d'Croz, D. M., Popp, A., Sands, R., Tabeau, A., van der Mensbrugghe, D., von Lampe, M., Wise, M., Blanc, E., Hasegawa, T., Kavallari, A. and Valin, H. (2014) 'Land-use change trajectories up to 2050: Insights from a global agro-economic model comparison', *Agricultural Economics (United Kingdom)*, 45(1), pp. 69–84. doi: 10.1111/agec.12090.
- Sheil, D., Ladd, B., Silva, L. C. R., Laffan, S. W. and Van Heist, M. (2016) 'How are soil carbon and tropical biodiversity related?', *Environmental Conservation*, 43(3), pp. 231–241. doi: 10.1017/S0376892916000011.
- Smith, P. (2013) 'Delivering food security without increasing pressure on land', *Global Food Security*. Elsevier, 2(1), pp. 18–23. doi: 10.1016/j.gfs.2012.11.008.
- Smith, P., Davis, S. J., Creutzig, F., Fuss, S., Minx, J., Gabrielle, B., Kato, E., Jackson, R. B., Cowie, A., Kriegler, E., Van Vuuren, D. P., Rogelj, J., Ciais, P., Milne, J., Canadell, J. G., McCollum, D., Peters, G., Andrew, R., Krey, V., Shrestha, G., Friedlingstein, P., Gasser, T., Grubler, A., Heidug, W. K., Jonas, M., Jones, C. D., Kraxner, F., Littleton, E., Lowe, J., Moreira, J. R., Nakicenovic, N., Obersteiner, M., Patwardhan, A., Rogner, M., Rubin, E., Sharifi, A., Torvanger, A., Yamagata, Y., Edmonds, J. and Yongsung, C. (2016) 'Biophysical and economic limits to negative CO₂ emissions', *Nature Climate Change*. Nature Publishing Group, 6(1), pp. 42–50. doi: 10.1038/nclimate2870.
- Smith, P., Price, J., Molotoks, A., Warren, R. and Malhi, Y. (2018) 'Impacts on terrestrial biodiversity of moving from a 2°C to a 1.5°C target', *Philosophical Transactions of the Royal Society A*, 376(20160456).
- Stehfest, E., Vuuren, D. van, Kram, T., Bouwman, L., Alkemade, R., Bakkenes, M., Biemans, H., Bouwman, A., Elzen, M. den, Janse, J., Lucas, P., Minnen, J. van, Müller, C. and Prins, A. (2014) *Integrated Assessment of Global Environmental Change with IMAGE 3.0. Model description and policy applications*, PBL Netherlands Environmental Assessment Agency, The Hague.
- Strassburg, B. B. N., Kelly, A., Balmford, A., Davies, R. G., Gibbs, H. K., Lovett, A., Miles, L., Orme, C. D. L., Price, J., Turner, R. K. and Rodrigues, A. S. L. (2010) 'Global congruence of carbon storage and biodiversity in terrestrial ecosystems', *Conservation Letters*, 3(2), pp. 98–105. doi: 10.1111/j.1755-263X.2009.00092.x.
- Titeux, N., Henle, K., Mihoub, J. B., Regos, A., Geijzenborffer, I. R., Cramer, W., Verburg, P. H. and Brotons, L. (2016) 'Biodiversity scenarios neglect future land-use changes', *Global change biology*, 22(7), pp. 2505–2515. doi: 10.1111/gcb.13272.

Tscharntke, T., Klein, A. M., Kruess, A., Steffan-Dewenter, I. and Thies, C. (2005) 'Landscape perspectives on agricultural intensification and biodiversity on ecosystem service management', *Ecology Letters*, 8(8), pp. 857–874. doi: 10.1111/j.1461-0248.2005.00782.x.

Visconti, P., Pressey, R. L., Giorgini, D., Maiorano, L., Bakkenes, M., Boitani, L., Alkemade, R., Falcucci, A., Chiozza, F. and Rondinini, C. (2011) 'Future hotspots of terrestrial mammal loss.', *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 366(1578), pp. 2693–2702. doi: 10.1098/rstb.2011.0105.

van Vuuren, D. P. and Carter, T. R. (2014) 'Climate and socio-economic scenarios for climate change research and assessment: Reconciling the new with the old', *Climatic Change*, 122(3), pp. 415–429. doi: 10.1007/s10584-013-0974-2.

van Vuuren, D. P., Stehfest, E., Gernaat, D. E. H. J., Doelman, J. C., van den Berg, M., Harmsen, M., de Boer, H. S., Bouwman, L. F., Daioglou, V., Edelenbosch, O. Y., Girod, B., Kram, T., Lassaletta, L., Lucas, P. L., van Meijl, H., Müller, C., van Ruijven, B. J., van der Sluis, S. and Tabeau, A. (2017) 'Energy, land-use and greenhouse gas emissions trajectories under a green growth paradigm', *Global Environmental Change*. Elsevier Ltd, 42, pp. 237–250. doi: 10.1016/j.gloenvcha.2016.05.008.

Warren, R., Vanderwal, J., Price, J., Welbergen, J. A., Atkinson, I., Ramirez-Villegas, J., Osborn, T. J., Jarvis, A., Shoo, L. P., Williams, S. E. and Lowe, J. (2013) 'Quantifying the benefit of early climate change mitigation in avoiding biodiversity loss', *Nature Climate Change*. Nature Publishing Group, 3(7), pp. 678–682. doi: 10.1038/nclimate1887.

West, P. C., Gibbs, H. K., Monfreda, C., Wagner, J., Barford, C. C., Carpenter, S. R. and Foley, J. A. (2010) 'Trading carbon for food: Global comparison of carbon stocks vs. crop yields on agricultural land', *Proceedings of the National Academy of Sciences*, 107(46), pp. 19645–19648. doi: 10.1073/pnas.1011078107.

Woltjer, G. B. and Kuiper, M. . (2014) 'The MAGNET model: Module description', Wageningen: LEI 877 Wageningen UR., p. 144. Available at: www.wageningenUR.nl/en/lei.

Zhao, C., Liu, B., Piao, S., Wang, X., Lobell, D. B., Huang, Y., Huang, M., Yao, Y., Bassu, S., Ciais, P., Durand, J.-L., Elliott, J., Ewert, F., Janssens, I. A., Li, T., Lin, E., Liu, Q., Martre, P., Müller, C., Peng, S., Peñuelas, J., Ruane, A. C., Wallach, D., Wang, T., Wu, D., Liu, Z., Zhu, Y., Zhu, Z. and Asseng, S. (2017) 'Temperature increase reduces global yields of major crops in four independent estimates', *Proceedings of the National Academy of Sciences*, 114(35), pp. 9326–9331. doi: 10.1073/pnas.1701762114.

Tables

Table 1.

Metric	Proportion of variance explained by		R ²
	Model	SSP	
AZE sites	63.5	21.4	0.849
Carbon loss from biomass	69.7	25.0	0.947
Carbon loss from soil	62.1	27.2	0.893
Amphibian spp. rich hotspots	63.5	23.0	0.864
Bird spp. rich hotspots	75.3	19.7	0.949
Mammal spp. rich hotspots	68.3	22.1	0.904
CI Hotspots	83.9	11.2	0.951

Figure and table captions

Table 1. The proportion of variance explained by the model and SSP for each of the biodiversity metrics considered. The R² value for the linear model for each metric is given. P-values are not used as linear models were not used to identify whether model or SSP has a statistically significant effect on the biodiversity metrics examined.

Figure 1. Panels a to c show the number of AZE sites impacted by cropland expansion between 2010-2050 for each region and model by socio-economic scenario (SSP1-3). Panel d shows a comparison between models at global level.

Figure 2. Projected cropland change between 2010 and 2050 in (a) bird, (b) mammal, and (c) amphibian species rich hotspots, and (d) CI hotspots across the different SSP scenarios and models. Species-rich regions are comprised of cells with a richness index ≥ 0.9 .

Figure 3. Spatial distribution of regions of threat; regions with high biodiversity under pressure from cropland expansion. Calculated in each 0.5° grid cell as the fraction of a grid cell converted to cropland between 2010 and 2050 multiplied by the summed richness index of birds, mammals, and amphibians. The different SSPs are displayed across the rows and the different models are displayed in the columns. Blue dashed lines delineate the tropics.

Figure 4. Panels a to c show carbon loss from soil and biomass as a result of cropland expansion between 2010-2050 for each region and model by socio-economic scenario (SSP1-3). Panel d shows a comparison between models at global level




Figure 1. Panels a to c show the number of AZE sites impacted by cropland expansion between 2010-2050 for each region and model by socio-economic scenario (SSP1-3). Panel d shows a comparison between models at global level.

254x190mm (300 x 300 DPI)




Figure 2. Projected cropland change between 2010 and 2050 in (a) bird, (b) mammal, and (c) amphibian species rich hotspots, and (d) CI hotspots across the different SSP scenarios and models. Species-rich regions are comprised of cells with a richness index ≥ 0.9 .

338x190mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 3. Spatial distribution of regions of threat; regions with high biodiversity under pressure from cropland expansion. Calculated in each 0.5° grid cell as the fraction of a grid cell converted to cropland between 2010 and 2050 multiplied by the summed richness index of birds, mammals, and amphibians. The different SSPs are displayed across the rows and the different models are displayed in the columns. Blue dashed lines delineate the tropics.

508x355mm (200 x 200 DPI)




Figure 4. Panels a to c show carbon loss from soil and biomass as a result of cropland expansion between 2010-2050 for each region and model by socio-economic scenario (SSP1-3). Panel d shows a comparison between models at global level.

254x190mm (300 x 300 DPI)