



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/154048/>

Version: Accepted Version

Article:

Cruz, S., Wimmer, K., Bender, P. C. et al. (2019) Single-particle structure of neutron-rich Sr isotopes via $^2\text{H}(^{94,95,96}\text{Sr}, p)$ reactions. *Physical Review C - Nuclear Physics*. 054321. ISSN: 2469-9993

<https://doi.org/10.1103/PhysRevC.100.054321>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Single-Particle Structure of Neutron-Rich Sr Isotopes Via $d(^{94,95,96}\text{Sr}, p)$ Reactions

S. Cruz,^{1,2} K. Wimmer,^{3,4,*} P.C. Bender,² R. Krücken,^{1,2} G. Hackman,² F. Ames,² C. Andreoiu,⁵ R.A.E. Austin,⁶ C.S. Bancroft,³ R. Braid,⁷ T. Bruhn,² W.N. Catford,⁸ A. Cheeseman,² A. Chester,⁹ D.S. Cross,⁵ C.Aa. Diget,¹⁰ T. Drake,¹¹ A.B. Garnsworthy,² R. Kanungo,^{2,6} A. Knapton,⁸ W. Korten,^{12,2} K. Kuhn,⁷ J. Lassen,² R. Laxdal,² M. Marchetto,² A. Matta,^{8,13} D. Miller,² M. Moukaddam,² N.A. Orr,¹³ N. Sachmpazidi,³ A. Sanetullaev,^{6,2} C.E. Svensson,¹⁴ N. Terpstra,³ C. Unsworth,² and P.J. Voss⁵

¹*Department of Physics and Astronomy, University of British Columbia, Vancouver, BC V6T 1Z4, Canada*

²*TRIUMF, Vancouver, BC V6T 2A3, Canada*

³*Department of Physics, Central Michigan University, Mt Pleasant, MI 48859, USA*

⁴*Department of Physics, The University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113-0033, Japan*

⁵*Department of Chemistry, Simon Fraser University, Burnaby, BC V5A 1S6, Canada*

⁶*Department of Astronomy and Physics, Saint Mary's University, Halifax, NS B3H 3C2, Canada*

⁷*Department of Physics, Colorado School of Mines, Golden, CO 80401, USA*

⁸*Department of Physics, University of Surrey, Guildford, Surrey, GU2 7XH, United Kingdom*

⁹*Department of Physics, Simon Fraser University, Burnaby, BC V5A 1S6, Canada*

¹⁰*Department of Physics, University of York, York, YO10 5DD, United Kingdom*

¹¹*Department of Physics, University of Toronto, Toronto, ON M5S 1A7, Canada*

¹²*IRFU, CEA, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France*

¹³*LPC, ENSICAEN, CNRS/IN2P3, UNICAEN, Normandie Université, 14050 Caen cedex, France*

¹⁴*Department of Physics, University of Guelph, Guelph, ON, N1G 2W1, Canada*

(Dated: September 22, 2019)

Background: The region around neutron number $N = 60$ in the neutron-rich Sr and Zr nuclei is one of the most dramatic examples of a ground state shape transition from (near) spherical below $N = 60$ to strongly deformed shapes in the heavier isotopes.

Purpose: The single-particle structure of $^{95-97}\text{Sr}$ approaching the ground state shape transition at ^{98}Sr has been investigated via single-neutron transfer reactions using the (d, p) reaction in inverse kinematics. These reactions selectively populate states with a large overlap of the projectile ground state coupled to a neutron in a single-particle orbital.

Method: Radioactive $^{94,95,96}\text{Sr}$ nuclei with energies of 5.5 AMeV were used to bombard a CD_2 target. Recoiling light charged particles and γ rays were detected using a quasi- 4π silicon strip detector array and a 12 element Ge array. The excitation energy of states populated was reconstructed employing the missing mass method combined with γ -ray tagging and differential cross sections for final states were extracted.

Results: A reaction model analysis of the angular distributions allowed for firm spin assignments to be made for the low-lying 352, 556 and 681 keV excited states in ^{95}Sr and a constraint has been placed on the spin of the higher-lying 1666 keV state. Angular distributions have been extracted for 10 states populated in the $d(^{95}\text{Sr}, p)^{96}\text{Sr}$ reaction, and constraints have been provided for the spins and parities of several final states. Additionally, the 0, 167 and 522 keV states in ^{97}Sr were populated through the $d(^{96}\text{Sr}, p)$ reaction. Spectroscopic factors for all three reactions were extracted.

Conclusions: Results are compared to shell model calculations in several model spaces and the structure of low-lying states in ^{94}Sr and ^{95}Sr is well-described. The spectroscopic strength of the 0^+ and 2^+ states in ^{96}Sr is significantly more fragmented than predicted. The spectroscopic factors for the $d(^{96}\text{Sr}, p)^{97}\text{Sr}$ reaction suggest that the two lowest lying excited states have significant overlap with the weakly deformed ground state of ^{96}Sr , but the ground state of ^{97}Sr has a different structure.

I. INTRODUCTION

An atomic nucleus can deform its shape in order to minimize its energy. This is observed across the nuclear landscape, both in ground states and excited states. Indeed, it seems that even a small number of valence protons and neutrons outside of a closed core can drive the whole nucleus into a deformed shape. The long-range attractive residual proton-neutron ($p-n$) interaction allows the nucleus to gain additional binding energy by

arranging the nucleons in certain ways across the valence orbitals, which in turn causes a departure from sphericity [1]. The expense of such re-arrangements is dependent on the size of the energy gaps between single-particle orbitals above the Fermi energy. If the energy spacing is small, the valence nucleons can scatter into valence orbitals which are above the Fermi energy and drive the nucleus into a low-energy deformed configuration. On the other hand, if the energy spacing is large, the valence nucleons are unable to scatter into higher orbitals and this favors spherical shapes. The size of these energy gaps is in turn dependent on the number of valence nucleons, due to the monopole component of the residual

* Corresponding author: wimmer@phys.s.u-tokyo.ac.jp

interaction. Clearly, the underlying shell structure of nuclei plays an important role in the propensity for nuclei to deform.

The evolution of ground state shapes across an isotopic chain is commonly observed to be a gradual process, although in some cases the shape can change dramatically with the addition of just a few nucleons. A striking example of this has been observed across the Sr and Zr isotopic chains, where an abrupt change of shape in the ground states takes place at $N \sim 60$. The ground state shape transition has been measured directly using laser spectroscopy, as a sudden increase in charge radii at $N = 60$ [2]. This is also evidenced by the sudden drop in 2_1^+ energies across the even-even isotopes at $N \geq 60$, which indicates that the ground state shape changes from a nearly spherical structure to a strongly deformed prolate ($\beta \sim 0.4$) structure [3]. Recent Coulomb excitation measurements have established that the ground state of ^{96}Sr and the 0_2^+ state in ^{98}Sr possess similar structures which, assuming axial symmetry, correspond to weakly deformed shapes with $\beta \sim 0.1$ [4]. In the $N = 56$ isotope ^{94}Sr , recent re-determination of the $B(E2; 2_1^+ \rightarrow 0_1^+)$ value from a lifetime measurement [5] supports the interpretation that the ground state in ^{94}Sr is close to spherical. Taken together, these measurements point towards a gradual evolution in shape up to $N \sim 58$ with $\beta \leq 0.1$ which then rapidly changes at $N = 60$ to $\beta \sim 0.4$ for the ground state. However, the degree of deformation in the ground state of the $N = 59$ nucleus ^{97}Sr is not well understood although the spin and parity of the ground state has been established as $1/2^+$, which is not expected within the spherical shell model. The magnetic moments of the $^{95,97}\text{Sr}$ ground states were reported to be very similar through laser spectroscopy [2] and deviate from the shell model expectation.

Also of interest is the emergence of shape-coexisting states in the vicinity of $N \sim 60$ and $Z \sim 40$. A very strong $E0$ transition between the 1229 and 1465 keV excited 0^+ states in ^{96}Sr , with $\rho^2(E0) = 0.185(50)$ [6] is a strong indicator of mixing between states which have different intrinsic deformations. Enhanced $E0$ transition strengths between low-lying 0^+ states have also been observed in the nearby nuclei ^{98}Sr , ^{98}Zr , ^{100}Zr , ^{100}Mo and ^{102}Mo [7].

The $N \sim 60$, $Z \sim 40$ region of the nuclear chart has been the subject of substantial interest theoretically for many years [8–27]. It has been shown that the emergence of deformed low-energy configurations can be explained in the shell model by the evolution of single-particle structure and the interaction between protons and neutrons in certain valence orbitals, namely the spin-orbit partner orbitals $\pi 0g_{9/2}$ and $\nu 0g_{7/2}$ [9, 10]. State-of-the-art beyond mean field calculations have been able to reproduce the observed shape transition at $N = 60$ in Sr, Zr and Mo [20, 21], although correctly predicting the ground state spins and parities of the odd-mass isotopes remains a challenge. Ultimately, advances in theoretical models are limited by the experimental data that is avail-

able. While numerous experiments have provided useful information on the Sr isotopes [2, 4, 28–34], a firm understanding of the underlying single-particle configurations of low-energy states is essential for a detailed description of this region. This situation motivated a series of single-neutron transfer reactions across the neutron-rich Sr isotopes $^{94,95,96}\text{Sr}$. The main results for the $d(^{95}\text{Sr}, p)$ reaction were already presented in [35]. The present paper discusses the details of the experiment and the analysis as well as further results.

II. EXPERIMENTAL SETUP AND CONDITIONS

The experiments were performed at the TRIUMF-ISAC-II facility [36]. The $d(^{94}\text{Sr}, p)$ and $d(^{95,96}\text{Sr}, p)$ measurements were the first high mass ($A > 30$) experiments with a re-accelerated secondary beam to be performed at TRIUMF. The Sr beams were produced by impinging a 480 MeV proton beam on a thick Uranium Carbide (UC_x) target. Sr atoms diffusing out of the UC_x target were selectively ionized into a singly charged (1^+) state using the TRIUMF Resonant Ionization Laser Ion Source [36] in order to enhance the extraction rate of the Sr species compared to surface-ionized contaminants, also produced within the production target. The cocktail beam was then sent through the ISAC mass separator [36] to produce a beam containing only isotopes of the same A (94, 95, 96). The beam was then transported to the Charge State Booster where the isotopes were charge-bred by an Electron Cyclotron Resonance plasma source to a higher charge state (see Table I for details). This was necessary so that the beam could next be sent to the Radio-Frequency Quadrupole (RFQ), which accepts a maximum mass-to-charge ratio (A/q) of 30 [36]. Inside the RFQ, time-dependent electric fields were tuned to accelerate the specific A/q of Sr ions. Contaminant isotopes in the beam were mismatched with the acceleration phase of the RFQ and so did not undergo any acceleration. Following the RFQ, these contaminants were deflected out of the beam using the bending dipole magnets in the accelerator chain. The beams were transported to the ISAC-II facility where their kinetic energy was increased to 5.5 A MeV using the superconducting linear accelerator [36]. Finally, the beams were transported to the experimental station where they impinged upon 0.5 mg/cm^2 deuterated polyethylene (CD_2) targets, mounted in the center of the SHARC silicon detector array [37].

SHARC (Silicon Highly-segmented Array for Reactions and Coulex) is a compact arrangement of double-sided silicon strip detectors which is optimized for high geometrical efficiency and excellent spatial resolution, with $\Delta\theta_{\text{lab}} \sim 1^\circ$ and ϕ coverage of approximately 90%. The SHARC array configuration consists of two double-sided silicon strip detector (DSSSD) box sections (DBOX and UBOX) and an annular DSSSD detector (UQQQ).

157 The downstream DBOX section, with the approximate
 158 angular range $35^\circ < \theta_{\text{lab}} < 80^\circ$, was configured us-
 159 ing a $\Delta E - E$ detector arrangement (140 μm DSSSDs
 160 and 1 mm thick unsegmented pad detectors) so that
 161 different ions could be identified (Fig. 1). For scat-
 162 tering angles $\theta_{\text{lab}} < 90^\circ$ elastic scattering of protons
 163 and deuterons overlaps with the kinematic lines of the
 164 transfer reactions requiring the particle identification. In
 165 the upstream UBOX ($95^\circ < \theta_{\text{lab}} < 140^\circ$) and UQQQ
 166 ($147^\circ < \theta_{\text{lab}} < 172^\circ$) sections, particle identification was
 167 not used as only protons are emitted with $\theta_{\text{lab}} > 90^\circ$ (as
 168 shown in Fig. 1). Background events arise from β decay
 169 of radioactive beam accidentally stopped in the scatter-
 170 ing chamber, and light particles emitted in fusion evap-
 171 oration reactions with carbon in the CD_2 target. The
 172 former can be suppressed by the particle identification
 173 cut as shown in the inset of Fig. 1 in laboratory forward
 174 direction and a cut on the detected energy in backward
 175 direction. Protons from fusion evaporation reactions con-
 176 tribute a continuous background to the excitation energy
 177 spectra. This background is more pronounced at labo-
 178 ratory forward angles due to the forward focusing of the
 179 reaction products. If unambiguous identification of the
 180 state populated in the reaction by γ -ray coincidences is
 181 possible the residual background is negligible.

182 The SHARC array was mounted in the center of the TI-
 183 GRESS γ -ray detector array [38]. In these experiments,
 184 TIGRESS was composed of 12 HPGe clover detectors
 185 arranged in a compact hemispherical arrangement with
 186 approximately 2π steradians geometrical coverage (see
 187 Fig. 2 of [39]). The individual crystals contain an elec-
 188 trical core contact and eight-fold electrical segmentation
 189 on the outer contact; four quadrants and a lateral di-
 190 vide, giving an overall 32-fold segmentation within each
 191 clover. This segmentation enhances the sensitivity to
 192 the emission angle of the γ ray to enable more precise
 193 Doppler reconstruction. For transitions from states with
 194 very short lifetimes the in-beam resolution after Doppler
 195 corrections amounts to 0.6 %. The segmented design also
 196 made it possible to improve the quality of the data taken
 197 in TIGRESS by using add-back to reconstruct full γ -ray
 198 energies from multiple scattering events. The Compton
 199 suppressor shields were not used in the present work.

200 The beam composition was measured at regular in-
 201 tervals during the experiment using a Bragg ionization
 202 detector [40], which was positioned on another beam-
 203 line adjacent to the TIGRESS experimental station. The
 204 beam composition in each experiment was also analyzed
 205 using β -decay data from the radioactive beam-like ions
 206 which were scattered onto the DQQQ (not instrumented
 207 in the present work). The primary contaminant in each
 208 beam were the isobars $^{94-96}\text{Rb}$. Contributions from non-
 209 isobaric A/q contaminants, originating from the ISAC
 210 CSB, were found to be negligible in the $A = 94$ and
 211 95 beams. However, substantial ^{17}O contamination was
 212 identified in the first half of the $A = 96$ beam-time due to
 213 challenges in beam tuning. Only the data taken during
 214 the second half of the $A = 96$ beam time was analyzed.

215 Further details regarding the beam are given in Table I.

Beam	Q [e]	Rate [s^{-1}]*	Duration [days]	Purity [%]
^{94}Sr	15^+	$\sim 3 \times 10^4$	~ 3	50(5)
^{95}Sr	16^+	$\sim 1.5 \times 10^6$	~ 2.5	95(3)
^{96}Sr	17^+	$\sim 1 \times 10^4$	~ 1	58(13)

*including contaminations

TABLE I. Summary of the $^{94,95,96}\text{Sr}$ beam properties.

III. ANALYSIS AND RESULTS

219 The SHARC and TIGRESS detectors were calibrated
 220 using standard sources. In the case of TIGRESS ^{60}Co
 221 and ^{152}Eu sources were used to obtain the energy and ef-
 222 ficiency calibrations of each detector. The ΔE detectors
 223 of SHARC were calibrated using a triple alpha source.
 224 The E detectors were calibrated using the proton and
 225 deuteron elastic scattering data, which was acquired si-
 226 multaneously with the $d(\text{Sr}, p)$ data. Fig. 1 shows the
 227 kinetic energy of measured protons and deuterons as a
 228 function of laboratory scattering angle for the ^{95}Sr beam
 229 incident on the CD_2 target. The total kinetic energy of




FIG. 1. Kinematics plot for ^{95}Sr incident on the CD_2 target, compared to calculated kinematics lines drawn for elastic scattering (black, dotted lines) and (d,p) transfer at 0, 2, 4 and 6 MeV excitation energy (red). In addition to uniquely identified particle in the DBOX, elastic scattered protons and deuterons are shown below the identification threshold of about 5000 keV identified by their kinematic $E(\theta_{\text{lab}})$ relation. The inset shows the particle identification plot for the DBOX section (see text), which was used to distinguish between protons and deuterons.

231 measured particles was reconstructed by adding calcu-
 232 lated energy losses using SRIM [41] in the target and Si
 233 detector dead layers to the energy deposited in SHARC.
 234 The energy loss correction amounted less than 100 keV
 235

for protons in laboratory forward direction as well as for scattering angles larger than 120° , and up to 500 keV for protons scattered close to 100° . Details of the calibration methods can be found in ref. [42]. The excitation energy (E_x) was reconstructed using the measured energy and scattering angle of the detected particles using the missing mass method. The excitation energy resolution of the DBOX, UBOX and UQQQ sections was determined to be approximately 550, 450 and 400 keV (FWHM) for the respective angular ranges. The primary contributions to the energy resolution were the energy loss of the beam and proton recoils in the thick target. For this reason, excited states which were less than approximately 500 keV apart could not be individually resolved. Excited states were thus identified using the de-excitation γ ray in addition to an E_x gate [43]. For low statistics cases, such as the ^{94}Sr and ^{96}Sr experiments, a constrained multi-peak fit was used to consistently extract the population strengths of unresolved adjacent states. This is discussed further in the subsequent sections.

The experimental angular distributions were compared to distorted wave Born approximation (DWBA) calculations that were carried out using the FRESKO code [44]. The optical model parameters used in the analysis were determined from fits to the elastic scattering data measured simultaneously. For the proton optical potential the data are not sensitive to the parameters and the parametrization of ref. [45] was used in the following. Several global optical model parameter sets [45–47] were compared to the (d, d) angular distributions and it was found that the parameters of Lohr and Haeberli [47], with some small adjustments, resulted in very good agreement with the combined (d, d) data for all three experiments. The combined fit for $d(^{94,95,96}\text{Sr}, d)$ can be seen in Fig. 2. It should be noted that the angular distributions shown




FIG. 2. Comparison of $d(^{94,95,96}\text{Sr}, d)$ angular distribution data to DWBA calculations using the optimized optical potential that is given in Table II. The inset shows the comparison of the $p(^{94,95,96}\text{Sr}, p)$ data to the global potential PP-76 [45] (see text).

in Fig. 2 include the contributions for the beam contamination (mainly Rb), however the parameters are expected

to vary slowly with A and Z . The parameters used in the analysis of the transfer reaction data are summarized in Table II. The overall normalization constant, required to convert the experimental cross sections into units of mb/sr, was also determined from the elastic scattering. The ratio of proton and deuteron elastic scattering in each experiment was used to determine the fraction of deuterons and protons within the CD_2 target, 96(2)%, 92(1)% and 96(2)% deuterons for the $^{94,95,96}\text{Sr}$ experiments, respectively. The uncertainties include statistical and reaction model uncertainties. The normalization constants were corrected for the beam purity and target deuteron content.

The $d(^{94,95,96}\text{Sr}, p)$ reactions were modeled as a single-step process where the transferred neutron populates an unoccupied valence orbital. By comparing the experimental cross section for each final state to the calculations, the spectroscopic factor can be extracted. In addition to the statistical uncertainty, these spectroscopic factors carry a theoretical systematic uncertainty arising from the choice of the reaction model, optical model parameters, and the potential used to calculate the nucleon bound-state wave function. By comparing different parametrizations, this uncertainty has been estimated to be 20 %. Relative spectroscopic factors are not affected by the uncertainty. In order to better gauge the uncertainty arising from the reaction modeling, adiabatic distorted wave approximation (ADWA) calculations were also performed. For the incoming channel global nucleon-nucleus optical model parameters from [48] evaluated at half the beam energy were used. The ADWA model takes the breakup of the loosely bound deuteron explicitly into account, but the reliability at the rather low beam energies of the present work is not well established. In general the ADWA results describe the shape of the angular distribution better as shown below, and result in smaller spectroscopic factors by about 15% compared to the DWBA.

By comparing the experimental angular distributions to reaction model calculations the most probable $\Delta\ell$ value was determined for each state using a χ^2 analysis. It was not possible to differentiate between the spin-orbit partner orbitals $1d_{5/2}$ and $1d_{3/2}$ (both $\Delta\ell = 2$), and so both are given as possible scenarios where applicable. The neutron $0h_{11/2}$ ($\ell = 5$) orbital was not considered here as the single-particle energy has been estimated as 3.5 MeV at ^{91}Zr [17, 22].

A. Results for the $d(^{94}\text{Sr}, p)^{95}\text{Sr}$ reaction

The γ rays and excitation energy of states in ^{95}Sr that were populated via the $d(^{94}\text{Sr}, p)$ reaction are shown in Fig. 3. Strong 329, 352 and 681 keV γ -ray lines can be seen in the E_x versus E_γ matrix. Fig. 4 shows the ^{95}Sr level scheme for states that were identified below 2 MeV. All states and transition energies were previously known. Substantial direct population of the 0, 352 and 681 keV

Data	R_c	V_0	R_0	A_0	W_D	R_D	A_D	V_{SO}	R_{SO}	A_{SO}
(d,d), This Work	1.30	109.45	1.07	0.86	10.42	1.37	0.88	7.00	0.75	0.50
(d,d), LH-74 [47]	1.30	109.45	1.05	0.86	10.42	1.43	0.77	7.00	0.75	0.50
(p,p), PP-76 [45]	1.25	58.73	1.25	0.65	13.50	1.25	0.47	7.50	1.25	0.47

TABLE II. Optical model parameters that were used to describe $^{94,95,96}\text{Sr}$ elastic scattering angular distributions in the DWBA calculations (Fig. 2). The global optical model parameters of Lohr and Haerberli (LH-74) [47], with some small adjustments were found to give the best fit to the combined (d, d) data. The global optical model parameters of Perey and Perey (PP-76) were used to describe the combined (p, p) data.




FIG. 3. Excitation energy versus γ -ray energy matrix (upper) and projected γ -ray spectrum (lower panel) for ^{95}Sr states populated via $d(^{94}\text{Sr}, p)$.

333 states was observed. There is also clear evidence for the
 334 direct population of the 1666 keV excited state through
 335 the observation of the 427 keV γ ray. This line is en-
 336 hanced in the spectrum if a gate on excitation energies
 337 $1 < E_x < 2$ MeV is placed. However, the statistics were
 338 too low for an angular distribution analysis. It is also
 339 apparent that excited states up to ~ 5 MeV were popu-
 340 lated through this reaction and decay via the 352 and
 341 681 keV states. However, it was not possible to identify
 342 any states above the 1666 keV state due to the limited
 343 statistics.

344 *The ground state of ^{95}Sr :* The ground, 352, and
 345 681 keV states were not clearly resolved in the excitation
 346 energy spectrum (Fig. 5). Therefore the angular distribu-
 347 tions were extracted simultaneously using a constrained
 348 three (Gaussian) peak-plus-exponential background fit of
 349 the excitation energy spectrum for each angular bin. An
 350 example fit is shown in Fig. 5. The peak widths and sep-




FIG. 4. Level scheme for ^{95}Sr states that were populated through $d(^{94}\text{Sr}, p)$. The 204 keV γ ray was not observed due to the 21.9(5) ns [3, 49] half-life of the 556 keV state (more details in the text).




FIG. 5. Excitation energy spectrum extracted from the recoiling proton energies and angles at a center of mass angle $\theta_{\text{cm}} = 30^\circ$. The continuous green line shows the constrained 3-peak fit of the 0, 352 and 681 keV ^{95}Sr states. The dashed line represents the continuous background.

352 arations between them were fixed using the known E_x
 353 resolution (determined with simulations and verified us-
 354 ing the the $d(^{95}\text{Sr}, p)$ data set [35]) and the energies of the
 355 states, respectively. The shape of the ground state angu-
 356 lar distribution (Fig. 6 (a)) is in good agreement with

357 the $\Delta\ell = 0$ reaction model calculations, with a spectro-
 358 scopic factor of 0.41(9) for the DWBA and 0.34(7) for the
 359 ADWA, respectively. Systematic uncertainties include
 360 the experimental sources discussed above and theoretical
 361 uncertainties arising from the optical model parameters
 362 used. Our results are thus consistent with the known
 363 $J^\pi = 1/2^+$ assignment for this state [50].




FIG. 6. Panels (a-c): Comparison of the reaction model calculations to the angular distributions for the 0, 352 and 681 keV states in ^{95}Sr . The experimental data has been obtained from the constrained 3-peak fit (Fig. 5). The solid lines are the best-fitting reaction model calculations using the DWBA (blue) and ADWA (green) methods. Panel (d): comparison of the two methods to extract the angular distribution for the 352 keV state (see text).

364

365

366 *The 352 keV state:* Two independent experimental angular
 367 distributions were produced for the 352 keV state;
 368 one was extracted using the three peak fit (see Fig. 5 (b))
 369 and a second was extracted by gating on the 352 keV γ -
 370 ray transition and the excitation energy (Fig. 6 (d)). The
 371 shape of both angular distributions are in clear agreement
 372 with the $\Delta\ell = 2$ calculation, constraining the spin and
 373 parity of this state to be $J^\pi = 3/2^+$ or $5/2^+$. Combin-
 374 ing the $\Delta\ell = 2$ angular distribution with the previously
 375 established $M1$ character of the 352 keV γ -ray transition
 376 to the ^{95}Sr ground state [3] allows a firm spin and parity
 377 assignment of $3/2^+$ for this state. The spectroscopic fac-
 378 tors for adding a neutron to the $1d_{3/2}$ orbital are 0.50(10)
 379 and 0.55(13), using the two methods respectively, using
 380 the DWBA reaction theory. The weighted average of the
 381 two spectroscopic factors is presented in Table III. As
 382 for the ground state the ADWA calculation results in a
 383 slightly lower spectroscopic factor of 0.45(7).

384 *The 556 keV state:* Although direct population of the
 385 long-lived 556 keV state ($T_{1/2} = 21.9(5)$ ns) in this ex-
 386 periment could not be confirmed owing to the low γ -ray
 387 detection efficiency due to its long lifetime, its spin and
 388 parity can be constrained by combining the $3/2^+$ assign-

389 ment for the 352 keV state from this work with previ-
 390 ous measurements. The 204 keV γ -ray transition from
 391 the 556 keV to the 352 keV state was previously deter-
 392 mined to have pure $E2$ character using conversion elec-
 393 tron spectroscopy [3]. Additionally no decay directly to
 394 the ground state has been observed in this or previous [3]
 395 work. This constrains the spin and parity of the 556 keV
 396 state to be $J^\pi = 7/2^+$. The $d(^{94}\text{Sr}, p)$ transfer reaction
 397 is not expected to populate $7/2^+$ states strongly as the
 398 large angular momentum transfer $\Delta\ell = 4$ suppresses the
 399 cross section. While no cross section or angular distri-
 400 bution could be extracted from the present data set, the
 401 spectrum in Fig. 5 shows that the direct population of
 402 this state must be small.

403 *The 681 keV state:* Three independent experimen-
 404 tal angular distributions were produced for the 681 keV
 405 state. In addition to the three peak fit result (shown
 406 in Fig. 6), angular distributions (not shown) were also
 407 produced for this state by gating on the 329 keV and
 408 681 keV transitions as well as the excitation energy. The
 409 shape of all three extracted angular distributions are in
 410 good agreement with each other and with the $\Delta\ell = 2$
 411 DWBA calculation, constraining the spin and parity of
 412 this state to be $J^\pi = 3/2^+$ or $5/2^+$. The absence of any
 413 $M1$ component in the 681 keV ground state transition [3]
 414 allows us to assign $J^\pi = 5/2^+$ to the 681 keV state. The
 415 spectroscopic factors for population of the $1d_{5/2}$ orbital
 416 that were extracted (with the DWBA calculations) using
 417 the three methods are 0.20(5), 0.14(5) and 0.14(7), re-
 418 spectively. The weighted average of these spectroscopic
 419 factors is presented in Table III. The ADWA analysis
 420 resulted in a weighted average spectroscopic factor of
 421 0.14(3).

422 *The 1666 keV state:* The observation of a 427 keV
 423 peak in Fig. 3, coincident with excitation energies in the
 424 range of $1 < E_x < 2$ MeV, establishes that the 1666 keV
 425 state was populated in the $d(^{94}\text{Sr}, p)$ reaction. This state
 426 was observed in ^{252}Cf spontaneous fission decay [51], a
 427 process which preferentially populates high spin states.
 428 In that work a tentative spin and parity of $11/2^+$ was as-
 429 signed based on the large branching ratio to the 1239 keV
 430 (tentative $9/2^+$) state. However, the population of the
 431 state in transfer makes this assignment unlikely. The ad-
 432 dition of a single neutron to the ^{94}Sr ground state via the
 433 $d(^{94}\text{Sr}, p)$ reaction can directly populate ^{95}Sr states with
 434 spins and parities of $1/2^+, 3/2^+, 5/2^+$, and $7/2^+$. The
 435 cross section for $11/2^-$ states with $\Delta\ell = 5$ is very low and
 436 is not further considered in this work. We therefore pro-
 437 pose a spin and parity of $(3/2, 5/2, 7/2)^+$ for the 1666 keV
 438 state. The angular distribution for this state could not be
 439 extracted, comparison of the integrated cross section with
 440 the DWBA and ADWA calculations suggests a spectro-
 441 scopic factor of $C^2S < 0.05$ for $\Delta\ell = 0, 2$ or $C^2S \approx 0.12$
 442 for $\Delta\ell = 4$ transfer to the $0g_{7/2}$ orbital.

B. Results for the $d(^{95}\text{Sr}, p)$ reaction

The γ rays and excitation energy of states in ^{96}Sr that were populated via the $d(^{95}\text{Sr}, p)$ reaction are shown in Fig. 7. The very strong 815 keV γ -ray line visible over




FIG. 7. Excitation energy versus γ -ray energy matrix (upper) and projected γ -ray spectrum (lower) for ^{96}Sr states populated via the $d(^{95}\text{Sr}, p)$ reaction.

the whole excitation energy range indicates that many excited states decay to the 815 keV 2_1^+ state. An angular distribution analysis was carried out for a total of 10 states in ^{96}Sr , up to and including a newly observed state at 3506(5) keV. Substantial population of states above this energy was observed as well, although it was not possible to identify individual states based on the measured γ rays. Fig. 8 shows the ^{96}Sr level scheme for states that were identified in this experiment.

The 0^+ states: The known 0, 1229 and 1465 keV 0^+ states were populated in the $d(^{95}\text{Sr}, p)$ experiment. The main results were already presented in ref. [35], here we just summarize the results for the 0^+ states. The ground state angular distribution was extracted by fitting the background of the excitation energy spectrum with a constrained exponential function ($\chi^2 \sim 1$) and taking the excess counts in the range $-0.5 < E_x < 0.5$ MeV. The 1229 keV 0_2^+ state angular distribution was produced by gating on the $0_2^+ \rightarrow 2_1^+$ 414 keV γ ray. Both angular distributions (Fig. 9) are in very good agreement with

the calculated $\Delta\ell = 0$ DWBA distributions. The spectroscopic factors for the 0 and 1229 keV 0^+ states were determined to be 0.19(3) and 0.22(3), respectively.

For the 1465 keV 0_3^+ state, it was not possible to extract an angular distribution by gating on the $0_3^+ \rightarrow 2_1^+$ 650 keV γ ray owing to its long half-life of 6.7(10) ns. The γ -ray detection efficiency of TIGRESS was simulated using GEANT4 [52] for both prompt and isomeric decays from a fast-moving ($\beta = 0.1$) ^{96}Sr ejectile. The simulations also take into account attenuation of the γ rays in the chamber and beam-line materials. The long half-life of the 1465 keV state results in a large decrease in γ -ray detection efficiency and poor Doppler reconstruction as it was not possible to determine the decay position of ^{96}Sr . The shape of the Doppler-reconstructed photo-peak was found to depend strongly on the position of the TIGRESS detectors, with clovers positioned at $\theta_{\text{lab}} > 120^\circ$ being the least affected. A γ -ray analysis was used to determine the relative population strengths of the two excited 0^+ states in ^{96}Sr by comparing counts in the 414 keV $0_2^+ \rightarrow 2_1^+$ and 650 keV $0_3^+ \rightarrow 2_1^+$ peaks under identical gate conditions. A 1 MeV excitation energy window was used so that both the 1229 and 1465 keV ^{96}Sr states could be fully included within the energy window, given the resolution of SHARC. This analysis was carried out using only the most downstream TIGRESS detectors positioned at $\theta_{\text{lab}} > 120^\circ$. The ratio of counts in the peaks (after correcting for the relative TIGRESS efficiency) was determined to be 0.22(4). This ratio was compared to the simulation results, which also take into account the indirect feeding of the 1229 keV state from the 1465 keV state via the $0_3^+ \rightarrow 0_2^+$ $E0$ transition and the branching ratio of the 650 keV transition. The experimentally measured relative population strengths are consistent with a scenario where the relative population of the 1465 to the 1229 keV state was 1.50(52). The spectroscopic factor for the 1465 keV state given in Table III is this relative population strength ratio multiplied by the 1229 keV state's spectroscopic factor as determined above. The DWBA calculations for both of these states predict the same integrated cross section within $\sim 3\%$, and so no excitation energy correction was applied.

The 815 keV state: It was not possible to extract an angular distribution for this state owing to the weak direct population, strong feeding from the 1229 keV state, and the E_x resolution. Instead, a γ -ray analysis was used to estimate the population strength. An energy gate of $0.4 < E_x < 1.2$ MeV in the upstream sections of SHARC was used so that all contributions from the 815 keV state were included. The indirect feeding from the 1229 keV state was subtracted based on the yield of the 414 keV transition, corrected for the TIGRESS efficiency. The 815 keV transition could not be resolved from the close-lying 813 keV transition originating from the 1628 keV state. The known branching ratio of the ground state decay allowed for the determination of the relative population of the 815 and 1628 keV states. The spectroscopic factor for the transfer to the 815 keV state listed in Ta-




FIG. 8. Level scheme of states in ^{96}Sr that were populated in the $d(^{95}\text{Sr}, p)$ reaction. The newly observed level at 3506 keV is indicated by a star.




FIG. 9. Angular distributions for $\Delta\ell = 0$ states in ^{96}Sr . The experimental data is presented alongside the fitted DWBA (blue) and ADWA (green) calculations, respectively.

ble III was then obtained using this ratio and the result for the 1628 keV state, see below, after correcting for the Q -value dependence of the calculated DWBA cross section for transfer to $1d_{3/2}$ neutron orbital.

The 1628 keV state: The 1628 keV state decays most strongly to the 2_1^+ state at 815 keV by the emission of a 813 keV γ ray. An angular distribution was thus extracted by double gating on both coincident 813 keV and 815 keV γ rays. The resulting angular distribution, shown in Fig. 10 (a), is in very good agreement with the $\Delta\ell = 2$ DWBA calculation. This, therefore, constrains the spin and parity to be 1^+ , 2^+ , or 3^+ . A suggested spin and parity of 2^+ was assigned to this state through β -decay studies of ^{96}Rb [28] using γ - γ angular correlations between the 813 keV and 815 keV transitions, although 1^+ could not be completely ruled out given the available statistics. Although weak, the branching ratios of this state to the $0_{1,2}^+$ states [28] make it highly unlikely that this state has spin and parity 3^+ . If this state were 1^+ , the decay to the $0_{1,2}^+$ states would be of pure $M1$ character. The single-particle Weisskopf estimates for the strength of these $M1$ transitions indicate that they would be similar in strength to the 813 keV transition, but they are measured to be only 12.2 and 5.3%, respectively. These observations favor a $J^\pi = 2^+$ assignment




FIG. 10. Angular distributions for $\Delta\ell = 2$ states in ^{96}Sr . The experimental data is presented alongside the fitted DWBA (blue) and ADWA (green) calculations, respectively.

for the 1628 keV state. The spectroscopic factor listed in Table III assumes transfer to the neutron $1d_{3/2}$ orbital, as the $1d_{5/2}$ orbital is considered to be fully occupied at $N = 56$.

The 1793 keV state: This state was weakly populated, with most of the observed γ -ray strength coming from indirect feeding from higher levels. Fig. 11 (a) shows the angular distribution for the 1793 keV state, which was produced by gating on the $4_1^+ \rightarrow 2_1^+$ 978 keV γ ray transition. The measured angular distribution, which was

565 best reproduced by a $\Delta\ell = 4$ DWBA calculation, is con-
566 sistent with the established spin of 4^+ [28].




FIG. 11. Angular distributions for $\Delta\ell = 4$ states in ^{96}Sr . The experimental data is presented alongside the fitted DWBA (blue) and ADWA (green) calculations, respectively. Potential contamination of the 2120 keV state angular distribution by the neighboring 2113 keV state has been neglected (see text).

567

568

569 *The 1995 keV state:* This state was strongly popu-
570 lated directly through the $d(^{95}\text{Sr}, p)$ transfer reaction,
571 with negligible indirect feeding. It can be clearly seen
572 in Fig. 7 as a strong 1180 keV γ ray in coincidence with
573 excitation energies in the range $1.6 < E_x < 2.4$ MeV.
574 The angular distribution, shown in Fig. 10 (b) was pro-
575 duced by gating on the 1180 keV γ ray. It shows clear
576 $\Delta\ell = 2$ character which constrains the spin and parity
577 to be 1^+ , 2^+ , or 3^+ . A spin and parity of 3^+ is un-
578 likely since decay to the ground and 0_2^+ states has been
579 observed. A $J^\pi = 1^+$ assignment was suggested based
580 on β -decay studies of ^{96}Rb [28] using γ - γ angular corre-
581 lations between the 1180 keV and 815 keV γ rays. For
582 completeness, Table III also lists the $1d_{3/2}$ spectroscopic
583 factor for the $J^\pi = 2^+$ assignment.

584 *The 2084 keV state:* This state was also strongly popu-
585 lated with negligible feeding from higher lying states.
586 The direct ground state decay can be clearly seen in Fig. 7
587 as a strong 2084 keV γ -ray line in coincidence with exci-
588 tation energies in the range $1.6 < E_x < 2.4$ MeV. The
589 angular distribution obtained by gating on this transition
590 (Fig. 10 (c)) shows clear a $\Delta\ell = 2$ character constraining
591 the spin and parity of this state to 1^+ , 2^+ or 3^+ . Using
592 similar arguments as for the 1995 keV level, the decay
593 branches to the $0_{1,2}^+$ states effectively rule out 3^+ . The
594 $\log ft$ value of the β -decay of the ^{96}Rb $2^{(-)}$ ground state
595 to the 2084 keV state suggests a first forbidden transition
596 which, together with the present result, constrains this
597 state to have spin and parity 1^+ or 2^+ .

598 *The 2120 keV state:* The main (91 %) decay branch
599 of this state is by a 1305 keV transition to the 2^+ state.
600 However, it cannot be resolved from the 1299 keV transi-
601 tion arising from the 2113 keV state given the TIGRESS
602 energy resolution after Doppler-correction. The 2113 keV
603 state also decays by 485 keV (branching ratio 22 %) and
604 607 keV (35%) γ rays which have been observed in the
605 excitation energy range $1.8 < E_x < 2.6$ MeV. This indi-
606 cates that the relative population strengths are 25(20)%

607 for the 2113 keV level and 75(20)% for the 2120 keV
608 state. The angular distribution gated on both the 1299
609 and 1305 keV γ -ray lines shown in Fig. 11 (b) is thus
610 dominated by the 2120 keV state. It is in best agree-
611 ment with $\Delta\ell = 4$ which is in accord with the tentative
612 assignment $J = 4$ from spontaneous fission studies of
613 ^{248}Cm [31]. The spectroscopic factor for transfer to the
614 $0g_{7/2}$ orbital given in Table III is an upper limit for the
615 2120 keV state ignoring the contribution of the 2113 keV
616 level to the angular distribution.

617 *The 2217 keV state:* The angular distribution shown
618 in Fig. 10 (d) was produced by gating on the 1402 keV
619 γ -ray transition depopulating this state and is well de-
620 scribed by a $\Delta\ell = 2$ calculation. Therefore $J^\pi = 2^+$ is
621 assigned to this state confirming the previous provisional
622 $J = 2$ assignment based on γ - γ angular correlation mea-
623 surements [28].

624 *The 2576 keV state:* The angular distribution for
625 this level (Fig. 10 (e)) was produced by gating on the
626 1761 keV γ -ray transition. It has previously been ob-
627 served only in β -decay of ^{96}Rb [3] and its strength sug-
628 gests a first-forbidden decay. This is in agreement with
629 the $\Delta\ell = 2$ angular distribution deduced here, which
630 constrains the spin and parity to be 1^+ , 2^+ or 3^+ . Spec-
631 troscopic factors assuming transfer to the $1d_{3/2}$ ($0g_{7/2}$)
632 neutron orbital for $J^\pi = 1^+, 2^+$ (3^+) are listed in Ta-
633 ble III.

634 *The 3506 keV state:* The 3506(6) keV transition is
635 newly observed in this work (inset of Fig. 7). The ex-
636 citation energy spectrum gated on this transition shows
637 that this is a direct ground state decay. The angular
638 distribution obtained by gating on this γ ray is shown in
639 Fig. 10 (f). The measured angular distribution is in good
640 agreement with the $\Delta\ell = 2$ DWBA calculation. No other
641 new or known transitions were observed when gating on
642 this excitation energy range, indicating that the branch-
643 ing ratio for the 3506 keV γ ray to the ground state is
644 100(10)%. This constrains the spin and parity to be 1^+
645 or 2^+ .

646 C. The $d(^{96}\text{Sr}, p)$ reaction

647 The γ rays and excitation energy of states in ^{97}Sr that
648 were populated via the $d(^{96}\text{Sr}, p)$ reaction are shown in
649 Fig. 12. The 167 and 355 keV γ rays in the energy range
650 $-0.5 < E_x < 1$ MeV indicate that both the known 167
651 and 522 keV excited states were populated in this ex-
652 periment. Fig. 13 shows the ^{97}Sr level scheme for states
653 that were identified in this work. No other excited states
654 could be unambiguously identified, owing to the limited
655 statistics. Given the small difference in energy between
656 the ground state and 167 keV first excited state, and
657 the E_x energy resolution, it was not possible to obtain
658 the cross sections and angular distributions based on the
659 excitation energy spectrum alone. The strength of the
660 ground state was thus derived by means of a constrained
661 three-peak fit for the 0, 167 and 522 keV states as dis-
662 cussed in the next section.




FIG. 12. Projected γ -ray spectrum for ^{97}Sr states populated via the $d(^{96}\text{Sr}, p)$ reaction. A cut on excitation energies below 1 MeV has been applied.




FIG. 13. Level scheme for ^{97}Sr states that were populated through $d(^{96}\text{Sr}, p)$.

664 cussed above for ^{95}Sr . Examples are shown in Fig. 14.

665

667 *The ground state:* The ground state was very weakly
 668 populated through the $d(^{96}\text{Sr}, p)$ reaction and the angular
 669 distribution shown in Fig. 15 (a) did not exhibit a
 670 clear shape as no data could be obtained for the smallest
 671 scattering angles ($\theta_{\text{cm}} < 20^\circ$). In this region the yield is
 672 expected to be very small and due to the small Q -value
 673 the background is high at low excitation energy. How-
 674 ever, the ground state is known to be $J^\pi = 1/2^+$ [2]
 675 and the angular distribution obtained is in accord with
 676 $\Delta\ell = 0$. The spectroscopic factor given in Table III has
 677 been extracted from the data shown in Fig. 15 (a) as well
 678 as a two-component fit of the summed angular distribu-
 679 tions of the ground and 167 keV states.

680 *The 167 keV state:* Two independent angular distri-
 681 butions were produced for the 167 keV state; one was
 682 extracted using the three peak fit (Fig. 15 (b)) and a
 683 second was derived by gating on the 167 keV γ ray and
 684 the excitation energy limiting the contribution from the
 685 522 keV state. The shape of both angular distributions
 686 are in good agreement with the $\Delta\ell = 2$ reaction model
 687 calculations, in agreement with the established spin and
 688 parity of $3/2^+$ [49]. The spectroscopic factors that were
 689 extracted for each of the methods are 0.25(7) and 0.24(8),
 690 respectively, assuming the addition of a neutron to the
 691 $1d_{3/2}$ orbital. The weighted average of the two spectro-
 692 scopic factors is given in Table III.

693 *The 522 keV state:* The small number of counts in the
 694 355 and 522 keV γ -ray peaks (shown in Fig. 12) did not
 695 allow for a γ -gated angular distribution for the 522 keV
 696 state, and so the spectroscopic factor for this state was
 697 determined by using the three-peak fit. The $\Delta\ell = 2$
 698 angular distribution shown in Fig. 15 (c) constrains the
 699




FIG. 14. Excitation energy spectrum extracted from the recoiling proton energies and angles at a center of mass angles $\theta_{\text{cm}} = 22, 26,$ and 30° . The continuous green line shows the constrained 3-peak fit of the 0, 167 and 522 keV states. The dashed line represents the continuous background.

700 J^π of this state to be $3/2^+$ or $5/2^+$, in agreement with the
 701 $M1$ multiplicities of the decay to the 167 keV state and
 702 also from the 687 keV $5/2^+$ state [49]. The population of
 703 this state by adding a neutron to the $1d_{3/2}$ orbital is most
 704 likely as the $1d_{5/2}$ orbital is expected to be fully occupied
 705 at $N = 59$ and the spectroscopic factor should be even
 706 lower than in ^{95}Sr . Consequently, $3/2^+$ is a more likely
 707 spin and parity for this state. For completeness, Table III
 708 includes the spectroscopic factors for both possibilities
 709 0.21(8) and 0.13(5) for $J^\pi = 3/2^+$ and $5/2^+$, respectively,
 710 using the DWBA calculations.

IV. DISCUSSION

713 The results obtained here can be used to gain insights
 714 into the underlying single-particle configurations of states
 715 in $^{95,96,97}\text{Sr}$. The results are compared in the following to
 716 shell model calculations to investigate the role of proton
 717 and neutron configurations in the low-lying states. While
 718 the present calculations are not well adapted to describe
 719 the deformed structures in ^{96}Sr and ^{97}Sr , the structure of
 720 ^{95}Sr before the shape transition should be well described,
 721 even in rather limited model spaces as will be discussed.

Nucleus	E_x [keV]	E_γ [keV]	J^π	$\Delta\ell$	C^2S (DWBA)	C^2S (ADWA)
^{95}Sr	0	fit	$\frac{1}{2}^+$	0	0.41(9)	0.34(7)
	352	fit, 352	$\frac{3}{2}^+$	2	0.53(8) [†]	0.45(7) [†]
	556	-	$\frac{7}{2}^+$	-	-	-
	681	fit, 329, 681	$\frac{5}{2}^+$	2	0.16(3) [†]	0.14(3) [†]
	1239	-	$\frac{3}{2}^+, \frac{5}{2}^+, \frac{7}{2}^+$	-	-	-
	1666	-	$\frac{3}{2}^+, \frac{5}{2}^+, \frac{7}{2}^+$	-	-	-
^{96}Sr	0	fit	0^+	0	0.19(3)	0.15(3)
	815	-	2^+	-	0.038(12)	0.034(12)
	1229	414	0^+	0	0.22(3)	0.19(3)
	1465	-	0^+	-	0.33(13)	0.29(12)
	1628	813 + 815	2^+	2	0.069(25)	0.056(23)
	1793	978	4^+	4	0.066(16)	0.058(17)
	1995	1180	$1^+, (2^+)$	2	0.20(3), (0.12(2))	0.18(3), (0.10(2))
	2084	2084	$1^+, 2^+$	2	0.24(5), 0.15(3)	0.21(4), 0.12(3)
	2120	1305	$4^+, (3^+)$	4	0.19(4), (0.21(4))	0.16(4), (0.21(4))
	2217	1402	2^+	2	0.047(8)	0.034(8)
	2576	1761	$1^+, 2^+, 3^+$	2	0.062(12), 0.037(7), 0.025(5)	0.049(9), 0.028(6), 0.019(5)
	3506(5)*	3506(5)	$1^+, 2^+$	2	0.047(9), 0.027(5)	0.034(8), 0.020(4)
^{97}Sr	0	fit	$\frac{1}{2}^+$	0	0.07(5)	0.06(5)
	167	fit, 167	$\frac{3}{2}^+$	2	0.11(10) [‡] 0.25(5) [†] 0.21(7) [‡]	0.07(7) [‡] 0.20(5) [†] 0.19(7) [‡]
	522	fit	$\frac{3}{2}^+, \frac{5}{2}^+$	2	0.21(8), 0.13(5)	0.17(7), 0.11(4)

[†] C^2S presented is the weighted average from multiple determinations

* new state

[‡]determined from the summed angular distribution of ground and 167 keV state

TABLE III. Results for $^{95,96,97}\text{Sr}$ states that were studied through the $d(^{94,95,96}\text{Sr}, p)$ reactions. Spectroscopic factors (C^2S) are given for all allowed J^π . J^π values in bold are new assignments or refined constraints. The method of angular distribution extraction, if any, for each state is presented under E_γ . Assignments and spectroscopic factors in parenthesis are alternative assignments that cannot be definitively ruled out by the present data, but are unlikely given previous experiments.

Shell model calculations for $^{94-97}\text{Sr}$ were carried out using NushellX [53] with the *glek* interaction [54] and several different model spaces. The single-particle energies of the interaction were adjusted so that the energies of low-lying states in the vicinity of $N \sim 56$ and $Z \sim 38$ were in good agreement with experiment [35]. In the present calculations the neutron $1d_{5/2}$, $2s_{1/2}$, $1d_{3/2}$ and $0g_{7/2}$ orbitals, outside an inert $N = 50$ core, were included. The higher-lying $0h_{11/2}$ orbital was not included as contributions from this orbital to low-lying positive parity states are expected to be small owing to the high single-particle energy [22].

Three different truncations of the proton valence space were investigated. In the smallest model space (a) the protons were frozen in a $(1p_{3/2})^4$ configuration so that the calculated states were built up using only the neutron configurations. Model space (b) included the $1p_{1/2}$

orbital and protons could be distributed across the $1p$ orbitals so that the effect of $(1p_{3/2})^{(4-x)}(1p_{1/2})^x$ configurations could be investigated. A third model space, (c), was used to investigate the effect of the proton $0g_{9/2}$ orbital on low-lying states. Up to two protons were allowed to occupy this orbital, so that configurations such as $(1p_{3/2})^2(0g_{9/2})^2$ and $(1p_{1/2})^2(0g_{9/2})^2$ were possible. This truncation was necessary due to the available computational resources. Proton seniority $\nu \neq 0$ configurations are expected to play a negligible role in the configurations of states that are strongly populated via the $d(\text{Sr}, p)$ reactions as single-step neutron transfer cannot break and re-couple proton pairs. Overall, additional proton degrees of freedom resulted in a lowering of the excitation energies, as correlations between complex configurations provide extra binding energy. This effect was evidenced by the increased mixing of the large number of configura-




FIG. 15. Fit of the reaction model calculations to the experimental data for the 167 and 522 keV states in ^{97}Sr . The solid lines are the best-fitting reaction model calculations using the DWBA (blue) and ADWA (green) methods. The fitting was restricted to the forward angles ($\theta_{\text{cm}} < 40^\circ$). For the 167 keV state the angular distribution extracted by gating on the 167 keV γ -ray transition and the excitation energy is also shown.

tions in the wave functions. The increased proton model space also impacted the predicted spectroscopic factors, as the mixed wave functions, unsurprisingly, tend to have smaller overlaps.

A. ^{95}Sr

In a shell model picture, low-lying states in ^{95}Sr can be approximated as simple excitations of the unpaired neutron into the different valence orbitals, which define the spins and parities of the low-lying states. The ground state spectroscopic factor (Table IV) is in good agreement with that calculated in the shell model for all three model spaces, although the substantial improvement in (b) indicates that proton pair excitations into the $1p_{1/2}$ orbital play an important role in the ground states of both ^{94}Sr and ^{95}Sr . The same is also true for the energy and spectroscopic factor of the $3/2^+$ first excited state: the calculated energy of this level drops substantially with the inclusion of the proton $1p_{1/2}$ orbital. As can be seen, a gradual reduction in spectroscopic strength is predicted for the ground state and 352 keV excited states as the proton degrees of freedom are increased. In each case, there were no other $1/2^+$ or $3/2^+$ states with substantial spectroscopic strength ($C^2S > 0.04$) predicted. On the other hand, each calculation predicted a low-energy $5/2^+$ state with $C^2S > 0.15$ at around ~ 600 keV (Table IV) which is dominated by a neutron $(1d_{5/2})^5(2s_{1/2})^2$ configuration in all of the calculations. The population of such a state in the one-neutron transfer suggests that

the $\nu 1d_{5/2}$ orbit is not fully occupied in the ground state of ^{94}Sr . The larger model spaces, which increase the neutron particle-hole configurations in the ^{94}Sr ground state, show an increase in the spectroscopic factor for the $5/2^+$ state. This also affirms the assignment of $5/2^+$ to the state seen at 681 keV. The spectroscopic factor and the excitation energy of the $7/2^+$ state strongly depends on the proton configurations. This demonstrates the effect of the Federman-Pittel mechanism [9, 10] whereby the mutual interaction of the $\pi 0g_{9/2}$ and $\nu 0g_{7/2}$ orbitals drives the deformation in this region. While the spectroscopic factor for this state could not be deduced, the observed yield (Fig. 5) suggests that this state has a small spectroscopic factor, at variance with the shell model calculations.

Figure 16 shows the experimental level energies and DWBA spectroscopic factors for ^{95}Sr states that were populated via the $d(^{94}\text{Sr}, p)$ reaction compared to the shell model calculations. Overall, the shell model cal-




FIG. 16. Comparison of experimental (exp) spectroscopic factors (C^2S) to those from shell model calculations carried out in model spaces (a), (b) and (c) – see text. States are labeled by the neutron single-particle orbital populated in the transfer reaction.

culations for proton model space (b) describe these low-lying states very well aside from the $7/2^+$ state. This suggests that the ground states of both ^{94}Sr and ^{95}Sr have similar and nearly spherical shapes and in agreement with $B(E2)$ [5, 30] and charge radii [2] measure-

Nucleus	J^π	exp.		SM (a)		SM (b)		SM (c)	
		E (keV)	C^2S	E (keV)	C^2S	E (keV)	C^2S	E (keV)	C^2S
^{95}Sr	$\frac{1}{2}^+$	0	0.41(9)	0	0.553	0	0.449	0	0.413
	$\frac{3}{2}^+$	352	0.53(8)	766	0.865	412	0.767	375	0.744
	$\frac{5}{2}^+$	681	0.16(3)	691	0.146	585	0.180	523	0.201
	$\frac{7}{2}^+$	556		1086	0.959	602	0.828	205	0.757
^{97}Sr	$\frac{1}{2}^+$	0	0.10(5)	1631	0.013	1279	0.024	417	0.002
	$\frac{3}{2}^+$	167	0.25(5)	0	0.881	0	0.804	117	0.713
	$\frac{5}{2}^+$	308		270	0.979	149	0.931	0	0.819
	$\frac{7}{2}^+$			1714	0.025	1336	0.042	57	0.000
	$\frac{9}{2}^+$	522	0.13(5)						

TABLE IV. Comparison of $d(^{94,96}\text{Sr}, p)$ spectroscopic factors to shell model calculations for low-lying states. The labels SM (a), (b) and (c) denote the three proton model spaces that were investigated (see text).

811 ments. It should be noted that a recent Monte-Carlo
 812 shell model calculation [27] predicts the onset of deforma-
 813 tion in the Sr nuclei too early. This is evident from the
 814 calculated spectra of the even-even Sr nuclei [34] as well
 815 as the level scheme of ^{95}Sr with 13 states below 1 MeV,
 816 some of them strongly deformed [55].

B. ^{96}Sr

817
 818 Table V compares the shell model results within
 819 each proton model space for the lowest states. In the
 821 $d(^{95}\text{Sr}, p)^{96}\text{Sr}$ reaction each state with $J > 0$ can be popu-
 822 lated by more than one value for the angular momentum
 823 transfer. The coupling of the $1/2^+$ ground state of ^{95}Sr
 824 to a valence neutron in $1d_{5/2}$ ($J = 2, 3$), $2s_{1/2}$ ($J = 0, 1$),
 825 $1d_{3/2}$ ($J = 1, 2$), and $0g_{7/2}$ ($J = 3, 4$) leads to various
 826 final states. The shell model calculations suggest that
 827 $1d_{5/2}$ dominates the $J = 2, 3$ states and the contribu-
 828 tion of $2s_{1/2}$ to the 1^+ states is negligible. Indeed the
 829 experimental angular distributions for the 1^+ candidates
 830 are well accounted for by $\Delta\ell = 2$ transfer as shown in
 831 Fig. 10. The results of the calculations are compared to
 832 the experimental data in Fig. 17.

834 According to the calculations, the ground state
 835 of ^{96}Sr is dominated ($> 60\%$) by a neutron
 836 $(1d_{5/2})^6(2s_{1/2})^2$ configuration with substantial ($\sim 15\%$)
 837 $(1d_{5/2})^4(2s_{1/2})^2(1d_{3/2})^2$ contributions in all of the model
 838 spaces. The transfer from the $1/2^+$ ground state of ^{95}Sr
 839 has, therefore, a large spectroscopic factor approaching
 840 that of the independent particle model ($C^2S = 2$). The
 841 result depends only weakly on the proton model space,
 842 reflecting the result obtained for ^{95}Sr where the spectro-
 843 scopic factor of the $1/2^+$ ground state (and the $3/2^+$
 844 first excited state) only weakly depend on the avail-
 845 able proton space. The predicted spectroscopic factor
 846 ($C^2S_{\text{SM}} \sim 1.5$) was found to be much larger than the
 847 experimental result ($C^2S_{\text{exp}} = 0.19(3)$), suggesting that
 848 the ground state of ^{96}Sr can not be well-described within
 849 the context of the spherical shell model. Assuming axial




FIG. 17. Comparison of experimental (exp) spectroscopic factors (C^2S) for $d(^{94}\text{Sr}, p)^{95}\text{Sr}$ to shell model calculations that were carried out in model spaces (a), (b) and (c) – see text. States are labeled by their spin and parity as well as the orbital populated in the transfer reaction. Open symbols label the 1^+ states populated by transfer to the $2s_{1/2}$ orbital, as well as transfer to the $1d_{5/2}$ orbital for $J^\pi = 2, 3^+$. Only states with $C^2S > 0.01$ are shown. For experiment $J^\pi = 2^+$ has been assumed for the 2084, 2576, and 3506 keV.

850 symmetry a Coulomb excitation experiment determining
 851 the quadrupole moment of the 2_1^+ state suggests a weakly

SM (a)			SM (b)			SM (c)		
J^π	E (keV)	C^2S	J^π	E (keV)	C^2S	J^π	E (keV)	C^2S
0_1^+	0	1.742	0_1^+	0	1.575	0_1^+	0	1.454
0_2^+	2271	0.056	0_2^+	1691	0.098	0_2^+	444	0.105
0_3^+	3066	0.001	0_3^+	2034	0.006	0_3^+	1483	0.052
1_1^+	2116	0.823	1_1^+	1961	0.725	1_1^+	2048	0.671
2_1^+	1959	0.829	2_1^+	1662	0.402	2_1^+	705	0.002
2_2^+	2307	0.001	2_2^+	1905	0.246	2_2^+	1442	0.061
2_3^+	2706	0.064	2_3^+	2155	0.035	2_3^+	1804	0.013
2_4^+	2884	0.014	2_4^+	2160	0.061	2_4^+	1883	0.378
3_1^+	2345	0.828	3_1^+	2078	0.699	3_1^+	1885	0.517
4_1^+	2250	0.134	4_1^+	2011	0.038	4_1^+	1326	0.002
4_2^+	2278	0.811	4_2^+	2120	0.720	4_2^+	1818	0.541

TABLE V. Comparison of $d(^{95}\text{Sr}, p)^{96}\text{Sr}$ spectroscopic factors and excitation energies from the shell model calculations. The labels SM (a), (b) and (c) denote the three proton model spaces that were investigated (see text).

852 deformed ($\beta \sim 0.1$) ground state [4, 33].

853 On the other hand, the experimental spectroscopic fac-
854 tors for the excited 0^+ states are substantially larger than
855 for the ground state. The 1229 and 1465 keV states
856 in ^{96}Sr are known to arise from the mixing between a
857 strongly deformed and a nearly spherical configuration,
858 as evidenced by the large $\rho^2(E0)$ transition strength be-
859 tween them [28]. The strongly deformed states should
860 not be populated directly in one-neutron transfer onto
861 the spherical ^{95}Sr ground state. Therefore, the spec-
862 troscopic factors of these states reflects their underlying
863 spherical component which is populated strongly by the
864 (d, p) reaction. This suggests the existence of three differ-
865 ent shapes in ^{96}Sr , with a weakly deformed, likely oblate,
866 ground state and strongly mixed spherical and well de-
867 formed (prolate with $\beta = 0.31(3)$) configurations in the
868 excited 0^+ states. This is discussed in more detail in
869 Ref. [35].

870 Given that the ground state of ^{96}Sr was not well
871 reproduced in any of the calculations, it is expected
872 that there will also be substantial discrepancies with
873 the low energy states of ^{96}Sr . The wave function for
874 the 2_1^+ state was predicted to be dominated by the
875 neutron $(1d_{5/2})^6(2s_{1/2})^1(1d_{3/2})^1$ configurations in shell
876 model calculation (a) (73%) and (b) (27%), which has
877 a large overlap with the ^{95}Sr ground state. Within the
878 model space of calculation (c), many additional contri-
879 butions were present in the lowest energy 2^+ state and
880 the spectroscopic factor (Table V) is very small. The
881 drop in energy of the 2^+ state to 705 keV in model (c)
882 reflects the lowering of the $7/2^+$ state in ^{95}Sr as excita-
883 tions to the proton $0g_{9/2}$ orbital become possible. The
884 large spectroscopic factor predicted for the 2_4^+ state re-
885 flects its wave function composition, which in this case
886 is similar to the 2_1^+ state of the other calculations. The
887 experimental 2084 keV state might be associated with

888 this level. In agreement with the experimental results,
889 the calculations in model space (c) predict small spec-
890 troscopic factors for the other 2^+ states. The first 2^+
891 state in $^{90-96}\text{Sr}$ was previously interpreted as a proton
892 spin-flip excitation from the $1p_{3/2}$ to the $1p_{1/2}$ orbital as
893 no indications of the neutron sub-shell closure are visible
894 at $N = 56$. The constant excitation energy can then ex-
895 plained by the quenching of the proton $1p_{3/2} - 1p_{1/2}$ spin-
896 orbit splitting as the neutron $1d_{5/2}$ orbital is filled [56].
897 Such configurations would not be populated here using
898 the (d, p) reaction. The small experimental spectroscopic
899 factor for the 2^+ state is consistent with a proton exci-
900 tation or with a non-spherical configuration that has a
901 small overlap with the ^{95}Sr ground state.

902 The main contributions to the wave func-
903 tion of the low-lying 4^+ states are the neutron
904 $(1d_{5/2})^5(2s_{1/2})^2(1d_{3/2})^1$ and $(1d_{5/2})^6(2s_{1/2})^1(0g_{7/2})^1$
905 configurations. The latter configuration can be popu-
906 lated directly via one-neutron transfer ($\Delta\ell = 4$), which
907 results in an enhancement of the spectroscopic factor
908 as seen in Table V. There is no strong evidence to
909 suggest that the structure of the 1793 keV 4_1^+ ^{96}Sr state
910 is well-described within any of the present calculations.
911 The 4^+ state at 2120 keV has a larger spectroscopic
912 factor, and may be associated with the calculated 4_1^+
913 state. Additionally, $\Delta\ell = 4$ strength has been observed
914 around $E = 3200$ keV, but could not be assigned
915 to a particular state [42]. A low-lying 3^+ state was
916 also predicted in each of the model spaces. The same
917 $(1d_{5/2})^6(2s_{1/2})^1(0g_{7/2})^1$ configuration was found to be
918 the primary component of this state, contributing 67%,
919 47% and 33% to the total wave function in model spaces
920 (a), (b), and (c), respectively. Experimentally, there
921 is no candidate for a 3^+ state with large spectroscopic
922 factor, although the 4^+ assignment of the 2120 keV state
923 is tentative, and could be a 3^+ state. Another state of

interest is the first 1^+ state, which appears at around 2 MeV in all of the calculations. This state originates from the neutron $(1d_{5/2})^6(2s_{1/2})^1(1d_{3/2})^1$ configuration which can be populated directly via $\Delta\ell = 2$ transfer. The calculations predict that this configuration makes up 78%, 68% and 61% of the total wave function in model spaces (a), (b), and (c), respectively. The 1^+ state at 1995 keV is a likely candidate for this configuration, as it was strongly populated in the $d(^{95}\text{Sr}, p)\text{Sr}$ reaction.

To summarize, the spectroscopic strength in ^{96}Sr is smaller and more fragmented than in the shell model calculations, in particular for the 0^+ and 2^+ states. The absolute spectroscopic factors are not reproduced, but the rather large spectroscopic factors for low-lying 1^+ and 4^+ states are overall in line with the calculations. The discrepancy for the 0^+ states, with the observation of the majority of the spectroscopic strength in the excited 0^+ states, suggests that the ground state of ^{96}Sr is not spherical, but rather weakly (oblate) deformed [35].

C. ^{97}Sr

The comparison of the experimental results with the shell model calculations in Table IV suggest that the structure of ^{97}Sr is more complicated than for ^{95}Sr . The ground state spin and parity $1/2^+$ [50] is unexpected in the framework of the spherical shell model, where the $2s_{1/2}$ orbital should be fully occupied at $N = 59$. Isotope shift measurements across the Sr chain indicate that the ground state of ^{97}Sr is either spherical or weakly deformed [2]. The magnetic moment of the ^{97}Sr ground state is close to the value of ^{95}Sr and much smaller than the Schmidt value. The close-lying $0g_{7/2}$ and $1d_{3/2}$ $K^\pi = 1/2^+$ orbitals could lead to substantial mixing even for weakly deformed states, and thus explain these results.

In addition to the excitation energies, the calculated spectroscopic factors for the $d(^{96}\text{Sr}, p)$ reaction are listed in Table IV. As discussed previously, the striking discrepancies between the calculated spectroscopic factors for the $d(^{95}\text{Sr}, p)$ reaction and our experimental results indicate that the shell model will not adequately describe the $d(^{96}\text{Sr}, p)$ reaction. A good description of the ^{96}Sr ground state wave function is essential for calculating the overlap with states in ^{97}Sr and the results from the $d(^{95}\text{Sr}, p)$ reaction make it clear that ^{94}Sr and ^{95}Sr ground states are well described by the shell model but the ^{96}Sr ground state is not. The interpretation of the spectroscopic factors is thus limited here to qualitative remarks.

From the weak population of the ^{97}Sr ground state in the $d(^{96}\text{Sr}, p)$ reaction we can conclude that it has a considerably different wave function than that of the weakly deformed ^{96}Sr ground state, although this does not necessarily imply that it is strongly deformed. Clearly, further experimental measurements must be made to elucidate the structure of this state. The largest spectroscopic factor is found here for the $3/2^+$ state, similar to ^{95}Sr , yet

this state does not necessarily have the same structure as the configuration of the even-even projectile affects the spectroscopic factor as well. Relatively strong population of a low-lying $5/2^+$ state via the $d(^{96}\text{Sr}, p)$ reaction indicates that there are substantial vacancies in the neutron $1d_{5/2}$ orbital in the ^{96}Sr ground state and this level could be regarded as the $N = 59$ analogue of the 681 keV $5/2^+$ ^{95}Sr state.

V. SUMMARY AND OUTLOOK

In summary, states in $^{95,96,97}\text{Sr}$ have been studied via the $d(^{94,95,96}\text{Sr}, p)$ reactions for the first time. In total, 16 angular distribution measurements and associated spectroscopic factors have been determined. Spectroscopic factors were deduced for an additional 2 states by using a relative γ -ray analysis. These spectroscopic factors were compared to shell model calculations using realistic effective interactions within several carefully chosen valence spaces.

In ^{95}Sr , firm spin and parity assignments of $3/2^+$, $7/2^+$ and $5/2^+$ have been made for the 352, 556 and 681 keV states, respectively. Further constraints on the spin and parity of the 1666 keV state have been made, based on predicted cross sections. Good agreement was observed between experiment and shell model calculations, which suggests that low-lying states in ^{95}Sr arise from relatively simple neutron configurations.

In ^{96}Sr , all angular distribution analyses that were carried out confirm and refine previous spin and parity assignments, and new spin and parity constraints of 1^+ , 2^+ , 3^+ have been made for the 2576 state. A state at 3506(5) keV has been newly identified, which is a candidate for a 1^+ or 2^+ level. It was found that the excited 0^+ states possess a larger overlap with the ground state of ^{95}Sr than the 0_1^+ state, as evidenced by the larger spectroscopic factors. This result is in contrast to the shell model calculations, which predict that almost all of the $\Delta\ell = 0$ strength is concentrated in the 0_1^+ state. A weakly deformed structure is suggested for the ^{96}Sr ground state. The results presented here also agree with the proposed proton configuration of the 2_1^+ state [56] which is not strongly populated in the present experiment.

In ^{97}Sr , substantial spectroscopic strength to the 167 and 522 keV states was observed while the ground state was very weakly populated. The angular distributions are in agreement with the established spins and parities of the 167 and 522 keV states, however no quantitative comparison with the shell model could be made as the ^{96}Sr ground state was not well-described within the calculations.

The results discussed here provide valuable information concerning the single-particle composition of states in $^{95,96,97}\text{Sr}$. By comparing the experimental spectroscopic factors to shell model calculations, we are able to gain an improved understanding of structural changes

that indicate a departure from simple shell structure for $N \geq 58$. In future, two-neutron transfer reactions should provide for a complementary examination of the underlying structure of the 0^+ states in the even-even neutron-rich Sr isotopes. Low-energy Coulomb excitation to characterize the deformation of excited states in the even-odd Sr nuclei could provide information complementary to the present work. Lastly, large-scale shell model calculations in larger valence spaces, which have been so far only applied to the neutron-rich Zr isotopes [22, 27], will provide an important addition to the present discussion.

ACKNOWLEDGMENTS

The efforts of the TRIUMF operations team in supplying the Sr beams are highly appreciated. We ac-

knowledge support from the Science and Technologies Facility Council (UK, grants EP/D060575/1 and ST/L005727/1), the National Science Foundation (US, grant PHY-1306297), the Natural Sciences and Engineering Research Council of Canada, the Canada Foundation for Innovation and the British Columbia Knowledge and Development Fund. TRIUMF receives funding via a contribution through the National Research Council Canada. N.A.O. acknowledges support from the CNRS PICS “PACIFIC”.

-
- [1] R. F. Casten, *Nuclear Structure from a Simple Perspective*, 2nd ed., edited by P. E. Hodgson (Oxford Science Publications, 2000).
- [2] F. Buchinger, E. B. Ramsay, E. Arnold, W. Neu, R. Neugart, *et al.*, *Phys. Rev. C* **41**, 2883 (1990).
- [3] “Evaluated nuclear structure data file,”.
- [4] E. Clément, M. Zielinska, A. Gørgen, W. Korten, S. Péru, *et al.*, *Phys. Rev. Lett.* **116**, 022701 (2016).
- [5] A. Chester, G. Ball, R. Caballero-Folch, *et al.*, *Phys. Rev. C* **96**, 011302(R) (2017).
- [6] G. Jung, “Nuclear Spectroscopy on Neutron Rich Rubidium With Even Mass Numbers,” PhD thesis, Justus Liebig-Universität, Giessen (1980).
- [7] J. Wood, E. Zganjar, C. D. Coster, and K. Heyde, *Nucl. Phys. A* **651**, 323 (1999).
- [8] D. Arseniev, A. Sobczewski, and V. Soloviev, *Nucl. Phys. A* **139**, 269 (1969).
- [9] P. Federman and S. Pittel, *Phys. Lett. B* **69**, 385 (1977).
- [10] P. Federman and S. Pittel, *Phys. Rev. C* **20**, 820 (1979).
- [11] A. Kumar and M. R. Gunye, *Phys. Rev. C* **32**, 2116 (1985).
- [12] S. Michiaki and A. Akito, *Nucl. Phys. A* **515**, 77 (1990).
- [13] J. Skalski, P.-H. Heenen, and P. Bonche, *Nucl. Phys. A* **559**, 221 (1993).
- [14] A. Baran and W. Höhenberger, *Phys. Rev. C* **52**, 2242 (1995).
- [15] G. Lalazissis and M. Sharma, *Nucl. Phys. A* **586**, 201 (1995).
- [16] J. Skalski, S. Mizutori, and W. Nazarewicz, *Nucl. Phys. A* **617**, 282 (1997).
- [17] A. Holt, T. Engeland, M. Hjorth-Jensen, and E. Osnes, *Phys. Rev. C* **61**, 064318 (2000).
- [18] H. Zhang, S. Im, J. Li, W. Zuo, Z. Ma, B. Chen, and W. Scheid, *Eur. Phys. Jour. A* **30**, 519 (2006).
- [19] T. Rzaca-Urban, K. Sieja, W. Urban, F. Nowacki, J. L. Durell, A. G. Smith, and I. Ahmad, *Phys. Rev. C* **79**, 024319 (2009).
- [20] R. Rodriguez-Guzman, P. Sarriguren, and L. M. Robledo, *Phys. Rev. C* **82**, 044318 (2010).
- [21] R. Rodriguez-Guzman, P. Sarriguren, L. Robledo, and S. Perez-Martin, *Phys. Lett. B* **691**, 202 (2010).
- [22] K. Sieja, F. Nowacki, K. Langanke, and G. Martínez-Pinedo, *Phys. Rev. C* **79**, 064310 (2009).
- [23] Y.-X. Liu, Y. Sun, X.-H. Zhou, Y.-H. Zhang, S.-Y. Yu, Y.-C. Yang, and H. Jin, *Nucl. Phys. A* **858**, 11 (2011).
- [24] H. Mei, J. Xiang, J. M. Yao, Z. P. Li, and J. Meng, *Phys. Rev. C* **85**, 034321 (2012).
- [25] J. Xiang, Z. Li, Z. Li, J. Yao, and J. Meng, *Nucl. Phys. A* **873**, 1 (2012).
- [26] A. Petrovici, *Phys. Rev. C* **85**, 034337 (2012).
- [27] T. Togashi, Y. Tsunoda, T. Otsuka, and N. Shimizu, *Phys. Rev. Lett.* **117**, 172502 (2016).
- [28] G. Jung, B. Pfeiffer, L. J. Alquist, H. Wollnik, P. Hungerford, *et al.*, *Phys. Rev. C* **22**, 252 (1980).
- [29] K. L. Kratz, A. Schröder, H. Ohm, M. Zendel, H. Gabelmann, *et al.*, *Zeitschrift für Physik A* **306**, 239 (1982).
- [30] H. Mach, F. Wahn, G. Molnar, K. Sistemich, J. C. Hill, *et al.*, *Nucl. Phys. A* **523**, 197 (1991).
- [31] C. Y. Wu, H. Hua, D. Cline, A. B. Hayes, R. Teng, *et al.*, *Phys. Rev. C* **70**, 064312 (2004).
- [32] J. Park, A. B. Garnsworthy, R. Krücken, C. Andreoiu, *et al.*, *Phys. Rev. C* **93**, 014315 (2016).
- [33] E. Clément, M. Zielinska, S. Péru, H. Goutte, S. Hilaire, *et al.*, *Phys. Rev. C* **94**, 054326 (2016).
- [34] J.-M. Régis, J. Jolie, N. Saed-Samii, N. Warr, M. Pfeiffer, *et al.*, *Phys. Rev. C* **95**, 054319 (2017).
- [35] S. Cruz, P. C. Bender, R. Krücken, K. Wimmer, *et al.*, *Phys. Lett. B* **786**, 94 (2018).
- [36] G. C. Ball, L. Buchmann, B. Davids, R. Kanungo, C. Ruiz, and C. E. Svensson, *J. Phys. G* **38**, 024003 (2011).
- [37] C. A. Diget *et al.*, *Jour. of Instr.* **6**, P02005 (2011).
- [38] G. Hackman and C. E. Svensson, *Hyp. Int.* **225**, 241 (2014).
- [39] A. Matta *et al.*, *Phys. Rev. C* **99**, 044320 (2019).
- [40] C. Nobs, “Simulating and testing the TRIUMF Bragg ionisation chamber,” Masters thesis, University of Surrey, Guildford (2013).
- [41] J. F. Ziegler, M. D. Ziegler, and J. P. Biersack, *Nucl. Instrum. Meth. B* **268** (2010).
- [42] S. Cruz, “Single particle structure of exotic strontium isotopes,” PhD thesis, Univeristy of British Columbia

- (2017).
- [43] W. N. Catford *et al.*, Phys. Rev. Lett. **104**, 192501 (2010).
- [44] I. Thompson, Comp. Phys. Rep. **7**, 167 (1988).
- [45] C. Perey and F. Perey, At. Data Nucl. Data Tab. **17**, 1 (1976).
- [46] W. W. Daehnick, J. D. Childs, and Z. Vrcelj, Phys. Rev. C **21**, 2253 (1980).
- [47] J. Lohr and W. Haerberli, Nucl. Phys. A **232**, 381 (1974).
- [48] A. Koning and J. Delaroche, Nucl. Phys. A **713**, 231 (2003).
- [49] K. L. Kratz, H. Ohm, A. Schröder, H. Gabelmann, W. Ziegert, *et al.*, Zeit. f. Phys. A **312**, 43 (1983).
- [50] F. Buchinger, E. B. Ramsay, R. E. Silverans, P. Lievens, E. Arnold, W. Neu, R. Neugart, K. Wendt, G. Ulm, and the ISOLDE Collaboration, Zeit. f. Phys. A **327**, 361 (1987).
- [51] J. K. Hwang, A. V. Ramayya, J. H. Hamilton, *et al.*, Phys. Rev. C **69**, 067302 (2004).
- [52] S. Agostinelli *et al.*, Nucl. Instr. Meth. A **506**, 250 (2003).
- [53] B. A. Brown, “Computer code NUSHELLX.”
- [54] H. Mach, E. K. Warburton, R. L. Gill, R. F. Casten, J. A. Becker, B. A. Brown, and J. A. Winger, Phys. Rev. C **41**, 226 (1990).
- [55] T. Togashi and T. Otsuka, priv. comm. (2018).
- [56] P. Federman, S. Pittel, and A. Etchegoyen, Phys. Lett. **140B**, 269 (1984).