

This is a repository copy of *Use of magnetic resonance angiography to select candidates with recently symptomatic carotid stenosis for surgery: systematic review.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/155/>

Article:

Kelly, S., Bamford, J.M., Airey, C.M. et al. (7 more authors) (2002) Use of magnetic resonance angiography to select candidates with recently symptomatic carotid stenosis for surgery: systematic review. BMJ. pp. 198-201. ISSN: 1756-1833

<https://doi.org/10.1136/bmj.324.7331.198>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Use of magnetic resonance angiography to select candidates with recently symptomatic carotid stenosis for surgery: systematic review

Marie E Westwood, Steven Kelly, Elizabeth Berry, John M Bamford, Michael J Gough, C Mark Airey, James F M Meaney, Linda M Davies, Jane Cullingworth and Michael A Smith

BMJ 2002;324:198-
doi:10.1136/bmj.324.7331.198

Updated information and services can be found at:
<http://bmj.com/cgi/content/full/324/7331/198>

These include:

References

This article cites 41 articles, 18 of which can be accessed free at:
<http://bmj.com/cgi/content/full/324/7331/198#BIBL>

8 online articles that cite this article can be accessed at:
<http://bmj.com/cgi/content/full/324/7331/198#otherarticles>

Rapid responses

One rapid response has been posted to this article, which you can access for free at:
<http://bmj.com/cgi/content/full/324/7331/198#responses>

You can respond to this article at:
<http://bmj.com/cgi/eletter-submit/324/7331/198>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Topic collections

Articles on similar topics can be found in the following collections

- [Diagnostics tests](#) (289 articles)
 - [Other Neurology](#) (2972 articles)
 - [Radiological diagnosis](#) (377 articles)
 - [Ischaemic heart disease](#) (1491 articles)
 - [Stroke](#) (467 articles)
 - [Vascular Surgery](#) (175 articles)
-

Notes

To order reprints of this article go to:
<http://bmj.bmjournals.com/cgi/reprintform>

To subscribe to *BMJ* go to:
<http://www.bmjournals.com/subscriptions>

Use of magnetic resonance angiography to select candidates with recently symptomatic carotid stenosis for surgery: systematic review

Marie E Westwood, Steven Kelly, Elizabeth Berry, John M Bamford, Michael J Gough, C Mark Airey, James F M Meaney, Linda M Davies, Jane Cullingworth, Michael A Smith

Abstract

Objective To determine if sufficient evidence exists to support the use of magnetic resonance angiography as a means of selecting patients with recently symptomatic high grade carotid stenosis for surgery.

Design Systematic review of published research on the diagnostic performance of magnetic resonance angiography, 1990-9.

Main outcome measures Performance characteristics of diagnostic test.

Results 126 potentially relevant articles were identified, but many articles failed to examine the performance of magnetic resonance angiography as a diagnostic test at the surgical decision thresholds used in major clinical trials on endarterectomy. 26 articles were included in a meta-analysis that showed a maximal joint sensitivity and specificity of 99% (95% confidence interval 98% to 100%) for identifying 70-99% stenosis and 90% (81% to 99%) for identifying 50-99% stenosis. Only four articles evaluated contrast enhanced magnetic resonance angiography.

Conclusions Magnetic resonance angiography is accurate for selecting patients for carotid endarterectomy at the surgical decision thresholds established in the major endarterectomy trials, but the evidence is not very robust because of the heterogeneity of the studies included. Research is needed to determine the diagnostic performance of the most recent developments in magnetic resonance angiography, including contrast enhanced techniques, as well as to assess the impact of magnetic resonance angiography on surgical decision making and outcomes.

Introduction

Patients with arteriosclerosis, particularly those with recent carotid territory stroke or transient ischaemic attack who might benefit from carotid endarterectomy, may be investigated with conventional angiography, ultrasonography, or magnetic resonance angiography.^{1 2} Magnetic resonance angiography was introduced in the 1980s. The technology continues to evolve rapidly, and three main approaches are

currently used: two dimensional time of flight methods, three dimensional time of flight methods, and three dimensional contrast enhanced techniques. Time of flight (or "inflow") methods are non-invasive and emphasise flowing blood because a different signal is given by material that has moved into the imaging volume. Contrast enhanced methods use gadolinium based contrast materials that are better tolerated than the iodinated media used for conventional angiography. Contrast enhancement overcomes some of the drawbacks of time of flight techniques, especially signal loss near stenoses. A greater volume of the body can be imaged in a shorter time, allowing evaluation of the vasculature from the aortic arch to the circle of Willis.

The use of magnetic resonance angiography and ultrasonography has expanded rapidly in recent years, and the use of conventional angiography has declined. Although conventional catheter angiography remains the definitive imaging technique, it is an invasive procedure for which the patient must be admitted to hospital, it involves the use of ionising radiation, and when used in the carotid circulation it is accompanied by serious complications, including a 0.5-2% risk of stroke.³ Magnetic resonance angiography may be more acceptable to patients and may be of particular use in patients not suitable for conventional angiography—for example, patients with an allergy to iodinated contrast medium, frail and elderly patients, and patients with severe peripheral vascular disease. These potential benefits may be offset by poor performance as a diagnostic test.

This systematic review examines the evidence on the performance of magnetic resonance angiography in evaluating patients with recently symptomatic internal carotid artery stenosis. The North American symptomatic carotid endarterectomy trial (NASCET) and the European carotid surgery trial (ECST) found a clear benefit of surgery in patients with recently symptomatic stenoses of 70-99% as measured by conventional angiography with the NASCET criteria.^{4 5} We therefore evaluated the evidence on the diagnostic performance of magnetic resonance angiography in comparison with conventional angiography at this threshold. NASCET and ECST found a smaller benefit of surgery in patients with symptomatic 50-99% steno-

Academic Unit of Medical Physics and Centre of Medical Imaging Research, University of Leeds, Leeds General Infirmary, Leeds LS1 3EX

Marie E Westwood
research fellow

Steven Kelly
research fellow

Elizabeth Berry
senior lecturer in medical physics

Michael A Smith
professor of medical physics

Vascular Surgery, Leeds Teaching Hospitals NHS Trust, Leeds General Infirmary
Michael J Gough
consultant vascular surgeon

Department of Radiology, Leeds Teaching Hospitals NHS Trust, Leeds General Infirmary
James F M Meaney
consultant radiologist
Jane Cullingworth
superintendent radiographer

Department of Neurology, Leeds Teaching Hospitals NHS Trust, St James's Hospital, Leeds LS9 7TF

John M Bamford
consultant neurologist

Division of Public Health, Nuffield Institute for Health, University of Leeds, Leeds LS2 9PL

C Mark Airey
senior research fellow in public health

continued over

BMJ 2002;324:1-5

Centre for Health Economics,
University of York,
York YO1 5DD
Linda M Davies
*senior research fellow
in health economics*

Correspondence to:
E Berry e.berry@leeds.ac.uk

sis; ECST also showed a clear downward trend in the benefit of surgery for stenoses less than 70%.⁵ The benefit in this group also depended on the age and sex of the patients. However, a recent Cochrane review concluded that surgery was beneficial for patients with 50-69% stenosis,⁶ so we also evaluated the evidence on the diagnostic performance of magnetic resonance angiography when these more moderate stenoses were included (that is, for patients with 50-99% stenosis).

For each diagnostic threshold, we aimed to answer the question "What are the sensitivity and specificity of magnetic resonance angiography, in comparison with the gold standard of conventional angiography, in distinguishing severely stenosed arteries suitable for surgery, from either occluded or minimally stenosed arteries?"

Methods

We sought evaluations of the performance and effectiveness of magnetic resonance angiography in carotid artery stenosis published between January 1990 and December 1999. This review was part of a larger one, in which we also sought articles discussing the use of magnetic resonance angiography in peripheral vessels. We searched the electronic databases Medline, Embase, HealthSTAR, Science Citation Index, Index to Scientific and Technical Proceedings, Cochrane Library, Inside (British Library), and Online Computer Library Centre by using the keywords magnetic resonance angiography and MRA (or accepted synonyms and abbreviations).⁷ We also conducted a hand search of 10 key journals in the fields of imaging and vascular disease. We examined the reference lists of all articles retrieved from the above sources.

We compiled a list of unique articles by eliminating duplicates identified from more than one database. We applied preliminary exclusion criteria within the bibliographic databases, and one reviewer then applied these and a second list of exclusion criteria (table 1). We retrieved the remaining articles, and two reviewers applied final inclusion criteria (table 2) independently, in a hierarchical manner from A to H. Inclusion criteria A-C and F identified articles that described robust primary research comparing the diagnostic performance of magnetic resonance angiography against conventional angiography (A) in terms of sensitivity and specificity (B) and at the specific thresholds of stenosis

described in our introduction (C, F). Criterion D avoids results from the same study being included twice. The remaining criteria are indicators of the validity of the results. The results from articles that did not satisfy criterion E may have had verification bias; if the study group included asymptomatic participants (G) then patient cohort bias may have been present; and disease progression bias could result from a large time period between magnetic resonance angiography and the gold standard (H).⁸ Checklists covering study design, study group characteristics, technical details, and potential biases in executing the study were completed for each article satisfying the inclusion criteria A to D.

We included articles satisfying criteria A-D in the quantitative meta-analysis. Primary data (true positive, true negative, false positive, and false negative values) were extracted independently by two reviewers, and agreed by consensus, for the following clinical decision thresholds measured by using the NASCET criteria or similar⁴: 70-99% stenosed vessel (suitable for carotid endarterectomy) versus 0-69% stenosed or 100% occluded vessel (not suitable for carotid endarterectomy); 50-99% stenosed vessel (suitable for carotid endarterectomy) versus 0-49% stenosed or 100% occluded vessel (not suitable for carotid endarterectomy). Results derived with the very different ECST criteria were excluded from the meta-analysis.⁵

We plotted results from the independent studies on sensitivity versus 1 – specificity axes to illustrate the scatter of results. We then combined the sensitivity and specificity results for independent studies into a summary receiver operating characteristic curve by using the method recommended by the Cochrane Screening and Diagnostic Tests Methods Working Group for meta-analysis of diagnostic test results.⁹⁻¹² We used a logistic function to transform the sensitivity and specificity data. To prevent undefined values on logistic transformation, we applied a contingency correction of 0.01 to all true positive, false negative, true negative, and false positive values if any one was zero. We fitted a straight line to the transformed results by using an unweighted least squares fit, performed an inverse transformation on the fitted line, and plotted the result as a summary receiver operating characteristic curve.

The summary receiver operating characteristic curve is an excellent graphical summary, but for comparison purposes we calculated a further statistic— Q^* and its 95% confidence interval.⁹ Q^* is the point on the summary receiver operating characteristic curve where sensitivity and specificity have the same value; it represents the maximal joint sensitivity and specificity. Q^* is a good summary value in this application as there is no particular disadvantage to sensitivity and specificity being equal: patients with false positive results needlessly undergo the risks of surgery, but patients with false negative results are denied the benefits of surgery. Q^* would not be a good choice for describing a screening test, where the aim is to have no false negatives but a few false positives are acceptable.

We then performed a multiple linear regression analysis at the line fitting phase of the summary receiver operating characteristic analysis, to determine if any of five covariates had a significant effect, at the 95% level, on the fitted summary receiver operating characteristic curve.¹³ The five covariates were tech-

Table 1 Number of articles excluded after application of the exclusion criteria; 206 out of 7183 articles remained after these processes, of which 126 were on carotid artery stenosis

Reason for exclusion	Excluded within bibliographic databases	Excluded by reviewer (after exclusions within databases)
Review	1230	74
Editorial	123	0
Letter	108	6
Case report	1552	0
Conference abstract	473	14
Non-human study	400	7
Not on magnetic resonance angiography or in the applications of interest	Not applied	2694
Technical evaluation	Not applied	159
Paediatric participants	Not applied	14
≤10 participants	Not applied	123
Total No excluded	3886	3091

Table 2 Exclusions on application of final inclusion criteria. Only eight articles remained when all criteria A-H were applied. When only criteria A-D were used, 26 articles satisfied the criteria.¹⁶⁻⁴¹ References for the excluded articles are available from the authors

Inclusion criterion	No of articles excluded
A Study compared magnetic resonance angiography with digital subtraction angiography or cut film angiography	32
B Sufficient data reported for the construction of a 2x2 contingency table	44
C Performance at 50-99% or 70-99% or 100% stenosis reported	20
D Not a duplicate study of same patient group. Where more than one study was found, the one using the largest patient group was included.	5
E All patients in study received selective carotid intra-arterial digital subtraction or cut film angiography	13
F The method used to determine the degree of stenosis was described	4
G No asymptomatic patients were included	5
H No time delays of over 1 month occurred between examinations	1
Total No of articles excluded	124




Fig 1 Summary receiver operating characteristic curve for magnetic resonance angiography: 70-99% stenosis is a positive result, and 0-69% stenosis and 100% occlusion are negative results. Straight line shows 95% confidence interval of Q^* . Enlargement of region of sensitivity and specificity 90-100% shown for clarity

nique of magnetic resonance angiography, inclusion of articles that did not satisfy the inclusion criteria E-H, the risk of test or diagnostic review bias, the risk of verification bias, and the risk of withdrawal bias.¹⁴ As five variables were tested, the P value corresponding to 95% significance was 0.01.¹⁵ Statistical analyses were performed with SPSS for Windows (release 9.0.0).

Results

We identified 16 185 articles with the initial broad search strategy. After we had removed duplicates, 7183 unique articles remained. The exclusion criteria reduced the number to 206 (table 1), and of 126 candidate articles on carotid artery stenosis 26 satisfied the inclusion criteria A to D.¹⁶⁻⁴¹ Only eight articles satisfied all the inclusion criteria A to H.^{17 20 22 23 25 35 36 39} Six of these eight papers included results for the 70-99% threshold,^{17 22 23 25 35 36} and two included results for the 50-99% threshold.^{20 39}

For the diagnosis of 70-99% stenosis (fig 1), four sets of results obtained by using contrast enhanced techniques were included,^{26 32 34 36} together with 11 sets of results (from nine articles^{19 23 25 27 29 34 35 37 41}) obtained by using three dimensional time of flight and 10 sets of results¹⁶⁻³⁸ obtained by using two dimensional time of flight techniques. Q^* was 99% (95% confidence interval

98% to 100%). None of the variables tested in the multiple linear regression, including magnetic resonance angiography technique, was significant at the 95% level.

For the diagnosis of 50-99% stenosis (fig 2), no results obtained by using contrast enhanced techniques were included. Results from four studies using three dimensional time of flight techniques were included,^{21 29 39 40} together with six sets of results (from five articles^{20 24 28 31 33}) obtained by using two dimensional time of flight. Q^* was 90% (81% to 99%). None of the variables tested in the multiple linear regression was significant at the 95% level.

The patient populations of studies included in the quantitative meta-analysis were heterogeneous. Patient numbers ranged from 11 to 101 (mean=40). In all articles where sex distribution was reported most patients were men; in these articles the proportion of men ranged from 55% to 100% (mean=69%). Six articles did not report sex distribution.^{17 22 26 29 30 35} The lower limit for age of patients ranged from 18 to 63 years (mean=43 years), and the upper limit ranged from 73 to 87 years (mean=80 years). Six articles did not report age range.^{25 27 29 34 35 38} Eight articles stated that asymptomatic patients were included,^{16 18 27-29 31 32 40} and 18 articles gave no information about symptoms.^{16 17 19-26 28-30 33-37}




Fig 2 Summary receiver operating characteristic curve for magnetic resonance angiography: 50-99% stenosis is a positive result, and 0-49% stenosis and 100% occlusion are negative results. Straight line shows 95% confidence interval of Q^*

Discussion

Available evidence

Although many articles have been written about magnetic resonance angiography and carotid disease, little rigorous research has been conducted on the performance of magnetic resonance angiography in evaluating carotid artery stenosis. Small numbers of participants and inadequate details of study design mean that many studies included in this review have a potential for bias, but none of the factors tested in the multiple linear regression analysis had a significant effect on the results. Further sources of heterogeneity in patient populations (age, sex distribution, presenting symptoms) may influence patients' suitability for carotid endarterectomy,^{5 42} and differences existed between the study groups included in the review, even among those that satisfied all the inclusion criteria.

Potential for bias in this review arose from the increasingly common practice of screening candidates for magnetic resonance angiography by using ultrasonography and proceeding only if this shows a stenosis greater than a predetermined value. Some of the results used in this review were probably obtained in groups of patients preselected in this way, but we were not able to determine this from the articles.

Ultimately, to show effectiveness, we need evidence of the impact of magnetic resonance angiography on clinical decision making and on outcomes in patients.^{43 44} Comparative studies, with patients randomised to magnetic resonance angiography or conventional angiography, could be used to gather evidence of the impact on decision making. In this review, however, we found no studies that compared magnetic resonance angiography with conventional angiography for surgical decision making or outcomes. Although high quality research is needed, especially full cost effectiveness studies and evaluation of new magnetic resonance angiography techniques, further large trials involving conventional angiography are unlikely to be undertaken. Modelling, using existing evidence, may be the way forward.

Degree of stenosis

To be able to determine whether a patient is a suitable candidate for carotid endarterectomy, a diagnostic test must distinguish severely (>70%) stenosed or moderately (>50%) stenosed arteries (which are suitable for carotid endarterectomy) from both minimally stenosed (0-69% or 0-49%) and occluded arteries (100%), which are not suitable for carotid endarterectomy.^{4 5} Candidate articles often failed to assess magnetic resonance angiography in these terms. We excluded 20 articles from the review because they did not classify operable carotid artery stenosis as 70-99% or 50-99%. Of these, seven articles also failed to report the diagnostic performance of magnetic resonance angiography in separating severely stenosed from occluded arteries. This can be an important distinction in selection for surgery, especially if ultrasonography has indicated a small amount of flow.

Some authors have recommended that the magnitude of the likely benefit of carotid endarterectomy in cases of moderate to severe stenosis should be assessed by using more restricted bands of stenosis.⁴⁵ Future trials with restricted bands might show which stenosis thresholds, as determined by magnetic resonance

What is already known on this topic

Carotid endarterectomy for recently symptomatic carotid stenosis is beneficial in patients with 70-99% stenosis as measured by conventional angiography

It is not known whether the less invasive imaging technique of magnetic resonance angiography can accurately identify patients who will benefit from surgery

What this study adds

Magnetic resonance angiography is highly sensitive and specific in diagnosing 70-99% carotid stenosis

However, the studies on which this conclusion is based are of low quality and high heterogeneity

angiography, are associated with surgical benefit, in addition to determining the reliability of magnetic resonance angiography in detecting patients suitable for surgery. This research might be an ideal testing ground for tracker studies that evaluate fast changing technologies.⁴⁶

Conclusion

Our review does not support the use of magnetic resonance angiography to select surgical candidates with 50-99% stenosis. The 95% confidence interval for Q^* extended from 81% to 99%, and only two of the articles whose results were included in the meta-analysis satisfied the inclusion criteria related to validity. It would be advisable for users of magnetic resonance angiography to ensure that rigorous training and audit are in place, including feedback from surgeons and continuing quality control comparisons with ultrasonography.

Our results indicate that magnetic resonance angiography is very effective for detecting 70-99% stenosis as defined by conventional angiography. Q^* was 99% (98% to 100%). Although there is a promising trend towards better performance from contrast enhanced methods, further research is essential as only four articles were included in this review and no significant difference was found between the results obtained by using the three main techniques.

We thank all members of the review team for their contributions to the conduct of the systematic review, to hand searching, and to translation.

Contributors: MEW wrote the first draft of the paper and, with SK, performed the literature search and data extraction and participated in designing the review and writing the paper. EB coordinated the review and the writing of the paper and performed the data analysis. She will be guarantor. JMB, MJG, CMA, JFMM, LMD, and JC participated in the design and execution of the review and in writing the paper. MAS participated in writing the paper.

Funding: Financial support from the secretary of state for health under the NHS Health Technology Assessment Programme (97/13/04). The views and opinions expressed do not necessarily reflect those of the secretary of state for health. Leeds Teaching Hospitals NHS Trust received funding from the NHS Executive; the views expressed in this publication are those of the authors and not necessarily those of the NHS Executive.

Competing interests: JFMM has been reimbursed for presenting material to meetings organised by the pharmaceutical industry (Schering); JFMM has received funds for research on magnetic resonance angiography from both Philips Medical

Systems and Schering UK; JFMM has received funding from Philips Medical Systems for a part time research assistant; MAS has collaborative links with Philips Medical Systems, who have provided research support for magnetic resonance angiography.

- 1 Edelman RR. MR angiography: present and future. *Am J Roentgenol* 1993;161:1-11.
- 2 Erdoes LS, Marek JM, Mills JL, Berman SS, Whitehill T, Hunter GC, et al. The relative contributions of carotid duplex scanning, magnetic resonance angiography, and cerebral arteriography to clinical decision-making: a prospective study in patients with carotid occlusive disease. *J Vasc Surg* 1996;23:950-6.
- 3 Davies KN, Humphrey PR. Complications of cerebral angiography in patients with symptomatic carotid territory ischaemia screened by carotid ultrasound. *J Neurol Neurosurg Psychiatry* 1993;56:967-72.
- 4 Barnett HJ, Taylor DW, Eliasziw M, Fox AJ, Ferguson GG, Haynes RB, et al. Benefit of carotid endarterectomy in patients with symptomatic moderate or severe stenosis. *N Engl J Med* 1998;339:1415-25.
- 5 Randomised trial of endarterectomy for recently symptomatic carotid stenosis: final results of the MRC European carotid surgery trial (ECST). *Lancet* 1998;351:1379-87.
- 6 Cina CS, Clase CM, Haynes RB. Carotid endarterectomy for symptomatic carotid stenosis. *Cochrane Database Syst Rev* 2000;(3):CD001081.
- 7 Berry E, Kelly S, Hutton J, Harris KM, Smith MA. Identifying studies for systematic reviews—an example from medical imaging. *Int J Technol Assess Healthcare* 2000;16:668-72.
- 8 Kelly S, Berry E, Roderick P, Harris KM, Cullingworth J, Gathercole L, et al. The identification of bias in studies of the diagnostic performance of imaging modalities. *Br J Radiol* 1997;70:1028-35.
- 9 Moses LE, Shapiro D, Littenberg B. Combining independent studies of a diagnostic test into a summary ROC curve: data-analytic approaches and some additional considerations. *Stat Med* 1993;12:1293-316.
- 10 Irwig L, Tosteson AN, Gatsonis C, Lau J, Colditz G, Chalmers TC, et al. Guidelines for meta-analyses evaluating diagnostic tests. *Ann Intern Med* 1994;120:667-76.
- 11 Vamvakas EC. Meta-analyses of studies of the diagnostic accuracy of laboratory tests—a review of the concepts and methods. *Arch Pathol Lab Med* 1998;122:675-86.
- 12 Cochrane Methods Working Group on Systematic Review of Screening and Diagnostic Tests. Recommended methods [updated 6 June 1996]. <http://som.flinders.edu.au/fusa/cochrane/cochrane/sadtdoc1.htm> (accessed 9 Nov 2001).
- 13 Altman DG. *Practical statistics for medical research*. London: Chapman and Hall, 1991.
- 14 Reid MC, Lachs MS, Feinstein AR. Use of methodological standards in diagnostic test research. Getting better but still not good. *JAMA* 1995;274:645-51.
- 15 Gore SM. *Statistics in practice*. London: British Medical Association, 1982.
- 16 Bianchi MC. Magnetic resonance angiography vs digital angiography in surgical selection of carotid stenosis. *Nuova Rivista Di Neurologia* 1995;5:229-34.
- 17 Dadachanji MC, Shroff MM, Modi D, Jankharia BG. Comparison of MR angiography with contrast angiography for the diagnosis of carotid artery stenosis. *J Assoc Physicians India* 1995;43:92-5.
- 18 Drevet D, Russier S, Age B, Lepine PM, Zabet JM, Joffre P. Study of atheromatous stenoses of carotid bifurcations by Doppler ultrasound, spiral angio-MRI, magnetic resonance angiography and comparison with arteriography. *J Radiol* 1997;78:1271-7.
- 19 Fellner C, Strotzer M, Fraunhofer S, Held P, Spies V, Seitz J, et al. MR angiography of the supra-aortic arteries using a dedicated head and neck coil: image quality and assessment of stenoses. *Neuroradiology* 1997;39:763-71.
- 20 Huston J, Lewis BD, Wiebers DO, Meyer FB, Riederer SJ, Weaver AL. Carotid artery: prospective blinded comparison of two-dimensional time-of-flight MR angiography with conventional angiography and duplex US. *Radiology* 1993;186:339-44.
- 21 Kido DK, Panzer RJ, Szumowski J, Hollander J, Ketonen LM, Monajati A, et al. Clinical evaluation of stenosis of the carotid bifurcation with magnetic resonance angiographic techniques. *Arch Neurol* 1991;48:484-9.
- 22 Laster REJ, Acker JD, Halford HH, Nauert TC. Assessment of MR angiography versus arteriography for evaluation of cervical carotid bifurcation disease. *Am J Neuroradiol* 1993;14:681-8.
- 23 Link J, Brinkmann G, Steffens JC, Graessner J, Muller-Hulsbeck S, Heller M. MR angiography of the carotid arteries using 3D TOF technique with sagittal double-volume acquisition using a new head-neck coil. *Rofo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr* 1996;165:544-50.
- 24 Litt AW, Eidelman EM, Pinto RS, Riles TS, McLachlan SJ, Schwartzberg S, et al. Diagnosis of carotid artery stenosis: comparison of 2DFT time-of-flight MR angiography with contrast angiography in 50 patients. *Am J Neuroradiol* 1991;12:149-54.
- 25 Magarelli N, Scarabino T, Simeone AL, Florio F, Carriero A, Salvolini U, et al. Carotid stenosis: a comparison between MR and spiral CT angiography. *Neuroradiology* 1998;40:367-73.
- 26 Martinat P, Leclerc X, Gauvrit JY, Giboreau F, Pruvo JP. Contribution of fast-sequence three-dimensional MRI angiography with Gadolinium injection in the evaluation of supra-aortic vessels. *J Radiol* 1998;79:673-80.
- 27 Mattle HP, Kent KC, Edelman RR, Atkinson DJ, Skillman JJ. Evaluation of the extracranial carotid arteries: correlation of magnetic resonance angiography, duplex ultrasonography, and conventional angiography. *J Vasc Surg* 1991;13:838-44.
- 28 Nicholas GG, Osborne MA, Jaffe JW, Reed JF. Carotid artery stenosis: preoperative noninvasive evaluation in a community hospital. *J Vasc Surg* 1995;22:9-16.
- 29 Ozaki CK, Irwin PB, Flynn TC, Huber TS, Seeger JM. Surgical decision making for carotid endarterectomy and contemporary magnetic resonance angiography. *Am J Surg* 1999;178:182-4.
- 30 Pavone P, Catalano C, Di Girolamo M, Albertini PG, Marsili L, Passariello R. Angiography with magnetic resonance of the carotid arteries. Evaluation of clinical results obtained with low magnetic field equipment. *Radiol Med (Torino)* 1993;86:579-86.
- 31 Polak JF, Bajakian RL, O'Leary DH, Anderson MR, Donaldson MC, Jolesz FA. Detection of internal carotid artery stenosis: comparison of MR angiography, color Doppler sonography, and arteriography. *Radiology* 1992;182:35-40.
- 32 Remonda L, Heid O, Schroth G. Carotid artery stenosis, occlusion, and pseudo-occlusion: first-pass, gadolinium-enhanced, three-dimensional MR angiography—preliminary study. *Radiology* 1998;209:95-102.
- 33 Riles TS, Eidelman EM, Litt AW, Pinto RS, Oldford F, Schwartzberg GW. Comparison of magnetic resonance angiography, conventional angiography, and duplex scanning. *Stroke* 1992;23:341-6.
- 34 Sardanelli F, Zandrino F, Parodi RC, De Caro G. MR angiography of internal carotid arteries: breath-hold Gd-enhanced 3D fast imaging with steady-state precession versus unenhanced 2D and 3D time-of-flight techniques. *J Comput Assist Tomogr* 1999;23:208-15.
- 35 Scarabino T, Carriero A, Magarelli N, Florio F, Giannatempo GM, Bonomo L, et al. MR angiography in carotid stenosis: a comparison of three techniques. *Eur J Radiol* 1998;28:117-25.
- 36 Scarabino T, Carriero A, Giannatempo GM, Marano R, De Matthaeis P, Bonomo L, et al. Contrast-enhanced MR angiography (CE MRA) in the study of the carotid stenosis: comparison with digital subtraction angiography (DSA). *J Neuroradiologie* 1999;26:87-91.
- 37 Sitzler M, Furst G, Fischer H, Siebler M, Fehlings T, Kleinschmidt A, et al. Between-method correlation in quantifying internal carotid stenosis. *Stroke* 1993;24:1513-8.
- 38 Turnipseed WD, Kennell TW, Turski PA, Acher CW, Hoch JR. Magnetic resonance angiography and duplex imaging: noninvasive tests for selecting symptomatic carotid endarterectomy candidates. *Surgery* 1993;114:643-8.
- 39 Uehara T, Tabuchi M, Ohsumi Y, Yoneda Y, Mori E. Usefulness of 3-dimensional time-of-flight mr-angiography for evaluation of carotid-artery bifurcation stenosis. *Cerebrovasc Dis* 1995;5:199-203.
- 40 Wilkerson DK, Keller I, Mezrich R, Schroder WB, Sebok D, Gronlund J, et al. The comparative evaluation of three-dimensional magnetic resonance for carotid artery disease. *J Vasc Surg* 1991;14:803-9.
- 41 Young GR, Humphrey PR, Shaw MD, Nixon TE, Smith ET. Comparison of magnetic resonance angiography, duplex ultrasound, and digital subtraction angiography in assessment of extracranial internal carotid artery stenosis. *J Neurol Neurosurg Psychiatry* 1994;57:1466-78.
- 42 Bamford J. Risk stratification and carotid surgery: new technology but old trials. *Brain* 2001;124:455-6.
- 43 Thornbury JR, Eugene W. Caldwell Lecture. Clinical efficacy of diagnostic imaging: love it or leave it. *Am J Roentgenol* 1994;162:1-8.
- 44 Mackenzie R, Dixon AK. Measuring the effects of imaging: an evaluative framework. *Clin Radiol* 1995;50:513-8.
- 45 Rothwell PM, Gutnikov SA, Eliasziw M, Fox AJ, Taylor W, Mayberg MR, et al. Overall results of a pooled analysis of individual patient data from trials of endarterectomy for symptomatic carotid stenosis. *Stroke* 2001;32:327.
- 46 Lilford RJ, Braunholtz DA, Greenhalgh R, Edwards SJ. Trials and fast changing technologies: the case for tracker studies. *BMJ* 2000;320:43-6.

(Accepted 12 October 2001)