



UNIVERSITY OF LEEDS

This is a repository copy of *Agronomic cultivation, chemical composition, functional activities and applications of Pereskia species – A mini review*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/155200/>

Version: Accepted Version

Article:

Maciel, VBV, Yoshida, CMP and Goycoolea, FM (2019) Agronomic cultivation, chemical composition, functional activities and applications of *Pereskia* species – A mini review. *Current Medicinal Chemistry*, 26 (24). pp. 4573-4584. ISSN 0929-8673

<https://doi.org/10.2174/0929867325666180926151615>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Agronomic Cultivation, Chemical Composition, Functional Activities and Applications of Pereskia Species – A Mini Review

Vinicius B.V. Maciel^{1,3}, Cristiana M.P. Yoshida², and Francisco M. Goycoolea^{3,*}

¹The University of São Paulo, Faculty of Animal Science and Food Engineering, Av. Duque de Caxias Norte, 225 - CEP 13635-900, Pirassununga, SP, Brazil;

²Federal University of São Paulo, Institute of Ambiental, Chemistry and Pharmaceutical Science, Rua São Nicolau, 210, CEP 09913-030, Diadema, SP, Brazil;

³The University of Leeds, School of Food Science and Nutrition, LS2 9JT, Leeds, United Kingdom

ABSTRACT

Background: The exploration of the plant biodiversity as a natural source to obtain sustainable food products and new bioactive pharmaceutical compounds has been growing significantly due to their abundance, safety and economy. Natural pharmaceutical and edible compounds present some advantages when compared to synthetic ones, such as being chemically inert and widely available.. In this sense, plants of the genus *Pereskia* belonging to the Cactaceae family, have been studied. It is an unconventional wild edible plant that contains a large amount of protein and minerals. Studies have demonstrated their biological activities and potential application in different areas such as pharmaceutical, medicinal and food.

Objective: This review is focused on the chemical composition, functional properties, applications on pharmaceutical, nutraceutical and food areas and formulation techniques to enhance stability and bioavailability of bioactive compounds from the underutilized wild edible plant known as ora-pro-nobis (*Pereskia aculeata* or *Pereskia grandifolia*).

Conclusion: The latest studies involving ora-pro-nobis demonstrated its great potential due to its biological activities, which could stimulate further investigations. The utilization of this plant as a natural source to supplement the diet, or to prepare new food formulations and pharmaceutical products is an attractive approach to explore and fully realise the potential of the rich biodiversity found in Brazil and in other countries.

Keywords: ora-pro-nobis, *Pereskia aculeata*, *Pereskia grandifolia*, Cactaceae, unconventional plant, nutritional value, medicinal food.

1. INTRODUCTION

Biodiversity is essential for food security and nutrition, and could offer options for sustainable livelihoods (FAO, 2018). The world population is increasing and the fast depletion of natural resources are essential factors to propel the search for diversification on agriculture. Unconventional wild edible plants are those that are available from their natural habitat, easy access, affordable and could be used as food source due to their high nutritional value (Bhati and Jain, 2016; Beluhan and Ranogajee, 2010). In general, the consumption of underutilized plants is local or regional (Akinnifesi et al., 2006). The less importance to unconventional wild plants than conventional agricultural commodities is associated to the production and market value (Eleazar and Cesoiu, 2012). The total dietary energy intake of humans has been focused on relatively few plants species obtained from domesticated species: Cereals (e.g., barley, maize, millet, rice, rye, sorghum, sugar cane and wheat), tubers (e.g., cassava, potato, sweet potato and yam) (Grivetti and Ogle, 2000). Many underutilized and neglected species are nutritionally rich (e.g., high levels of essential minerals, vitamins, proteins, etc.) and adapted to low input agriculture. These plants can make an important contribution to a better and sustainable diet for local communities (Salvi and Ss, 2016).

The integration of indigenous trees of fruits and unconventional edible plants into farming systems could provide people's need for food and nutritional security. This implies a better knowledge of the nutritional quality of unconventional foods produced in agroforestry farming systems (Leaky et al., 1999). Many species of unconventional wild plants could be an effective way to help a diverse and healthy diet to combat micro- and macro-nutrients deficiencies in vulnerable social groups, particular in developing countries (Salvi and Ss, 2016). Leterme et al. (2006) studied the mineral composition of a large number of fruits and unconventional foods including nuts, leaves and tubers, produced by agroforestry programmes in the Andes and the rain forests of Colombia. They observed that the leaves were outstanding mineral sources, mainly calcium and iron. It was found that the cultivation conditions (e.g., soil fertility, pH, water supply, climate and seasonal variation) resulted in nutritional composition variability.

Tropical and subtropical countries concentrate a great diversity of plants including unconventional wild plants (Barreira et al., 2015). It has become increasingly evident that the nutritional potential of these unconventional wild plants could improve the macro- and micronutrients health status in human diets. The therapeutic potential of many indigenous plant species has also been studied and the discovery of new bioactive

compounds in unconventional wild and traditional medicinal plants is an active field of research (Azam et al., 2014).

Collaborative efforts among scientists and regional consumers should be stimulated to establish and develop innovative production and exploitation systems of promising unconventional wild plants. In this way, it could be possible to improve the availability of feed resources, while protecting the biodiversity (FAO, 2012). This is particularly relevant given the current climate change challenges and the increasing need to identify resilient crops.

A unique complete published list or online database with the total edible plants available in the world is not easy to find. Kunkel (1984) listed about 12,500 species with edible potential. Rapport & Drausal (2001) proposed that there are 27,000 species. Kinupp & Lorenzi (2014) listed 600 species of unconventional edible plants, including 351 most promising species found in the Brazilian flora, such as *Pereskia aculeata*, *Xanthosoma taioaba*, *Talinum paniculatum*, *Smallanthus sonchifolius*, *Acanthosyris spinescens*, *M. fluminensis* and *Vasconcellea quercifolia*, among others.

There are reports of the potential use of *Pereskia* species in different areas, food, pharmaceutical, medicinal, others. However, our knowledge of *Pereskia* and many other unconventional wild plants is scarce, and there is an urgent need to shift the focus of interest towards these type of plants, given their environmental and potential public health impact (Mattila et al., 2018). In this review we aim to present an overview of the current state of knowledge about *Pereskia*. We have kept the focus on its botanic, taxonomic and agronomical aspects, chemical composition, as well as its documented biological activities and current uses in pharmacy and foods.

2. THE GENUS *Pereskia*

Pereskia, a genus with 17 species, presents regular leaf development and function, generally viewed as representing the “ancestral cactus”. It includes *P. aculeata*, *P. aureiflora*, *P. bahiensis*, *P. bleo*, *P. diaz-romeroana*, *P. guamacho*, *P. grandifolia*, *P. horrida*, *P. lychnidiflora*, *P. marcanoi*, *P. nemorosa*, *P. portulacifolia*, *P. quisqueyana* Alain, *P. sacharosa*, *P. stenantha*, *P. weberiana* and *P. zinniiflora* (Leuenberger, 1986). In the present classification, it has been proposed the division of subfamily Pereskioideae in two genera only: *Pereskia* (17 spp.) and *Maihuenia* (2 spp.) (Hunt, 2016).

These species are primarily distributed in dry and arid regions of Caribbean, Central and South America (Edwards et al., 2005). It is believed that the northwest region of South America was the original centre of the diversification and spread of the genus *Pereskia* (Butterworth and Wallace, 2005). In Brazil, the main species commonly found are *P. aculeata* and *P. grandifolia* (Pinto and Scio, 2014).

Members of *Pereskia* genus have been described as having superior to inferior ovaries, broad, flattened leaves with C₃ photosynthesis, areoles with leaf production, dense, fibrous wood, a simple cortex without cortical bundles, poorly developed stem epidermal and hypodermal layers, non-succulent tissues or partially succulent leaves, and as inhabiting relatively mesic environments (Mauseth and Landrum, 1997).

Pereskia aculeata Miller and *Pereskia grandifolia* Haw.

P. aculeata Miller and *P. grandifolia* Haw., known as ora-pro-nobis or Barbados gooseberry (Pinto and Scio, 2014) are an unconventional wild edible plants. Both species are climbing vines plants that can reach up to 10 m in length (Figure 1a), with long branches and thorns (Figure 1b) in the armpit of the elliptical and succulent leaves (Duarte and Hayashi, 2005; Lorenzi and Souza, 1995). They are found in Brazil from the northeastern (Bahia) to the southern region (Minas Gerais, Paraná, Santa Catarina and Rio Grande do Sul) of the country (Conceição et al., 2014). Ora-pro-nobis, from the Latin means “pray of us”, is a species that belongs to the kingdom Plantae, class Magnoliopsida, order Caryophyllales, family Cactaceae and genus *Pereskia* (Almeida and Corrêa, 2012). Its natural habitat is the tropical forest, but it can be grown domestically in rural and urban areas (Souza et al., 2016a). *P. grandifolia* and *P. aculeata* are perennial plants that have mucilaginous leaves and present different colour of their flowers: in *P. aculeata* the flower is white (Figure 1c) and in *P. grandifolia* is pink (Figure 1d) (Kinupp and Lorenzi, 2014). They produce edible fruits with few seeds and which are dispersed by birds. The propagation of these type of plants can be carried out sexually or asexually, although the first method is minimally employed. It occurs due to the scarce information and expertise about the propagation. Despite the production time beings lower in sexual propagation, plants with desirable morphological characteristics are needed to ensure sustainable cultivation and identify the existing genetic diversity in these species (Souza et al., 2016b; Gibson and Nobel, 1990).




Figure 1. *Pereskia* (Cactaceae) species: (a) *Pereskia aculeata* Miller cultivated experimentally (6 months after cultivation) in University of São Paulo campus (Pirassununga, Brazil); (b) Detail of vegetative branch in *Pereskia aculeata* Miller, evidencing the leaves arrangement and presence of thorns (Duarte and Hayashi, 2005); (c) *Pereskia aculeata* Miller flower¹ (Santos et al., 2012); (d) *Pereskia grandifolia* Haw. Flower (image captured on 3rd May of 2018 and available in <https://www.flickr.com/photos/marcusviniciuslameiras/4894377091>).

Duarte and Hayashi (2005) developed a thorough anatomical study of the leaf and stem of *Pereskia aculeata*. They observed that leaves are simple and elliptical, and has uniseriate epidermis, paracytic stomata on both surfaces, dorsiventral-like mesophyll, several druses of calcium oxalate and large isodiametric cells. In the midrib a collateral bundle in open arc is embedded. The petiole exhibits a plain-convex transection and a collateral bundle in closed arc shape. In the stem part, in early secondary growth, has uniseriate epidermis, angular collenchyma, cortical parenchyma with many amiloplasts, perivascular fibre caps and collateral vascular organization. In the cortex and pith, numerous cavities containing mucilage and druses of calcium oxalate are seen, the latter also in the phloem.

There are isolated studies on growth characteristics and agronomic management of the *P. aculeata* and *P. grandifolia* species, but these are still scarce. Leuenberger (1992)

addressed general aspects of *Pereskia* species growth. Toffaneli and Resende (2011) studied the growth of *P. aculeata* in three different conditions: No support – free growth; espalier in the double string on the first thread; and espalier in the double string on the second thread. They observed that the treatment with no support presented the most promising results. Queiroz et al. (2015) evaluated the growth response of the *P. aculeata* under periodic drought through controlled reductions in the substrate matric potential. They found smaller dry mass accumulation in leaves (reduction of 21.4 %) than in stems (reduction of 48.1 %) and roots (reduction of 63.7 %). These results were noteworthy because the leaves are the main commercial part of the plant. Souza et al. (2016a) studied the effect of nitrogen fertilization on mineral, protein and nitrate contents, as well as the yield of *P. aculeata* leaves and verified that doses up to 400 kg ha⁻¹ of nitrogen ensure adequate leaf yield, protein and mineral contents within the desired range for the species, being a food rich in proteins, iron and calcium.

The form of propagation and multiplication using stem cuttings is the commonly applied. However, few studies have been published involving *ora-pro-nobis* cultivation. Souza et al. (2016b) studied the thermal effects on the germination of the *P. aculeata* and *P. grandifolia* species at different temperatures (24, 27, 30, 33 and 36 °C). The best germination response for *P. aculeata* and *P. grandifolia* occurred at 30 and 33 °C, respectively, with greater germination strength and fewer days to attain 63.21 % of germinations. Campos et al. (2017) evaluated the profile of buds of *ora-pro-nobis* in substrate of carbonized rice husk containing three treatments: apical, medial and basal cuttings. They concluded that medium and basal cuttings of the plant branches were more viable and showed more shoots at 37 days.

2.2 Chemical composition of *Pereskia aculeata* and *Pereskia grandifolia*

Nutritional compounds

Unconventional wild plants such as *Pereskia* species show, typically, higher nutrient levels (e.g., protein, minerals, vitamins) than the conventional ones. However, its production and consumption potentials are reduced and sometimes has been neglected (Oliveira et al., 2013). Thereby, the studies involving its nutritive potential are based on material obtained from institutional gardens or private yards (usually native plants) (Almeida and Corrêa 2012; Takeiti et al., 2009; Almeida Filho and Cambraia, 1974).

Leaves of *P. aculeata* are consumed as a vegetable, although its fruit is also edible (Souza et al., 2016a). The leaves present high levels of protein and minerals (Takeiti et

al., 2009; Almeida Filho and Cambraia 1974), and hydrocolloids (Conceição et al., 2014; Lima Junior et al. 2013). Furthermore, the significant content of fiber and the absence of leaf toxicity (Almeida Filho and Cambraia, 1974) of this species makes it a useful and attractive food source (Lima Junior et al., 2013).

The first report about the protein content of the ora-pro-nobis leaves was given by Almeida Filho and Cambraia (1974) as being high (17.4-25.4%) when compared with other common vegetables sources, such as corn (7.6-10 %, w/w), beans (18-20 %, w/w), spinach (2.2 %, w/w), kale (1.6 %, w/w) and lettuce (1.3 %, w/w). They observed that ora-pro-nobis leaves are extremely rich in high-quality proteins. Studies conducted on its leaves showed protein contents ranging between 17.4 g kg⁻¹ (Almeida Filho and Cambraia, 1974) to 28.9 g kg⁻¹ of dry matter (Almeida et al., 2014) and a high digestibility (85 %, Lima Junior et al., 2013). In addition to presenting a well-balanced composition, the leaves have an exceptionally high content of specific essential amino acids, particularly L-lysine. According to Mercê et al. (2001), the lysine constitutes 5.4 % w/w of the total protein content, which is 2–23 times the content found in any other previous cited vegetables. These authors affirmed that this observation is noteworthy, since lysine is an essential amino acid in animal nutrition, and it is deficient in cereals, which are the main source of food in many low-income communities (Pinto and Scio, 2014). The protein and essential amino acid levels reported are substantially higher than the minimum amount recommended by the Food and Agriculture Organization of the United Nations (FAO) as necessary for human consumption (Sierakowski et al., 1987). Furthermore, the protein content of ora-pro-nobis is similar or higher than those observed in taro (*Xanthosoma sagittifolium*, 27.59 %; Pinto et al., 2001), nightingale (*Urtica circularis*, 28.00 %), mestruz (*Coronopus didymus*, 28.17 %) and nightshade (*Solanum americanum*, 29.90 %) (Kinupp and Barros, 2008).

Takeiti et al. (2009) performed a complete study with ora-pro-nobis leaves evaluating the nutritional components regarding the proximal chemical composition, minerals, vitamins, amino acids, protein content and digestibility. The amino acids content, general composition and mineral contents are given in Tables 1, 2 and 3, respectively.

Table 1. Amino acid composition, FAO/WHO (1990) recommended allowances for preschool children (2-5 years old), in vitro protein digestibility of *Pereskia aculeata* leaves (Takeiti et al., 2009).

Amino acid	g/100 g (dry basis)^a	% of total	FAO/WHO (1990)
Essential			
Arginine	1.44 ± 0.02	5.32	
Histidine	0.59 ± 0.01	2.17	114
Isoleucine	1.07 ± 0.01	3.95	141
Leucine	2.00 ± 0.02	7.40	112
Lysine	1.43 ± 0.05	5.29	91
Methionine	0.23 ± 0.01	0.85	
Phenylalanine	1.27 ± 0.01	4.71	
Threonine	1.00 ± 0.01	3.71	109
Valine	1.28 ± 0.01	4.75	136
Tryptophan	5.52 ± 0.19	20.46	1860
∑ subtotal	15.83	59.61	181
Non-essential			
Aspartic acid	1.71 ± 0.22	6.32	
Serine	1.00 ± 0.01	3.71	
Glutamic acid	2.67 ± 0.03	9.90	
Proline	1.11 ± 0.01	4.10	
Cystine	0.35 ± 0.02	1.28	
Glycine	1.31 ± 0.01	4.86	
Alanine	1.36 ± 0.01	5.04	
Tyrosine	1.21 ± 0.03	4.49	
∑ subtotal	11.13	39.7	
Total sulphur amino acids (Met + Cys)	0.58	2.13	85
Total aromatic amino acids (Phe + Tyr)	2.48	9.2	146
In vitro protein digestibility (%)	75.90 ± 0.02		

^a Values are means of duplicate determinations carried out by Takeiti et al. (2009).

The ora-pro-nobis leaves contain high amounts of dietary fibre, minerals, such as calcium, magnesium, potassium, zinc and iron, besides vitamins C, B9 and E. Many of these nutritional components, mainly protein and minerals, are detected in higher proportions than those observed in other conventional leafy vegetables, such as kale, cabbage and spinach (Oliveira et al., 2013; Takeiti et al., 2009; Mercê et al., 2001; Morton, 1987; Almeida Filho and Cambraia, 1974).

Table 2. General composition of *Pereskia aculeata* leaves (g 100⁻¹) (Takeiti et al., 2009).

	P. aculeata (leaves)^a	Spinacia oleracea (spinach)^b	I. batatas poir (sweet potato leaves)^c
Moisture, fresh leaves	85.9 ± 0.2	90.7	87.1
Total protein ^d	28.4 ± 0.4 (3.1) ^e	3.2	3.8
Lipids	4.1 ± 0.3 (0.4) ^e	0.3	0.3
Ash	16.1 ± 0.1 (1.7) ^e	0.7	1.9
Crude fibre	9.8 ± 0.2 (1.0) ^e	0.6	-
Soluble dietary fibre	5.2 (0.5 ± 0.02) ^e	-	6.8 (0.9) ^e
Insoluble dietary fibre	33.9 (3.3 ± 0.07) ^e	-	39.1 (5.1) ^e
Total dietary fibre	39.1 (3.8 ± 0.06) ^e	3.5	45.9 (5.9) ^e

^aValues are means in triplicate determinations. Values expressed on a dry basis, except for moisture.

^bValues refer to g/100 g raw leaves according to Salazar et al. (2006).

^cValues refer to g/100 g raw leaves according to Ishida et al. (2000).

^d6.25_N g/100 g.

^eValues in parentheses are related to fresh green leaves.

Oliveira et al. (2013) determined the ascorbic acid (vitamin C) content in ora-pro-nobis leaves and compared it to the others unconventional wild plants. They found similar values of vitamin C for ora-pro-nobis and taro (*Xanthosoma sagittifolium*) ranging between 192.21 and 198.33 mg 100⁻¹ (fresh weight), respectively. Comparing these results with others observed in vegetables commonly used as food, such as kale (96.7 mg 100 g⁻¹) and rocket (46.3 mg 100 g⁻¹), it was concluded, that ora-pro-nobis leaves could be a good alternative source to supply the daily requirement of vitamin C.

Table 3. Mineral composition of fresh green *Pereskia aculeata* leaves (mg 100 g⁻¹) (Takeiti et al., 2009).

Minerals	P. aculeata (ora-pro-nobis leaves)^a	Spinacia oleracea (spinach)^b	I. batatas poir (sweet potato leaves)^c
Calcium	3420	106	187
Magnesium	1900	62	79
Potassium	1632	662	639
Phosphorus	156	51	68
Manganese	46.4	-	-
Zinc	26.7	0.2	0.8
Iron	14.2	3.1	5.4
Boron	5.55	-	-

Copper	1.4	0.2	0.4
--------	-----	-----	-----

^aValues are expressed in triplicate determinations (wet weight basis).

^bValues refer to mg/100 g raw leaves according to Salazar et al. (2006).

^cValues refer to mg/100 g raw leaves according to Ishida et al. (2000).

3. BIOLOGICAL ACTIVITIES

Anti-inflammatory activity

An intense chronic and acute topical anti-inflammatory activity against irritant agents were observed in hexane fractions of methanol extract of *P. aculeata* leaves applied in ear dermatitis in mice. This effect was associated to the presence of significant amounts of phytosterols (taraxerol, taraxasterol and phytol) (Pinto et al., 2015a). The authors concluded that the acute and chronic anti-inflammatory activity of *P. aculeata* leaves found is very promising and corroborates the need to understand their ethnopharmacological applications better.. Salt et al. (1987) identified sitosterol in *P. aculeata*, *P. bleo* and *P. grandifolia* leaves, which could indicate anti-inflammatory, antibacterial and antifungal characteristic. Purified plant sterols, including sitosterol, are known to have similar structure and functions as cholesterol. They are used in commercial functional foods in Europe (e.g., Benecol[®] and Pro-Activ[®]), with approved health claims to reduce cholesterol (Katan et al., 2003). Whether *Pereskia* leaves, either as a direct food source or its extracts, could have potential in these regards or not, is yet to be established.

Antioxidant activity

The antioxidant effect was observed in extracts of *P. grandifolia* leaves (Sim et al., 2010), *P. bleo* (Wahab et al., 2009) and *P. aculeata* (Pinto et al., 2012), and it was associated to the presence of phenolic compounds. The data obtained in their testing systems establish the antioxidant potency of *Pereskia* species. Souza et al. (2014) identified antioxidant activity with respect to DPPH radical in *P. aculeata* extracts prepared in acetone (80 %) and ethanol (70 %). They mentioned that the leaves of this species could be considered as a potential source of antioxidant compounds. According to Diplock et al. (1998), the presence of antioxidants could prevent and/or reduce the risk of major diseases such as heart diseases, arteriosclerosis, some cancers, Alzheimer and diabetes, associated with an excess of reactive free radicals.

Antimicrobial activity

Antimicrobial activities were found applying ethyl acetate extracts of *P. grandifolia*

leaves against *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Bacillus subtilis* (Philip et al., 2009). A moderate activity against *P. aeruginosa* and *Salmonella choleraesuis* was observed for *P. bleo* hexane and methanol leaves extract (Wahab et al., 2009). The ether extract of *P. sacharosa* leaves indicated an antimicrobial activity against *Helicobacter pylori*, a bacterium associated with gastric ulcers (Uyub et al., 2010).

Cytotoxicity

Recent studies have addressed the cytotoxic activity of *Pereskia aculeata* ethyl acetate extracts against different human cell lines (Carvalho et al., 2014; Nurestri et al., 2009). For example, nasopharyngeal epidermoid carcinoma cell line (KB), cervical carcinoma cell line (CasKi), colon carcinoma cell line (HCT 116), hormone-dependent breast carcinoma cell line (MCF-7), lung carcinoma cell line (A549) and the non-cancer human fibroblast cell line (MRC-5) were identified using in vitro cytotoxicity assay. The results indicated that *P. grandifolia* possessed remarkable cytotoxicity effect against KB and MCF-7 cell lines. The authors also identified the compounds β -sitosterol, vitamin E, phytone, 2,4-di-tert-butylphenol and a mixture consisting of 2,4-ditertbutylphenol, methyl palmitate, methyl oleate and methyl stearate, isolated from the extract of *P. grandifolia* in ethyl acetate. The observed cytotoxic activity against KB cells was attributed to the compound 2,4-di-tert-butylphenol. Malek et al. (2009, 2008) also found the presence this compound in *P. bleo* extracts and confirmed the cytotoxicity against KB cells. By contrast, in our own studies (Maciel et al., 2018), we have shown that *P. aculeata* aqueous extracts administered to Caco-2 cells did not show any cytotoxic effects within the dose range studied (185-890 ng mL⁻¹). The results seem to indicate that the type of phytochemicals present in the aqueous extracts of *ora-pro-nobis* differ notably from those present in the less polar extracts and are far less cytotoxic.

Diuretic and hypotensive activities

Pereskia grandifolia is well recognized in Brazilian traditional medicine as a diuretic agent. Kazama et al. (2012) reported diuretic activity in aqueous extract of *P. grandifolia* leaves tested in rats. They associated the results to the inhibition of arginine vasopressin release. The authors also evaluated the ethanolic extract from *P. grandifolia* leaves. They observed that besides of the low capacity of this extract in to induce potassium and chloride excretion, without showing signs of acute toxicity. The results indicated that its bioactive components have a high potential as an adjuvant in some

kidney diseases and disorders of arginine-vasopressin secretion.

Antinociceptive activity

This bioactivity has not been reported in the more common *P. aculeata* and *P. grandifolia* extracts. However, in *P. bleo* was studied by Abdul-Wahab et al. (2012). They found reasonable antinociceptive activity in mice from the leaves used to produce different extracts and suggested this effect to peripheral mechanisms of pain control. Guilhon et al. (2012) investigated the antinociceptive activity of the ethanolic extract and fractions obtained from *P. bleo* leaves. The authors suggested that the results could be mediated, probably, by the central mechanisms related to the opioid system. Pinto et al. (2015b), evaluated the antinociceptive activity of the hydromethanolic fraction obtained from the methanol extract of *P. aculeata* leaves. The results suggested peripheral and central antinociception devoid of an opioid effect. According to those authors, it was the first time that this bioactivity (analgesic potential) is reported for *P. aculeata*.

4. APPLICATIONS IN PHARMACEUTICAL, NUTRACEUTICALS AND FOOD

Unconventional wild plants commonly serve multiple functions on dietary and therapeutic uses, such as dye, dietary fibre, food, medicine and oil (Grivetti and Ogle, 2000). The applications of *Pereskia* species are usually limited and need more research to explore the species of this genus. The published data indicates the *Pereskia* species potential as active biological agent and nutritional supplements (Pinto and Scio, 2014).

Pharmaceutical uses

Plants produce metabolites, and many of them show versatile pharmacological activities (Wahab et al., 2009). The chemical compounds of *Pereskia* species are of potential pharmaceutical application relevance. The fruits of *P. aculeata* are rich in carotenoids (namely, α -carotene, β -carotene, lutein, α -cryptoxanthin/zeinoxanthin, and β -cryptoxanthin) (Agostini-Costa et al., 2012). The high levels of α -carotene and zeaxanthin found in berries and leaves, respectively, could be carotenoid biomarker characteristics for *Pereskia* species, which may be important sources of functional compounds for health-promotion (Agostini-Costa et al., 2014).

The fruits of *P. aculeata* and *P. grandifolia* are used in folk medicine as antisiphilitics and expectorants, while leaves of *P. aculeata* are applied as an emollient, treatment of inflammation and skin diseases (Duarte and Hayashi, 2005; Farago et al.,

2004; Cruz, 1995). The leaves of both species are used in regional cuisine by natives in some regions of Brazil (Pinto and Scio, 2014; Almeida and Corrêa, 2012), and *P. grandifolia* is cultivated mainly as an ornamental plant and to form hedges (Moran and Zimmermann, 1991). Malaysia is another country that use largely *P. grandifolia*, where the plant was introduced since ~50-60 years ago. It is employed by traditional medicine to treat several types of diseases: cancer, headache, stomachache, gastric ulcers, haemorrhoids, dermatitis, diabetes, hypertension, rheumatism, inflammation, and as a tonic to revitalize the body (Sim et al., 2010). Murilo et al. (2010) indicated the use of *P. bleo* in traditional medicine in Panama to treat gastrointestinal disorders and for human nutrition.

Wound healing activity has been attributed to the mucilage of cactus leaves (Thornfeldt, 2005; Aburjai and Natsheh, 2003). However, this property has never been tested under experimental conditions (Carvalho et al., 2014). Wound healing is a complex process involving steps of cellular migration and proliferation, especially, but not only, of fibroblast cells (Krishnan, 2006). It is expected that the compound to be tested should be effective in promoting the cellular proliferation and migration of fibroblast, as indication of wound healing property. In studies performed by Carvalho et al. (2014) with hydroethanolic extract obtained from *P. aculeata* leaves, it was evidenced the beneficial effect of the mucilage containing extract on the fibroblast cell culture. It corroborates the folk use of *P. aculeata* for wound healing properties. Sartor et al. (2010) also observed the healing capacity of the crude extract of *P. aculeata* leaves applied directly on skin wound of rats. Pinto et al. (2016) investigated the topical wound healing activity of gels containing the methanol extract and hexane fractions of the *P. aculeata* leaves in a model of excisional wound healing in mice. The wound healing process was considerably accelerated, mainly by the hexane fraction gel formulation.

Nutraceutical and food application

Consumers' concerns about quality and healthy foods are increasing. Sobrinho et al. (2015) incorporated 1 and 2 % (w/w) of *ora-pro-nobis* (*P. aculeata*) leaves flour in sausage formulations aiming to make this meat product healthier. This addition promoted an increase of the protein and fibre contents in sausages, improving the sensory properties, particularly colour and texture of the product. Fermented milk beverages were obtained adding mucilage extract from *P. aculeata* as a gelling or emulsifier ingredient. The final product showed higher protein content, firmness and adhesiveness (Amaral et al., 2018).

Fortified bread was developed adding 6 % (w/w) of ora-pro-nobis flour (*P. aculeata*), increasing the fibre and protein content with good sensory global acceptance (Martinevski et al., 2013). Rocha et al. (2008) developed a pasta containing 1.0, 1.5 and 2.0 % of ora-pro-nobis flour, increasing the protein, fibre and mineral contents when compared the product with the control pasta. Zem et al. (2017) produced flour of *P. aculeata* and investigated its effects on a diet applied in rats. They concluded that it might not be adequate to warrant satisfactory growth levels of the animals when provided in the diet a single protein source (*P. aculeata* flour). However, the authors suggested that due to few limiting factors, it may be targeted for protein supplementation with other vegetal sources and thus it may provide for the protein needs in human diets.

Prajapati et al. (2013) elaborated a review involving the pharmaceutical applications of various natural gums, mucilages and their modified forms. They mentioned that a large number of gum and mucilage plant-based pharmaceutical and cosmetic excipients are available today. These excipients are attractive for a wide range of formulations, given their natural abundance, safety, and amenability for “bio” or “organic” products.. Natural materials have advantages over synthetic ones since they are chemically inert, nontoxic, less expensive, biodegradable and widely available. Recent trend toward the use of plant based and natural products demands the replacement of synthetic additives with natural ones. Some recent works has been published in this context involving *P. aculeata*. Conceição et al. (2014) and Lima Junior et al. (2013) investigated the extraction process of the gum and hydrocolloids from the extract of *P. aculeata* leaves. They proved that this plant could be an alternative source of viable hydrocolloids, founding the final product rich in protein and minerals. The authors also evaluated the potential use of the powdered product as an emulsifying and stabilizing agent in food applications. Studies of the complex nature of the biopolymers such as arabinogalactans, extracted from ora-pro-nobis leaves and their interactions with Co^{2+} , Cu^{2+} , Mn^{2+} and Ni^{2+} regarding the thermal stability of the metallic compounds were performed by Sierakowski et al. (1990) and Mercê et al. (2001). In their results, it was suggested the potential use of the hydrocolloids in food and pharmaceutical industries. Silva et al. (2017) also produced mucilage derived from *P. aculeata* evaluating different temperatures (30, 45 and 60 °C) to obtain the extract.

Barbalho et al. (2016) studied the *P. aculeata* flour on metabolic profile and intestinal motility of Wistar rats, and observed a total decreased of lipids, weight gain, visceral fat and increased levels of HDL-c (high density lipoprotein). They found that the

flour ingestion reduced the risk of cardiovascular disease by 67.1 %. The beneficial effects of *P. aculeata* flour were attributed to the higher content of vitamin C, carotenoids and fibres. Almeida and Corrêa (2012) studied rats fed with hypercaloric diet and concluded that the daily ingestion of *P. grandifolia* flour ingestion could prevent obesity and cardiovascular diseases due to the effectiveness reduced weight gain, body mass index, glycemia and lower levels of triglycerides.

5. FORMULATION TECHNIQUES TO ENHANCE STABILITY AND BIOAVAILABILITY OF BIOACTIVE COMPOUNDS

The functionality of plant extracts and pure compounds isolated from medicinal plants is highly dependent on the stability, appearance, taste and flavour, formulation and active ingredient availability. Often, active ingredients such as food supplements (e.g. omega-3 fatty acids) or micronutrients (e.g., iron), can often cause detrimental changes in product formulation. These can be associated with their inherent properties or with their interactions with other ingredients. Colloidal delivery systems are used to achieve an adequate balance between the solubility and dispersibility of active ingredients to overcome the technical challenges associated with chemical stability, taste, bioavailability, sedimentation and solubility (Velikov, 2012).

Studies in our laboratories (Maciel et al., 2018), have aimed to develop a colloidal carrier system loaded with ora-pro-nobis (*P. aculeata*) aqueous extract, as a potential formulation for iron food fortification. To this end, we prepared chitosan/pectin polyelectrolyte electrostatic self-assembled complexes and evaluated their physicochemical and cellular iron bioavailability in Caco-2 cells. The thus obtained particles were in the diameter size range of ~1.0 to 7.0 μm , depending on the relative molar charge ratio (n^+/n^-) of oppositely charged chitosan (positive) and pectin (negative). We gained proof-of-concept that the systems comprised by an excess of chitosan ($n^+/n^- > 1.0$) led to higher cellular ferritin levels than, although these were lower than for the free ora-pro-nobis extract (6.81 ± 0.36 vs. 174.96 ± 15.44 ng/mg cell protein, respectively). In other studies (Omwenga et al., 2018; Qin et al., 2018), we have also demonstrated the effectiveness of associating lipophilic flavonoid compounds such as quercetin, baicalein and cinnamaldehyde in oil-core nanocapsules coated with chitosan. These studies show consistently, that the encapsulation of these bioactive compounds leads to enhanced bioactivity against bacterial quorum sensing and biofilm formation. Also, the encapsulation of the flavonoids quercetin and baicalein, has a cytoprotective effect.

Whether some of the bioactive lipophilic compounds present in ora-pro-nobis non-polar extracts would be amenable for nanoencapsulation to modulate their bioactivity, is yet to be established in future studies.

6. CONCLUSION

Few reports on the identified phytochemical compounds, nutrients and functional properties suggest that the ora-pro-nobis (*Pereskia aculeata* and *Pereskia grandifolia*), an underutilized indigenous edible plant, should be better investigated. From the ongoing research worldwide, and with the current available database it was evidenced that the ora-pro-nobis possess high nutritional value and still presents interesting potential for the development of supplementation and fortification formulation of minerals, vitamins and protein, as well as other health benefits that are yet to be discovered. This review has sought to present an updated account of the current knowledge on the different applications of ora-pro-nobis. It is clear that further investigations should be conducted so that the potential uses observed in pharmaceutical, nutraceutical and food area of this plant are fully realised. In the years to come, there will be continued interest in both novel and traditional natural plant sources and their uses in the development of novel materials as health enhancers.

8. REFERENCES

- Abdul-Wahab, I.R.; Guilhon, C.C.; Fernandes, P.D.; Boylan, F. Antinociceptive activity of *Pereskia bleo* Kunth (Cactaceae) leaves extracts. *J. Ethnopharmacol.*, **2012**, 144(3), 741-746.
- Aburjai, T.; Natsheh, F. Plants used in cosmetics. *Phytother. Res.*, **2003**, 17(9), 987-1000.
- Agostini-Costa, T.S.; Pêsoa, G.K.A.; Silva, D.B.; Gomes, I.S.; Silva, J.P. Carotenoid composition of berries and leaves from a Cactaceae – *Pereskia* sp. *J. Funct. Foods*, **2014**, 11, 178-184.
- Agostini-Costa, T.S.; Wondracek, D.C.; Rocha, W.S.; Silva, D.B. Carotenoids profile and total polyphenols in fruits of *Pereskia aculeata* Miller. *Rev. Bras. Frutic.*, **2012**, 34(1), 234-238.
- Almeida, M.E.F.; Junqueira, A.M.B.; Simão, A.A.; Corrêa, A.D. Caracterização química das hortaliças não-convencionais conhecidas como ora-pro-nobis. *Biosci. J.*, **2014**, 30(1), 431-439,

Almeida Filho, J.; Cambraia, J. Estudo do valor nutritivo do “Ora-pro-nobis” (*Pereskia aculeata* Mill.). *Rev. Ceres*, **1974**, 21(114), 105-111.

Almeida, M.E.F.; Corrêa, A.D. Utilização de cactáceas do gênero *Pereskia* na alimentação humana em um município de Minas Gerais. *Ciênc. Rural*, **2012**, 42(4), 751-756.

Amaral, T.N.; Junqueira, L.A.; Prado, M.E.T.; Cirillo, M.A.; Abreu, L.R.; Costa, F.F.; Resende, J.V. Blends of *Pereskia aculeata* Miller mucilage, guar gum, and gum Arabic added to fermented milk beverages. *Food Hydrocoll.*, **2018**, 79, 331-342.

Azam, F.M.S.; Biswas, A.; Mannan, A.; Afsana, N.A.; Jahan, R.; Rahmatullah, M. Are famine food plants also ethnomedicinal plants? An Ethnomedicinal appraisal of famine food plants of two districts of Bangladesh. *Evid. Based Complement. Alternat. Med.*, **2014**, Article ID 741712.

Barbalho, S.M.; Guiguer, E.L.; Marinelli, P.S.; Do Santos Bueno, P.C.; Pescinini-Salzedas, L.M.; Dos Santos, M.C.; Oshiiwa, M.; Mendes, C.G.; de Menezes, M.L.; Nicolau, C.C.; Otoboni, A.M.; de Alvares Goulart, R. *Pereskia aculeata* Miller flour: metabolic effects and composition. *J. Med. Food*, **2016**, 19(9), 890-894.

Barreira, T.F.; Paula Filho, G.X.; Rodrigues, V.C.C.; Andrade, F.M.C.; Santos, R.H.S.; Priore, S.E.; Pinheiro-Sant’ana, H.M. Diversity and equivalence of unconventional food plants in rural zone of Viçosa, Minas Gerais, Brazil. *Rev. Bras. Plant. Med.*, **2015**, 17(4), 964-974.

Beluhan, S.; Ranogajee, A. Chemical composition and nonvolatile components of *Crotial* wild edible mushrooms, *Food Chem.*, **2010**, 124, 1076-1082.

Bhati, D.; Jain, S. Micro nutrient composition of unconventional wild fruits and vegetables. *The Biascan*, **2016**, 11(4), 2405-2409

Butterworth, C.A.; Wallace, R.S. Molecular phylogenetics of the leafy cactus genus *Pereskia* (Cactaceae). *Syst. Bot.*, **2005**, 30(4), 800-808.

Campos, J.A.; Oliveira, N.J.F.; Chamba, J.S.V.; Colen, F.; Costa, C.A.; Filho, A.S.S. Shoot growth of ora-pro-nobis in alternative substrate of carbonized rice husk. *HOLOS*, **2017**, 7, 148-167.

Carvalho, E.G.; Soares, C.P.; Blau, L.; Menegon, R.F.; Joaquim, W.M. Wound healing properties and mucilage content of *Pereskia aculeata* from different substrates. *Rev. Bras. Farmacogn.*, **2014**, 24, 677-682.

Conceição, M.C.; Junqueira, L.A.; Silva, K.C.G.; Prado, M.E.T.; Resende, J.V. Thermal and microstructural stability of a powdered gum derived from *Pereskia aculeata* Miller

leaves. *Food Hydrocolloid.*, **2014**, 40, 104-114.

Cruz, G.L. *Dicionário das plantas úteis do Brasil*, 5th ed.; Bertrand Brasil:Rio de Janeiro, **1995**.

Diplock, A.T.; Charleux, J.L.; Crozier-Willi, G.; Kok, F.J.; Rice-Evans, C.; Roberfroid, M.; Stahl, W.; Viña-Ribes, J. Functional food science and defence against reactive oxidative. *Br. J. Nut.*, **1998**, 80(S1), S77-S112.

Duarte, M.R.; Hayashi, S.S. Estudo anatômico de folha e caule de *Pereskia aculeata* Mill. (Cactaceae). *Rev. Bras. Farmacogn.*, **2005**, 15(2), 103-109.

Edwards, E.J.; Nyffeler, R.; Donoghue, M.J. Basal cactus phylogeny: implications of *Pereskia* (Cactaceae) paraphyly for the transition to the cactus life form. *Am. J. Bot.*, **2005**, 92(7), 1177-1188.

Eleazar, N.P.; Cesoi. Recognition of importance of underutilized fruit crops. *R & D Notes. Bar Res. Develop. Not.* **2012**, 14(4), 3.

FAO. *Biodiversity: FAO's role in biodiversity*. (Available from: <http://www.fao.org/biodiversity/en/>), **2018**.

FAO. Use of lesser-known plants and plant parts as animal feed resources in tropical regions, by E.S., Quansah.; H.P.S., Makkar. *Animal Production and Health Working Paper*, **2012**.

Farago, P.V.; Takeda, I.J.M.; Budel, J.M.; Duarte, M.R. Análise morfo-anatômica de folhas de *Pereskia grandifolia* Haw. Cactaceae. *Lat. Am. J. Pharm.*, **2004**, 23(3), 323-327.

Gibson, A.C.; Nobel, P.S. *The Cactus Primer*, Harvard University Press: Cambridge, **1990**.

Grivetti, L.E.; Ogle, B.M. Value of traditional foods in meeting macro-and micronutrient needs: the wild plant connection. *Nut. Res. Rev.*, **2000**, 13(1), 31-46.

Guilhon, C.C.; Wahab, I.R.A.; Boylan, F.; Fernandes, P.D. Evaluation of antinociceptive activity of *Pereskia bleo* Kunth. *Plant. Med.*, **2012**, 78(11), 1098.

Hunt, D. *Cites cactaceae checklist*, 3rd ed.; Royal Botanic Gardens: Kew (England), **2016**.

Katan, M.B.; Grundy, S.M.; Jones, P.; Law, M.; Miettinen, T.; Paoletti, R. Efficacy and safety of plant stanols and sterols in the management of blood cholesterol levels. *Mayo Clin. Proc.*, **2003**, 78(8), 965-978.

Kazama, C.C.; Uchida, D.T.; Canzi, K.N.; Souza, P.; Crestani, S.; Junior, A.G.; Junior, A.L. Involvement of arginine-vasopressin in the diuretic and hypotensive effects of

Pereskia grandifolia Haw. (Cactaceae). *J. Ethopharmacol.*, **2012**, 144(1), 86-93.

Kinupp, V.F.; Barros, I.B.I. Teores de proteína e minerais de espécies nativas, potenciais hortaliças e frutas. *Food Sci. Technol.*, **2008**, 28(4), 846-857.

Kinupp, V.F.; Lorenzi, H. Plantas alimentícias não convencionais (PANC) no Brasil, 1st ed.; Plantarum: Nova Odessa, **2014**.

Krishnan, P. The scientific study of herbal wound healing therapies: Current state of play. *Curr. Anaesth. Crit. Care.*, **2006**, 17(1-2), 21-27.

Kunkel, G. Plants for human consumption: a noted checklist of the edible phanerogams and ferns. Koeltz Scientific Books: Koenigstein, **1984**.

Leakey, R. Potential for novel food products from agroforestry trees: a review. *Food Chem.*, **1999**, 66(1), 1-14.

Leterme, P.; Buldgen, A.; Estrada, F.; Londono, A.M. Mineral content of tropical fruits and unconventional foods of the Andes and the rain forest of Colombia. *Food Chem.*, **2006**, 95(4), 644-852.

Leuenberger, B.E. *Pereskia* (Cactaceae). *Mem. N. Y. Bot. Gard.*, **1986**, 41, 1-141.

Leuenberger, B.E. Leaf-bearing cacti (genus *Pereskia*) in cultivation. *Cact. Succ. J.*, **1992**, 64(5), 247-261.

Lorenzi H.; Souza, H.M. Plantas ornamentais no Brasil: arbustivas, herbáceas e trepadeiras, 1st ed.; Plantarum: Nova Odessa, **1995**.

Maciel, V.; Yoshida, C.; Boesch, C.; Goycoolea, F.; Carvalho, R. Iron uptake by Caco-2 cells from a Brazilian natural plant extract loaded into chitosan/pectin nano- and micro-particles. *Proc. Nut. Soc.*, **2018**, 77(OCE2), E45. DOI: 10.1017/S0029665118000393.

Malek, S.N.A.; Shin, S.K.; Wahab, N.A.; Yaacob, H. Cytotoxic components of *Pereskia bleo* (Kunth) DC. (Cactaceae) leaves. *Mol.*, **2009**, 14, 1713-1724.

Malek, S.N.A.; Wahab, N.A.; Yaacob, H.; Shin, S.K.; Lai, H.S.; Serm, L.G.; Rahman, S.N.S.A. Cytotoxic activity of *Pereskia bleo* (Cactaceae) against selected human cell lines. *Int. J. Canc. Res.*, **2008**, 4(1), 20-27.

Martinevski, C.S.; Oliveira, V.R.; Rios, A.O.; Flores, S.H.; Venzke, J.G. Utilização de bertalha (*Anrederacordifolia*(ten.) Steenis) e ora-pro-nobis (*Pereskia aculeata* Mill.) na elaboração de pães. *Alim. Nutr.*, **2013**, 24(3), 1-6.

Matilla, P.; Makinen, S.; Euroola, M.; Jalava, T.; Pihlava, J-M; Hellstrom, J.; Pihlanto, A. Nutritional value of commercial protein-rich plant products. *Plant Foods Hum. Nut.*, **2018**, 1-8. Available from: <https://link.springer.com/content/pdf/10.1007%2Fs11130-018-0660-7.pdf>.

Mauseth, J.D.; Landrum, J.V. Relictual vegetative anatomical characters in Cactaceae: the genus *Pereskia*. *J. Plant Res.*, **1997**, 110(1), 55-64.

Mercê, A.L.R.; Landaluze, J.S.; Mangrich, A.S.; Szpoganicz, B.; Sierakowski, M.R. Complexes of arabinogalactan of *Pereskia aculeata* and Co^{2+} , Cu^{2+} , Mn^{2+} , and Ni^{2+} . *Bioresour. Technol.*, **2001**, 76(1), 29-37.

Moran, V.C.; Zimmermann, H.G. Biological control of cactus weeds of minor importance in South Africa. *Agric. Ecosyst. Environ.*, **1991**, 37, 37-55.

Murilo, E.; Meléndez-Matínez, A.J.; Portugal, F. Screening of vegetables and fruits from Panama for rich sources of lutein and zeaxanthin. *Food Chem*, **2010**, 122(1), 167-172.

Nurestri, A.M.S.; Sim, K.S.; Norhanom, A.W. Phytochemical and cytotoxic investigations of *Pereskia grandifolia* Haw (Cactaceae) leaves. *J. Biol. Sci.*, **2009**, 9(5), 488-493.

Oliveira, D.C.S.; Wobeto, C.; Zanuzo, M.R.; Severgnini, C. Composição mineral e teor de ácido ascórbico nas folhas de quatro espécies olerícolas não-convencionais. *Hort. Bras.*, **2013**, 31, 472-475.

Omwenga E.O.; Hensel A.; Shitandi A.; Goycoolea F.M. Chitosan nanoencapsulation of flavonoids enhances their quorum sensing and biofilm formation inhibitory activities against an *E. coli* Top 10 biosensor. *Colloids Surf. B Biointerfaces*, **2018**, 164, 125-133.

Philip, K.; Malek, S.N.A.; Sani, W.; Shin, S.K.; Kumar, S.; Lai, H.S.; Serm, L.G.; Rahman, S.N.S.A. Antimicrobial activity of some medicinal plants from Malaysia. *Am J Appl. Sci.*, **2009**, 6(8), 1613-1617.

Pinto, N.A.V.D.; Carvalho, V.D.; Corrêa, A.D.; Rios, A.O. Avaliação de fatores antinutricionais das folhas da taioba (*Xanthosoma sagittifolium* Schoot). *Ciênc, Agrotec.*, **2001**, 25(3), 601-604.

Pinto, N.A.V.D.; Duque, A.P.N.; Pacheco, N.R.; Mendes, R.F.; Silva Motta, E.V.S.; Bellozi, P.M.Q.; Ribeiro, A.; Salvador, M.J.; Scio, E. *Pereskia aculeata*: A plant food with antinociceptive activity. *Pharm. Biol.*, **2015b**, 53(12), 1780-1785.

Pinto, N.A.V.D.; Machado, D.C.; Silva, J.M.; Conegundes, J.L.M.; Gualberto, A.C.M.; Gameiro, J.; Chedier, L.M.; Castañón, M.C.M.N.; Scio, E. *Pereskia aculeata* Miller leaves present in vivo topical anti-inflammatory activity in models of acute and chronic dermatitis. *J. Ethnopharmacol.*, **2015a**, 173, 330-337.

Pinto, N.C.C.; Santos, R.C.; Machado, D.C.; Florêncio, J.R.; Fagundes, E.M.Z.; Antinarelli, L.M.R.; Coimbra, E.S.; Ribeiro, A.; Scio, E. Cytotoxic and antioxidant activity of *Pereskia aculeata* Miller. *Pharm. On.*, **2012**, 3, 6369.

- Pinto, N.C.C.; Scio, E. The biological activities and chemical composition of *Pereskia* species (Cactaceae)—A Review. *Plant Foods Hum. Nutr.*, **2014**, 69, 189-195.
- Pinto, N.C.C.; Cassini-Vieira, P.; Souza-Fagunde, E.M.; Barcelos, L.S.; Castañón, M.C.M.N.; Scio, E. *Pereskia aculeata* Miller leaves accelerate excisional wound healing in mice. *J. Ethnopharmacol.*, **2016**, 194, 131-136.
- Prajapati, V.D.; Jani, G.K.; Moradiya, N.G.; Randeria, N.P. Pharmaceutical applications of various natural gums, mucilages and their modified forms. *Carbohydr. Polym.*, **2013**, 92, 1685-1699.
- Qin X.; Kräft T.; Goycoolea F.M. Chitosan encapsulation modulates the effect of trans-cinnamaldehyde on AHL-regulated quorum sensing activity. *Colloids Surf. B Biointerfaces*, **2018**, in press, Available online 24 May 2018. DOI: doi.org/10.1016/j.colsurfb.2018.05.054.
- Queiroz, C.R.A.A.; Andrade, R.R.; Morais, S.A.L.; Pavani, L.C. Growing *Pereskia aculeata* under intermittent irrigation according to levels of matric potential reduction. *Pesq. Agropec. Trop.*, **2015**, 45(1), 1-8.
- Rapport, E.H.; Drausal, B.S. Edible plants. In: *Encyclopedia of biodiversity*; S. Levi, Ed.; Academic Press:New York; **2001**, Vol. 3, pp. 375-382.
- Rocha, D.R.C.; Pereira Junior, G.A.; Vieira, G.; Pantoja, L.; Santos, A.S.; Pinto, N.A.D. Macarrão adicionado de ora-pro-nobis (*Pereskia aculeata* Miller) desidratado. *Alim. Nut.*, **2008**, 19(4), 459-465.
- Salt, T.A.; Tocker, J.E.; Adler, J.H. Dominance of Δ^5 -sterols in eight species of the Cactaceae. *Phytochem.*, **1987**, 26(3), 731-733.
- Salvi, J.; SS, K. A review: Underutilized wild edible plants as a potential source of alternative nutrition. *Int. J. Bot. Stud.*, **2016**, 1(4), 32-36.
- Santos, I.C.; Pedrosa, M.W.; Carvalho, O.C.; Guimarães, C.D.C.; Silva, L.S. Ora-pro-nóbis: da cerca à mesa. Empresa de Pesquisa Agropecuária de Minas Gerais – EPAMIG, **2012**, 177. Available from: <http://studylibpt.com/doc/3195858/circular-t%C3%A9n%C2%BA177hot->.
- Sartor, C.F.P.; Amaral, V.; Guimarães, H.E.T.; Barros, K.N.; Felipe, D.F.; Cortez, L.E.R.; Veltrini, V.C. Study of the healing action of *Pereskia aculeata* leaves. *Rev. Saúde Pesq.*, **2010**, 3(2), 149-154.

- Sierakowski, M.R.; Corin, P.A.J.; Reicher, F.; Corrêa, J.B.C. Some structural features of a heteropolysaccharide from the leaves of the cactus *Pereskia aculeata*. *Phytochem.*, **1987**, 26(6), 1709-1713.
- Sierakowski, M. R., Gorin, P. A. J., Reicher, F., & Corrêa, J. B. C. (1990). Location of O-acetyl groups in the heteropolysaccharide of the cactus *Pereskia aculeata*. *Carbohydr. Res.*, **1990**, 201(2), 277-284.
- Silva, K.C.G.; Amaral, T.N.; Junqueira, L.A.; Leite, N.O.; Resende, J.V. Adsorption of protein on activated carbon used in the filtration of mucilage derived from *Pereskia aculeata* Miller. *S. Afr. J. Chem. Eng.*, **2017**, 23, 42-49.
- Sim, K.S.; Nurestri, A.M.S.; Sinniah, S.K.; Kim, K.H.; Norhanom, A.W. Acute oral toxicity of *Pereskia bleo* and *Pereskia grandifolia* in mice. *Pharmacogn. Mag.*, **2010**, 6(21), 67-70.
- Sobrinho, S.S.; Costa, L.L.; Gonçalves, C.A.A.; Campagnol, P.C.B. Emulsified cooked sausages enriched with flour from ora-pro-nobis leaves (*Pereskia aculeata* Miller). *Int. Food Res. J.*, **2015**, 22(1), 318-323.
- Sousa, R.M.F; Lira, C.S.; Rodrigues, A.O.; Morais, S.A.L.; Queiroz, C.R.A.A.; Chang, R.; Aquino, F.J.T.; Muñoz, R.A.A.; Oliveira, A. Atividade antioxidante de extratos de folhas de ora-pronóbis (*Pereskia aculeata* Mill.) usando métodos espectrofotométricos e voltamétricos in vitro. *Biosci. J.*, 30(1), 448-457.
- Souza, M.R.M.; Pereira, P.R.G.; Magalhães, I.P.B.; Sedyama, M.A.N.; Vidigal, S.M.; Milagres, C.S.F.; Baracat-Pereira, M.C. Mineral, protein and nitrate contents in leaves of *Pereskia aculeata* subjected to nitrogen fertilization. *Pesq. Agropec. Trop.*, **2016a**, 46(1), 43-50.
- Souza, L.F.; Gasparetto, B.F.; Lopes, R.R.; Barros, I.B.I. Temperature requirements for seed germination of *Pereskia aculeata* and *Pereskia grandifolia*. *J. Therm. Biol.*, **2016b**, 57, 6-10.
- Takeiti, C.Y.; Antônio, G.C.; Motta, E.M.P.; Collares-Queiroz, F.P.; Park, K. J. Nutritive evaluation of a non-conventional leafy vegetable (*Pereskia aculeata* Miller). *Int. J. Food Sci. Nut.*, **2009**, 60(1), 148-160.
- Tofanelli, M.B.D.; Rezende, S.G. Sistemas de condução na produção de folhas de ora-pro-nobis. *Pesq. Agropec. Trop.*, **2011**, 41(3), 466-469.
- Thornfeldt, C. Cosmeceuticals containing herbs: fact, fiction, and future. *Dermatol. Surg.*, **2005**, 31(7 Pt 2), 873-880.
- Uyub, A.M.; Nwachukwu, I.N.; Azlan, A.A.; Fariza, S.S. In-vitro antibacterial activity

and cytotoxicity of selected medicinal plant extracts from Penang Island Malaysia on metronidazole-resistant *Helicobacter pylori* and some path. *Ethnobot. Res. Appl.*, **2010**, 8, 95-106.

Velikov K.P. Colloidal emulsions and particles as micronutrient and nutraceutical delivery systems. In: *Functional Foods - Concept to Product*; M. Saarela, Ed.; Woodhead Publishing; **2011**, 2nd Edition, pp. 319-391. DOI: doi.org/10.1533/9780857095909.3.319

Zem, L.B.; Helm, C.V.; Henriques, G.S.; Cabrini, D.A.; Zuffellato-Ribas, K.C. *Pereskia aculeata*: biological analysis on wistar rats. *Food Sci. Technol.*, **2017**, 37(1), 42-47.

Wahab, S.I.A.; Abdul, A.B.; Mohan, S.M.; Al-Zubairi, A.S.; Elhassan, M.M.; Ibrahim, M.Y. Biological activities of *Pereskia bleo* extracts. *Int. J. Pharm.*, **2009**, 5(1), 71-75.