



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/156004/>

Version: Published Version

Article:

Hemmings, N. and Evans, S. (2020) Unhatched eggs represent the invisible fraction in two wild bird populations. *Biology Letters*, 16 (1). 20190763. ISSN: 1744-9561

<https://doi.org/10.1098/rsbl.2019.0763>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Research



Cite this article: Hemmings N, Evans S. 2020 Unhatched eggs represent the invisible fraction in two wild bird populations. *Biol. Lett.* **16**: 20190763.
<http://dx.doi.org/10.1098/rsbl.2019.0763>

Received: 18 October 2019
Accepted: 2 December 2019

Subject Areas:
ecology, evolution

Keywords:
fertility, hatching failure, missing fraction, lifetime reproductive success, prenatal mortality

Author for correspondence:
Simon Evans
e-mail: s.r.evans@exeter.ac.uk

Population ecology

Unhatched eggs represent the invisible fraction in two wild bird populations

Nicola Hemmings¹ and Simon Evans^{2,3}

¹Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, UK

²Edward Grey Institute, Department of Zoology, University of Oxford, Oxford OX1 3SZ, UK

³Centre for Ecology and Conservation, University of Exeter, Cornwall Campus, Penryn TR10 9FE, UK

NH, 0000-0003-2418-3625; SE, 0000-0001-5812-4039

Prenatal mortality is typically overlooked in population studies, which biases evolutionary inference by confounding selection and inheritance. Birds represent an opportunity to include this ‘invisible fraction’ if each egg contains a zygote, but whether hatching failure is caused by fertilization failure versus prenatal mortality is largely unknown. We quantified fertilization failure rates in two bird species that are popular systems for studying evolutionary dynamics and found that overwhelming majorities (99.9%) of laid eggs were fertilized. These systems thus present opportunities to eliminate the invisible fraction from life-history data.

1. Introduction

Study populations of wild animals offer great insight into the ecological and evolutionary processes operating under natural conditions, based on the ability to observe sampled individuals throughout their lives [1]. However, theoretical biologists have long warned that population sampling should occur at the inter-generational boundary for valid inference [2], since sampling at later ages creates an ‘invisible fraction’ [3], a demographic group composed of individuals that died before the sampling age. This invisible fraction can be very large, potentially representing the majority of a conception cohort. For example, more than three-quarters of human conceptions are naturally aborted [4] and this prenatal mortality is phenotypically non-random [5]. A direct consequence of the invisible fraction is that sampled offspring are more similar to their parents than is the complete conception cohort because sampled offspring have successfully negotiated early-life selection. This similarity will be attributed to inheritance but it actually results from selection. Clearly, then, demographic and evolutionary dynamics will be misrepresented in the presence of an invisible fraction.

As such, individual-level life-history records that incorporate prenatality would contribute significantly to our understanding of the evolutionary dynamics of wild animal populations. However, in viviparous taxa such as mammals, inferring the population size at the prenatal stage is extremely challenging: non-invasive observation can quantify only late-term abortions since less developed prenatals are typically resorbed by their mother [6]. Studying oviparous taxa, such as birds, overcomes this difficulty because zygotes are rapidly and individually externalized in discrete vessels (i.e. eggs). Birds also lay their eggs in predictable locations (i.e. nests), so it is practical to count the egg production of marked individuals throughout their lives. Indeed, there are many long-term population studies of birds for which such data have been routinely recorded, sometimes for decades, but uncertain fertilization success [7] means that each egg may not necessarily contain a zygote, in which case egg counts could not be used to census a conception cohort.

Across a broad diversity of wild bird species, roughly one-tenth of eggs fail to hatch [7–9]. Hatching failure also occurs in commercially important, domesticated birds [10–14], with average rates of 8–15%, even in breeds selected for efficient chick production [15]. Hatching failure in both wild and domestic birds results from either fertilization failure (i.e. the egg formed in the absence of a zygote) or prenatal mortality, and researchers typically rely on visual inspection of eggs' contents to distinguish between these fates (e.g. [16–22]). The ovum is usually fertilized 24 h before oviposition, and cell division begins 6–8 h after fertilization [23,24], so by the time the egg is laid, the blastoderm consists of *ca.* 10 000 cells (measured for domestic fowl: [25]) (see [26] for a passerine comparison). However, the early stages of embryonic development (when most prenatal mortality occurs: [11]) are invisible to the naked eye, particularly once the egg has started to deteriorate [26,27], and microscopic examination requires tissue staining to confidently diagnose fertilization failure [26,28,29]. Thus, a large proportion of expired prelates go unobserved by macroscopic inspection, upwardly biasing estimates of the fertilization failure rate in both wild populations and commercial breeding flocks. Here, we present data on prenatal mortality from study populations of two bird species that are popular and influential systems for studying the ecology and evolution of wild animal populations [30–33], and from which hundreds of unhatched eggs were collected across 3 years to determine fertilization status and—where applicable—age-specific mortality of embryos.

2. Material and methods

(a) Study site and systems

The long-term monitoring of the breeding populations of great tits (*Parus major*) and blue tits (*Cyanistes caeruleus*) in Wytham Woods, Oxfordshire, represents one of the longest-running ecological studies of individually marked animals in the world [1], having started in 1947 [34]. Wytham Woods is a 388-hectare mixed deciduous woodland containing 1207 nest-boxes. Both species readily adopt nest-boxes as nesting sites and an excess of nest-boxes allows almost all breeding attempts to be monitored [35]. As previously described [31], nest-boxes were visited regularly from early April to look for signs of nest building and to count eggs. When hatching was expected, the nest-box was checked more regularly and hatching date estimated based on chicks' weight and appearance.

From 2008 to 2010, all eggs that failed to hatch were collected 14 days after the nest's date of first hatching and stored in a refrigerator for up to one month (in 2010, unhatched eggs were collected from blue tit nests only). It is unlikely that any eggs were removed by parents prior to collection: although mentioned in the literature [36], this behaviour is extremely unusual. Eggs that failed to hatch owing to desertion by the parents (i.e. when parents abandoned the clutch or died during the incubation period) were excluded from the analysis, as were eggs damaged prior to or during collection. Across the study period (blue tits: 2008–2010; great tits: 2008 and 2009), 4.2% (419 of 10 047) of blue tit eggs and 9.9% (673 of 6788) of great tit eggs failed to hatch. Of these, 416 unhatched blue tit eggs (from 254 clutches) and 375 unhatched great tit eggs (from 197 clutches) were examined following the methods described in Birkhead *et al.* [26]. A smaller proportion of unhatched great tit eggs was examined because many more were damaged during collection. However, the occurrence of damage followed no obvious spatial or temporal aggregation,

so examined eggs were considered to be a random subset of those that failed.

(b) Egg examination

Sampled eggs were opened and the contents emptied into phosphate-buffered saline (PBS) solution. If a germinal disc or embryo was observed, it was cleaned in PBS solution and examined under a stereomicroscope to identify the developmental stage at which death occurred, based on a modified version of Hamburger & Hamilton's normal stages of chick development [37], adapted for application to passerines by Hemmings & Birkhead [28]. The majority of embryos found in eggs of both species (blue tit: 392 of 407, 96.3%; great tit: 319 of 370, 86.2%) were 'staged' (assigned to one of the 40 identifiable phases of embryonic development: [28,37]) in this way. Missing staging scores resulted from egg contents being too disintegrated to allow accurate staging (despite some prenatal development being discernible). Developmental stages were used to calculate each prelate's 'age' at death, given known rates of prenatal development in these species [28].

If the egg yolk was disintegrated, the egg's contents were thoroughly examined by stereomicroscope to search for indicative material that might be invisible to the naked eye. If neither a germinal disc nor an embryo was observed, or if development appeared to be at such an early stage that fertility of the egg remained uncertain, then the perivitelline layer of the yolk was stained with Hoechst 33342 fluorescent DNA stain (0.05 mg ml⁻¹), as was any assumed embryonic or germinal disc tissue. Stained tissues were examined under a fluorescence microscope with a BP 340–380 excitation filter, LP 425 suppression filter, dark-field optics and a 20× objective lens, to confirm the presence of (a) nuclei from embryonic tissue, (b) sperm trapped in the perivitelline layer of the ovum and (c) penetration holes made by sperm that had entered the ovum [26]. Fertilization success was determined primarily from criterion (a), with (b) and (c) providing additional evidence: eggs were assumed to be unfertilized if cell nuclei indicative of embryonic tissue could not be found.

Data are archived in the Dryad Digital Repository at <https://doi.org/10.5061/dryad.0d6h6j8> [38].

3. Results

Of all unhatched eggs examined across the 3 years, 2.2% (9 of 416) of blue tit eggs and 1.3% (5 of 375) of great tit eggs were unfertilized. Unfertilized blue tit eggs came primarily from two clutches (three unfertilized eggs in one and four in another, both in 2009), whereas the five unfertilized great tit eggs came from five different clutches. Assuming these rates of fertilization failure are consistent with those of unhatched eggs that were not examined, we estimate that 9 of 10 047 (0.1%) eggs laid by blue tits, and 9 of 6788 (0.1%) eggs laid by great tits over the course of the study were unfertilized. The overwhelming majority of eggs thus contained a zygote that either survived to hatch (95.8% of blue tit eggs; 90.1% of great tit eggs) or died during prenatal development (4.1% of blue tit eggs; 9.8% of great tit eggs). Importantly, of the individuals suffering prenatal mortality, 50.4% (205/407) of blue tits and 31.9% (118/370) of great tits died prior to chick developmental stage 15, which is the earliest stage at which embryo development can be reliably discerned without using the specialized methods we employed [20]. If we had relied on non-microscopic examination of egg contents for diagnosis, we would have found that approximately 52% of unhatched blue tit eggs and 33% of unhatched great tit eggs were unfertilized.

4. Discussion

In two bird species that are widely used study systems for studying the evolutionary ecology of wild animals, we show that the vast majority of unhatched eggs are fertilized, with hatching failure attributable to prenatal mortality. This includes a large share of eggs for both study populations (a slight majority in the case of blue tits) for which macroscopic inspection would conclude fertilization failure was the cause. In reality, a negligible proportion (0.1%) of the eggs laid in our blue tit and great tit populations were unfertilized.

For both species, the majority of prenatal deaths occurred in the first half of the incubation period, consistent with the observation that mortality risk typically declines through each stage of the life cycle [39]. Moreover, approximately one-third of prenatal deaths in great tits and half of those in blue tits occurred before chick developmental stage 15 [37], the earliest stage at which an embryo is observable by alternative methods (i.e. egg candling or macroscopic post-mortem examination; [28]). These frequencies of apparently undeveloped eggs are broadly similar to those reported for other passerine species, suggesting that similar rates of early embryo mortality may be found across songbirds. For example, in a population of Eurasian reed warblers (*Acrocephalus scirpaceus*), 42% of 152 unhatched eggs lacked a visible embryo and fertility status was not determined [20]. Similarly, 25.3% of 387 hihi (*Notiomystis cincta*) eggs were undeveloped and assumed infertile based on macroscopic examination [40].

It is uncertain whether the patterns we report here, in two closely related and ecologically similar species, are more broadly generalizable across bird species. Very few previous studies have accurately distinguished between fertilization failure and prenatal mortality in wild birds, and these are mostly focused on small or threatened populations with higher than average levels of hatching failure, which may have specific reproductive problems [41,42]. Despite this, the results of these studies generally reflect the low rates of infertility we observed here: of 40 unhatched tree sparrow (*Passer montanus*) eggs collected in a single season, all were fertilized [26]; of 120 unhatched wild hihi (*N. cincta*) lacking macroscopic indication of development (previously classified as unfertilized), 88% were fertilized [41]; of 10 undeveloped yellow-shouldered Amazon parrot (*Amazona barbadensis*) eggs, also collected from the wild, all were fertilized [41]; and of 518 wild house sparrow (*Passer domesticus*) eggs,

98.9% were fertilized [43]. These findings, combined with our results, indicate that it is unreliable to assume undeveloped eggs are unfertilized. Therefore, at least in the systems studied so far, the most accurate method of identifying cases of prenatal mortality in the absence of microscopic examination may be to assume that all eggs contained a zygote. An important future objective will be to assess how the incidence of fertilization failure relative to early embryo mortality changes depending on environmental factors, particularly in the presence of environmental pollutants, which have long been linked to reduced fertility [44].

Given that (a) the occurrence of multiple ova per egg is exceedingly rare in birds [45,46], (b) non-surviving individuals remain in the nest as unhatched eggs and (c) post-fledging survival and breeding success of all hatchlings is monitored, our results demonstrate that—at least for two popular avian study species—it is feasible to observe individual survival from almost immediately post-conception in the wild. It is thereby possible to extend empirical consideration of the life histories of wild animals beyond the ‘cradle to grave’ perspective that dominates popular notions of what a lifespan represents. Incorporating prenatality would have an impact within the fields of behavioural ecology, population biology, conservation biology and evolutionary ecology, all disciplines where individual-level life-history data are the basis of empirical analysis.

Ethics. Unhatched eggs were collected during routine fieldwork on the blue tit and great tit populations in Wytham Woods. Failed eggs were taken during chick ringing (14 days post-hatching) to minimize disturbance.

Data accessibility. The data supporting this article are available at the Dryad Digital Repository (<https://doi.org/10.5061/dryad.0d6h6j8>) [38].

Authors' contributions. N.H. carried out all field and laboratory work and revised the manuscript. S.E. conceived the idea for the manuscript and wrote the first draft. Both authors gave final approval for publication and agree to be held accountable for the work performed herein.

Competing interests. We declare we have no competing interests.

Funding. This article was funded by the Natural Environment Research Council (grant no. NE/F008775/1) and the Royal Society (grant no. DH160200) (to N.H.).

Acknowledgements. We thank all those who helped with fieldwork in Wytham Woods, Tim Birkhead for advice during data collection, Barbara Tschirren and two anonymous reviewers for comments on earlier versions of the manuscript. The fieldwork was inspired by Robin McCleery, who suggested a study of hatching failure in these populations.

References

- Clutton-Brock T, Sheldon BC. 2010 Individuals and populations: the role of long-term, individual-based studies of animals in ecology and evolutionary biology. *Trends Ecol. Evol.* **25**, 562–573. (doi:10.1016/j.tree.2010.08.002)
- Arnold SJ, Wade MJ. 1984 On the measurement of natural and sexual selection: applications. *Evolution* **38**, 720–734. (doi:10.1111/j.1558-5646.1984.tb00345.x)
- Grafen A. 1988 On the uses of data on lifetime reproductive success. In *Reproductive success* (ed. TH Clutton-Brock), pp. 454–471. Chicago, IL: University of Chicago Press.
- Roberts CJ, Lowe CR. 1975 Where have all the conceptions gone? *Lancet* **305**, 498–499. (doi:10.1016/S0140-6736(75)92837-8)
- Babler WJ. 1978 Prenatal selection and dermatoglyphic patterns. *Am. J. Phys. Anthropol.* **48**, 21–27. (doi:10.1002/ajpa.1330480105)
- Frazer JFD. 1955 Foetal death in the rat. *J. Embryol. Exp. Morphol.* **3**, 13–29.
- Koenig WD. 1982 Ecological and social factors affecting hatchability of eggs. *Auk* **99**, 526–536.
- Morrow EH, Arnqvist G, Pitcher TE. 2002 The evolution of infertility: does hatching rate in birds coevolve with female polyandry? *J. Evol. Biol.* **15**, 702–709. (doi:10.1046/j.1420-9101.2002.00445.x)
- Spottiswoode C, Møller AP. 2004 Genetic similarity and hatching success in birds. *Proc. R. Soc. Lond. B* **271**, 267–272. (doi:10.1098/rspb.2003.2605).
- Christensen VL. 2001 Factors associated with early embryonic mortality. *World's Poultr. Sci. J.* **57**, 359–372. (doi:10.1079/WPS20010025)
- Romanoff AL, Romanoff AJ. 1972 *Pathogenesis of the avian embryo: an analysis of causes of malformations and prenatal death*. New York, NY: Wiley-Interscience.

12. Lerner SP, French N, McIntyre D, Baxter-Jones C. 1993 Age-related changes in egg production, fertility, embryonic mortality, and hatchability in commercial turkey flocks. *Poult. Sci.* **72**, 1025–1039. (doi:10.3382/ps.0721025)
13. Meijerhof R. 1992 Pre-incubation holding of hatching eggs. *World's Poult. Sci. J.* **48**, 57–68. (doi:10.1079/WPS19920006)
14. Scott TA, Mackenzie CJ. 1993 Incidence and classification of early embryonic mortality in broiler breeder chickens. *Br. Poult. Sci.* **34**, 459–470. (doi:10.1080/00071669308417601)
15. Beaumont C, Millet N, Le Bihan-Duval E, Kipi A, Dupuy V. 1997 Genetic parameters of survival to the different stages of embryonic death in laying hens. *Poult. Sci.* **76**, 1193–1196. (doi:10.1093/ps/76.9.1193)
16. Cordero PJ, Aparicio JM, Veiga JP. 2004 Parental genetic characteristics and hatching success in the spotless starling, *Sturnus unicolor*. *Anim. Behav.* **67**, 637–642. (doi:10.1016/j.anbehav.2003.06.005)
17. Rothstein SI. 1973 Variation in the incidence of hatching failure in the cedar waxwing and other species. *Condor* **75**, 164–169. (doi:10.2307/1365863)
18. Hooson S, Jamieson IG. 2004 Variation in breeding success among reintroduced island populations of South Island saddlebacks *Philesturnus carunculatus carunculatus*. *Ibis* **146**, 417–426. (doi:10.1111/j.1474-919X.2004.00275.x)
19. Potti J, Merino S. 1996 Causes of hatching failure in the pied flycatcher. *Condor* **98**, 328–336. (doi:10.2307/1369151)
20. Orłowski G, Hałupka L, Pokorny P, Klimczuk E, Sztwiertnia H, Dobicki W. 2016 The effect of embryonic development on metal and calcium content in eggs and eggshells in a small passerine. *Ibis* **158**, 144–154. (doi:10.1111/ibi.12327)
21. Seel DC. 1968 Clutch-size, incubation and hatching success in the house sparrow and tree sparrow *Passer* spp. at Oxford. *Ibis* **110**, 270–282. (doi:10.1111/j.1474-919X.1968.tb00038.x)
22. Wetton JH, Parkin DT. 1991 An association between fertility and cuckoldry in the house sparrow, *Passer domesticus*. *Proc. R. Soc. Lond. B* **245**, 227–233. (doi:10.1098/rspb.1991.0114)
23. Eyal-Giladi H, Kochav S. 1976 From cleavage to primitive streak formation: a complementary normal table and a new look at the first stages of the development of the chick: I. General morphology. *Dev. Biol.* **49**, 321–337. (doi:10.1016/0012-1606(76)90178-0)
24. Watt JM, Pettite JN, Etches RJ. 1993 Early development of the chick embryo. *J. Morphol.* **215**, 165–182. (doi:10.1002/jmor.1052150205)
25. Kochav S, Ginsburg M, Eyal-Giladi H. 1980 From cleavage to primitive streak formation: a complementary normal table and a new look at the first stages of the development of the chick: II. Microscopic anatomy and cell population dynamics. *Dev. Biol.* **79**, 296–308. (doi:10.1016/0012-1606(80)90117-7)
26. Birkhead TR, Hall J, Schut E, Hemmings N. 2008 Unhatched eggs: methods for discriminating between infertility and early embryo mortality. *Ibis* **150**, 508–517. (doi:10.1111/j.1474-919X.2008.00813.x)
27. Croyle KE, Durrant BS, Jensen T. 2015 Detection of oocyte perivitelline membrane-bound sperm: a tool for avian collection management. *Conserv. Physiol.* **3**, cou060. (doi:10.1093/conphys/cou060)
28. Hemmings N, Birkhead TR. 2016 Consistency of passerine embryo development and the use of embryonic staging in studies of hatching failure. *Ibis* **158**, 43–50. (doi:10.1111/ibi.12336)
29. Kosin IL. 1944 Macro- and microscopic methods of detecting fertility in unincubated hen's eggs. *Poult. Sci.* **23**, 266–269. (doi:10.3382/ps.0230266)
30. Lack D. 1966 *Population studies of birds*. Oxford, UK: Oxford University Press.
31. Perrins CM. 1965 Population fluctuations and clutch size in the great tit, *Parus major* L. *J. Anim. Ecol.* **34**, 601–647. (doi:10.2307/2453)
32. Charmantier A, McCleery RH, Cole LR, Perrins C, Kruuk LEB, Sheldon BC. 2008 Adaptive phenotypic plasticity in response to climate change in a wild bird population. *Science* **320**, 800–803. (doi:10.1126/science.1157174)
33. Kempenaers B, Verheyen GR, den Broeck MV, Burke T, Broeckhoven CV, Dhondt A. 1992 Extra-pair paternity results from female preference for high-quality males in the blue tit. *Nature* **357**, 494–496. (doi:10.1038/357494a0)
34. Lack D. 1964 A long-term study of the great tit (*Parus major*). *J. Anim. Ecol.* **33**, 159–173. (doi:10.2307/2437)
35. Greenwood PJ, Harvey PH, Perrins CM. 1979 The role of dispersal in the great tit (*Parus major*): the causes, consequences and heritability of natal dispersal. *J. Anim. Ecol.* **48**, 123–142. (doi:10.2307/4105)
36. Perrins C. 1979 *British tits*. London, UK: Collins.
37. Hamburger V, Hamilton HL. 1951 A series of normal stages in the development of the chick embryo. *J. Morphol.* **88**, 49–92. (doi:10.1002/jmor.1050880104)
38. Hemmings N, Evans S. 2019 Data from: Unhatched eggs represent the invisible fraction in two wild bird populations. Dryad Digital Repository. (doi:10.5061/dryad.0d6h6j8)
39. Low M, Pärt T. 2009 Patterns of mortality for each life-history stage in a population of the endangered New Zealand stitchbird. *J. Anim. Ecol.* **78**, 761–771. (doi:10.1111/j.1365-2656.2009.01543.x)
40. Brekke P, Bennett PM, Wang J, Pettorelli N, Ewen JG. 2010 Sensitive males: inbreeding depression in an endangered bird. *Proc. R. Soc. B* **277**, 3677–3684. (doi:10.1098/rspb.2010.1144)
41. Hemmings N, West M, Birkhead TR. 2012 Causes of hatching failure in endangered birds. *Biol. Lett.* **8**, 964–967. (doi:10.1098/rsbl.2012.0655)
42. Briskie JV, Mackintosh M. 2004 Hatching failure increases with severity of population bottlenecks in birds. *Proc. Natl Acad. Sci. USA* **101**, 558–561. (doi:10.1073/pnas.0305103101)
43. Birkhead TR, Veiga JP, Fletcher F. 1995 Sperm competition and unhatched eggs in the house sparrow. *J. Avian Biol.* **26**, 343–345. (doi:10.2307/3677051)
44. Fry DM. 1995 Reproductive effects in birds exposed to pesticides and industrial chemicals. *Environ. Health Perspect.* **103**, 165–171. (doi:10.1289/ehp.95103s7165)
45. Griffith SC, Stewart R. 1998 Genetic confirmation of non-identical embryonic twins in the house sparrow *Passer domesticus*. *J. Avian Biol.* **29**, 207–208. (doi:10.2307/3677202)
46. Betuel AM, Tuttle EM, Gosner RA. 2015 Genetic verification of dizygotic twin embryos in the white-throated sparrow (*Zonotrichia albicollis*). *Wilson J. Ornithol.* **127**, 498–502. (doi:10.1676/14-090.1)