

This is a repository copy of *Are intestinal parasites associated with obesity in Mexican children and adolescents?*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/156105/>

Version: Accepted Version

---

**Article:**

Zavala, Gerardo A [orcid.org/0000-0002-9825-8725](https://orcid.org/0000-0002-9825-8725), Doak, Colleen M, Portrait, France et al. (6 more authors) (2019) Are intestinal parasites associated with obesity in Mexican children and adolescents? *Parasitology International*. pp. 126-131. ISSN: 1873-0329

<https://doi.org/10.1016/j.parint.2019.04.002>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

**Are intestinal parasitic infections associations with obesity in Mexican children and adolescents ?**

Gerardo A. Zavala<sup>a,b,\*</sup>, Colleen M. Doak<sup>a</sup>, France Portrait<sup>a</sup>, Jaap C. Seidell<sup>a</sup>, Olga P. García<sup>b</sup>, Jorge L. Rosado<sup>b</sup>, Tatiana Jarquín<sup>b</sup>, Katja Polman<sup>a,c</sup>, Maiza Campos-Ponce<sup>a</sup>

<sup>a</sup> Faculty of Earth and Life Sciences, VU Amsterdam University. De Boelelaan 1105, 1081 HV, Amsterdam, The Netherlands.

<sup>b</sup> Facultad de Ciencias Naturales, Universidad Autónoma de Querétaro. Avenida de las Ciencias s/n, Juriquilla, 76230, Queretaro, Mexico

<sup>c</sup> Department of Biomedical Sciences, Unit of Medical Helminthology, Institute of Tropical Medicine. Kronenburgstraat 43, 2000 Antwerpen, Belgium.

Running title: Parasitic infection and obesity

\*Corresponding author: Gerardo A. Zavala Gomez

E-mail: [g Zavala@gmail.com](mailto:g Zavala@gmail.com) [g.a.zavalagomez@vu.nl](mailto:g.a.zavalagomez@vu.nl)

De Boelelaan 1105, 1081 HV Amsterdam W&N building room 526

**Running title: are intestinal parasites associated with obesity ?**

## 1. INTRODUCTION

Obesity and associated morbidities are rapidly growing public health challenges that many low and middle income countries are facing [1, 2]. Often in these countries, infections are common as well, especially in children [3, 4]. Recent studies have shown that certain viral and bacterial infections are associated with obesity (described as “infectobesity”) [5-8]. However, studies on the association between parasite infections and obesity are still scarce [9-11].

We hypothesize that intestinal parasites may be associated with obesity by two plausible mechanisms. The first possibility is that both obesity and parasitic infections are positively associated with poverty, as reported previously in different low and middle income countries [12, 13]. However, a second alternative may be that intestinal parasites have an effect on the metabolism by the alteration of the gut microbiome composition.

It is assumed that “stress factors” or “insults” such as infectious diseases and under-nutrition presented during a “critical window” of development (i.e. childhood or puberty) can lead to changes in the gut microbiome composition [14-16]. These changes may be reflected as alterations in metabolism and may lead to fat deposition over time [17-20]. For instance, modified gut microbiota can increase caloric uptake from the diet and can modulate host genes that affect energy deposition in adipocytes and thereby increase the risk of diet-induced obesity [19].

According to the last health and nutrition survey of Mexico (ENSANUT 2012) Mexico has a combined prevalence of overweight and obesity of approximately 70% in adults and 30% in children [21, 22]. Approximately 50% of the total population is infected with one or more species of intestinal parasites [23, 24] with *A. lumbricoides* being the most common intestinal helminth

infection, and *E. coli* the most prevalent intestinal protozoan infection [11, 25]. In a previous study, we found intestinal protozoa infection, particularly *Entamoeba coli* (*E. coli*) infection was associated with a higher percentage of body fat and food intake, while *Ascaris lumbricoides* (*A. lumbricoides*) infection was associated with a lower food intake in children [11]. The objective of this ecological study is to test if children living in states with high (reported) incidence of intestinal parasitic infection have higher BMI<sub>z</sub> and higher BMI<sub>z</sub> later in life.

## **2. Materials and methods**

### *2.1 Datasets:*

We used individual and state-wide data (32 states) collected by three different federal organizations of Mexico. Individual level data on height, weight and age was obtained from the 2012 National Health and Nutrition Survey (ENSANUT 2012). This survey of over 50,000 households is representative for the Mexican population at national and state level. The data is available at <http://ensanut.insp.mx> and the methodology is described elsewhere [26]. Statewide data on intestinal helminths and protozoa in 2012, 2006 and 2000 was obtained from the National System for Epidemiological Surveillance (SINAVE) [27]. The data is available at: <http://www.epidemiologia.salud.gob.mx/anuario/html/anuarios.html> Finally we used state-wide data on demographic and socioeconomic variables from the National Institute of Statistics and Geography (INEGI) 2012 report, available at <http://www.inegi.org.mx/> [28].

### *2.2 Dependent Variable: BMI for age z-score*

Using data from ENSANUT 2012, we calculated the BMI for age z-score (BMI<sub>z</sub>) for all individuals. BMI<sub>z</sub> for individuals above 19 years was calculated using the last available point of

the WHO grow charts for children (228 months). The calculations were made using the World Health Organization (WHO) SPSS anthroplus macro for children 5-19 years (WHO, Geneva, Switzerland). The macro is publicly available at <http://www.who.int/childgrowth/software/en/>.

### *2.3 Independent variables: infection with intestinal parasites*

Information on intestinal helminths and protozoa was extracted from SINAVE, which is the only publicly available data source on intestinal parasite infection across the country. In this system, data is reported annually as the incidence per 100 000 person-years in different age groups (i.e. less than 1y, 1-5y, 6-10y, 11-19y) following the same procedures in each of the 32 states of Mexico [27]. *A. lumbricoides* infection, the most common helminth infection in Mexico, is reported as the incidence of *A. lumbricoides* infection. For protozoa infection the only available data was on “intestinal protozoa infection”: that is calculated by SINAVE as the grouped incidence of *Balantidium coli*, *Cryptosporidium* and unspecified intestinal protozoa infection.

We used the SINAVE incidence data as a proxy for the probability of infection, in function of the state and the age of the subject at a particular time point. For instance, the probability of infection in 2012 of an individual of 15 years old was approximated by the incidence of the intestinal parasite infection in his/her state of residence in 2012 for his/her corresponding age group. In addition, the same individual’s probability of infection in 2006 and in 2000, was determined by the incidence of each intestinal parasitic infection in his/her state and for his/her age group at that given year.

## 2.4 Additional covariates

We included both individual-level covariates from the ENSANUT 2012 survey and state-level covariates from the INEGI survey in 2012. The individual-level covariates used in the analysis were: sex (male/female), age (years), place of residence (rural/urban) and “marginality” (marginalized/not-marginalized). Marginality indicates whether a child is from a marginalized socioeconomic status or not based on indicators such as parents education level, access to sanitation facilities, access to drinking water, income and population size of the community, as described in detail elsewhere [26]. State-level covariates were: population with health coverage (%), education level of adults (mean number of years in school), households without sanitation facilities (%) and the poverty rate (population living in poverty). The poverty rate was determined by the “poverty index”, which indicates whether a household is poor or not. In addition to income, the poverty index also takes into consideration access to healthcare, social security, material of the roof, floor and walls of the household, access to basic services and food, according to standard procedures which are described elsewhere [29].

## 2.5 Statistical analysis

As shown in figure 1 the population was stratified into three age groups depending on the age of individuals in 2000, 2006 and 2012. For the cross sectional analysis on the association between infection and obesity in 2012 we extracted data on individuals aged 1-5y (n=8,927), 6-10y (n=16,347) and 11-19y (n=13,992) in 2012. For the analysis on the association between infection in 2006 and obesity in 2012 we selected those individuals from the ENSANUT survey of 2012 who were aged 1-5y (n=9,523), 6-10y (n=13,025), and 11-19y (n=7,845) in 2006. Likewise

for the analysis concerning infection in 2000 we selected those individuals from the ENSANUT survey of 2012 who were aged 1-5y (n=6,625), 6-10y (n=7,580), and 11-19y (n=5,623) in 2000.

Linear regression models were used to determine the association between the probability of infection (2000, 2006 and 2012) with BMIz or in 2012. Associations were estimated for *A. lumbricoides* and intestinal protozoa for each of the three age groups separately. In order to facilitate interpretation of the results, the incidence of infection (proximate probability of infection) for the regression analysis was transformed from new cases in 100 000 person-years to new cases per 100 person-years. Findings below p value of 0.05 were considered significant.

Poverty rate and sex were explored as a possible effect modifiers [30]. For this purpose we performed the same analyses as described above, including an interaction term between sex or poverty rate and the probability of infection in each model. Models with statistically significant interaction terms ( $p < 0.05$ ) were stratified in two groups, one above and one below the median of poverty rate.

In order to have a visual overview of the combined prevalence of overweight and obesity and the incidence of each parasitic infection in 2012, we mapped each variable stratified in quintiles. The unit of mapping was “the state” the largest administrative unit of Mexico, and the maps were generated using Arc GIS V10.1 (Redlands, CA).

### **3. Results**

In 2012, the combined prevalence of overweight and obesity in Mexico was 10% for children aged 1-5y, 35% for children aged 6-10y, 36% for the age group of 11-19y (Figure 2). In total, 47% percent of the population lived in poverty. The health coverage was 62% and the rate

of households without sanitation facilities was 12% (Table 1). The incidence of *A. lumbricoides* and intestinal protozoan infection decreased from 2000 to 2006 and from 2006 to 2012 for all age groups (Table 1).

### 3.1 *Ascaris lumbricoides*

Table 2 shows a positive association between the probability of *A. lumbricoides* infection in 2000 and 2006 with BMIz in 2012. In the adjusted model, an increase of 1% in the probability of infection in 2006 was associated with an increase of 0.13 in the BMIz in 2012 for age group 1-5y, 0.27 for age group 6-10y and 0.50 in BMIz for age group 11-19y. Furthermore, an increase of 1% in the probability of infection with *A. lumbricoides* in 2000 was associated with an increase of 0.10 in the BMIz in 2012 for age group 1-5y, 0.11 for age group 6-10y, and 0.25 for the 11-19y age group. In contrast table 3 shows that a higher probability of being infected with *A. lumbricoides* in 2012 was associated with a decrease of 0.32 in the BMIz for age group 1-5y and a decrease of 0.21 for age group 6-10y.

Table 4 shows the results stratified by poverty rate. In the states with low poverty rates in 2012, *A. lumbricoides* infection was associated with an increased BMIz in 2012 in all age groups. In states with high poverty rates *A. lumbricoides* infection differed between age strata; the probability of infection with *A. lumbricoides* was associated with a lower BMIz for age group 1-5y, no association in BMIz for age group 6-10y and a higher BMIz for age group 11-19y. Neither sex nor poverty rate were modifiers for the associations between the probability of *A. lumbricoides* infection in 2006 or 2000 with BMIz in 2012 for any of the age groups. Therefore no stratified analysis were performed for these years.



### 3.2 Intestinal Protozoa

Table 2 shows the associations between the probability of infection with intestinal protozoa in 2000 and 2006 and BMIz in 2012 across the three studied age groups. The probability of infection with protozoa in 2006 was associated with a higher BMIz in 2012 in the 6-10y and 11-19y age groups. An increase of 1% in the probability of infection in 2000 was associated with an increase in the BMIz in 2012 of 0.47 for age group 1-5y, 0.61 for age group 6-10y and 0.99 for age group 11-19y.

Table 3 shows the associations between the proximate probability of intestinal protozoan infection in 2012 and BMIz in 2012 for every age group. In the adjusted model an increase of 1% in the probability of protozoan infection was associated with an increase of 0.6 in the BMIz for age group 6-10y and an increase of 0.9 for age group 11-19y

Neither sex nor poverty rate were effect modifiers for the associations between the probability of protozoan infection in 2012, 2006 or 2000 with BMIz in 2012 for any of the age groups. Therefore no stratified analysis were performed.

## 4. Discussion

Our results indicate that children with a higher probability of *A. lumbricoides* or intestinal protozoan infection (*Balantidium coli*, *Cryptosporidium*, *unspecified intestinal protozoa infection*) are more likely to have a higher BMIz in the same year, 6, and 12 years later in life. This finding is consistent with other studies showing early child “insults” including infections to be associated

to later overweight and obesity [31-33]. This finding could be related to changes in the gut microbiota and inflammatory reactions due to parasitic infection that may lead to changes in appetite, food intake and thereby BMIz [34-36]. In line with this hypothesis, we recently found that *E. coli* infection was associated with a higher percentage of body fat and food intake in children [11, 37]. Similarly Schilder et al., in a firefly model, observed that an intestinal protozoa common in insects caused fat deposition in the thorax which is comparable to obesity in mammals [9, 10]. Longitudinal studies are needed to assess the temporal association of intestinal parasites on obesity over time.

While a higher probability of intestinal protozoan infection in 2012 was associated with a higher BMIz in the same year, we found the opposite for *A. lumbricoides* infection. If the associations would have been explained purely by poverty, the same trend and direction on the associations would have been observed for both parasitic infection incidence, which was not the case either in the crude or adjusted model. We also found an association between BMIz and *A. lumbricoides* for the age groups 1-5y and 6-10y, and not for the oldest age group (11-19y). The difference between age groups might be explained by the fact that *A. lumbricoides* infection-related symptoms are more common in younger children [38]. Children infected with *A. lumbricoides* may experience abdominal pain, nausea and discomfort, which may lead to a lower food intake and therefore lower BMIz [39]. The results of our previous study in Mexican schoolchildren supports this hypothesis, as we found a negative association between *A. lumbricoides* infection and food intake[37].

We found opposite associations between BMIz and *A. lumbricoides* incidence in states with high and low poverty rates. These differences may be explained with previous studies, as

shown in a review by Guerrant et. al [33] in which children living in poverty were more likely to be malnourished, but also more likely to have stronger symptoms when infected.

Our findings should be interpreted in the context of this being an ecological analysis and not an estimate on any causal effect of parasitic infection on obesity at individual level (ecological fallacy). However, we intended to minimize this issue using individual level data on BMIz and specific covariates. Although we adjusted for potential confounders, we cannot control for unknown or unmeasured factors such as food availability, diet, and physical activity, therefore the outcomes of this study should be interpreted with caution. In addition, it was not possible to take changes in the state of residence over the years in consideration, but according to INEGI, migration between states was relatively low. In the year 2000: 3,584 957 persons migrated between states, representing 3.6% of the population and in 2006: 2,406 454 persons migrated, corresponding to 2.3% of the population at that time [40]. We used incidence data of the studied parasites as a measure for the probability of infection [41], and the true prevalence of parasitic infection is most likely underestimated. A major strength of our study is that ENSANUT and INEGI surveys are representative of the Mexican population at national and state level. In addition the parasite infection data of the SINAVE is collected following the same procedures nationwide and is therefore a good measuring tool for comparison purposes.

## **5. Conclusions**

Our results suggest that children living in states with a higher probability of infection with intestinal parasites have a higher BMIz later in life. The association between current intestinal

parasite infection and BMI is less straightforward, and seems to be opposite for *A. lumbricoides* and intestinal protozoa. Further research is needed to confirm these ecological associations and study possible mechanisms underlying the short-term and long-term consequences of intestinal parasite infections on health. These findings may have important implications for Mexico, given the context of a high prevalence of parasitic infection and obesity.

## Acknowledgements

This study has been partially funded by Consejo Nacional de Ciencia y Tecnología (CONACyT) México, that provided the PhD grant 218666

## Disclosure statement

the authors report no conflicts of interest


## References

- [1] T. Kelly, W. Yang, C.-S. Chen, K. Reynolds, J. He, Global burden of obesity in 2005 and projections to 2030, *International journal of obesity* 32(9) (2008) 1431-1437.
- [2] A. Must, J. Spadano, E.H. Coakley, A.E. Field, G. Colditz, W.H. Dietz, The disease burden associated with overweight and obesity, *Jama* 282(16) (1999) 1523-1529.
- [3] C.L.F. Walker, J. Perin, M.J. Aryee, C. Boschi-Pinto, R.E. Black, Diarrhea incidence in low-and middle-income countries in 1990 and 2010: a systematic review, *BMC public health* 12(1) (2012) 1.
- [4] K.E. Jones, N.G. Patel, M.A. Levy, A. Storeygard, D. Balk, J.L. Gittleman, P. Daszak, Global trends in emerging infectious diseases, *Nature* 451(7181) (2008) 990-993.
- [5] R.J. Schilder, J.H. Marden, Metabolic syndrome and obesity in an insect, *Proceedings of the National Academy of Sciences of the United States of America* 103(49) (2006) 18805-9.
- [6] Z. Yang, V. Grinchuk, A. Smith, B. Qin, J.A. Bohl, R. Sun, L. Notari, Z. Zhang, H. Sesaki, J.F. Urban, Jr., T. Shea-Donohue, A. Zhao, Parasitic nematode-induced modulation of body weight and associated metabolic dysfunction in mouse models of obesity, *Infection and immunity* 81(6) (2013) 1905-14.
- [7] R.L. Atkinson, Could viruses contribute to the worldwide epidemic of obesity?, *International journal of pediatric obesity : IJPO : an official journal of the International Association for the Study of Obesity* 3 Suppl 1 (2008) 37-43.
- [8] N.V. Dhurandhar, A framework for identification of infections that contribute to human obesity, *The Lancet infectious diseases* 11(12) (2011) 963-969.


- [9] R.J. Schilder, J.H. Marden, Metabolic syndrome in insects triggered by gut microbes, *Journal of diabetes science and technology* 1(5) (2007) 794-6.
- [10] R.J. Schilder, J.H. Marden, Metabolic syndrome and obesity in an insect, *Proceedings of the National Academy of Sciences* 103(49) (2006) 18805-18809.
- [11] G.A. Zavala, O.P. Garcia, M. Campos-Ponce, D. Ronquillo, M.C. Caamano, C.M. Doak, J.L. Rosado, Children with moderate-high infection with *Entamoeba coli* have higher percentage of body and abdominal fat than non-infected children, *Pediatric obesity* (2015).
- [12] A.M. Prentice, The emerging epidemic of obesity in developing countries, *International journal of epidemiology* 35(1) (2006) 93-99.
- [13] J. Nematian, E. Nematian, A. Gholamrezanezhad, A.A. Asgari, Prevalence of intestinal parasitic infections and their relation with socio-economic factors and hygienic habits in Tehran primary school students, *Acta tropica* 92(3) (2004) 179-186.
- [14] M.A. Hanson, P.D. Gluckman, Developmental origins of health and disease--global public health implications, *Best practice & research. Clinical obstetrics & gynaecology* 29(1) (2015) 24-31.
- [15] N. Cameron, E.W. Demerath, Critical periods in human growth and their relationship to diseases of aging, *American journal of physical anthropology Suppl* 35 (2002) 159-84.
- [16] R.S. Liu, D.P. Burgner, M.A. Sabin, C.G. Magnussen, M. Cheung, N. Hutri-Kahonen, M. Kahonen, T. Lehtimäki, E. Jokinen, T. Laitinen, L. Taittonen, T. Dwyer, J.S. Viikari, M. Kivimäki, O.T. Raitakari, M. Juonala, Childhood Infections, Socioeconomic Status, and Adult Cardiometabolic Risk, *Pediatrics* 137(6) (2016).
- [17] D.J. Barker, Developmental origins of adult health and disease, *Journal of epidemiology and community health* 58(2) (2004) 114-5.
- [18] P. Klimek, M. Leitner, A. Kautzky-Willer, S. Thurner, Effect of fetal and infant malnutrition on metabolism in older age, *Gerontology* 60(6) (2014) 502-7.
- [19] V. Gangarapu, K. Yildiz, A.T. Ince, B. Baysal, Role of gut microbiota: Obesity and NAFLD, *The Turkish journal of gastroenterology : the official journal of Turkish Society of Gastroenterology* 25(2) (2014) 133-140.
- [20] M. Sanchez, S. Panahi, A. Tremblay, Childhood Obesity: A Role for Gut Microbiota?, *International journal of environmental research and public health* 12(1) (2014) 162-175.
- [21] G. Olaiz-Fernandez, J. Rivera-Dommarco, T. Shamah-Levy, R. Rojas, S. Villalpando-Hernández, M. Hernández-Ávila, J. Sepúlveda-Amor, Encuesta Nacional de Salud y Nutrición in: I.N.d.S. Publica (Ed.) Cuernavaca, Mexico, 2006.
- [22] T.R. McCarty, J.A. Turkeltaub, P.J. Hotez, Global progress towards eliminating gastrointestinal helminth infections, *Current opinion in gastroenterology* 30(1) (2014) 18-24.
- [23] E. Morales-Espinoza, J. Sanchez.Perez, M. Garcia-Gil, G. Vargas-Morales, J. MEndez.Sanchez, M. Perez.Ramirez, Intestinal parasites in children in highly deprived areas in the border region of Chiapas, Mexico, *salud publica de mexico* 45(5) (2003) 379-289.
- [24] J. Gutierrez-Jimenez, M.G. Torres-Sanchez, L.P. Fajardo-Martinez, M.A. Schlie-Guzman, L.M. Luna-Cazares, A.R. Gonzalez-Esquincá, S. Guerrero-Fuentes, J.E. Vidal, Malnutrition and the presence of intestinal parasites in children from the poorest municipalities of Mexico, *Journal of infection in developing countries* 7(10) (2013) 741-7.
- [25] L. Quihui, M.E. Valencia, D.W. Crompton, S. Phillips, P. Hagan, G. Morales, S.P. Díaz-Camacho, Role of the employment status and education of mothers in the prevalence of intestinal parasitic infections in Mexican rural schoolchildren, *BMC public health* 6(1) (2006) 225.
- [26] J. Gutierrez, J. Rivera-Dommarco, T. Shamah-Levy, S. Villalpando-Hernandez, A. Franco, L. Cuevas-Nasu, M. Romero-Martínez, M. Hernández-Ávila, Encuesta Nacional de Salud y Nutrición 2012. Resultados Nacionales, Instituto Nacional de Salud Pública (MX), Cuernavaca, México, 2012.

- [27] R. Tapia-Conyer, P. Kuri-Morales, L. Gonzalez-Urban, E. Sarti, Evaluation and reform of Mexican National Epidemiological Surveillance System, *American journal of public health* 91(11) (2001) 1758-60.
- [28] g.e.i. Instituto nacional de estadística, Anuario de estadísticas por entidad federativa, Instituto nacional de estadística, geografía e informática 2012.
- [29] Metodología para la medición multidimensional de la pobreza en México, CONEVAL, México, DF, 2014.
- [30] R.L. Guerrant, M.D. DeBoer, S.R. Moore, R.J. Scharf, A.A. Lima, The impoverished gut—a triple burden of diarrhoea, stunting and chronic disease, *Nature Reviews Gastroenterology and Hepatology* 10(4) (2013) 220-229.
- [31] C. Pico, M. Palou, T. Priego, J. Sanchez, A. Palou, Metabolic programming of obesity by energy restriction during the perinatal period: different outcomes depending on gender and period, type and severity of restriction, *Frontiers in physiology* 3 (2012) 436.
- [32] R.M. Anguita, D.M. Sigulem, A.L. Sawaya, Intrauterine food restriction is associated with obesity in young rats, *The Journal of nutrition* 123(8) (1993) 1421-8.
- [33] R.L. Guerrant, M.D. DeBoer, S.R. Moore, R.J. Scharf, A.A. Lima, The impoverished gut--a triple burden of diarrhoea, stunting and chronic disease, *Nature reviews. Gastroenterology & hepatology* 10(4) (2013) 220-9.
- [34] D. Stark, S. Van Hal, D. Marriott, J. Ellis, J. Harkness, Irritable bowel syndrome: a review on the role of intestinal protozoa and the importance of their detection and diagnosis, *International journal for parasitology* 37(1) (2007) 11-20.
- [35] M. Kalliomäki, E. Isolauri, Role of intestinal flora in the development of allergy, *Current opinion in allergy and clinical immunology* 3(1) (2003) 15-20.
- [36] F. Bäckhed, H. Ding, T. Wang, L.V. Hooper, G.Y. Koh, A. Nagy, C.F. Semenkovich, J.I. Gordon, The gut microbiota as an environmental factor that regulates fat storage, *Proceedings of the National Academy of Sciences of the United States of America* 101(44) (2004) 15718-15723.
- [37] G.A. Zavala, J.L. Rosado, C.M. Doak, M. del Carmen Caamaño, M. Campos-Ponce, D. Ronquillo, K. Polman, O.P. García, Energy and food intake are associated with specific intestinal parasitic infections in children of rural Mexico, *Parasitology international* 66(6) (2017) 831-836.
- [38] P. O'Lorcain, C.V. Holland, The public health importance of *Ascaris lumbricoides*, *Parasitology* 121 Suppl (2000) S51-71.
- [39] J.C. Dunn, H.C. Turner, A. Tun, R.M. Anderson, Epidemiological surveys of, and research on, soil-transmitted helminths in Southeast Asia: a systematic review, *Parasites & vectors* 9(1) (2016) 31.
- [40] A. Ahmed, H.M. Al-Mekhlafi, S.H. Choy, I. Ithoi, A.H. Al-Adhroey, A.M. Abdulsalam, J. Surin, The burden of moderate-to-heavy soil-transmitted helminth infections among rural malaysian aborigines: an urgent need for an integrated control programme, *Parasites & vectors* 4 (2011) 242.
- [41] B. Speich, H. Marti, S.M. Ame, S.M. Ali, Bogoch, II, J. Utzinger, M. Albonico, J. Keiser, Prevalence of intestinal protozoa infection among school-aged children on Pemba Island, Tanzania, and effect of single-dose albendazole, nitazoxanide and albendazole-nitazoxanide, *Parasites & vectors* 6 (2013) 3.

328 **Figure1.** Diagram of the study design



**Figure 2.** Map of combined prevalence of overweight and obesity and the incidence of *Ascaris lumbricoides* and protozoa in Mexico, 2012





346

347

**Table 1.** General Characteristics of the population

		Mean	S.D.	Minimum	Maximum	Source
<i>A. lumbricoides</i> 1 to 5 years						
Incidence in 2012	94	± 129.3	3	625	SINAVE 2012	
Incidence in 2006	404	± 461.1	5	1925	SINAVE 2006	
Incidence in 2000	1177	± 1151.2	99	4559	SINAVE 2000	
<i>A. lumbricoides</i> 6 to 10 years						
Incidence in 2012	67	± 108.2	0	526	SINAVE 2012	
Incidence in 2006	212	± 301.8	3	1822	SINAVE 2006	
Incidence in 2000	907	± 999.8	45	3990	SINAVE 2000	
<i>A. lumbricoides</i> 11 to 18 years						
Incidence in 2012	27	± 42.7	0	219	SINAVE 2012	
Incidence in 2006	90	± 140.0	1	757	SINAVE 2006	
Incidence in 2000	367	± 421.5	18	1711	SINAVE 2000	
Protozoa 1 to 5 years						
Incidence in 2012	93	± 104.0	2	545	SINAVE 2012	
Incidence in 2006	239	± 275.2	9	1498	SINAVE 2006	
Incidence in 2000	351	± 215.5	73	905	SINAVE 2000	
Protozoa 6 to 10 years						
Incidence in 2012	78	± 93.7	1	568	SINAVE 2012	
Incidence in 2006	150	± 200.7	4	1590	SINAVE 2006	
Incidence in 2000	212	± 154.8	43	731	SINAVE 2000	
Protozoa 11 to 18 years						
Incidence in 2012	48	± 65.4	0	354	SINAVE 2012	
Incidence in 2006	78	± 97.5	2	902	SINAVE 2006	
Incidence in 2000	113	± 88.0	23	420	SINAVE 2000	
Prevalence of ow/ob (1-5 y) 2012	10.14	± 2.6	4.91	14.83	ENSANUT 2012	
Prevalence of ow/ob (6-10 y) 2012	34.54	± 6.3	22.52	51.53	ENSANUT 2012	
Prevalence of ow/obs (11-18 y) 2012	35.96	± 5.0	27.80	47.38	ENSANUT 2012	
Females (%)	53.4	2.2	50.6	56	ENSANUT 2012	
High marginality (%)	42.5	5.3	36.2	43.5	ENSANUT 2012	
Poverty (%)	46.6	± 13.3	21.0	78.5	INEGI. 2012	
Extreme Poverty (%)	11.2	± 8.7	1.8	38.3	INEGI. 2012	
Years in school	8.6	± 0.9	6.3	10.6	INEGI. 2012	
Health coverage (%)	62.4	± 10.9	39.9	81.0	INEGI. 2012	

ow/ob: overweight/obesity

348

**Table 2.** Linear regression model between the proximate probability of infection in 2012 with BMI for age z-score in 2012

	<i>Ascaris lumbricoides</i>									Protozoa																		
	$\beta$			95% C.I.			$p$	$\beta$			95% C.I.			$p$	$\beta$			95% C.I.			$p$							
Incidence in 2012	Crude model						Adjusted model						Crude model						Adjusted model									
1 to 5 years (n=8927)	-0.17	(	-0.18	-	-0.16	)	<0.01	-0.32	(	-0.33	-	0.31	)	<0.01	0.02	(	0.01	-	0.03	)	0.74	0.08	(	0.06	-	0.10	)	0.34
6 to 10 years (n=16347)	-0.15	(	-0.16	-	-0.14	)	<0.01	-0.21	(	-0.22	-	0.19	)	0.01	0.19	(	0.18	-	0.21	)	0.02	0.61	(	0.59	-	0.63	)	<0.01
11 to 18 years (n=13992)	0.17	(	0.16	-	0.19	)	0.05	0.16	(	0.13	-	0.18	)	0.23	0.43	(	0.42	-	0.45	)	0.00	0.85	(	0.83	-	0.88	)	<0.01

Adjusted by: urban/rural strata, age, sex, marginality, poverty, health-coverage. Incidence per 100 person-year.

349

350

**Table 3.** Linear regression model between the proximate probability of infection in 2000 and 2006 with BMI for age z-score in 2012

	<i>Ascaris lumbricoides</i>											protozoa																																
	$\beta$		95% C.I.			p		$\beta$		95% C.I.			p		$\beta$		95% C.I.			p		$\beta$		95% C.I.			p																	
Incidence in 2006	Crude model											Adjusted model											Crude model											Adjusted model										
1 to 5 years (n=9523)	-0.03	(	-0.04	-	-0.03	)	0.16	0.13	(	0.12	-	0.13	)	<0.01	0.00	(	0.00	-	0.01	)	0.79	0.00	(	0.00	-	0.01	)	0.93																
6 to 10 years (n=13025)	0.04	(	0.04	-	0.05	)	0.14	0.27	(	0.26	-	0.28	)	<0.01	0.07	(	0.06	-	0.08	)	0.14	0.13	(	0.12	-	0.13	)	0.01																
11 to 18 years (n=7845)	0.33	(	0.32	-	0.34	)	<0.01	0.50	(	0.49	-	0.52	)	<0.01	0.24	(	0.23	-	0.26	)	0.00	0.16	(	0.15	-	0.18	)	0.03																
Incidence in 2000																																												
1 to 5 years (n=6625)	0.04	(	0.04	-	0.05	)	<0.01	0.10	(	0.09	-	0.10	)	<0.01	0.29	(	0.28	-	0.30	)	0.00	0.47	(	0.46	-	0.48	)	<0.01																
6 to 10 years (n=7580)	0.08	(	0.08	-	0.09	)	<0.01	0.11	(	0.11	-	0.11	)	<0.01	0.58	(	0.56	-	0.60	)	0.00	0.61	(	0.59	-	0.63	)	<0.01																
11 to 18 years (n=5623)	0.19	(	0.18	-	0.19	)	<0.01	0.25	(	0.24	-	0.26	)	<0.01	0.88	(	0.85	-	0.90	)	0.00	0.99	(	0.96	-	1.02	)	<0.01																

Adjusted by: urban/rural strata, age, sex, marginality, poverty, health-coverage. Incidence per 100 person-year.

351

352

**Table 4.** Linear regression model between the proximate probability of infection with *A. lumbricoides* in 2012 with BMI for age z-score in 2012 according to statewide poverty rates

	Low Poverty					$\beta$	High Poverty							
	B	95% C.I.			p		95% C.I.			p				
<i>A. lumbricoides</i> Incidence in 2012														
1 to 5 years	0.33	(	0.31	-	0.35	)	<0.01	0.28	(	0.29	-	0.27	)	<0.01
6 to 10 years	0.51	(	0.48	-	0.53	)	<0.01	0.05	(	0.07	-	0.04	)	0.42
11 to 18 years	1.14	(	1.11	-	1.17	)	<0.01	0.34	(	0.31	-	0.36	)	0.01

Adjusted by: urban/rural strata, age, sex, marginality, poverty and health-coverage. Incidence per 100 person-year.