



UNIVERSITY OF LEEDS

This is a repository copy of *The Behavior of Low Molecular Weight Organic Carbon-14 Containing Compounds in Contaminated Groundwater During Denitrification and Iron-Reduction*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/156869/>

Version: Supplemental Material

Article:

Boylan, AA, Stewart, DI orcid.org/0000-0001-5144-1234, Graham, JT et al. (1 more author) (2020) *The Behavior of Low Molecular Weight Organic Carbon-14 Containing Compounds in Contaminated Groundwater During Denitrification and Iron-Reduction*. *Geomicrobiology Journal*, 37 (5). pp. 486-495. ISSN 0149-0451

<https://doi.org/10.1080/01490451.2020.1728442>

© 2020 Informa UK Limited, trading as Taylor & Francis Group. This is an author produced version of a paper published in *Geomicrobiology Journal*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

**Supporting Information for The behaviour of low molecular weight
organic carbon-14 containing compounds in contaminated groundwater
during denitrification and iron-reduction.**

Aislinn A. Boylan^a, Douglas I. Stewart^b, James T. Graham^c, Ian T. Burke^{a*}

This consists of five sections, S1, S2, S3, S4 and S5.

S1. Method for ^{14}C measurement in solid fraction

In to a 250 mL reaction vessel $1 \pm 0.01\text{g}$ of sediment was added with a magnetic stirrer. The system was sparged with $\text{N}_2(\text{g})$ prior to and during the extraction process, gas flow rate was set to 40 mL/min. Once sparged the reaction vessel was attached to two 125 mL gas washing bottles, the first of which contained 100 mL of Carbo-Sorb E and the second 100 mL of 1 M NaOH (acting as a safety bottle to ensure no release of $^{14}\text{CO}_2(\text{g})$). 20 mL of 2 M H_2SO_4 was added to the reaction vessel to achieve a pH ~ 3 when buffered by sediment. It was left to react for 30 minutes before disconnecting the gas washing bottles and exchanging for duplicate wash bottles. For the oxidation step 20 mL of 5% potassium persulfate and 4 mL 4% AgNO_3 were added to the container and heated to 80-90 °C. Further additions of $\text{K}_2\text{S}_2\text{O}_8$ and AgNO_3 at the same concentration and volume were made after one and two hours of reaction. The system was left for another hour to give a total oxidation reaction time of 3 hours. Triplicate samples of 1 mL were collected from wash bottles containing Carbo-Sorb E and mixed with 10 mL of PermaFluor E (Ahn et al., 2003; Magnusson et al., 2007). All samples were left to dark adjust for 24 hours prior to counting on the liquid scintillation counter.

S2. Experimental results for complete run

Experiments were run until a minimum of 80% of any ^{14}C -labelled LMWO was removed from solution. The complete data sets from addition of LMWO to the completion of the experiments for each condition are shown below.

S2.1 Nitrate amended microcosms, 7 to 22 days.




Figure S1 The results of the complete run of denitrification experiments (7 to 22 days) a) the percentage of ^{14}C in solution; b) pH; c) concentration of nitrite in solution; d) concentration of nitrate in solution (original nitrate addition = 10 mM, t=7 indicates time of ^{14}C addition). Error bars denote one standard deviation of triplicate measurements; where not shown error bars are less than the size of the symbol used.

S2.2 Iron-reducing microcosms, 28 to 60 days




Figure S2 The results of the complete run of iron-reducing experiments (28 to 60 days) a) the percentage of ^{14}C in solution; b) pH; c) fraction of Fe(II) in solids; d) concentration of Fe^{2+} in solution. Error bars denote one standard deviation of triplicate measurements; where not shown error bars are less than the size of the symbol used.

S3. Results of control experiments

Control experiments were run concurrently with active microcosms with the same incubation time to establish the desired redox condition. After the incubation time (7 days and 28 days for denitrification and iron reduction respectively) the microcosms were autoclaved at 120° for 30 minutes to eliminate microbial activity and the LMWO was added. The experiments were then run over the subsequent seven days to determine any abiotic processes which may affect the aqueous ¹⁴C concentration of the LMWO. The results of these experiments are shown below.

S3.1 Nitrate amended control microcosms




Figure S3 The results of the denitrification control experiments a) the percentage of ¹⁴C in solution; b) pH; c) concentration of nitrite in solution; d) concentration of nitrate in solution (original nitrate addition = 10 mM, t=7 indicates time of LMWO addition). Experiment duration from day 7 to day 14.

S3.2 Iron-reducing control microcosms




Figure S4 The results of the iron-reducing control experiments a) the percentage of ^{14}C in solution; b) pH; c) fraction of Fe(II) in solids; d) concentration of Fe^{2+} in solution.

Experiment duration from day 28 to day 35.

S4. OTU table

Table S1 Summary OTU table with all identified phylum and significant orders (*italics*) for all samples

	Unaltered			Denitrification				Iron-reducing			
	A	B	C	Acetate	Formate	Formaldehyde	Methanol	Acetate	Formate	Formaldehyde	Methanol
Acidobacteria	28.30	29.99	26.01	34.88	35.45	28.08	7.69	24.69	24.29	10.60	19.23
Actinobacteria	14.54	6.48	10.10	9.56	9.07	14.66	6.63	11.93	14.34	9.11	3.79
Armatimonadetes	1.23	1.06	2.03	1.01	1.00	0.86	0.09	0.23	0.28	0.45	2.28
Bacteroidetes	1.02	2.43	5.76	1.55	1.58	5.50	5.38	3.81	3.36	4.27	3.91
Chlamydiae	0.01	0.01	0.26	0.01	0.02	0.17	0.01	0.01	0.02	0.01	0.04
Chloroflexi	1.14	1.08	1.69	2.04	1.39	1.95	0.91	2.19	2.06	0.56	1.90
Deinococcus-Thermus	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Gemmatimonadetes	2.61	0.89	2.16	2.92	3.26	2.07	0.77	2.01	2.29	1.72	2.66
Planctomycetes	7.37	10.48	10.80	7.95	7.94	5.75	2.87	6.79	7.52	4.85	5.23
Alphaproteobacteria	9.99	12.64	12.78	10.98	11.05	9.92	6.22	13.16	11.78	13.88	9.40
Betaproteobacteria	5.67	9.16	6.76	11.52	13.10	12.19	34.77	21.51	21.60	37.62	28.58
<i>Burkholderiales</i>	1.47	1.72	2.29	6.44	7.33	8.15	32.16	14.60	10.64	34.41	9.48
<i>Methylophilales</i>	0.00	0.01	0.12	0.00	0.00	0.18	0.01	0.00	0.00	0.11	0.11
<i>Rhodocyclales</i>	0.01	0.01	0.10	0.01	0.00	0.01	0.13	0.19	0.27	0.01	0.02
Deltaproteobacteria	1.51	2.08	2.32	0.90	0.94	1.16	0.22	0.69	0.52	0.63	1.06
Gammaproteobacteria	3.12	2.61	3.87	3.39	3.08	4.19	30.39	2.81	2.25	7.87	10.86
<i>Chromatiales</i>	0.02	0.01	0.09	0.12	0.08	0.07	0.01	0.06	0.02	0.02	0.05
<i>Legionellales</i>	0.05	0.07	0.24	0.03	0.07	0.19	0.02	0.04	0.03	0.12	0.19
<i>Pseudomonadales</i>	0.46	0.52	0.63	0.59	0.45	0.55	0.25	0.38	0.40	0.40	0.26
<i>Xanthomonadales</i>	0.23	0.62	1.15	0.39	0.42	1.69	29.69	0.77	0.62	5.75	4.42
Proteobacteria - unclassified	3.92	2.88	3.19	2.82	2.70	2.04	0.82	2.29	1.86	0.99	1.64
Spiro	0.01	0.03	0.02	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00
Synergistetes	0.17	0.06	0.11	0.33	0.24	0.27	0.08	0.22	0.28	0.12	0.19
Verrucomicrobia	13.38	12.27	6.84	6.29	5.44	6.09	1.24	3.34	3.14	4.59	5.16
BRC1	0.05	0.12	0.08	0.02	0.02	0.02	0.00	0.00	0.00	0.07	0.01
candidate_division_WPS-1	0.37	0.50	0.55	0.24	0.25	0.39	0.07	0.10	0.10	0.18	0.35
candidate_division_WPS-2	0.34	0.64	0.63	0.44	0.33	0.49	0.08	0.20	0.21	0.22	0.31

S5. Siderite precipitation

Table S2 Input solution composition for siderite precipitation model

Solution composition	g/L in DIW
pH	6.7
Pe	0
KCl	0.006
MgSO ₄ ·7H ₂ O	0.976
MgCl ₂ ·6H ₂ O	0.081
NaCl	0.0094
Fe(II)	0.02
C as CO ₃ ²⁻	0.02

Geochemical modelling was undertaken using the PHREEQC (version 3) geochemical speciation program (Parkhurst and Appelo, 2013) and the Hatches database (version 18)(Cross and Ewart, 1991). Siderite saturation index predicted to be +0.02 under these conditions.

References

Ahn, H.J., Song, B.C., Sohn, S.C., Lee, M.H., Song, K., Jee, K.Y., 2013. Application of a wet oxidation method for the quantification of ³H and ¹⁴C in low-level radwastes. *Applied Radiation and Isotopes* 81, 62-66.

Cross, J. E., & Ewart, F. T. (1991). HATCHES—a thermodynamic database and management system. *Radiochimica Acta* 52(2), 421-422.

Magnusson, Å., Stenström, K., Aronsson, P.-O., 2007. ¹⁴C in spent ion-exchange resins and process water from nuclear reactors: A method for quantitative determination of organic and inorganic fractions. *Journal of Radioanalytical and Nuclear Chemistry* 275, 261-273.

Parkhurst, D. L., & Appelo, C. A. J. (2013). *Description of input and examples for PHREEQC version 3: a computer program for speciation, batch-reaction, one-dimensional transport, and inverse geochemical calculations* (No. 6-A43). US Geological Survey.