



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/158093/>

Version: Accepted Version

---

**Article:**

He, H., Reynolds, C., Hadjikakou, M. et al. (2020) Quantification of indirect waste generation and treatment arising from Australian household consumption: a waste input-output analysis. *Journal of Cleaner Production*, 258. 120935. ISSN: 0959-6526

<https://doi.org/10.1016/j.jclepro.2020.120935>

---

Article available under the terms of the CC-BY-NC-ND licence  
(<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

1 Quantification of indirect waste generation and  
2 treatment arising from Australian household  
3 consumption: a waste input-output analysis

4

5 He He <sup>a,e,g</sup>, Christian John Reynolds <sup>b,e,f</sup>, Michalis Hadjikakou <sup>c</sup>, Nicholas Holyoak <sup>d</sup>, John Boland <sup>e</sup>

6

7 <sup>a</sup> Department of Civil and Environmental Engineering, Imperial College London, Skempton Building, London,  
8 SW7 2AZ, United Kingdom

9 <sup>b</sup> Department of Geography, Faculty of Social Sciences, The University of Sheffield

10 <sup>c</sup> School of Life and Environmental Sciences, Deakin University

11 <sup>d</sup> School of Computer Science, Engineering and Mathematics, Flinders University

12 <sup>e</sup> Centre for Industrial and Applied Mathematics, Mawson Lakes Campus, University of South Australia,  
13 Mawson Lakes Boulevard, Mawson Lakes, SA 5095, Australia

14 <sup>f</sup>, Centre for Food Policy, City, University of London; Northampton Square, London EC1V0HB, UK

15 <sup>g</sup>, School of Engineering, University of Edinburgh, Edinburgh, United Kingdom

16

17 E-mail addresses: HE HE (h.he@imperial.ac.uk), Christian John Reynolds (c.reynolds@sheffield.ac.uk),  
18 m.hadjikakou@deakin.edu.au (Michalis Hadjikakou), nicholas.holyoak@flinders.edu.au (Nicholas Holyoak),  
19 John.Boland@unisa.edu.au (John Boland)

20

21

22

23

24

25

26

27 Abstract

28

29 Waste input-output (WIO) model is a suitable method to explore the nexus between economic  
30 activities and waste management. Contemporary research that typically explores this nexus follows  
31 two main aspects: either they consider Final demand as a whole, or they identify the nexus between  
32 households, with different types of socio-demographic indicators and household waste generation.  
33 However, it is complex to apply the WIO model from the perspective of household consumption— a  
34 major component of Final demand — because of a lack of economic and environmental data related  
35 to household consumption. This paper proposes a new perspective, applying the WIO model to  
36 assess the nexus between different patterns of household consumption and indirect waste  
37 generation and treatment. This novelty is to combine macro- and micro- economic and  
38 environmental data related to Australian industrial sectors, different patterns of household  
39 consumption (Mosaic data), and direct waste generation into the WIO model for exploring this nexus  
40 in two scenarios. Results indicate that the total amount of indirect waste generation caused by B05  
41 (couples without children who spend the majority of their time at the office) are 99.24 kg more than  
42 that of D16 (couples without children who are retired and stay at home) for scenario I. The  
43 correlation coefficients for differences of output of economy and indirect waste generation between  
44 B05 and D16 are 0.9796 and 0.9773 in scenarios I and II, respectively. Sensitivity analysis indicates  
45 the change of the amount of direct waste generation in a reasonable range cannot dramatically  
46 affect the major economic activities and waste generation. This research suggests a different  
47 perspective of household consumption to estimate indirect waste generation through a WIO model  
48 to provide more reliable information for waste management in the supply chain.

49

50 **Keywords:** waste input-output, household consumption, Australian economy, waste footprint

51

52

53

54

55

# 56 1. Introduction

57

58 Humans consume an increasing variety of goods and services produced by industrial sectors, which  
59 cause direct and indirect waste generation. Humans are the principal factor for driving production,  
60 consumption, and subsequently, the resulting waste generation (Karak et al. 2012). Direct waste  
61 generation of household consumption refers to the waste generated from household members at  
62 home (Ponis et al. 2017). Indirect waste generation in this research indicates that the waste  
63 generated in the supply chain of the economic system caused by household consumption. With the  
64 sharp growth of population, the amount of waste generation is forecasted to increase by about 70%  
65 globally in 2050 (The World Bank 2018).

66 Developed countries, such as United States, United Kingdom (UK), and Australia generate more  
67 waste per capita per day than developing countries (such as Bangladesh, Vietnam, and Malaysia)  
68 (Mmerekı et al. 2016). For example, the average amount of waste per capita per day in Australia  
69 (7.40 kilograms) (Department of the Environmental and Energy 2017) is about 10 times more than  
70 that the global average (0.74 kilograms) (The World Bank 2018) in 2016. The amount of Australian  
71 waste generation is forecasted to increase by approximately 60% by 2050 (Big Australia 2018).  
72 Developed countries have higher rates of recycling than developing countries. For instance, 60%  
73 waste generated in Australia was recycled in 2014–15 (Department of the Environmental and Energy  
74 2017) while only 22% of the total waste generated in Malaysia have been estimated to be recycled  
75 (Moh et al. 2014). There are considerable variations and complexities in waste generation due to  
76 different patterns of household consumption, the industrialized degree of countries, and the ability  
77 of treating waste.

78 One of the major global waste reduction goals is Sustainable Development Goal (SDG) 12, which  
79 seeks to substantially reduce waste generation through prevention, reduction, recycling and reuse  
80 (SDG 12.5), and halves per capita global food waste at the retail and consumer levels (SDG 12.3) by  
81 2030 (United Nations Statistics Division 2018). In order to achieve SDG 12, substantially reducing  
82 waste generation, and comprehending the complexity of waste management, there must be a  
83 systematic analysis of how waste generation and treatment interact with human consumption  
84 within an economic system on two fronts. First, how different patterns of human consumption  
85 activities affect waste generation and treatment should be analysed. Second, the nexus between the  
86 economy and waste generation must be assessed.

87 Studies regarding how/where waste is generated in the supply chain, and how waste is treated by  
88 different waste treatment methods have been proliferated. One popular research framework for  
89 displaying and analysing the complexity of waste production and treatment has been input-output  
90 (IO) modelling: a type of quantitative macroeconomic accounting that represents the  
91 interdependencies between different branches of a national economy or different regional  
92 economies. A short summary of IO modelling and waste follows: Joosten et al. (2000) used national  
93 supply–use tables to explore the nexus between plastic products and intermediate sectors in the  
94 Netherlands in 1990. Kagawa et al. (2004) delivered a simple multiregional IO model for waste  
95 analysis to estimate intraregional and interregional effects of industrial wastes caused by regional  
96 final consumptions. They have provided the analysis for the nexus between economic activities and  
97 waste management, but failed to account for detail analysis for patterns of consumptions and waste  
98 types. Nakamura and Kondo (2002) have linked waste types with treatment methods via the  
99 development of the waste input-output (WIO) model, which allows different types of waste  
100 treatment methods to “treat” or “dispose” of multiple types of waste. The WIO model has been  
101 developed further into a waste supply-use tables (WSUTs) that allows this complexity of treatment  
102 and waste flow to be seen in a single table (Lenzen and Reynolds 2014). The WSUT framework was  
103 used to demonstrate how Australian waste generation is affected by the intermediate sectors and  
104 waste treatment sectors (Reynolds et al. 2014). Based on the WSUTs, a multi-regional WSUT was  
105 developed to analyse the indicators of waste generation, such as the waste footprint and sectoral  
106 waste production intensity (Fry et al. 2015). Saleem et al. (2016) developed the first version of  
107 the UK WIO model table to analyse direct and indirect waste arising across the supply chain. He et al.  
108 (2017) compiled an Australian WIO table based on data from the Australian Bureau of Statistics  
109 (ABS). In terms of WIO model, the nexus between the economic system and waste management has  
110 been analysed to offer effective information (e.g. waste footprint and sectoral waste production  
111 intensity) for environmental policy-makers. Alabi et al. (2017) applied IO multiplier methods to  
112 develop an understanding of demand drivers of physical waste. Zeller et al. (2018) developed  
113 regional waste supply and use tables in terms of regional waste statistics and national input-output  
114 tables. These were used to quantify waste generation from households. Nakamura et al. (2018)  
115 extended function of the WIO model from a static model to a dynamic model, which covers the issue  
116 of quality in recycling that involves mixing, dissipation, and contamination. He et al. (2018)  
117 investigated the effect of the Household sector as an ‘endogenous’ factor on waste generation and  
118 treatment based on the environmentally-extend input-output model. Liao et al. (2015) have  
119 analysed the effect of household consumption on waste generation. Ruiz-Peñalver et al. (2019) have  
120 estimated total waste generation throughout the supply chain in Spain. These studies adjusted the

121 basic structure of IO table to explore the nexus between economic activities and waste generation and  
122 treatment based on the Final demand (Final demand being made up of consumption by households  
123 and government as well as capital formation, inventory and exports). However, there remains a lack  
124 of detailed analysis on 1) the effect of changing components within Final demand (ie household  
125 consumption), or 2) the socio-demographic sub-population effects of consumption on waste  
126 generation. Therefore, it is necessary to conduct detailed analysis from the perspective of different  
127 socio-demographic sub-population types to identify what effects the different patterns of  
128 consumption have on waste generation and treatment in the economic system.

129

130 Consumption is the major driver for waste generation (Wilson 2007). Consumption can be  
131 disaggregated into different types via different socio-demographic characteristics, such as income,  
132 education, and household size. Some studies explored the nexus between household consumption  
133 and waste generation (see (Parfitt et al., 2010, Song et al., 2015)), or waste composition (see  
134 (Daskalopoulos et al., 1998, Edjabou et al., 2015)). How different demographic indicators of  
135 consumption, such as income (Johnstone and Labonne 2004, Bandara et al., 2007, Aparcana 2017),  
136 household size (Dennison et al., 1996, Triguero et al., 2016), and education (Barr 2007, Benítez et al.,  
137 2008, Han et al., 2018) affected waste generation has been illustrated. Although the above-  
138 mentioned studies analyse the direct effect of the household consumption on waste generation,  
139 little attention has previously been paid to the effect of household consumption on indirect waste  
140 generation and treatment in the economy. The analysis of indirect waste generation in the supply  
141 chain can give a description of how human consumption patterns affect waste generation and  
142 treatment, which benefits decision-making for waste management. This lack of publications in this  
143 area is mainly due to the limitation of supply chain level waste data – rather than any conceptual or  
144 mathematical restriction. In addition, it is difficult to compare the results of how the different  
145 patterns of household consumption affect indirect waste generation and treatment across the  
146 regions and time frames because of inconsistencies of the scope and the substantial gaps in the  
147 available information (Thyberg et al., 2015, Reutter et al., 2017).

148

149 This paper aims at filling this knowledge gap through providing a novel perspective and case study to  
150 analyse the effects of different patterns of household consumption on the indirect waste generation  
151 and treatment in Australia, along with two novel variants to overcome limitations shown in the  
152 literature: (i) a version of Mosaic data describing the information of different patterns of  
153 consumption categorised by different socio-demographic data for the household; and (ii) on-site  
154 collected data for household waste generation and treatment corresponding to different patterns of

155 consumption to refine and compile the WIO table. This linking of Mosaic data and on-site waste  
156 collection data to a WIO table is a novel and new contribution to the literature.

157

158 The paper is structured as follows: the Method section gives information on the method of WIO  
159 model, data sources, sensitivity analysis, and the design of different scenarios. The Results section  
160 shows the effects of different patterns of household consumption on indirect waste generation and  
161 treatment and the comparative analysis between different scenarios with the sensitivity analysis.  
162 The Discussion section indicates the major findings based on different patterns of household  
163 consumption and scenarios with the discussed sensitivity analysis. The Conclusions section displays  
164 the novelty of the research, the advantages and disadvantages of the comparative analysis, and  
165 future research.

166

## 167 2. Method

168

### 169 2.1 Method of the WIO model

170

171 The basic method and notation of the WIO model was introduced in Nakamura and Kondo (2002).

172 The WIO model in balanced form from He et al. (2017) is written as

$$\begin{pmatrix} K_{I,I} & K_{I,II} \\ PG_{,I} & PG_{,II} \end{pmatrix} + \begin{pmatrix} X_{I,F} \\ PW_{,F} \end{pmatrix} = \begin{pmatrix} x_I \\ x_{II} \end{pmatrix} \quad (1)$$

173 where  $K_{I,I} \in \mathbb{R}^{N^I \times N^I}$  represents intermediate sectors' matrix for  $N^I$  goods and service-producing  
174 sectors, the components of  $K_{I,II} \in \mathbb{R}^{N^I \times N^{II}}$  mean the monetary inputs from per intermediate industry  
175 into  $N^{II}$  waste treatment sectors,  $P$  is an  $N^{II} \times N^w$  nonnegative matrix for  $N^w$  waste types, and the  
176  $p_{ij}$  in the matrix represents the proportion of waste  $j$  treated by waste treatment method  $i$ ,  $G_{,I}$  is  
177 defined as an  $N^w \times N^I$  matrix for the category of waste generated by intermediate sectors,  $G_{,II}$   
178 represents an  $N^w \times N^{II}$  matrix that the waste is generated by  $N^{II}$  waste treatment sectors. A Final  
179 demand matrix for  $N^I$  goods and service-producing sectors is defined as  $X_{I,F}$  for  $N^F$  sectors, and  $W_{,F}$   
180 is the waste generated by Final demand.  $x_I \in \mathbb{R}^{N^I \times 1}$  refers to a gross output vector for  $N^I$  goods and  
181 service-producing sectors, and  $x_{II} \in \mathbb{R}^{N^{II} \times 1}$  presents the total amount of waste to be treated by  $N^{II}$   
182 waste treatment sectors.

183 The coefficient matrix of WIO model can be expressed

$$\begin{pmatrix} A_{I,I} & A_{I,II} \\ B_{I,I} & B_{II,II} \end{pmatrix} \begin{pmatrix} X_I \\ X_{II} \end{pmatrix} + \begin{pmatrix} X_{I,F} \\ PW_{,F} \end{pmatrix} = \begin{pmatrix} X_I \\ X_{II} \end{pmatrix} \quad (2)$$

184 where the research defines input coefficients matrices  $A_{I,I} = K_{I,I} \hat{x}_I^{-1}$  (million \$AUD/million \$AUD),  
 185  $A_{I,II} = K_{I,II} \hat{x}_{II}^{-1}$  (million \$AUD/ tonne),  $B_{I,I} = PG_{,I} \hat{x}_I^{-1}$  (tonne/million \$AUD), and  $B_{II,II} = PG_{,II} \hat{x}_{II}^{-1}$   
 186 (tonne/tonne), where the “hat” over a vector  $x$  denotes a diagonal matrix with the elements of the

187 vector along the main diagonal. For instance, if  $x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$ , then  $\hat{x} = \begin{bmatrix} x_1 & 0 & 0 \\ 0 & x_2 & 0 \\ 0 & 0 & x_3 \end{bmatrix}$ .

188 The solution of Equation (2) is given by

$$\begin{pmatrix} X_I \\ X_{II} \end{pmatrix} = \left( I - \begin{pmatrix} A_{I,I} & A_{I,II} \\ B_{I,I} & B_{II,II} \end{pmatrix} \right)^{-1} \begin{pmatrix} X_{I,F} \\ PW_{,F} \end{pmatrix} \quad (3)$$

189 The research considers household consumption (HC) as the Final demand ( $X_{I,F}$  (million \$AUD)) in the  
 190 WIO model. Data collection and aggregation of household consumption and its direct waste

191 generation and treatment  $\begin{pmatrix} X_{I,F} \\ PW_{,F} \end{pmatrix}$  will be discussed in Section 2.2. Then, the total output will be

192 calculated in terms of Equation (3). Finally, the input matrix by industrial sectors of the WIO model

193  $\begin{pmatrix} K_{I,I} & K_{I,II} \\ PG_{,I} & PG_{,II} \end{pmatrix}$  can be obtained from Equation (4). The process of calculation is written as:

$$\begin{pmatrix} K_{I,I} & K_{I,II} \\ PG_{,I} & PG_{,II} \end{pmatrix} = \begin{pmatrix} A_{I,I} & A_{I,II} \\ B_{I,I} & B_{II,II} \end{pmatrix} \hat{x}^{-1} \quad (4)$$

## 195 2.2 Data sources

196

197 The Australian WIO tables in 2009–2010 and 2010–11 can be obtained in the research of He et al.

198 (2017), which are illustrated in Tables A.1 and A. 2, available in the supplementary file. To

199 summarise, the model is a 1 region, 8 intermediate sector model, that is based on domestic

200 technology consumption. This section describes the data collection regarding household

201 consumption ( $X_{I,F}$ ) and direct waste generation by household consumption ( $PW_{,F}$ ). Two types of

202 data source were employed for obtaining  $X_{I,F}$  and  $PW_{,F}$ :

203 (a) The household consumption ( $X_{I,F}$ ) were derived from the Mosaic Index of Mosaic data

204 (Nicholas 2016). Australian households are categorised into 13 Mosaic Groups and 49

205 Mosaic Types according to a series of socio-demographic variables, such as annual income,

206 education, and employment status. The Mosaic data contains information about weekly  
207 household consumption on goods and services (e.g. Fish and seafood and Bedroom furniture  
208 repair) for different Mosaic groups and Mosaic types. Each Mosaic Group includes three,  
209 four, or five Mosaic Types. These Mosaic Types are defined by the Grand index including 398  
210 variables according to the three following categories: (1) Who We Are, (2) Where We Live,  
211 and (3) What We Do (Nicholas 2016). D16 refers to couples without children who are retired  
212 and stay at home while B05 refers to couples without children who spend the majority of  
213 their time at the office. In Australia, these categories of Mosaic types D16 and B05 have  
214 accounted for an important proportion representing about 38% of all Australian families  
215 (ABS 2017) in the 2016 Census. There is a similar proportion for all Finnish households in  
216 2009 with 28% for couples without children (Katajajuuri et al. 2014). The proportion of these  
217 two Mosaic categories are expected to grow to more than 40% of the Australian families by  
218 2036 (ABS 2015).

219 (b) The amounts of waste generation ( $PW_{,F}$ ) of different types of Australian households were  
220 collected from an on-site experiment in the Lochiel Park, which that has recently been  
221 completed with approximately 110 homes in the north-east of Adelaide (Land Management  
222 Corporation, South Australian Government 2008) and is a living laboratory of CRC Low  
223 Carbon Living. The process of collecting data regarding  $PW_{,F}$  is described below. First, on-  
224 site survey about the amount of waste generated in eight households at Lochiel Park Green  
225 Village in South Australia has been conducted to collect data from households' waste bins  
226 (Ethics Approval, Application ID: 0000032810). In these eight households, seven of them  
227 belonged to D16 while one of them belonged to B05 of Mosaic types. The classifications of  
228 these eight households are based on the file of CRC Study Zone SA1 (Nicholas 2016). The  
229 research weighed these households' bins every week for a total of 14 times from December  
230 2nd 2015 to March 2nd 2016 by using an instrument marked in standard units. The three bin  
231 Kerbside Waste Collection service at Lochiel Park indicates the function of the blue bin for  
232 general waste, the yellow bin for recyclables, and the green bin for green organics  
233 (Campbelltown City Council 2017). The municipal solid waste collected from these  
234 households' bins are considered as the direct waste generation, which excludes bulky waste,  
235 Waste Electrical and Electronic Equipment recycling waste, and construction waste.  
236 Corresponding to the waste treatment sectors (the Landfill sector and the Recovery sector  
237 (resource recovery)) in the Australian WIO table, the research considered that the amount of  
238 waste in the blue bin were treated by the Landfill sector and that in the yellow and green  
239 bins was treated by the Recovery sector. Due to data limitations, it is not possible to

240 distinguish between energy recovery and material recovery ( material recycling). The  
 241 amounts of waste ( $PW_{,F}$ ) generated by D16 and B05 and treated by the Landfill sector and  
 242 the Recovery sector are shown in Tables 1 and 2, respectively. The household consumption  
 243 per week on 755 types of goods and services are shown in the sheet of DollarUnique of HES  
 244 Mosaic Index (Nicholas 2016). These 755 types of goods and services were aggregated into 8  
 245 types of goods and services ( $X_{I,F}$ ) corresponding to the number of intermediate sectors in  
 246 the Australian WIO table, which are shown in Table 3. The data of  $X_{I,F}$  and  $PW_{,F}$  were  
 247 multiplied by 52 to obtain the annual household consumption and waste generation and  
 248 treatment because the period of data of  $X_{I,F}$  and  $PW_{,F}$  was weekly.

249 The exact composition of the waste was not recorded by the survey, only the bin destination  
 250 (recycling or landfill). Though we do not have individual waste composition analysis, we provide  
 251 supplemental data of South Australian and Adelaide population level municipal waste compositional  
 252 analysis.

253 The source of the waste data the wider economy activities was sourced from He et al. (2017), which  
 254 was based on the Australian Bureau of Statistics waste accounts (2009–2010 and 2010–2011), (ABS  
 255 2013, 2014). The scope of waste from economic activities that is covered by Paper & Cardboard,  
 256 Glass, Plastics, Metals, Organics (e), Masonry, Electrical & Electronic, Solid Hazardous Waste, Leather  
 257 & Textiles, Tyres & Other Rubber, Timber & Wood Products, and Inseparable/Unknown. These waste  
 258 accounts were cross checked with the previous estimates of Reynolds et al (2015a). Both included  
 259 estimates of household waste. these were used to validate the results of our fieldwork.

260 **Table 1** The average amount of waste per week for D16 (couples without children who are retired and stay at  
 261 home).

Address	Mosaic Type	The amount of waste landfilled (kg)	The amount of waste recycled (kg)
1	D16	2.97	7.63
2	D16	9.04	5.43
3	D16	6.06	18.39
4	D16	2.74	5.14
5	D16	5.92	8.54
6	D16	4.82	7.12
7	D16	8.01	7.71
Average	D16	5.77	8.60
Total (52 weeks)	D16	300.04	447.2

262

263 **Table 2** The average amount of waste per week for B05 (couples without children who spend the majority of  
 264 their time at the office).

Address	Mosaic Type	The amount of waste landfilled (kg)	The amount of waste recycled (kg)
8	B05	3.83	7.85
Total (52 weeks)	B05	199.16	408.20

265

266 **Table 3** Aggregated household consumption per week on intermediate sectors.

Mosaic types	Ag	Mi	Ma	EGW	Co	Pa	AOI	WMS	Total
(\$AUD)									
D16	59.21	0	533.61	91.23	0	18.43	674.01	0.54	1377.03
B05	54.65	0	533.49	79.85	0	23.93	807.48	0.74	1500.14

267 Note: Agriculture, forestry, and fishing = Ag; Mining = Mi; Manufacturing = Ma; Electricity, gas, and water = EGW; Waste  
 268 management services = WMS; Construction = Co; Public administration = Pa; All other industry = AOI.

269

## 270 2.3 Method of the sensitivity analysis

271

272 The sensitivity analysis is a technique of assessing how the uncertainty in the output of a model can  
 273 be apportioned to different sources of uncertainty in the model input (Doubilet et al. 1985, Saltelli  
 274 2002, Hamby. 1994 ). Sensitivity analysis has been widely applied into IO model to identify the  
 275 uncertainty of the model output. As one of the major methods for sensitivity analysis, Monte Carlo  
 276 method has been used for the assessment of uncertainty relating to the level of aggregation (Bullard  
 277 and Sebald 1988), total CO<sub>2</sub> emission intensities (Hondo et al. 2002), multi-regional IO model to  
 278 convert currencies (Lenzen et al. 2010), impacts of the model on eco-efficiency assessment (Egilmez  
 279 et al. 2016), and technical waste and primary input coefficients (Yazan et al. 2016). Sensitivity  
 280 analysis has also applied to other environmental models. Clavreul et al. (2012) developed a general  
 281 method with a sequence of four steps for quantitative uncertainty assessment of life cycle analysis  
 282 (LCA) of waste management systems. Salemdeeb et al. (2017) applied the similar method to conduct  
 283 sensitivity analysis for the parameter values in the hybrid LCA approach. Due to the limitations of  
 284 household waste, this section conducts a sensitivity analysis to determine how the amount of  
 285 household waste (HW) collected on-site impacts the major variables in the WIO tables under a given  
 286 set of assumptions.

287 The first step is to generate a random sample of the amount of HW based on the sample which have  
 288 been collected on-site. The research first calculated the mean and standard deviation of the amount  
 289 of HW and then followed the method of Pollard (1979) to calculate the adjusted means and  
 290 deviations using a truncated normal distribution using the iterative Equations 5, 6, 7, and 8. The  
 291 truncated normal distribution is used as the amount of waste cannot be negative, and the authors  
 292 have assumed the maximum amount of waste to be the largest amount of on-site HW. The suffix n  
 293 denotes the nth approximation to the maximum likelihood estimate, the B and C are the maximum  
 294 and minimum numbers of the collected sample, and the symbols  $\phi$  and  $\Phi$  refer to the ordinate and  
 295 cumulative area of the unit normal curve,  $\bar{X}$  is the average of the collected data, S is the standard  
 296 deviation of the collected data. We found that after 5 iterations, stable estimates of the population  
 297 mean and standard deviation,  $\mu_n$  and  $\sigma_n$ , were reached:

$$298 \quad \alpha_n = (B - \mu_n)/\sigma_n \quad (5)$$

$$299 \quad \beta_n = (C - \mu_n)/\sigma_n \quad (6)$$

$$300 \quad \mu_{n+1} = \bar{X} + \sigma_n(\phi(\beta_n) - \phi(\alpha_n))/(\Phi(\beta_n) - \Phi(\alpha_n)) \quad (7)$$

$$301 \quad \sigma_{n+1}^2 = S^2 + (\bar{X} - \mu_{n+1}) + \sigma_n^2(\beta_n\phi(\beta_n) - \alpha_n\phi(\alpha_n))/(\Phi(\beta_n) - \Phi(\alpha_n)) \quad (8)$$

302 The second step is to apply the stable  $\mu_n$  and  $\sigma_n$  to obtain a random sample of the amount of HW  
 303 and input the sample into the WIO model to obtain the major indicators for waste generation and  
 304 treatment. This research chose 10 values of the sample as the amount of HW for sensitivity analysis.  
 305 This research also applies the same method to analyse 15 variables. It kept the variables of the  
 306 column of the Household sector in the WIO table constant and input these 10 values of the amount  
 307 of HW to obtain the corresponding indicators, such as the most inputs from the All other industry  
 308 (AOI) sector and the most amount of waste generated in the Manufacturing sector. The last step is  
 309 to calculate the coefficient of variation of the values of HW and indicators as well as identify  
 310 whether the accuracy of indicators is affected by the uncertainty of the values of HW or not.

311

## 312 2.4 Design of different scenarios

313

314 Due to the different periods of data sources, such as  $X_{I,F}$  in 2013,  $PW_{,F}$  in 2015–16, and Australian  
 315 WIO tables in 2009–10 and 2010–11, the research built two comparative scenarios. The comparative  
 316 analysis based on these two scenarios was conducted for illustrating the differences of indirect

317 waste generation caused by household consumption of B05 and D16. The main reason for  
318 conducting these two scenarios was to assess the effects of different years' economic situations on  
319 indirect waste generation and treatment. The two scenarios are:

320 1) Scenario I – the year of input coefficient and Leontief matrix is 2009–10, the year  $X_{I,F}$  is  
321 2013, and  $PW_{,F}$  is 2015–16.

322 2) Scenario II – the year of input coefficient and Leontief matrix is 2010–11, the year  $X_{I,F}$  is  
323 2013, and  $PW_{,F}$  is 2015–16.

324

## 325 3. Results

326

### 327 3.1 Waste footprint (Indirect) of two scenarios

328

329 This section presents waste footprints for household consumption (D16 & B05) with the focus on the  
330 share that is indirect generated in each industrial sector. Waste 'footprint' includes the waste people  
331 dispose of directly (direct), plus all the waste produced upstream (indirect) during the production of  
332 goods and services to satisfy human demand (Fry et al. 2015).

333 The (indirect) waste footprint caused by the household consumption of D16 (couples without  
334 children who are retired and stay at home) in Scenario I is shown in Figure 1. From the left to right,  
335 the diagram shows the amount of household consumption by D16, the amount of indirect waste  
336 generation in industrial sectors, and finally the amount of waste treated by waste treatment sectors.  
337 The amount of household consumption (left) is calculated by summarising the products and services  
338 of D16 consumption by retrieving the  $X_{I,F}$ . The amount of indirect waste generation in industrial  
339 sectors (middle) are found by retrieving the  $PG_{,I}$ . The amount of waste treated by the Landfill and  
340 Recovery sectors (right) are found by retrieving the  $PG_{,I}$  and  $PG_{,II}$ . In the supplementary appendix,  
341 Figures 2, 3, and 4 have the similar expressions for D16 in scenario II, B05 (couples without children  
342 who spend the majority of their time at the office) in scenarios I and II, respectively. In summary, the  
343 analysis of waste generation based on a WIO model from 2010 instead of 2009 results in a reduction  
344 of indirect waste generation of around 8%.

345

346 The amount of direct waste generation for D16 (747.24 kg) and B05 (607.36 kg) in Tables 1 and 2  
347 are less than that of indirect waste generation for D16 (scenario I: 2129.90 kg and scenario II:

348 1960.76 kg) and B05 (scenario I: 2230.14 kg and scenario II: 2046.23 kg). Each industrial sector  
349 generates indirect waste in scenario I more than that in Scenario II, except for the Mining sector. For  
350 example, the most amount of indirect waste caused by household consumption of D16 are  
351 generated from the Manufacturing sector in scenarios I (1033.81 kg) and II (1026.26 kg). As for waste  
352 treatment methods, the amount of indirect waste (1148.43 kg) treated by the Landfill sector is  
353 greater than that (981.47 kg) treated by the Recovery sector in scenario I. Scenario II shows the  
354 similar situation with the Landfill sector treating 1049.55 kg and the Recovery sector treating 911.21  
355 kg indirect waste. The exact composition of the waste in each stream/treatment method was not  
356 analysed due to lack of data.

357

358 The B05 has generated more indirect waste than the D16 in scenarios I (B05: 2229.14 kg and D16:  
359 2129.90 kg) and II (B05: 2044.23 kg and D16: 1960.76 kg) with more household consumption (B05:  
360 \$AUD 77,968.80 and D16: \$AUD 71577.48). The indirect waste generation and treatment caused by  
361 the household consumption of B05 has similar analysis with that by that of D16, shown in Figures 3  
362 and 4.

363

## 364 3.2 Comparative analysis of different types of households on 365 indirect waste generation in Australian economy with 366 different scenarios

367


368 This section compares indirect waste generated in each industrial sector, which are caused by the  
369 household consumption of B05 and D16 in different scenarios. The differences of indirect waste  
370 generation between B05 and D16 are calculated as the amount of indirect waste generation from  
371 the consumption of B05 minus that from that of D16.

372 Fig. 5 shows the comparative analysis of indirect waste generation for differences between B05 and  
373 D16 for scenarios I and II. The total amount of indirect waste generation caused by the consumption  
374 of B05 are 99.24 kg more than that of D16 in scenario I. The differences of indirect waste generation  
375 in all industrial sectors between B05 and D16 are positive, except for the Agriculture, forestry, and  
376 fishing sector and the Electricity, gas, and water sector. The All other industry sector has the largest  
377 positive difference for indirect waste generation in scenario I, amounting to 82.6 kg. The total  
378 amount of the differences of indirect waste generation (99.24 kg) between B05 and D16 in scenario I

379 are more than that (83.45 kg) in scenario II. The situations for the differences of indirect waste  
380 generation in each industrial sector caused by household consumption of B05 and D16 in scenario II  
381 are similar to that in scenario I.

382

383 Comparative analysis of indirect waste generation for differences between scenarios I and II for B05  
384 and D16 has been presented in Fig. 6. The differences of indirect waste generation between  
385 scenarios I and II are calculated as the amount of indirect waste generation in scenario II minus that  
386 scenario I for B05 and D16, respectively. The total amount of the indirect waste generation in  
387 industrial sectors caused by the household consumption of D16 between scenarios I and II amounts  
388 to -169.14 kg. Of this, the largest components were the All other industry (-86.88 kg), the  
389 Construction sector (-41.44 kg), and the Electricity, gas, and water sector (-31.26 kg). Difference of  
390 indirect waste generation in the Mining sector is the only positive number (8.62 kg). Compared with  
391 the data of D16, the household consumption of B05 results in a larger difference of indirect waste  
392 generation (-184.91 kg). The differences of indirect waste generation between scenarios I and II for  
393 B05 is similar to that for D16.



395

396

397



398

399

400


**Fig. 1.** Waste footprint of industrial sectors in Scenario I; indirect waste generation (industrial sectors) (middle) driven by the household of D16 (left) and treated by the Landfill and Recovery sectors (right). Note: Agriculture, forestry, and fishing = Ag; Mining = Mi; Manufacturing = Ma; Electricity, gas, and water = EGW; Waste management services = WMS; Construction = Co; Public administration = Pa; All other industry = AOI; Household consumption = HC; Household waste = HW.

401



402

403 **Fig. 5.** Comparative analysis of indirect waste generation for differences between B05 and D16 for scenarios I and II (kg). Note: Agriculture, forestry, and fishing  
 404 = Ag; Mining = Mi; Manufacturing = Ma; Electricity, gas, and water = EGW; Waste management services = WMS; Construction = Co; Public administration = Pa;  
 405 All other industry = AOI; Household consumption = HC.  
 406



407

408 **Fig. 6.** Comparative analysis of indirect waste generation for differences of scenarios I and II for B05 and D16 (kg). Note: Agriculture, forestry, and fishing = Ag;  
409 Mining = Mi; Manufacturing = Ma; Electricity, gas, and water = EGW; Waste management services = WMS; Construction = Co; Public administration = Pa; All  
410 other industry = AOI; Household consumption = HC.

411 **3.3 Correlation analysis between output of economy and**  
 412 **indirect waste generation in industrial sectors**

413  
 414 This section aims at exploring the nexus between the output of economy and waste generation in  
 415 industrial sectors. The objectives were to perform a) correlation analysis between the output of  
 416 economy and indirect waste generation in industrial sectors with the consumption of B05 or D16 in  
 417 scenarios I or II, b) correlation analysis between differences of the output of economy and differences of  
 418 indirect waste generation in industrial sectors for B05 and D16 in scenarios I or II, and c) correlation  
 419 analysis between differences of the output of economy and differences of indirect waste generation in  
 420 industrial sectors in scenarios I and II for B05 or D16. The correlation coefficients between the output of  
 421 economy and indirect waste generation in industrial sectors are strong and positive (Table 4). Table 5  
 422 shows that correlation coefficients between differences of output of economy and differences of  
 423 indirect waste generation in industrial sectors for B05 and D16 are 0.9796 in scenario I and 0.9773  
 424 scenario II, respectively. It is higher than correlation coefficients between the output of economy and  
 425 indirect waste generation in industrial sectors in scenario I for B05 (0.7515) or D16 (0.7470) and in  
 426 scenario II for B05 (0.6886) or D16 (0.6887), respectively (Table 4). Table 6 shows that there are weak  
 427 nexuses between differences of the output of economy and differences of indirect waste generation in  
 428 industrial sectors for B05 and D16 between scenarios I and II.

429 **Table 4** Correlation coefficient for B05 and D16 in scenarios I and II.

Scenarios	Household types	Total output for indirect waste generation
Scenario I	B05	0.7515
	D16	0.7470
Scenario II	B05	0.6886
	D16	0.6887

430

431 **Table 5** Correlation coefficient for differences between B05 and D16 in Scenario I or II.

Scenarios	Household types	Differences of total output for indirect waste generation
-----------	-----------------	-----------------------------------------------------------

---

Scenario I	Between B05 and D16	0.9796
Scenario II	Between B05 and D16	0.9773

---

432

433

434 **Table 6** Correlation coefficient for differences for B05 and D16 between scenarios I and II.

Household types	Scenarios	Differences of total output for differences of indirect waste generation
B05	Between scenarios I and II	-0.04365
D16	Between scenarios I and II	-0.06789

435

436 

### 3.4 Results of sensitivity analysis

437

438 In this section, the results based on the calculation of three steps mentioned in Section 2.3 have been  
 439 displayed. HW treated by the Landfill sector is considered as an example to illustrate the calculation in  
 440 detail. After 5 iterations from the equations 5 to 8, a stable  $\mu_n = 6.22$  and  $\sigma_n = 3.74$  based on the 7  
 441 samples of the Landfill sector was reached. Table 7 shows the 10 values of the amount of HW for the  
 442 Landfill sector and the Recovery sector selected from the random sample for  $\mu_n = 6.22$  and  $\sigma_n = 3.74$   
 443 based on normal distributions for D16. Figure 7 displays the Box-and-Whisker Plots of the estimated  
 444 amount of HW landfilled and recovered for D16.


445 **Table 7** The estimated amount of HW for the Landfill and Recovery sectors for D16 (kg per week).

Number	1	2	3	4	5	6	7	8	9	10
Landfill (Kilograms)	3.50	9.69	0.91	12.34	6.04	7.60	15.84	6.89	4.58	0.52
Recovery (Kilograms)	5.21	13.53	1.72	17.09	8.62	10.71	21.79	9.76	6.66	1.20

---

446

447



448

449

**Fig. 7.** Box-and-Whisker Plots of the estimated amount of HW landfilled and recovered for D16.

450

The amount of HW for the Landfill sector for D16 in scenario I were replaced by the values in Table 7.

451

The other values for the Household sector were kept constant for D16 in scenario I. The 10 WIO tables

452

for D16 corresponding to the 10 values of the amount of HW for the Landfill sector were then

453

constructed. This operation was carried out to allow analysis of the uncertainty around the amount of

454

HW treated by the Landfill sector. Specifically how changes to HW for D16 affects two major indicators:

455

1) the value of monetary inputs (for the All other industry sector) is linked to demand from households ;

456

and 2) changes to the value of waste caused by the shifting demand (and waste generation) of the

457

Household sector.

458

Table 8 reports the coefficients of variation of the amount of HW landfilled and recovered, the most

459

monetary inputs from the All other industry sector, and the most amount of waste generation by the

460

Manufacturing sector in 2009–10 and 2010–11. As there was only one value for B05, there was no

461

formal method to estimate standard deviation with certainty. In order to perform sensitivity analysis,

462

the research used the standard deviation of the sample of D16 to analyse the sample of B05. Table 9

463

shows the similar results of B05 in 2009–10 and 2010–11. Coefficient of variation in Tables 8 and 9 is a

464

measure of relative variability. It is the ratio of the standard deviation of the numbers of estimated

465

waste generation to the mean of the numbers of estimated waste generation. The coefficients of

466

variation for 10 variables and 15 variables show the similar result. This indicates that there is no need to

467

add more variables since adding extra 5 variables did not make any difference for estimated results.

468

469 **Table 8** The coefficients of variation of the amount of HW landfilled and recovered and major indicators for D16.

Year	Waste treatment methods	Indicators	Coefficients of variation
2009–10	Landfilled	The amount of HW	0.7143
		The monetary inputs from the All other industry sector	0.0001
		The waste generation by the Ma sector	0.0000
	Recovery	The amount of HW	0.6776
		The monetary inputs from the All other industry sector	0.0001
		The waste generation by the Ma sector	0.0000
2010–11	Landfilled	The amount of HW	0.7143
		The monetary inputs from the All other industry sector	0.0001
		The waste generation by the Ma sector	0.0000
	Recovery	The amount of HW	0.6776
		The monetary inputs from the All other industry sector	0.0001
		The waste generation by the Ma sector	0.0000

470

471

472

473

474

475

476

477

478

479

480 **Table 9** The coefficients of variation of the amount of HW landfilled and recovered and major indicators for B05.

Year	Waste treatment methods	Indicators	Coefficients of variation
2009–10	Landfilled	The amount of HW	0.7124
		The monetary inputs from the All other industry sector	0.0001
		The waste generation by the Ma sector	0.0000
	Recovery	The amount of HW	0.7790
		The monetary inputs from the All other industry	0.0001
		The waste generation by the Ma sector	0.0000
2010–11	Landfilled	The amount of HW	0.7124
		The monetary inputs from the All other industry sector	0.0001
		The waste generation by the Ma sector	0.0000
	Recovery	The amount of HW	0.7790
		The monetary inputs from the All other industry I sector	0.0001
		The waste generation by the Ma sector	0.0000

481

482

483 

## 4. Discussion

484 This research has proposed a new perspective to assess the nexus between the Australian household  
485 consumption and indirect waste generation and treatment. This is explored further using a case study  
486 based on the existing Australian WIO model, the Mosaic data for household consumption, and on-site  
487 collection data for direct waste generation and treatment. These have different impact perspectives on  
488 the research outcomes. Household consumption is the major factor to result in indirect waste  
489 generation. The design of different types of scenarios (existed Australian WIO model) provides more  
490 details for better understanding the effect of economy on indirect waste generation. The on-site

491 collection for direct household waste data has been integrated in the WIO table and make the table as a  
492 whole.

493 The strength of this research lies in its somewhat detailed description of economic activities and waste  
494 treatment caused by different patterns of household consumption through the combination of macro-  
495 and micro- economic and environmental data related to Australian industrial sectors, different types of  
496 household consumption, and waste generation. Most previous research relating to the IO model  
497 quantified direct waste generation from average or aggregated household consumption (Wiedmann et  
498 al. 2006, Reutter et al. 2017, Reutter et al. 2017) and ignored the fact that household consumption can  
499 be categorized into different types depending on different socio-demographic indicators. Although some  
500 researchers have analysed the indirect waste generation caused by the household in the supply chain  
501 (Parfitt et al. 2010), there is a lack of a detailed analysis about how the patterns of household  
502 consumption affect indirect waste generated in the supply chain. Now having shown that this method is  
503 feasible, a detailed analysis can be undertake using high resolution input-output models (Lenzen et al  
504 2014).

505 The proportion of direct waste generation from the household has been about 25% in the total amount  
506 of waste while that of indirect waste generation is about 75%. Similar results have been illustrated by  
507 Kim et al. (2017), which have shown that percentage of industrial waste is high with 76.3%, and lower  
508 household waste ( 26.5%). It is worthwhile indicating that this relationship between direct and indirect  
509 waste is based only on the scope and availability of data, if the scope of inclusion changes, then this  
510 relationship may change. The waste footprints show how the different patterns of household  
511 consumption caused indirect waste generated in industrial sectors and treated by waste treatment  
512 sectors. As for the same pattern of household consumption, the indirect waste generated in scenario II  
513 (the year of input coefficient and Leontief matrix is 2010–11) is less than that in scenario I (the year of  
514 input coefficient and Leontief matrix is 2009–10). This change could illustrate the developments that  
515 have been found in other studies, for example, cleaner technologies (Ahamed et al. 2016, Yong et al.  
516 2016), waste management (Zaman 2015), and economic model (Geissdoerfer et al. 2017, Haupt et al.  
517 2017). The Manufacturing sector generated the most amount of indirect waste. This corresponds to the  
518 results of other studies generation (Priefer et al. 2016, Van Ewijk and Stegemann 2016) that identified  
519 goods, such as food, clothes and electronic (e-) waste, consumed by the household and made by this  
520 sector as the main source of indirect waste generation. The waste footprints have illustrated that the  
521 total waste generated along the supply chain is about 3 times higher than what households dispose of

522 directly, which is similar to the result from Fry, Lenzen et al. (2016). The B05 has generated more  
523 indirect waste than the D16 in the same scenario with more household consumption. The result has  
524 been connected with that the higher level of household consumption tends to generate more amounts  
525 of waste (Dyson and Chang 2005, Sjöström and Östblom 2010, Suthar and Singh 2015). In the UK the  
526 amount of waste generation, such as food, clothing, paper, plastics, electronic (e-) waste, and glass, had  
527 an important growth due to the growth in the overall household expenditure: from around £34 billion in  
528 1971 to £795 billion in 2006 (Tudor et al. 2011). Although the amount of indirect waste treated by the  
529 Landfill sector is more than that by the Recovery sector, the differences between them becomes smaller  
530 from scenario I to II. It gives an insight that the Landfill sector will be not a major solution for waste  
531 management in Australia (ABC NEWS 2018).

532 The comparative analysis between household consumption of B05 and D16 shows the effects of  
533 different patterns of household consumption on indirect waste generation in industrial sectors. The  
534 comparative analysis between B05 and D16 for scenarios I and II indicates the largest differences of  
535 indirect waste generation occur in the All other industry sector, which contains a series of service  
536 sectors including Accommodation and Food services, Rental, Hiring and Real Estate Services,  
537 Professional, Scientific and Technical Services, and Arts and Recreation Services (ABS 2008). It indicates  
538 that after satisfying the basic human demand for living products, people who spend more money on  
539 services, such as education (Smyth et al. 2010, Fagnani and Guimarães 2017), health (Thakur et al. 2015,  
540 Almeida et al. 2017), and tourism (Arbulú et al. 2015), can generate more indirect waste. The increase of  
541 indirect waste generation in the All other industry sector for scenarios I and II could be explained in part  
542 by the growth of food waste in the Food services sector: for example, Reynolds et al. 2015b found an  
543 increase in the number of meals eaten outside the home in the context of the increase of the income.  
544 For example, the increase of the amount of the UK household waste generation is partly related to a  
545 2.1% increase in the number of meals eaten outside the home between 2012 and 2015 (The  
546 Environment, Food and Rural Affairs Committee 2017).

547 The negative numbers of indirect waste generation in the comparative analysis based on the same  
548 pattern of household consumption between scenarios I and II could indicate that more clean production  
549 technologies and environmental strategies have been applied in most industrial sectors to reduce the  
550 waste generation. For example, the Construction sector generated less waste in scenario II than scenario  
551 I, which can be partly attributed to the technology of Pre-Fabricated Construction (Sandanayake et al.  
552 2016) and 'Construction and demolition waste guide – recycling and re-use across the supply chain'

553 (Australian Government Department of Sustainability, Environment, Water, Population and  
554 Communities 2011). The only positive number of the differences for the indirect waste generation  
555 appears in the Mining sector. It illustrates that as the single largest producer of solid waste in Australia,  
556 the Mining sector has resulted in the cumulative solid waste legacy of mining due to the wide  
557 application of large-scale open cut mining (Worrall et al. 2009, Mudd 2010).

558 The correlation analysis shows that the differences of household consumption between B05 and D16  
559 has strong nexus with the differences of indirect waste generation in scenarios I and II. It further  
560 indicates that the household consumption has significant effect on indirect waste generation (Sjöström  
561 and Östblom 2010, Suthar and Singh 2015).

562 The results of the sensitivity analysis indicate that the change of the amount of household waste  
563 generation in a reasonable range cannot dramatically affect the major economic activities and indirect  
564 waste generation. For example, the change in the amount of HW for D16 (The coefficient of variation is  
565 0.7143.) has a slight significant change of the monetary inputs from the All other industry sector (The  
566 coefficient of variation is  $5.5 \times 10^{-5}$ ), and the amount of waste generation from the Manufacturing sector  
567 (The coefficient of variation is  $2.7 \times 10^{-5}$ ). It also indirectly reflects that most of waste in the Australian  
568 economy caused by the Household consumption is generated in the supply chain, rather than at the  
569 household level. Therefore, the data of the amount of waste weighed from on-site audit only performs a  
570 benchmark value for the WIO analysis.

571 Finally, our application of WIO analyses the effect of different patterns of household consumption on  
572 indirect waste generation by incorporating different types of scenarios. This allows macro- and micro-  
573 economic data to be integrated with waste data. This quantitative method would benefit the study of  
574 effects of household consumption on waste management in a national scale. It could also be used to  
575 assist in the design of environmental policies for different households in terms of environmental impacts  
576 related to their different consumption and waste generation patterns.

577 This ability to investigate indirect waste generation from the perspective of different patterns of  
578 household consumption is particularly important when considering the rapidly changing demography of  
579 Australia, the UK and many other countries globally. The methods proposed in this paper allow the  
580 waste management implications of this demographic change to be investigated at a higher level of detail  
581 than under previous WIO or other traditional methods.

582

## 583 5. Conclusions

584

585 The novelty of this research is to analyse the indirect waste generation in the supply chain from the  
586 perspective of patterns of household consumption. We combine macro- and micro- economic and  
587 environmental data related to Australian industrial sectors, different patterns of household  
588 consumption (Mosaic data), and direct waste generation into the WIO model for exploring this nexus in  
589 two scenarios. This research has demonstrated it is possible to analyse indirect waste generation and  
590 treatment arising from household consumption, this is typically hidden within the Australian economic  
591 and waste data. However, the scale of the IO model (8 sector), and the aggregated waste data make this  
592 paper more of a proof-of-concept study than a detailed investigation.

593 Results show that indirect waste generation is hidden from the end-user of products and services. We  
594 find the level of indirect waste generation for B05 (couples without children who spend most of their  
595 time at the office) and D16 (couples without children who are retired and stay at home) is greater than  
596 that of direct waste generation for D16 and B05. 75% of Australian household waste generation is  
597 related to indirect waste generation. It indicates that the waste generated in the supply chain is much  
598 more than the waste generated in the household. Due to this result we encourage the consideration of  
599 waste management strategies that conduct waste minimisation across the supply chain, rather than just  
600 at the consumer level. The amount of indirect waste generated by B05 is more than that by D16. There  
601 are two reasons for this result: 1) B05 spends more time at the office and eat more food outside; and 2)  
602 B05 spends more money for their living. Both of them can generate more indirect waste from the supply  
603 chain. Policy-makers should levy extra fee for waste generation in terms of household consumption. Our  
604 Correlation analysis indicates a clear nexus between household consumption and indirect waste  
605 generation. This indicates that technologies and policies published by Australian governments and  
606 aimed at reducing waste generation should focus on the supply chain or upstream processes in addition  
607 to on-site disposal.

608 However, the patchy economic and waste data regarding household consumption and waste  
609 management are a significant hurdle for analysis. First, in relation to the accuracy of the research,  
610 although the data of waste generation from on-site collection have limited effects on the research  
611 findings, we could further enhance the accuracy with data of waste from more samples of different  
612 types of household. Second, if the base data can be matched to the same period(s), the modelling will

613 obtain more accurate and useful results for policy-makers. Finally, this research does not intend to  
614 represent an entire class of people nationwide, but calculates for two large groups of the Australian  
615 population, how much waste will be indirectly generated in Australian industrial sectors due to their  
616 consumption and waste generation.

617 Indeed, a weakness of this paper is that the volumes of waste modelled have been presented and  
618 investigated in aggregate with no differentiation or waste composition analysis besides treatment  
619 destination. As Mmereki et al. (2016), states "the composition of solid waste varies greatly from country  
620 to country and changes significantly with time", this means that: 1) future changes to consumption and  
621 production will change the composition of the waste and thus environmental and economic impacts. 2)  
622 If this study was conducted in another geography and with other demographic groups the waste  
623 compositions will vary due to cultural and geographic factors. This complexity (between aggregated and  
624 disaggregated waste, destinations of waste, and cultural and geographic determinates) should be  
625 investigated in future research. A method to disaggregate generic waste volumes into component parts  
626 using IO tables has been proposed by Reynolds et al ( 2015a). In future work this method could be  
627 extended and linked with the scenarios presented in this paper, or compared with other geographies.

628 Furthermore, the IOT used in this study contains only one aggregated service sector (and only eight  
629 aggregated intermediate sectors in total). This aggregation means that differences in service sector  
630 consumption between B05 and D16 are not explored fully in our analysis, and the indirect waste  
631 generation differences cannot be explored fully. A disaggregated IOT and waste account, along with a  
632 full compositional audit of the waste would provide a much richer data source to model. A further  
633 weakness of the paper is that the IO tables it uses as a foundation are based on the assumption of  
634 domestic technology being identical to global technology (and vice versa). This has some major  
635 implications for our findings validation, as indirect waste generation will be embedded in both the global  
636 and Australian supply chains, and waste generation and production efficiency differences between  
637 supply chains is not currently taken into account.

638 With the above weaknesses in mind, the research presented here integrates the patterns of household  
639 consumption into a WIO model (albeit a model with only 8 intermediate sectors, and a single aggregated  
640 waste sector) . By using the micro-economic information of household consumption as the major driving  
641 force of indirect waste generation, it regards part of the operation of the supply chain as a function to  
642 generate waste in order to fulfil the different needs for human well-being. It is important from  
643 environmental management perspective to understand indirect waste generation with corresponding

644 implication for the nexus between economy and environmental issues. This research can be used as a  
645 bridge between different future household consumption scenarios. Further studies' directions based on  
646 this research for waste management are:

- 647 • Collect (disaggregated and detailed) waste data in terms of household types of Mosaic data to  
648 form a series of (detailed global or multiregional) WIO models to comprehensively explore the  
649 nexus between economic activities and waste generation caused by different types of  
650 household types, and

651 apply this method to analyse the indirect energy consumption and greenhouse gas emissions from  
652 different types of household consumptions. This research provides a method for obtaining more  
653 information regarding nexus between household consumption and indirect waste generation, allowing  
654 us to understand the effects of household consumption on indirect waste generated from industrial  
655 sectors. It is hoped this new capacity can 1) revitalize discussions around waste management and  
656 sustainable development from the perspective of household consumption, and 2) guide future data  
657 collection efforts.

658

## 659 References

660

661 ABC NEWS 2018, 'Landfill is not a long-term solution for waste management in Australia, viewed 27  
662 October 2018, <[https://www.abc.net.au/news/2018-11-18/now-is-the-time-to-scrap-landfill-in-](https://www.abc.net.au/news/2018-11-18/now-is-the-time-to-scrap-landfill-in-australia/10487726)  
663 [australia/10487726](https://www.abc.net.au/news/2018-11-18/now-is-the-time-to-scrap-landfill-in-australia/10487726)>.

664

665 Ahamed, Yin, Ng, Ren, Chang, and Wang. (2016). "Life cycle assessment of the present and proposed  
666 food waste management technologies from environmental and economic impact perspectives." Journal  
667 of Cleaner Production **131**: 607-614.

668

669 Alabi, Oluwafisayo Titilope and Turner, Karen and Allan, Grant and Swales, Kim. (2017). "Examining the  
670 nature and structure of a local pollutant: An illustrative case of physical waste generation in Scotland  
671 using environmental input-output accounting methods." University of Strathclyde.

672

673 Almeida, C., F. Agostinho, D. Giannetti, and B. Huisingh. (2017). "Cleaner production towards a  
674 sustainable transition." Journal of Cleaner Production **142**: 1-7.

675 Aparcana, S. (2017). "Approaches to formalization of the informal waste sector into municipal solid  
676 waste management systems in low-and middle-income countries: Review of barriers and success  
677 factors." Waste Management 61: 593-607.

678

679 Arbulú, I., J. Lozano, and Rey-Maqueira. (2015). "Tourism and solid waste generation in Europe: A panel  
680 data assessment of the Environmental Kuznets Curve." Waste Management **46**: 628-636.

681

682 Australian Bureau of Statistics (ABS) 2008, 1292.0 Australian and New Zealand Standard Industrial  
683 Classification (ANZSIC), 2006 (Revision 1.0), viewed 10 March 2018,  
684 <[http://www.abs.gov.au/AUSSTATS/abs@.nsf/Lookup/1292.0Main+Features12006%20\(Revision%201.0](http://www.abs.gov.au/AUSSTATS/abs@.nsf/Lookup/1292.0Main+Features12006%20(Revision%201.0)?OpenDocument)  
685 )?OpenDocument>.

686

687 ABS. 2013. 4602.055.005, Waste Account, Australia, Experimental Estimates, 2013; Australian Bureau of  
688 Statistics: Canberra, Australia, 2013, viewed 27 February 2020,  
689 <<https://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/4602.0.55.0052013?OpenDocument>>.

690

691 ABS. 2014. 4602.0.55.006—Waste Account, Australia, 2010–11; Australian Bureau of Statistics:  
692 Canberra, Australia, 2014, viewed 27 February 2020,  
693 <[http://www.abs.gov.au/Ausstats/abs@.nsf/0/682A9D1590A1168DCA257CCA00194E7E?OpenDocume](http://www.abs.gov.au/Ausstats/abs@.nsf/0/682A9D1590A1168DCA257CCA00194E7E?OpenDocument)  
694 nt>.

695

696 ABS. 2015. 3236.0 - Household and Family Projections, Australia, 2011 to 2036, viewed 27 October 2017,  
697 <<http://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/3236.02011%20to%202036?OpenDocument>  
698 >.

699

700 ABS. 2017. 2024.0 - Census of Population and Housing: Australia Revealed, 2016, viewed 27 October  
701 2017, <<http://www.abs.gov.au/ausstats/abs@.nsf/mf/2024.0>>.

702

703 Australian Government Department of Sustainability, Environment, Water, Population and Communities  
704 2011, Construction and demolition waste guide – recycling and re-use across the supply chain, viewed  
705 15 February 2018, <[http://www.environment.gov.au/protection/national-waste-](http://www.environment.gov.au/protection/national-waste-policy/publications/construction-and-demolition-waste-guide)  
706 [policy/publications/construction-and-demolition-waste-guide](http://www.environment.gov.au/protection/national-waste-policy/publications/construction-and-demolition-waste-guide)>.

707

708 Bandara, Nilanthi, J. Hettiaratchi, G. Wirasinghe, and Patrick Pilapiiya. (2007). "Relation of waste  
709 generation and composition to socio-economic factors: a case study." Environmental Monitoring and  
710 Assessment 135(1-3): 31-39.

711

712 Barr, Stewar. (2007). "Factors influencing environmental attitudes and behaviors: A UK case study of  
713 household waste management." Environment and Behavior 39(4): 435-473.

714

715 Bullard, C. W. and A. V. Sebald (1988). "Monte Carlo sensitivity analysis of input-output models." The  
716 Review of Economics and Statistics: 708-712.

717

718 Clavreul, J., D. Guyonnet and T. H. Christensen (2012). "Quantifying uncertainty in LCA-modelling of  
719 waste management systems." Waste Management 32(12): 2482-2495.

720

721 Daskalopoulos, Badr, and Probert. (1998). "Municipal solid waste: a prediction methodology for the  
722 generation rate and composition in the European Union countries and the United States of America."  
723 Resources, Conservation & Recycling 24(2): 155-166.

724

725 Dennison, Dodd, and Whelan. (1996). "A socio-economic based survey of household waste  
726 characteristics in the city of Dublin, Ireland—II. Waste quantities." Resources, Conservation and  
727 Recycling 17(3): 245-257.

728

729 Doubilet, P., Begg, Weinstein, Braun, and Mcneil. (1985). "Probabilistic sensitivity analysis using Monte  
730 Carlo simulation: a practical approach." Medical Decision Making 5(2): 157-177.

731

732 Dyson, B. and N.-B. Chang (2005). "Forecasting municipal solid waste generation in a fast-growing urban  
733 region with system dynamics modeling." Waste Management 25(7): 669-679.

734

735 Edjabou, Jensen, Götze, Pivnenko, Petersen, Scheutz, and Astrup. (2015). "Municipal solid waste  
736 composition: Sampling methodology, statistical analyses, and case study evaluation." Waste  
737 Management 36: 12-23.

738

739 Egilmez, G., S. Gumus, M. Kucukvar and O. Tatari (2016). "A fuzzy data envelopment analysis framework  
740 for dealing with uncertainty impacts of input–output life cycle assessment models on eco-efficiency  
741 assessment." Journal of Cleaner Production 129: 622-636.

742

743 Fagnani, E. and J. R. J. J. o. C. P. Guimarães (2017). "Waste management plan for higher education  
744 institutions in developing countries: The Continuous Improvement Cycle model." Journal of Cleaner  
745 Production 147: 108-118.

746

747 Fry, J., M. Lenzen, D. Giurco and S. Pauliuk (2015). "An Australian multi - regional waste supply - use  
748 framework." Journal of Industrial Ecology 20(6): 1295-1305.

749

750 Geissdoerfer, Savaget, Bocken, and Hultink. (2017). "The Circular Economy—A new sustainability  
751 paradigm?" . " Journal of Cleaner Production 143: 757-768.

752

753 Hamby, D. (1994). "A review of techniques for parameter sensitivity analysis of environmental models."  
754 Environmental Monitoring and Assessment 32(2): 135-154.

755

756 He, He ; Reynolds, Christian John ; Boland, John. (2018). "Assessment of solid waste generation and  
757 treatment in the Australian economic system: A Closed Waste Supply-Use model." Waste Management  
758 78: 346-355.

759

760 He He ; Christian John Reynolds ; Julia Piantadosi ; John Boland. (2017). "Effects of Australian Economic  
761 Activities on Waste Generation and Treatment." Recycling 2(3): 12.

762

763 Hondo, H., S. Sakai and S. Tanno (2002). "Sensitivity analysis of total CO<sub>2</sub> emission intensities estimated  
764 using an input–output table." Applied Energy 72(3-4): 689-704.

765

766 Han, Z., et al. (2018). "Influencing factors of domestic waste characteristics in rural areas of developing  
767 countries." Waste Management 72: 45-54.

768

769 Johnstone, N. and J. J. L. E. Labonne (2004). "Generation of household solid waste in OECD countries: an  
770 empirical analysis using macroeconomic data." Land Economics 80(4): 529-538.

771

772 Joosten, L., M. Hekkert, and E. Worrell. (2000). "Assessment of the plastic flows in The Netherlands using  
773 STREAMS." Resources, Conservation and Recycling 30(2): 135-161.

774

775 Kagawa, Shigemi, Hajime Inamura, and Yuichi Moriguchi. (2004). "A simple multi-regional input–output  
776 account for waste analysis." Economic Systems Research 16(1): 1-20.

777

778 Karak, Tanmoy, R.M. Bhagat, and Pradip Bhattacharyya. (2012). "Municipal solid waste generation,  
779 composition, and management: the world scenario." Critical Reviews in Environmental Science and  
780 Technology 42(15): 1509-1630.

781

782 Katajajuuri, J.-M., K. Silvennoinen, H. Hartikainen, L. Heikkilä and A. Reinikainen (2014). "Food waste in  
783 the Finnish food chain." Journal of Cleaner Production 73: 322-329.

784

785 Kim, W.-G., O. Nam-Chol and H.-S. Pak (2017). "Waste Composition for Solid Waste Management and Its  
786 Characteristic Analysis, a Case Study." Landscape Architecture and Regional Planning 2(3): 72.

787

788 Lenzen, M., R. Wood and T. Wiedmann (2010). "Uncertainty analysis for multi-region input–output  
789 models—a case study of the UK's carbon footprint." Economic Systems Research 22(1): 43-63.

790

791 Lenzen, M., et al. (2014). "Compiling and using input–output frameworks through collaborative virtual  
792 laboratories." Science of the Total Environment 485: 241-251.

793

794 Liao, M.-i., P.-c. Chen, H.-w. Ma and S. Nakamura (2015). "Identification of the driving force of waste  
795 generation using a high-resolution waste input–output table." Journal of Cleaner Production 94: 294-  
796 303.

797

798 Mmereki, Daniel, Andrew Baldwin, and Baizhan Li. (2016). "A comparative analysis of solid waste  
799 management in developed, developing and lesser developed countries." Environmental Technology  
800 Reviews 5(1): 120-141.

801

802 Moh, and Abd Manaf. (2014). "Overview of household solid waste recycling policy status and challenges  
803 in Malaysia." Resources, Conservation & Recycling 82: 50-61.

804

805 Mudd, G. M. J. R. P. (2010). "The environmental sustainability of mining in Australia: key mega-trends  
806 and looming constraints." Resources Policy 35(2): 98-115.

807

808 Nakamura, S. and Y. Kondo (2002). "Input\_Output Analysis of Waste Management." Journal of Industrial  
809 Ecology 6(1): 39-63.

810

811 Parfitt, Julian, Mark Barthel, and Sarah Macnaughton. (2010). "Food waste within food supply chains:  
812 quantification and potential for change to 2050." Philosophical Transactions of the Royal Society B  
813 365(1554): 3065-3081.

814

815 Pollard, J. H. (1979). A handbook of numerical and statistical techniques: with examples mainly from the  
816 life sciences. Cambridge: Cambridge UP, 1977.

817

818 Ponis, S. T., P.-A. Papanikolaou, P. Katimertzoglou, A. C. Ntalla and K. I. Xenos (2017). "Household food  
819 waste in Greece: A questionnaire survey." Journal of Cleaner Production **149**: 1268-1277.

820

821 Priefer, Jörissen & Bräutigam. (2016). "Food waste prevention in Europe—A cause-driven approach to  
822 identify the most relevant leverage points for action." Resources, Conservation & Recycling **109**: 155-  
823 165.

824

825 Reutter, Lant, Reynolds, and Lane. (2017). "Food waste consequences: Environmentally extended input-  
826 output as a framework for analysis." Journal of Cleaner Production **153**: 506-514.

827

828 Reutter, Lant, and Lane. (2017). "The challenge of characterising food waste at a national level—An  
829 Australian example." Environmental Science & Policy **78**: 157-166.

830

831 Reynolds, C. J., et al. (2014). "Estimating informal household food waste in developed countries: The  
832 case of Australia." Waste Management & Research **32(12)**: 1254-1258.

833 Reynolds, C., Geschke, A., Piantadosi, J., Boland, J., (2015a). Estimating industrial solid waste and  
834 municipal solid waste data at high resolution using economic accounts: an input–output approach with  
835 Australian case study. *J. Mater. Cycles Waste Manag.* **18**, 677–686. doi:10.1007/s10163-015-0363-1

836 Reynolds, C. J., J. Piantadosi, J. D. Buckley, P. Weinstein and J. Boland (2015b). "Evaluation of the  
837 environmental impact of weekly food consumption in different socio-economic households in Australia  
838 using environmentally extended input–output analysis." Ecological Economics **111**: 58-64.

839

840 Ruiz-Peñalver, S. M., M. Rodríguez and J. A. Camacho (2019). "A waste generation input output analysis:  
841 The case of Spain." Journal of Cleaner Production 210: 1475-1482.

842

843 Salemdeeb, R., E. K. zu Ermgassen, M. H. Kim, A. Balmford and A. Al-Tabbaa (2017). "Environmental and  
844 health impacts of using food waste as animal feed: a comparative analysis of food waste management  
845 options." Journal of Cleaner Production 140: 871-880.

846

847 Saltelli, A. (2002). "Sensitivity analysis for importance assessment." Risk Analysis 22(3): 579-590.

848

849 Sandanayake, M, Zhang, G, Setunge, S and Li, C. (2016). Environmental Emissions in Building  
850 Construction—Two Case Studies of Conventional and Pre-Fabricated Construction Methods in Australia.  
851 SCMT4: Fourth International Conference on Sustainable Construction Materials and Technologies,  
852 Sustainable Construction Materials and Technologies.

853

854 Sjöström, M. and G. J. E. E. Östblom (2010). "Decoupling waste generation from economic growth—A  
855 CGE analysis of the Swedish case." Ecological Economics 69(7): 1545-1552.

856

857 Smyth, Fredeen, and Booth. (2010). "Reducing solid waste in higher education: The first step towards  
858 'greening' a university campus." Resources, Conservation & Recycling 54(11): 1007-1016.

859

860 Song, Guobao M., Mingjing Li, Henry Musoke Semakula, and Shushen Zhang. (2015). "Food consumption  
861 and waste and the embedded carbon, water and ecological footprints of households in China." Science  
862 of the Total Environment 529: 191-197.

863

864 Suthar, S. and P. Singh (2015). "Household solid waste generation and composition in different family  
865 size and socio-economic groups: A case study." Sustainable Cities and Society 14: 56-63.

866

867 The Environment, Food and Rural Affairs Committee, 2017. Food waste in England, viewed 17 June  
868 2019, <<https://publications.parliament.uk/pa/cm201617/cmselect/cmenvfru/429/429.pdf>>.

869

870 Thyberg, Krista L, David J Tonjes, and Jessica Gurevitch. (2015). "Quantification of food waste disposal in  
871 the United States: a meta-analysis." Environmental Science & Technology 49(24): 13946-13953.

872

873 Triguero, A., et al. (2016). "Factors influencing willingness to accept different waste management  
874 policies: empirical evidence from the European Union." Journal of Cleaner Production 138: 38-46.

875

876 Tudor, T., G. M. Robinson, M. Riley, S. Guilbert and S. W. Barr (2011). "Challenges facing the sustainable  
877 consumption and waste management agendas: perspectives on UK households." Local Environment  
878 16(1): 51-66.

879

880 United Nations Statistics Division, 2018. SUSTAINABLE DEVELOPMENT GOAL 12—Ensure sustainable  
881 consumption and production patterns, viewed 15 May 2019, <[https://sdg-tracker.org/sustainable-](https://sdg-tracker.org/sustainable-consumption-production)  
882 [consumption-production](https://sdg-tracker.org/sustainable-consumption-production)>.

883

884 Van Ewijk, S. and J. J. J. o. C. P. Stegemann (2016). "Limitations of the waste hierarchy for achieving  
885 absolute reductions in material throughput." Journal of Cleaner Production 132: 122-128.

886

887 Wiedmann, Minx, Barrett, and Wackernagel. (2006). "Allocating ecological footprints to final  
888 consumption categories with input–output analysis." Ecological Economics 56(1): 28-48.

889

890 Worrall, Neil, Brereton, and Mulligan. (2009). "Towards a sustainability criteria and indicators  
891 framework for legacy mine land." Journal of Cleaner Production 17(16): 1426-1434.

892

893 Yazan, D. M., V. A. Romano and V. Albino (2016). "The design of industrial symbiosis: an input–output  
894 approach." Journal of Cleaner Production 129: 537-547.

895

896 Yong, Klemeš, Varbanov, and Huisinigh.(2016). "Cleaner energy for cleaner production: modelling,  
897 simulation, optimisation and waste management." Journal of Cleaner Production 111: 1-16.

898


899 Zaman, A. U. J. J. o. C. P. (2015). "A comprehensive review of the development of zero waste  
900 management: lessons learned and guidelines." Journal of Cleaner Production 91: 12-25.

901

902 Supplementary Appendix  
903 SA and Adelaide waste composition  
904

905 Sankey diagrams of additional scenarios


906



907

908 **Fig. 2.** Waste footprint of industrial sectors in Scenario II; indirect waste generation (industrial sectors) (middle) driven by the household of D16 (left) and  
909 treated by the Landfill and Recovery sectors (right). Note: Agriculture, forestry, and fishing = Ag; Mining = Mi; Manufacturing = Ma; Electricity, gas, and water =  
910 EGW; Waste management services = WMS; Construction = Co; Public administration = Pa; All other industry = AOI; Household consumption = HC.


911



912

913 **Fig. 3.** Waste footprint of industrial sectors in Scenario I; indirect waste generation (industrial sectors) (middle) driven by the household of B05 (left) and  
914 treated by the Landfill and Recovery sectors (right). Note: Agriculture, forestry, and fishing = Ag; Mining = Mi; Manufacturing = Ma; Electricity, gas, and water =  
915 EGW; Waste management services = WMS; Construction = Co; Public administration = Pa; All other industry = AOI; Household consumption = HC.

916



917  
 918 **Fig. 4.** Waste footprint of industrial sectors in Scenario II; indirect waste generation (industrial sectors) (middle) driven by the household of B05 (left) and  
 919 treated by the Landfill and Recovery sectors (right). Note: Agriculture, forestry, and fishing = Ag; Mining = Mi; Manufacturing = Ma; Electricity, gas, and water =  
 920 EGW; Waste management services = WMS; Construction = Co; Public administration = Pa; All other industry = AOI; Household consumption = HC.