



This is a repository copy of *Visualisation and quantification of wheat root system architecture and soil moisture distribution in an aggregated soil using neutron computed tomography*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/158929/>

Version: Published Version

Proceedings Paper:

Mawodza, T., Menon, M. orcid.org/0000-0001-5665-7464, Casson, S. et al. (1 more author) (2020) Visualisation and quantification of wheat root system architecture and soil moisture distribution in an aggregated soil using neutron computed tomography. In: Proceedings of EGU2020: Sharing Geoscience Online. EGU General Assembly 2020, 04-08 May 2020, Vienna, Austria. Copernicus GmbH .

<https://doi.org/10.5194/egusphere-egu2020-1196>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



Visualisation and quantification of wheat root system architecture and soil moisture distribution in an aggregated soil using neutron computed tomography

Tinashe Mawodza^{1,2}, Manoj Menon¹, Stuart Casson², and Genoveva Burca³

¹ Sheffield University, Geography, United Kingdom of Great Britain and Northern Ireland (t.mawodza@sheffield.ac.uk)

² Sheffield University, Department of Molecular Biology and Biotechnology, Sheffield, United Kingdom of Great Britain and Northern Ireland

³ Science and Technology Facilities Council, MAT facility, Harwell, United Kingdom of Great Britain and Northern Ireland

Sustainably intensifying global crop production in a world of diminishing natural resources is paramount for the attainment of zero hunger worldwide (a United Nations sustainable development goal). Key to this sustainable intensification is a deep understanding of the dynamics and complexities of plant-soil interactions for optimisation of plant productivity. Neutron computed radiography and tomography are powerful, non-invasive tools that enable the characterisation of plant-soil systems in situ. They also enable the visualisation and quantification of water distribution and movement within plant-soils systems. In this novel study, we use high resolution neutron computed tomography to investigate root system architectural differences in two different genotypes (Wild type vs TaEPF1-OE1-water use efficient mutant line) of bread wheat (*Triticum aestivum*). We further investigated how wheat roots interact with the heterogeneously distributed soil moisture. For this investigation, plants were grown in an aggregated sandy loamy soil with moderate amounts of organic matter (4%) for 13 days prior to imaging. We were able to produce a detailed three dimensional visualisation of the root architectural distribution of the two different genotypes imaged. These did not show significant differences between the two genotypes under investigation. We were also able to visualise relative soil moisture distribution and made inferences to how the roots of the wheat plants under investigation interact with the heterogeneously distributed soil moisture. Our results showed increased lateral root growth in regions with finer soil aggregates that had an estimated lower moisture content as compared to larger soil aggregates that retained higher amounts of moisture. This study demonstrates that detailed investigations into plant-soil interactions using neutron imaging techniques can be done successfully even in aggregated soils with considerable amounts of organic matter. This is a departure from the majority of neutron imaging experiments that predominantly use disaggregated sand soils devoid of organic matter as a growth medium.