

This is a repository copy of *Leishmaniasis immunopathology-impact on design and use of vaccines, diagnostics and drugs*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/159211/>

Version: Accepted Version

---

**Article:**

Kaye, Paul M [orcid.org/0000-0002-8796-4755](https://orcid.org/0000-0002-8796-4755), Cruz, Israel, Picado, Albert et al. (2 more authors) (2020) *Leishmaniasis immunopathology-impact on design and use of vaccines, diagnostics and drugs*. *Seminars in immunopathology*. pp. 1-18. ISSN 1863-2300

<https://doi.org/10.1007/s00281-020-00788-y>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# Seminars in Immunopathology

## Leishmaniasis immunopathology – impact on design and use of vaccines, diagnostics and drugs --Manuscript Draft--

<b>Manuscript Number:</b>	
<b>Full Title:</b>	Leishmaniasis immunopathology – impact on design and use of vaccines, diagnostics and drugs
<b>Article Type:</b>	Review Article (invited)
<b>Keywords:</b>	leishmaniasis; immunopathology; vaccines; diagnostics, drugs
<b>Corresponding Author:</b>	Simon Croft, PhD London School of Hygiene and Tropical Medicine Department of Global Health and Development UNITED KINGDOM
<b>Corresponding Author Secondary Information:</b>	
<b>Corresponding Author's Institution:</b>	London School of Hygiene and Tropical Medicine Department of Global Health and Development
<b>Corresponding Author's Secondary Institution:</b>	
<b>First Author:</b>	Simon Croft, PhD
<b>First Author Secondary Information:</b>	
<b>Order of Authors:</b>	Simon Croft, PhD
	Paul M. Kaye, University of York
	Israel Cruz
	Albert Picado
	Katrien Van Bocxlaer
<b>Order of Authors Secondary Information:</b>	
<b>Funding Information:</b>	
<b>Abstract:</b>	Leishmaniasis is a disease complex caused by 20 species of protozoan parasites belonging to the genus Leishmania. In humans the it has two main clinical forms, visceral leishmaniasis (VL) and cutaneous or tegumentary leishmaniasis (CL), as well as several other cutaneous manifestations in a minority of cases. In the mammalian host Leishmania infect different populations of macrophages where they multiply and survive in the phagolysosomal compartment. The progression of both VL and CL depends on the maintenance of a parasite-specific immunosuppressive state based upon this host macrophage infection. The complexity and variation of immune responses and immunopathology in humans and the different host interactions of the different Leishmania species has an impact upon the effectiveness of vaccines, diagnostics and drugs.

[Click here to view linked References](#)

# **Leishmaniasis immunopathology – impact on design and use of vaccines, diagnostics and drugs**

**Paul M. Kaye<sup>1</sup>, Israel Cruz<sup>2,4</sup>, Albert Picado<sup>2</sup>, Katrien Van Bocxlaer<sup>1</sup>, Simon L. Croft<sup>3</sup>**

<sup>1</sup>York Biomedical Research Institute, Hull York Medical School and Dept. of Biology, University of York, U.K.

<sup>2</sup>Foundation for Innovative New Diagnostics (FIND). Geneva, Switzerland.

<sup>3</sup>Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, U.K.

<sup>4</sup>National School of Public Health. Instituto de Salud Carlos III. Madrid, Spain

Correspondence: [simon.croft@lshtm.ac.uk](mailto:simon.croft@lshtm.ac.uk)

## 1: Introduction

Leishmaniasis is a disease complex with two main clinical presentations – visceral leishmaniasis (VL, or kala-azar) and cutaneous leishmaniasis (CL). Both VL and CL have a worldwide distribution in nearly 100 endemic countries. Overall the number of potentially fatal VL cases has decreased during the past decade with an estimated annual incidence of less than 50,000 (2017 figures, <https://www.who.int/leishmaniasis/burden/en> accessed 01.10.19) with seven countries: Brazil, Ethiopia, India, Kenya, Somalia, South Sudan and Sudan having the major burden. The decrease in VL case numbers in the Indian subcontinent (ISC) is coincident with the introduction of a regional elimination programme launched in 2005, with improved access to diagnosis and treatment. The naturally cyclical pattern of transmission intensity of this anthroponotic disease may also contribute to this decrease in the ISC. In contrast, the number of cases in East Africa has not fallen and this endemic area is now a major focus for control programmes. The estimated worldwide annual incidence of CL is between 0.7 to 1.2 million cases [1] with Central and South America, the Middle East, Ethiopia and North Africa being the major endemic areas. Far less attention has been paid to the control and treatment of CL, a predominantly zoonotic and a slow self-curing disease (3 – 18 months in most cases) with disfigurement and stigma being the main corollaries of infection. In addition, there are rarer variants of CL: mucocutaneous leishmaniasis (MCL), diffuse cutaneous leishmaniasis (DCL), disseminated cutaneous leishmaniasis (DsCL), leishmaniasis recidivans (LR) and post-kala-azar dermal leishmaniasis (PKDL). With CL, these variant skin manifestations are often referred to collectively as the tegumentary leishmaniases (TL). PKDL is a perplexing clinical presentation which follows cure of VL (5-10.% cases in Asia, up to 50% cases in East Africa, [2,3]).

The protozoan parasites that cause VL and TL belong to at least 20 distinct species of the genus *Leishmania* (with new species being identified in the past decade) that are transmitted between mammalian hosts by female phlebotomine sandflies. Different species of *Leishmania* are largely responsible for the diverse clinical manifestations (Table 1) which are generated by the interplay between the parasite, host factors and vector biology. However, clinical cases reports suggest these

species-dependent clinical outcomes are not distinct. In the sand fly gut *Leishmania* parasites exist as flagellated promastigote forms, while in the mammalian host the parasites invade macrophages losing their flagellum and transforming into the intracellular amastigote. The amastigote multiplies and survives in the phagolysosomal compartment of the macrophage. It has long been known that promastigotes differentiate within the sandfly to generate infective “metacyclic” forms [4-6] and recent evidence suggests that there may also be heterogeneity in the intracellular amastigote form [7-9]. *Leishmania* parasites actively manipulate both of their hosts. In the sandfly, transmission efficiency of metacyclic promastigotes is enhanced by secretion of a proteophosphoglycan-rich, mucin-like gel (Promastigote Secretory Gel, PSG), which accumulates in the sand fly gut and mouthparts [10]. Mammalian infection is established in the skin following the inoculation of metacyclic promastigotes that possess a lipophosphoglycan coat that enables them to resist complement and attach to and invade host cells. Peptides in sand fly saliva (for example, maxadilan) cause vasodilation and erythema that also aids the establishment of the infection in macrophages in the dermal layer of the skin [11]. Early responses to infection involve neutrophil infiltration and invasion of resident macrophages. Progress of the disease depends on the parasite species and host responses. For both VL and CL, disease progression depends on the maintenance of a parasite-specific immunosuppressive state including host cell macrophages. A key difference between VL and CL is the propensity of VL-causing species (Table 1) to establish disease at systemic sites, invading macrophages of the liver, spleen and bone marrow, with minimal skin pathology. Restoration of macrophage function, either through the development of an appropriate immune response or therapeutic intervention, can lead to amastigote killing in macrophages by nitric oxide and oxygen radicals. Resolution of disease following the activation of macrophages is enhanced by T helper 1 (Th1) - like responses mediated through the interaction of antigen-presenting cells (e.g., dendritic cells, macrophages) with CD4<sup>+</sup> and CD8<sup>+</sup> T cells and subsequent secretion of pro-inflammatory cytokines, including interferon- $\gamma$  (IFN $\gamma$ ) and tumor necrosis factor (TNF). In clinical forms of VL and in DCL, Th2 – like cell responses (including IL-4 and IL-13) are also evident [12,13] and inhibition of macrophage activation and immune dysregulation may be mediated through IL-10 and transforming growth factor- $\beta$  (TGF $\beta$ ) [14]. Immune responses and immunoregulatory pathways have been extensively defined in experimental inbred mice that show some but not all the characteristics of

**Commented [KVB1]:** Establish instead of cause to avoid word repetition?

human disease [15,16]. Immune responses directed against the parasite may also play a pathogenic role and contribute to tissue damage and clinical severity (see below).

The prevention, treatment and control of both CL and VL depends on drugs, diagnostics and vaccines, as described in recent comprehensive reviews [2,17]. The complexity and variation of immune responses and immunopathology in humans and the different host interactions of the *Leishmania* species has an impact upon the effectiveness of these tools (Fig. 1). Recent studies have significantly increased our understanding of these interactions and suggested new strategies for discovery and development, and such data are the focus of this review.

## **2. Key features of immunopathology in leishmaniasis**

Virulence factors allow pathogenic microbes to establish infection, colonise, replicate and disseminate throughout the host, evade anti-microbial immunity and facilitate transmission. A plethora of *Leishmania* derived molecules have been described and reviewed elsewhere that have the cardinal properties of bona fide virulence factors [18]. Although some pathogens liberate virulence factors that directly cause tissue damage e.g. gangrene due to collagenases and phospholipases produced by *Clostridium perfringens*, similar factors have not yet been described for *Leishmania*. Rather, clinical symptoms in leishmaniasis are, for the most part, the result of inappropriate immune responses leading to collateral damage to host tissues. Hence, for its clinically relevant features, leishmaniasis represents a disease driven by immunopathology. Hence, the “appropriateness” of an inflammatory / immune response, defined by its ability to strike the delicate balance between removal of *Leishmania* parasites and the minimization of pathology, may be defined by its quality, quantity or kinetics as well as by the parasite species under study. Some of the key features of the immunopathology associated with leishmaniasis are described below.

**Macrophage activation:** Macrophage activation is a central tenet of anti-leishmanial immunity, and the capacity of macrophages to polarise their functions towards the generation anti-microbial effector molecules such as NO and O<sub>2</sub><sup>-</sup> under the influence of type 1 cytokines (classical activation) or

towards the generation of arginine and polyamines under the influence of type 2 cytokines (alternate activation) is well established. However, the simple assumption that type I cytokines are “protective” and type 2 cytokines are “detrimental” is not always borne out by the literature. For example, IL-4, the prototypic Th-2 cytokine, plays an important early role in the generation of host protective CD4<sup>+</sup> IFN $\gamma$ <sup>+</sup> Th-1 cells in murine models of CL [19] and IL-4R signalling is required for optimal protection in primary resistance and vaccine-induced immunity in experimental VL [20-22]. Similarly, chemotherapy with sodium stibogluconate (SSG) in models of VL also requires IL-4R signalling for optimal effectiveness [22]. More recent evidence indicates that cytokines influence macrophage metabolic states, with consequences for the amastigote’s access to required nutrients. Specific metabolic countermeasures may have evolved to combat these stresses and ensure long lived parasitism [23].

Tissue macrophages have two origins. Resident tissue macrophages originate from embryonic yolk sac-derived progenitors, fully differentiate to acquire their mature characteristics under the direction of tissue-specific transcription factors / tissue-derived cues and have the capacity of self-renewal. In contrast, relatively short-lived tissue macrophages originate from hematopoietic stem cells via monocyte precursors [24,25]. Resident tissue macrophages play many of the key roles in tissue homeostasis and function, and they have been identified as niches for survival of *L. major* in the dermis [26] and *L. donovani* in the liver and spleen [27] and it is likely some of their housekeeping properties are impacted on by infection. Monocytes and monocyte-derived macrophages may also be targets of infection [28-30] though their relatively short life span may suggest that there is greater dynamics in the parasite populations they contain. Recent evidence suggests that the bone marrow becomes a target of collateral T cell-mediated damage during VL [31,32] but the consequences of this for the dynamics of *L. donovani* infection in these diverse macrophage pools has yet to be fully established. As discussed below, it is also possible that each of these macrophage / monocyte populations will have distinct relationships with anti-leishmanial drugs (see section 5), in addition to their roles in tissue repair, remodelling and restoration of homeostasis, that are central to the processes underpinning clinical cure.

*Granulomatous inflammation:* Granulomatous inflammation is a focal mononuclear cell-rich response to poorly degradable foreign material that seeks to wall off the insult and /or focus immune responses. Granuloma form and function in experimental, canine and human leishmaniasis have been reviewed in detail elsewhere [33]. Granulomatous inflammation is a hallmark of VL and correlates well in experimental models with the degree of immune response. Failure of granulomatous inflammation may contribute to the severity of HIV-VL coinfection [34]. Although over-exuberant, granulomatous inflammation may have directly pathologic consequences, e.g. in schistosomiasis, this has rarely been described in VL [35]. In TL, granulomas are more variably observed, and may reflect the time of biopsy in relation to disease evolution and causative agent. For example, a recent study in Tunisia demonstrated that both granuloma formation and the extent of the dermal lymphocytic infiltrate were minimal in *L. major* zoonotic CL compared to sporadic CL due to *L. infantum* [36]. Granulomas have also been observed in a majority of cases of non-ulcerating atypical CL due to *L. infantum* in Honduras [37] and are a valuable diagnostic characteristic of CL caused by *L. donovani* in Sri Lanka [38]. Although vaccine induced immunity in a primate models of VL was accompanied with enhancement of granuloma formation [39], it is not known whether this is a pre-requisite for vaccine-induced protection. In human vaccine trials, histopathological responses to vaccination have not been studied to date, and the histopathologic response documented following leishmanization was not documented [40]. The role of granulomatous inflammation in modifying the response to chemotherapy (see section 5 below) has not been formally addressed experimentally or clinically, largely because appropriate models do not exist and the invasive nature of tissue biopsies. It is possible that the intensity of granulomatous inflammation alters drug pharmacokinetics properties, and that macrophages at the core of such structures represent a “privileged” site for amastigotes, but this remains to be formally evaluated.

*Alterations in tissue architecture:* At homeostasis, each tissue has its own characteristic microanatomy, dedicated to performing its main function, be that of barrier or as a site of immune



response induction. By definition, tissue homeostasis is disrupted during clinical disease and this has been most well characterised in the context of remodelling of lymphoid tissue architecture during experimental and canine VL. Immune tissues are highly compartmentalised, with discrete B and T cell zones organised on a framework of stromal cells and fed by lymphatics and blood vessels [41]. During normal immune responses, these tissues adapt to allow for clonal retention and proliferation, but these are usually short-lived changes. Leishmaniasis poses a long-lived insult to lymphoid tissues and these respond by exaggerated degree of remodelling that may negatively impact on immune function. In the spleen of mice, dogs and humans infected with *L. donovani* / *L. infantum*, these changes include loss of stromal elements (follicular dendritic cells and fibroblastic reticular cells), dissolution of the marginal zone, white pulp atrophy and various degrees of admixing of splenic leucocyte populations [42-44].

*Systemic changes in metabolic and immune pathways:* Transcriptomics has recently been applied to a variety of experimental models and to human samples in order to provide a more holistic view of the pathology associated with both VL and CL. In human VL, comparisons between the whole blood transcriptome of healthy endemic controls with active cases, drug-cured cases and asymptomatic individuals have been reported from Brazil [45] and India [46]. Transcriptomic analysis of the splenic, hepatic and blood response to infection over time has been performed in mice [47] and for spleen only in hamsters [48]. High level analysis of these various reports indicates (i) IFN signatures are a prominent feature of infection; (ii) Th2 responses are variably represented with a bias to more severe disease; (iii) changes associated with cell cycle, lipid metabolism, angiogenesis and hematological disturbances are evident in active disease and reduced after treatment; (iv) treated patients, at least over the time period of study, fail to fully return to a homeostatic state. Not surprisingly, each study also identified unique aspects to the transcriptomic profile. A comprehensive cross-organ analysis in the mouse revealed only a limited concordance of DE genes between spleen, blood and liver and a clear lack of concordance across species [47]. Although such

comparative studies require a degree of caution (due to variations in analytical approach) they serve as a reminder of the importance of further research on the pathophysiology of human disease.

*CL pathogenesis:* In contrast to VL, where studies on pathology have focused on systemic sites, CL immunopathology has largely focused on the skin, though many of the changes observed in chronic VL lymphoid tissue may also occur in the lymph nodes draining CL lesion sites. Precise mapping of the nature of the inflammatory lesion and its development over time has been reported in murine models of CL, most notably due to *L. major* infection, but also with *L. mexicana* and *L. braziliensis*, with an emphasis on identifying determinants of host protection (reviewed in [49]). More recent studies spurred on by clinical observations in human *L. braziliensis* infection [50] and the identification of *Leishmania* viruses in metastatic strains of *L. guyanensis* [51] have focused on determinants of excessive host pathology. In *L. braziliensis* infection, a pathogenic role for CD8<sup>+</sup> T cells has now been firmly established. At a mechanistic level, highly cytotoxic CD8<sup>+</sup> T cells are believed to induce the release of cellular DAMPS that trigger inflammasome activation that perpetuates the inflammatory cascade. These studies provide opportunities for novel forms of immunotherapy to minimise tissue damage [52]. Viral infections are well known as stimulators of the type I interferon response and it has been shown that type I interferons either driven by endogenous symbiont *Leishmania* viruses or through concomitant bystander viral infection can lead enhance the metastatic potential of *L. guyanensis* in mouse models [53]. These data require further substantiation in a clinical setting, however, but also pose questions regarding novel therapeutic approaches.

### **3. Vaccines**

The case supporting development of a vaccine for leishmaniasis from a scientific and public health perspective has been made many times elsewhere [54]. Barriers to successful development of a vaccine to date have included i) lack of translational funding; ii) lack of correlates of immunity, iii) over-dependence on animal models; and iv) lack of a coherent programme of advocacy.

Nevertheless, recent years have seen excellent progress that has brought four candidate vaccines to or near to the clinic. These include a recombinant fusion protein delivered with strong Th1-inducing adjuvants (LEISHF3+ GLA-SE; [55]), a naked multi-epitope DNA vaccine (LeishDNAvac; [56]), an adenovirus-based vaccine (ChAd63-KH; [57]) and a live genetically attenuated vaccine (*L. major*/*L. donovani* centrin; [58]). Each has taken a different approach to vaccine antigen identification and vaccine delivery, providing a potential rich environment in the future for understanding the determinants of vaccine-induced immunity in humans.

Prophylactic vaccines by definition are used in non-infected individuals to prevent the development of disease. Hence, immunopathology associated with *Leishmania* infection does not play a determining effect on vaccine design or efficacy. This does not rule out, however, that a subset of the target population for vaccination in an endemic setting may harbour sub-clinical and undetectable infections (using the diagnostic tools discussed below) that nevertheless have some local pathologic consequence. More likely, however is the scenario whereby co-infections or other intrinsic (e.g. nutrition) or extrinsic (e.g. UV radiation) environmental factors impact on vaccine-induced lymphocyte activation, memory cell generation and /or effector and regulatory cell balance. Due attention should be paid to these possible factors when designing and evaluating future clinical trials.

In contrast, therapeutic vaccines require enhancement of the state of immunity in those already with leishmaniasis pathology. The extent to which ongoing pathology impacts the efficacy of therapeutic vaccines for CL is presently unknown and will require carefully designed clinical trials in which patients can be stratified according to pathologic criteria. In animal models of VL, therapeutic vaccination can overcome the immunosuppressive state induced by VL [59] and post-exposure prophylactic vaccination in *L. infantum*-infected dogs was shown to reduce progression to symptomatic VL [60]. Prevention of PKDL by vaccination of previously treated VL patients poses an interesting challenge, given that the immune status of treated VL patients is likely to have been improved but not normalised [46]. A clinical trial to assess this approach is in development

(clinicaltrials.gov; NCT04107961). Combination studies of vaccines deployed with additional immunomodulators to overcome pathology-induced immune regulation should be actively considered.

#### **4: Diagnostics**

As the leishmaniasis are characterized by their clinical pleomorphism confirmation based solely on clinical grounds is a challenge. The wide variety of skin and mucosal lesions in CL involves an extensive differential diagnosis. In addition, diseases like malaria, infectious mononucleosis and malignancies, among others, present with fever, hepatomegaly and splenomegaly as does VL, often accompanied by cachexia and malnutrition in chronic courses.

Parasite demonstration by microscopy of Giemsa stained tissue smears is the gold standard in the diagnosis of leishmaniasis; this is not always accessible and has variable sensitivity. Thus laboratory confirmation rates before treating CL can be as low as 5% in some settings, with patients being put on treatment with potentially toxic drugs without receiving a confirmatory diagnosis [61]. For VL invasive tissue sampling is required, usually from bone marrow or spleen and to a lesser extent lymph node. This requires expertise and specialised facilities for managing potential complications, so VL suspects are usually referred to specialised treatment centres for diagnosis. Because of this WHO recommended that a strict case definition for VL, i.e. Fever for more than 2 weeks plus hepatosplenomegaly and /or wasting; with this a positive serology enables treatment initiation [62].

##### Diagnosis for case management

A common feature in VL is a polyclonal hypergammaglobulinaemia with a marked increase in serum IgG level, including IgG with specificity for *Leishmania* [63]. Therefore, identifying anti-*Leishmania* specific antibodies is currently the cornerstone of VL diagnosis. Rapid diagnostic tests (RDTs) based on the recombinant antigen rK39, from a *Leishmania infantum* strain from Latin America (syn. *L. chagasi*), are widely used and they show very good performance in the Indian subcontinent (ISC): 97% sensitivity (95% CI: 90.0 - 99.5), 90.2% specificity (95% CI: 76.1 - 97.7). In eastern Africa the specificity is similar 91.1% (95% CI: 80.3 - 97.3), but the sensitivity drops to 85.3% (95% CI: 74.5 -

93.2) [64]. This inferior performance requires that in eastern Africa it is necessary to include the direct agglutination test (DAT) in the diagnostic algorithm (Fig. 2). This is a robust test based on whole *Leishmania* antigen and it has been extensively validated. It cannot be considered an RDT as it requires some degree of laboratory expertise and capacity, and results are obtained after an overnight incubation [65].

The peculiarities of the antibody response against specific *Leishmania* antigens are key in the performance of serological tests. VL elicits a different immune response and different levels of anti-*Leishmania* IgG across regions. rK39-based RDTs are based on a kinesin sequence from an American strain of *L. infantum*, this can explain the different performance of these tests in eastern Africa, where there is a high molecular diversity of the rK39 homologous sequences among regional *L. donovani* strains [66]. Another explanation for this is that there can be a different potency in the immune response, exemplified by the finding that anti-*Leishmania* IgG levels in VL patients from Sudan were significantly lower than in patients from India, independently of the geographic origin of the leishmanial antigen used to assess this [67]. A new RDT, based on the recombinant antigen rK28, may contribute to overcome this problem. The rK28 is a synthetic polypeptide containing rK39 repeats from a Sudanese *L. donovani* strain, flanked by HASPB1 repeats and the HASPB2 open reading frame from an Ethiopian strain [68]. Some preliminary studies claim that rK28 RDTs show better performance in eastern Africa than those using the rK39 antigen; however, data are not definitive, as studies comparing rK28 RDTs with the rK39 RDT IT-Leish (Bio-Rad), which is the one recommended in most eastern African guidelines for VL, are limited. These studies are either non-prospective or use just serum or plasma from well characterized controls and cases (confirmed by parasitology) [69-71]. A prospective large scale evaluation of an rK28 RDT will be conducted by the AfriKADIA consortium in Ethiopia, Kenya, Sudan and Uganda; results will be available by the end of 2020 [<https://www.afrikadia.org/>].

Unlike in VL, CL is not characterized by an elevated production of anti-*Leishmania* antibodies, and the level of these is also quite variable between the different forms of TL. Therefore, antibody detection tests are not useful in the diagnosis of these forms of leishmaniasis. Anti-*Leishmania*

antibodies are barely detectable in CL and to some extent these can be detected in ML and DCL but with limited and variable performance [62]. New opportunities for the serological diagnosis of CL may be brought by the detection of anti- $\alpha$ -Gal antibodies, with promising preliminary results for Old World leishmaniasis, nevertheless this field requires further research [72].

In immunosuppressed HIV+ patients, antibody-detection tests have limited sensitivity and cannot be used to rule out VL [73,74]. Also, the detection of total anti-*Leishmania* antibodies is not very helpful in the diagnosis of PKDL or VL relapses, as circulating antibodies from previous VL episodes remain for long periods even after successful treatment [75]. It seems however that exploring specific IgG types can be of help in assessing post-treatment cure and the risk of relapse, as well as in supporting the diagnosis of PKDL. The qualitative and quantitative detection of IgG1 in blood or serum/plasma samples, by means of ELISA or RDT using whole or recombinant rK39 leishmanial antigen, have shown the potential of IgG1 as a biomarker of post-chemotherapeutic relapse, as demonstrated with samples from Sudanese and Indian patients [76-78].

Antigen-detection tests can help in resolving some of the problems described above, these tests should be more specific and distinguish active from past infections. Most of the approaches used up to date for the detection of leishmanial antigens in VL patients have targeted urine [79-81]. A latex agglutination test has shown a highly variable sensitivity (36%-100%) and specificity (64%-99%) across endemic regions and different groups of patients, including HIV-positive, thus this has not been fully implemented [64]. Different antigen detection ELISA tests have been developed, and they have shown high sensitivity and specificity in a preliminary study using samples from VL patients from different endemic regions, as well as utility in monitoring treatment efficacy [82]. Lack of further evaluation, the need to process the sample and the technical requirements of an ELISA test have precluded putting them into routine practice. A more practical approach would be to detect leishmanial antigens in blood or serum/plasma. A prototype immunochromatographic test showed high sensitivity (96%) and specificity (99%) in the diagnosis of VL in Chinese patients using blood or serum [83]; unfortunately, further validation of this test and its results in other settings has not been pursued.

Commented [KVB2]: Remove 'can'?

A point-of-care (POC) test for early diagnosis of CL is much needed to benefit patients and communities, its implementation will reduce morbidity and transmission (where this is anthroponotic). A target product profile (TPP) for a POC for CL was developed by consultation among experts with a consensus that this POC test should target *Leishmania* antigens [84]. A promising option for the point-of-care diagnosis of CL that fits this TPP is an RDT detecting the leishmanial amastigote antigen peroxidoxin. This test has shown variable performance across endemic regions, but for CL due to *L. major* or *L. tropica* the sensitivity and specificity obtained in different studies in Tunisia, Morocco and Afghanistan is acceptable [85-87].

#### Diagnostics in the elimination and post-elimination of visceral leishmaniasis

WHO aims at eliminating VL as a public health problem in the Indian subcontinent (ISC) by 2020. The implementation of test-and-treat strategies, relying on the use of antibody-detection RDTs has been pivotal in moving towards this target [88]. This effort will need to be sustained in the next few years to avoid resurgence and move towards the interruption of *Leishmania* transmission by 2030 [89]. However the current diagnostic tools may not be adequate to support this endeavour.

More specific tests will be needed in the near and post-VL elimination phases, since the positive predictive value of antibody-detection RDTs decreases in a context of low endemicity [88,90]. Only VL suspects with symptoms (e.g. fever, splenomegaly) for more than two weeks are tested by the current RDTs, which precludes early diagnosis and treatment of acute cases who remain infectious in the community. Reducing the time between onset of symptoms and diagnosis-treatment has been identified as a key parameter for achieving and sustaining elimination [91]. But current antibody-detection RDTs do not allow this, as they are not specific for acute VL and need to be interpreted along with a clinical case definition [64]. Detection of *Leishmania* antigens may be a more specific approach [92,93,82,83], and these can be detected earlier than anti-*Leishmania* antibodies [81,94]. Thus antigen-detection RDTs would allow diagnosis of VL cases without a time constraint or strict clinical case definition (e.g. < 2 weeks of fever); as described above several approaches for leishmanial antigens detection in clinical samples have been developed, but for different reasons these

have not been implemented. A recent target product profile describes the features of an antigen detection RDT that can be used to assist in VL elimination [95]. Antigen detection tests would also allow diagnosing VL in groups of patients in which antibody detection tests underperform such as HIV-VL co-infected patients, relapses, as well as in PKDL. All these cases are gaining relevance in the last stages of the VL elimination in the ISC, as they are an important reservoir of *Leishmania* and are a serious threat to VL elimination [96,97,90].

Diagnosis of PKDL is a key factor in elimination efforts as well as in patient management, for PKDL treatments are long (up to 12 weeks). PKDL diagnosis is challenging as case confirmation requires microscopy and/or molecular tests in skin biopsies. As these diagnostic procedures are difficult to perform, diagnosis in endemic areas is often made clinically [98]. Antibody detection RDTs, which are part of the PKDL diagnosis algorithm in the ISC, have limited value and alternatives such as point-of-care molecular tests with promising performance (e.g. loop-mediated isothermal amplification [LAMP]) should be evaluated [99,90].

There have been few if any studies directly seeking to evaluate the impact of pathology on the design and / or performance of diagnostic tests for VL or CL. Although there may be limitations in translating such knowledge, a consideration of the role of pathology in determining the specificity and sensitivity of current and future tests may be warranted. For example, perturbations in the regulation of antibody responses are linked to disease progression [100] and antibody and antigen half-lives will be affected by changes in plasma or lesion protein binding capacity, glomerular filtration rate (e.g. secondary to immune complex deposition [101]) and or the degree of enteropathy [102]. Hence, each of the parameters highlighted in Fig. 3 will be influenced by the degree of pathology observed during the early pre-clinical, clinical and post-treatment phases of the disease. As above in the case of vaccine efficacy, these pathologic changes might occur directly as a consequence of ongoing or previous leishmaniasis, but equally so may reflect the indirect immunopathological consequence of coinfections or other immune insults.



## 5. Drugs

There has been considerable progress in the development of treatments for leishmaniasis over the past two decades. A number of new drugs have been registered for VL (miltefosine, the liposomal formulation AmBisome, paromomycin) with some taking a lead role in disease control and elimination programmes [103,2]. At the same time there is an inadequacy of drugs for CL [104,105]. However, for the first time in history the pipeline of novel oral drugs for the treatment of VL and CL is more promising with five novel compounds for VL [106-108] and a renewed focus to find novel treatments for CL [109,110]. In this section, two aspects of the drug – immune response interaction will be discussed (i) immunosuppression and immunomodulation, and (ii) the impact of immunopathology on pharmacokinetics (Fig. 4).

### Immunosuppression and immunomodulation

Interactions between drugs and the immune response have been known, but poorly understood, since the early 20<sup>th</sup> century. It still provides a strong focus for research: a recent study on antibiotics that showed how host cell metabolites both reduce drug activity whilst also enhancing host cell phagocytic activity provides an example of such work [111]. For leishmaniasis, as for other infectious diseases, there have been two directions of study: (i) to understand why drugs do not work in patients and disease models and (ii) the development of a strategy to stimulate/modulate the immune response to overcome any immunosuppression or accelerate cure.

Drugs used for clinical treatment of VL have to work effectively in most endemic regions within the context of the immunosuppression provoked by this disease. With the more profound immunosuppression associated with HIV-VL co-infections, the standard drugs become significantly less effective and only co-administrations have been shown to improve treatment [112]. The low efficacy of three of the drugs (liposomal amphotericin B, paromomycin and miltefosine) in eastern African VL [113,114] that are so effective in the Indian subcontinent, has not been explained. Clinical isolates from both regions show similar susceptibility to these drugs within standard *in vitro* assays, so

the clear implication is that host factors (immunology, pathology, pharmacokinetics [PKs]) are the responsible for this difference. The potential importance of PKs has been illustrated by Dorlo and colleagues [115] who have determined different miltefosine exposures in adults and children in eastern Africa. There are further clinical and experimental observations, one confirming, one confounding: (a) several of the drugs that work effectively as VL treatments in immunosuppressed VL patients (*ibid*) have significant dependence on cellular immune responses in mouse models of infection, either immuno-suppressed or deficient [116-118], (b) all the drugs, whether immune or non-immune dependent in these studies, have well described interactions with the immune system (see reviews by Dorlo (2012) for miltefosine [119], Cohen (2016) for amphotericin B [120] and Murray (2005) for pentavalent antimonials [121]). It is clear that more detailed studies are required to understand these interactions.

For CL, drugs work within the context of a slow self-cure immune response; exceptions include diffuse cutaneous leishmaniasis (DCL) where there is an absence of an effective cellular immune response and absence of effective treatment. The objectives for drug treatment, and combinations with immunomodulators, are to both reduce the parasite burden and at the same time accelerate self-cure. Much of the area lacks clear clinical data due to the absence of robust clinical studies [104,105].

The central role of the macrophage in the survival of *Leishmania* parasites has been the driving force for most studies on immunomodulation. Have we progressed much further than a statement from Bernard Shaw's play Doctor's Dilemma (1906) "There is at bottom only one genuinely scientific treatment for all diseases, and that is to stimulate the phagocytes"? BCG, which has well characterised stimulatory interactions with macrophages [122], was the immunodulatory component of treatment with pentavalent antimonials and the recommended treatment for CL in Venezuela [123]. It was also used in combination with alum and pentavalent antimonials in clinical trials for the treatment of post-kala-azar dermal leishmaniasis [124]. Bacterial cell wall components, muramyl dipeptide and trehalose dimycolate have also been used in clinical studies [125] on CL and experimental VL [126]. Based on clearer understanding of cytokine-macrophage interaction both IFN $\gamma$  [127] and GM-CSF [128] have been used in the treatment of both VL and CL. The rationale behind the potential application of GM-CSF for CL is threefold: (a) GM-CSF can promote

proliferation, activation and differentiation of various myeloid cells (macrophages, dendritic cells and neutrophils) and their progenitors, (b) clinical trials have reported increased wound healing of diverse wound types (including CL) upon topical GM-CSF application, and (iii) CM-CSF is able to modulate the tissue immune response by increasing the levels of the anti-inflammatory cytokine IL-10. Randomized controlled trials in Brazil have indicated that GM-CSF both after oral [129] and topical [130] administration reduced the healing time of CL lesions when given in combination with intravenous antimonials. In addition to its anti-inflammatory role, IL-10 also inhibits IFN $\gamma$  induced macrophage killing of *Leishmania* and Murray and colleagues [131] demonstrated in a model of VL that blockade of IL-10 using anti-IL-10R antibodies resulted in significant increase in antimonial drug activity. A clinical trial to evaluate combining AmBisome with anti-IL-10R antibodies in VL patients was proposed but later withdrawn due to unavailability of the required antibody (clinicaltrials.gov; NCT01437020).

Immunomodulator molecules which are also TLR agonists are being pursued for CL treatment. For example, CpG-D35, a D-type CpG TLR9 agonist, is currently undergoing Phase-I clinical trials after showing it was able to curtail lesion development upon administration to macaque monkeys infected with *L. major* [132]. Earlier studies indicated that this TLR9 agonist induces maturation of plasmacytoid dendritic cells and secretion of IFN $\alpha$  and  $\gamma$  without direct activation of B cells [133]. A small molecule, the imidazoline imiquimod a TLR7 agonist, was shown by Matlashewski and colleagues to stimulate macrophage functions including signal transduction and NO production in experimental models, sufficient to kill intra-macrophage amastigotes [134]. Subsequent clinical studies using imiquimod as a topical adjunct therapy to pentavalent antimonials were inconclusive. Three of these randomized controlled trials (RCT) [135-137] showed no significant increased cure rate for the combination of imiquimod plus pentavalent antimonials versus antimonials alone. Efforts continue to find more active analogues [138]. A different approach was adopted by Smith et al (2000) with the small molecule, tucaresol (in clinical trials for sickle cell disease), an orally bioavailable compound that enhances T-helper-cell activity, with the induction of increased IL-2 and IFN $\gamma$  levels in mice and humans and

provides a costimulatory signal between macrophages and T-cells [139]. In a mouse model of VL tucareosol gave a 60% reduction in parasite burden similar to that achieved by INF $\gamma$  in the same model.

Other recent approaches have also added considerably to understanding drug-immune interactions and possible new routes forward to immunomodulation and successful cure. Long-term clinical research on South American CL [52] showed that NLRP3 inflammasome is activated by CD8<sup>+</sup> T cell-mediated cytotoxicity and drives disease progression. Based on these observations, the group studied a number of small molecule inhibitors of the inflammasome, for example, MCC950 and the diabetes drug glyburide. In mouse models of CL, they showed that mice treatment with compounds that inhibit NLRP3 inflammasome activation, MCC950 or glyburide, failed to develop the severe disease seen in untreated mice. In VL, more detailed knowledge of the immune response has also been exploited. Based on knowledge of the immunopathology of the spleen, the receptor tyrosine kinase inhibitor sunitinib maleate was shown to induce restoration of splenic microarchitecture. Although this drug did not possess inherent anti-leishmanial activity, it afforded a significant dose sparing effect for subsequently administered antimonials [140].

CL is also characterised by local inflammation. A combination of the anti-inflammatory agent, pentoxifylline, with antimonial drugs has been shown to be an important adjunct therapy for some forms of CL and mucocutaneous leishmaniasis. In Brazil, RCTs have shown a synergetic effect of pentoxifylline in conjunction to antimonial therapy in ML patients [141] whereas this was not observed in patients suffering CL [142,143]. The difference in pentoxifylline efficacy as part of a combination therapy between New World CL and ML might be due to distinct differences in cytokine and macrophage populations present in the lesion [144] – an observation that merits further exploration as it could help elucidate immunomodulatory processes. It is worth noting that when tested in Old World CL, the combined approach of pentoxifylline with an antileishmanial agent showed enhanced healing in comparison to antileishmanial treatment only [145]. For further information on the activity of immunomodulators, see reviews by Dalton and Kaye (2010), Taslimi et al., (2018) and Adriaensen et al (2017) [140,146,147].

#### PK/PD relationships, immunopathology in drug monitoring and targets for host directed therapy.

Besides the direct involvement in the disease progression and outcome, the immunopathology also impacts drug pharmacodynamics (PD) and/or pharmacokinetics (PK). Impact on the latter has been shown in mice with CL, where the presence of parasites and the associated inflammation in the skin disturbed the cutaneous barrier function and physiology [148,149]. Interestingly, the permeation of hydrophilic compounds increased to a higher extent in comparison to lipophilic compounds which was hypothesized to be in part due to the inflammatory hydrophilic environment in CL skin. Whilst these findings are particularly important for topical drug administration where the drug is applied locally, increased drug levels were also observed in *Leishmania*-infected skin upon systemic drug administration [150,151] where they were associated with enhanced capillary leakiness and increased macrophage infiltration both of which could potentially be attributed to local inflammation. How *Leishmania* infection alters the skin tissue microenvironment in humans is yet unknown and is subject to further research.

During the development of VL, the liver, a major drug metabolism site for drugs, undergoes morphological and functional changes; hence, it seems evident that VL impacts drug pharmacokinetics. Studies demonstrated a deterioration of the capacity of host hepatic microsomal membranes to metabolise xenobiotics in VL infected mice and hamsters [152,153]. This is important especially for drugs metabolised by cytochrome P450 isoenzymes as this can lead to reduced drug efficacy (metabolites are active) or toxicity when increased amounts linger around in the systemic circulation. Drug distribution also appears to be affected. For amphotericin B, for example, a significantly lower concentration of drug was found in liver and spleen of VL-infected mice as compared to uninfected mice, which was hypothesised to be due to a reduced phagocytic activity in infected macrophages [155, 157]. These findings were not investigated in humans but it seems plausible that the PK of certain drugs is altered by the disruption of the liver function as observed in VL. Pathology is also known to affect drug ADME (absorption, distribution, metabolism and excretion) processes by alterations in protein binding which is important given that hypoalbuminemia is observed in both human and experimental VL. Low serum albumin levels have been associated with alterations in the degree of protein binding

of highly protein-bound drugs such as miltefosine and thus could potentially impact its PK-PD relationship [154,155].

Another phenomenon which remains unexplored in VL in patients and experimental models is the influence of granuloma formation on drug PK and thus efficacy. In tuberculosis, a granuloma forms a shielded lesion compartment that harbours bacteria and makes it difficult for drugs to permeate [156]; hence, the long and intense treatment regimen composed of a cocktail of four different drugs. Whilst difference in the extent of granulomatous inflammation are clear between leishmaniasis and TB [33], the altered local cellularity of a granuloma may nevertheless be of importance.

Of further importance is the impact of the pathology on drug PD which involves processes best reflected by “what the drug does to the body”. Most chemotherapeutics exert their activity by stimulating or inhibiting enzymes that are involved in pathways essential for parasite survival. Recent research describes the presence of different metabolically active parasite populations in CL lesions [8]. Kloehn et al (2015) measured semi-quiescent *L. mexicana* parasites in non-healing CL lesions in mice [7]. This reduced metabolic state is characterised by low transcription rates and protein turnover and might thus contribute to a reduced drug efficacy.

## Summary

Here we have illustrated how the complexity and variation of human immune responses and immunopathology and the different host interactions to twenty *Leishmania* species has an impact upon the effectiveness of vaccines, diagnostics and drugs (Fig. 1). Although rodent models have proved to be a useful tool for experimental studies on these interactions, well described limitations of (a) absence of models for important species (*L. aethiopica*, *L. tropica*, *L. braziliensis*), (b) differences between mouse and human immune responses (for example, to TLR agonists, [157]) and serological indicators, and (c) differences in pharmacokinetics between mouse and humans [158]), underpin most of this work. The application of transcriptomics, genomic data bases and methodologies to determine pharmacokinetics in infected tissues (not just plasma) in future studies in human subjects must now be

used to inform future work so tools for treatment, control and elimination can be used with more understanding and effectiveness.

## Acknowledgements

PK is supported by a Wellcome Trust Investigator Award (WT1063203); KVB received a fellowship award from the Research Council United Kingdom Grant Challenges Research Funder under grant agreement 'A Global Network for Neglected Tropical Diseases' grant number MR/P027989/1.

## References

1. Alvar J, Velez ID, Bern C, Herrero M, Desjeux P, Cano J, Jannin J, den Boer M (2012) Leishmaniasis Worldwide and Global Estimates of Its Incidence. *PloS one* 7 (5):e35671
2. Burza S, Croft SL, Boelaert M (2018) Leishmaniasis. *Lancet* 392 (10151):951-970. doi:10.1016/s0140-6736(18)31204-2
3. Mukhopadhyay D, Dalton JE, Kaye PM, Chatterjee M (2014) Post kala-azar dermal leishmaniasis: an unresolved mystery. *Trends Parasitol* 30 (2):65-74. doi:10.1016/j.pt.2013.12.004
4. Sacks DL (1989) Metacyclogenesis in *Leishmania* promastigotes *Experimental Parasitology* 69 (1):100-103
5. Dostalova A, Volf P (2012) *Leishmania* development in sand flies: parasite-vector interactions overview. *Parasites & vectors* 5:276. doi:10.1186/1756-3305-5-276
6. Serafim TD, Coutinho-Abreu IV, Oliveira F, Meneses C, Kamhawi S, Valenzuela JG (2018) Sequential blood meals promote *Leishmania* replication and reverse metacyclogenesis augmenting vector infectivity. *Nature Microbiology* 3 (5):548-555. doi:10.1038/s41564-018-0125-7
7. Kloehn J, Saunders EC, O'Callaghan S, Dagley MJ, McConville MJ (2015) Characterization of Metabolically Quiescent *Leishmania* Parasites in Murine Lesions Using Heavy Water Labeling. *PLOS Pathogens* 11 (2):e1004683. doi:10.1371/journal.ppat.1004683
8. Mandell MA, Beverley SM (2017) Continual renewal and replication of persistent *Leishmania* major parasites in concomitantly immune hosts. *Proceedings of the National Academy of Sciences of the United States of America* 114 (5):E801-e810. doi:10.1073/pnas.1619265114
9. Burchmore RJS, Barrett MP (2001) Life in vacuoles – nutrient acquisition by *Leishmania* amastigotes. *International Journal for Parasitology* 31 (12):1311-1320. doi:[https://doi.org/10.1016/S0020-7519\(01\)00259-4](https://doi.org/10.1016/S0020-7519(01)00259-4)
10. Rogers ME, Chance ML, Bates PA (2002) The role of promastigote secretory gel in the origin and transmission of the infective stage of *Leishmania mexicana* by the sandfly *Lutzomyia longipalpis*. *Parasitology* 124:495-507
11. Giraud E, Svobodova M, Muller I, Volf P, Rogers ME (2019) Promastigote secretory gel from natural and unnatural sand fly vectors exacerbate *Leishmania* major and *Leishmania tropica* cutaneous leishmaniasis in mice. *Parasitology*:1-7. doi:10.1017/s0031182019001069

12. Pirmez C, Yamamura M, Uyemura K, Paes-Oliveira M, Conceicao-Silva F, Modlin RL (1993) Cytokine patterns in the pathogenesis of human leishmaniasis. *The Journal of clinical investigation* 91 (4):1390-1395. doi:10.1172/jci116341
13. Nylen S, Sacks D (2007) Interleukin-10 and the pathogenesis of human visceral leishmaniasis. *Trends in immunology* 28 (9):378-384. doi:10.1016/j.it.2007.07.004
14. Caldas A, Favali C, Aquino D, Vinhas V, van Weyenbergh J, Brodskyn C, Costa J, Barral-Netto M, Barral A (2005) Balance of IL-10 and interferon-gamma plasma levels in human visceral leishmaniasis: implications in the pathogenesis. *BMC infectious diseases* 5:113-113. doi:10.1186/1471-2334-5-113
15. Kaye P, Scott P (2011) Leishmaniasis: complexity at the host-pathogen interface. *Nature Reviews Microbiology* 9 (8):604-615
16. Scott P, Novais FO (2016) Cutaneous leishmaniasis: immune responses in protection and pathogenesis. *Nat Rev Immunol* 16 (9):581-592. doi:10.1038/nri.2016.72
17. Torres-Guerrero E, Quintanilla-Cedillo MR, Ruiz-Esmenjaud J, Arenas R (2017) Leishmaniasis: a review. *F1000Research* 6:750. doi:10.12688/f1000research.11120.1
18. Atayde VD, Hassani K, da Silva Lira Filho A, Borges AR, Adhikari A, Martel C, Olivier M (2016) Leishmania exosomes and other virulence factors: Impact on innate immune response and macrophage functions. *Cellular immunology* 309:7-18. doi:10.1016/j.cellimm.2016.07.013
19. Biedermann T, Zimmermann S, Himmelrich H, Gummy A, Egeter O, Sakrauski AK, Seegmuller I, Voigt H, Launois P, Levine AD, Wagner H, Heeg K, Louis JA, Rocken M (2001) IL-4 instructs TH1 responses and resistance to Leishmania major in susceptible BALB/c mice. *Nat Immunol* 2 (11):1054-1060. doi:10.1038/ni725
20. Stager S, Alexander J, Kirby AC, Botto M, Rooijen NV, Smith DF, Brombacher F, Kaye PM (2003) Natural antibodies and complement are endogenous adjuvants for vaccine-induced CD8+ T-cell responses. *Nature medicine* 9 (10):1287-1292. doi:10.1038/nm933
21. McFarlane E, Mokgethi T, Kaye PM, Hurdial R, Brombacher F, Alexander J, Carter KC (2019) IL-4 Mediated Resistance of BALB/c Mice to Visceral Leishmaniasis Is Independent of IL-4Rα Signaling via T Cells. *Frontiers in Immunology* 10 (1957). doi:10.3389/fimmu.2019.01957
22. Alexander J, Carter KC, Al-Fasi N, Satoskar A, Brombacher F (2000) Endogenous IL-4 is necessary for effective drug therapy against visceral leishmaniasis. *Eur J Immunol* 30 (10):2935-2943. doi:10.1002/1521-4141(200010)30:10<2935::aid-immu2935>3.0.co;2-q
23. Sernee MF, Ralton JE, Nero TL, Sobala LF, Kloehn J, Vieira-Lara MA, Cobbold SA, Stanton L, Pires DEV, Hanssen E, Males A, Ward T, Bastidas LM, van der Peet PL, Parker MW, Ascher DB, Williams SJ, Davies GJ, McConville MJ (2019) A Family of Dual-Activity Glycosyltransferase-Phosphorylases Mediates Mannogen Turnover and Virulence in Leishmania Parasites. *Cell host & microbe* 26 (3):385-399.e389. doi:10.1016/j.chom.2019.08.009
24. Yona S, Kim KW, Wolf Y, Mildner A, Varol D, Breker M, Strauss-Ayali D, Viukov S, Guillemins M, Misharin A, Hume DA, Perlman H, Malissen B, Zelzer E, Jung S (2013) Fate mapping reveals origins and dynamics of monocytes and tissue macrophages under homeostasis. *Immunity* 38 (1):79-91. doi:10.1016/j.immuni.2012.12.001
25. Guillemins M, Scott CL (2017) Does niche competition determine the origin of tissue-resident macrophages? *Nat Rev Immunol* 17 (7):451-460. doi:10.1038/nri.2017.42
26. Lee SH, Charmoy M, Romano A, Paun A, Chaves MM, Cope FO, Ralph DA, Sacks DL (2018) Mannose receptor high, M2 dermal macrophages mediate nonhealing Leishmania major infection in a Th1 immune environment. *The Journal of experimental medicine* 215 (1):357-375. doi:10.1084/jem.20171389
27. Beattie L, Sawtell A, Mann J, Frame TCM, Teal B, de Labastida Rivera F, Brown N, Walwyn-Brown K, Moore JWJ, MacDonald S, Lim EK, Dalton JE, Engwerda CR, MacDonald KP, Kaye PM (2016) Bone marrow-derived and resident liver macrophages display unique transcriptomic signatures but similar biological functions. *Journal of hepatology* 65 (4):758-768. doi:10.1016/j.jhep.2016.05.037
28. Heyde S, Philipsen L, Formaglio P, Fu Y, Baars I, Hobbel G, Kleinholz CL, Seiss EA, Stettin J, Gintschel P, Dudeck A, Bouso P, Schraven B, Muller AJ (2018) CD11c-expressing Ly6C+CCR2+ monocytes constitute a reservoir for efficient Leishmania proliferation and cell-to-cell transmission. *PLoS Pathog* 14 (10):e1007374. doi:10.1371/journal.ppat.1007374



29. Romano A, Carneiro MBH, Doria NA, Roma EH, Ribeiro-Gomes FL, Inbar E, Lee SH, Mendez J, Paun A, Sacks DL, Peters NC (2017) Divergent roles for Ly6C+CCR2+CX3CR1+ inflammatory monocytes during primary or secondary infection of the skin with the intra-phagosomal pathogen *Leishmania major*. *PLoS Pathog* 13 (6):e1006479. doi:10.1371/journal.ppat.1006479
30. Terrazas C, Varikuti S, Oghumu S, Steinkamp HM, Ardic N, Kimble J, Nakhasi H, Satoskar AR (2017) Ly6C(hi) inflammatory monocytes promote susceptibility to *Leishmania donovani* infection. *Scientific reports* 7 (1):14693. doi:10.1038/s41598-017-14935-3
31. Pinto AI, Brown N, Preham O, Doehl JSP, Ashwin H, Kaye PM (2017) TNF signalling drives expansion of bone marrow CD4+ T cells responsible for HSC exhaustion in experimental visceral leishmaniasis. *PLoS Pathog* 13 (7):e1006465. doi:10.1371/journal.ppat.1006465
32. Abidin BM, Hammami A, Stager S, Heinonen KM (2017) Infection-adapted emergency hematopoiesis promotes visceral leishmaniasis. *PLoS Pathog* 13 (8):e1006422. doi:10.1371/journal.ppat.1006422
33. Kaye PM, Beattie L (2016) Lessons from other diseases: granulomatous inflammation in leishmaniasis. *Seminars in immunopathology* 38 (2):249-260. doi:10.1007/s00281-015-0548-7
34. Angarano G, Maggi P, Rollo MA, Larocca AM, Quarto M, Scalone A, Gradoni L (1998) Diffuse necrotic hepatic lesions due to visceral leishmaniasis in AIDS. *The Journal of infection* 36 (2):167-169. doi:10.1016/s0163-4453(98)80007-8
35. Yazici P, Yeniay L, Aydin U, Tasbakan M, Ozutemiz O, Yilmaz R (2008) Visceral leishmaniasis as a rare cause of granulomatous hepatitis: a case report. *Turkiye parazitolojii dergisi* 32 (1):12-15
36. Boussoffara T, Boubaker MS, Ben Ahmed M, Mokni M, Guizani I, Ben Salah A, Louzir H (2019) Histological and immunological differences between zoonotic cutaneous leishmaniasis due to *Leishmania major* and sporadic cutaneous leishmaniasis due to *Leishmania infantum*. *Parasite (Paris, France)* 26:9. doi:10.1051/parasite/2019007
37. Sandoval Pacheco CM, Araujo Flores GV, Favero Ferreira A, Sosa Ochoa W, Ribeiro da Matta VL, Zuniga Valeriano C, Pereira Corbett CE, Dalastra Laurenti M (2018) Histopathological features of skin lesions in patients affected by non-ulcerated or atypical cutaneous leishmaniasis in Honduras, Central America. *Int J Exp Pathol* 99 (5):249-257. doi:10.1111/iep.12295
38. Thilakarathne IK, Ratnayake P, Vithanage A, Sugathadasa DP (2019) Role of Histopathology in the Diagnosis of Cutaneous Leishmaniasis: A Case-Control Study in Sri Lanka. *The American Journal of dermatopathology* 41 (8):566-570. doi:10.1097/dad.0000000000001367
39. Grimaldi G, Jr., Teva A, Porrozzio R, Pinto MA, Marchevsky RS, Rocha MG, Dutra MS, Bruna-Romero O, Fernandes AP, Gazzinelli RT (2014) Clinical and parasitological protection in a *Leishmania infantum*-macaque model vaccinated with adenovirus and the recombinant A2 antigen. *PLoS Negl Trop Dis* 8 (6):e2853. doi:10.1371/journal.pntd.0002853
40. Khamesipour A, Dowlati Y, Asilian A, Hashemi-Fesharki R, Javadi A, Noazin S, Modabber F (2005) Leishmanization: use of an old method for evaluation of candidate vaccines against leishmaniasis. *Vaccine* 23 (28):3642-3648. doi:10.1016/j.vaccine.2005.02.015
41. Mebius RE, Kraal G (2005) Structure and function of the spleen. *Nat Rev Immunol* 5 (8):606-616. doi:10.1038/nri1669
42. Kaye PM (2018) Stromal Cell Responses in Infection. *Advances in experimental medicine and biology* 1060:23-36. doi:10.1007/978-3-319-78127-3\_2
43. Stanley AC, Engwerda CR (2007) Balancing immunity and pathology in visceral leishmaniasis. *Immunology and cell biology* 85 (2):138-147. doi:10.1038/sj.icb7100011
44. Hermida MD, de Melo CVB, Lima IDS, Oliveira GGS, Dos-Santos WLC (2018) Histological Disorganization of Spleen Compartments and Severe Visceral Leishmaniasis. *Frontiers in cellular and infection microbiology* 8:394. doi:10.3389/fcimb.2018.00394
45. Gardinassi LG, Garcia GR, Costa CH, Costa Silva V, de Miranda Santos IK (2016) Blood Transcriptional Profiling Reveals Immunological Signatures of Distinct States of Infection of Humans with *Leishmania infantum*. *PLoS Negl Trop Dis* 10 (11):e0005123. doi:10.1371/journal.pntd.0005123
46. Fakiola M, Singh OP, Syn G, Singh T, Singh B, Chakravarty J, Sundar S, Blackwell JM (2019) Transcriptional blood signatures for active and amphotericin B treated visceral leishmaniasis in India. *PLoS Negl Trop Dis* 13 (8):e0007673. doi:10.1371/journal.pntd.0007673
47. Ashwin H, Seifert K, Forrester S, Brown N, MacDonald S, James S, Lagos D, Timmis J, Mottram JC, Croft SL, Kaye PM (2018) Tissue and host species-specific transcriptional changes in models of

experimental visceral leishmaniasis. *Wellcome open research* 3:135. doi:10.12688/wellcomeopenres.14867.2

48. Kong F, Saldarriaga OA, Spratt H, Osorio EY, Travi BL, Luxon BA, Melby PC (2017) Transcriptional Profiling in Experimental Visceral Leishmaniasis Reveals a Broad Splenic Inflammatory Environment that Conditions Macrophages toward a Disease-Promoting Phenotype. *PLoS Pathog* 13 (1):e1006165. doi:10.1371/journal.ppat.1006165

49. Loria-Cervera EN, Andrade-Narvaez FJ (2014) Animal models for the study of leishmaniasis immunology. *Rev Inst Med Trop Sao Paulo* 56 (1):1-11. doi:10.1590/s0036-46652014000100001

50. Carvalho LP, Passos S, Schriefer A, Carvalho EM (2012) Protective and pathologic immune responses in human tegumentary leishmaniasis. *Front Immunol* 3:301. doi:10.3389/fimmu.2012.00301

51. Ives A, Ronet C, Prevel F, Ruzzante G, Fuertes-Marraco S, Schutz F, Zangger H, Revaz-Breton M, Lye LF, Hickerson SM, Beverley SM, Acha-Orbea H, Launois P, Fasel N, Masina S (2011) Leishmania RNA virus controls the severity of mucocutaneous leishmaniasis. *Science (New York, NY)* 331 (6018):775-778. doi:10.1126/science.1199326

52. Novais FO, Carvalho AM, Clark ML, Carvalho LP, Beiting DP, Brodsky IE, Carvalho EM, Scott P (2017) CD8+ T cell cytotoxicity mediates pathology in the skin by inflammasome activation and IL-1 $\beta$  production. *PLoS Pathog* 13 (2):e1006196. doi:10.1371/journal.ppat.1006196

53. Rossi M, Castiglioni P, Hartley MA, Eren RO, Prevel F, Desponds C, Utzschneider DT, Zehn D, Cusi MG, Kuhlmann FM, Beverley SM, Ronet C, Fasel N (2017) Type I interferons induced by endogenous or exogenous viral infections promote metastasis and relapse of leishmaniasis. *Proceedings of the National Academy of Sciences of the United States of America* 114 (19):4987-4992. doi:10.1073/pnas.1621447114

54. Alvar J, Croft SL, Kaye P, Khamesipour A, Sundar S, Reed SG (2013) Case study for a vaccine against leishmaniasis. *Vaccine* 31 Suppl 2:B244-249. doi:10.1016/j.vaccine.2012.11.080

55. Cecilio P, Perez-Cabezas B, Fernandez L, Moreno J, Carrillo E, Requena JM, Fichera E, Reed SG, Coler RN, Kamhawi S, Oliveira F, Valenzuela JG, Gradoni L, Glueck R, Gupta G, Cordeiro-da-Silva A (2017) Pre-clinical antigenicity studies of an innovative multivalent vaccine for human visceral leishmaniasis. *PLoS Negl Trop Dis* 11 (11):e0005951. doi:10.1371/journal.pntd.0005951

56. Das S, Freier A, Boussoffara T, Das S, Oswald D, Losch FO, Selka M, Sacerdoti-Sierra N, Schonian G, Wiesmuller KH, Seifert K, Schroff M, Juhls C, Jaffe CL, Roy S, Das P, Louzir H, Croft SL, Modabber F, Walden P (2014) Modular multiantigen T cell epitope-enriched DNA vaccine against human leishmaniasis. *Science translational medicine* 6 (234):234ra256. doi:10.1126/scitranslmed.3008222

57. Osman M, Mistry A, Keding A, Gabe R, Cook E, Forrester S, Wiggins R, Di Marco S, Colloca S, Siani L, Cortese R, Smith DF, Aebischer T, Kaye PM, Lacey CJ (2017) A third generation vaccine for human visceral leishmaniasis and post kala azar dermal leishmaniasis: First-in-human trial of ChAd63-KH. *PLOS Neglected Tropical Diseases* 11 (5):e0005527. doi:10.1371/journal.pntd.0005527

58. Gannavaram S, Torcivia J, Gasparyan L, Kaul A, Ismail N, Simonyan V, Nakhasi HL (2017) Whole genome sequencing of live attenuated *Leishmania donovani* parasites reveals novel biomarkers of attenuation and enables product characterization. *Scientific reports* 7 (1):4718. doi:10.1038/s41598-017-05088-4

59. Maroof A, Brown N, Smith B, Hodgkinson MR, Maxwell A, Losch FO, Fritz U, Walden P, Lacey CN, Smith DF, Aebischer T, Kaye PM (2012) Therapeutic vaccination with recombinant adenovirus reduces splenic parasite burden in experimental visceral leishmaniasis. *The Journal of infectious diseases* 205 (5):853-863. doi:10.1093/infdis/jir842

60. Toepf A, Larson M, Wilson G, Grinnage-Pulley T, Bennett C, Leal-Lima A, Anderson B, Parrish M, Anderson M, Fowler H, Hinman J, Kontowicz E, Jefferies J, Beeman M, Buch J, Saucier J, Tyrrell P, Gharpure R, Cotter C, Petersen C (2018) Randomized, controlled, double-blinded field trial to assess *Leishmania* vaccine effectiveness as immunotherapy for canine leishmaniosis. *Vaccine* 36 (43):6433-6441. doi:10.1016/j.vaccine.2018.08.087

61. WHO (2016) Leishmaniasis in high-burden countries: an epidemiological update based on data reported in 2014 *WHO Weekly epidemiological record* 22 (91):285-296

62. Organization WECotCotLWH (2010) Control of the leishmaniasis: report of a meeting of the WHO Expert Committee on the Control of Leishmaniasis vol 949. World Health Organization, Geneva
63. Ghose AC, Haldar JP, Pal SC, Mishra BP, Mishra KK (1980) Serological investigations on Indian kala-azar. *Clin Exp Immunol* 40 (2):318-326
64. Boelaert M, Verdonck K, Menten J, Sunyoto T, van Griensven J, Chappuis F, Rijal S (2014) Rapid tests for the diagnosis of visceral leishmaniasis in patients with suspected disease. *The Cochrane database of systematic reviews* (6):Cd009135. doi:10.1002/14651858.CD009135.pub2
65. Chappuis F, Sundar S, Hailu A, Ghalib H, Rijal S, Peeling RW, Alvar J, Boelaert M (2007) Visceral leishmaniasis: what are the needs for diagnosis, treatment and control? *Nature Reviews Microbiology* 5 (11):873-882
66. Bhattacharyya T, Boelaert M, Miles MA (2013) Comparison of visceral leishmaniasis diagnostic antigens in African and Asian *Leishmania donovani* reveals extensive diversity and region-specific polymorphisms. *PLoS Negl Trop Dis* 7 (2):e2057. doi:10.1371/journal.pntd.0002057
67. Bhattacharyya T, Bowes DE, El-Safi S, Sundar S, Falconar AK, Singh OP, Kumar R, Ahmed O, Boelaert M, Miles MA (2014) Significantly lower anti-*Leishmania* IgG responses in Sudanese versus Indian visceral leishmaniasis. *PLoS Negl Trop Dis* 8 (2):e2675. doi:10.1371/journal.pntd.0002675
68. Patabhi S, Whittle J, Mohamath R, El-Safi S, Moulton GG, Guderian JA, Colombara D, Abdoon AO, Mukhtar MM, Mondal D, Esfandiari J, Kumar S, Chun P, Reed SG, Bhatia A (2010) Design, development and evaluation of rK28-based point-of-care tests for improving rapid diagnosis of visceral leishmaniasis. *PLoS Negl Trop Dis* 4 (9). doi:10.1371/journal.pntd.0000822
69. Bezuneh A, Mukhtar M, Abdoun A, Teferi T, Takele Y, Diro E, Jemaneh A, Shiferaw W, Wondimu H, Bhatia A, Howard RF, Ghalib H, Ireton GC, Hailu A, Reed SG (2014) Comparison of point-of-care tests for the rapid diagnosis of visceral leishmaniasis in East African patients. *The American journal of tropical medicine and hygiene* 91 (6):1109-1115. doi:10.4269/ajtmh.13-0759
70. Mukhtar M, Abdoun A, Ahmed AE, Ghalib H, Reed SG, Boelaert M, Menten J, Khair MM, Howard RF (2015) Diagnostic accuracy of rK28-based immunochromatographic rapid diagnostic tests for visceral leishmaniasis: a prospective clinical cohort study in Sudan. *Trans R Soc Trop Med Hyg* 109 (9):594-600. doi:10.1093/trstmh/trv060
71. Mukhtar M, Ali SS, Boshara SA, Albertini A, Monnerat S, Bessell P, Mori Y, Kubota Y, Ndung'u JM, Cruz I (2018) Sensitive and less invasive confirmatory diagnosis of visceral leishmaniasis in Sudan using loop-mediated isothermal amplification (LAMP). *PLoS Negl Trop Dis* 12 (2):e0006264. doi:10.1371/journal.pntd.0006264
72. Subramaniam KS, Austin V, Schocker NS, Montoya AL, Anderson MS, Ashmus RA, Mesri M, Al-Salem W, Almeida IC, Michael K, Acosta-Serrano A (2018) Anti-alpha-Gal antibodies detected by novel neoglycoproteins as a diagnostic tool for Old World cutaneous leishmaniasis caused by *Leishmania major*. *Parasitology* 145 (13):1758-1764. doi:10.1017/s0031182018000860
73. Monge-Maillou B, Norman FF, Cruz I, Alvar J, Lopez-Velez R (2014) Visceral leishmaniasis and HIV coinfection in the Mediterranean region. *PLoS Negl Trop Dis* 8 (8):e3021. doi:10.1371/journal.pntd.0003021
74. Diro E, Lynen L, Ritmeijer K, Boelaert M, Hailu A, van Griensven J (2014) Visceral Leishmaniasis and HIV coinfection in East Africa. *PLoS Negl Trop Dis* 8 (6):e2869. doi:10.1371/journal.pntd.0002869
75. Gidwani K, Picado A, Ostyn B, Singh SP, Kumar R, Khanal B, Lejon V, Chappuis F, Boelaert M, Sundar S (2011) Persistence of *Leishmania donovani* antibodies in past visceral leishmaniasis cases in India. *Clinical and vaccine immunology : CVI* 18 (2):346-348. doi:10.1128/cvi.00473-10
76. Bhattacharyya T, Ayandeh A, Falconar AK, Sundar S, El-Safi S, Gripenberg MA, Bowes DE, Thunissen C, Singh OP, Kumar R, Ahmed O, Eisa O, Saad A, Silva Pereira S, Boelaert M, Mertens P, Miles MA (2014) IgG1 as a potential biomarker of post-chemotherapeutic relapse in visceral leishmaniasis, and adaptation to a rapid diagnostic test. *PLoS Negl Trop Dis* 8 (10):e3273. doi:10.1371/journal.pntd.0003273
77. Marlais T, Bhattacharyya T, Singh OP, Mertens P, Gilman Q, Thunissen C, Hinckel BCB, Pearson C, Gardner BL, Ains S, de la Roche M, Hayes K, Hafezi H, Falconar AK, Eisa O, Saad A, Khanal B, Bhattarai NR, Rijal S, Boelaert M, El-Safi S, Sundar S, Miles MA (2018) Visceral Leishmaniasis IgG1 Rapid Monitoring of Cure vs. Relapse, and Potential for Diagnosis of Post Kala-

- Azar Dermal Leishmaniasis. *Frontiers in cellular and infection microbiology* 8:427. doi:10.3389/fcimb.2018.00427
78. Mollett G, Bremer Hinckel BC, Bhattacharyya T, Marlais T, Singh OP, Mertens P, Falconar AK, El-Safi S, Sundar S, Miles MA (2019) Detection of Immunoglobulin G1 Against rK39 Improves Monitoring of Treatment Outcomes in Visceral Leishmaniasis. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 69 (7):1130-1135. doi:10.1093/cid/ciy1062
79. Kohanteb J, Ardehali SM, Rezai HR (1987) Detection of *Leishmania donovani* soluble antigen and antibody in the urine of visceral leishmaniasis patients. *Trans R Soc Trop Med Hyg* 81 (4):578-580. doi:10.1016/0035-9203(87)90414-7
80. De Colmenares M, Portus M, Riera C, Gallego M, Aisa MJ, Torras S, Munoz C (1995) Short report: detection of 72-75-kD and 123-kD fractions of *Leishmania* antigen in urine of patients with visceral leishmaniasis. *The American journal of tropical medicine and hygiene* 52 (5):427-428. doi:10.4269/ajtmh.1995.52.427
81. Azazy AA, Chance ML, Devaney E (1997) A time-course study of circulating antigen and parasite-specific antibody in cotton rats infected with *Leishmania donovani*. *Ann Trop Med Parasitol* 91 (2):153-162. doi:10.1080/00034983.1997.11813125
82. Vallur AC, Tutterrow YL, Mohamath R, Pattabhi S, Hailu A, Abdoun AO, Ahmed AE, Mukhtar M, Salam MA, Almeida ML, Almeida RP, Mondal D, Albertini A, Ghalib H, Duthie MS, Reed SG (2015) Development and comparative evaluation of two antigen detection tests for Visceral Leishmaniasis. *BMC Infect Dis* 15:384. doi:10.1186/s12879-015-1125-3
83. Gao CH, Yang YT, Shi F, Wang JY, Steverding D, Wang X (2015) Development of an Immunochromatographic Test for Diagnosis of Visceral Leishmaniasis Based on Detection of a Circulating Antigen. *PLoS Negl Trop Dis* 9 (6):e0003902. doi:10.1371/journal.pntd.0003902
84. Cruz I, Albertini A, Barbeitas M, Arana B, Picado A, Ruiz-Postigo JA, Ndung'u JM (2019) Target Product Profile for a point-of-care diagnostic test for dermal leishmaniases. *Parasite epidemiology and control* 5:e00103. doi:10.1016/j.parepi.2019.e00103
85. Ben Salah A, Zaatour A, Gharbi A, Bettaieb J, Ghawar W, Khedher A Clinical evaluation of CL detect TM rapid test for cutaneous leishmaniasis: Performance characteristics when compared to smear microscopy at multiple test sites. In: *ASTMH, New Orleans, 2014*.
86. Bennis I, Verdonck K, El Khalfaoui N, Riyad M, Fellah H, Dujardin JC, Sahibi H, Bouhout S, Van der Auwera G, Boelaert M (2018) Accuracy of a Rapid Diagnostic Test Based on Antigen Detection for the Diagnosis of Cutaneous Leishmaniasis in Patients with Suggestive Skin Lesions in Morocco. *The American journal of tropical medicine and hygiene* 99 (3):716-722. doi:10.4269/ajtmh.18-0066
87. Vink MMT, Nahzat SM, Rahimi H, Buhler C, Ahmadi BA, Nader M, Zazai FR, Yousufzai AS, van Loenen M, Schallig H, Picado A, Cruz I (2018) Evaluation of point-of-care tests for cutaneous leishmaniasis diagnosis in Kabul, Afghanistan. *EBioMedicine* 37:453-460. doi:10.1016/j.ebiom.2018.10.063
88. Selvapandiyan A, Croft SL, Rijal S, Nakhasi HL, Ganguly NK (2019) Innovations for the elimination and control of visceral leishmaniasis. *PLoS Negl Trop Dis* 13 (9):e0007616. doi:10.1371/journal.pntd.0007616
89. WHO (2017) Integrating neglected tropical diseases into global health and development: fourth WHO report on neglected tropical diseases. World Health Organisation,
90. Rijal S, Sundar S, Mondal D, Das P, Alvar J, Boelaert M (2019) Eliminating visceral leishmaniasis in South Asia: the road ahead. *BMJ (Clinical research ed)* 364:k5224. doi:10.1136/bmj.k5224
91. Medley GF, Hollingsworth TD, Olliaro PL, Adams ER (2015) Health-seeking behaviour, diagnostics and transmission dynamics in the control of visceral leishmaniasis in the Indian subcontinent. *Nature* 528 (7580):S102-S108. doi:10.1038/nature16042
- [http://www.nature.com/nature/journal/v528/n7580\\_supp\\_custom/abs/nature16042.html#supplementary-information](http://www.nature.com/nature/journal/v528/n7580_supp_custom/abs/nature16042.html#supplementary-information)
92. Rijal S, Boelaert M, Regmi S, Karki BM, Jacquet D, Singh R, Chance ML, Chappuis F, Hommel M, Desjeux P, Van der Stuyt P, Le Ray D, Koirala S (2004) Evaluation of a urinary antigen-based

- latex agglutination test in the diagnosis of kala-azar in eastern Nepal. *Tropical medicine & international health* : TM & IH 9 (6):724-729. doi:10.1111/j.1365-3156.2004.01251.x
93. Salam MA, Khan MG, Mondal D (2011) Urine antigen detection by latex agglutination test for diagnosis and assessment of initial cure of visceral leishmaniasis. *Trans R Soc Trop Med Hyg* 105 (5):269-272. doi:10.1016/j.trstmh.2010.12.007
94. Attar ZJ, Chance ML, el-Safi S, Carney J, Azazy A, El-Hadi M, Dourado C, Hommel M (2001) Latex agglutination test for the detection of urinary antigens in visceral leishmaniasis. *Acta Trop* 78 (1):11-16. doi:10.1016/s0001-706x(00)00155-8
95. Picado A, Cruz I, Sampath R, Ndung'u JM (2019) An antigen detectin rapid diagnostic test (RDT) to acelerate control and elimination of visceral leishmaniasis. 11th European Congress on Tropical Medicine and International Health 113. doi:10.1093/trstmh/trz0
96. Molina R, Gradoni L, Alvar J (2003) HIV and the transmission of *Leishmania*. *Ann Trop Med Parasitol* 97 Suppl 1:29-45. doi:10.1179/000349803225002516
97. Molina R, Ghosh D, Carrillo E, Monnerat S, Bern C, Mondal D, Alvar J (2017) Infectivity of Post-Kala-azar Dermal Leishmaniasis Patients to Sand Flies: Revisiting a Proof of Concept in the Context of the Kala-azar Elimination Program in the Indian Subcontinent. *Clinical infectious diseases* : an official publication of the Infectious Diseases Society of America 65 (1):150-153. doi:10.1093/cid/cix245
98. Zijlstra EE (2019) Biomarkers in Post-kala-azar Dermal Leishmaniasis. *Frontiers in cellular and infection microbiology* 9:228. doi:10.3389/fcimb.2019.00228
99. Verma S, Avishek K, Sharma V, Negi NS, Ramesh V, Salotra P (2013) Application of loop-mediated isothermal amplification assay for the sensitive and rapid diagnosis of visceral leishmaniasis and post-kala-azar dermal leishmaniasis. *Diagnostic microbiology and infectious disease* 75 (4):390-395. doi:10.1016/j.diagmicrobio.2013.01.011
100. Silva-Barrios S, Stager S (2019) Hypergammaglobulinemia sustains the development of regulatory responses during chronic *Leishmania donovani* infection in mice. *Eur J Immunol* 49 (7):1082-1091. doi:10.1002/eji.201847917
101. da Silva Junior GB, Guardão Barros EJ, De Francesco Daher E (2014) Kidney involvement in leishmaniasis—a review. *The Brazilian Journal of Infectious Diseases* 18 (4):434-440. doi:<https://doi.org/10.1016/j.bjid.2013.11.013>
102. Cello JP, Day LW (2009) Idiopathic AIDS Enteropathy and Treatment of Gastrointestinal Opportunistic Pathogens. *Gastroenterology* 136 (6):1952-1965. doi:10.1053/j.gastro.2008.12.073
103. Aronson N, Herwaldt BL, Libman M, Pearson R, Lopez-Velez R, Weina P, Carvalho E, Ephros M, Jeronimo S, Magill A (2017) Diagnosis and Treatment of Leishmaniasis: Clinical Practice Guidelines by the Infectious Diseases Society of America (IDSA) and the American Society of Tropical Medicine and Hygiene (ASTMH). *The American journal of tropical medicine and hygiene* 96 (1):24-45. doi:10.4269/ajtmh.16-84256
104. González U, Pinart M, Reveiz L, Alvar J (2008) Interventions for Old World cutaneous leishmaniasis. *The Cochrane database of systematic reviews* (4):111. doi:10.1002/14651858.CD005067.pub3
105. González U, Pinart M, Rengifo-Pardo M, Macaya A, Alvar J, Tweed JA (2009) Interventions for American cutaneous and mucocutaneous leishmaniasis. *Cochrane Database Systematic Reviews* 15 (2 (CD004834)):1-171
106. DNDi (2019) DNDi - Disease projects - Portfolio. *Drugs for Neglected Diseases Initiative*. <https://www.dndi.org/diseases-projects/portfolio/>. Accessed 7/11/2019
- 2019
107. Khare S, Nagle AS, Biggart A, Lai YH, Liang F, Davis LC, Barnes SW, Mathison CJN, Myburgh E, Gao M-Y, Gillespie JR, Liu X, Tan JL, Stinson M, Rivera IC, Ballard J, Yeh V, Groessl T, Federe G, Koh HXY, Venable JD, Bursulaya B, Shapiro M, Mishra PK, Spraggon G, Brock A, Mottram JC, Buckner FS, Rao SPS, Wen BG, Walker JR, Tuntland T, Molteni V, Glynne RJ, Supek F (2016) Proteasome inhibition for treatment of leishmaniasis, Chagas disease and sleeping sickness. *Nature* 537:229. doi:10.1038/nature19339
- <https://www.nature.com/articles/nature19339#supplementary-information>

108. Thompson AM, O'Connor PD, Marshall AJ, Blaser A, Yardley V, Maes L, Gupta S, Launay D, Braillard S, Chatelain E, Wan B, Franzblau SG, Ma Z, Cooper CB, Denny WA (2018) Development of (6 R)-2-Nitro-6-[4-(trifluoromethoxy)phenoxy]-6,7-dihydro-5 H-imidazo[2,1- b][1,3]oxazine (DNDI-8219): A New Lead for Visceral Leishmaniasis. *J Med Chem* 61 (6):2329-2352. doi:10.1021/acs.jmedchem.7b01581
109. Van Bocxlaer K, Caridha D, Black C, Vesely B, Leed S, Sciotti RJ, Wijnant GJ, Yardley V, Braillard S, Mowbray CE, Ioset JR, Croft SL (2019) Novel benzoxaborole, nitroimidazole and aminopyrazoles with activity against experimental cutaneous leishmaniasis. *International journal for parasitology Drugs and drug resistance*. doi:10.1016/j.ijpddr.2019.02.002
110. Caridha D, Vesely B, van Bocxlaer K, Arana B, Mowbray CE, Rafati S, Uliana S, Reguera R, Kreishman-Deitrick M, Sciotti R, Buffet P, Croft SL (2019) Route map for the discovery and pre-clinical development of new drugs and treatments for cutaneous leishmaniasis. *International journal for parasitology Drugs and drug resistance*. doi:10.1016/j.ijpddr.2019.06.003
111. Yang JH, Bhargava P, McCloskey D, Mao N, Palsson BO, Collins JJ (2017) Antibiotic-Induced Changes to the Host Metabolic Environment Inhibit Drug Efficacy and Alter Immune Function. *Cell host & microbe* 22 (6):757-765.e753. doi:10.1016/j.chom.2017.10.020
112. Burza S, Mahajan R, Sinha PK, van Griensven J, Pandey K, Lima MA, Sanz MG, Sunyoto T, Kumar S, Mitra G, Kumar R, Verma N, Das P (2014) Visceral leishmaniasis and HIV co-infection in Bihar, India: long-term effectiveness and treatment outcomes with liposomal amphotericin B (AmBisome). *PLoS Negl Trop Dis* 8 (8):e3053. doi:10.1371/journal.pntd.0003053
113. Adam AAO, Dafalla MMM, Mohammed HA, Elamin MY, Younis BM, Elfaki ME, Musa AM, Elhassan AM, Khalil EAG (2014) Visceral leishmaniasis-hepatitis B/C coinfections: a rising necessity to triage patients for treatment. *Ann Saudi Med* 34 (2):143-146. doi:10.5144/0256-4947.2014.143
114. Hailu W, Weldegebreel T, Hurissa Z, Tafes H, Omollo R, Yifru S, Balasegaram M, Hailu A (2010) Safety and effectiveness of meglumine antimoniate in the treatment of Ethiopian visceral leishmaniasis patients with and without HIV co-infection. *Trans R Soc Trop Med Hyg* 104 (11):706-712. doi:10.1016/j.trstmh.2010.07.007
115. Mbui J, Olobo J, Omollo R, Solomos A, Kip AE, Kirigi G, Sagaki P, Kimutai R, Were L, Omollo T, Egondi TW, Wasunna M, Alvar J, Dorlo TPC, Alves F (2019) Pharmacokinetics, Safety, and Efficacy of an Allometric Miltefosine Regimen for the Treatment of Visceral Leishmaniasis in Eastern African Children: An Open-label, Phase II Clinical Trial. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 68 (9):1530-1538. doi:10.1093/cid/ciy747
116. Murray HW (2000) Suppression of posttreatment recurrence of experimental visceral Leishmaniasis in T-cell-deficient mice by oral Miltefosine. *Antimicrobial Agents and Chemotherapy* 44 (11):3235-3236
117. Murray HW, Delph-Etienne S (2000) Visceral leishmanicidal activity of hexadecylphosphocholine (miltefosine) in mice deficient in T cells and activated macrophage microbicidal mechanisms. *J Infect Dis* 181 (2):795-799
118. Escobar P, Yardley V, Croft SL (2001) Activities of hexadecylphosphocholine (miltefosine), AmBisome, and sodium stibogluconate (Pentostam) against *Leishmania donovani* in immunodeficient scid mice. (0066-4804 (Print))
119. Dorlo TP, Balasegaram M, Beijnen JH, de Vries PJ (2012) Miltefosine: a review of its pharmacology and therapeutic efficacy in the treatment of leishmaniasis. *Journal of Antimicrobial Chemotherapy* 67 (11):2576-2597
120. Cohen BE (2016) The Role of Signaling via Aqueous Pore Formation in Resistance Responses to Amphotericin B. *Antimicrobial Agents and Chemotherapy* 60 (9):5122-5129. doi:10.1128/AAC.00878-16
121. Murray HW, Berman JD, Davies CR, Saravia NG (2005) Advances in leishmaniasis. *The Lancet* 366 (9496):1577
122. Chavez-Galan L, Vesin D, Martinvalet D, Garcia I (2016) Low Dose BCG Infection as a Model for Macrophage Activation Maintaining Cell Viability *%J Journal of Immunology Research*. 2016:17. doi:10.1155/2016/4048235

123. Convit J, Ulrich M, Zerpa O, Borges R, Aranzazu N, Valera M, Villarroel H, Zapata Z, Tomedes I (2003) Immunotherapy of american cutaneous leishmaniasis in Venezuela during the period 1990-99. *Trans R Soc Trop Med Hyg* 97 (4):469-472. doi:10.1016/s0035-9203(03)90093-9
124. Musa AM, Khalil EAG, Mahgoub FAE, Elgawi SHH, Modabber F, Elkadaru AEMY, Aboud MH, Noazin S, Ghalib HW, El-Hassan AM, Group/Sudan TLR (2008) Immunochemotherapy of persistent post-kala-azar dermal leishmaniasis: a novel approach to treatment. *Transactions of The Royal Society of Tropical Medicine and Hygiene* 102 (1):58-63. doi:10.1016/j.trstmh.2007.08.006 %J Transactions of The Royal Society of Tropical Medicine and Hygiene
125. Cohen HA (1979) Induction of delayed-type sensitivity to *Leishmania* parasite in a case of leishmaniasis cutanea diffusa with BCG and cord-factor (Trehalose-6-6' dimycolate). *Acta dermato-venereologica* 59 (6):547-549. doi:10.2340/0001555559547549
126. Adinolfi LE, Bonventre PF, Vander Pas M, Eppstein DA (1985) Synergistic effect of glucantime and a liposome-encapsulated muramyl dipeptide analog in therapy of experimental visceral leishmaniasis. *Infect Immun* 48 (2):409-416
127. Badaro R, Johnson WD, Jr. (1993) The role of interferon-gamma in the treatment of visceral and diffuse cutaneous leishmaniasis. *The Journal of infectious diseases* 167 Suppl 1:S13-17. doi:10.1093/infdis/167.supplement\_1.s13
128. Badaro R, Nascimento C, Carvalho JS, Badaro F, Russo D, Ho JL, Reed SG, Johnson WD, Jr., Jones TC (1994) Granulocyte-macrophage colony-stimulating factor in combination with pentavalent antimony for the treatment of visceral Leishmaniasis. *European journal of clinical microbiology & infectious diseases* : official publication of the European Society of Clinical Microbiology 13 Suppl 2:S23-28. doi:10.1007/bf01973598
129. Almeida R, D'Oliveira A, Jr., Machado P, Bacellar O, Ko AI, de Jesus AR, Mobashery N, Brito Santos J, Carvalho EM (1999) Randomized, double-blind study of stibogluconate plus human granulocyte macrophage colony-stimulating factor versus stibogluconate alone in the treatment of cutaneous Leishmaniasis. *The Journal of infectious diseases* 180 (5):1735-1737. doi:10.1086/315082
130. Santos JB, de Jesus AR, Machado PR, Magalhaes A, Salgado K, Carvalho EM, Almeida RP (2004) Antimony plus recombinant human granulocyte-macrophage colony-stimulating factor applied topically in low doses enhances healing of cutaneous Leishmaniasis ulcers: a randomized, double-blind, placebo-controlled study. *The Journal of infectious diseases* 190 (10):1793-1796. doi:10.1086/424848
131. Murray HW (2005) Interleukin 10 receptor blockade--pentavalent antimony treatment in experimental visceral leishmaniasis. *Acta Trop* 93 (3):295-301. doi:10.1016/j.actatropica.2004.11.008
132. Flynn B, Wang V, Sacks DL, Seder RA, Verthelyi D (2005) Prevention and treatment of cutaneous leishmaniasis in primates by using synthetic type D/A oligodeoxynucleotides expressing CpG motifs. *Infect Immun* 73 (8):4948-4954. doi:10.1128/iai.73.8.4948-4954.2005
133. Verthelyi D, Ishii KJ, Gursel M, Takeshita F, Klinman DM (2001) Human peripheral blood cells differentially recognize and respond to two distinct CPG motifs. *Journal of immunology (Baltimore, Md : 1950)* 166 (4):2372-2377. doi:10.4049/jimmunol.166.4.2372
134. Buates S, Matlashewski G (1999) Treatment of experimental leishmaniasis with the immunomodulators imiquimod and S-28463: efficacy and mode of action. *J Infect Dis* 179:1485-1494
135. Arevalo I, Tulliano G, Quispe A, Spaeth G, Matlashewski G, Llanos-Cuentas A, Pollack H (2007) Role of imiquimod and parenteral meglumine antimoniate in the initial treatment of cutaneous leishmaniasis. *Clinical Infectious Diseases* 44 (12):1549-1554
136. Miranda-Verastegui C, Llanos-Cuentas A, Arevalo I, Ward BJ, Matlashewski G (2005) Randomized, double-blind clinical trial of topical imiquimod 5% with parenteral meglumine antimoniate in the treatment of cutaneous leishmaniasis in Peru. *Clinical Infectious Diseases* 40 (10):1395-1403
137. Firooz A, Khamesipour A, Ghoorchy MH, Nassiri-Kashani M, Eskandari SE, Khatami A, Hooshmand B, Gorouhi F, Rashighi-Firoozabadi M, Dowlati Y (2006) Imiquimod in combination with meglumine antimoniate for cutaneous leishmaniasis. *Archives of Dermatology* 142 (12):1575-1579
138. El Hajj R, Bou Youness H, Lachaud L, Bastien P, Masquefa C, Bonnet PA, El Hajj H, Khalifeh I (2018) EAPB0503: An Imiquimod analog with potent in vitro activity against cutaneous leishmaniasis

- caused by *Leishmania major* and *Leishmania tropica*. *PLoS Negl Trop Dis* 12 (11):e0006854. doi:10.1371/journal.pntd.0006854
139. Smith AC, Yardley V, Rhodes J, Croft SL (2000) Activity of the novel immunomodulatory compound tucareosol against experimental visceral leishmaniasis. *Antimicrob Agents Chemother* 44 (6):1494-1498. doi:10.1128/aac.44.6.1494-1498.2000
  140. Dalton JE, Kaye PM (2010) Immunomodulators: use in combined therapy against leishmaniasis. *Expert review of anti-infective therapy* 8 (7):739-742. doi:10.1586/eri.10.64
  141. Machado PR, Lessa H, Lessa M, Guimaraes LH, Bang H, Ho JL, Carvalho EM (2007) Oral pentoxifylline combined with pentavalent antimony: a randomized trial for mucosal leishmaniasis. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 44 (6):788-793. doi:10.1086/511643
  142. Brito G, Dourado M, Polari L, Celestino D, Carvalho LP, Queiroz A, Carvalho EM, Machado PR, Passos S (2014) Clinical and immunological outcome in cutaneous leishmaniasis patients treated with pentoxifylline. *The American journal of tropical medicine and hygiene* 90 (4):617-620. doi:10.4269/ajtmh.12-0729
  143. Brito G, Dourado M, Guimaraes LH, Meireles E, Schriefer A, de Carvalho EM, Machado PRL (2017) Oral Pentoxifylline Associated with Pentavalent Antimony: A Randomized Trial for Cutaneous Leishmaniasis. *The American journal of tropical medicine and hygiene* 96 (5):1155-1159. doi:10.4269/ajtmh.16-0435
  144. Faria DR, Gollob KJ, Barbosa J, Jr., Schriefer A, Machado PR, Lessa H, Carvalho LP, Romano-Silva MA, de Jesus AR, Carvalho EM, Dutra WO (2005) Decreased in situ expression of interleukin-10 receptor is correlated with the exacerbated inflammatory and cytotoxic responses observed in mucosal leishmaniasis. *Infect Immun* 73 (12):7853-7859. doi:10.1128/iai.73.12.7853-7859.2005
  145. Sadeghian G, Nilforoushzadeh MA (2006) Effect of combination therapy with systemic glucantime and pentoxifylline in the treatment of cutaneous leishmaniasis. *Int J Dermatol* 45 (7):819-821. doi:10.1111/j.1365-4632.2006.02867.x
  146. Taslimi Y, Zahedifard F, Rafati S (2018) Leishmaniasis and various immunotherapeutic approaches. *Parasitology* 145 (4):497-507. doi:10.1017/s003118201600216x
  147. Adriaenssen W, Dorlo TPC, Vanham G, Kestens L, Kaye PM, van Griensven J (2017) Immunomodulatory Therapy of Visceral Leishmaniasis in Human Immunodeficiency Virus-Coinfected Patients. *Front Immunol* 8:1943. doi:10.3389/fimmu.2017.01943
  148. Sakthianandeswaren A, Elso CM, Simpson K, Curtis JM, Kumar B, Speed TP, Handman E, Foote SJ (2005) The wound repair response controls outcome to cutaneous leishmaniasis. *Proceedings of the National Academy of Sciences of the United States of America* 102 (43):15551-15556
  149. Van Bocxlaer K, Yardley V, Murdan S, Croft SL (2016) Drug permeation and barrier damage in *Leishmania*-infected mouse skin. *The Journal of antimicrobial chemotherapy* 71 (6):1578-1585. doi:10.1093/jac/dkw012
  150. Wijnant GJ, Van Bocxlaer K, Fortes Francisco A, Yardley V, Harris A, Alavijeh M, Murdan S, Croft SL (2018) Local Skin Inflammation in Cutaneous Leishmaniasis as a Source of Variable Pharmacokinetics and Therapeutic Efficacy of Liposomal Amphotericin B. *Antimicrob Agents Chemother* 62 (10):e00631-00618. doi:10.1128/aac.00631-18
  151. Wijnant GJ, Van Bocxlaer K, Yardley V, Harris A, Alavijeh M, Silva-Pedrosa R, Antunes S, Mauricio I, Murdan S, Croft SL (2018) Comparative efficacy, toxicity and biodistribution of the liposomal amphotericin B formulations Fungisome((R)) and AmBisome((R)) in murine cutaneous leishmaniasis. *International journal for parasitology Drugs and drug resistance* 8 (2):223-228. doi:10.1016/j.ijpddr.2018.04.001
  152. Coombs GH, Wolf CR, Morrison VM, Craft JA (1990) Changes in hepatic xenobiotic-metabolizing enzymes in mouse liver following infection with *Leishmania donovani*. *Mol Biochem Parasitol* 41 (1):17-24. doi:10.1016/0166-6851(90)90092-z
  153. Mukhopadhyay R, Madhubala R (1994) Antileishmanial activity and modification of hepatic xenobiotic metabolizing enzymes in golden hamster by 2(3)-tert-butyl-4-hydroxyanisole following infection with *Leishmania donovani*. *Biochem Pharmacol* 47 (2):253-256. doi:10.1016/0006-2952(94)90014-0
  154. Castro MdM, Cossio A, Velasco C, Osorio L (2017) Risk factors for therapeutic failure to meglumine antimoniate and miltefosine in adults and children with cutaneous leishmaniasis in



Colombia: A cohort study. *PLOS Neglected Tropical Diseases* 11 (4):e0005515.

doi:10.1371/journal.pntd.0005515

155. Voak AA, Standing JF, Sepulveda N, Harris A, Croft SL, Seifert K (2018) Pharmacodynamics and cellular accumulation of amphotericin B and miltefosine in *Leishmania donovani*-infected primary macrophages. *The Journal of antimicrobial chemotherapy* 73 (5):1314-1323.

doi:10.1093/jac/dky014

156. Dartois V (2014) The path of anti-tuberculosis drugs: from blood to lesions to mycobacterial cells. *Nature reviews Microbiology* 12 (3):159-167. doi:10.1038/nrmicro3200

157. Bryant CE, Monie TP (2012) Mice, men and the relatives: cross-species studies underpin innate immunity. *Open biology* 2 (4):120015. doi:10.1098/rsob.120015

158. Zhao M, Lepak AJ, Andes DR (2016) Animal models in the pharmacokinetic/pharmacodynamic evaluation of antimicrobial agents. *Bioorg Med Chem* 24 (24):6390-6400.

doi:10.1016/j.bmc.2016.11.008

Figure 4




Figure 3





Figure 2



**VL diagnostic algorithm.** Source: National guidelines from Ethiopia, Kenya, Sudan and Uganda

\* Parasitological diagnosis may also be conducted in tertiary hospitals and research centres

\*\*National guidelines in Sudan also consider the use of rK28 RDTs

Figure 1

