

This is a repository copy of A missense mutation of Leu74Pro of OGR1 found in familial amelogenesis imperfecta actually causes the loss of the pH-sensing mechanism.

White Rose Research Online URL for this paper: https://eprints.whiterose.ac.uk/159875/

Version: Supplemental Material

Article:

Sato, K, Mogi, C, Mighell, AJ orcid.org/0000-0002-9624-6923 et al. (1 more author) (2020) A missense mutation of Leu74Pro of OGR1 found in familial amelogenesis imperfecta actually causes the loss of the pH-sensing mechanism. Biochemical and Biophysical Research Communications, 526 (4). pp. 920-926. ISSN 0006-291X

https://doi.org/10.1016/j.bbrc.2020.04.005

© 2020, Elsevier. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/.

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: https://creativecommons.org/licenses/

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/

Highlights

- A mutation at Leu74Pro of OGR1 found in amelogenesis imperfecta was characterized.
- Either a wild-type or a mutant OGR1 receptor was transfected in HEK-293 cells.
- Both receptors are distributed in plasma membranes.
- The wild-type receptor, but not the mutant receptor, was internalized by protons and Ni²⁺.
- Ca^{2+} response to ligands was lost in the mutant receptor-expressing cells.