



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/160045/>

Version: Accepted Version

Article:

Turk, F. and Waller, G. (2020) Is self-compassion relevant to the pathology and treatment of eating and body image concerns? A systematic review and meta-analysis. *Clinical Psychology Review*, 79. 101856. ISSN: 0272-7358

<https://doi.org/10.1016/j.cpr.2020.101856>

Article available under the terms of the CC-BY-NC-ND licence
(<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Is self-compassion relevant to the pathology and treatment of eating and body image concerns? A systematic review and meta-analysis

Fidan Turk^{a,*} and Glenn Waller^a

^aDepartment of Psychology, University of Sheffield, Sheffield

*Corresponding author: Fidan Turk, Department of Psychology, University of Sheffield, Cathedral Court, 1 Vicar Lane, Sheffield S1 2LT, UK

E-mail address: fturk1@sheffield.ac.uk

Telephone: +44-114-222-6568

Role of funding sources:

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Contributors:

Both authors contributed to the conceptualization and methodology. FT conducted the formal analysis, literature search, data management, and prepared the original draft. GW reviewed and edited drafts and supervised the process. Both authors approved the final version.

Conflict of interest:

The authors declare they have no conflict of interest.

Is self-compassion relevant to the pathology and treatment of eating and body image concerns? A systematic review and meta-analysis

Abstract

Eating disorders are severe mental health conditions, with substantial consequences for health and quality of life. Such disorders are strongly associated with body image concerns. It is important to consider treatments that might enhance our ability to treat such cases. Recently, there has been a growing body of research on self-compassion in relation to such problems. However, we are not yet clear about the extent of such effects, given the range of studies and methodologies used. Therefore, a systematic literature review was carried out using four key databases. Meta-analysis was used to reach conclusions about the size of the effects and moderators. Random-effects meta-analyses were conducted with 59 studies. Higher self-compassion was associated with lower eating pathology, reduced body image concerns, and greater positive body image, with medium to strong effect sizes (respectively, $r = -0.34$, $r = -0.45$, $r = 0.52$). Furthermore, self-compassion interventions for eating pathology and body image were effective, and superior to control groups (respectively, $g = 0.58$, $g = 0.39$). These findings support the role of self-compassion in understanding and addressing eating and body image concerns. Self-compassion appears to be an adaptive emotion regulation strategy in eating disorders and body image.

Key words: eating pathology, self-compassion, body image, meta-analysis, intervention

Is self-compassion relevant to the pathology and treatment of eating and body image concerns? A systematic review and meta-analysis

Eating disorders can be severe mental health conditions, associated with significant physical and psychological impairment. The lifetime prevalence of eating disorders varies across cultures, ranging from 0.2% to 13.1 % (Cho et al., 2007; Preti et al., 2009; Solmi, Hotopf, Hatch, Treasure, & Micali, 2016; Stice, Marti, & Rohde, 2013). The existing literature has identified a number of potential risk factors, with body image concerns as a key predictor (Polivy & Herman, 2002). Eating pathology and body image are critical elements of eating disorders (Waller & Mountford, 2015).

Various treatments are effective in the treatment of eating disorders (National Institute for Health and Care Excellence [NICE], 2017). Recent studies have reported a remission rate of about 45 to 70% (e.g., Couturier, Kimber & Szatmari, 2013; Fairburn et al., 2015; Södersten, Bergh, Leon, Brodin, & Zandian, 2017). However, as there are still individuals who do not achieve remission, treatment developments are still necessary. A key area suggested for such development is to address the affective component of eating disorders directly (e.g., Engel et al., 2013; Goss & Allan, 2014; Wonderlich et al., 2008). The evidence underpinning such an approach comes from a number of functional accounts of eating disorders (Haynos & Fruzzetti, 2011; Heatherton & Baumeister, 1991; Lavender et al., 2014; Leehr et al., 2015; McCarthy, 1990; McManus & Waller, 1995; Smyth et al., 2007), suggesting that disordered eating behaviours can be a coping response to negative emotions (e.g., suppress/block/distract).

Both escape theory (Heatherton & Baumeister, 1991) and affect regulation models (Lavender et al., 2014; McCarthy, 1990) assume that binge-purge episodes can serve the function of coping with negative affect, by producing short-term relief. A functional analysis of binge-eating by McManus and Waller (1995) suggested that binging acts to block negative

emotions. Similarly, the transactional model of emotion dysregulation in anorexia nervosa suggests that disordered eating behaviours (e.g., restriction, extreme exercise) might function to suppress negative emotions or distract from negative emotions (Haynos & Fruzzetti, 2011; Smyth et al., 2007). Neurobiological theories are also relevant, as they emphasize the role of hormones and neurotransmitters in how negative emotions might trigger binge eating (Leehr et al., 2015). Thus, a range of theoretical models stress the critical role of emotions in eating disorders.

Self-compassion has been proposed to be an adaptive way of regulating such emotions (Gilbert, 2019; Neff, 2003a). Self-compassion is conceptualized by Neff (2003a) as understanding one's own pain in a non-judgmental way and seeing suffering as a part of a shared human experience. It involves three components: self-kindness (vs self-judgment); common-humanity (vs isolation); and mindfulness (vs over-identification) (Neff, 2003a). In contrast, Gilbert (2009) conceptualizes 'compassion' as "a sensitivity to suffering in self and others with a commitment to try to alleviate and prevent it" (p.13). In Gilbert's evolutionary theory, compassion consists of six elements - sensitivity, sympathy, empathy, motivation/caring, distress tolerance, and non-judgement.

Self-compassion might reduce the use of dysfunctional attempts to regulate emotions in eating disorders. Neff (2003a) suggested that self-compassionate individuals are less likely to experience their emotions adversely. With self-compassion, instead of escaping from negative emotions or pushing them away, those emotions are acknowledged as valid and important. Therefore, self-compassionate people are less likely to engage in avoidance/escape or suppression of emotions. Indeed, compassion training can activate brain areas that have been associated with positive emotions (Klimecki, Leiberg, Ricard, & Singer, 2014). Gilbert's model suggests that individuals who are less compassionate have difficulties generating and activating self-soothing emotions. These individuals are not able to regulate

threat-based emotions, and therefore they overeat as a way of calming emotional states.

Recent studies have also suggested that self-compassion can: buffer against risk factors for eating and body concerns (e.g., Ferreira, Matos, Duarte & Pinto-Gouvei, 2014; Stutts & Blomquist, 2018); prevent the initial occurrence of risk factors (e.g., Ferreira, Pinto, & Duarte, 2013; Marta-Simões, Ferreira, & Mendes, 2016); and reduce existing eating and body image problems (e.g., Breines, Toole, Tu & Chen, 2014; Kelly & Carter, 2015). However, the variability in treatment targets (e.g., body image, eating behaviours, eating attitudes) and methodologies means that it is difficult to draw clear conclusions from the literature. Braun, Park and Gorin (2016) undertook a systematic review of the role of self-compassion in eating and body image issues, and concluded that self-compassion may protect against eating and body image concerns. However, given the research that has emerged since then and the need to quantify the effects of self-compassion, it is appropriate to update that systematic review and to present a meta-analysis of the degree to which self-compassion is associated with and impacts on eating and body image concerns.

The first aim of this systematic review and meta-analysis was to investigate the association of self-compassion with levels of eating and body image concerns. The second aim was to determine whether self-compassion-related interventions are effective in enhancing eating and body image concerns. Based on the literature, it is hypothesized that:

1. Low self-compassion will be associated with high eating pathology
2. Self-compassion interventions will reduce eating pathology
3. Self-compassion will be positively associated with positive body image
4. Self-compassion will be negatively associated with body image concerns
5. Self-compassion interventions will positively enhance healthy body image.

Method

Identification and selection of studies

A literature search was conducted (up to 19th December 2019), using four electronic databases (PsycINFO, PubMed, ProQuest, Web of Science). The search terms (see Table 1) were used in a three-component strategy (Body Experience Terms; Compassion Terms; Eating and Body Terms). Additionally, the reference lists of identified studies were manually screened and previous reviews of self-compassion in the context of eating and body concerns were searched, in order to identify any other relevant studies.

Eligibility criteria

Studies were included if: (i) they used quantitative designs (e.g., experimental, correlational); (ii) they were written in English; and (iii) they were published in peer-reviewed journals. Papers for the correlational component of the analyses had to have used a validated or standardised measure of self-compassion and eating and/or body image variables.

Eating pathology refers to unhealthy eating attitudes and behaviours (e.g., bingeing and purging), and it can be measured by a number of validated tools (e.g., Eating Disorder Examination - Questionnaire [EDE-Q; Fairburn & Beglin, 1994]). Body image is the individual's subjective evaluation of their own physical appearance (Thompson, Heinbeg, Altabe & Tantleff-Dunn, 1999). Positive and negative body image appear to be distinct constructs, both dimensionally and qualitatively (Tylka & Wood-Barcalow, 2015). Positive body image involves protective attitudes and behaviours regarding one's own body (e.g., appreciation and acceptance - Tylka & Wood-Barcalow, 2015), whereas negative body image centres on negative subjective evaluation of one's physical body (Stice & Shaw, 2002).

Table 1

Research terms used for the literature review: “AND” was used to show that papers required having one term from each column, while “OR” was used to show that any of those keywords is enough for eligibility

(Body Experience Terms)	AND	(Compassion Terms)	AND	(Eating and Body Terms)
Objectification		Compassion		Eating
OR		OR		OR
Self-objectification		Self-compassion		Eating pathology
OR		OR		OR
Objectified		Self-warmth		Eating disorders
OR		OR		OR
Body surveillance		Self-kindness		Disordered eating
OR		OR		OR
Objectified body		Self-compassionate		Eating symptoms
OR				OR
Body preoccupation				Eating symptomatology
OR				OR
Body monitoring				Anorexia
OR				OR
Body checking				Bulimia
OR				OR
Body shame				Bulimic
OR				OR
Body appreciation				Binge
OR				OR
Body comparison				Bingeing
				OR
				Binge-eating
				OR
				Purging
				OR
				Purge
				OR
				Restriction
				OR
				Diet
				OR
				Restrained eating
				OR
				OSFED
				OR
				EDNOS
				OR
				Compulsive exercise
				OR
				Exercise
				OR
				Body dissatisfaction
				OR
				Body image

Intervention studies had to use an identified self-compassion intervention, and address eating and/or body image. No age or gender restriction was imposed. Both clinical and non-clinical samples were included. Papers were excluded if: they did not measure either self-compassion or eating and/or body image outcome variables; they primarily focused on body dysmorphic disorder; or the patients had identified neurological or psychotic disorders. Book chapters, reviews, and qualitative papers were also excluded. Finally, studies were excluded if they did not provide sufficient data to calculate effect sizes.

Data extraction

For the meta-analysis, reported correlations of self-compassion with eating pathology and body image were extracted to test hypothesis 1. To test hypotheses 3 and 4, positive body image variables (e.g., body appreciation or body satisfaction) and body image concerns (e.g., body shame or body dissatisfaction) were extracted from each paper where these were reported. The following data were also extracted from each study, to test for moderators - year of publication, design, and sample characteristics (gender, age, and clinical status and BMI).

To test hypotheses 2 and 5, outcome data from self-compassion interventions (means, standard deviations/standard errors and sample sizes) were extracted for eating pathology and body image. Studies were included if they used either the Self-Compassion Scale (SCS; Neff, 2003b) or the Self-Compassion Scale-Short Form (SCS-SF; Raes, Pommier, Neff, & Van Gucht, 2011), as they are the only validated measures of self-compassion. Any psychometrically validated measure of eating pathology or/and body image concerns was used.

Quality assessment

The quality of included studies was assessed, in order to inform the critique of current

literature and to inform areas for future directions (rather than to remove any studies from the review). Methodological quality was examined using the Effective Public Health Practice Project (EPHPP) assessment tool for quantitative studies. The EPHPP has demonstrated good inter-rater reliability and construct validity (Thomas, Ciliska, Dobbins, & Micucci, 2004). All studies were checked for a range of components - selection bias, study design, confounders, blinding, data collection method, and attrition. Each component was rated as weak, moderate, strong, or not applicable. Each study was given an overall score based on the number of components meeting criteria. A paper rating of ‘strong’ indicated no weak component ratings, ‘moderate’ indicated one weak component rating, and ‘weak’ indicated two or more weak component ratings).

Data analyses

Effect size coding. Effect sizes (r values) were directly obtained from correlational studies to examine the association between self-compassion and eating pathology and body image. Effect sizes were interpreted according to Cohen’s (1992) guidelines: $r = .10$ as small, $r = .30$ as medium, and $r = .50$ as large. In the case of one study (Barnett & Sharp, 2016 – study 2), the scoring of the effect size was reversed, as examination of the scoring criteria used suggested that it had been scored incorrectly in the original paper. Removal of this study did not reduce the overall effect size.

For intervention studies, when studies had both within-subject comparisons and between-subject comparisons, we included only the between-subject comparisons. A single group pre-post study shows the effect of an intervention on a group, whereas randomized and non-randomized group comparison studies indicate the effects of the intervention on the experimental group compared to the control group. Therefore, meta-analyses were conducted separately based on research design. Effect sizes were taken directly from the study, where they were provided. Otherwise, for single group pre-post studies, the following formula was

used to calculate the effect size (d):

$$d = \frac{|m_1 - m_2|}{\sqrt{s_1^2 + s_2^2 - (2rs_1s_2)}}$$

Where not provided in the studies, the value of ' r ' (correlation between the two conditions) was assumed conservatively to be 0.7 (Rosenthal, 1993). If studies reported only the Standard Error (SE) instead of Standard Deviation (S), S was obtained from SE by multiplying by the square root of the sample size.

For group comparison studies, effect sizes were calculated by using the following formulas¹. If the two groups had an equal number of participants

$$d = \frac{|m_1 - m_2|}{\sqrt{\frac{s_1^2 + s_2^2}{2}}}$$

Or, if the two groups differed in sample size;

$$d = \frac{|m_1 - m_2|}{\sqrt{\frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}}}$$

Effect sizes were calculated from first available post-intervention time point. Only one out of five group comparison studies for eating pathology had an active control group. Therefore, all such effect sizes were calculated based on the wait-list control group. Similarly, effect sizes were calculated using the wait-list control group for body image studies, due to only six studies having had active control groups.

Obtained Cohen's d effect sizes were converted to Hedge's g , to correct for small sample bias (Hedges & Olkin, 1985). Effect sizes were interpreted using Cohen's convention, where 0.2 is indicative of a small effect, 0.5 a moderate effect, and 0.8 a large effect (Cohen, 1992).

Independence of effect sizes. Some of the included studies reported multiple effect

¹ Effect sizes were calculated by using following website:
https://memory.psych.mun.ca/models/stats/effect_size.shtml

sizes for outcome measures. Multiple effect sizes within the same study violate the assumption of independence in meta-analytic modelling. Therefore, if studies reported more than one measure of eating and/or body concerns, we selected and used the primary key outcome measure. The most widely used self-report measures (e.g., EDE-Q for eating psychopathology) took precedence. For example, Kelly and Stephen (2016) reported The Body Appreciation Scale (BAS) and The Body Areas Satisfaction Scale as measures of body image. Most other included studies used the BAS (e.g., Andrew, Tiggemann, & Clark, 2016; Homan, & Tylka, 2015; Kelly, Miller, & Stephen, 2016). Therefore, the BAS was used, to provide consistency across studies. In cases where studies did not use those common outcome measures, we selected the measure that was most strongly related to self-compassion. For instance, Stutt and Blomquists (2018) reported eating pathology using nine items relating to compensatory behaviors in eating pathology, drawn from the ADD Health Survey (<http://www.cpc.unc.edu/projects/addhealth>) and eight items from the Eating Loss of Control Scale (ELOCS; Blomquist, Roberto, Barnes, White, Masheb, & Grilo, 2014). As self-compassion has been suggested as a healthy emotional regulation practice (Berking & Whitley, 2014), the ELOCS was chosen for meta-analysis as it is associated with greater emotion dysregulation (Blomquist et al., 2014).

Meta-analytic model. Meta-analysis was carried out using *Meta-Essentials* (Suurmond, van Rhee, & Hak, 2017). Due to a wide range of the characteristics of samples and the interventions included in this meta-analysis, a random-effects model was used to take into consideration between-study and within-study variance. Heterogeneity of effect sizes was examined using the Q -statistic with a p -value and I^2 statistic. The Q -statistic is a measure of variation around the average. The I^2 statistic indicates the proportion of total variance attributable to between-study variation. As benchmark values, 25%, 50%, and 75% were used to identify low, moderate and high heterogeneity, respectively (Higgins, Thompson, Deeks &

Altman, 2003).

Follow-up subgroup and moderator analyses were conducted to explore potential sources of heterogeneity. Subgroup analyses were used to examine categorical variables (e.g., the clinical status of participants, different measures of outcomes), while the characteristics of the included population samples (e.g., BMI, mean of age, percentage of female participants, quality of the studies) and duration of the body image interventions were investigated as potential moderators to explain between-study variation. Subgroup analyses were conducted if each subgroup had a minimum of four studies (Fu et al., 2011).

Publication bias. Meta-Essentials has multiple ways to examine publication bias. Funnel plots were used to indicate standard error (sample size) against reported effect size, where symmetrical distribution of samples shows the absence of publication bias. Additionally, a trim and fill imputation procedure was used to produce an estimate of the number of the studies missing due to publication bias (Duval & Tweedie, 2000).

Results

Study selection

The search initially identified 4333 publications. The search results from each database were imported into Mendeley reference management software. All studies were then combined and duplicates removed. After removing duplicates, 4036 studies remained. Having examined of abstracts of those studies, 200 studies were retained for further consideration. Of those 200, 133 were excluded (reasons outlined in Figure 1), leaving a total of 67 studies meeting the inclusion criteria for the review. Only 59 of those studies were included in this meta-analysis, due to their having sufficient data to calculate effect sizes.




Figure 1. PRISMA flowchart of inclusion of studies

Study characteristics

Table 2 presents an overview of the characteristics of the included studies. The sample size of those studies ranged from 9 to 1306. The majority used only a female sample ($n = 48$) or a predominantly female sample ($n = 5$). Fourteen studies recruited eating disorder patient samples (four assessed only binge-eating disorder patients; one targeted bulimic symptoms). Twenty-nine studies recruited student participants. The included studies came from a number of countries: USA ($n = 26$), Canada ($n = 13$), Portugal ($n = 13$), Australia ($n = 4$), UK ($n = 2$), and Thailand ($n = 1$). The majority of studies ($n = 43$) were published after 2014, supporting the need to update on the Braun et al. (2016) systematic review.

Table 2

Characteristics of included studies

Study	Design	Measure	Construct Measure	Sample
Correlational Studies				
<i>Eating Pathology and Self-compassion</i>				
Fresnics, Wag & Borders, 2019	Cross-sectional Correlational	<i>Self-Compassion Scale-Short Form</i> <i>Eating Disorder Examination Questionnaire</i> Freiburg Mindfulness Inventory Ruminative Response Scale	Self-compassion Eating pathology	190 female undergraduate
Pullmer, Zaitsoff & Coelho, 2019	Cross-sectional Correlational	<i>Self-Compassion Scale-Short Form</i> <i>Eating Disorder Examination Questionnaire</i> Hopkins Symptom Checklist	Self-compassion Eating pathology	58 female ED patient
Gouveia, Canavarro & Moreira, 2019	Cross-sectional Correlational	Child and Adolescent Mindfulness Measure <i>Self-compassion Scale short form</i> Difficulties in Emotion regulation Scale <i>Emotional eating subscale of the Dutch Eating Behaviour Questionnaire</i>	Self-compassion Eating pathology	245 Adolescent (120F/ 125M)
Kelly & Tasca, 2016	Longitudinal	<i>Self-Compassion Scale-Short Form</i> Experiences of Shame Scale <i>Eating Disorder Examination Questionnaire</i>	Self-compassion Eating pathology	78 ED patient (76F/ 2M)
Taylor, Daiss, & Krietsch, 2015	Cross-sectional Correlational	<i>Self-compassion scale short form</i> Mindful Eating Questionnaire <i>Eating Attitudes Test-26</i>	Self-compassion Eating pathology	150 college student (128F/ 22M)
Tylka, Russel, & Neal, 2015	Cross-sectional Correlational	Perceived Sociocultural Pressures Scale <i>Self-compassion Scale—Short Form</i> Internalization subscale of the Sociocultural Attitudes towards Appearance Questionnaire <i>Eating Attitudes Test-26</i>	Self-compassion Eating pathology	435 community women
Ferreira, Matos, Duarte, & Pinto-Gouveia, 2014	Cross-sectional Correlational	<i>Eating Disorder Examination 16.0D</i> Shame Experiences Interview Impact of Event Scale—Revised Centrality of Event Scale <i>Self-Compassion Scale</i>	Self-compassion Eating pathology	34 ED patient
Kelly & Carter, 2014	Longitudinal Correlational	<i>Eating Disorder Examination—Questionnaire</i> Experience of Shame Scale Social Safeness and Pleasure Scale Social Provisions Scale <i>Self-Compassion Scale-Short Form</i>	Self-compassion Eating pathology	89 ED patient (86 F/ 3 M)
Kelly, Carter, Zuroff & Borairi, 2013	Cross-sectional Correlational	<i>Eating Disorder Examination Questionnaire</i> Experiences of Shame Scale <i>Self-Compassion Scale-Short Form</i> Fear of self-compassion	Self-compassion Eating pathology	74 ED patient (72F/ 2M)
<i>Body Image and Self-compassion</i>				
Lonergan et al, 2019	Cross-sectional Correlational	Photo Manipulation and Investment Scales <i>Body Shape Satisfaction Scale</i> <i>Self-Compassion Scale Short-Form</i>	Body image Self-compassion	184 students (95F/ 89M)
Modica, 2019	Cross-sectional Correlational	Facebook use The Facebook Questionnaire Physical Appearance Comparison Scale Facebook Intensity Scale <i>Body Esteem Scale</i> <i>Body Surveillance subscale of Objectified Body Consciousness Scale</i> Appearance subscale of the Contingencies of Self-Worth Scale <i>Self-compassion Scale</i>	Body surveillance Self-compassion Body esteem	232 women
Schmidt, Raque-Bogdan & Hollern, 2019	Cross-sectional Correlational	<i>Self-Compassion Scale</i> <i>Body Appreciation Scale</i> <i>Multidimensional Body-self Relations Questionnaire</i> Body Image Quality of Life Inventory Positive and Negative Affect Schedule	Self-compassion Body appreciation	152 female undergraduate students

Barnett & Sharp, 2016 (Study, 1)	Cross-sectional Correlational	The Almost Perfect Scale – Revised Short Form <i>The Body Image Satisfaction Scale</i> <i>The Self-Compassion Scale</i>	Self-compassion Body satisfaction	580 female student
Marta-Simões, Ferreira, & Mendes, 2016	Cross-sectional Correlational	Other as Shamer Scale <i>Self-Compassion Scale</i> <i>Body Appreciation Scale</i>	Self-compassion Body appreciation	155 individuals (111 F/ 44 M)
Raque-Bogdan, Piontkowski, Hui, Ziemer & Garriott, 2016	Cross-sectional Correlational	Experiences in Close Relationships – Relationship Structures scale <i>Self-Compassion Scale</i> <i>Body Appreciation Scale</i>	Self-compassion Body appreciation	1306 female
Webb, Fiery, & Jafari, 2016	Cross-sectional Correlational	Anti-Fat Attitudes questionnaire The Fat Talk Scale <i>Objectified Body Consciousness Scale</i> <i>Self-Compassion Scale</i>	Self-compassion Body shame	309 undergraduate women
Andrew, Tiggeman & Clark, 2016	Cross-sectional Correlational	<i>Body Appreciation Scale</i> Media consumption Body Acceptance by Others Scale <i>Self-Compassion Scale-Short Form</i> Emotional Autonomy Scale <i>Body Surveillance Subscale of the Objectified Body Consciousness Scale</i> Physical Appearance Comparison Scale Internalization subscale of the Sociocultural Attitudes Towards Appearance Questionnaire	Self-compassion Body appreciation Body surveillance	266 undergraduate women
Homan & Tylka, 2015	Cross-sectional Correlational	<i>Self-Compassion Scale Short Form</i> The Body Comparison Orientation subscale from the Body, Eating, and Exercise Comparison Orientation Measure <i>Body Appreciation Scale</i> Contingencies of Self-Worth Scale	Self-compassion Body appreciation Body comparison	263 Female
Duarte, Ferreira, Trindade & Pinto-Gouveia, 2015	Cross-sectional Correlational	<i>Figure Rating Scale</i> Physical Appearance Scale <i>Self-Compassion Scale</i> World Health Organization Brief Quality of Life Assessment Scale	Self-compassion Body dissatisfaction	662 female student
Breines, Toole, Tu & Chen, 2014 (study 2)	Lab based*	State appearance-related self-compassion Objectified Body Consciousness scale Disordered eating scale adapted from Eisenberg and Neumark-Sztainer	Self-compassion Body surveillance Eating pathology	158 female undergraduate
Daye, Webb & Jafari, 2014	Cross-sectional Correlational	The Caregiver Eating Messages Scale <i>Objectified body consciousness</i> <i>Self-Compassion Scale</i>	Body shame Self-compassion	322 college women
Pinto-Gouveia, Ferreira & Duarte, 2014	Cross-sectional Correlational	Other as Shamer Scale Striving to Avoid Inferiority Scale Social Comparison through Physical Appearance Scale The Forms of Self-Criticizing and Self-Reassuring Scale <i>Self-Compassion Scale</i> <i>Body dissatisfaction scale of Eating Disorder Inventory</i> Eating Disorder Examination 16.OD	Self-compassion Body dissatisfaction	123 women non clinical 102 patient with ED
Pisitsungkagarn, Taephant & Attasaranya, 2014	Cross-sectional Correlational	<i>Body Appreciation Scale</i> Rosenberg's Self-esteem Scale <i>Self-compassion Scale</i>	Self-compassion Body appreciation	302 Thai female undergraduates
Wasylikiw, MacKinnon, & MacLennan, 2012 (Study 1)	Cross-sectional correlational	Rosenberg Self-Esteem Scale <i>Self-Compassion Scale</i> <i>Body Shape Questionnaire</i> <i>Body Appreciation Scale</i> Weight concern subscale from the Body Esteem Scale	Self-compassion Body appreciation Body dissatisfaction	142 female undergraduates
Mosewich et al, 2011	Cross-sectional Correlational	<i>Self-Compassion Scale</i> Rosenberg Self-Esteem Scale Self-Conscious Affect for Adolescents Social Physique Anxiety Scale Obligatory Exercise Questionnaire <i>Objectified body consciousness</i> Performance Failure Appraisal Inventory Fear of Negative Evaluation Scale	Self-compassion Body surveillance Body shame	151 women athletes

Eating Pathology and Body Image and Self-compassion

Kramer & Cuccolo, 2019	Intervention*	<i>Body Appreciation Scale-2</i> <i>Self-Compassion Scale</i> <i>Eating Disorder Examination Questionnaire</i> <i>Multidimensional Body-Self Relations Questionnaire</i> <i>Five Facet Mindfulness Questionnaire</i>	Self-compassion Eating pathology Body appreciation Body dissatisfaction	99 students (76 F/ 23M)
Pullmer, Coelho, & Zaitsoff, 2019	Cross-sectional Longitudinal	<i>Body Areas Satisfaction Scale</i> <i>Self-Compassion Scale</i> <i>Eating Disorder Examination Questionnaire</i> Hopkins Symptom Checklist	Self-compassion Body satisfaction Eating pathology	238 adolescent student (134 F/ 104 M)
Gouvenia, Canavarro, & Moreira, 2018	Cross-sectional Correlational	<i>Interpersonal Mindfulness in Parenting Scale</i> <i>Self-Compassion Scale</i> <i>Experience of Shame Scale</i> <i>Emotional Eating subscale of the Dutch Eating Behavior Questionnaire</i>	Self-compassion Eating pathology Body shame	572 dyads mother/father and adolescent (445 F/127 M)
Marta-Simões & Ferreira, 2018	Cross-sectional Correlational	<i>Early Memories of Warmth and Safeness Scale—Peers Version</i> <i>Self-Compassion Scale</i> <i>Social Safeness and Pleasure Scale</i> <i>Body Appreciation Scale-2</i> <i>Eating Disorder Examination Questionnaire</i>	Self-compassion Body appreciation Eating Pathology	387 women
Stutts & Blomquist, 2018	Longitudinal Correlational	<i>Self-Compassion Scale</i> <i>Weight concern and shape concern subscales of the Eating Disorder Examination Questionnaire</i> ADD Health Survey <i>Eating Loss of Control Scale</i> Perceived Stress Scale	Self-compassion Body dissatisfaction Eating pathology	765 student (535 F/ 230 M)
Ferreira, Oliveira, & Mendes, 2017	Cross-sectional Correlational	<i>Early Memories of Warmth and Safeness Scale</i> <i>Self-Compassion Scale</i> <i>Body Appreciation Scale</i> <i>Eating Disorder Examination Questionnaire</i>	Self-compassion Body appreciation Eating pathology	490 women general population
Barnett & Sharp, 2016 (Study 2)	Cross-sectional Correlational	<i>The Almost Perfect Scale – Revised Short Form</i> <i>The Body Image Satisfaction Scale</i> <i>Self-Compassion Scale – Short Form</i> <i>Eating Attitudes Test</i>	Self-compassion Body satisfaction Eating pathology	398 female students
Kelly, Miller, & Stephen, 2016	Daily surveys	<i>Self-Compassion Scale-Short Form</i> <i>Intuitive Eating Scale-2</i> <i>Body Appreciation Scale</i> <i>three items derived from the Eating Disorder Examination Questionnaire</i> <i>the Positive and Negative Affect Schedule</i>	Self-compassion Body appreciation Body dissatisfaction Eating attitudes	92 undergraduate female
Webb & Hardin, 2016	Cross-sectional Correlational	<i>Modified Weight Bias Internalization Scale</i> <i>Body Image Shame Scale</i> <i>Body Image Acceptance and Action Questionnaire</i> <i>Self-Compassion Scale</i> <i>Intuitive Eating Scale-2</i>	Self-compassion Body shame Body acceptance Eating attitudes	333 college women
Kelly & Stephen, 2016	Daily Dairy	<i>Self-compassion Scale Short form</i> <i>The Rosenberg Self-Esteem Inventory</i> <i>The Intuitive Eating Scale-2</i> <i>1 item of Restraint subscale of the Eating Disorders Examination Questionnaire</i> <i>The Body Appreciation Scale</i> <i>The Body Areas Satisfaction Scale</i> <i>Body Image States Scale</i>	Self-compassion Body appreciation Eating Pathology	92 female college students
Kelly, Vimalakanthan, & Miller, 2014	Cross-sectional Correlational	<i>Self-compassion scale</i> <i>Rosenberg Self-esteem inventory</i> <i>Body-Image Acceptance and Action Questionnaire</i> <i>EDE-Q</i>	Self-compassion Body acceptance Body dissatisfaction Eating Pathology	153 female student
Ferreira, Pinto-Gouveia, & Duarte, 2013	Cross-sectional correlational	<i>Self-Compassion Scale</i> <i>Other as Shamer Scale</i> <i>Depression, Anxiety and Stress Scales</i> <i>Eating Disorder Inventory</i> <i>Eating Disorder Examination 16.0D</i>	Self-compassion Eating pathology Body dissatisfaction	102 ED female patient 123 female general population
Schoenefeld & Webb, 2013	Cross-sectional Correlational	<i>Self-compassion Scale</i> <i>Distress Tolerance Scale</i> <i>Body Image-Acceptance and Action Questionnaire</i>	Self-compassion Eating attitudes Body acceptance	322 female undergraduate students

		Intuitive Eating Scale Rosenberg Self-Esteem Scale		
Wasyliw, MacKinnon, & MacLennan, 2012	Study 2: Cross-sectional correlational	Rosenberg Self-Esteem Scale <i>Self-Compassion Scale</i> <i>Body Shape Questionnaire</i> <i>Rigid Restraint Scale</i> The Center for Epidemiologic Studies Depression Scale	Self-compassion Body dissatisfaction Eating pathology	189 female undergraduates

Intervention Studies

Eating Pathology and Self-compassion

Kelly, Wisniewski, Martin-Wagar, & Hoffman, 2017	Group-Based CFT RCT- 12 weeks	Eating Disorder Examination Questionnaire Self-compassion Scale Fears of Compassion Scale Experience of Shame Scale	Self-compassion Eating pathology	22 ED patient
Williams, Tsivos, Brown, Whitelock, & Sampson, 2017	Retrospective study CFT intervention- 12 months	Eating Disorder Examination Questionnaire	Self-compassion Eating pathology	9 female
Kelly & Carter, 2015	CFT based self- compassion RCT- 3 weeks	Eating Disorder Examination Questionnaire Binge Eating Frequency Self-compassion Scale The Center of Epidemiological Studies for Depression Fears of Compassion Scale Credibility/Expectancy Questionnaire Homework Rating Scale	Self-compassion Eating pathology	41 individuals with BED
Gale, Gilbert, Read & Goss, 2014	Repeated measures design CFT intervention- 16 weeks	The Eating Disorder Examination Questionnaire The Stirling Eating Disorder Scale The Clinical Outcomes in Routine Evaluation – Outcome Measure	Eating Pathology Self-compassion	139 ED patient
Adams & Leary, 2007	Experimental manipulation of self- compassion	Revised Rigid Restraint Scale	Eating pathology	84 female student

Body Image and Self-compassion

Ziemer, Lamphere, Raque-Bogdan & Schmidt, 2019	Randomized controlled study of writing intervention- 3 weeks	Body Appreciation Scale 2 Body Image Quality of Life Inventory Positive and Negative Affect Schedule Self-Compassion Scale	Self-compassion Body image	152 female student
Moffitt, Neumann, & Williamson, 2018	Mixed experimental design- 3 minutes	Self-Compassion Scale Rosenberg Self-Esteem Scale Body Dissatisfaction subscale from the Eating Disorders Inventory Visual analogue Scales Self-improvement motivation	Self-compassion Body Dissatisfaction	153 female undergraduate
Rodgers et al., 2018	Bodimojo: grounded in self-compassion mobile- based intervention- 6 weeks	The Self-Compassion Scale Appearance Esteem subscale of the Body Esteem Scale Physical Appearance Comparison Scale Body Image-Acceptance and Action Questionnaire Positive and Negative Affect Schedule 10- Children	Self-compassion Body image	274 adolescent (71 Male/ 203 Female)
Stern & Engeln, 2018	Study 1: experimental manipulation- 15 min	Positive and Negative Affect Schedule Body Image States Scale	Body satisfaction	251 female undergraduate
	Study 2: experimental manipulation- 15 min	Positive and Negative Affect Schedule Body Image States Scale	Body satisfaction	240 undergraduate female
	Study 3: online intervention- 3 min	Positive and Negative Affect Schedule Body Image States Scale	Body satisfaction	158 sorority female
Seekis, Bradley & Duffy, 2017	Experimental manipulation- 5 min	State Body Appreciation Scale-2 Body Image States Scale The Physical Appearance State and Trait Anxiety Scale – state version	Body Image	96 female university student
Toole & Craighead, 2016	Online self-compassion intervention- 2 weeks	The Self-Compassion Scale The Body Appreciation Scale Rosenberg Self-Esteem Scale	Self-compassion Body	87 undergraduate women

		Body Surveillance subscale of the Objectified Body Consciousness Scale Body Shame subscale of the OBCS Contingencies of Self-Worth Scale-Appearance Subscale Body Shape Questionnaire	dissatisfaction Body surveillance Body shame	
Albertson, Neff & Dill-Shacklefor, 2015	RCT brief meditation intervention- 3 weeks	The Self-compassion Scale Body Shape Questionnaire Body Shame subscale of the Objectified Body Consciousness Scale Body Appreciation Scale The Contingencies of Self-Worth Scale	Self-compassion Body dissatisfaction Body shame Body image	228 adult women

Eating Pathology and Body Image and Self-compassion

Pinto-Gouveia et al, 2019	Non controlled BEfree intervention	Binge eating symptomatology (BES) EDE 16.0D The Acceptance and Action Questionnaire-II Cognitive Fusion Questionnaire-Body Image The Engage Living Scale Other as Shamer Scale Forms of Self-Criticising/Attacking & Self Reassuring Scale Self-Compassion Scale Five Facet Mindfulness Questionnaire-15	Self-compassion Eating Pathology Body Image	31 women with BED and overweight/ obese
Voelker, Petrie, Huang, & Chandran, 2019	Bodies in motion intervention- 4 weeks	Weight Pressures in Sport for Females Perceived Sociocultural Pressures Scale Sociocultural Attitudes Towards Appearance Questionnaire-4 Concerns about Weight and Concerns about Shape subscales from the EDE-Q Body Shame Scale Body Appreciation Scale-2 Body Parts Satisfaction Scale-Revised Positive and Negative Affect Schedule bulimic symptomatology score from nine items on the EDE-Q Dietary Intent Scale Frieberg Mindfulness Inventory-Short Form Self-Compassion Scale-Short Form	Self-compassion Body image Eating pathology	97 athletes
Kelly & Waring, 2018	Self-compassionate letter-writing intervention- 2 weeks	Self-Compassion Scale Fear of Compassion Scale Other as Shamer Scale Experience of Shame Scale Autonomous and Controlled Motivation for Treatment Questionnaire Readiness Ruler Eating Disorder Examination Questionnaire	Self-compassion Eating pathology Body shame	40 nontreatment seeking female
Duarte, Pinto-Gouveia & Stubbs, 2017	Low-intensity 4-week intervention	Eating Disorder Examination The Binge Eating Scale Body Image Shame Scale Depression, Anxiety, and Stress Scale Cognitive Fusion Questionnaire for food craving Body Image Acceptance and Action Questionnaire Five-Facet Mindfulness Questionnaire Compassionate Engagement and Action Scales Self-Compassion Scale Self-Criticism and Self-Reassurance Scale	Self-compassion Body image Eating pathology	20 BED women
Palmeira, Cunha & Pinto-Gouveia, 2017	Kg-free: an acceptance-mindfulness, compassion- Intervention-12 weeks	Weight Self-Stigma Questionnaire Obesity Related Well-Being Questionnaire Three Factor Eating Questionnaire-21R Acceptance and Action Questionnaire for Weight-Related Difficulties-Revised Other as Shamer Scale Self-Compassion Scale Five Facet Mindfulness Questionnaire	Self-compassion Eating pathology	53 overweight/ obese women
Pinto-Gouveia et al., 2017	Nonrandomized Controlled Longitudinal design-12 weeks	EDE 16.0D Binge Eating Scale Beck Depression Inventory-I Other as Shamer Scale Obesity-Related Well-Being Questionnaire	Self-compassion Body image Eating pathology	36 BED female overweight/ obese

Body Image-Acceptance and Action
Questionnaire
Cognitive Fusion Questionnaire-Body Image
The Engaged Living Scale
Self-Compassion Scale
Five Facet Mindfulness

Note. Measures in italic font indicate chosen measures for meta-analysis *: only correlational data used in meta-analysis

The majority of studies measured self-compassion using the Self-Compassion Scale (Neff, 2003b; $n = 35$), some used the Self-Compassion Scale-Short Form (Raes et al, 2011; $n = 14$), and the remainder did not measure self-compassion directly but used a self-compassion intervention ($n = 10$). The most commonly used measures of eating pathology were the EDE-Q ($n = 19$), the Eating Attitudes Test (Garner, Olmsted, Bohr, & Garfinkel, 1982; $n = 4$), and the Dutch Eating Behavior Questionnaire (Van Strien, Frijters, Bergers, & Defares, 1986, $n = 2$). In terms of body image, the majority of the studies used the Body Appreciation Scale (Avalos, Tylka, & Wood-Barcalow, 2005; $n = 12$) or the Body Appreciation Scale-2 (Tylka & Wood-Barcalow, 2015; $n = 5$). Other studies used the Objectified Body Consciousness Scale (McKinley & Hyde, 1996; $n = 8$), the Body Shape Questionnaire (Evans & Dolan, 1993; $n = 3$), or the Body Esteem Scale (Franzoi & Shields, 1984; $n = 3$).

Nature of the correlational studies. Of the 59 studies, 39 were correlational designs. Four of the 39 were longitudinal studies and two were daily diary studies. The remainder were correlational studies. Nine addressed only eating pathology, 16 addressed only body image, and 14 measured both.

Nature of intervention studies. The intervention studies identified included a variety of designs, such as randomized controlled trials, case series, nonrandomized control studies. Of the 20 studies examining the effectiveness of self-compassion interventions, 16 had a control group (three addressed eating pathology, nine body image, and four both). Only four studies were with a single group, examining the effectiveness of uncontrolled interventions

on eating pathology.

Of the 16 studies with control groups, a variety of treatment methods were used to address body image and eating pathology, including writing interventions, self-compassion meditation, and a number of proprietary programs (e.g., Bodies in Motion, BEFree). Of the four studies that lacked a comparison condition, four were delivered in a group format, and the other delivered compassion-focused therapy individually.

Quality assessment

Quality rating of studies is presented in Table 3. Attrition could not be considered in understanding the quality of cross-sectional correlational studies because there was only one data point (meaning that attrition could not take place). Confounders and blinding do not apply in non-intervention studies within the EPHPP assessment tool. Therefore, cross-sectional correlational studies were assessed using only the remaining three criteria. Of the 59 studies, only five received strong quality score ratings, 21 were rated as moderate, and 33 were rated as weak.

The main limitation across studies was selection bias. Most of the studies ($n = 37$) failed to recruit a representative sample, which limits the generalizability of the findings. The majority of the samples were based on young college women. However, body image issues and eating pathology are more prevalent among younger adult females (e.g., Pritchard, 2008), so that might not be a weakness per se. Another methodological weakness was study design. Thirty-nine studies used correlational designs, which preclude causal inferences. The major strength was the data collection method, as almost all studies ($n = 56$) used reliable and valid measures. However, the self-report questionnaires used in those studies are subject to response bias (Van de Mortel, 2008). Intervention studies most commonly received weak ratings due to the presence of confounders ($n = 7$). Regarding blinding, most intervention studies failed to state whether assessors were aware of the exposure status of participants.

However, participants were not aware of the research questions ($n = 15$).

Table 3

*Quality Assessment of Included Studies Using the EPHPP tool**

Author (date)	Component Rating						Overall Rating
	Selection Bias	Study Design	Confounders	Blinding	Data Collection Method	Attrition	
Correlational Studies							
<i>Eating Pathology and Self compassion</i>							
Fresnics, Wag & Borders, 2019	W	M	M	NA	S	M	Moderate
Pullmer, Zaitsoff & Coelho, 2019	M	W	NA	NA	S	NA	Moderate
Gouvenia, Canavarro & Moreira, 2019	M	W	NA	NA	S	NA	Moderate
Kelly & Tasca, 2016	M	M	NA	NA	S	M	Strong
Taylor, Daiss, & Krietsch, 2015	W	W	NA	NA	S	NA	Weak
Tylka, Russel, & Neal, 2015	W	W	NA	NA	S	NA	Weak
Ferreira, Matos, Duarte, & Pinto-Gouveia, 2014	M	W	NA	NA	S	NA	Moderate
Kelly & Carter, 2014	M	W	NA	NA	S	NA	Weak
Kelly, Carter, Zuroff & Borairi, 2013	M	M	NA	NA	S	M	Strong
<i>Body Image and Self-compassion</i>							
Lonergan et al, 2019	W	W	NA	NA	S	NA	Weak
Modica, 2019	W	W	NA	NA	M	NA	Weak
Schmidt, Raque-Bogdan & Hollern, 2019	W	W	NA	NA	S	NA	Weak
Barnett & Sharp, 2016 (Study 1)	W	W	NA	NA	M	NA	Weak
Marta-Simões, Ferreira, & Mendes, 2016	M	W	NA	NA	S	NA	Moderate
Raque-Bogdan, Piontkowski, Hui, Ziemer & Garriott, 2016	W	W	NA	NA	S	NA	Weak
Webb, Fiery, & Jafari, 2016	W	W	NA	NA	S	NA	Weak
Andrew, Tiggeman, & Clark, 2016	W	W	NA	NA	S	NA	Weak
Homan & Tylka, 2015	W	W	NA	NA	S	NA	Weak
Duarte, Ferreira, Trindade & Pinto-Gouveia, 2015	W	W	NA	NA	S	NA	Weak
Breines, Toole, Tu & Chen, 2014 (study 2)	W	M	W	M	S	NA	Weak
Daye, Webb, & Jafari, 2014	W	W	NA	NA	S	NA	Weak
Pinto-Gouveia, Ferreira, & Duarte, 2014	M	W	NA	NA	S	NA	Moderate

Pisitsungkagarn, Taephant, & Attasaranya, 2014	W	W	NA	NA	S	NA	Weak
Wasylikiw, MacKinnon, & MacLennan, 2012 (Study 1)	W	W	NA	NA	S	NA	Weak
Mosewich et al., 2011	M	W	NA	NA	S	NA	Moderate

Eating Pathology and Body Image and Self-compassion

Kramer & Cuccolo, 2019	W	M	M	NA	S	M	Moderate
Pullmer, Coelho, & Zaitsoff, 2019	M	M	NA	NA	S	S	Strong
Gouvenia, Canavarro, & Moreira, 2018	M	W	NA	NA	S	NA	Moderate
Marta-Simões & Ferreira, 2018	W	W	NA	NA	S	NA	Weak
Stutts & Blomquist, 2018	W	M	NA	NA	S	M	Moderate
Ferreira, Oliveira, & Mendes, 2017	W	W	NA	NA	S	NA	Weak
Barnett & Sharp, 2016 (Study 2)	W	W	NA	NA	S	NA	Weak
Kelly, Miller, & Stephen, 2016	W	W	NA	NA	S	NA	Weak
Webb & Hardin, 2016	W	W	NA	NA	S	NA	Weak
Kelly & Stephen, 2016	W	W	NA	NA	S	NA	Weak
Kelly, Vimalakanthan, & Miller, 2014	W	W	NA	NA	S	NA	Weak
Ferreira, Pinto-Gouveia, & Duarte, 2013	M	W	NA	NA	S	NA	Weak
Schoenefeld & Webb, 2013	W	W	NA	NA	S	NA	Weak
Wasylikiw, MacKinnon, & MacLennan, 2012 (Study 2)	W	W	NA	NA	S	NA	Weak

Interventional Studies

Eating Pathology and Self-compassion

Kelly, Wisniewski, Martin-Wagar, & Hoffman, 2017	M	S	S	M	S	W	Moderate
Williams et. al (2017)	M	M	W	W	S	W	Weak
Kelly & Carter (2015)	M	S	S	M	S	M	Strong
Gale, Gilbert, Read & Goss (2014)	M	M	W	W	S	M	Weak
Adams & Leary (2007)	W	M	W	M	S	NA	Weak

Body Image and Self-compassion

Ziemer, Lamphere, Raque-Bogdan & Schmidt, 2019	W	S	M	M	S	S	Moderate
Moffitt, Neumann, & Williamson, 2018	W	S	S	M	S	S	Moderate
Rodgers et al., 2018	M	S	M	M	S	S	Strong
Stern & Engeln, 2018 (Study 1)	W	S	M	M	S	S	Moderate
(Study 2)	W	S	M	M	S	S	Moderate

(Study 3)	W	S	W	M	S	M	Weak
Seekis, Bradley, & Duffy, 2017	W	S	S	M	S	M	Moderate
Toole & Craighead, 2016	W	S	S	M	S	S	Moderate
Albertson, Neff & Dill-Shacklefor, 2015	M	S	S	M	S	W	Moderate
<i>Eating Pathology and Body Image and Self-compassion</i>							
Pinto-Gouveia et al., 2019	M	M	M	M	S	W	Moderate
Voelker, Petrie, Huang, & Chandran, 2019	W	S	S	W	S	M	Weak
Kelly & Waring, 2018	W	S	S	M	S	W	Weak
Duarte, Pinto-Gouveia, & Stubbs, 2017	M	S	M	M	S	W	Moderate
Palmeira, Cunha, & Pinto-Gouveia, 2017	M	M	W	M	S	M	Moderate
Pinto-Gouveia et al., 2017	M	M	S	W	S	W	Weak

Note: *, EPHPP, Effective Public Health Practice Project. S: Strong, no weak component rating; M: Moderate, one weak component rating; W: Weak, two or more weak component ratings. NA: not applicable

Meta-analysis of association between eating pathology and self-compassion (hypothesis

1)

Figure 2 displays a forest plot for eating pathology, with each individual effect size representing a correlation between eating pathology and self-compassion. The combined random effects estimate for the association between eating pathology and self-compassion was $r = -0.34$ (95% CI = -0.40 to -0.28, $Z = -10.58$; $p < 0.001$), based on 5132 participants taken from 22 studies. This overall effect (r) was of a medium size (Cohen, 1992), showing that greater self-compassion is associated with lower levels of eating pathology. There was substantial variability across the studies ($Q = 82.81$, $p < 0.001$), with a high level of heterogeneity ($I^2 = 74.64\%$).

Subgroup analysis and meta-regression results are presented in Table 4. The correlation between self-compassion and eating pathology was not significantly affected by the type of self-compassion measure ($p = 0.19$). The effect size for the eating disorder group ($r = -0.49$) tended to be larger than for the non-clinical group ($r = -0.31$). However, a note of caution is necessary here, since each subgroup has high heterogeneity.




Figure 2. Forest plot of the correlations between self-compassion and eating pathology

Table 4

Subgroup and meta-regression analysis of relationship between eating pathology and self-compassion

	Number of Studies	Correlation (<i>r</i>)	95% CI	I ²	<i>p</i>
Clinical status					.04*
ED	5	-0.49*	-0.69 to -0.23	62.92	
Non ED	17	-0.31*	-0.37 to -0.25	73.58	
Independent measure					0.19
SCS	10	-0.31*	-0.38 to -0.23	59.56	
SCS-SF	12	-0.38*	-0.48 to -0.26	80.57	
Moderator	Number of Studies	B-Coefficient	95% CI	SE	<i>p</i>
Age (mean)	19	-0.10	-0.01 to 0.01	0.01	.65
Percentage of female participants	21	-0.03	-0.55 to 0.47	0.24	.88
BMI (mean)	13	-0.16	-0.04 to 0.03	0.01	.56
Study Quality Rating (0-3 criteria)	22	-0.53	-0.20 to -0.03	0.04	.005*

Note. *significant at *p* < .05, BMI: Body Mass Index; CI: confidence interval; DEBQ: Dutch Eating Behavior Questionnaire; EAT; Eating Attitudes Test-26; EDE-Q: Eating Disorder Examination Questionnaire; ED: eating disorder group; SE; standard error; SCS: Self-Compassion Scale, SCS-SF: Self-compassion Scale-Short Form

Medium to large heterogeneity was observed in the subgroups. Moderating effects for age, gender, and BMI were non-significant. Therefore, the relationship between self-compassion and eating pathology did not differ according to participants' age, gender, or BMI. Only the quality of the studies was associated with the relationship between self-compassion and eating pathology. Self-compassion was more strongly linked to eating pathology in the higher-quality studies.

The funnel plot (see Figure 3) suggested a slightly asymmetric distribution of study findings. The blue dots indicate the observed studies, and the green circles show the missing studies imputed by the trim-and-fill method. Trim and fill bias analysis imputed two studies. When the meta-analysis was adjusted for this potential bias, the new effect size slightly reduced ($r = -0.33$, 95% CI -0.42 to -0.25).




Figure 3. Funnel plot for the relationship between self-compassion and eating pathology. CES: Combined Effect Size.

Meta-analysis of effects of self-compassion interventions on eating pathology (hypothesis 2)

Studies with no comparison group. The effects of self-compassion intervention

studies with a single group from baseline to post-intervention were examined. There were four studies with a total sample size of 187. The combined effect size in this group was large and significant ($g = 1.30$, 95 % CI = 0.25 to 0.50; $Z = 5.18$; $p < 0.001$), showing that self-compassion interventions reduced eating pathology substantially from baseline to post-intervention (see Figure 4). Between-study variance was large and significant ($Q = 20.10$, $p < 0.001$, $I^2 = 85.15\%$). There were too few studies to test moderators or publication bias.




Figure 4. Forest plot showing the effect of the self-compassion interventions on eating pathology in studies with no comparison group

Studies with a comparison group. The effects of the self-compassion-related intervention on eating pathology were compared with a control group in six studies ($N = 153$ in self-compassion interventions; $N = 133$ in control groups). Self-compassion interventions had a significant impact on eating pathology compared to the control conditions (only Kelly et al. [2007] had an active control group, in the form of treatment as usual), with an effect size of 0.58 (95 % CI = 0.16 to 0.20; $Z = 3.22$; $p < 0.001$). The distribution of effect sizes is displayed in Figure 5. I^2 indicates that 40.48 % of the variation was attributable to between-study variance ($Q = 10.08$ $p = 0.12$). There were insufficient studies to examine publication bias.




Figure 5. Forest plot showing effect sizes for changes in eating pathology compared to the control group

Summary. Overall, higher levels of self-compassion were associated with lower levels of eating pathology, with a medium effect size. Similarly, self-compassion-related interventions are effective in reducing eating pathology compared to controls, again with a medium effect size.

Meta-analysis of association of positive body image and self-compassion (hypothesis 3)

Twenty studies examined the relationship between self-compassion and positive body image variables, including a total of 6230 participants. Figure 6 shows that there was a large combined effect for the association between higher self-compassion and greater positive body image, $r = 0.52$, (95% CI = 0.46 to 0.57, $Z = 16.28$; $p < 0.001$). Across the studies, heterogeneity was significant and large ($Q = 115.66$, $p < 0.001$, $I^2 = 83.57\%$).




Figure 6. Forest plot for the correlation between positive body image and self-compassion

Subgroup analysis examined the association between the effect sizes and the different measures of self-compassion. The differences in effect sizes between SCS ($r = 0.52$) and SCS-SF ($r = 0.51$) was not significant ($p = 0.82$; see Table 5). However, that conclusion should be treated with caution, because these subgroups are not sufficiently homogenous.

The results of the meta-regression analyses indicate that gender and quality of studies were not significant predictors of effect size. However, the moderating effects of age and

BMI were significant, indicating that the association between positive body image and self-compassion is higher in individuals with higher BMI and a greater age (see Table 5).

Table 5 - Subgroup and meta-regression analysis of relationship between positive body image and self-compassion

	Number of Studies	Correlation (<i>r</i>)	95% CI	<i>I</i> ²	<i>p</i>
Independent Measure					0.82
SCS	14	0.52*	0.46 to 0.58	83.56	
SCS-SF	6	0.51*	0.33 to 0.65	85.80	
Moderator	Number of Studies	B-Coefficient	95% CI	SE	<i>p</i>
Age (mean)	18	0.47	0.00 to 0.02	0.01	0.03
Percentage of female participants	20	-0.5	-0.65 to 0.55	0.29	0.86
BMI (mean)	13	0.71	0.02 to 0.09	0.01	0.01
Study Quality Rating (0-3 criteria)	20	0.24	-0.06 to 0.20	0.06	0.27

Note. BMI: Body Mass Index; CI: confidence interval; SE: standard error, SCS: Self-Compassion Scale; SCS-SF: Self-compassion Scale-Short Form

Considering publication bias, the inspection of the funnel plot suggests symmetric distribution of results (see Figure 7). After adjustment for omitted studies ($n = 1$), the effect size changed slightly to an r of 0.55 (95% CI 0.47 to 0.63).

Meta-analysis of association of body image concerns and self-compassion (hypothesis 4)

As shown in Figure 8, the results of 21 studies ($N = 5966$) showed that the combined uncorrected random effects estimate for the relationship between body image concerns and self-compassion was $r = -0.45$ (95% CI = -0.50 to -0.39; $Z = -14.37$; $p < .0001$). This represents a medium effect size, indicating that higher levels of body image concerns are related to lower levels of self-compassion. There was significantly high heterogeneity ($Q = 130.29$, $p < 0.001$, $I^2 = 84.65$), with 84 % of the variance in the effect size attributable to between-study variance.

Subgroup analysis shows that the correlation between self-compassion and body image concerns was not significantly affected by the type of the body image measure ($p = 0.26$) or the type of self-compassion measure ($p = 0.44$; see Table 6). However, the high heterogeneity in those two subgroups means that these conclusions are not reliable.




Figure 7. Funnel plot for relationship between self-compassion and positive body image. CES: Combined Effect Size.




Figure 8. Forest plot for the relationship between self-compassion and body image concerns

The results of the meta-regression analyses showed that the moderating effects of age, gender, BMI and study quality were not significant (see Table 6). Visual inspection of the

funnel plot (see Figure 9) suggested a relatively symmetric distribution of study findings. Additionally, the trim-and-fill analysis showed that there were no missing studies.

Table 6

Subgroup and meta-regression analysis of relationship between body image concerns and self-compassion

	Number of Studies	Correlation (<i>r</i>)	95% CI	I ²	<i>p</i>
Variables					0.26
Body Dissatisfaction	11	-0.43*	-0.53 to -0.32	88.79	
Body Shame	6	-0.49*	-0.52 to -0.45	00.01	
Independent measure					0.44
SCS	14	-0.47*	-0.51 to -0.42	63.60	
SCS-SF	7	-0.41*	-0.57 to -0.23	91.51	
Moderator	Number of Studies	B-Coefficient	95% CI	SE	<i>p</i>
Age (mean)	18	-0.09	-0.02 to 0.01	0.01	0.74
Percentage of female participants	21	-0.12	-0.63 to 0.38	0.24	0.60
BMI (mean)	13	0.24	-0.03 to 0.07	0.02	0.37
Study Quality Rating (0-3 criteria)	21	-0.18	-0.22 to 0.10	0.08	0.42

Note. *significant at $p < .05$, BMI: Body Mass Index; CI: confidence interval; SE; standard error; SCS: Self-Compassion Scale; SCS-SF: Self-compassion Scale-Short Form




Figure 9. Funnel plot for relationship between self-compassion and body image concerns. CES: Combined Effect Size.

Meta-analysis of impact of self-compassion interventions on body image (hypothesis 5)

The effects of self-compassion-related interventions were compared with a control group in 13 studies, totalling $N = 1714$ ($N = 819$ in intervention group; $N = 895$ in control). In Figure 10, studies 1 to 7 had active control groups, while studies 8 to 13 were studies with waitlist control groups. The overall effect was small to moderate at $g = 0.39$ (95 % CI = 0.22 to 0.55; $Z = 5.02$; $p < 0.001$), indicating that the self-compassion group had more improvement in body image than the control group. Between-study heterogeneity was low but significant ($I^2 = 49.06$ %; $Q = 23.56$, $p < 0.05$).




Figure 10. Forest plot showing the effect of self-compassion interventions compared to the control group.

The effect sizes were similar for studies involving active control groups ($g = 0.39$) and waitlist control groups ($g = 0.38$), though Meta-Essentials cautions that such analyses are not meaningful. There was no significant moderator effect of the duration of interventions ($b = -0.08$, 95% CI [0.00, 0.00], $p = .80$). Therefore, longer self-compassion interventions were no more effective than brief ones.

Inspection of the funnel plot and trim and fill procedure identified an asymmetric distribution of the study results (see Figure 11). After adjusting for missing studies ($n = 3$), the effect size dropped from $g = 0.39$ to $g = 0.29$ (95 % CI = 0.12 to 0.46), representing a small effect size in favour of self-compassion interventions over controls.




Figure 11. Forest plot of body image effect sizes for self-compassion group versus the control group. CES: Combined Effect Size.

Summary. Self-compassion interventions resulted in more positive body image and lower body image concerns, with medium effect sizes. This outcome demonstrates that the relationship between self-compassion and body image variables is causal, rather than simply being correlational. Thus, self-compassion interventions are effective in enhancing healthy

body image.

Discussion

The objective of this meta-analysis was to examine the relationship of self-compassion with eating concerns and body image. Where possible (particularly where RCTs were used), the causality of that relationship was addressed in order to understand the effectiveness of self-compassion related interventions. The findings support the hypotheses throughout, showing significant correlations and causal effects of self-compassion on eating pathology and body image. They broadly support previous reviews and meta-analyses on self-compassion in relation to mental health (e.g., Ferrari et al., 2019; Kirby, Tellegen & Steindl, 2017; Macbeth & Gumley, 2012; Marsh, Chan & Macbeth, 2018; Zessin, Dickhäuser, & Garbade, 2015).

The relationship of self-compassion with eating concerns and body image was characterized by a medium to high level of heterogeneity. Follow-up meta-regression analysis indicated that the association between self-compassion and eating pathology was higher in better quality studies, indicating the importance of stronger studies in this field. Regarding positive body image, BMI and age were significant moderators, indicating that self-compassion is more related to positive self-perception if individuals are older and bigger in size, suggesting that these findings need to be understood in the context of demographic factors. Considering body image interventions, it is important to note that there was no moderator effect of the duration of interventions, suggesting that short self-compassion interventions are still effective.

Limitations of this review

The current review is the first meta-analysis to examine the relationship of self-compassion with eating pathology and body image, and shows evidence of both correlational and causal links. However, this current review also has limitations. First, only studies

published in the English language were included. Thus, the findings might be an under-representation of non-Western cultures, potentially resulting in limited generalizability or inflation of the findings due to positive findings being more likely to be published in English language journals. The current review is also limited by the quality of existing studies, which rely heavily on correlational designs.

The combined effect sizes here must be interpreted with caution. A recent meta-analysis indicated that the association between psychopathology and negative indicators of self-compassion is stronger than the link with positive indicators (Muris & Petrocchi, 2017). Therefore, the protective role of self-compassion against eating and body image concerns could be overestimated due to the negative subscales (self-judgment, isolation and over-identification) that have already been shown to be related to psychopathology.

It is also important to note that the small number of samples in several analyses mean that it was not possible to interpret heterogeneity fully. When larger numbers of interventions have been reported, it will become possible to undertake the necessary moderator and subgroup analyses to account for the heterogeneity. Similarly, larger numbers of intervention studies with active control groups would ensure that effect sizes were more reliable (Cuijpers, Weitz, Cristea, & Twisk, 2017; Cunningham, Kypri & McCambridge, 2013). Finally, many of the self-compassion interventions were part of a wider therapy, meaning that the apparent effect size might have been affected. ‘Pure’ self-compassion studies need to be compared with other treatment approaches.

It is also important to note that positive and negative body image have been identified as being qualitatively different constructs (Tylka & Wood-Barcalow, 2015). As most studies in this review used either positive or negative body image measures, it is not possible to compare them directly. Future research should consider both constructs in order to distinguish the impact of self-compassion on these two different indicators of body image.

Research and clinical implications

As it is clear that self-compassion plays a role in understanding eating concerns and body image, it is important to consider the underlying mechanism that explain why and how self-compassion has its impact. Further research is needed to determine such mechanisms. For example, self-compassion has been suggested to be a moderator (e.g., Lonergan et al, 2019; Pisitsungkagarn et al, 2014) which interacts with other risk factors (e.g., body surveillance, thin-ideal internalization) to drive eating and body image concerns. Others have proposed self-compassion as a potential mediator, where lower levels of self-compassion partially or fully account for the effect of a range of risk factors (e.g., shame, perfectionism) on eating pathology and body image (e.g., Marta-Simões et al, 2016; Raque-Bogdan, Piontkowski, Hui, Ziemer, & Garriott, 2016). However, it will be particularly important to consider self-compassion as a potential causal factor, given that the evolutionary theory of compassion suggests that individuals who do not develop self-compassion in their early childhood might be prone to shame, self-criticism and guilt, leading to increased eating and body image concerns (Gilbert, 2014). Such models need to be tested in the field of eating disorders, and considered within prevention as well as treatment approaches.

Given the role of negative emotions in eating pathology (as outlined above), it can be hypothesised that self-compassion operates by providing a more adaptive means of coping with emotion (e.g., Neff, 2003a). Self-compassion requires mindful awareness of negative emotions, so that unwanted feelings are not avoided or suppressed (Neff, 2003a). Therefore, it is possible that self-compassion enables individuals to confront distressing emotions (Finlay-Jones, Rees & Kane, 2015), rather than coping with them maladaptively by engaging in pathological eating behaviours (e.g., restriction, bingeing, vomiting). Future research should more closely monitor the psychological and biological correlates of self-compassion that might reflect such a process.

As raised above, self-compassion is likely to work via its impact on emotion regulation in eating disorders. One could argue that Gilbert's (2014) approach is relevant to understanding the aetiology of eating pathology and body image, given its focus on evolutionary theory. On the other hand, Germer and Neff's (2013) approach might be more useful in understanding the maintenance of eating pathology and body image, and provides more immediate methods of directing clinical change (e.g., self-compassion mediation/letter writing). However, the data to date do not allow us to differentiate between the models in such a way. Therefore, future research and clinical work should look at which models are related to self-compassion.

Conclusion

The results of the present meta-analysis suggest that self-compassion is causally linked to eating pathology and body image, with broadly moderate effect sizes. Self-compassion approaches therefore provide a potentially useful tool for addressing emotionally-driven behaviours. This effect of self-compassion is robust, with little influence of moderators or demographics. Self-compassion approaches are relatively new, and their mechanisms of change not yet fully understood. Therefore, this treatment approach merits further attention and development in research, therapy and prevention settings, particularly where eating and body concerns are emotionally-driven and maintained. To determine its optimum use, self-compassion should be considered both as a stand-alone therapy and in combination with existing evidence-based approaches.

References

- Adams, C. E., & Leary, M. R. (2007). Promoting self-compassionate attitudes toward eating among restrictive and guilty eaters. *Journal of Social and Clinical Psychology, 26*, 1120–1144. doi: 10.1521/jscp.2007.26.10.1120
- Albertson, E. R., Neff, K. D., & Dill-Shackleford, K. E. (2015). Self-compassion and body dissatisfaction in women: a randomized controlled trial of a brief meditation intervention. *Mindfulness, 6*, 444–454. doi: 10.1007/s12671-014-0277-3
- Andrew, R., Tiggemann, M., & Clark, L. (2016). Predicting body appreciation in young women: An integrated model of positive body image. *Body Image, 18*, 34-42. doi: 10.1016/j.bodyim.2016.04.003
- Avalos, L., Tylka, T. L., & Wood-Barcalow, N. (2005). The Body Appreciation Scale: Development and psychometric evaluation. *Body Image, 2*, 285-297. doi: 10.1016/j.bodyim.2005.06.002
- Barnett, M. D., & Sharp, K. J. (2016). Maladaptive perfectionism, body image satisfaction, and disordered eating behaviors among U.S. college women: The mediating role of self-compassion. *Personality and Individual Differences, 99*, 225–234. doi: 10.1016/j.paid.2016.05.004
- Berking, M., & Whitley, B. (2014). *Affect Regulation Training*. New York, NY: Springer.
- Blomquist, K. K., Roberto, C. A., Barnes, R. D., White, M. A., Masheb, R. M., & Grilo, C. M. (2014). Development and validation of the Eating Loss of Control Scale. *Psychological Assessment, 26*, 77-89. doi: 10.1037/a0034729
- Braun, T. D., Park, C. L., & Gorin, A. (2016). Self-compassion, body image, and disordered eating: A review of the literature. *Body Image, 17*, 117-131. doi: 10.1016/j.bodyim.2016.03.003
- Breines, J., Toole, A., Tu, C., & Chen, S. (2014). Self-compassion, body image, and self-

- reported disordered eating. *Self and Identity*, 13, 432-448. doi: 10.1080/15298868.2013.838992
- Cho, M. J., Kim, J. K., Jeon, H. J., Suh, T., Chung, I. W., Hong, J. P., ... & Lee, C. K. (2007). Lifetime and 12-month prevalence of DSM-IV psychiatric disorders among Korean adults. *The Journal of Nervous and Mental Disease*, 195, 203-210. doi: 10.1097/01.nmd.0000243826.40732.45
- Cohen, J. (1992). A power primer. *Psychological Bulletin*, 112, 155–159. doi: 10.1037//0033-2909.112.1.155
- Couturier, J., Kimber, M., & Szatmari, P. (2013). Efficacy of family-based treatment for adolescents with eating disorders: A systematic review and meta-analysis. *International Journal of Eating Disorders*, 46, 3-11. doi: 10.1002/eat.22042
- Cuijpers, P., Weitz, E., Cristea, I. A., & Twisk, J. (2017). Pre-post effect sizes should be avoided in meta-analyses. *Epidemiology and Psychiatric Sciences*, 26, 364-368. doi: 10.1017/S2045796016000809
- Cunningham, J. A., Kypri, K., & McCambridge, J. (2013). Exploratory randomized controlled trial evaluating the impact of a waiting list control design. *BMC Medical Research Methodology*, 13, 150. doi: 10.1186/1471-2288-13-150
- Daye, C. A., Webb, J. B., & Jafari, N. (2014). Exploring self-compassion as a refuge against recalling the body-related shaming of caregiver eating messages on dimensions of objectified body consciousness in college women. *Body Image*, 11, 547–556. doi: 10.1016/j.bodyim.2014.08.001
- Duarte, C., Ferreira, C., Trindade, I. A., & Pinto-Gouveia, J. (2015). Body image and college women's quality of life: The importance of being self-compassionate. *Journal of Health Psychology*, 20, 754–764. doi: 10.1177/1359105315573438
- Duarte, C., Pinto-Gouveia, J., & Stubbs, R. J. (2017). Compassionate Attention and

- Regulation of Eating Behaviour: A pilot study of a brief low-intensity intervention for binge eating. *Clinical Psychology & Psychotherapy*, 24, O1437–O1447. doi: 10.1002/cpp.2094
- Duval, S., & Tweedie, R. (2000). Trim and fill: a simple funnel-plot–based method of testing and adjusting for publication bias in meta-analysis. *Biometrics*, 56, 455-463. doi: 10.1111/j.0006-341x.2000.00455.x
- Evans, C., & Dolan, B. (1993). Body Shape Questionnaire: derivation of shortened “alternate forms”. *International Journal of Eating Disorders*, 13, 315-321. doi: 10.1002/1098-108x(199304)13:3<315::aid-eat2260130310>3.0.co;2-3
- Engel, S. G., Wonderlich, S. A., Crosby, R. D., Mitchell, J. E., Crow, S., Peterson, C. B., ... & Gordon, K. H. (2013). The role of affect in the maintenance of anorexia nervosa: Evidence from a naturalistic assessment of momentary behaviors and emotion. *Journal of Abnormal Psychology*, 122, 709-719. doi: 10.1037/a0034010
- Fairburn, C. G., Bailey-Straebl, S., Basden, S., Doll, H. A., Jones, R., Murphy, R., ... & Cooper, Z. (2015). A transdiagnostic comparison of enhanced cognitive behaviour therapy (CBT-E) and interpersonal psychotherapy in the treatment of eating disorders. *Behaviour Research and Therapy*, 70, 64-71. doi: 10.1016/j.brat.2015.04.010
- Fairburn, C. G., & Beglin, S. J. (1994). Assessment of eating disorders: Interview or self-report questionnaire? *International Journal of Eating Disorders*, 16, 363-370. doi: 10.1002/1098-108X(199412)16:4<363::AID-EAT2260160405>3.0.CO;2-%23
- Ferreira, C., Matos, M., Duarte, C., & Pinto-Gouveia, J. (2014). Shame memories and eating psychopathology: The buffering effect of self-compassion. *European Eating Disorders Review*, 22, 487-494. doi: 10.1002/erv.2322
- Ferreira, C., Oliveira, S., & Mendes, A. L. (2017). Kindness toward one’s self and body: exploring mediational pathways between early memories and disordered eating. *Spanish*

Journal of Psychology, 23, 1–8. doi: 10.1017/sjp.2017.50

- Ferreira, C., Pinto-Gouveia, J., & Duarte, C. (2013). Self-compassion in the face of shame and body image dissatisfaction: Implications for eating disorders. *Eating Behaviors*, 14, 207–210. doi: 10.1016/j.eatbeh.2013.01.005
- Ferrari, M., Hunt, C., Harrysunker, A., Abbott, M. J., Beath, A. P., & Einstein, D. A. (2019). Self-compassion interventions and psychosocial outcomes: A meta-analysis of RCTs. *Mindfulness*, 10, 1455–1473. doi: 10.1007/s12671-019-01134-6
- Finlay-Jones, A. L., Rees, C. S., & Kane, R. T. (2015). Self-compassion, emotion regulation and stress among Australian psychologists: Testing an emotion regulation model of self-compassion using structural equation modelling. *PloS One*, 10, 1–19. doi: 10.1371/journal.pone.0133481
- Franzoi, S. L., & Shields, S. A. (1984). The Body Esteem Scale: Multidimensional structure and sex differences in a college population. *Journal of Personality Assessment*, 48, 173–178. doi: 10.1207/s15327752jpa4802_12
- Fresnics, A. A., Wang, S. B., & Borders, A. (2019). The unique associations between self-compassion and eating disorder psychopathology and the mediating role of rumination. *Psychiatry Research*, 274, 91–97. doi: 10.1016/j.psychres.2019.02.019
- Fu, R., Gartlehner, G., Grant, M., Shamliyan, T., Sedrakyan, A., Wilt, T. J., ... & Santaguida, P. (2011). Conducting quantitative synthesis when comparing medical interventions: AHRQ and the Effective Health Care Program. *Journal of Clinical Epidemiology*, 64, 1187–1197. doi: 10.1016/j.jclinepi.2010.08.010
- Gale, C., Gilbert, P., Read, N., & Goss, K. (2014). An evaluation of the impact of introducing compassion focused therapy to a standard treatment programme for people with eating disorders. *Clinical Psychology and Psychotherapy*, 21, 1–12. doi: 10.1017/sjp.2017.50
- Garner, D. M., Olmsted, M. P., Bohr, Y., & Garfinkel, P. E. (1982). The Eating Attitudes Test:

- psychometric features and clinical correlates. *Psychological Medicine*, *12*, 871-878. doi: 10.1017/s0033291700049163
- Germer, C. K., & Neff, K. D. (2013). Self-compassion in clinical practice. *Journal of Clinical Psychology*, *69*, 856-867. doi: 10.1002/jclp.22021
- Gilbert, P. (2019). Psychotherapy for the 21st Century: An integrative, evolutionary, contextual, biopsychosocial approach. *Psychology and Psychotherapy: Theory, Research and Practice*, *92*, 164-189. doi: 10.1111/papt.12226
- Gilbert, P. (2014). The origins and nature of compassion focused therapy. *British Journal of Clinical Psychology*, *53*, 6-41. doi: 10.1111/bjc.12043
- Gilbert, P. (2009). *The compassionate mind: A new approach to life's challenges*. Oakland, CA: New Harbinger.
- Goss, K., & Allan, S. (2014). The development and application of compassion-focused therapy for eating disorders (CFT-E). *British Journal of Clinical Psychology*, *53*, 62-77. doi: 10.1111/bjc.12039
- Goss, K., & Allan, S. (2009). Shame and pride in eating disorders. *Clinical Psychology and Psychotherapy*, *16*, 303– 316. doi: 10.1002/cpp.627
- Gouveia, M. J., Canavarro, M. C., & Moreira, H. (2019). Associations between mindfulness, self-compassion, difficulties in emotion regulation, and emotional eating among adolescents with overweight/obesity. *Journal of Child and Family Studies*, *28*, 273–285. doi: 10.1007/s10826-018-1239-5
- Gouveia, M. J., Canavarro, M. C., & Moreira, H. (2018). Is mindful parenting associated with adolescents' emotional eating? The mediating role of adolescents' self-compassion and body shame. *Frontiers in Psychology*, *9*. doi: 10.3389/fpsyg.2018.02004
- Haynos, A. F., & Fruzzetti, A. E. (2011). Anorexia nervosa as a disorder of emotion dysregulation: Evidence and treatment implications. *Clinical Psychology: Science and*

- Practice*, 18, 183-202. doi: 10.1111/j.1468-2850.2011.01250.x
- Heatherton, T. F., & Baumeister, R. F. (1991). Binge eating as escape from self-awareness. *Psychological Bulletin*, 110, 86-108. doi: 10.1037/0033-2909.110.1.86
- Hedges, L. V., & Olkin, I. (1985). *Statistical methods for meta-analysis*. San Diego, CA: Academic Press.
- Higgins, J. P., Thompson, S. G., Deeks, J. J., & Altman, D. G. (2003). Measuring inconsistency in meta-analyses. *British Medical Journal*, 327, 557-560. doi: 10.1136/bmj.327.7414.557
- Homan, K. J., & Tylka, T. L. (2015). Self-compassion moderates body comparison and appearance self-worth's inverse relationships with body appreciation. *Body Image*, 15, 1-7. doi: 10.1016/j.bodyim.2015.04.007
- Kelly, A. C., & Carter, J. C. (2014). Eating disorder subtypes differ in their rates of psychosocial improvement over treatment. *Journal of Eating Disorders*, 2, 1–10. doi: 10.1186/2050-2974-2-2
- Kelly, A. C., & Carter, J. C. (2015). Self-compassion training for binge eating disorder: A pilot randomized controlled trial. *Psychology and Psychotherapy: Theory, Research and Practice*, 88, 285-303. doi: 10.1111/papt.12044
- Kelly, A. C., Carter, J. C., Zuroff, D. C., & Borairi, S. (2013). Self-compassion and fear of self-compassion interact to predict response to eating disorders treatment: A preliminary investigation. *Psychotherapy Research*, 23, 252–264. doi 10.1080/10503307.2012.717310
- Kelly, A. C., Miller, K. E., & Stephen, E. (2016). The benefits of being self-compassionate on days when interactions with body-focused others are frequent. *Body Image*, 19, 195-203. doi: 10.1016/j.bodyim.2016.10.005
- Kelly, A. C., & Stephen, E. (2016). A daily diary study of self-compassion, body image, and eating behavior in female college students. *Body Image*, 17, 152–160. doi:

10.1016/j.bodyim.2016.03.006

Kelly, A. C., & Tasca, G. A. (2016). Within-persons predictors of change during eating disorders treatment: An examination of self-compassion, self-criticism, shame, and eating disorder symptoms. *International Journal of Eating Disorders, 49*, 716–722. doi: 10.1002/eat.22527

Kelly, A. C., & Waring, S. V. (2018). A feasibility study of a 2-week self-compassionate letter-writing intervention for nontreatment seeking individuals with typical and atypical anorexia nervosa. *International Journal of Eating Disorders, 51*, 1005–1009. doi: 10.1002/eat.22930

Kelly, A. C., Vimalakanthan, K., & Miller, K. E. (2014). Self-compassion moderates the relationship between body mass index and both eating disorder pathology and body image flexibility. *Body Image, 11*, 446–453. doi: 10.1016/j.bodyim.2014.07.005

Kelly, A. C., Wisniewski, L., Martin-Wagar, C., & Hoffman, E. (2017). Group-based compassion-focused therapy as an adjunct to outpatient treatment for eating disorders: A pilot randomized controlled trial. *Clinical Psychology & Psychotherapy, 24*, 475-487. doi: 10.1002/cpp.2018

Kirby, J. N., Tellegen, C. L., & Steindl, S. R. (2017). A meta-analysis of compassion-based interventions: Current state of knowledge and future directions. *Behavior Therapy, 48*, 778-792. doi: 10.1016/j.beth.2017.06.003

Klimecki, O. M., Leiberg, S., Ricard, M., & Singer, T. (2014). Differential pattern of functional brain plasticity after compassion and empathy training. *Social Cognitive and Affective Neuroscience, 9*, 873-879. doi: 10.1093/scan/nst060

Kramer, R., & Cuccolo, K. (2019). Yoga practice in a college sample: associated changes in eating disorder, body image, and related factors over time. *Eating Disorders, 1-19*. doi: 10.1080/10640266.2019.1688007

- Lavender, J. M., Wonderlich, S. A., Peterson, C. B., Crosby, R. D., Engel, S. G., Mitchell, J. E., ... & Berg, K. C. (2014). Dimensions of emotion dysregulation in bulimia nervosa. *European Eating Disorders Review*, *22*, 212-216. doi: 10.1002/erv.2288.
- Lehr, E. J., Krohmer, K., Schag, K., Dresler, T., Zipfel, S., & Giel, K. E. (2015). Emotion regulation model in binge eating disorder and obesity-a systematic review. *Neuroscience & Biobehavioral Reviews*, *49*, 125-134. doi: 10.1016/j.neubiorev.2014.12.008
- Lonergan, A. R., Bussey, K., Mond, J., Brown, O., Giffiths, S., Muray, S. B., & Mitchison, D. (2019). Me, my selfie, and I: The relationship between editing and posting selfies and body dissatisfaction in men and women. *Body Image*, *28*, 39-43. doi: 10.1016/j.bodyim.2018.12.001
- MacBeth, A., & Gumley, A. (2012). Exploring compassion: a meta-analysis of the association between self-compassion and psychopathology. *Clinical Psychology Review*, *32*, 545-552. doi: 10.1016/j.cpr.2012.06.003
- Marsh, I. C., Chan, S. W., & MacBeth, A. (2018). Self-compassion and psychological distress in adolescents-a meta-analysis. *Mindfulness*, *9*, 1011-1027. doi: 10.1007/s12671-017-0850-7
- Marta-Simões, J., & Ferreira, C. (2018). Self-to-others and self-to-self relationships: paths to understanding the valence of body image and eating attitudes in emerging adult women. *Eating and Weight Disorders*, *0*, 1-8. doi: 10.1007/s40519-018-0612-0
- Marta-Simões, J., Ferreira, C., & Mendes, A. L. (2016). Exploring the effect of external shame on body appreciation among Portuguese young adults: The role of self-compassion. *Eating Behaviors*, *23*, 174-179. doi: 10.1016/j.eatbeh.2016.10.006
- McCarthy, M. (1990). The thin ideal, depression and eating disorders in women. *Behaviour Research and Therapy*, *28*, 205-214. doi: 10.1016/0005-7967(90)90003-2
- McKinley, N. M., & Hyde, J. S. (1996). The objectified body consciousness scale:

- Development and validation. *Psychology of Women Quarterly*, *20*, 181-215. doi: 10.1111/j.1471-6402.1996.tb00467.x
- McManus, F., & Waller, G. (1995). A functional analysis of binge-eating. *Clinical Psychology Review*, *15*, 845-863. doi: 10.1016/0272-7358(95)00042-9
- Modica, C. (2019). Facebook, body esteem, and body surveillance in adult women: The moderating role of self-compassion and appearance-contingent self-worth. *Body Image*, *29*, 17–30. doi: 10.1016/j.bodyim.2019.02.002
- Moffitt, R. L., Neumann, D. L., & Williamson, S. P. (2018). Comparing the efficacy of a brief self-esteem and self-compassion intervention for state body dissatisfaction and self-improvement motivation. *Body Image*, *27*, 67–76. doi: 10.1016/j.bodyim.2018.08.008
- Mosewich, A. D., Kowalski, K. C., Sabiston, C. M., Sedgwick, W. A., & Tracy, J. L. (2011). Self-compassion: a potential resource for young women athletes. *Journal of Sport & Exercise Psychology*, *33*, 103-123. doi: 10.1123/jsep.33.1.103
- Muris, P., & Petrocchi, N. (2017). Protection or vulnerability? A meta-analysis of the relations between the positive and negative components of self-compassion and psychopathology. *Clinical Psychology & Psychotherapy*, *24*, 373-383. doi: 10.1002/cpp.2005
- Neff, K. (2003a). Self-compassion: An alternative conceptualization of a healthy attitude toward oneself. *Self and identity*, *2*, 85-101. doi: 10.1080/15298860390129863
- Neff, K. D. (2003b). Development and validation of a scale to measure self-compassion. *Self and Identity*, *2*, 223-250. doi: 10.1080/15298860390209035
- National Institute for Health and Care Excellence (2017). *Eating disorders: Recognition and treatment*. London, UK: National Institute for Health and Care Excellence.
- Palmeira, L., Cunha, M., & Pinto-Gouveia, J. (2017). Processes of change in quality of life, weight self-stigma, body mass index and emotional eating after an acceptance-

- mindfulness- and compassion-based group intervention (Kg-Free) for women with overweight and obesity. *Journal of Health Psychology*, *24*, 1056-1069. doi: 10.1177/1359105316686668
- Pinto-Gouveia, J., Carvalho, S. A., Palmeira, L., Castilho, P., Duarte, C., Ferreira, C., ... Costa, J. (2017). BEfree: A new psychological program for binge eating that integrates psychoeducation, mindfulness, and compassion. *Clinical Psychology and Psychotherapy*, *24*, 1090–1098. doi: 10.1002/cpp.2072
- Pinto-Gouveia, J., Carvalho, S. A., Palmeira, L., Castilho, P., Duarte, C., Ferreira, C., ... & Costa, J. (2019). Incorporating psychoeducation, mindfulness and self-compassion in a new programme for binge eating (BEfree): Exploring processes of change. *Journal of Health Psychology*, *24*, 466-479. doi: 10.1177/1359105316676628
- Pinto-Gouveia, J., Ferreira, C., & Duarte, C. (2014). Thinness in the pursuit for social safeness: an integrative model of social rank mentality to explain eating psychopathology. *Clinical Psychology & Psychotherapy*, *21*, 154–165. doi: 10.1002/cpp.1820
- Pisitsungkarn, K., Taephant, N., & Attasaranya, P. (2014). Body image satisfaction and self-esteem in Thai female adolescents: the moderating role of self-compassion. *International Journal of Adolescent Medicine and Health*, *26*, 333-338. doi: 10.1515/ijamh-2013-0307
- Polivy, J., & Herman, C. P. (2002). Causes of eating disorders. *Annual Review of Psychology*, *53*, 187-213. doi: 10.1146/annurev.psych.53.100901.135103
- Preti, A., de Girolamo, G., Vilagut, G., Alonso, J., de Graaf, R., Bruffaerts, R., ... & ESEMeD-WMH Investigators. (2009). The epidemiology of eating disorders in six European countries: results of the ESEMeD-WMH project. *Journal of Psychiatric Research*, *43*, 1125-1132. doi: 10.1016/j.jpsychires.2009.04.003
- Pritchard, M. (2008). Disordered eating in undergraduates: Does gender role orientation

- influence men and women the same way? *Sex Roles*, *59*, 282-289. doi: 10.1007/s11199-008-9449-8
- Pullmer, R., Coelho, J. S., & Zaitsoff, S. L. (2019). Kindness begins with yourself: The role of self-compassion in adolescent body satisfaction and eating pathology. *International Journal of Eating Disorders*, *52*, 809-816. doi.org: 10.1002/eat.23081
- Pullmer, R., Zaitsoff, S. L., & Coelho, J. S. (2019). Self-compassion and eating pathology in female adolescents with eating disorders: the mediating role of psychological distress. *Mindfulness*, *10*, 2716-2723. doi: 10.1007/s12671-019-01254-z
- Raes, F., Pommier, E., Neff, K. D., & Van Gucht, D. (2011). Construction and factorial validation of a short form of the Self-Compassion Scale. *Clinical Psychology & Psychotherapy*, *18*, 250-255. doi: 10.1002/cpp.702
- Raque-Bogdan, T. L., Piontkowski, S., Hui, K., Ziemer, K. S., & Garriott, P. O. (2016). Self-compassion as a mediator between attachment anxiety and body appreciation: An exploratory model. *Body Image*, *19*, 28-36. doi: 10.1016/j.bodyim.2016.08.001
- Rodgers, R. F., Donovan, E., Cousineau, T., Yates, K., McGowan, K., Cook, E., ... Franko, D. L. (2018). BodiMojo: efficacy of a mobile-based intervention in improving body image and self-compassion among adolescents. *Journal of Youth and Adolescence*, *47*, 1363–1372. doi: 10.1007/s10964-017-0804-3
- Rosenthal, R. (1993). *Meta-analytic procedures for social research*. Newbury Park, CA: Sage
- Schmidt, C. K., Raque-Bogdan, T. L., & Hollern, E. A. (2019). Self-compassion, affect, and body image in college women. *Journal of College Counseling*, *22*, 152-163. doi: 10.1002/jocc.12127
- Schoenefeld, S. J., & Webb, J. B. (2013). Self-compassion and intuitive eating in college women: Examining the contributions of distress tolerance and body image acceptance and action. *Eating Behaviors*, *14*, 493–496. doi: 10.1016/j.eatbeh.2013.09.001

- Seekis, V., Bradley, G. L., & Duffy, A. (2017). The effectiveness of self-compassion and self-esteem writing tasks in reducing body image concerns. *Body Image, 23*, 206–213. doi: 10.1016/j.bodyim.2017.09.003
- Smyth, J. M., Wonderlich, S. A., Heron, K. E., Sliwinski, M. J., Crosby, R. D., Mitchell, J. E., & Engel, S. G. (2007). Daily and momentary mood and stress are associated with binge eating and vomiting in bulimia nervosa patients in the natural environment. *Journal of Consulting and Clinical Psychology, 75*, 629-638. doi: 10.1037/0022-006X.75.4.629
- Solmi, F., Hotopf, M., Hatch, S. L., Treasure, J., & Micali, N. (2016). Eating disorders in a multi-ethnic inner-city UK sample: prevalence, comorbidity and service use. *Social Psychiatry and Psychiatric Epidemiology, 51*, 369-381. doi: 10.1007/s00127-015-1146-7
- Stern, N. G., & Engeln, R. (2018). Self-compassionate writing exercises increase college women's body satisfaction. *Psychology of Women Quarterly, 42*, 326-341. Doi: 10.1177/0361684318773356
- Stice, E., Marti, C. N., & Rohde, P. (2013). Prevalence, incidence, impairment, and course of the proposed DSM-5 eating disorder diagnoses in an 8-year prospective community study of young women. *Journal of Abnormal Psychology, 122*, 445-457. doi: 10.1037/a0030679
- Stice, E., & Shaw, H. E. (2002). Role of body dissatisfaction in the onset and maintenance of eating pathology: A synthesis of research findings. *Journal of Psychosomatic Research, 53*, 985–993. doi: 10.1016/S0022-3999(02)00488-9
- Stutts, L. A., & Blomquist, K. K. (2018). The moderating role of self-compassion on weight and shape concerns and eating pathology: A longitudinal study. *International Journal of Eating Disorders, 51*, 879-889. doi: 10.1002/eat.22880
- Södersten, P., Bergh, C., Leon, M., Brodin, U., & Zandian, M. (2017). Cognitive behavior therapy for eating disorders versus normalization of eating behavior. *Physiology & Behavior, 174*, 178-190. doi: 10.1016/j.physbeh.2017.03.016

- Suurmond, R., van Rhee, H., & Hak, T. (2017). Introduction, comparison, and validation of Meta-Essentials: A free and simple tool for meta-analysis. *Research Synthesis Methods, 8*, 537-553. doi: 10.1002/jrsm.1260
- Taylor, M. B., Daiss, S., & Krietsch, K. (2015). Associations among self-compassion, mindful eating, eating disorder symptomatology, and body mass index in college students. *Translational Issues in Psychological Science, 1*, 229–238. doi: 10.1037/tps0000035
- Thomas, B. H., Ciliska, D., Dobbins, M., & Micucci, S. (2004). A process for systematically reviewing the literature: Providing the research evidence for public health nursing interventions. *Worldviews on Evidence-Based Nursing, 1*, 176-184. doi: 10.1111/j.1524-475X.2004.04006.x
- Thompson, J. K., Heinberg, L. J., Altabe, M., & Tantleff-Dunn, S. (1999). *Exacting beauty: Theory, assessment, and treatment of body image disturbance*. Washington, DC: American Psychological Association.
- Toole, A. M., & Craighead, L. W. (2016). Brief self-compassion meditation training for body image distress in young adult women. *Body Image, 19*, 104–112. doi: 10.1016/j.bodyim.2016.09.001
- Tylka, T. L., Russell, H. L., & Neal, A. A. (2015). Self-compassion as a moderator of thinness-related pressures' associations with thin-ideal internalization and disordered eating. *Eating Behaviors, 17*, 23–26. doi: 10.1016/j.eatbeh.2014.12.009
- Tylka, T. L., & Wood-Barcalow, N. L. (2015). What is and what is not positive body image? Conceptual foundations and construct definition. *Body Image, 14*, 118-129. doi: 10.1016/j.bodyim.2015.04.001
- Van de Mortel, T. F. (2008). Faking it: social desirability response bias in self-report research. *Australian Journal of Advanced Nursing, 25*, 40-48.
- Van Strien, T., Frijters, J. E., Bergers, G. P., & Defares, P. B. (1986). The Dutch Eating

- Behavior Questionnaire (DEBQ) for assessment of restrained, emotional, and external eating behavior. *International Journal of Eating Disorders*, *5*, 295-315. doi.org: 10.1002/1098-108X(198602)5:2<295::AID-EAT2260050209>3.0.CO;2-T
- Voelker, D. K., Petrie, T. A., Huang, Q., & Chandran, A. (2019). Bodies in Motion: An empirical evaluation of a program to support positive body image in female collegiate athletes. *Body Image*, *28*, 149–158. doi: 10.1016/j.bodyim.2019.01.008
- Waller, G., & Mountford, V. A. (2015). Weighing patients within cognitive-behavioural therapy for eating disorders: How, when and why. *Behaviour Research and Therapy*, *70*, 1-10. doi: 10.1016/j.brat.2015.04.004
- Wasylikiw, L., MacKinnon, A. L., & MacLellan, A. M. (2012). Exploring the link between self-compassion and body image in university women. *Body Image*, *9*, 236–245. doi: 10.1016/j.bodyim.2012.01.007
- Webb, J. B., Fiery, M. F., & Jafari, N. (2016). “You better not leave me shaming!”: Conditional indirect effect analyses of anti-fat attitudes, body shame, and fat talk as a function of self-compassion in college women. *Body Image*, *18*, 5–13. doi: 10.1016/j.bodyim.2016.04.009
- Webb, J. B., & Hardin, A. S. (2016). An integrative affect regulation process model of internalized weight bias and intuitive eating in college women. *Appetite*, *102*, 60–69. doi: 10.1016/j.appet.2016.02.024
- Williams, M. O., Tsivos, Z., Brown, S., Whitelock, N. S., & Sampson, M. (2017). Compassion-focused therapy for bulimia nervosa and bulimic presentations: a preliminary case series. *Behaviour Change*, *34*, 199–207. doi: 10.1017/bec.2017.13
- Wonderlich, S. A., Engel, S. G., Peterson, C. B., Robinson, M. D., Crosby, R. D., Mitchell, J. E., ... & Strauman, T. J. (2008). Examining the conceptual model of integrative cognitive-affective therapy for BN: Two assessment studies. *International Journal of Eating*

Disorders, 41, 748-754. doi: 10.1002/eat.20551

Zessin, U., Dickhäuser, O., & Garbade, S. (2015). The relationship between self-compassion and well-being: A meta-analysis. *Applied Psychology: Health and Well-Being, 7*, 340-364. doi: 10.1111/aphw.12051

Ziemer, K. S., Lamphere, B. R., Raque-Bogdan, T. L., & Schmidt, C. K. (2019). A randomized controlled study of writing interventions on college women's positive body image. *Mindfulness, 10*, 66–77. doi: 10.1007/s12671-018-0947-7