



This is a repository copy of *Cell-specific conditional deletion of interleukin-1 (IL-1) ligands and its receptors : a new toolbox to study the role of IL-1 in health and disease.*

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/161835/>

Version: Published Version

Article:

Pinteaux, E., Abdulaal, W.H., Mufazalov, I.A. et al. (5 more authors) (2020) Cell-specific conditional deletion of interleukin-1 (IL-1) ligands and its receptors : a new toolbox to study the role of IL-1 in health and disease. *Journal of Molecular Medicine*, 98 (7). pp. 923-930. ISSN 0946-2716

<https://doi.org/10.1007/s00109-020-01928-5>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



Cell-specific conditional deletion of interleukin-1 (IL-1) ligands and its receptors: a new toolbox to study the role of IL-1 in health and disease

Emmanuel Pinteaux¹ · Wesam H Abdulaal^{1,2} · Ilgiz A Mufazalov³ · Neil E Humphreys^{1,4} · Maj Simonsen-Jackson¹ · Sheila Francis⁵ · Werner Müller¹ · Ari Waisman³

Received: 7 April 2020 / Revised: 15 May 2020 / Accepted: 20 May 2020
© The Author(s) 2020

Abstract

The pro-inflammatory cytokine interleukin-1 (IL-1) plays a key role in many physiological processes and during the inflammatory and immune response to most common diseases. IL-1 exists as two agonists, IL-1 α and IL-1 β that bind to the only signaling IL-1 type 1 receptor (IL-1R1), while a second decoy IL-1 type 2 receptor (IL-1R2) binds both forms of IL-1 without inducing cell signaling. The field of immunology and inflammation research has, over the past 35 years, unraveled many mechanisms of IL-1 actions, through in vitro manipulation of the IL-1 system or by using genetically engineered mouse models that lack either member of the IL-1 family in ubiquitous constitutive manner. However, the limitation of global mouse knockout technology has significantly hampered our understanding of the precise mechanisms of IL-1 actions in animal models of disease. Here we report and review the recent generation of new conditional mouse mutants in which exons of *Il1a*, *Il1b*, *Il1r1*, and *Il1r2* genes flanked by loxP sites (^{fl/fl}) can be deleted in cell-/tissue-specific constitutive or inducible manner by Cre recombinase expression. Hence, IL-1 α ^{fl/fl}, IL-1 β ^{fl/fl}, IL-1R1^{fl/fl}, and IL-1R2^{fl/fl} mice constitute a new toolbox that will provide a step change in our understanding of the cell-specific role of IL-1 and its receptor in health and disease and the potential development of targeted IL-1 therapies.

Keywords Inflammation · Immunity · IL-1 · IL-1 receptors · Cre/loxP · Conditional deletion

Werner Müller and Ari Waisman are joint senior authors

✉ Emmanuel Pinteaux
emmanuel.pinteaux@manchester.ac.uk

¹ Faculty of Biology, Medicine and Health, University of Manchester, AV Hill Building, Oxford Road, Manchester M13 9PT, United Kingdom

² Biochemistry Department, Faculty of Sciences, King Abdulaziz University, P.O.BOX 80203, Jeddah 21589, Kingdom of Saudi Arabia

³ Institute for Molecular Medicine, University Medical Center of the Johannes Gutenberg-University of Mainz, Langenbeckstrasse 1, Building 308A, 55131 Mainz, Germany

⁴ Epigenetics and Neurobiology Unit, Adriano Buzzati-Traverso Campus, EMBL-Rome, Via Ramarini, 3200015 Monterotondo, RM, Italy

⁵ Department of Infection, Immunity & Cardiovascular Disease, Medical School, University of Sheffield, S10 2RX, Sheffield, United Kingdom

Introduction

Interleukin-1 is a master pro-inflammatory cytokine implicated in a wide range of physiological processes including development [1], regulation of neuroimmune and neuroendocrine functions [2], and central processes such as sleep and memory [3] and plays a key role in the initiation and orchestration of the inflammatory response to most, if not all, pathological inflammatory conditions, including infections and non-communicable diseases such as atherosclerosis or stroke (see [4] for review). The IL-1 family comprises two IL-1 agonists (IL-1 α and IL-1 β) and the naturally IL-1 receptor antagonist (IL-1Ra) that bind primarily to the IL-1 type 1 receptor (IL-1R1) and IL-1 receptor accessory protein (IL-1RAcP) leading to cell signaling (reviewed in [5]), while a second IL-1 type 2 receptor (IL-1R2) binds both IL-1 isoforms without inducing cell signaling, acting therefore as a decoy receptor [6].

Since their discovery, IL-1 α and IL-1 β are believed to have similar and often overlapping biological functions, since

they bind to the same receptor inducing similar cell signaling mechanisms. However, there are marked differences in the regulation of expression and mechanisms of actions of these two cytokines. Although both isoforms require enzymatic cleavage for generation of their mature forms, IL-1 β is the main secreted isoform, whereas IL-1 α remains cytoplasmic but can also be released during cell death or by mechanisms that are different from that of IL-1 β (reviewed in [7]). Furthermore, both IL-1 α and IL-1 β have been thought to exert similar biological activities primarily through binding to IL-1R1. However, several previous studies have demonstrated differential actions of both cytokines in various paradigms of inflammation; for instance, IL-1 α and IL-1 β exert differential potency at inducing fever when administered exogenously [8], while IL-1 α , but not IL-1 β , triggers sepsis lethality in mouse [9] and is required for T cell activation during allergen-induced hypersensitivity [10]. Further, IL-1 α , but not IL-1 β , induces brain cells to generate the LG3 neuroprotective protein fragment of the extracellular matrix component perlecan, a prominent component of the blood-brain barrier [11]. Of interest, polymorphisms in the human *Il1a*, but not *Il1b* gene, is associated with higher incidence of vascular malformation and/or higher risk of ischemic stroke [12, 13]. In contrast, IL-1 β , but not IL-1 α , activates IL-6 expression in neurons [14], selectively mediates the response to vascular injury [15], while IL-1 α - and IL-1 β -specific actions have also been identified in acute colon inflammation in mice [16]. Taken together, these observations suggest that IL-1 α and IL-1 β may be differentially expressed during inflammation and may exert non-overlapping ligand-specific differential actions dependent on the disease paradigm.

Mouse genetic models to understand the role of IL-1 α and IL-1 β in disease

For decades, the field of inflammation research has unraveled key mechanisms of IL-1 actions using traditional global gene targeting knockout technology in animal models. Indeed, IL-1 α -deficient ($^{-/-}$), IL-1 β $^{-/-}$, and IL-1 α/β $^{-/-}$ (as well as IL-1Ra $^{-/-}$) mice generated by Horai and collaborators in 1998 [17] have proven useful to identify some selective mechanisms of actions of both isoforms in some pathological conditions. In those genetic models, disruption of the *Il1a* and *Il1b* genes was achieved by deletion of the NH₂-terminal coding region for mature IL-1 α (exon 5–intron 5) and IL-1 β (exon 3–5), leading to ubiquitous constitutive inhibition of expression of either genes. These genetic models have been used widely in many disease models and have subsequently led to the identification of some IL-1 α - and IL-1 β -specific mechanisms as described above. Further, IL-1R1 $^{-/-}$ mice, originally generated by Immunex by targeted deletion of exon 1 and 2 of the *Il1r1* gene [18], showed that most, but not all,

IL-1 actions are mediated by IL-1R1 (see [19] for review). Indeed, studies using IL-1R1 $^{-/-}$ mice in animal models of gut infection with helminth *Trichuris muris* [20] and experimental stroke [21] found that IL-1 β can function in an IL-1R1-independent manner, while IL-1 β exacerbates neuronal apoptosis caused by status epilepticus through a mechanism independent of IL-1R1 [22]. Further, some neuroprotective actions of IL-1 are believed to be triggered independently of IL-1R1 via activation of the neuroprotective PI3K/Akt signaling pathway [23], while we have reported IL-1R1-independent IL-1 actions in glial cells [24]. Those IL-1R1-independent actions, primarily observed in the original IL-1R1 $^{-/-}$ mice, are known to be mediated through a spliced variant of the *Il1r1* gene leading to a truncated IL-1R1 isoform still expressed upon exon 1–2 deletion, due to the activation of an additional internal promoter positioned upstream of exon 1–2 [25]. This truncated isoform of the receptor has been fully characterized and lacks part of the extracellular IL-1 binding region but is still capable of inducing an intracellular signal in response to IL-1 that is known to mediate the neuroprotective actions of IL-1 in the brain via activation of the PI3K/Akt pathways [25]. Ubiquitous *Il1rap* gene deletion, that encodes IL-1RAcP, has also been achieved in mice, by targeting exon D1 and part of exon D2 that encode the first Ig-like and part of the second Ig-like extracellular domains, resulting in complete inhibition of IL-1 signaling in response to IL-1 α and IL-1 β [26]. In accordance with the phenotypic responses observed in IL-1R1 $^{-/-}$ mice, IL-1RAcP $^{-/-}$ mice show reduced neuroimmune and febrile responses to IL-1 [27, 28]. Finally, IL-1R2 $^{-/-}$ mice in which exon 2–4 are deleted using conventional gene targeting method have also been generated [29]. These mice show increased susceptibility to collagen-induced arthritis, while IL-1 β -induced cytokine response was enhanced in macrophages. In agreement with its inhibitory function, IL-1Ra $^{-/-}$ mice develop spontaneous autoimmune arthritis [30] and psoriasis-like cutaneous inflammation [31] and show increased brain injury to experimental stroke [32] and atherosclerotic lesion in experimental atherosclerosis [33]. Taken together, these observations demonstrate the complexity of the IL-1 system and point to important, yet undiscovered, mechanisms of actions of IL-1 ligands and their receptors, which cannot be explored by using classical pharmacological or genetic approaches.

Generation of a new toolbox to allow cell-specific conditional deletion of IL-1 ligands and their receptors

Germline gene deletion in mice has yielded important discoveries regarding the role of IL-1 ligands and their receptors in various inflammatory paradigms. However, this approach has important limitations such as viability and fertility of progeny,

subtle phenotypic changes, and/or compensatory mechanisms that may alter steady-state immune responses. In relation to the IL-1 system, IL-1 $\alpha^{-/-}$, IL-1 $\beta^{-/-}$, IL-1R1 $^{-/-}$, IL-1RAcP $^{-/-}$, and IL-1R2 $^{-/-}$ mice have all been reported to be viable with no obvious altered fertility. However, some reports suggested that IL-1 regulates ovulation, oocyte maturation, and early embryonic development [34, 35], which could lead to long-term significant phenotypic changes. Indeed, normal bone growth and remodeling are altered in IL-1R1 $^{-/-}$ and IL-1 $\alpha/\beta^{-/-}$ mice [36, 37], whereas decrease body fat mass is reduced in IL-1Ra $^{-/-}$ mice [38], strongly suggesting that significant phenotypic changes occur after ubiquitous deletion of IL-1 family members. Finally, significant compensatory changes are known to occur after ubiquitous gene deletion, and microarray analysis demonstrated that expression of several genes is altered in IL-1R1 $^{-/-}$ mice [24]. To overcome these limitations, the Cre/loxP system that allows selective/conditional deletion of targeted genes was recently used to generate new mouse mutant lines to allow cell-specific conditional deletion of IL-1 ligands and its receptors in a Cre recombinase-dependent manner (loxP-flanked, abbreviated as fl/fl). To this end, we have recently reported the generation and characterization of new IL-1 $\alpha^{fl/fl}$ and IL-1 $\beta^{fl/fl}$ mouse lines [39, 40] generated from *Il1a*^{tm1a(EUCOMM)Wtsi} (clone EPD0822-4-H02) or *Il1b*^{tm1a(EUCOMM)Hmgu} (clone HEPD0840-8-E03) embryonic stem cells purchased from the European Mouse Mutant Cell Repository (EuMMCR). The full description of the gene targeting strategies for both IL-1 $\alpha^{fl/fl}$ and IL-1 $\beta^{fl/fl}$ mice as well as experimental procedure from initial culturing and microinjection of ES cells leading to the generation of mice allowing for the conditional deletion of IL-1 α and IL-1 β are published [39, 40]. In these new alleles, exon 4 of the *Il1a* gene (for IL-1 $\alpha^{fl/fl}$ mice) or exon 4–5 of the *Il1b* gene (for IL-1 $\beta^{fl/fl}$ mice) flanked with loxP sites can be deleted by Cre recombinase, leading to exon 4 or 4–5 deletion and generation of cell-specific IL-1 α and IL-1 β -deficient allele, respectively (Fig. 1A and B).

Recently, two new mouse mutants, allowing for the conditional deletion of *Il1r1* (IL-1R1^{fl/fl}), have been described. Robson and collaborators [41] have generated a new IL-1R1^{fl/fl}, in which exon 3–4 of the *Il1r1* gene (encoding part of the extracellular binding region) can be deleted by Cre recombination and demonstrated ubiquitous inhibition of IL-1R1 signaling by the crossing of the conditional allele to the CMV-Cre mice, which mediated recombination in early embryogenesis. Concomitantly, we have generated a new IL-1R1^{fl/fl} mouse (developed by Taconic, Cologne, Germany), in which exon 5 that also encodes part of the extracellular binding region of the receptor is flanked by LoxP sites [42] (Fig. 1C). In those two new IL-1R1 mutants, deletion of exon 3–4 or exon 5 inactivates the two previously described functional IL-1R1 gene transcripts (including the full-length IL-1R1 and truncated IL-1R3) upon Cre-mediated




Fig. 1 Generation of IL-1 $\alpha^{fl/fl}$, IL-1 $\beta^{fl/fl}$, and IL-1R1^{fl/fl} mice. **A** Exon 4 of the *Il1a* gene (for IL-1 $\alpha^{fl/fl}$), **B** exon 4–5 of the *Il1b* gene (for IL-1 $\beta^{fl/fl}$ mice), or **C** exon 5 of the *Il1r1* gene (for IL-1R1^{fl/fl}) flanked with loxP sites is excised upon Cre recombination, resulting in cell-specific IL-1 α -, IL-1 β -, or IL-1R1-deficient allele, respectively

recombination [25]. In our study, we have also reported the generation of a new ubiquitous IL-1R1 $^{-/-}$ mouse as well as myeloid cell-specific IL-1R1-deficient mice by crossing IL-1R1^{fl/fl} with mice expressing Cre recombinase under the promoter of keratin 14 (K14-Cre) and the Vav promoter, respectively [42]. Of importance, an advanced genetic tool based on restoration of *Il1r1* gene expression has been developed by Liu and collaborators [43]. In this advanced model, excision of a disruptive intronic sequence in the *Il1r1* gene under Cre recombination in a global IL-1R1 $^{-/-}$ background allows functional restoration of IL-1 signaling under cell-specific promoters and has been important in the discovery of mechanisms of IL-1 signaling in the brain in the broad context IL-1-driven central inflammation [44]. Finally, generation of IL-1R2^{fl/fl} targeting exon 3 of the *Il1r2* gene was also reported [45], and we now report in our hand the generation of a similar IL-1R2^{fl/fl} mouse that targets exon 3 of *Il1r2* gene and further generation of IL-1R2 $^{-/-}$ by crossing IL-1R2^{fl/fl} with mice expressing Cre recombinase under the promoter of keratin 14 (K14-Cre) (Fig. 2A).

Generation of IL-1R2^{fl/fl} and IL-1R2 $^{-/-}$ mice

IL-1R2 conditional (IL-1R2^{fl/fl}) mice were generated at Taconic (Cologne, Germany) by gene targeting using BAC clones as the targeting vector from the C57BL/6J RPCI-23 BAC library encoding two loxP sites flanked exon 3 of the




Fig. 2 Generation of IL-1R2^{fl/fl} and IL-1R2^{-/-} mice. **A** Genetic approach to generate IL-1R2^{fl/fl} mice was designed to induce deletion of exon 3 encoding part of the extracellular binding domain, generating a frameshift from exon 4 to all downstream exons leading to genetic inhibition of IL-1R2. **B** Genotyping identification of IL-1R2^{fl/fl} mice was carried out by PCR using the following primers: Forward, TGTCTCCATCAGAC TGACTTTAGG, depicted (1), and reverse, ACCATGTCTGCCTG TTCACC, depicted (2) on genomic DNA. Amplification product size obtained was as follows: wild type (228 bp) and IL-1R2^{fl/fl} (347 bp). **C** Genotypic identification of exon 3 deletion in IL-1R2^{-/-} mice (obtained by crossing IL-1R2^{fl/fl} mice with mice expressing Cre recombinase under a keratin 14 promoter) was carried out by PCR on isolated genomic DNA using the following primers: Forward, GTAGTGGGCAATCA GATGGAC, depicted (3), and reverse, ACCATGTCTGCCTG TTCACC, depicted (2). Amplification product size obtained was 300 bp in the IL-1R2^{-/-} mice after Cre recombination

Il1r2 gene and subsequent homologous recombination in C57BL/6 N embryonic stem (ES) cells. From targeted C57BL/6 N ES cells, as verified by southern blotting, chimeric mice were generated and bred to C57BL/6 females. Germline transmission was identified by genotyping PCR sample analysis using a Caliper LabChip GX device (details are available upon request). Genotyping identification of IL-1R2^{fl/fl} mice was carried out on genomic DNA by PCR (see Fig. 2B) (details of primers used in Fig. 2 legend and protocol of DNA amplification available upon request). Amplification product size obtained were as follows: wild type (228 bp) and IL-1R2^{fl/fl} (347 bp).

A new ubiquitous IL-1R2^{-/-} mouse was generated by crossing the IL-1R2^{fl/fl} mice with mice expressing Cre recombinase under the control of the human keratin 14 promoter in oocytes as described [46], leading to genetic deletion of exon 3 in all tissues. The deletion of exon 3 causes a frame shift from exon 4 to all downstream exons. Genotypic

identification of exon 3 deletion in IL-1R2^{-/-} mice was carried out by PCR on isolated genomic DNA (see Fig. 2C) (details of primers used in figure legend and protocol of DNA amplification available upon request).

Advanced discoveries in mechanisms of IL-1 actions in disease using the toolbox

The new toolbox comprising IL-1 α ^{fl/fl}, IL-1 β ^{fl/fl}, IL-1R1^{fl/fl}, and IL-1R2^{fl/fl} mice allows for the first time the generation of new mouse lines in which IL-1 and its receptors can be deleted in a cell/tissue specific manner. To date, IL-1 α ^{fl/fl}, IL-1 β ^{fl/fl}, IL-1R1^{fl/fl}, and IL-1R2^{fl/fl} mice have been crossed with specific Cre drivers leading to cell- or tissue-specific deletion of either gene in a constitutive (Cre) or inducible (Cre-ER) manner, revealing new mechanisms of IL-1 actions. Table 1 provides a list of the cell-/tissue-specific deficient mice in IL-1 α , IL-1 β , IL-1R1, and IL-1R2 generated to date that have been tested under different inflammatory paradigms. While the generation of cell-specific IL-1 α and IL-1 β -deficient lines is limited, due to the recent generation of IL-1 α ^{fl/fl} and IL-1 β ^{fl/fl} mice, the first studies using those models have demonstrated the critical role of microglial IL-1 β in the establishment of pain in the context of complex regional pain syndrome [39], while cardiomyocyte-derived IL-1 α has been found not to contribute to tissue remodeling during myocardial infarction [40]. In contrast, IL-1R1^{fl/fl} mice have generated various cell-/tissue-specific IL-1R1-deficient lines, most studies showing a critical role for IL-1 signaling in immune cell activation and vascular activation in various models of peripheral infection and chronic inflammation (see Table 1). For instance, IL-1 signaling in cells of the hematopoietic lineage is required for the IL-17 and IL-22 response to gut infection by the nematode *Trichuris Muris* [42], whereas IL-1 signaling in T cells [50] and in GM-CSF producing cells [52] plays a critical role in experimental autoimmune encephalomyelitis. Furthermore, IL-1 signaling in T cells plays a key role in the systemic immune response to injection of CD3 antibodies [49]. Inhibition of IL-1 signaling by ColVI-Cre driver in intestinal mesenchymal cells showed that IL-1R1 in these cells has no important role in the development of intestinal carcinogenesis [60]. Furthermore, no direct role of IL-1 signaling in CD45⁺ hematopoietic was found in IL-1-mediated resistance to *Mycobacterium tuberculosis* [62]. In contrast, IL-1R1 on hepatocytes reduces liver injury in a model of acute liver failure [61], while deletion of IL-1R1 in pancreatic cells alters glucose homeostasis and triggers β -cell de-differentiation [57]. Finally, a study using Ly6G-Cre mice found that IL-1R1 in neutrophils plays a key role in reducing the tumorigenic effects of IL-1 [51]. In peripheral vascular beds, cadherin-Cre mediated IL-1R1 deletion in endothelial cells contributes to the anti-tumor function of adoptively transferred T cells

Table 1 List of cell-/organ-specific deficient mice for IL-1 isoforms or their receptors and main effects observed

Gene	Cre driver	Cell/tissue targeted	Main effects	References
Il1a	Myh6-Cre	Cardiomyocytes	No effect on cardiac tissue remodeling after MI	Bageghni et al. [40]
Il1b	CMV-Cre	Ubiquitous deletion	Reduces bone metastasis during breast cancer	Tulotta et al. [47]
	CX3CR1-CreER	Microglial cell	Reduces the development of pain	[39]
Il1r1	K14-Cre	Ubiquitous deletion	Mediates peripheral immune response to <i>T. Muris</i> infection Inhibits melanoma inflammatory niche	[42] Young et al. [48]
	CMV-Cre	Ubiquitous deletion	Decreases inflammatory responses to systemic challenges NI	Mufazalov et al. [49]; Mufazalov et al. [50] Robson et al. [41]
	PGK-Cre	Ubiquitous deletion	Regulates cardiac tissue remodeling after MI	Bageghni et al. [40]
	Col1a2-CreER	Fibroblasts	Regulates cardiac tissue remodeling after MI	Bageghni et al. [40]
	CD4-Cre	T cells	Regulates immune response to CD3 antibody injection Regulates neuroinflammation in EAE	Mufazalov et al. [49] Mufazalov et al. [50]
	Vav-Cre	Myeloid cells	Mediates peripheral immune response to T. Muris infection	Adbulaal et al. [42]
	Ly6G-Cre	Neutrophils	Reduces the tumorigenic effect of IL-1	Dmitrieva-Posocco et al. [51]
	Csf2-Cre	GM-CSF positive cells	Regulates inflammation after EAE	Komuczki et al. [52]
	CX3CR1-CreER	Microglial cells	Reduces renewal of microglial population	Bruttger et al. [53]
			Regulates microglial activation after CNS inflammation	Zhu et al. [54]*
			No effect on febrile response to IL-1	Knoll et al. [55]*
	Myh11-CreER	Smooth muscle cells	Reduces atheroprotective effect of IL-1 after atherosclerosis	Gomez et al. [56]
	Pdx1-Cre	Pancreatic cells	Alters glucose homeostasis and triggers β -cell de-differentiation	Burke et al. [57]
	Slco1c1-CreER	Brain endothelial cells	Reduces CNS inflammation and brain damage after stroke	Wong et al. [58]
			Reduces fever response to IL-1	Matsuwaki et al. [59]
	Nestin-Cre	Neuronal cells	No effect on febrile response to IL-1	Matsuwaki et al. [59]
			Reduces brain damage after stroke	Wong et al. [58]
	Trpv1-Cre	Nociceptor sensory neurons	No effect on febrile response to IL-1	Matsuwaki et al. [59]
	htPA-Cre	Neural crest cells	No effect on febrile response to IL-1	Matsuwaki et al. [59]
	ChAT-Cre	Catecholaminergic neurons	Decreases brain damage after stroke	Wong et al. [58]
	PF4-Cre	Platelets	No effect on brain damage after stroke	Wong et al. [58]
	ColVI-CreER	Intestinal mesenchymal cells	No effect on development of intestinal cancer	Koliaraki et al. [60]
	Hep-Cre	Hepatocytes	Reduces liver injury after acute liver failure	Gehrke et al. [61]
	CD45-Cre	Leukocytes	No role in IL-1-mediated resistance to <i>Mycobacterium tuberculosis</i>	Bohrer et al. [62]
	Cdh5-Cre	Vascular endothelial cells	Mediates the anti-tumor properties of T cells. Regulates IL-1-induced brain inflammation	Lee et al. [63]
	Tie2-Cre	Endothelial cells	Decreases IL-1-induced brain inflammation	Liu et al. [44] [#]
Il1r2	CMV-Cre	Ubiquitous deletion	Reduces inflammation after arthritis	Martin et al. [45]
	K14-Cre	Ubiquitous deletion	NI	Unpublished**

For IL-1R1^{fl/fl} mice, all cell-/tissue-specific IL-1R1^{-/-} mice have been generated by IL-1R1^{fl/fl} mouse from Abdulaal et al. [42], except those marked (*), generated by IL-1R1^{fl/fl} mouse from Robson et al. [41]. [#]In the study of Liu and collaborators (2019), the following mouse lines have also been generated: IL-1R1^{fl/fl} x LysM-Cre, IL-1R1^{fl/fl} x CX3CR1-Cre, IL-1R1^{fl/fl} x Camk2a-Cre, IL-1R1^{fl/fl} x Vglut2-Cre, and IL-1R1^{fl/fl} x GFAP-Cre. **IL-1R2^{fl/fl} mice crossed with K14-Cre mice are reported in the present publication. *Cre*, constitutive deletion by Cre drivers; *Cre-ER*, inducible deletion by Cre-ER drivers; *EAE*, experimental autoimmune encephalomyelitis; *MI*, myocardial infarction; *NI*, not indicated

regulated by IL-1 β [63], whereas IL-1R1 signaling in smooth muscle cells contributes to the atheroprotective effect of IL-1 in advanced atherosclerotic lesions [56].

In the brain, IL-1 signaling in microglia is required for the renewal properties of microglial progenitor cells [53]. Brain endothelial IL-1R1 is essential in the initiation of the fever

response elicited by IL-1, whereas deletion of IL-1R1 on central or peripheral neurons (including catecholaminergic neurons and nociceptor sensory neurons) had no noticeable effect on the febrile response [59]. An interesting work by Knoll and collaborators [55] confirmed that endothelial IL-1R1 signaling is critical in the establishment of the febrile response to IL-1, whereas IL-1R1 signaling in microglia of the brain parenchyma has no role. Further, endothelial IL-1R1 is essential for endothelial activation in the context of IL-1-driven brain inflammation [44] and a further study using microglial-specific IL-1R1-deficient mice showed that IL-1 actions on brain endothelia triggers the production of endothelial-derived factors that are able to activate microglial cells [54]. In the context of stroke, brain endothelial IL-1R1, but also neuronal IL-1R1, is critical in mechanisms of cerebrovascular inflammation and brain damage after experimental cerebral ischemia, whereas no involvement of IL-1 signaling in peripheral cells, including platelets on stroke outcome was observed [58].

Finally, little work has been conducted regarding IL-1R2^{fl/fl} mice, and to date, only ubiquitous constitutive deletion of IL-1R2 (IL-1R2^{-/-}) has been achieved, including that of our work. The only work reporting the use of IL-1R2^{-/-} in disease is that of Martin and collaborators [45], who demonstrated that IL-1R2 deletion plays an important inhibitory role on IL-1-regulated inflammation in a model of arthritis, in accordance with its known inhibitory function, as recently reviewed [64].

Concluding remarks and future directions

IL-1 is a key cytokine regulating many physiological processes as well as the inflammatory responses to infection or injury, and global constitutive IL-1-deficient mouse models, which is a fairly recent approach, have to date helped unraveling key mechanisms of IL-1 actions in disease but have significant limitations. The recent generation of new mouse mutants allowing conditional deletion of IL-1 and its receptors has led to the discovery of unexpected new mechanisms of inflammation regulated by IL-1. To date, a limited number of cell-/tissue-specific IL-1-deficient mice have been generated, and this is mainly due to the fact that IL-1 α ^{fl/fl}, IL-1 β ^{fl/fl}, IL-1R1^{fl/fl}, and IL-1R2^{fl/fl} mice have only been recently produced. However, several projects targeting the IL-1 system in other cells/tissues are currently ongoing. Importantly, to the best of our knowledge, IL-1Ra^{fl/fl} and IL-1RAcP^{fl/fl} mice have not been generated yet, and future generation of new lines in which all IL-1 ligands and their receptors can be targeted in other cell/tissue and other disease models will lead to new mechanisms to be discovered, providing a step change in our understanding of IL-1 actions disease and the potential development of new targeted IL-1 therapies.

Acknowledgements The authors would like to thank the Transgenic Facility of the University of Manchester and Dr. Elena Redondo-Castro for helping with the culturing of ES cells for the generation of IL-1 α ^{fl/fl} and IL-1 β ^{fl/fl} mice.

Funding information E.P. and S.F. received funding from the British Heart Foundation (BHF), grant PG/13/55/30365 for the generation of IL-1 α ^{fl/fl} and IL-1 β ^{fl/fl} mice. A.W. is a member of the Research Center for Immunotherapy (FZI) Mainz and was supported by the Deutsche Forschungsgemeinschaft (DFG) grants CRC/TRR128 and CRC1292. W.A. received funding from King Abdulaziz University, Jeddah, Kingdom of Saudi Arabia. W.M. received funding from the European Union (FP7/2012–2017) under grant agreement n° 305,564, SysmedIBD. I.A.M. received intramural funding (Stufe1) provided by the University of Mainz.

Compliance with ethical standards

Conflict interests The authors declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Blomberg L, Hashizume K, Viebahn C (2008) Blastocyst elongation, trophoblastic differentiation, and embryonic pattern formation. *Reproduction* 135:181–195
2. Gadek-Michalska A, Tadeusz J, Rachwalska P, Bugajski J (2013) Cytokines, prostaglandins and nitric oxide in the regulation of stress-response systems. *Pharmacol Rep* 65:1655–1662
3. Marshall L, Born J (2002) Brain-immune interactions in sleep. *Int Rev Neurobiol* 52:93–131
4. Mantovani A, Dinarello CA, Molgora M, Garlanda C (2019) Interleukin-1 and related cytokines in the regulation of inflammation and immunity. *Immunity* 50:778–795
5. Dinarello CA (2018) Introduction to the interleukin-1 family of cytokines and receptors: drivers of innate inflammation and acquired immunity. *Immunol Rev* 281:5–7
6. McMahan CJ, Slack JL, Mosley B et al (1991) A novel IL-1 receptor, cloned from B cells by mammalian expression, is expressed in many cell types. *EMBO J* 10:2821–2832
7. Brough D, Denes A (2015) Interleukin-1 α and brain inflammation. *IUBMB Life* 67:323–330
8. Anforth HR, Bluthé RM, Bristow A et al (1998) Biological activity and brain actions of recombinant rat interleukin-1 α and interleukin-1 β . *Eur Cytokine Netw* 9:279–288
9. Benjamin JT, Moore DJ, Bennett C, van der Meer R, Royce A, Loveland R, Wynn JL (2018) Cutting edge: IL-1 α and not IL-1 β drives IL-1R1-dependent neonatal murine sepsis lethality. *J Immunol* 201:2873–2878

10. Nakae S, Naruse-Nakajima C, Sudo K et al (2001) IL-1 alpha, but not IL-1 beta, is required for contact-allergen-specific T cell activation during the sensitization phase in contact hypersensitivity. *Int Immunol* 13:1471–1478
11. Saini MG, Pinteaux E, Lee B, Bix GJ (2011) Oxygen-glucose deprivation and interleukin-1 α trigger the release of perlecan LG3 by cells of neurovascular unit. *J Neurochem* 119:760–771
12. Um JY, Moon KS, Lee KM, Yun JM, Cho KH, Moon BS, Kim HM (2003a) Association of interleukin-1 alpha gene polymorphism with cerebral infarction. *Mol Brain Res* 115:50–54
13. Um JY, Moon KS, Lee KM, Kim HM (2003b) Interleukin-1 gene cluster polymorphisms in cerebral infarction. *Cytokine* 23:41–46
14. Tsakiri N, Kimber I, Rothwell NJ, Pinteaux E (2008) Differential effects of interleukin-1 alpha and beta on interleukin-6 and chemokine synthesis in neurons. *Mol Cell Neurosci* 38:259–265
15. Chamberlain J, Evans D, King A, Dewberry R, Dower S, Crossman D, Francis S (2006) Interleukin-1beta and signaling of interleukin-1 in vascular wall and circulating cells modulates the extent of neointima formation in mice. *Am J Pathol* 168:1396–1403
16. Bersudsky M, Luski L, Fishman D, White RM, Ziv-Sokolovskaya N, Dotan S, Rider P, Kaplanov I, Aychek T, Dinarello CA, Apte RN, Voronov E (2014) Non-redundant properties of IL-1 α and IL-1 β during acute colon inflammation in mice. *Gut* 63:598–609
17. Horai R, Asano M, Sudo K, Kanuka H, Suzuki M, Nishihara M, Takahashi M, Iwakura Y (1998) Production of mice deficient in genes for interleukin (IL)-1 α , IL-1 β , IL-1 α/β , and IL-1 receptor antagonist shows that IL-1 β is crucial in turpentine-induced fever development and glucocorticoid secretion. *J Exp Med* 187:1463–1475
18. Glaccum MB, Stocking KL, Charrier K et al (1997) Phenotypic and functional characterization of mice that lack the type I receptor for IL-1. *J Immunol* 159:3364–3371
19. Boutin H, Kimber I, Rothwell NJ, Pinteaux E (2003) The expanding interleukin-1 family and its receptors: do alternative IL-1 receptor/signaling pathways exist in the brain? *Mol Neurobiol* 27:239–248
20. Humphreys NE, Grecnis RK (2009) IL-1-dependent, IL-1R1-independent resistance to gastrointestinal nematodes. *Eur J Immunol* 39:1036–1045
21. Touzani O, Boutin H, Lefeuvre R et al (2002) Interleukin-1 influences ischemic brain damage in the mouse independently of the interleukin-1 type I receptor. *J Neurosci* 22:38–43
22. Rincón-López C, Tlapa-Pale A, Medel-Matus J-S, Martínez-Quiroz J, Rodríguez-Landa JF, López-Meraz ML (2017) Interleukin-1 β increases neuronal death in the hippocampal dentate gyrus associated with status epilepticus in the developing rat. *Neurologia* 32:587–594
23. Diem R, Hobom M, Grötsch P, Kramer B, Bähr M (2003) Interleukin-1 β protects neurons via the interleukin-1 (IL-1) receptor-mediated Akt pathway and by IL-1 receptor-independent decrease of transmembrane currents in vivo. *Mol Cell Neurosci* 22:487–500
24. Andre R, Moggs JG, Kimber I, Rothwell NJ, Pinteaux E (2006) Gene regulation by IL-1beta independent of IL-1R1 in the mouse brain. *Glia* 53:477–483
25. Qian J, Zhu L, Li Q, Belevych N, Chen Q, Zhao F, Herness S, Quan N (2012) Interleukin-1R3 mediates interleukin-1-induced potassium current increase through fast activation of Akt kinase. *Proc Natl Acad Sci U S A* 109:12189–12194
26. Cullinan EB, Kwee L, Nunes P et al (1998) IL-1 receptor accessory protein is an essential component of the IL-1 receptor. *J Immunol* 161:5614–5620
27. Liège S, Layé S, Li KS, Moze E, Neveu PJ (2000) Interleukin 1 receptor accessory protein (IL-1RAcP) is necessary for centrally mediated neuroendocrine and immune responses to IL-1 β . *J Neuroimmunol* 110:134–139
28. Zetterström M, Lundkvist J, Malinowsky D, Eriksson G, Bartfai T (1998) Interleukin-1-mediated febrile responses in mice and interleukin-1 beta activation of NF κ B in mouse primary astrocytes, involves the interleukin-1 receptor accessory protein. *Eur Cytokine Netw* 9:131–138
29. Shimizu K, Nakajima A, Sudo K, Liu Y, Mizoroki A, Ikarashi T, Horai R, Kakuta S, Watanabe T, Iwakura Y (2015) IL-1 receptor type 2 suppresses collagen-induced arthritis by inhibiting IL-1 signal on macrophages. *J Immunol* 194:3156–3168
30. Akitsu A, Ishigame H, Kakuta S, Chung SH, Ikeda S, Shimizu K, Kubo S, Liu Y, Umemura M, Matsuzaki G, Yoshikay Y, Saijo S, Iwakura Y (2015) IL-1 receptor antagonist-deficient mice develop autoimmune arthritis due to intrinsic activation of IL-17-producing CCR2 + V γ 6 + γ δ T cells. *Nat Commun* 6:7464
31. Shepherd J, Little MC, Nicklin MJH (2004) Psoriasis-like cutaneous inflammation in mice lacking interleukin-1 receptor antagonist. *J Invest Dermatol* 122:665–669
32. Pinteaux E, Rothwell NJ, Boutin H (2006) Neuroprotective actions of endogenous interleukin-1 receptor antagonist (IL-1ra) are mediated by glia. *Glia* 53:551–556
33. Isoda K, Sawada S, Ishigami N, Matsuki T, Miyazaki K, Kusuhara M, Iwakura Y, Ohsuzu F (2004) Lack of interleukin-1 receptor antagonist modulates plaque composition in apolipoprotein E-deficient mice. *Arterioscler Thromb Vasc Biol* 24:1068–1073
34. Gérard N, Caillaud M, Martoriati A et al (2004) The interleukin-1 system and female reproduction. *J Endocrinol* 180:203–212
35. Caillaud M, Duchamp G, Gérard N (2005) In vivo effect of interleukin-1 beta and interleukin-1 RA on oocyte cytoplasmic maturation, ovulation, and early embryonic development in the mare. *Reprod Biol Endocrinol* 3:26
36. Simsa-Maziel S, Zaretsky J, Reich A, Koren Y, Shahar R, Monsonego-Oman E (2013) IL-1RI participates in normal growth plate development and bone modeling. *Am J Physiol Endocrinol Metab* 305:E15–E21
37. Lee YM, Fujikado N, Manaka H, Yasuda H IY IL-1 plays an important role in the bone metabolism under physiological conditions. - PubMed - NCBI. <https://www.ncbi.nlm.nih.gov/pubmed/?term=20679512>. Accessed 15 May 2020
38. Somm E, Henrichot E, Pemin A, Juge-Aubry CE, Muzzin P, Dayer JM, Nicklin MJH, Meier CA (2005) Decreased fat mass in interleukin-1 receptor antagonist-deficient mice: impact on adipogenesis, food intake, and energy expenditure. *Diabetes* 54:3503–3509
39. Helyes Z, Tékus V, Szentés N, Pohóczky K, Botz B, Kiss T, Kemény Á, Környei Z, Tóth K, Lénárt N, Ábrahám H, Pinteaux E, Francis S, Sensi S, Dénes Á, Goebel A (2019) Transfer of complex regional pain syndrome to mice via human autoantibodies is mediated by interleukin-1-induced mechanisms. *Proc Natl Acad Sci U S A* 116:13067–13076
40. Bageghni SA, Hemmings KE, Yuldasheva NY et al (2019) Fibroblast-specific deletion of interleukin-1 receptor-1 reduces adverse cardiac remodeling following myocardial infarction. *JCI insight* 5
41. Robson MJ, Bin ZC, Quinlan MA et al (2016) Generation and characterization of mice expressing a conditional allele of the interleukin-1 receptor type 1. *PLoS One* 11:e0150068
42. Abdulaal WH, Walker CR, Costello R, Redondo-Castro E, Mufazalov IA, Papaemmanouil A, Rothwell NJ, Allan SM, Waisman A, Pinteaux E, Müller W (2016) Characterization of a conditional interleukin-1 receptor 1 mouse mutant using the Cre/LoxP system. *Eur J Immunol* 46:912–918
43. Liu X, Yamashita T, Chen Q et al (2015) Interleukin 1 type 1 receptor restore: a genetic mouse model for studying interleukin 1 receptor-mediated effects in specific cell types. *J Neurosci* 35:2860–2870. h

44. Liu X, Nemeth DP, McKim DB et al (2019) Cell-type-specific interleukin 1 receptor 1 signaling in the brain regulates distinct neuroimmune activities. *Immunity* 50:317–333.e6
45. Martin P, Palmer G, Rodriguez E, Seemayer CA, Palomo J, Talbot-Ayer D, Gabay C (2017) Deficiency in IL-1 receptor type 2 aggravates K/BxN serum transfer-induced arthritis in mice but has no impact on systemic inflammatory responses. *J Immunol* 198:2916–2926
46. Hafner M, Wenk J, Nenci A, Pasparakis M, Scharffetter-Kochanek K, Smyth N, Peters T, Kess D, Holtkötter O, Shephard P, Kudlow JE, Smola H, Haase I, Schippers A, Krieg T, Müller W (2004) Keratin 14 Cre transgenic mice authenticate keratin 14 as an oocyte-expressed protein. *Genesis* 38:176–181
47. Tulotta C, Lefley DV, Freeman K, et al (2019) Endogenous production of IL-1beta by breast cancer cells drives metastasis and colonization of the bone microenvironment. *Clin Cancer Res* 25:2769–2782
48. Young HL, Rowling EJ, Bugatt M et al (2017) An adaptive signaling network in melanoma inflammatory niches confers tolerance to MAPK signaling inhibition. *J Exp Med* 214:1691–1710
49. Mufazalov IA, Regen T, Schelmbauer C, Kuschmann J, Muratova AM, Nikolaev A, Müller W, Pinteaux E, Waisman A (2016) Generation of a novel T cell specific interleukin-1 receptor type 1 conditional knock out mouse reveals intrinsic defects in survival, expansion and cytokine production of CD4 T cells. *PLoS One* 11:e0161505
50. Mufazalov IA, Schelmbauer C, Regen T, Kuschmann J, Wanke F, Gabriel LA, Hauptmann J, Müller W, Pinteaux E, Kurschus FC, Waisman A (2017) IL-1 signaling is critical for expansion but not generation of autoreactive GM-CSF+ Th17 cells. *EMBO J* 36:102–115
51. Dmitrieva-Posocco O, Dzutsev A, Posocco DF et al (2019) Cell-type-specific responses to interleukin-1 control microbial invasion and tumor-elicited inflammation in colorectal cancer. *Immunity* 50:166–180.e7
52. Komuczki J, Tuzlak S, Friebe E et al (2019) Fate-mapping of GM-CSF expression identifies a discrete subset of inflammation-driving T helper cells regulated by cytokines IL-23 and IL-1β. *Immunity* 50:1289–1304.e6
53. Bruttger J, Karram K, Wörtge S, Regen T, Marini F, Hoppmann N, Klein M, Blank T, Yona S, Wolf Y, Mack M, Pinteaux E, Müller W, Zipp F, Binder H, Bopp T, Prinz M, Jung S, Waisman A (2015) Genetic cell ablation reveals clusters of local self-renewing microglia in the mammalian central nervous system. *Immunity* 43:92–106
54. Zhu L, Liu X, Nemeth DP, DiSabato DJ, Witcher KG, Mckim DB, Oliver B, le X, Gorantla G, Berdysz O, Li J, Ramani AD, Chen Z, Wu D, Godbout JP, Quan N (2019) Interleukin-1 causes CNS inflammatory cytokine expression via endothelia-microglia bi-cellular signaling. *Brain Behav Immun* 81:292–304
55. Knoll GJ, Krasnow SM, Marks DL (2017) Interleukin-1β signaling in fenestrated capillaries is sufficient to trigger sickness responses in mice. *J Neuroinflammation* 14: 219
56. Gomez D, Baylis RA, Durgin BG, Newman AAC, Alencar GF, Mahan S, St. Hilaire C, Müller W, Waisman A, Francis SE, Pinteaux E, Randolph GJ, Gram H, Owens GK (2018) Interleukin-1β has atheroprotective effects in advanced atherosclerotic lesions of mice. *Nat Med* 24:1418–1429
57. Burke SJ, Batdorf HM, Burk DH, Martin TM, Mendoza T, Stadler K, Alami W, Karlstad MD, Robson MJ, Blakely RD, Mynatt RL, Collier JJ (2018) Pancreatic deletion of the interleukin-1 receptor disrupts whole body glucose homeostasis and promotes islet β-cell de-differentiation. *Mol Metab* 14:95–107
58. Wong R, Lénárt N, Hill L, Toms L, Coutts G, Martinecz B, Császár E, Nyiri G, Papaemmanouil A, Waisman A, Müller W, Schwaninger M, Rothwell N, Francis S, Pinteaux E, Denés A, Allan SM (2019) Interleukin-1 mediates ischaemic brain injury via distinct actions on endothelial cells and cholinergic neurons. *Brain Behav Immun* 76:126–138
59. Matsuwaki T, Shionoya K, Ilnatko R, Eskilsson A, Kakuta S, Dufour S, Schwaninger M, Waisman A, Müller W, Pinteaux E, Engblom D, Blomqvist A (2017) Involvement of interleukin-1 type 1 receptors in lipopolysaccharide-induced sickness responses. *Brain Behav Immun* 66:165–176
60. Koliaraki V, Chalkidi N, Henriques A et al (2019) Innate sensing through mesenchymal TLR4/MyD88 signals promotes spontaneous intestinal tumorigenesis. *Cell Rep* 26:536–545.e4
61. Gehrke N, Hövelmeyer N, Waisman A, Straub BK, Weinmann-Menke J, Wörns MA, Galle PR, Schattenberg JM (2018) Hepatocyte-specific deletion of IL1-RI attenuates liver injury by blocking IL-1 driven autoinflammation. *J Hepatol* 68:986–995
62. Bohrer AC, Tocheny C, Assmann M, Ganusov VV, Mayer-Barber KD (2018) Cutting edge: IL-1R1 mediates host resistance to mycobacterium tuberculosis by trans-protection of infected cells. *J Immunol* 201:1645–1650
63. Lee PH, Yamamoto TN, Gurusamy D, Sukumar M, Yu Z, Hu-Li J, Kawabe T, Gangaplara A, Kishton RJ, Henning AN, Vodnala SK, Germain RN, Paul WE, Restifo NP (2019) Host conditioning with IL-1β improves the antitumor function of adoptively transferred T cells. *J Exp Med* 216:2619–2634
64. Schlüter T, Schelmbauer C, Karram K, Mufazalov IA (2018) Regulation of IL-1 signaling by the decoy receptor IL-1R2. *J Mol Med (Berl)* 96:983–992

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.