



This is a repository copy of *Quasi-parallel propagation of solitary waves in magnetised non-relativistic electron–positron plasmas*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/162822/>

Version: Accepted Version

---

**Article:**

Ruderman, M.S. [orcid.org/0000-0003-2324-8466](https://orcid.org/0000-0003-2324-8466) (2020) Quasi-parallel propagation of solitary waves in magnetised non-relativistic electron–positron plasmas. *Journal of Plasma Physics*, 86 (3). 905860311. ISSN 0022-3778

<https://doi.org/10.1017/s0022377820000483>

---

This article has been published in a revised form in *Journal of Plasma Physics* [<https://doi.org/10.1017/S0022377820000483>]. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works. © The Author(s).

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# Quasi-parallel propagation of solitary waves in magnetised non-relativistic electron-positron plasmas

MICHAEL S. RUDERMAN<sup>1,2,3</sup>

<sup>1</sup>School of Mathematics and Statistics, University of Sheffield, Hicks Building, Hounsfield Road, Sheffield S3 7RH, UK

<sup>2</sup>Space Research Institute (IKI), Russian Academy of Sciences, Moscow, Russia

<sup>3</sup>Moscow Center for Fundamental and Applied Mathematics

email: m.s.ruderman@sheffield.ac.uk

(Received 28 April 2020)

**Abstract.** We study the propagation of nonlinear waves in non-relativistic electron-positron plasmas. The waves are assumed to propagate at small angles with respect to the equilibrium magnetic field. We derive the equation describing the wave propagation under the assumption that the waves are weakly dispersive and also can weakly depend on spatial variables orthogonal to the equilibrium magnetic field. We obtain solutions of the derived equation describing solitons. Then we study the stability of solitons with respect to transverse perturbations.

---

## 1. Introduction

The problem of wave propagation in electron-positron plasmas attracted the attention of theorists for a few decades first of all in relation with astrophysical applications. It is believed that in astrophysics electron-positron plasmas exist in pulsar magnetospheres (Sturrock 1971, Ruderman and Sutherland 1975, Chian and Kennel 1983, Arons and Barnard 1986, Aharonian et al. 2012, Cerutti and Beloborodov 2017), active galactic nuclei (Ruffini et al. 2010, El-Labany et al. 2013, Kawakatu et al. 2016) and early universe (Gailis et al. 1995, Shukla 2003, Tatsuno et al. 2003). It is believed that large-amplitude low-frequency waves play an important role in such astrophysical processes as slowing down of pulsars, pulsar radiation, and cosmic ray acceleration.

The linear theory of wave propagation in electron-positron plasmas was developed using both hydrodynamic as well as kinetic description (Arons and Barnard 1986, Sakai and Kawata 1980a, Stewart and Laing 1992). The nonlinear theory of waves in electron-positron plasmas has been also developed. The Nonlinear Schrödinger (NLS) equation was derived and used to study the modulational instability and envelope solitons (Chian and Kennel 1983, Cattaert et al. 2005, Rajib et al. 2015). The Korteweg-de Vries (KdV) and modified Korteweg-de Vries (mKdV) equations were obtained and the dependence of width and amplitude of solitons described by these equations on parameters of an unperturbed state was studied (Verheest and Lakhina 1996, Lakhina and Verheest 1997, Rajib et al. 2015).

We aim to study the propagation of nonlinear waves that is quasi-parallel with respect to the equilibrium magnetic field. In the case of electron-ion plasmas this problem was intensively studied during a few decades. It was shown that the one-dimensional quasi-parallel propagation of nonlinear waves is described by the Derivative Nonlinear Schrödinger (DNLS) equation (Rogister 1971, Mjølhus 1976, Mio et al. 1976a, Ruderman 2002). This equation was used to study the modulational instability of circularly polarised Alfvén waves (Mjølhus 1976, Mio et al. 1976b). The DNLS equation describes a few kinds of solitons as well as the generation of rogue waves (Ichikawa et al. 1980, Mjølhus and Hada 1997, Fedun et al. 2008). It was shown that the DNLS equation is completely integrable, the Lax pair for this equation was found, and the inverse scattering method was used to obtain exact solutions (Kaup and Newell 1978, Kawata and Inoue 1978).

Later an extension of the DNLS equations to two and three dimensions (3D DNLS) was derived (Mjølhus and Wyller 1986, Ruderman 1987, Mjølhus and Hada 1997). This extension is similar to that obtained by Kadomtsev and Petviashvili (1970) (KP equation) for the KdV equation. The 3D DNLS was used to study the stability of solitons of the DNLS equation with respect to transvers perturbations (Ruderman 1987, Mjølhus and Hada 1997).

The propagation of large-amplitude Alfvén waves parallel to the external magnetic field has been also studied in an electron-positron plasma (Sakai and Kawata 1980a, b, Mikhailovskii et al. 1985a, b, c, Verheest 1996, Lakhina and Verheest 1997). It was shown that, in contrast to the electron-proton plasma, nonlinear waves propagating parallel to the magnetic field are described by the vector form of the mKdV equation.

In this paper we aim to extend this vector mKdV equation to two and three dimensions. First studies of waves in electron-positron plasmas were related to astrophysical applications. However, then the progress of experimental physics opened the possibility of creation of electron-positron plasmas in laboratory (Surko et al. 1989, Surko and Murphy 1990, Greaves et al. 1994, Liang et al. 1998, Gahn et al. 2000, Bell and Kirk 2008, Chen et al. 2009, Sarri et al. 2013). Another example is the semi-conductor plasma, where holes behave like positive charges with the mass equal to that of electrons (Shukla et al. 1986). Although in astrophysical applications an electron-positron plasma is almost always relativistic, a non-relativistic electron-positron plasma is also of astrophysical interest. It can radiate very effectively by the cyclotron emission. As a result, it cools and eventually becomes non-relativistic. As for laboratory plasmas, in many cases they can be describing in the non-relativistic approximation. This observation inspired Iwamoto (1993) and Zank and Greaves (1995) to study waves in non-relativistic electron-positron plasmas.

The propagation of nonlinear waves and, in particular, solitons, electron-ion plasmas were extensively studied in laboratory experiments (e.g. Ikezi 1973, Tran 1979, Lonngren 1983). To our knowledge up to now there have been no experimental studies of waves in electron-positron plasmas. Apparently it is related to substantial difficulty of creating electron-positron plasmas in laboratory. Hence, theorists are ahead of experimentalists in studying waves in these plasmas. The state of affairs here is the same as was in the case of electron-ion plasmas where nonlinear waves were studied theoretically much earlier than experimentally. There is no doubt that waves in electron-positron plasmas will be studied experimentally because they are

of great importance for understanding physical phenomena both in astrophysical as well as in laboratory plasmas.

In this article we also use the non-relativistic approximation that strongly simplifies the derivation of the multi-dimensional generalisation of the mKdV equation. The article is organised as follows. In the next section we formulate the problem and present the governing equations. In Sect. 3 we briefly discuss the linear theory. In Sect. 4 we derive the equation describing small-amplitude weakly dispersive quasi-three-dimensional nonlinear waves. In Sect. 5 we obtain the solutions describing planar one-dimensional solitons. In Sect. 6 we study the soliton stability with respect to transvers perturbations. Section 7 contains the summary of the obtained results and conclusion.

## 2. Problem formulation and governing equations

We consider the propagation of nonlinear waves along the equilibrium magnetic field in a plasma that consists of electrons and positrons. We treat the electron and positron components as two charged fluids. We do not consider the annihilation or pair creation meaning that the particle number is conserved. We use the non-relativistic approximation meaning that the velocities of the two fluids are much smaller than the speed of light  $c$ , and the pressure of each fluid is much smaller than the density times  $c^2$ . We also assume that the phase speed of propagation of small perturbations is much smaller than  $c$ . The plasma motion is describing by the mass conservation and momentum equations:

$$\frac{\partial n_s}{\partial t} + \nabla \cdot (n_s \mathbf{v}_s) = 0, \quad (2.1a)$$

$$\frac{\partial \mathbf{v}_s}{\partial t} + (\mathbf{v}_s \cdot \nabla) \mathbf{v}_s + \frac{\nabla p_s}{m n_s} = \frac{q_s}{m} (\mathbf{E} + \mathbf{v}_s \times \mathbf{B}). \quad (2.1b)$$

In these equations  $n_s$  is the number density,  $\mathbf{v}_s$  the velocity,  $p_s$  the pressure,  $m$  the electron mass, and  $s = +$  and  $s = -$  refers to the positrons and electrons, respectively;  $\mathbf{E}$  is the electrical field,  $\mathbf{B}$  is the magnetic field,  $q_+ = q$ ,  $q_- = -q$ , and  $q$  is the elementary charge. We assume that the motion is adiabatic and take

$$p_s = p_0 \left( \frac{n_s}{n_0} \right)^\kappa, \quad (2.2)$$

where  $n_0$  and  $p_0$  are the unperturbed number density and pressure (the same for the electrons and positrons), and  $\kappa (= 5/3)$  is the adiabatic exponent. Equations (2.1a)–(2.2) must be supplemented with the Maxwell equations. Since we use the non-relativistic approximation, we can neglect the displacement current and write the Maxwell equations as

$$\nabla \cdot \mathbf{E} = \frac{\rho}{\varepsilon_0}, \quad (2.3a)$$

$$\nabla \cdot \mathbf{B} = 0, \quad (2.3b)$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}, \quad (2.3c)$$

$$\nabla \times \mathbf{B} = \mu_0 \mathbf{j}, \quad (2.3d)$$

where  $\varepsilon_0$  is the permittivity of free space,  $\mu_0$  is the permeability of free space, and the total electrical charge and current densities are determined by

$$\rho = \rho_+ + \rho_- = q(n_+ - n_-), \quad (2.4a)$$

$$\mathbf{j} = \mathbf{j}_+ + \mathbf{j}_- = q(n_+ \mathbf{v}_+ - n_- \mathbf{v}_-). \quad (2.4b)$$

Recall that  $\varepsilon_0 \mu_0 = c^{-2}$ .

We assume that in the equilibrium  $n_+ = n_- = n_0$ ,  $\mathbf{v}_+ = \mathbf{v}_- = 0$ ,  $\mathbf{E} = 0$ , and  $\mathbf{B} = B_0 \mathbf{e}_x$ , where  $\mathbf{e}_x$  is the unit vector along the  $x$ -axis of Cartesian coordinates  $x, y, z$ .

### 3. Linear theory

Here we briefly describe the linear theory of wave propagation because below we use it as a guide for scaling when deriving the equation governing the propagation of nonlinear waves. Since below we study the nonlinear wave propagation along the magnetic field, we only consider linear wave propagation in the equilibrium magnetic field direction. We linearise Eqs. (2.1) and (2.2) and then take perturbations of all quantities proportional to  $\exp[i(kx - \omega t)]$ . As a result, we obtain two disconnected systems of algebraic equations. The first system is for the perturbations of the number density, pressure, and  $x$ -components of the velocity and electric field. It describes the longitudinal wave mode. We do not study this mode in detail and only state that in the long wavelength approximation its phase speed is  $a_0 = (\kappa p_0 / m n_0)^{1/2}$ . This speed can be considered as the sound speed.

The second system is for the  $y$  and  $z$ -components of the velocity, electric field, and magnetic field perturbation. It describes transversal wave modes. Below we derive the equation describing the nonlinear transversal waves. Hence, here we present a more detailed study of this wave mode. The transversal waves are described by

$$\omega m \mathbf{v}_{\perp s} = i q_s (\mathbf{E}_{\perp} - B_0 \mathbf{e}_x \times \mathbf{v}_{\perp s}), \quad (3.1a)$$

$$k \mathbf{E}_{\perp} = -\omega \mathbf{e}_x \times \mathbf{B}_{\perp}, \quad (3.1b)$$

$$i k \mathbf{e}_x \times \mathbf{B}_{\perp} = \mu_0 q n_0 (\mathbf{v}_{\perp +} - \mathbf{v}_{\perp -}), \quad (3.1c)$$

where

$$\mathbf{v}_{\perp s} = (0, v_{ys}, v_{zs}), \quad (3.2a)$$

$$\mathbf{E}_{\perp} = (0, E_y, E_z), \quad \mathbf{B}_{\perp} = (0, B_y, B_z). \quad (3.2b)$$

Introducing the plasma bulk velocity and electrical current,

$$\mathbf{v}_{\perp} = \frac{1}{2} (\mathbf{v}_{\perp +} + \mathbf{v}_{\perp -}), \quad \mathbf{j} = q n_0 (\mathbf{v}_{\perp +} - \mathbf{v}_{\perp -}), \quad (3.3)$$

we obtain from Eqs. (3.1a) and (3.1c)

$$\omega m n_0 \mathbf{v}_{\perp} = -\frac{i}{2} B_0 \mathbf{e}_x \times \mathbf{j}, \quad (3.4)$$

$$\mathbf{E}_{\perp} = B_0 \mathbf{e}_x \times \mathbf{v}_{\perp} - \frac{i \omega m \mathbf{j}}{2 n_0 q^2}, \quad (3.5)$$

$$i k \mathbf{e}_x \times \mathbf{B}_{\perp} = \mu_0 \mathbf{j}. \quad (3.6)$$

Equation (3.5) is the Ohm's law. The second term on the right-hand side is similar to

the Hall term in the Ohm's law for the electron-ion plasma. However, the Hall term would be proportional to  $\mathbf{e}_x \times \mathbf{j}$  rather than  $\mathbf{j}$  as in Eq. (3.5). This difference is related to the fact that the masses of positively and negatively charged particles are the same in the electron-positron plasma, while the mass of positively charged particles is much larger than the mass of negatively charged particles in the electron-ion plasma. The dispersion of waves propagating along the magnetic field in electron-ion plasma is related to the account of ion inertia in the induction equation, while the electron inertia is neglected. In contrast, in an electron-positron plasma the inertia of both electrons and positrons is accounted.

Equations (3.1b) and (3.4)–(3.6) constitute the system of linear homogeneous algebraic equation. It only has non-trivial solutions when its determinant is zero. This condition gives the dispersion equation

$$m\omega^2(mk^2 + 2\mu_0q^2n_0) = q^2k^2B_0^2. \quad (3.7)$$

For small values of  $k$  this dispersion equation reduces to the approximate form

$$\omega = kV(1 - k^2\ell^2), \quad (3.8)$$

where

$$V = \frac{B_0}{\sqrt{2\mu_0mn_0}}, \quad \ell = \frac{1}{2q}\sqrt{\frac{m}{\mu_0n_0}}. \quad (3.9)$$

The wave dispersion is related to the presence of the second term in Eq. (3.5). If we neglect this term, then the dispersion relation reduces to  $\omega = kV$ .

The condition that  $k$  is small is written as  $k\ell \ll 1$ . In the non-relativistic approximation we must have the phase speed much smaller than the speed of light,  $V \ll c$ . This condition reduces to  $B^2/\mu_0 \ll mn_0c^2$ , that is the magnetic energy density is much smaller than the rest density of the plasma. We note that the term describing the wave dispersion (the second term in the brackets in Eq. (3.8)) is proportional to  $k^2$ . In the case of electron-ion plasma it is proportional to  $k$ .

#### 4. Derivation of equation for small-amplitude nonlinear waves

We consider nonlinear waves propagating along the equilibrium magnetic field. We expect that the equation describing the nonlinear wave propagation will be similar to the 3D DNLS equation describing quasi-parallel propagation of nonlinear waves in an ion-electron plasma with the only difference that the term describing the wave dispersion will be different. This difference arises from the fact that, as we have already seen, the term describing the dispersion of waves in an electron-positron plasma is proportional to  $k^2$ , while it is proportional to  $k$  in the electron-ion plasma.

To derive the nonlinear equation describing the longitudinal propagation of nonlinear waves we use the reductive perturbation method (Taniuti and Wei 1968, Kakutani et al. 1968). In accordance with this method we introduce the dimensionless amplitude of the order of  $\epsilon \ll 1$ . In the linear theory the characteristic time is  $L/V$ , where  $L$  is the characteristic length of perturbation and  $V$  is the phase speed of very long waves. We assume that the ratio of  $L/\ell$  is  $\epsilon^{-1}$ . The characteristic time of variation of the perturbation shape caused by the nonlinearity and dispersion is  $\epsilon^{-2}L/V$ . We also consider weak dependence of perturbations on  $y$  and  $z$  with the characteristic scale  $\epsilon^{-2}L$ . On the time scale much smaller than  $\epsilon^{-2}L/V$  a perturbation propagates as a wave with permanent shape with all variables only depending

on  $x - Vt$ . In accordance with the above analysis we introduce stretched variables:

$$\xi = \epsilon(x - Vt), \quad \eta = \epsilon^2 y, \quad \zeta = \epsilon^2 z, \quad \tau = \epsilon^3 t. \quad (4.1)$$

With the aid of Eqs. (2.4) we transform Eqs. (2.1) and (2.3) in the new variables to

$$\epsilon^2 \frac{\partial n_s}{\partial \tau} - V \frac{\partial n_s}{\partial \xi} + \frac{\partial(n_s v_{xs})}{\partial \xi} + \epsilon \nabla_{\perp} \cdot (n_s \mathbf{v}_{\perp s}) = 0, \quad (4.2a)$$

$$\begin{aligned} \epsilon^2 \frac{\partial v_{xs}}{\partial \tau} - V \frac{\partial v_{xs}}{\partial \xi} + v_{xs} \frac{\partial v_{xs}}{\partial \xi} + \epsilon \mathbf{v}_{\perp s} \cdot \nabla_{\perp} v_{xs} \\ + \frac{1}{mn_s} \frac{\partial p_s}{\partial \xi} = \epsilon^{-1} \frac{q_s}{m} [E_x + \mathbf{e}_x \cdot (\mathbf{v}_{\perp s} \times \mathbf{B}_{\perp})], \end{aligned} \quad (4.2b)$$

$$\begin{aligned} \epsilon^2 \frac{\partial \mathbf{v}_{\perp s}}{\partial \tau} - V \frac{\partial \mathbf{v}_{\perp s}}{\partial \xi} + v_{xs} \frac{\partial \mathbf{v}_{\perp s}}{\partial \xi} + \epsilon (\mathbf{v}_{\perp s} \cdot \nabla_{\perp}) \mathbf{v}_{\perp s} + \epsilon \frac{\nabla_{\perp} p_s}{mn_s} \\ = \epsilon^{-1} \frac{q_s}{m} [\mathbf{E}_{\perp} + \mathbf{e}_x \times (v_{xs} \mathbf{B}_{\perp} - B_x \mathbf{v}_{\perp s})], \end{aligned} \quad (4.2c)$$

$$\frac{\partial E_x}{\partial \xi} + \epsilon \nabla_{\perp} \cdot \mathbf{E}_{\perp} = \epsilon^{-1} \frac{q}{\epsilon_0} (n_+ - n_-), \quad (4.2d)$$

$$\frac{\partial B_x}{\partial \xi} + \epsilon \nabla_{\perp} \cdot \mathbf{B}_{\perp} = 0, \quad (4.2e)$$

$$\epsilon^2 \frac{\partial B_x}{\partial \tau} - V \frac{\partial B_x}{\partial \xi} = -\epsilon \mathbf{e}_x \cdot \nabla_{\perp} \times \mathbf{E}_{\perp}, \quad (4.2f)$$

$$\epsilon^2 \frac{\partial \mathbf{B}_{\perp}}{\partial \tau} - V \frac{\partial \mathbf{B}_{\perp}}{\partial \xi} = -\mathbf{e}_x \times \left( \frac{\partial \mathbf{E}_{\perp}}{\partial \xi} - \epsilon \nabla_{\perp} E_x \right), \quad (4.2g)$$

$$\epsilon \mathbf{e}_x \cdot (\nabla_{\perp} \times \mathbf{B}_{\perp}) = \epsilon^{-1} q \mu_0 (n_+ v_{x+} - n_- v_{x-}), \quad (4.2h)$$

$$\mathbf{e}_x \times \left( \frac{\partial \mathbf{B}_{\perp}}{\partial \xi} - \epsilon \nabla_{\perp} B_x \right) = \epsilon^{-1} q \mu_0 (n_+ \mathbf{v}_{\perp +} - n_- \mathbf{v}_{\perp -}), \quad (4.2i)$$

where

$$\nabla_{\perp} = \left( 0, \frac{\partial}{\partial \eta}, \frac{\partial}{\partial \zeta} \right). \quad (4.3)$$

Now we look for the solution in the form of expansions in the power series with

respect to  $\epsilon$ ,

$$\begin{aligned}
p_s &= p_0 + \epsilon p_s^{(1)} + \epsilon^2 p_s^{(2)} + \epsilon^3 p_s^{(3)} + \dots, \\
n_s &= n_0 + \epsilon n_s^{(1)} + \epsilon^2 n_s^{(2)} + \epsilon^3 n_s^{(3)} + \dots, \\
v_{xs} &= \epsilon v_{xs}^{(1)} + \epsilon^2 v_{xs}^{(2)} + \epsilon^3 v_{xs}^{(3)} + \dots, \\
\mathbf{v}_{\perp s} &= \epsilon \mathbf{v}_{\perp s}^{(1)} + \epsilon^2 \mathbf{v}_{\perp s}^{(2)} + \epsilon^3 \mathbf{v}_{\perp s}^{(3)} + \dots, \\
B_x &= B_0 + \epsilon B_x^{(1)} + \epsilon^2 B_x^{(2)} + \epsilon^3 B_x^{(3)} + \dots, \\
\mathbf{B}_{\perp} &= \epsilon \mathbf{B}_{\perp}^{(1)} + \epsilon^2 \mathbf{B}_{\perp}^{(2)} + \epsilon^3 \mathbf{B}_{\perp}^{(3)} + \dots, \\
E_x &= \epsilon E_x^{(1)} + \epsilon^2 E_x^{(2)} + \epsilon^3 E_x^{(3)} + \dots, \\
\mathbf{E}_{\perp} &= \epsilon \mathbf{E}_{\perp}^{(1)} + \epsilon^2 \mathbf{E}_{\perp}^{(2)} + \epsilon^3 \mathbf{E}_{\perp}^{(3)} + \dots.
\end{aligned} \tag{4.4}$$

We impose the boundary conditions at  $\xi \rightarrow \infty$ ,

$$\begin{aligned}
n_s \rightarrow n_0, \quad p_s \rightarrow p_0, \quad v_{xs} \rightarrow 0, \quad B_x \rightarrow B_0, \\
E_x \rightarrow 0, \quad \mathbf{v}_{\perp s} \rightarrow 0, \quad \mathbf{B}_{\perp} \rightarrow 0, \quad \mathbf{E}_{\perp} \rightarrow 0.
\end{aligned} \tag{4.5}$$

It follows from Eq. (4.5) that all quantities with the upper indices 1, 2, and so on tend to zero as  $\xi \rightarrow \infty$ .

#### 4.1. The zero-order approximation

Substituting the expansions given by Eq. (4.4) in Eqs. (4.2) and collecting terms of the order of unity in Eqs. (4.2b)–(4.2d), (4.2h), and (4.2i) we easily obtain

$$n_{+}^{(1)} = n_{-}^{(1)} = n^{(1)}, \quad v_{x+}^{(1)} = v_{x-}^{(1)} = v_x^{(1)}, \tag{4.6a}$$

$$\mathbf{v}_{\perp+}^{(1)} = \mathbf{v}_{\perp-}^{(1)} = \mathbf{v}_{\perp}^{(1)}, \tag{4.6b}$$

$$E_x^{(1)} = 0, \quad \mathbf{E}_{\perp}^{(1)} = B_0 \mathbf{e}_x \times \mathbf{v}_{\perp}^{(1)}. \tag{4.6c}$$

#### 4.2. The first-order approximation

Collecting terms of the order of  $\epsilon$  in Eqs. (2.4a) and (4.2), and using Eqs. (4.6) yields

$$V \frac{\partial n^{(1)}}{\partial \xi} = n_0 \frac{\partial v_x^{(1)}}{\partial \xi}, \quad p_s^{(1)} = \kappa p_0 \frac{n^{(1)}}{n_0}, \tag{4.7a}$$

$$\frac{1}{mn_0} \frac{\partial p_s^{(1)}}{\partial \xi} - V \frac{\partial v_x^{(1)}}{\partial \xi} = \frac{q_s}{m} [E_x^{(2)} + \mathbf{e}_x \cdot (\mathbf{v}_{\perp}^{(1)} \times \mathbf{B}_{\perp}^{(1)})], \tag{4.7b}$$

$$V \frac{\partial \mathbf{v}_{\perp}^{(1)}}{\partial \xi} = -\frac{q_s}{m} [\mathbf{E}_{\perp}^{(2)} + \mathbf{e}_x \times (v_x^{(1)} \mathbf{B}_{\perp}^{(1)} - B_x^{(1)} \mathbf{v}_{\perp}^{(1)} - B_0 \mathbf{v}_{\perp s}^{(2)})], \tag{4.7c}$$

$$n_{+}^{(2)} = n_{-}^{(2)} = n^{(2)}, \quad v_{x+}^{(2)} = v_{x-}^{(2)} = v_x^{(2)}, \tag{4.7d}$$

$$V \frac{\partial \mathbf{B}_{\perp}^{(1)}}{\partial \xi} = \mathbf{e}_x \times \frac{\partial \mathbf{E}_{\perp}^{(1)}}{\partial \xi}, \quad \frac{\partial B_x^{(1)}}{\partial \xi} = 0, \tag{4.7e}$$

$$\mathbf{e}_x \times \frac{\partial \mathbf{B}_{\perp}^{(1)}}{\partial \xi} = qn_0 \mu_0 (\mathbf{v}_{\perp+}^{(2)} - \mathbf{v}_{\perp-}^{(2)}). \tag{4.7f}$$



Equation (4.7b) represents two equations, one for  $s = +$ , and the other for  $s = -$ . Adding and subtracting these equations we obtain

$$\frac{1}{mn_0} \frac{\partial p_s^{(1)}}{\partial \xi} = V \frac{\partial v_x^{(1)}}{\partial \xi}, \quad (4.8a)$$

$$E_x^{(2)} = -\mathbf{e}_x \cdot (\mathbf{v}_\perp^{(1)} \times \mathbf{B}_\perp^{(1)}). \quad (4.8b)$$

It follows from Eqs. (4.7a), (4.8a), the second equation in Eq. (4.7e), and the boundary conditions Eq. (4.5) that

$$n^{(1)} = 0, \quad p_s^{(1)} = 0, \quad v_x^{(1)} = 0, \quad B_x^{(1)} = 0. \quad (4.9)$$

Equation (4.7c) also represents two equations, one for  $s = +$ , and the other for  $s = -$ . Adding and subtracting these equations and using Eq. (4.9) we obtain

$$\frac{\partial \mathbf{v}_\perp^{(1)}}{\partial \xi} = \frac{qB_0}{2mV} \mathbf{e}_x \times (\mathbf{v}_{\perp+}^{(2)} - \mathbf{v}_{\perp-}^{(2)}), \quad (4.10a)$$

$$2\mathbf{E}_\perp^{(2)} = B_0 \mathbf{e}_x \times (\mathbf{v}_{\perp+}^{(2)} + \mathbf{v}_{\perp-}^{(2)}). \quad (4.10b)$$

It follows from the first equation in Eq. (4.7e) and the last boundary condition in Eq. (4.5) that

$$\mathbf{E}_\perp^{(1)} = -V \mathbf{e}_x \times \mathbf{B}_\perp^{(1)}. \quad (4.11)$$

Using the second equation in Eq. (4.6c) and Eq. (4.11) yields

$$B_0 \mathbf{v}_\perp^{(1)} + V \mathbf{B}_\perp^{(1)} = 0. \quad (4.12)$$

Substituting Eq. (4.12) in Eqs. (4.8b) and (4.10a) yields

$$E_x^{(2)} = 0, \quad \mathbf{v}_{\perp+}^{(2)} - \mathbf{v}_{\perp-}^{(2)} = \frac{2mV^2}{qB_0^2} \mathbf{e}_x \times \frac{\partial \mathbf{B}_\perp^{(1)}}{\partial \xi}. \quad (4.13)$$

Equation (4.7f) and the second equation in Eq. (4.13) constitute a linear homogeneous system of equations for  $\partial \mathbf{B}_\perp^{(1)} / \partial \xi$  and  $\mathbf{v}_{\perp+}^{(2)} - \mathbf{v}_{\perp-}^{(2)}$ . It only has non-trivial solutions when its determinant is zero. This condition determines that  $V$  is given by Eq. (3.9).

### 4.3. The second-order approximation

Now we collect terms of the order of  $\epsilon^2$  in Eqs. (2.2) and (4.2). As a result, we obtain

$$p_s^{(2)} = \kappa p_0 \frac{n^{(2)}}{n_0}, \quad V \frac{\partial n^{(2)}}{\partial \xi} = n_0 \frac{\partial v_x^{(2)}}{\partial \xi} + n_0 \nabla_\perp \cdot \mathbf{v}_\perp^{(1)}, \quad (4.14a)$$

$$\frac{a_0^2}{n_0} \frac{\partial n^{(2)}}{\partial \xi} - V \frac{\partial v_x^{(2)}}{\partial \xi} = \frac{q_s}{m} [E_3^{(3)} + \mathbf{e}_x \cdot (\mathbf{v}_\perp^{(1)} \times \mathbf{B}_\perp^{(2)} + \mathbf{v}_{\perp s}^{(2)} \times \mathbf{B}_\perp^{(1)})], \quad (4.14b)$$

$$V \frac{\partial \mathbf{v}_{\perp s}^{(2)}}{\partial \xi} = -\frac{q_s}{m} [\mathbf{E}_\perp^{(3)} - \mathbf{e}_x \times (B_0 \mathbf{v}_{\perp s}^{(3)} - v_x^{(2)} \mathbf{B}_\perp^{(1)} + B_x^{(2)} \mathbf{v}_\perp^{(1)})], \quad (4.14c)$$

$$\frac{\partial E_x^{(2)}}{\partial \xi} + \nabla_\perp \cdot \mathbf{E}_\perp^{(1)} = \frac{q}{\epsilon_0} (n_+^{(3)} - n_-^{(3)}), \quad (4.14d)$$

$$\frac{\partial B_x^{(2)}}{\partial \xi} + \nabla_{\perp} \cdot \mathbf{B}_{\perp}^{(1)} = 0, \quad V \frac{\partial B_x^{(2)}}{\partial \xi} = \mathbf{e}_x \cdot \nabla_{\perp} \times \mathbf{E}_{\perp}^{(1)}, \quad (4.14e)$$

$$V \frac{\partial \mathbf{B}_{\perp}^{(2)}}{\partial \xi} = \mathbf{e}_x \times \frac{\partial \mathbf{E}_{\perp}^{(2)}}{\partial \xi}, \quad (4.14f)$$

$$\mathbf{e}_x \cdot \nabla_{\perp} \times \mathbf{B}_{\perp}^{(1)} = q\mu_0 n_0 (v_{x+}^{(3)} - v_{x-}^{(3)}) \quad (4.14g)$$

$$\mathbf{e}_x \times \frac{\partial \mathbf{B}_{\perp}^{(2)}}{\partial \xi} = q\mu_0 n_0 (\mathbf{v}_{\perp+}^{(3)} - \mathbf{v}_{\perp-}^{(3)}). \quad (4.14h)$$

Using Eq. (4.12) we transform the second equation in Eq. (4.14a) to

$$\frac{\partial v_x^{(2)}}{\partial \xi} - \frac{V}{n_0} \frac{\partial n^{(2)}}{\partial \xi} = \frac{V}{B_0} \nabla_{\perp} \cdot \mathbf{B}_{\perp}^{(1)}. \quad (4.15)$$

Equation (4.14b) represents two equations, one for  $s = +$ , and the other for  $s = -$ . Adding these equations we obtain

$$\frac{a_0^2}{n_0} \frac{\partial n^{(2)}}{\partial \xi} - V \frac{\partial v_x^{(2)}}{\partial \xi} = \frac{q}{2m} \mathbf{e}_x \cdot (\mathbf{v}_{\perp+}^{(2)} - \mathbf{v}_{\perp-}^{(2)}) \times \mathbf{B}_{\perp}^{(1)}. \quad (4.16)$$

Using Eq. (4.13) we transform this equation to

$$\frac{a_0^2}{n_0} \frac{\partial n^{(2)}}{\partial \xi} - V \frac{\partial v_x^{(2)}}{\partial \xi} = -\frac{V^2}{2B_0^2} \frac{\partial |\mathbf{B}_{\perp}^{(1)}|^2}{\partial \xi}. \quad (4.17)$$

We find from Eqs. (4.15) and (4.17)

$$\frac{\partial n^{(2)}}{\partial \xi} = \frac{n_0 V^2}{B_0 (V^2 - a_0^2)} \left( \frac{1}{2B_0} \frac{\partial |\mathbf{B}_{\perp}^{(1)}|^2}{\partial \xi} - \nabla_{\perp} \cdot \mathbf{B}_{\perp}^{(1)} \right), \quad (4.18a)$$

$$\frac{\partial v_x^{(2)}}{\partial \xi} = \frac{V}{V^2 - a_0^2} \left( \frac{V^2}{2B_0^2} \frac{\partial |\mathbf{B}_{\perp}^{(1)}|^2}{\partial \xi} - \frac{a_0^2}{B_0} \nabla_{\perp} \cdot \mathbf{B}_{\perp}^{(1)} \right). \quad (4.18b)$$

Using Eq. (4.11) and the first equation in Eq. (4.13) we obtain from Eq. (4.14d)

$$q(n_+^{(3)} - n_-^{(3)}) = \varepsilon_0 V \mathbf{e}_x \cdot \nabla_{\perp} \times \mathbf{B}_{\perp}^{(1)}. \quad (4.19)$$

Finally, Eq. (4.14c) represents two equations, one for  $s = +$ , and the other for  $s = -$ . Subtracting the second equation from the first one yields

$$\begin{aligned} 2\mathbf{E}_{\perp}^{(3)} - \mathbf{e}_x \times [B_0(\mathbf{v}_{\perp+}^{(3)} + \mathbf{v}_{\perp-}^{(3)}) - 2v_x^{(2)}\mathbf{B}_{\perp}^{(1)} \\ + 2B_x^{(2)}\mathbf{v}_{\perp}^{(1)}] = -\frac{mV}{q} \frac{\partial(\mathbf{v}_{\perp+}^{(2)} - \mathbf{v}_{\perp-}^{(2)})}{\partial \xi}. \end{aligned} \quad (4.20)$$

#### 4.4. The third-order approximation

In the third-order approximation we collect the terms of the order of  $\epsilon^3$  in Eqs. (4.2c), (4.2g), and (4.2i) to obtain

$$\begin{aligned} \frac{\partial \mathbf{v}_{\perp}^{(1)}}{\partial \tau} + v_x^{(2)} \frac{\partial \mathbf{v}_{\perp}^{(1)}}{\partial \xi} - V \frac{\partial \mathbf{v}_{\perp s}^{(3)}}{\partial \xi} + (\mathbf{v}_{\perp}^{(1)} \cdot \nabla_{\perp}) \mathbf{v}_{\perp}^{(1)} + \frac{a_0^2}{n_0} \nabla_{\perp} n^{(2)} = \frac{q_s}{m} [\mathbf{E}_{\perp}^{(4)} \\ + \mathbf{e}_x \times (v_x^{(2)} \mathbf{B}_{\perp}^{(2)} + v_{xs}^{(3)} \mathbf{B}_{\perp}^{(1)} - B_0 \mathbf{v}_{\perp s}^{(4)} - B_x^{(2)} \mathbf{v}_{\perp s}^{(2)} - B_x^{(3)} \mathbf{v}_{\perp}^{(1)})], \end{aligned} \quad (4.21a)$$

$$\frac{\partial \mathbf{E}_\perp^{(3)}}{\partial \xi} = \mathbf{e}_x \times \left( \frac{\partial \mathbf{B}_\perp^{(1)}}{\partial \tau} - V \frac{\partial \mathbf{B}_\perp^{(3)}}{\partial \xi} \right) + \nabla_\perp E_x^{(2)}, \quad (4.21b)$$

$$\mathbf{e}_x \times \left( \frac{\partial \mathbf{B}_\perp^{(3)}}{\partial \xi} - \nabla_\perp B_x^{(2)} \right) = q\mu_0 [n_0(\mathbf{v}_{\perp+}^{(4)} - \mathbf{v}_{\perp-}^{(4)}) + n^{(2)}(\mathbf{v}_{\perp+}^{(2)} - \mathbf{v}_{\perp-}^{(2)})]. \quad (4.21c)$$

Equation (4.21a) represents two equations, one for  $s = +$ , and the other for  $s = -$ . Adding these equations we obtain

$$\begin{aligned} & \frac{\partial \mathbf{v}_\perp^{(1)}}{\partial \tau} + v_x^{(2)} \frac{\partial \mathbf{v}_\perp^{(1)}}{\partial \xi} - \frac{V}{2} \frac{\partial(\mathbf{v}_{\perp+}^{(3)} + \mathbf{v}_{\perp-}^{(3)})}{\partial \xi} + \frac{a_0^2}{n_0} \nabla_\perp n^{(2)} + (\mathbf{v}_\perp^{(1)} \cdot \nabla_\perp) \mathbf{v}_\perp^{(1)} \\ &= \frac{q}{2m} \mathbf{e}_x \times [\mathbf{B}_\perp^{(1)}(v_{x+}^{(3)} - v_{x-}^{(3)}) - B_x^{(2)}(\mathbf{v}_{\perp+}^{(2)} - \mathbf{v}_{\perp-}^{(2)}) - B_0(\mathbf{v}_{\perp+}^{(4)} - \mathbf{v}_{\perp-}^{(4)})]. \end{aligned} \quad (4.22)$$

Using Eqs. (4.7f), (4.11), (4.13), and (4.14g) we transform Eqs. (4.20) and (4.21b)–(4.22) to

$$\begin{aligned} 2\mathbf{e}_x \times \frac{\partial \mathbf{E}_\perp^{(3)}}{\partial \xi} + B_0 \frac{\partial(\mathbf{v}_{\perp+}^{(3)} + \mathbf{v}_{\perp-}^{(3)})}{\partial \xi} &= \frac{mV}{qn_0\mu_0} \frac{\partial^3 \mathbf{B}_\perp^{(1)}}{\partial \xi^3} \\ &+ \frac{2}{B_0} \frac{\partial}{\partial \xi} [\mathbf{B}_\perp^{(1)}(VB_x^{(2)} + B_0v_x^{(2)})], \end{aligned} \quad (4.23a)$$

$$\frac{\partial \mathbf{E}_\perp^{(3)}}{\partial \xi} + V\mathbf{e}_x \times \frac{\partial \mathbf{B}_\perp^{(3)}}{\partial \xi} = \mathbf{e}_x \times \frac{\partial \mathbf{B}_\perp^{(1)}}{\partial \tau}, \quad (4.23b)$$

$$\frac{\partial \mathbf{B}_\perp^{(3)}}{\partial \xi} + q\mu_0 n_0 \mathbf{e}_x \times (\mathbf{v}_{\perp+}^{(4)} - \mathbf{v}_{\perp-}^{(4)}) = \frac{n^{(2)}}{n_0} \frac{\partial \mathbf{B}_\perp^{(1)}}{\partial \xi} + \nabla_\perp B_x^{(2)}, \quad (4.23c)$$

$$\begin{aligned} & \frac{qB_0}{2m} \mathbf{e}_x \times (\mathbf{v}_{\perp+}^{(4)} - \mathbf{v}_{\perp-}^{(4)}) - \frac{V}{2} \frac{\partial(\mathbf{v}_{\perp+}^{(3)} + \mathbf{v}_{\perp-}^{(3)})}{\partial \xi} = \frac{V^2}{B_0^2} \left[ B_x^{(2)} \frac{\partial \mathbf{B}_\perp^{(1)}}{\partial \xi} \right. \\ & \left. + (\mathbf{e}_x \times \mathbf{B}_\perp^{(1)}) \mathbf{e}_x \cdot \nabla_\perp \times \mathbf{B}_\perp^{(1)} \right] + \frac{V}{B_0} \frac{\partial \mathbf{B}_\perp^{(1)}}{\partial \tau} + \frac{Vv_x^{(2)}}{B_0} \frac{\partial \mathbf{B}_\perp^{(1)}}{\partial \xi} \\ & - \frac{a_0^2}{n_0} \nabla_\perp n^{(2)} - \frac{V^2}{B_0^2} (\mathbf{B}_\perp^{(1)} \cdot \nabla_\perp) \mathbf{B}_\perp^{(1)}. \end{aligned} \quad (4.23d)$$

The system of Eqs. (4.23) is the system of linear inhomogeneous algebraic equations for  $\partial \mathbf{E}_\perp^{(3)}/\partial \xi$ ,  $\partial \mathbf{B}_\perp^{(3)}/\partial \xi$ ,  $\partial(\mathbf{v}_{\perp+}^{(3)} + \mathbf{v}_{\perp-}^{(3)})/\partial \xi$ , and  $\mathbf{v}_{\perp+}^{(4)} - \mathbf{v}_{\perp-}^{(4)}$ . Using the expression for  $V$  it is straightforward to show that the determinant of this system is zero. Then the system of Eqs. (4.23) has non-trivial solution only if the compatibility condition is satisfied. This condition is

$$\begin{aligned} & \frac{\partial \mathbf{B}_\perp^{(1)}}{\partial \tau} + V\ell^2 \frac{\partial^3 \mathbf{B}_\perp^{(1)}}{\partial \xi^3} + \frac{1}{2} \mathbf{B}_\perp^{(1)} \left( \frac{\partial v_x^{(2)}}{\partial \xi} + \frac{V}{B_0} \frac{\partial B_x^{(2)}}{\partial \xi} \right) + \frac{\partial \mathbf{B}_\perp^{(1)}}{\partial \xi} \left( \frac{V}{B_0} B_x^{(2)} \right. \\ & \left. - \frac{Vn^{(2)}}{2n_0} + v_x^{(2)} \right) - \frac{V}{2} \nabla_\perp B_x^{(2)} - \frac{a_0^2 B_0}{2n_0 V} \nabla_\perp n^{(2)} \\ & + \frac{V}{2B_0} (\mathbf{e}_x \times \mathbf{B}_\perp^{(1)}) \mathbf{e}_x \cdot \nabla_\perp \times \mathbf{B}_\perp^{(1)} - \frac{V}{2B_0} (\mathbf{B}_\perp^{(1)} \cdot \nabla_\perp) \mathbf{B}_\perp^{(1)} = 0. \end{aligned} \quad (4.24)$$

The following identities can be verified by the direct calculation:

$$(\mathbf{e}_x \times \mathbf{B}_\perp^{(1)}) \mathbf{e}_x \cdot \nabla_\perp \times \mathbf{B}_\perp^{(1)} = -\mathbf{B}_\perp^{(1)} \times (\nabla_\perp \times \mathbf{B}_\perp^{(1)}), \quad (4.25a)$$

$$(\mathbf{B}_\perp^{(1)} \cdot \nabla_\perp) \mathbf{B}_\perp^{(1)} = \frac{1}{2} \nabla_\perp |\mathbf{B}_\perp^{(1)}|^2 - \mathbf{B}_\perp^{(1)} \times (\nabla_\perp \times \mathbf{B}_\perp^{(1)}). \quad (4.25b)$$

Using Eqs. (4.14e), (4.18a), and (4.18b) we obtain

$$\frac{\partial v_x^{(2)}}{\partial \xi} + \frac{V}{B_0} \frac{\partial B_x^{(2)}}{\partial \xi} = \frac{V^3}{B_0(V^2 - a_0^2)} \left( \frac{1}{2B_0} \frac{\partial |\mathbf{B}_\perp^{(1)}|^2}{\partial \xi} - \nabla_\perp \cdot \mathbf{B}_\perp^{(1)} \right), \quad (4.26a)$$

$$v_x^{(2)} - \frac{V n^{(2)}}{2n_0} + \frac{V}{B_0} B_x^{(2)} = \frac{V^3 |\mathbf{B}_\perp^{(1)}|^2}{4B_0^2(V^2 - a_0^2)} - \frac{V^3 \Phi}{2B_0(V^2 - a_0^2)}, \quad (4.26b)$$

where  $\Phi$  is defined by

$$\frac{\partial \Phi}{\partial \xi} = \nabla_\perp \cdot \mathbf{B}_\perp^{(1)}, \quad \Phi \rightarrow 0 \text{ as } \xi \rightarrow \infty. \quad (4.27)$$

Using Eqs. (4.25)–(4.27) we transform Eq. (4.24) to

$$\begin{aligned} \frac{\partial \mathbf{B}_\perp^{(1)}}{\partial \tau} + \alpha \frac{\partial}{\partial \xi} [\mathbf{B}_\perp^{(1)} (|\mathbf{B}_\perp^{(1)}|^2 - 2B_0\Phi)] \\ - \alpha B_0 \nabla_\perp (|\mathbf{B}_\perp^{(1)}|^2 - 2B_0\Phi) + V\ell^2 \frac{\partial^3 \mathbf{B}_\perp^{(1)}}{\partial \xi^3} = 0, \end{aligned} \quad (4.28)$$

where

$$\alpha = \frac{V^3}{4B_0^2(V^2 - a_0^2)}. \quad (4.29)$$

Introducing the notation

$$\mathbf{b} = \epsilon \mathbf{B}_\perp^{(1)}, \quad \tilde{\nabla}_\perp = \left( 0, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right), \quad \varphi = \epsilon^2 \Phi, \quad (4.30)$$

returning to the original independent variables, and dropping the tilde we rewrite Eqs. (4.27) and (4.28) as

$$\frac{\partial \varphi}{\partial x} = \nabla_\perp \cdot \mathbf{b}, \quad \varphi \rightarrow 0 \text{ as } x \rightarrow \infty, \quad (4.31)$$

$$\frac{\partial \mathbf{b}}{\partial t} + V \frac{\partial \mathbf{b}}{\partial x} + \alpha \frac{\partial}{\partial x} [\mathbf{b}(b^2 - 2B_0\varphi)] - \alpha B_0 \nabla_\perp (b^2 - 2B_0\varphi) + V\ell^2 \frac{\partial^3 \mathbf{b}}{\partial x^3} = 0. \quad (4.32)$$

This equation only differs from the 3D DNLS equation describing quasi-parallel propagation of MHD waves in an ion-electron plasma derived by Mjølhus and Wyller (1986) and Ruderman (1987) by the last term describing the dispersion. This difference is related to the difference in the dispersion relations for ion-electron and electron-positron plasmas as was pointed out in Sect. 3.

When  $\mathbf{b}$  is independent of  $y$  and  $z$  Eq. (4.32) reduces to the vector mKdV equation in a complete agreement with the result obtained by Verheest (1996) and Lakhina and Verheest (1997). In this equation the coefficient at the nonlinear term is  $\alpha$ . Khanna and Rajaram (1982) derived the DNLS equation in a collisionless electron-ion plasma with anisotropic pressure. They used the Chew, Goldberger and Low equations (1956) modified by including the account of Hall current in the induction equations and terms related to the finite Larmor radius in the momentum equation (Yajima 1966). While the general form of the equation remains the same, the expressions for its coefficients are quite different. In particular, while  $\alpha < 0$  when  $a_0 < V$ , in the case of plasma with anisotropic pressure the coefficient at

the nonlinear term is negative only in a relatively narrow interval of parameters. It is possible that the account of plasma pressure anisotropy can cause a similar modification of the coefficient at the nonlinear term in the vector mKdV equation.

We emphasise that the system of Eqs. (4.31) and (4.32) was derived under the assumption that the perturbations decay as  $|x| \rightarrow \infty$ . A natural question that arises is if this system of equations also describes perturbations periodic with respect to  $x$ . One-dimensional nonlinear sound waves are described by a very simple equation sometimes called the inviscid Burgers' equation (e.g. Whitham 1974, Rudenko and Soluyan 1977). This equation also describes magnetosonic waves propagating at not very small angles with respect to the equilibrium magnetic field. It is valid both for perturbations decaying at infinity as well as for periodic perturbations. The same is true for its multi-dimensional generalisation, the Khokhlov-Zabolotskaya equation (Zabolotskaya and Khokhlov 1969). The generalisations of the inviscid Burgers' and Khokhlov-Zabolotskaya equation taking into account either dissipation or dispersion, which are the Burgers', KdV, and KP equation, also describe both perturbations decaying at infinity as well as spatially periodic perturbations. The general and very important property of all these equations is that the nonlinearity that they describe is quadratic.

In contrast, magnetohydrodynamic waves propagating either along or at small angles with respect to the equilibrium magnetic field are characterised by cubic nonlinearity. In the one-dimensional case they are described in the framework of ideal MHD by the Cohen-Kulsrud equation (Cohen and Kulsrud 1974). Although this equation describing periodic waves is slightly different from that describing waves decaying at infinity, the former equation is easily reduced to the latter by a simple change of independent variables. The situation is the same with the extension of this equation to dissipative media, the so-called Cohen-Kulsrud-Burgers' equation, and to dispersive media, which is the DNLS equation. Hence, we conclude that in the one-dimensional case both the periodic waves as well as the waves decaying at infinity are described by the same equation.

The situation is drastically different in the multi-dimensional case. Ruderman (1986) studied the quasi-longitudinal propagation of MHD waves in the multi-dimensional case. In this case the mean over the period of the transverse magnetic field magnitude squared cannot be eliminated from the equation describing the evolution of the magnetic field perturbation because, in general, this mean varies in the transverse direction. As a result, the equation describing periodic perturbations differs substantially from that describing perturbations decaying at infinity. Passot and Sulem (1993) investigated a similar problem, but using the Hall MHD. As a result they obtained the analog of the 3D DNLS equation valid for periodic perturbations. If we neglect the last term in the equation derived by Passot and Sulem (1993) (see their equation (2.33)), then their equation can be reduced to the equation similar to one derived by Ruderman (1986). However, this reduction is not straightforward. The problem is that Ruderman (1986) considered the spatial variation of waves. He assumed that they are driven at  $x = 0$  and propagate in the positive  $x$ -direction. Passot and Sulem (1993) considered the temporal evolution of the waves. However, the equation derived by them with the term describing dispersion dropped looks very similar to the equation derived by Ruderman (1986). On the basis of this similarity we can make a conjecture that we can obtain an analog of Eq. (4.32) by changing the term describing dispersion in equation (2.33) in

the paper by Passot and Sulem (1993). However, to prove this conjecture a formal derivation is needed.

## 5. Obliquely propagating solitary waves

We look for solitary waves propagating at a small angle with respect to the equilibrium magnetic field. In accordance with this we look for solutions to Eq. (4.32) that depends of  $X = x + k_y y + k_z z - (C + V)t$ , where  $C$  is a constant, and  $|k_y| \ll 1$  and  $|k_z| \ll 1$ . It follows from Eq. (4.31) that

$$\varphi = \mathbf{k}_\perp \cdot \mathbf{b}, \quad \mathbf{k}_\perp = (0, k_y, k_z). \quad (5.1)$$

Using this result and the condition that  $\mathbf{b} \rightarrow 0$  as  $X \rightarrow \infty$  we obtain from Eq. (4.32)

$$V\ell^2 \mathbf{b}'' = C\mathbf{b} - \alpha(b^2 - 2B_0 \mathbf{k}_\perp \cdot \mathbf{b})(\mathbf{b} - B_0 \mathbf{k}_\perp), \quad (5.2)$$

where the prime indicates the derivative with respect to  $X$ . We can write down this equation in the Hamiltonian form,

$$g'_y = -\frac{\partial \mathcal{H}}{\partial b_y}, \quad g'_z = -\frac{\partial \mathcal{H}}{\partial b_z}, \quad h'_y = \frac{\partial \mathcal{H}}{\partial g_y}, \quad h'_z = \frac{\partial \mathcal{H}}{\partial g_z}, \quad (5.3)$$

where  $\mathbf{b} = (b_y, b_z)$ ,  $g_y = b'_y$ ,  $g_z = b'_z$ , and the Hamiltonian  $\mathcal{H}$  is given by

$$\mathcal{H} = \frac{1}{2}(g_y^2 + g_z^2) + \frac{1}{4V\ell^2} [\alpha(b^2 - 2B_0 \mathbf{k}_\perp \cdot \mathbf{b})^2 - 2Cb^2]. \quad (5.4)$$

Below we only consider solutions to the system of Eq. (5.3) describing planar solitary waves. In these solutions  $\mathbf{b} \parallel \mathbf{k}_\perp$ . In accordance with this we write

$$\mathbf{b} = \frac{\mathbf{k}_\perp}{k_\perp} h. \quad (5.5)$$

Since  $\mathcal{H}$  is independent of  $X$  it follows that the energy equal to  $\mathcal{H}$  is conserved. Since  $\mathbf{b} \rightarrow 0$  and  $\mathbf{b}' \rightarrow 0$  as  $X \rightarrow \infty$ , the energy conservation law is  $\mathcal{H} = 0$ . Then in the case of plane solitary waves we obtain

$$2V\ell^2 h'^2 = h^2 [2C - \alpha(h - 2B_0 k_\perp)^2]. \quad (5.6)$$

In the one-dimensional planar case Eq. (4.32) reduces to the modified Korteweg-de Vries equation, which is completely integrable. This implies that planar solitary waves are solitons (recall that solitons are solitary waves that are solutions of completely integrable nonlinear equations). The integral curves of Eq. (5.4) corresponding to solitons must start and end at  $h = 0$ , which is a critical point in the phase plane. In addition,  $h$  must take either maximum or minimum value, which implies that there should be the second critical point where the right-hand side of Eq. (5.4) is zero. The necessary condition of the existence of a solution to Eq. (5.6) describing a soliton is that its right-hand side must be non-negative when  $|h|$  varies from zero to its maximum, which is defined by the condition that the right-hand side is zero. When  $\alpha > 0$  this condition reduces to

$$C > 2\alpha B_0^2 k_\perp^2, \quad \alpha > 0, \quad (5.7)$$

while when  $\alpha < 0$  it reduces to

$$2\alpha B_0^2 k_\perp^2 < C < 0, \quad \alpha < 0. \quad (5.8)$$

For  $\alpha > 0$  there are two solitons. In one of them  $h > 0$  and we call it the bright

soliton, while in the other  $h < 0$  and we call it the dark soliton. These solitons are described by

$$h = \frac{\pm 2(C - 2\alpha B_0^2 k_\perp^2)}{\sqrt{2\alpha C} \cosh(X/L + \Theta) \mp 2\alpha B_0 k_\perp}, \quad (5.9)$$

where the upper and lower signs correspond to the bright and dark soliton, respectively. When  $\alpha < 0$  there is only one soliton, so we do not use the notion “bright” or “dark” in this case. It is described by Eq. (5.9) with the upper sign. The characteristic soliton thickness is given by

$$L = \ell \sqrt{\frac{V}{C - 2\alpha B_0^2 k_\perp^2}}. \quad (5.10)$$

Equation (5.10) is valid both for  $\alpha > 0$  as well as for  $\alpha < 0$ . The phase shift  $\Theta$  is defined by

$$\tanh \Theta = \begin{cases} \sqrt{1 - \frac{2\alpha B_0^2 k_\perp^2}{C}}, & \alpha > 0, \\ \sqrt{1 - \frac{C}{2\alpha B_0^2 k_\perp^2}}, & \alpha < 0. \end{cases} \quad (5.11)$$

The soliton amplitude is given by

$$A = \max |h| = \left| \sqrt{\frac{2C}{\alpha}} \pm 2B_0 k_\perp \right|, \quad (5.12)$$

where for  $\alpha > 0$  the upper and lower signs correspond to the bright and dark soliton, respectively. For  $\alpha < 0$  the bright soliton amplitude is given by Eq. (5.12) with the lower sign.

In this section we only obtained the solutions describing planar solitons. Although, at present, there is no rigorous study of the existence of non-planar solitary waves, we expect that there should be a whole three-parametric family of non-planar solitary waves. The two parameters are the same as in the planar solitons, that are  $C$  and  $k_\perp$ . The third parameter is the angle between the plane defined by  $\mathbf{k}_\perp$  and  $\mathbf{e}_x$  and the integral curve near the critical point corresponding to  $|X| \rightarrow \infty$ .

Verheest (1996) studied solitary waves of the vector mKdV equation with  $\alpha > 0$ . He showed that only a planar soliton exists. Below we will call this soliton the standard mKdV soliton. However, Verheest considered solitary waves propagating exactly along the equilibrium magnetic field. His proof is not valid in the case of oblique propagation. It is straightforward to verify that both bright and dark solitons tend to the standard mKdV soliton as  $k_\perp \rightarrow 0$ .

Since the vector mKdV equation has some similarities with the DNLS equation it is expedient to compare solitons of the two equation. The DNLS equation possesses not only solitons that only depend on the linear combination of the spatial variable and time, but also solitons in the form of an envelope with the magnetic field vector rotating inside this envelope with constant angular velocity. Below we only consider the first type of solitons. There are no solitons of this type propagating exactly along the equilibrium magnetic field. All of them propagate at some angle with respect to this field. And, in addition, all these solutions are non-planar. The family of solitons is three-parametric (Ruderman 1987). The two parameters are  $k_\perp$  determining the propagation direction, the propagation velocity  $C$ . The third

parameter,  $\vartheta$ , determines the type of soliton. When  $0 < \vartheta < \pi/2$  the component of the magnetic field orthogonal to the equilibrium magnetic field makes one full turn about the equilibrium magnetic field direction in the positive direction when  $\alpha > 0$  and in the negative direction when  $\alpha < 0$ . In accordance with the nomenclature introduced by Ruderman this soliton is called the compression Alfvén soliton. When  $\pi/2 < \vartheta < 2\pi/3$  the component of the magnetic field orthogonal to the equilibrium magnetic field makes one full turn about the equilibrium magnetic field direction in the negative direction when  $\alpha > 0$  and in the positive direction when  $\alpha < 0$ . In accordance with the nomenclature introduced by Ruderman this soliton is called the rarefaction Alfvén soliton. Finally, when  $2\pi/3 < \vartheta < \pi$  the component of the magnetic field orthogonal to the equilibrium magnetic field rotates by some angle and then returns back to the initial position. This soliton is called magnetosonic, fast when  $\alpha > 0$  and slow when  $\alpha < 0$ .

We see that the properties of solitons of the DNLS equation are very much different from those of solitons of the vector mKdV equation. Ruderman (1987) showed that compression Alfvén solitons are stable with respect to transverse perturbations, while rarefaction Alfvén solitons and magnetosonic solitons are unstable.

## 6. Soliton stability

In this section we study the stability of solitons described in the previous section with respect to transverse perturbations. This study is similar to those carried out for the stability of the KdV solitons by Kadomtsev and Petviashvili (1970) and for the stability of the DNLS solitons by Ruderman (1987). We write

$$\mathbf{b} = \mathbf{b}_s + \tilde{\mathbf{b}}, \quad \varphi = \mathbf{k}_\perp \cdot \mathbf{b}_s + \tilde{\varphi}, \quad (6.1)$$

where  $\mathbf{b}_s$  corresponds to the soliton defined by Eqs. (5.5) and (5.9). It describes either the bright or dark soliton. We substitute Eq. (6.1) in Eq. (4.32) and then linearise the obtained equation with respect to  $\tilde{\mathbf{b}}$  and  $\tilde{\varphi}$ . This gives

$$\begin{aligned} \frac{\partial \tilde{\mathbf{b}}}{\partial t} + V \frac{\partial \tilde{\mathbf{b}}}{\partial x} + \alpha \frac{\partial}{\partial x} \left[ \tilde{\mathbf{b}} (h^2 - 2B_0 k_\perp h) + \frac{2h}{k_\perp} \mathbf{k}_\perp \left( \frac{h}{k_\perp} \mathbf{k}_\perp \cdot \tilde{\mathbf{b}} - B_0 \tilde{\varphi} \right) \right] \\ - 2\alpha B_0 \nabla_\perp \left( \frac{h}{k_\perp} \mathbf{k}_\perp \cdot \tilde{\mathbf{b}} - B_0 \tilde{\varphi} \right) + V \ell^2 \frac{\partial^3 \tilde{\mathbf{b}}}{\partial x^3} = 0. \end{aligned} \quad (6.2)$$

Equation (4.31) is transformed to

$$\frac{\partial \tilde{\varphi}}{\partial x} = \nabla_\perp \cdot \tilde{\mathbf{b}}, \quad \tilde{\varphi} \rightarrow 0 \text{ as } x \rightarrow \infty. \quad (6.3)$$

Equation (4.32) was derived under the assumption that the ratio of characteristic spatial scale in the  $y$  and  $z$ -direction to that in the  $x$ -direction is  $\epsilon^{-1}$ . Now we assume that this ratio is even larger and is equal to  $(\epsilon\delta)^{-1}$ , where  $\delta \ll 1$ . We also study the stability with respect to normal modes and take  $\tilde{\mathbf{b}} \propto \exp(\lambda t + i\delta K_y y + i\delta K_z z)$ . Finally, we use the variable  $X$  instead of  $x$ . As a result, we transform Eqs. (6.2) and (6.3) to

$$\begin{aligned} \frac{d}{dX} \mathcal{L}(\tilde{\mathbf{b}}) = -\lambda \tilde{\mathbf{b}} + 2\alpha B_0 \left\{ \mathbf{k}_\perp \left[ \frac{d}{dX} \left( \frac{h}{k_\perp} (\tilde{\varphi} - \mathbf{k}_\perp \cdot \tilde{\mathbf{b}}) \right) \right. \right. \\ \left. \left. - i\delta B_0 (\mathbf{K} \cdot \tilde{\mathbf{b}}) \right] + i\delta \mathbf{K} \left( \frac{h}{k_\perp} (\mathbf{k}_\perp \cdot \tilde{\mathbf{b}}) - B_0 \tilde{\varphi} \right) \right\}, \end{aligned} \quad (6.4)$$



$$\frac{d\tilde{\varphi}}{dX} = i\delta\mathbf{K} \cdot \tilde{\mathbf{b}} + \mathbf{k}_\perp \cdot \frac{d\tilde{\mathbf{b}}}{dX}, \quad (6.5)$$

where  $\mathbf{K} = (0, K_y, K_z)$ , and

$$\mathcal{L}(\tilde{\mathbf{b}}) = \tilde{\mathbf{b}}[\alpha h(h - 2k_\perp B_0) - C] + \frac{2\alpha}{k_\perp^2} \mathbf{k}_\perp (\mathbf{k}_\perp \cdot \tilde{\mathbf{b}})(h - k_\perp B_0)^2 + V\ell^2 \frac{d^2\tilde{\mathbf{b}}}{dX^2}. \quad (6.6)$$

We look for the solution to Eqs. (6.4) and (6.5) in the form of expansions

$$\tilde{\mathbf{b}} = \mathbf{b}_0 + \delta\mathbf{b}_1 + \delta^2\mathbf{b}_2 + \dots, \quad \tilde{\varphi} = \varphi_0 + \delta\varphi_1 + \delta^2\varphi_2 + \dots, \quad \lambda = \delta\lambda_1 + \delta^2\lambda_2 + \dots \quad (6.7)$$

### 6.1. The zero-order approximation

Substituting Eq. (6.7) in Eqs. (6.4) and (6.5) and using the condition that  $\tilde{\mathbf{b}} \rightarrow 0$  as  $X \rightarrow -\infty$  we obtain in the zero-order approximation

$$\mathcal{L}(\mathbf{b}_0) = 0, \quad \varphi_0 = \mathbf{k}_\perp \cdot \mathbf{b}_0. \quad (6.8)$$

Differentiating Eq. (5.2) and using the second equation in Eq. (6.8) we obtain that

$$\mathbf{b}_0 = \ell \frac{\mathbf{k}_\perp}{k_\perp} \frac{dh}{dX}, \quad \varphi_0 = \ell k_\perp \frac{dh}{dX}. \quad (6.9)$$

The multiplier  $\ell$  is introduced in the expression for  $\mathbf{b}_0$  to have the same dimension of the left and right sides. We obtain the general solution to the first equation in Eq. (6.8) multiplying this expression by an arbitrary constant. Since we solve a linear problem we can take this constant equal to unity without loss of generality.

### 6.2. The first-order approximation

Now we collect the terms of the order of  $\delta$  in Eq. (6.4) and (6.5) to obtain

$$\begin{aligned} \frac{d}{dX} \mathcal{L}(\mathbf{b}_1) = & -\lambda_1 \mathbf{b}_0 + 2\alpha B_0 \left\{ \mathbf{k}_\perp \left[ \frac{d}{dX} \left( \frac{h}{k_\perp} (\varphi_1 - \mathbf{k}_\perp \cdot \mathbf{b}_1) \right) \right. \right. \\ & \left. \left. - iB_0 (\mathbf{K} \cdot \mathbf{b}_0) \right] + i\mathbf{K} \left( \frac{h}{k_\perp} (\mathbf{k}_\perp \cdot \mathbf{b}_0) - B_0 \varphi_0 \right) \right\}, \end{aligned} \quad (6.10a)$$

$$\frac{d\varphi_1}{dX} = i\mathbf{K} \cdot \mathbf{b}_0 + \mathbf{k}_\perp \cdot \frac{d\mathbf{b}_1}{dX}. \quad (6.10b)$$

Using Eq. (6.9) we transform Eq. (6.10b) to

$$\varphi_1 - \mathbf{k}_\perp \cdot \mathbf{b}_1 = \frac{i\ell h}{k_\perp} (\mathbf{k}_\perp \cdot \mathbf{K}). \quad (6.11)$$

With the aid of Eqs. (6.9) and (6.11) we transform Eq. (6.10a) to

$$\mathcal{L}(\mathbf{b}_1) = -\ell\lambda_1 h \frac{\mathbf{k}_\perp}{k_\perp} + i\ell\alpha B_0 h \left[ \frac{2\mathbf{k}_\perp}{k_\perp^2} (h - k_\perp B_0) (\mathbf{k}_\perp \cdot \mathbf{K}) + \mathbf{K} (h - 2k_\perp B_0) \right]. \quad (6.12)$$

Differentiating Eq. (5.2) with respect to  $C$  and  $\mathbf{k}_\perp$  we obtain

$$\mathcal{L} \left( \frac{\mathbf{k}_\perp}{k_\perp} \frac{\partial h}{\partial C} \right) = \frac{\mathbf{k}_\perp}{k_\perp} h, \quad (6.13a)$$

$$\mathcal{L} \left( \left( \mathbf{K} \cdot \frac{\partial}{\partial \mathbf{k}_\perp} \right) \frac{\mathbf{k}_\perp}{k_\perp} h \right) = \alpha B_0 h \left[ \frac{2\mathbf{k}_\perp}{k_\perp^2} (h - B_0 k_\perp) (\mathbf{k}_\perp \cdot \mathbf{K}) + \mathbf{K} (h - 2B_0 k_\perp) \right]. \quad (6.13b)$$

It follows from Eqs. (6.13a) and (6.13b) that the solution to Eq. (6.12) is given by

$$\mathbf{b}_1 = -\ell\lambda_1 \frac{\mathbf{k}_\perp}{k_\perp} \frac{\partial h}{\partial C} + i\ell \left( \mathbf{K} \cdot \frac{\partial}{\partial \mathbf{k}_\perp} \right) \frac{\mathbf{k}_\perp}{k_\perp} h. \quad (6.14)$$

### 6.3. The second-order approximation

Collecting the terms of the order of  $\delta^2$  in Eqs. (6.4) and (6.5), and using Eqs. (6.9) and (6.11), yields

$$\begin{aligned} \frac{d}{dX} \mathcal{L}(\mathbf{b}_2) = & -\lambda_1 \mathbf{b}_1 - \lambda_2 \ell \frac{\mathbf{k}_\perp}{k_\perp} \frac{dh}{dX} + 2\alpha B_0 \left\{ \mathbf{k}_\perp \left[ \frac{d}{dX} \left( \frac{h}{k_\perp} (\varphi_2 - \mathbf{k}_\perp \cdot \mathbf{b}_2) \right) \right. \right. \\ & \left. \left. - iB_0 (\mathbf{K} \cdot \mathbf{b}_1) \right] + \frac{\mathbf{K}}{k_\perp} [i(h - B_0 k_\perp)(\mathbf{k}_\perp \cdot \mathbf{b}_1) + \ell B_0 h(\mathbf{k}_\perp \cdot \mathbf{K})] \right\}, \end{aligned} \quad (6.15a)$$

$$\frac{d\varphi_2}{dX} = i\mathbf{K} \cdot \mathbf{b}_1 + \mathbf{k}_\perp \cdot \frac{d\mathbf{b}_2}{dX}. \quad (6.15b)$$

The homogeneous counterpart of Eq. (6.15a) has a non-trivial solution

$$\mathbf{b}_2 = \ell \frac{\mathbf{k}_\perp}{k_\perp} \frac{dh}{dX}.$$

This implies that Eq. (6.15a) has solutions only if its right-hand side satisfies the compatibility condition. To obtain this condition we take the scalar product of Eq. (6.15a) with  $(\mathbf{k}_\perp/k_\perp)h$  and integrate with respect to  $X$ . Using the integration by parts we obtain that the left-hand side is zero, which implies that the right-hand side must be equal to zero. Then, using Eq. (6.15b) and the integration by parts to transform the term containing  $\varphi_2$  we obtain the compatibility condition

$$\begin{aligned} \frac{\lambda_1}{k_\perp} \int_{-\infty}^{\infty} h(\mathbf{k}_\perp \cdot \mathbf{b}_1) dX = & \alpha B_0 \int_{-\infty}^{\infty} h \left[ i(h - 2k_\perp B_0)(\mathbf{K} \cdot \mathbf{b}_1) \right. \\ & \left. + \frac{2i}{k_\perp^2} (\mathbf{k}_\perp \cdot \mathbf{K})(h - k_\perp B_0)(\mathbf{k}_\perp \cdot \mathbf{b}_1) + \frac{2\ell}{k_\perp^2} B_0 (\mathbf{k}_\perp \cdot \mathbf{K})^2 h \right] dX. \end{aligned} \quad (6.16)$$

Now we introduce the notation

$$I_1 = \int_{-\infty}^{\infty} h^2 dX, \quad I_2 = \int_{-\infty}^{\infty} h^3 dX. \quad (6.17)$$

Then, using Eq. (6.14) and the identity

$$\begin{aligned} \mathbf{K} \cdot \frac{\partial}{\partial \mathbf{k}_\perp} \left[ \frac{\mathbf{k}_\perp \cdot \mathbf{K}}{k_\perp} (I_2 - 2k_\perp B_0 I_1) \right] = & (\mathbf{k}_\perp \cdot \mathbf{K}) \left( \frac{1}{k_\perp} \mathbf{K} \cdot \frac{\partial I_2}{\partial \mathbf{k}_\perp} \right. \\ & \left. - 2B_0 \mathbf{K} \cdot \frac{\partial I_1}{\partial \mathbf{k}_\perp} \right) - 2B_0 K^2 I_1 + \left( K^2 - \frac{(\mathbf{k}_\perp \cdot \mathbf{K})^2}{k_\perp^2} \right) \frac{I_2}{k_\perp}, \end{aligned} \quad (6.18)$$

we transform Eq. (6.16) to

$$\begin{aligned} \lambda_1^2 \frac{\partial I_1}{\partial C} - i\lambda_1 \left[ \mathbf{K} \cdot \frac{\partial I_1}{\partial \mathbf{k}_\perp} + \frac{2\alpha}{k_\perp} B_0 (\mathbf{k}_\perp \cdot \mathbf{K}) \left( \frac{\partial I_2}{\partial C} - 2k_\perp B_0 \frac{\partial I_1}{\partial C} \right) \right] \\ - 2\alpha B_0 \mathbf{K} \cdot \frac{\partial}{\partial \mathbf{k}_\perp} \left[ \frac{\mathbf{k}_\perp \cdot \mathbf{K}}{k_\perp} (I_2 - 2k_\perp B_0 I_1) \right] = 0. \end{aligned} \quad (6.19)$$

When the discriminant of quadratic equation Eq. (6.19) is positive it has two complex roots, and one of these roots has the positive real part. This implies that in

this case the soliton is unstable. On the other hand, when the discriminant is negative, Eq. (6.17) has two purely imaginary roots and the soliton is neutrally stable. Hence, the instability condition is written as

$$8\alpha B_0 \frac{\partial I_1}{\partial C} \mathbf{K} \cdot \frac{\partial}{\partial \mathbf{k}_\perp} \left[ \frac{\mathbf{k}_\perp \cdot \mathbf{K}}{k_\perp} (I_2 - 2k_\perp B_0 I_1) \right] > \left[ \mathbf{K} \cdot \frac{\partial I_1}{\partial \mathbf{k}_\perp} + \frac{2\alpha}{k_\perp} B_0 (\mathbf{k}_\perp \cdot \mathbf{K}) \left( \frac{\partial I_2}{\partial C} - 2k_\perp B_0 \frac{\partial I_1}{\partial C} \right) \right]^2. \quad (6.20)$$

Now we consider two cases, one with  $\alpha > 0$ , and the other with  $\alpha < 0$ . First we assume that  $\alpha > 0$ . It is shown in Appendix A that  $I_1$  and  $I_2$  are given by

$$I_1 = 4\ell B_0 k_\perp \sqrt{\frac{2V}{\alpha}} F_1(\sigma), \quad I_2 = 4\ell B_0^2 k_\perp^2 \sqrt{\frac{2V}{\alpha}} F_2(\sigma), \quad (6.21)$$

where  $\sigma = C(2\alpha B_0^2 k_\perp^2)^{-1}$  and

$$F_1(\sigma) = \sqrt{\sigma - 1} \pm \frac{\pi}{2} + \arctan \frac{1}{\sqrt{\sigma - 1}}, \quad (6.22a)$$

$$F_2(\sigma) = (\sigma + 2) \left( \pm \frac{\pi}{2} + \arctan \frac{1}{\sqrt{\sigma - 1}} \right) + 3\sqrt{\sigma - 1}. \quad (6.22b)$$

Using Eqs. (6.21) and (6.22) we transform Eq. (6.20) to

$$D_\pm \equiv [k_\perp^2 K^2 - (\mathbf{k}_\perp \cdot \mathbf{K})^2] Q_\pm(\sigma) - (\mathbf{k}_\perp \cdot \mathbf{K})^2 S_\pm(\sigma) > 0, \quad (6.23)$$

where

$$Q_\pm(\sigma) = \sigma \left( \pm \frac{\pi}{2} + \arctan \frac{1}{\sqrt{\sigma - 1}} \right) + \sqrt{\sigma - 1}, \quad (6.24a)$$

$$S_\pm(\sigma) = \frac{2\sigma}{\sqrt{\sigma - 1}} \left( \pm \frac{\pi}{2} + \arctan \frac{1}{\sqrt{\sigma - 1}} \right)^2 + 2\sqrt{\sigma - 1}. \quad (6.24b)$$

Obviously  $Q_+(\sigma) > 0$  meaning that  $D_+ > 0$  when  $k_\perp^{-1} K^{-1} |\mathbf{k}_\perp \cdot \mathbf{K}|$  is sufficiently small. This implies that the bright soliton is unstable.

Now we note that  $Q_-(0) = 0$  and

$$\frac{dQ_-}{d\sigma} = \arctan \frac{1}{\sqrt{\sigma - 1}} - \frac{\pi}{2} < 0, \quad (6.25)$$

which implies that  $Q_-(\sigma) < 0$ . Since  $S_-(\sigma) > 0$  it follows that  $D_- < 0$  implying that the dark soliton is stable.

Next we proceed to the case where  $\alpha < 0$ . It is shown in Appendix A that now  $I_1$  and  $I_2$  are given by

$$I_1 = 4\ell B_0 k_\perp \sqrt{\frac{2V}{|\alpha|}} G_1(\sigma), \quad I_2 = 4\ell B_0^2 k_\perp^2 \sqrt{\frac{2V}{|\alpha|}} G_2(\sigma), \quad (6.26)$$

where

$$G_1(\sigma) = \frac{1}{2} \ln \frac{1 + \sqrt{1 - \sigma}}{1 - \sqrt{1 - \sigma}} - \sqrt{1 - \sigma}, \quad (6.27a)$$

$$G_2(\sigma) = \left( 1 + \frac{\sigma}{2} \right) \ln \frac{1 + \sqrt{1 - \sigma}}{1 - \sqrt{1 - \sigma}} - 3\sqrt{1 - \sigma}. \quad (6.27b)$$

Using Eqs. (6.26) and (6.27) we transform Eq. (6.21) to

$$D \equiv [k_{\perp}^2 K^2 - (\mathbf{k}_{\perp} \cdot \mathbf{K})^2] Q(\sigma) - (\mathbf{k}_{\perp} \cdot \mathbf{K})^2 S(\sigma) > 0, \quad (6.28)$$

where

$$Q(\sigma) = \sigma \ln \frac{1 - \sqrt{1 - \sigma}}{1 + \sqrt{1 + \sigma}} + 2\sqrt{1 - \sigma}, \quad (6.29a)$$

$$S(\sigma) = \left( \ln \frac{1 - \sqrt{1 - \sigma}}{1 + \sqrt{1 + \sigma}} \right)^2 + 4\sqrt{1 - \sigma}. \quad (6.29b)$$

Since  $Q(0) = 0$  and

$$\frac{dQ}{d\sigma} = \ln \frac{1 - \sqrt{1 - \sigma}}{1 + \sqrt{1 + \sigma}} < 0, \quad (6.30)$$

it follows that  $Q(\sigma) > 0$ . This implies that  $D > 0$  when  $k_{\perp}^{-1} K^{-1} |\mathbf{k}_{\perp} \cdot \mathbf{K}|$  is sufficiently small. Consequently, the soliton existing when  $\alpha < 0$  is unstable.

As we have already point out in Sect. 5, both the bright and dark soliton become the standard mKdV soliton propagating exactly along the equilibrium magnetic field when  $\kappa_{\perp} \rightarrow 0$ . This soliton only exists when  $\alpha > 0$ . It is obvious that the previous stability analysis is not valid for  $\kappa_{\perp} = 0$ . Hence, the stability of solitons propagating along the equilibrium magnetic field must be studied separately. However, while the expression describing the standard mKdV soliton is simpler than those describing the obliquely propagating soliton, the study of stability of this soliton with respect to transverse perturbations turns out to be much more involved. The complexity of this study is related to the fact that, while obliquely propagating solitons are two-parametric, the standard soliton is only one-parametric. As a result, while we can obtain the relation similar to Eq. (6.13a) for the standard mKdV soliton, we cannot obtain an analog of Eq. (6.13b). Hence, we cannot get a relatively simple expression for  $\mathbf{b}_1$  similar to one given by Eq. (6.14). To calculate  $\mathbf{b}_1$  we need to solve a second order ordinary differential equation with variable coefficients. At present it is even not clear that the analytical expression for  $\mathbf{b}_1$  can be obtained. Quite possible that this problem can be only solved numerically.

## 7. Summary and conclusions.

In this article we studied the propagation of nonlinear waves along the equilibrium magnetic field in a non-relativistic electron-positron plasma. We assumed that the waves can weakly depend on the spatial coordinates orthogonal to the equilibrium magnetic field. Using the reductive perturbation method we derived the three-dimensional generalisation of the vector modified Korteweg-de Vries (mKdV) equation. We call this equation the 3D vector mKdV equation.

We obtained solutions to the 3D vector mKdV equation in the form of one-dimensional planar solitons propagating at a small angle with respect to the equilibrium magnetic field. The propagation direction is defined by the vector  $\mathbf{e}_x + \mathbf{k}_{\perp}$ , where  $\mathbf{e}_x$  is the unit vector in the direction of the equilibrium magnetic field,  $\mathbf{k}_{\perp} \perp \mathbf{e}_x$ , and  $k_{\perp} \ll 1$ . In planar solitons the magnetic field perturbation is everywhere in the direction of  $\mathbf{k}_{\perp}$ . We found that in the case where the Alfvén speed  $V$  is larger than the sound speed  $a_0$  there are two kinds of solitons, bright and dark. In the bright solitons the magnetic field perturbation is parallel to  $\mathbf{k}_{\perp}$ , and in the

dark solitons it is antiparallel to  $\mathbf{k}_\perp$ . In the case where  $V < a_0$  there is only one kind of solitons with the magnetic field parallel to  $\mathbf{k}_\perp$ .

We used the 3D vector mKdV equation to study the soliton stability with respect to transverse perturbations similar to that carried out by Kadomtsev and Petviashvili (1970) for solitons described by the KdV equation. We found that only the dark solitons are stable, while both the bright solitons in the case where  $V > a_0$  as well as solitons in the case where  $V < a_0$  are unstable.

## Appendix A. Calculation of $I_1$ and $I_2$

In this appendix we calculate  $I_1$  and  $I_2$ . We start from the case where  $\alpha > 0$ . Using Eq. (5.9) we obtain

$$I_1 = \int_{-\infty}^{\infty} \frac{4(C - 2\alpha B_0^2 k_\perp^2)^2 dX}{[\sqrt{2\alpha C} \cosh(X/L + \Theta) \pm 2\alpha B_0 k_\perp]^2}. \quad (\text{A } 1)$$

Using the variable substitution

$$u = \exp(X/L + \Theta) \pm B_0 k_\perp \sqrt{2\alpha/C} \quad (\text{A } 2)$$

we transform Eq. (A 1) to

$$I_1 = \frac{8L(C - 2\alpha B_0^2 k_\perp^2)^2}{\alpha C} \left( \int_{\pm B_0 k_\perp (2\alpha/C)^{1/2}}^{\infty} \frac{u du}{(u^2 + 1 - 2\alpha B_0^2 k_\perp^2 / C)^2} \mp B_0 k_\perp \sqrt{\frac{2\alpha}{C}} \int_{\pm B_0 k_\perp (2\alpha/C)^{1/2}}^{\infty} \frac{du}{(u^2 + 1 - 2\alpha B_0^2 k_\perp^2 / C)^2} \right). \quad (\text{A } 3)$$

We easily obtain

$$\int_{\pm B_0 k_\perp (2\alpha/C)^{1/2}}^{\infty} \frac{u du}{(u^2 + 1 - 2\alpha B_0^2 k_\perp^2 / C)^2} = \frac{1}{2}. \quad (\text{A } 4)$$

Next we calculate the second integral in Eq. (A 3). Using the variable substitution

$$w = u \sqrt{\frac{C}{C - 2\alpha B_0^2 k_\perp^2}} \quad (\text{A } 5)$$

we transform it to

$$\begin{aligned} & \int_{\pm B_0 k_\perp (2\alpha/C)^{1/2}}^{\infty} \frac{du}{(u^2 + 1 - 2\alpha B_0^2 k_\perp^2 / C)^2} \\ &= \left( \frac{C}{C - 2\alpha B_0^2 k_\perp^2} \right)^{3/2} \int_{\pm B_0 k_\perp \sqrt{\frac{C}{C - 2\alpha B_0^2 k_\perp^2}}}^{\infty} \frac{dw}{(1 + w^2)^2}. \end{aligned} \quad (\text{A } 6)$$

Then the integration by parts yields

$$\begin{aligned} \int \frac{dw}{(1 + w^2)^2} &= \int \frac{dw}{1 + w^2} - \int \frac{w^2 dw}{(1 + w^2)^2} = \frac{w}{2(1 + w^2)} \\ &- \frac{1}{2} \int \frac{dw}{1 + w^2} = \frac{w}{2(1 + w^2)} + \frac{1}{2} \arctan w, \end{aligned} \quad (\text{A } 7)$$

where we dropped the arbitrary constant. Using Eqs. (A 4), (A 6), and (A 7) we obtain from Eq. (A 3)

$$I_1 = \frac{4L(C - 2\alpha B_0^2 k_\perp^2)^2}{\alpha} \left[ 1 \mp \sqrt{\frac{2\alpha B_0^2 k_\perp^2}{C - 2\alpha B_0^2 k_\perp^2}} \left( \frac{\pi}{2} \mp \arctan \sqrt{\frac{2\alpha B_0^2 k_\perp^2}{C - 2\alpha B_0^2 k_\perp^2}} \right) \right]. \quad (\text{A } 8)$$

Substituting  $C = 2\alpha\sigma B_0^2 k_\perp^2$  in this expression we eventually obtain the first expression in Eq. (6.21).

Now we proceed to the calculation of  $I_2$ . Using Eq. (5.9) we obtain

$$I_2 = \pm \int_{-\infty}^{\infty} \frac{8(C - 2\alpha B_0^2 k_\perp^2)^3 dX}{[\sqrt{2\alpha C} \cosh(X/L + \Theta) \mp 2\alpha b_0 k_\perp]^3}. \quad (\text{A } 9)$$

Using the variable substitution defined by Eq. (A 2) we transform Eq. (A 9) to

$$I_2 = \pm \frac{64L(C - 2\alpha B_0^2 k_\perp^2)^3}{(2\alpha C)^{3/2}} \int_{\pm B_0 k_\perp (2\alpha/C)^{1/2}}^{\infty} \frac{u^2 \mp 2uB_0 k_\perp \sqrt{2\alpha/C} + 2\alpha B_0^2 k_\perp^2 / C}{(u^2 + 1 - 2\alpha B_0^2 k_\perp^2 / C)^3} du. \quad (\text{A } 10)$$

We easily obtain

$$\int_{\pm B_0 k_\perp (2\alpha/C)^{1/2}}^{\infty} \frac{u du}{(u^2 + 1 - 2\alpha B_0^2 k_\perp^2 / C)^3} = \frac{1}{4}. \quad (\text{A } 11)$$

Using the variable substitution defined by Eq. (A 5) yields

$$\begin{aligned} & \int_{\pm B_0 k_\perp (2\alpha/C)^{1/2}}^{\infty} \frac{u^2 + 2\alpha B_0^2 k_\perp^2 / C}{(u^2 + 1 - 2\alpha B_0^2 k_\perp^2 / C)^3} du = \left( \frac{C}{C - 2\alpha B_0^2 k_\perp^2} \right)^{3/2} \\ & \times \int_{\pm B_0 k_\perp \sqrt{\frac{C}{C - 2\alpha B_0^2 k_\perp^2}}}^{\infty} \frac{w^2 + 2\alpha B_0^2 k_\perp^2 (C - 2\alpha B_0^2 k_\perp^2)^{-1}}{(1 + w^2)^3} dw. \end{aligned} \quad (\text{A } 12)$$

Using the integration by parts and Eq. (A 7) yields

$$\begin{aligned} & \int \frac{w^2 + 2\alpha B_0^2 k_\perp^2 (C - 2\alpha B_0^2 k_\perp^2)^{-1}}{(1 + w^2)^3} dw = \frac{2\alpha B_0^2 k_\perp^2}{C - 2\alpha B_0^2 k_\perp^2} \int \frac{dw}{(1 + w^2)^2} \\ & - \frac{C - 4\alpha B_0^2 k_\perp^2}{C - 2\alpha B_0^2 k_\perp^2} \int \frac{w^2 dw}{(1 + w^2)^3} = -\frac{C - 4\alpha B_0^2 k_\perp^2}{4(C - 2\alpha B_0^2 k_\perp^2)} \frac{w}{(1 + w^2)^2} \\ & + \frac{C + 4\alpha B_0^2 k_\perp^2}{4(C - 2\alpha B_0^2 k_\perp^2)} \int \frac{dw}{(1 + w^2)^2} = \frac{C + 4\alpha B_0^2 k_\perp^2}{8(C - 2\alpha B_0^2 k_\perp^2)} \arctan w \\ & + \frac{w[(C + 4\alpha B_0^2 k_\perp^2)w^2 - C + 12\alpha B_0^2 k_\perp^2]}{8(C - 2\alpha B_0^2 k_\perp^2)(1 + w^2)^2}. \end{aligned} \quad (\text{A } 13)$$

With the aid of this result and Eqs. (A 11)–(A 13) we obtain from Eq. (A 10)

$$\begin{aligned} I_2 = \frac{2\ell}{\alpha} \sqrt{\frac{2V}{\alpha}} \left[ (C + 4\alpha B_0^2 k_\perp^2) \left( \pm \frac{\pi}{2} - \arctan \sqrt{\frac{2\alpha B_0^2 k_\perp^2}{C - 2\alpha B_0^2 k_\perp^2}} \right) \right. \\ \left. - 3\sqrt{2\alpha B_0^2 k_\perp^2 (C - 2\alpha B_0^2 k_\perp^2)} \right]. \end{aligned} \quad (\text{A } 14)$$

Substituting  $C = 2\alpha\sigma B_0^2 k_\perp^2$  in this expression we arrive at the second expression in Eq. (6.21).

Now we consider the case where  $\alpha < 0$ .  $I_1$  is given by Eq. (A1) with the upper sign. Using the variable substitution

$$u = e^{X/L} + \frac{1}{\sqrt{\sigma}} \quad (\text{A15})$$

we transform the expression for  $I_1$  to

$$I_1 = -\frac{8\zeta^3\sqrt{-VC}}{\alpha} \int_{1/\sqrt{\sigma}}^{\infty} \frac{u - 1/\sqrt{\sigma}}{(u^2 - \zeta^2)^2} du, \quad (\text{A16})$$

where  $\zeta = \sqrt{1/\sigma - 1}$ . Using the expansion

$$\begin{aligned} \frac{u - \sqrt{\sigma}}{(u^2 - \zeta^2)^2} &= \frac{1}{4\zeta^3\sqrt{\sigma}} \left( \frac{1}{u - \zeta} - \frac{1}{u + \zeta} \right) + \left( \frac{1}{4\zeta} - \frac{1}{4\zeta^3\sqrt{\sigma}} \right) \frac{1}{(u - \zeta)^2} \\ &\quad - \left( \frac{1}{4\zeta} + \frac{1}{4\zeta^3\sqrt{\sigma}} \right) \frac{1}{(u + \zeta)^2} \end{aligned} \quad (\text{A17})$$

we obtain

$$\int_{1/\sqrt{\sigma}}^{\infty} \frac{u - \sqrt{\sigma}}{(u^2 - \zeta^2)^2} du = \frac{1}{4\zeta^3\sqrt{\sigma}} \ln \frac{1 + \sqrt{1 - \sigma}}{1 - \sqrt{1 - \sigma}} - \frac{1}{2\zeta^2}. \quad (\text{A18})$$

Using Eqs. (A11), (A16), and (A18), and the expression for  $C$  and  $\zeta$  in terms of  $\sigma$  we obtain the first expression in Eq. (6.26).

Now we proceed to calculating  $I_2$ .  $I_2$  is given by Eq. (A9) with the upper sign. Using the variable substitution defined by Eq. (A15) we transform it to

$$I_2 = \frac{16\ell C\zeta^5}{\alpha} \sqrt{\frac{2V}{|\alpha|}} \int_{1/\sqrt{\sigma}}^{\infty} \frac{(u - 1/\sqrt{\sigma})^2}{(u^2 - \zeta^2)^3} du. \quad (\text{A19})$$

Using the expansion

$$\begin{aligned} \frac{(u - 1/\sqrt{\sigma})^2}{(u^2 - \zeta^2)^3} &= \left(1 - \frac{3}{\zeta^3\sqrt{\sigma}}\right) \left(\frac{1}{u + \zeta} - \frac{1}{u - \zeta}\right) + \frac{1}{16\zeta^2} \left(1 - \frac{2}{\zeta\sqrt{\sigma}}\right. \\ &\quad \left. - \frac{3}{\zeta^2}\right) \frac{1}{(u + \zeta)^2} + \frac{1}{16\zeta^2} \left(1 + \frac{2}{\zeta\sqrt{\sigma}} - \frac{3}{\zeta^2}\right) \frac{1}{(u - \zeta)^2} \\ &\quad - \frac{1}{8\zeta} \left(1 + \frac{1}{\zeta\sqrt{\sigma}}\right)^2 \frac{1}{(u + \zeta)^3} + \frac{1}{8\zeta} \left(1 - \frac{1}{\zeta\sqrt{\sigma}}\right)^2 \frac{1}{(u - \zeta)^3} \end{aligned} \quad (\text{A20})$$

we obtain

$$\begin{aligned} \int_{1/\sqrt{\sigma}}^{\infty} \frac{(u - 1/\sqrt{\sigma})^2}{(u^2 - \zeta^2)^3} du &= \frac{1}{16\zeta^3} \left(1 - \frac{3}{\sigma\zeta^2}\right) \ln \frac{1 - \zeta\sqrt{\sigma}}{1 + \zeta\sqrt{\sigma}} \\ &\quad + \frac{\sqrt{\sigma}}{16\zeta^2} \left(1 - \frac{2}{\zeta\sqrt{\sigma}} - \frac{3}{\zeta^2}\right) \frac{1}{1 + \zeta\sqrt{\sigma}} + \frac{\sqrt{\sigma}}{16\zeta^2} \left(1 + \frac{2}{\zeta\sqrt{\sigma}} - \frac{3}{\zeta^2}\right) \frac{1}{1 - \zeta\sqrt{\sigma}} \\ &\quad - \frac{\sigma}{16\zeta} \left(1 + \frac{1}{\zeta\sqrt{\sigma}}\right)^2 \frac{1}{(1 + \zeta\sqrt{\sigma})^2} + \frac{\sigma}{16\zeta} \left(1 - \frac{1}{\zeta\sqrt{\sigma}}\right)^2 \frac{1}{(1 - \zeta\sqrt{\sigma})^2}. \end{aligned} \quad (\text{A21})$$

Using Eqs. (A19) and (A21), and the expression for  $C$  and  $\zeta$  in terms of  $\sigma$  we obtain the second expression in Eq. (6.26).

## References

- Aharonian, F. A., Bogovalov, S. V., and Khangulyan, D. 2012 Abrupt acceleration of a ‘cold’ ultrarelativistic wind from the Crab pulsar. *Nature* **482**, 507-509.
- Arons, J. and Barnard, J. J. 1986 Wave propagation in pulsar magnetospheres: Dispersion relations and normal modes of plasmas in superstrong magnetic fields. *Astrophys. J.* **302**, 120-155.
- Bell, A. R. and Kirk, J. G. 2008 Possibility of prolific pair production with high-power lasers. *Phys. Rev. Lett.* **101**, 200403.
- Cattaert, T., Kourakis, I., and Shukla, P. K. 2005 Envelope solitons associated with electromagnetic waves in a magnetized pair plasma. *Phys. Plasmas* **12**, 012319
- Cerutti, B. and Beloborodov, A. M. 2017 Electrodynamics of pulsar magnetospheres. *Space Sci. Rev.* **207**, 111-136.
- Chen, H., Wilks, S. C., Bonlie, J. D., Liang, E. P., Myatt, J., Price, D. F., Meyerhofer, D. D., and Beiersdorfer, P. 2009 Making relativistic positrons using ultraintense short pulse lasers. *Phys. Rev. Lett.* **102**, 105001.
- Chian, A. C.-L. and Kennel, C. F. 1983 Self-modulational formation of pulsar microstructures. *Astrophys. Space Sci.* **97**, 9-18.
- Chew, G. F., Goldberger, M. L., and Low, F. E. 1956 The Boltzmann equation and the one-fluid hydromagnetic equations in the absence of particle collisions. *Proc. Roy. Soc. A* **236**, 112-118.
- Cohen, R. H. and Kulsrud, R. M. 1974 Nonlinear evolution of parallel-propagating hydromagnetic waves. *Phys. Fluids* **17**, 2215-2225.
- El-Labany, S. K., El-Shamy, E. F., Sabry, R., and Khedr, D. M. 2013 The interaction of two nonplanar solitary waves in electron-positron-ion plasmas: An application in active galactic nuclei. *Phys. Plasmas* **20**, 012105.
- Fedun, V. M., Ruderman, M. S., and Erdélyi, R. 2008 Generation of short-lived large-amplitude magnetohydrodynamic pulses by dispersive focusing. *Phys. Lett. A* **372**, 6107-6110.
- Gailis, R. M., Frankel, N. E., and Dettmann, C. P. 1995 Magnetohydrodynamics in the expanding Universe. *Phys. Rev. D* **52**, 6901-6917.
- Gahn, C., Tsakiris, G. D., Pretzler, G., Witte, K. J., Delfin, C., Wahlström, C. G., and Habs, D. 2000 Generating positrons with femtosecond-laser pulses. *Appl. Phys. Lett.* **77**, 2662.
- Greaves, R. G., Tinkle, M. D., and Surko, C. M. 1994 Creation and uses of positron plasmas. *Phys. Plasmas* **1**, 1439-1446.
- Ichikawa, Y.-H., Konno, K., Wadati, M., and Sanuki, H. 1980 Spiky soliton in circular polarized Alfvén wave. *J. Phys. Soc. Jpn.* **48**, 279-286.
- Ikezi, H. 1973 Experiment on ion-acoustic solitary waves. *Phys. Fluids* **16**, 1668-1675.
- Iwamoto, N. 1993 Collective modes in nonrelativistic electron-positron plasmas. *Phys. Rev. E* **47**, 604-611.
- Kadomtsev, B. B. and Petviashvili, V. I. 1970 On the stability of solitary waves in weakly dispersing media. *Sov. Phys. Dokl.* **15**, 539-541. [Translated from Russian: Kadomtsev, B. B. and Petviashvili, V. I. 1970 Ob ustoychivosti uedinyonnyh voln v slabo dispergiruyushchih sredah. *Doklady Akademii Nauk SSSR* **192**, 753-756.]
- Kakutani, T., Ono, H., Taniuti, T., and Wei, C.-C. 1968 Reductive perturbation method in nonlinear wave propagation II. Application to hydromagnetic waves in cold plasma. *J. Phys. Soc. Japen* **24**, 1159-1166.
- Kaup, D. J. and Newell, A. C. 1978 An exact solution for a derivative nonlinear Schrödinger equation. *J. Math. Phys.* **19**, 798-801.
- Kawakatu, N., Kino, M., and Takahara, F., 2016 Evidence for a significant mixture of electron/positron pairs in FR II jets constrained by cocoon dynamics. *Month. Not. Roy. Astr. Soc.* **457**, 1124-1136.
- Kawata, T. and Inoue, H. 1978 Exact solutions of the derivative nonlinear Schrödinger equation under the nonvanishing conditions. *J. Phys. Soc. Jpn.* **44**, 1968-1976.
- Khanna, M. and Rajaram, R. 1982 Evolution of nonlinear Alfvén waves propagating along



- the magnetic field in a collisionless plasma. *J. Plasma Phys.* **28**, 459-468.
- Lakhina, G. S. and Verheest, F. 1997 Alfvénic solitons in ultrarelativistic electron-positron plasmas. *Astrophys. Space Sci.* **253**, 97-106.
- Liang, E. P., Wilks, S. C., and Tabak, M. 1998 Pair production by ultraintense lasers. *Phys. Rev. Lett.* **91**, 4887-4890.
- Lonngren, K. E. 1983 Soliton experiments in plasma physics. *Plasma Phys. Contr. Fusion* **25**, 943-982.
- Mikhailovskii, A. B., Onishchenko, O. G., and Smolyakov, A. I. 1985a Theory of low-frequency electromagnetic solitons in a relativistic electron-positron plasma. *Sov. J. Plasma Phys.* **11**, 215-219. [Translated from Russian: Mikhailovskii, A. B., Onishchenko, O. G., and Smolyakov, A. I. 1985a Teoriya nizkochastotnykh elektromagnitnykh solitonov v relativistskoi elektronno-positronnoy plasme. *Fizika Plazmy* **11**, 369-375.]
- Mikhailovskii, A. B., Onishchenko, O. G., and Tatarinov, E. G. 1985b Alfvén solitons in a relativistic electron-positron plasma. I. Hydrodynamic theory. *Plasma Phys. Contr. Fusion* **27**, 527-537.
- Mikhailovskii, A. B., Onishchenko, O. G., and Tatarinov, E. G. 1985c Alfvén solitons in a relativistic electron-positron plasma. II. Kinetic theory. *Plasma Phys. Contr. Fusion* **27**, 539-556.
- Mio, K., Ogino, T., Minami, K., and Takeda, S. 1976a Modified nonlinear Schrödinger equation for Alfvén waves propagating along the magnetic field in cold plasmas. *J. Phys. Soc. Japen.* **41**, 265-271.
- Mio, K., Ogino, T., Minami, K., and Takeda, S. 1976b Modulational instability and envelope-solutions for nonlinear Alfvén waves propagating along magnetic-field in plasmas. *J. Phys. Soc. Japen.* **41**, 667-673.
- Mjølhus, E. 1976 On the modulational instability of hydromagnetic waves parallel to the magnetic field. *J. Plasma Phys.* **16**, 321-334.
- Mjølhus, E. and Hada, T. 1997 in: T. Hada, H. Matsumoto (Eds.), Nonlinear waves and chaos in space plasmas. *Terrapub, Tokio*, p. 121.
- Mjølhus, E. and Wyller, J. 1986 Alfvén solitons. *Phys. Scripta* **33**, 442-451.
- Passot, T. and Sulem, P. -L. 1993 Multidimensional modulation of Alfvén waves. *Phys. Rev. E* **48**, 2966-2974.
- Rajib, T. I., Sultana, S., and Mamun, A. A. 2015 Solitary waves in rotational pulsar magnetosphere. *Astrophys. Space Sci.* **357**, 52.
- Rogister, A. 1971 Parallel propagation of nonlinear low-frequency waves in high- $\beta$  plasma. *Phys. Fluids* **14**, 2733-2739.
- Rudenko, O. V. and Soluyan, S. I. 1977 *Theoretical foundations of nonlinear acoustics*. New York: Plenum Publishing Corporation, Consultants Bureau.
- Ruderman, M. S. 1986 Stability of quasilongitudinally propagating narrow beams of nonlinear magnetohydrodynamic waves. *Fluid Dynamics* **21**, 524-530. [Translated from Russian: Ruderman, M. S. 1986 Qvaziparallelnoe raspropstranenie uzkih puchkov nelineinykh magnitogidrodinamicheskikh voln. *Izvestiya Akademii Nauk SSSR, Mekhanika Zhidkosti i Gaza*, No. 4, 21-28.]
- Ruderman, M. S. 1987 Quasilongitudinally propagating solitons in a plasma with Hall dispersion. *Fluid Dynamics* **22**, 299-305. [Translated from Russian: Ruderman, M. S. 1987 Ustoichivost' kvaziprodol'no rasprostranyayushchihsya solitonov v plasme s hollovskoi dispersiey. *Izvestiya Akademii Nauk SSSR, Mekhanika Zhidkosti i Gaza*, No. 2, 159-165.]
- Ruderman, M. S. 2002 DNLS equation for large-amplitude solitons propagating in an arbitrary direction in a high- $\beta$  Hall plasma. *J. Plasma Phys.* **67**, 271-276.
- Ruderman, M. A. and Sutherland, P. G. 1975 Theory of pulsars: polar gaps, sparks, and coherent microwave radiation. *Astrophys. J.* **196**, 51-72.
- Ruffini, R., Vereshchagin, G., and Xue, S.-S. 2010 Electron-positron pairs in physics and astrophysics: From heavy nuclei to black holes. *Physics Reports* **487**, 1-140.
- Sakai, J. and Kawata, T. 1980a Waves in an ultra-relativistic electron-positron plasma. *J. Phys. Soc. Japan* **49**, 747-752.

- Sakai, J. and Kawata, T. 1980b Non-linear Alfvén-wave in an ultra-relativistic electron-positron plasma. *J. Phys. Soc. Japan* **49**, 753-758.
- Sarri, G., Schumaker, W., di Piazza, A., Vargas, M., Dromey, B., Dieckmann, M. E., Chvykov, V., Maksimchuk, A., Yanovsky, V., He, Z. H. 2013 Table-top laser-based source of femtosecond, collimated, ultrarelativistic positron beams. *Phys. Rev. Lett.* **110**, 255002.
- Shukla, R. K. 2003 Generation of magnetic fields in the early universe. *Phys. Lett. A* **310**, 182-186.
- Shukla, P. K., Rao, N. N., Yu, M. Y., and Tsintsadze, N. L. 1986 Relativistic nonlinear effects in plasmas. *Phys. Rep.* **135**, 1-149.
- Stewart, G. A. and Laing, E. W. 1992 Wave propagation in equal-mass plasmas. *J. Plasma Phys.* **47**, 295-319.
- Sturrock, P. A. 1971 A model of pulsars. *Astrophys. J.* **164**, 529-556.
- Surko, C. M. and Murphy, T. 3. 1990 Use of the positron as a plasma particle. *Phys. Fluids B* **2**, 1372-1375.
- Surko, C. M., Leventhal, M., and Passner, A. 1989 Positron plasma in the laboratory. *Phys. Rev. Lett.* **62**, 901-904.
- Taniuti, T. and Wei, C.-C. 1968 Reductive perturbation method in nonlinear wave propagation. I. *J. Phys. Soc. Japan* **24**, 941-946.
- Tatsuno, T., Berezhiani, V. I., Pekker, M., and Mahajan, S. M. 2003 Angular momenta creation in relativistic electron-positron plasma. *Phys. Rev. E* **68**, 016409.
- Tran, M. Q. 1979 Ion-acoustic solitons in a plasma-review of their experimental properties and related theories. *Phys. Scripta* **20**, 317-327.
- Verheest, F. 1996 Solitary Alfvén modes in relativistic electron-positron plasmas. *Phys. Lett. A* **213**, 177-182.
- Verheest, F. and Lakhina, G. S. 1996 Oblique solitary Alfvén modes in relativistic electron-positron plasmas. *Astrophys. Space Sci.* **240**, 215-224.
- Whitham, G. B. 1974 *Linear and nonlinear waves*. New York: Willey-Interscience Publication.
- Yajima, N. 1966 Effect of finite ion Larmor radius on propagation of magnetoacoustic waves. *Prog. Theor. Phys.* **36**, 1-16.
- Zabolotskaya, E. A. and Khokhlov, R. V. 1969 Quasi-plane waves in the nonlinear acoustics of confined beams. *Soviet Phys. Acoust. - USSR* **15**, 35-42. [Translated from Russian: Zabolotskaya, E. A. and Khokhlov, R. V. 1969 Kvazi-ploskie volny v nonlineinoi akustike ogranichennyh puchkov. *Akoust. Zh.*, **15**, 40-47.]
- Zank, G. P. and Greaves, R. G. 1995 Linear and nonlinear modes in nonrelativistic electron-positron plasmas. *Phys. Rev. E* **51**, 6079-6090.