



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/163212/>

Version: Accepted Version

---

**Article:**

Butler, RWH, Lickorish, HW, Vinnels, J et al. (2020) Untangling the Annot sand fairway: structure and stratigraphy of the Eastern Champsaur Basin (Eocene-Oligocene), French Alps. *Journal of the Geological Society*. jgs2020-015. ISSN: 0016-7649

<https://doi.org/10.1144/jgs2020-015>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

1 **Untangling the Annot sand fairway: structure and stratigraphy**  
2 **of the Eastern Champsaur Basin (Eocene-Oligocene), French**  
3 **Alps.**

4

5 Robert W.H. Butler<sup>1</sup>, Henry W. Lickorish<sup>2</sup>, Jamie Vinnels<sup>3</sup> & William D.  
6 McCaffrey<sup>2</sup>

7

8 1 Geology and Geophysics, School of Geosciences, University of Aberdeen,  
9 Aberdeen AB24 3UE, UK

10 2 School of Earth and Environment, University of Leeds, Leeds LS2 9JT, UK

11 3 Equinor, 2107 City West Blvd, Houston, Texas, 77042, USA

12

13

14 Orchid: RWHB, 0000-0002-7732-9686

15 Correspondence: [rob.butler@abdn.ac.uk](mailto:rob.butler@abdn.ac.uk)

16

17 **ABSTRACT**

18 Early foredeep successions can yield insight on tectonic processes operating  
19 adjacent to and ahead of fledgling orogenic belts but are commonly deformed  
20 by the same orogens. We develop a workflow towards stratigraphic  
21 understanding of these deformed basins, applied to the Eastern Champsaur  
22 Basin of the French Alps. This contains a down-system correlative of the  
23 southern-sourced (Eocene-Oligocene) Annot turbidites. These strata are  
24 deformed by arrays of W-facing folds that developed beneath the Embrunais-  
25 Ubaye tectonic allochthon. The folds vary in geometry through the  
26 stratigraphic multilayer. Total shortening in the basin is around 4 km and the  
27 restored (un-decompacted) stratal thickness exceeds 980m. The turbidites  
28 are generally sand-rich and bed-sets can be correlated through the entire fold  
29 train. The succession shows onlap and differential thickening indicating  
30 deposition across palaeobathymetry that evolved during active basement  
31 deformation, before being over-ridden by the allochthon. The sand system  
32 originally continued over what is now the Ecrins basement massif that, while  
33 contributing to basin floor structure, only served to confine and potentially

34 focus further sediment transport to the North. Deformation ahead of the main  
35 Alpine orogen appears to have continued progressively, and the past  
36 definition of distinct “phases” (“pre-“ and “post-Nummulitic”) is an artefact of  
37 the stratigraphic record.

38

39 Keywords: turbidites, structural restoration, foreland basin,

40

41 Turbidite systems can offer important constraints on bathymetric continuity of  
42 arrays of sedimentary basins (e.g. Smith 2004; Puigdefabregas *et al.* 2004)  
43 and thus inform models of basin evolution. For ancient foredeep systems the  
44 challenge is to unravel useful stratigraphic information from successions that  
45 are involved in orogenic belts and then deformed. If stratigraphic information  
46 can be extracted from deformed systems, the insights so gained can then not  
47 only inform models of orogenic evolution but also impact on greater tectono-  
48 stratigraphic understanding of turbidite systems. This is especially important  
49 as many of the key ideas of turbidite sedimentation come from studies of  
50 successions in syn-orogenic sedimentary basins. The aims of this paper are  
51 two-fold. The first is to document a workflow for developing stratigraphic  
52 knowledge from highly folded successions. The second is to provide a linked  
53 stratigraphic and structural reconstruction of the Eastern Champsaur Basin –  
54 a deformed part of the Eocene-Oligocene Annot Sandstone system of SE  
55 France (the Grès d’Annot in Alpine literature: e.g. Sinclair 1997; Joseph &  
56 Lomas 2004) that has become incorporated into the Western Alpine orogen.  
57 While an extensive re-examination of the Annot system in the light of our  
58 reconstructions lies beyond the scope of this paper, we do explore some  
59 implications for the extent of the system, how crustal-scale deformation  
60 migrates through time and how tectonic evolution is represented in the  
61 stratigraphic record.

62

### 63 **Geological context and motivation**

64

65 The Annot system of SE France as a whole occupies a pivotal position  
66 in turbidite research, with outcrops providing test-beds for models of deep-  
67 water sedimentation and analogues for hydrocarbon reservoirs in the

68 subsurface. Bouma's pioneering descriptions of turbidite facies come from  
69 proximal parts of the Annot system (e.g. Bouma 1959; Stanley & Bouma  
70 1964). This turbidite system was deposited in a broadly north-south trending  
71 foredeep basin, adjacent to the ancestral Alps and fed from eroding granitic  
72 basement to the south (Maures-Esterel massif, Fig. 1a) together with Corsica  
73 that, at the time, lay adjacent to the south coast of mainland France. Much of  
74 the early research is cited by Joseph & Lomas (2004) and papers that follow  
75 this introduction greatly expand upon these works. More recent studies  
76 include Mulder *et al.* (2010), Etienne *et al.* (2012), Salles *et al.* (2014), and  
77 Cunha *et al.* (2017).

78 Existing work has almost entirely focussed on sites that have  
79 experienced rather little deformation. The system is known to continue into  
80 more strongly deformed regions to the north and northeast (e.g. Sinclair  
81 1997). By increasing understanding of the Annot system, insight is gained on  
82 the continuity and early deformation of the Alpine "foreland". As subaqueous  
83 gravity flows, turbidity currents seek low bathymetry. Therefore, the continuity  
84 of sand fairways, the geological record of the routes taken by their causative  
85 flows, provides evidence for relative bathymetry in front of the evolving Alpine  
86 orogen. The various relationships between the turbidites and their substrate  
87 chart not only the syndepositional deformation of the foredeep basin floor but  
88 also the deformation that predated basin subsidence. This understanding is  
89 important for evaluating the tectonic significance of unconformities associated  
90 with orogens and their implications for evaluating the progression of  
91 continental deformation. By increasing understanding of the Annot system,  
92 insight is gained on the continuity and early deformation of the Alpine  
93 "foreland".

94 Our case study here focusses on the south-eastern flank of the  
95 external Alpine basement massif of the Ecrins (Fig. 1a). The Eocene-  
96 Oligocene litho-stratigraphy of the area, in common with the main Annot  
97 system (Fig 2; e.g. Joseph & Lomas 2004), consists of three main lithofacies  
98 (Debelmas *et al.* 1980) that comprise the "Nummulitic trilogy". Collectively this  
99 represents the foredeep megasequence developed during this early stage of  
100 Alpine orogenesis. The megasequence oversteps a substrate of crystalline  
101 basement, continuous with the neighbouring Ecrins massif, together with

102 remnant patches of the Triassic-Jurassic cover to the massif (e.g. Debelmas  
103 *et al.* 1980). The eastern and southern margins to the basin are now buried  
104 beneath the tectonic allochthon of the Embrunais-Ubaye thrust sheets.

105         The first part of the Nummulitic trilogy is the Nummulitic Limestone, a  
106 shallow-water transgressive unit here of Priabonian age (Debelmas *et al.*  
107 1980; Dumont *et al.* 2012). Locally, the Nummulitic Limestone is underlain by  
108 conglomerates derived from the adjacent crystalline basement, interpreted to  
109 be the fills to subaerially-incised palaeovalleys (e.g. Gupta 1997). The second  
110 part of the Nummulitic trilogy are the Blue Marls (“Marnes à Globigérines” on  
111 published maps, e.g. Debelmas *et al.* 1980). These calcareous mudstones  
112 chart palaeo-bathymetries increasing to several hundred metres depth. The  
113 final part of the trilogy are turbidites – the Champsaur Sandstone (“Grès du  
114 Champsaur” on published maps, e.g. Debelmas *et al.* 1980). Locally, there is  
115 a facies intermediate between these turbidites and the Blue Marls, the Brown  
116 Marls (e.g. Stanbrook & Clark 2004). These are generally inferred to have  
117 accumulated on local intra-basin highs from dispersed suspension clouds  
118 associated with the turbidity currents that deposited the Champsaur (and main  
119 Annot) Sandstone.

120         The Nummulitic trilogy is capped by a shaley olistostrome termed the  
121 “Schistes à Blocs” that lies directly under the Embrunais-Ubaye thrust sheets.  
122 It is classically inferred to have been shed off the advancing thrust sheets as  
123 the foredeep basin was closed (Kerckhove 1969). Detailed sedimentological  
124 investigation of the Schistes à Blocs in the Eastern Champsaur Basin is  
125 strongly inhibited by the intense penetrative deformation within this unit, which  
126 presumably reflects distributed shear deformation in the footwall to the  
127 overlying thrust sheets. Consequently, it is generally difficult to establish  
128 whether the unit represents a series of amalgamated debris flow deposits or is  
129 the product of a single submarine mass-wasting event.

130         Although given the single designator “Champsaur Sandstone”, early  
131 Oligocene siliciclastic sandstones along the southern flank of the Ecrins  
132 massif have two distinct compositions and, by inference, provenances, lying in  
133 distinct depocentres (Debelmas *et al.* 1980). Following our previous usage  
134 (Vinnels *et al.* 2010; Butler 2017), we term these depocentres the Western  
135 and Eastern Champsaur Basins. They are separated by a zone of

136 deformation generally termed the Selle Fault (e.g. Tricart 1981; Ford 1996  
137 and others since). To date, sedimentological research in the region has  
138 mainly focussed on the westerly-derived, volcanoclastic turbidites that outcrop  
139 in the Western Champsaur Basin, with detailed descriptions not only of  
140 depositional architectures (e.g. Brunt *et al.* 2007) but also of the discrete fold-  
141 thrust structures that have deformed the successions (Butler & McCaffrey  
142 2004). Separated from this western basin by the Selle Fault Zone, the Eastern  
143 Champsaur Basin is significantly more deformed than its western counterpart  
144 (Fig. 3). It is characterised by folds on various wavelengths and amplitudes  
145 that generally have tight hinges and straight limbs, approximating to chevron  
146 geometry. This style of deformation is typical of successions with a  
147 pronounced planar mechanical anisotropy such as is provided by inter-  
148 bedded sandstones and shales (Ramsay 1974). The folds face westwards  
149 with axial surfaces inclined moderately to the east. The structures were  
150 described by Kerckhove *et al.* (1978) and studied in detail by Bürgisser & Ford  
151 (1998). These studies show that the folds developed beneath, and locally  
152 incorporate, the Embrunais Ubaye thrust sheets which overran both the  
153 Eastern and Western Champsaur Basins. The thrust sheets provided tectonic  
154 burial sufficient for peak temperatures in Eastern Champsaur of c 300 °C  
155 (Bellanger *et al.* 2015).

156 Provenance of the Eastern Champsaur Sandstone has been disputed.  
157 Sinclair (1997) depicts their turbidites to be the down-system continuation of  
158 the Annot system otherwise preserved in the SW Alps of Provence. However,  
159 in their regional overview, Joseph & Lomas (2004) indicate a northern  
160 provenance, sourced from the an elevated, proto-Ecrins massif. The  
161 argument is resolved by Vinnels *et al.* (2010) who present palaeocurrent data  
162 to show a southerly derivation and document that the sandstones of the  
163 Eastern Champsaur basin have the same composition as the Annot  
164 Sandstone in its type area to the south. Thus Sinclair's (1997) view is  
165 supported. Vinnels *et al.* (2010) note that, although deflected by subtle basin-  
166 floor topography, the causative turbidity currents continued over what is now  
167 the uplifted Ecrins basement massif. However, Vinnels *et al.* only focused on  
168 the lower stratigraphic intervals, establishing facies variations and palaeoflow  
169 deflections along the basal onlap surface of turbidites onto the Blue Marls that

170 drape uplifted substrate within the basin. The current paper focusses  
171 exclusively on the Eastern Champsaur Basin, building on our previous work  
172 (Vinnels *et al.* 2010). The regional structural setting is discussed extensively  
173 by Butler 2017). The stratigraphic relationships between the eastern and  
174 western Champsaur Basins and the role of the Selle Fault in partitioning  
175 distinct deep-water systems lie beyond the scope of the current paper and are  
176 reserved for later contributions.

177

## 178 **A workflow**

179

180 Geologists working in mountain belts have long had to grapple with the  
181 complexities of deformation when attempting to build stratigraphic knowledge.  
182 However, the methods used for creating this understanding are rarely  
183 documented, making it difficult to assess uncertainties in the larger-scale  
184 basin reconstructions. Furthermore, there has been a reluctance for many  
185 researchers investigating sedimentary basins associated with mountain belts  
186 to examine their more-deformed components. In presenting a workflow here  
187 we hope not only to promote research within these settings but also to  
188 encourage clearer documentation of how deformed stratigraphic sections are  
189 reconstructed.

190 The Champsaur area exhibits just under 2 km of topographic relief with  
191 cliff sections that are of limited accessibility, attributes that are typical of much  
192 of the Alpine mountain ranges and of young orogenic belts in general. A  
193 virtue of the relief in our Champsaur study area is that major hillsides trend  
194 east-west, perpendicular to the trend of fold axes. They therefore provide  
195 ideal cross-sections, akin to in-lines in grids of seismic reflection data.  
196 Accessible routes up these hillsides provide excellent vertical sections,  
197 equivalent to boreholes for subsurface examples. Consequently, we apply a  
198 workflow similar to that adopted for surface stratigraphic-structural mapping,  
199 tying detailed stratigraphic observations from vertical sections (boreholes) to  
200 remotely sensed cross-sections (in this case by direct observation from  
201 opposite valley-sides, equivalent to seismic profiles).

202 The first decision lies in selecting the best vertical section from which to  
203 build a stratigraphic succession. The west face of Le Piquet provides a strike

204 section, 3.5 km across, that runs from up from the base-Nummulitic  
205 unconformity to a small klippe of Embrunais-Ubaye thrust sheets (Fig. 5a).  
206 Only the upper part of this hillside provides continuous outcrop that is also  
207 amenable to direct observation – and this yields stratigraphic logs (Fig. 5a).  
208 Logged units include a c 380m interval that is sandstone-dominated (labelled  
209 A6 on Fig. 5a), capped by Schistes à Blocs. However, the broad bed-set  
210 characteristics of the underlying succession may be established by remote  
211 observation. These include shale-dominated levels that are more recessive on  
212 hillsides (A3 and A5 on Fig. 5a).

213 The boundaries between the bed-set units can be traced through the  
214 hillsides on Le Piquet and across the Dourmillouse Valley onto the dip section  
215 provided by its northern slopes (Fig. 5b). The upper part of the valley side,  
216 including the ridge line, provides near continuous, if largely inaccessible,  
217 outcrop within which the main folds can be identified. The fold axial traces can  
218 be mapped down dip into the Dourmillouse Valley. Then the bed-set  
219 stratigraphic units of the Champsaur sandstone can be traced through the fold  
220 train. This section is moderately accessible and locally provides accessible  
221 sections for detailed stratigraphic logging.

222 The axial traces of the folds provide the key linking structures to tie  
223 adjacent transects. These can be traced over the ridge line bounding the  
224 Dourmillouse Valley to the north and into the adjacent Fornel Valley (Fig. 4).  
225 The northern side of the ridge-line is much steeper than its southern  
226 counterpart so that accessibility is severely restricted. However, fold axial  
227 traces may be readily traced through the cliffs and the bed-set stratigraphy  
228 defined and similarly tracked through the fold array (Fig. 5c). Note that there is  
229 some polyharmonic folding but the axial surfaces for the main fold closures  
230 may be traced from the ridge-line down to the Blue Marls.

231

### 232 **The bed-set stratigraphy**

233

234 Following the workflow laid out above it is possible to map out not only  
235 the axial traces of the principal folds but also the bed-set stratigraphic units  
236 within the Champsaur sandstone (Fig. 4). Although structural deformation  
237 hinders identification of individual beds across the whole basin, thicker

238 intervals can be laterally traced continuously for several kilometres and  
239 probably extend much further. Vertical stratigraphic sections display  
240 significant heterogeneity in sandstone to shale ratio. These are well-illustrated  
241 by the cliff sections of Le Piquet (Fig. 5a),

242         As noted above, it is the variety in sandstone-shale abundance that  
243 provides a framework, on the bed-set scale, for dividing the basin stratigraphy  
244 into mappable units (Fig. 5). These units are illustrated on arrays of composite  
245 logs (Fig. 6). Principally, our sedimentary logs were measured from the fold  
246 limbs to avoid hinge areas, where beds show significant layer parallel  
247 shortening strain manifested by cleavage, and tracts of strongly overturned  
248 strata where bed thinning is likely. Strains in both of these structural settings  
249 are localised in the shales – sandstone beds retain unmodified sedimentary  
250 structures suggesting penetrative strains in these units are very low. We  
251 estimate an uncertainty of less than 10% on the normally-compacted  
252 thickness of sandstones and perhaps up to 20% for the shales on the values  
253 reported in Fig. 6, as we cannot rule out deformation even in the normal limbs  
254 of folds. The bed-set sequences may still be identified regardless of strain  
255 variations and we now use these to erect a basin-wide lithostratigraphy.

256         The Champsaur Sandstone onlaps its substrate towards the western  
257 basin margin (Vinnels *et al.* 2010 and references therein): the oldest  
258 preserved parts of the sandstone sequence are only present on the vegetated  
259 lower slopes of the far eastern exposures of the basin. In the Narreyroux  
260 valley (log location 4 on Fig. 4), a 670m thick section of Champsaur  
261 Sandstone (A4 on Fig. 6) is underlain to the east by considerably more  
262 stratigraphy but precise correlation has proven elusive. The higher parts of  
263 the stratigraphy are more tractable. Two levels are especially sandstone-poor  
264 and form key markers (A3 and A5 in Fig. 6). The lower of these two shaley  
265 intervals is approximately 50m thick and contains a distinctive couplet of 2 m  
266 thick sandstone beds that can be found in all sections that include this  
267 stratigraphic interval. There are a few thin sandstone beds present in the  
268 higher shale-dominated interval (A5 in Fig. 4) but these are laterally  
269 discontinuous. Between the two shaley sections lies an interval with  
270 abundant thick sandstone beds (A4 in Fig 4). The section below the lower  
271 shaley interval also has higher sandstone content; its upper portion contains

272 beds that are 3-5 m thick while the lower portion is bedded on the 50 cm – 1  
273 m scale. These distinctions provide confirmations of stratigraphic correlations  
274 to elsewhere within the Eastern Champsaur district.

275 The lower shaley interval (A3) can be readily correlated between the  
276 Narreyroux, Fournel and Dourmillouse valleys (Fig. 4) to parts of the study  
277 area where more complete sections of the underlying sandstone-rich intervals  
278 are exposed. In these places we can separate a thicker-bedded upper  
279 component from a thinner-bedded lower component (A2 and A1 respectively  
280 in Fig. 6). The thicker-bedded interval (A2) is about 40 m thick. The logs from  
281 the Fournel and Dourmillouse valleys (Fig. 6) only demonstrate about 60 m of  
282 this lower portion. However, in the Narreyroux valley it achieves a thickness in  
283 excess of 150 m with a further 200 m represented by the poorly exposed and  
284 deformed terrain deeper in the section.

285 The upper shaley horizon defined in the Narreyroux valley (A5 on Fig.  
286 6) can be correlated across the Dourmillouse valley (Fig. 4) to the cliffs of Le  
287 Piquet. This is the “Thick Shale” of Vinnels *et al.* (2010). Overlying this, 380  
288 m of sandstone-rich turbidies (A6 on Fig. 4) are capped by the “Schistes a’  
289 Blocs” that directly underlie the Embrunais-Ubaye thrust sheets. The  
290 sandstone rich interval (A6) can be traced through the fold structures of the  
291 upper Dourmillouse and Fournel Valleys, and constitute the upper 60m of the  
292 logged section in the Narreyroux valley (Fig. 4). In the upper Fournel Valley  
293 the thickness of this interval exceeds 400 m (log 5 on Fig. 6). Individual beds  
294 may be traced through folds and across the ridge line between the upper  
295 parts of the Dourmillouse and Fournel valleys. For example, a prominent  
296 shale interval (5-15 m thick; X on logs 5 and 6 on Fig. 6), at the base of  
297 interval A6b) can be traced between sections.

298 The Eastern Champsaur Basin continues over the watershed at the  
299 head of the Dourmillouse Valley into drainage of the Champoleon Valley (Fig.  
300 4). Individual beds may be traced along the ridge line (Pointe des Estaris -  
301 Pointe des Pisses: log locations 8 and 9, Fig. 4) to complete the upper  
302 stratigraphy of this part of the Champsaur Sandstone up to the Schistes à  
303 Blocs (Fig. 6).

304 By combining the logs, we estimate that about 950m of Champsaur  
305 Sandstone stratigraphy (A6 – A7) overlies the prominent upper shaley horizon

306 (A5) in the ground between the upper Dourmillouse valley and Pointe des  
307 Pisses (Fig. 4). The equivalent interval at Le Piquet is just 380m thick (log 1  
308 on Fig. 6). Both sections are capped by the Schistes à Blocs so the thickness  
309 difference is not due to post-depositional tectonic truncation. The difference is  
310 also too great to be due to variations in distributed deformation or differential  
311 compaction and there are significant differences in the bed-set stratigraphy  
312 between these locations. We interpret the thickness variation to result from  
313 differential stratigraphic growth across the Eastern Champsaur Basin.

314

### 315 Depositional architecture and growth

316         Although folding permits construction of substantial stratigraphic  
317 sections and deformation has not been sufficient to prevent individual bed-  
318 tracing through the high Alpine landscape, it does inhibit the recognition of  
319 elements of depositional architecture that might be expressed at the km-scale  
320 (such as the development of compensating turbidite lobes or, depending on  
321 the scale, of erosional or constructional channels). In the Western Champsaur  
322 basin, adjacent to our study area (but forming a distinct turbidite system),  
323 extensive deep incision and former submarine canyons have been described  
324 (e.g. Brunt *et al.* 2007). However, there is no evidence of incision within the  
325 Eastern Champsaur Basin, beyond local scours (and associated layers of  
326 mudstone rip-up clasts) and bed-scale amalgamation, nor of channel-levee  
327 development. The 4-5 km strike section on the western face of Le Piquet  
328 (located on Fig. 4, illustrated on Fig. 5a) reveals very low-angle discordances  
329 within parts of the upper sandstone interval (A6) consistent with an expansion  
330 of stratigraphic thickness from north to south (Vinnels *et al.* 2010). Individual  
331 beds appear to thin northwards while bed-sets abut onto underlying beds –  
332 relationships we interpret as internal onlap. Collectively these relationships  
333 are plausibly interpreted as representing syn-depositional tilting within the  
334 Eastern Champsaur Basin elevating Le Piquet area relative to the Pointe des  
335 Pisses area during deposition of these upper sandstone intervals. Vinnels *et*  
336 *al.* (2010) suggest that these thickness changes reflect differential growth of  
337 an anticline in the basin floor. We infer that the intrabasinal slopes during  
338 deposition of at least this part of the Champsaur Sandstone were relatively  
339 gentle, there being no evidence for significant mass-wasting within the

340 succession. Therefore, evolving basin floor structures were largely swamped  
341 by concurrent sedimentation.

342

### 343 **The post-depositional structure of the basin**

344

345 The stratigraphic template of mappable bed-sets permits full correlation of  
346 units through the fold systems. These correlations are illustrated on our  
347 summary map (Fig. 4) and underpin two cross-sections through the Eastern  
348 Champsaur Basin (Fig. 7). The sections are necessarily simplified at the scale  
349 reproduced here, and no attempt is made here to illustrate detail of the  
350 thickness changes in the younger units (A6-A7), nor the precise geometry of  
351 polyharmonic folding deeper in the stratigraphic pile (A1).

352 The two cross-sections (Fig. 7) show a similar structural style: a train of  
353 asymmetric, west-facing folds with axial surfaces dipping towards the east.  
354 The folds detach downwards into the Blue Marls (Bürgisser & Ford 1998).  
355 Thrust faults are rare, show bed offsets of a few metres and are restricted to  
356 fold hinge zones. The attitude of the fold axial surfaces and fold interlimb  
357 angles vary through the region and, according to Bürgisser & Ford (1998),  
358 these variations reflect heterogeneous simple shear strains distributed into the  
359 footwall of the Embrunais-Ubaye thrust sheets. Folds occur in clusters,  
360 separated by segments where stratigraphic sections are only weakly  
361 deformed. Within the clusters the folds have wavelengths of 200-500m and  
362 amplitudes of around 200m. The larger folds are broadly harmonic, with axial  
363 surfaces that can be traced through the visible succession of the Champsaur  
364 Sandstone. Prime examples are those folds that cross the ridge-line on the  
365 north side of the Dourmillouse valley near the summit of Pic Felix Neff (Figs  
366 5b,c, 7). The distribution of these larger folds is consistent with a model that  
367 considers folds as ductile equivalents to imbricate thrusts (e.g. Pfiffner 1985;  
368 Butler 1992). Qualitatively, the general form of the folds appears to be  
369 influenced, at least in part, by variations in bed-thickness in the Champsaur  
370 Sandstone. The thicker, sandstone rich sequences (e.g. Fig. 2a; A6 on Figs 4  
371 and 5) appear to act as “control units”, in the sense of Price & Cosgrove  
372 (1990), determining the wavelength of the main fold sets (Fig. 8). The older,  
373 thinner-bedded units (Fig. 2; A2-A4 on Figs 4 and 5) deform poly-harmonically

374 with respect to the younger, thick-bedded stratigraphy (e.g. unit A6), and are  
375 characterised by folding at short-wavelength. Together with this layer-  
376 dependence, the location of folds in the Champsaur Sandstone may relate to  
377 variations in the properties of the basal detachment zone in the Blue Marls, as  
378 is well-known in other detachment systems (e.g. Cotton & Koyi 2000 and  
379 many others). Less effective detachment would promote folding in the  
380 overlying strata. These aspects of the deformation await more detailed  
381 investigations.

382 The cross-sections (Fig. 7) demonstrate the regional discordance  
383 between the lithostratigraphic units in the Champsaur Sandstone and the Blue  
384 Marls. As noted by Vinnels *et al.* (2010, but well-known informally), this  
385 represents a regional onlap surface, albeit modified by the tectonic  
386 detachment. It charts deformation of the basin floor during the accumulation of  
387 the Nummulitic trilogy. Vinnels *et al.* (2010) treat the onlap angle qualitatively.  
388 To consider the angle of this onlap in profile, and to establish an estimate of  
389 tectonic shortening in the basin, we present a balanced and restored section  
390 (Fig. 8) for the Dourmillouse transect (Fig. 7b).

391 The restoration assumes that the sand-rich parts of the succession  
392 deformed largely by concentric folding, without appreciable changes in bed  
393 thickness by distortional strain. To track this assumption, we chart the  
394 thicknesses of bed-sets through the cross-section. These thicknesses are  
395 reported on both the final state and restored cross-sections (thin bed-  
396 perpendicular rulings). With the assumption of concentric folding, restorations  
397 can be performed simply by restoring the length of strata, conserving the  
398 sinuous lengths measured on the final state section. This is line-length  
399 restoration, as proposed by Dahlstrom (1969). Note that in this analysis we do  
400 not consider the impact of broad arching of the enveloping surfaces of the  
401 folds with in the Champsaur Sandstone. This longer-wavelength deformation  
402 couples with the underlying basement and it is the architecture of the basin fill  
403 that we wish to reconstruct.

404 The graphical restoration (Fig. 8b) is hung from the younger  
405 stratigraphic levels (A6), the top of which forms a horizontal datum, or “target  
406 horizon”, for determining the progress of deformation. This interval, together  
407 with those below, retain bed-length and bed-thickness from the deformed

408 state. A complete restoration might be expected to yield strata that are planar  
409 and parallel – features that are not evident here. There are various possible  
410 explanations for these discrepancies. First, distortional strains, especially  
411 within the mudstone-rich intervals, have not been considered in our method,  
412 which assumes ideally concentric folding. Certainly, these units are cleaved,  
413 implying that they have experienced such distortions. There may also be  
414 errors in the drafting of the cross-section, unavoidable given variations in  
415 outcrop quality and uncertainties in the precise position of stratigraphic  
416 boundaries. Errors arising from these effects amplify away from the “target  
417 horizon”, creating significant short-wavelength relief on the base of section.  
418 These are regarded here to be restoration artefacts (selected examples  
419 identified in in Fig. 8b), as the top-basement surface is readily mapped and  
420 shows no such abrupt changes in relief where observed in the field.

421 Notwithstanding the short-wavelength artefacts, the restoration (Fig.  
422 8b) can be used to calculate the angular relationship between the depositional  
423 units in the Champsaur Sandstone and the underlying Blue Marls. The  
424 restoration implies onlap of 980m of Champsaur Sandstone over 9.4 km –  
425 implying that the average angle of the onlap slope was 6 degrees. Note that  
426 this angle would have been significantly higher at the time of deposition as we  
427 have not allowed for vertical compaction during burial.

428 Tectonic shortening for the upper stratal levels (A6) is the difference in  
429 length between deformed and undeformed state, 4 km. However, the  
430 mismatches in restored lengths of the underlying strata indicate that the  
431 section is not balanced (the bed-lengths and implied longitudinal strain are not  
432 equal in layers, so that the restoration is not complete). We resolve this by  
433 inferring significant “top-to-the-west” shear distributed through the Champsaur  
434 Sandstone, as invoked by Bürgisser & Ford (1998). Variations in layer-parallel  
435 shortening can account for the open folding of underlying sandstone units and  
436 in the substrate.

437

#### 438 **Discussion – implications for tectonostratigraphic evolution**

439

440 A palaeogeographic sketch is provided (Fig. 9), extending that of Joseph &  
441 Lomas 2004, their Fig. 5) which integrates our findings from the Eastern

442 Champsaur Basin into a semi-regional context. Following our earlier work  
443 (Vinnels *et al.* 2010), we consider its turbiditic basin fill to be a down-system  
444 continuation of the Annot sand fairway. We show the main Annot turbidite  
445 system to be derived from a southern source area and then routed along the  
446 sinuous Provencal basins defined by the underlying deformation within the  
447 broad foredeep basin (e.g. Salles *et al.* 2012). This system is distinct from a  
448 secondary turbidite system that feeds the Western Champsaur Basin.

449

#### 450 Down-system continuity of the Annot Sand Fairway

451 The total (compacted) stratal thickness of the Champsaur Sandstone in  
452 the Western Champsaur Basin approaches 1km. Stratal patterns together  
453 with onlap at the base and within the Champsaur Sandstone indicate active  
454 deformation of the underlying basement during deposition. We propose that it  
455 is this deformation that provided the basin floor structure that served to  
456 confine and guide turbidity currents, facilitating sediment transport further  
457 down-system. The high proportion of coarse sandstone compared to finer-  
458 grained fractions, coupled with palaeocurrent data (Vinnels *et al.* 2010)  
459 implies that there was significant bypass through the Eastern Champsaur  
460 Basin. The Annot sand fairway therefore continued north from Champsaur,  
461 with no ponding behind the ancestral Ecrins massif (cf. Joseph & Lomas  
462 2004; Fig. 9); turbidity currents are inferred here to have transited the region  
463 and fed down-system depocentres (Fig. 9). Certainly, the area of the Ecrins  
464 massif did not form a bathymetric high during deposition of the Annot system  
465 (cf. Ford *et al.*, 1999) – it must have been deeper than the upstream Annot  
466 fairway. Future studies should address the sedimentology of the likely  
467 correlatives of the Annot turbidites further down-system, in the strongly  
468 deformed sections along the eastern part of the Ecrins and north in the  
469 Aiguilles d'Arves (Fig. 1).

470

#### 471 Eastern confinement of the Annot system

472 For Joseph & Lomas (2004, their fig.5) and references therein) the  
473 eastern edge of the Annot sand system was confined by the Embrunais-  
474 Ubaye thrust sheets, an exotic tectonic allochthon emplaced from the internal  
475 Alpine domain onto the foreland domain (see also Ford *et al.* 2006). The

476 sheets were sufficiently thick to have buried the southern Ecrins district so  
477 that it reached temperatures of c 300°C (Bellanger *et al.* 2015). Presumably  
478 they created significant relief within the foredeep. Olistostromes were shed  
479 from this relief during thrust sheet emplacement, manifest as the Schistes à  
480 Blocs (e.g. Kerckhove 1969) which cap the turbidites in the Eastern  
481 Champsaur Basin and elsewhere in the Annot system. Ford & Lickorish  
482 (2004) suggest that it was the emplacement of the Embrunais-Ubaye thrust  
483 sheets that closed off the Annot basin system. The implication is that this was  
484 a progressive process, with the advance of thrust sheets gradually restricting  
485 the pathways available for turbidity currents. However, the relationship  
486 between the olistostomes and the Champsaur Sandstone is not consistent  
487 with this implication. Muddy debris flows, such as carried the Schistes à  
488 Blocs, would be expected to have had significant run-out distances, potentially  
489 forming obstructions within the main flow paths for the Annot turbidity  
490 currents. Deposits accumulating from axial flows in the case of the turbidity  
491 currents, and from coeval basin flank-derived olistostromes, should be  
492 interleaved. No such interleaving has been recognised by us in the Eastern  
493 Champsaur Basin, nor elsewhere in the Annot System. Therefore, we deduce  
494 that the olistostromes and Annot turbidites are not coeval – rather that the  
495 Schistes à Blocs forms a distinct, younger depositional unit that overran (and  
496 entrained) dark-shales. The corollary is that the flux of Annot turbidity currents  
497 had terminated for reasons unrelated to the emplacement of the Embrunais-  
498 Ubaye thrust sheets and that some other feature must have provided the  
499 lateral (eastern) confinement to Annot flows.

500         If lateral confinement was not provided by the tectonic allochthon of the  
501 Embrunais-Ubaye thrust sheets, then presumably it was provided by weakly  
502 inverted parts of the ancestral rifted margin of Europe adjacent to the suture  
503 of the closed Ligurian Tethys seaway. These may have formed additional  
504 basement ridges, equivalent to the Ecrins but now buried beneath Alpine  
505 thrust sheets. Alternatively, the Annot system may have abutted against  
506 deformed Briançonnais and Sub-Briançonnais units, as illustrated on Fig. 9.  
507 Both units contain sedimentary successions that are broadly time-equivalent  
508 to the Annot turbidites – the so-called “Flysch Noir”. This is a succession of  
509 pelites and thin, fine sandstones of mid-late Eocene age (e.g. Debelmas

510 1989). We tentatively suggest that the Flysch Noir represents an early part of  
511 the Annot turbidite system and that its pathway was elevated as its substrate  
512 of Brianconnais and Sub-Brianconnais units became deformed. Note that, by  
513 the early Oligocene, the eastern edge of the Brianconnais in this sector of the  
514 Alps had been incorporated into the Alpine orogen and was experiencing blue  
515 schist metamorphism, and maybe even greater burial (e.g. Michard *et al.*  
516 2004; Dumont *et al.* 2012). Regardless of the nature of the eastern retaining  
517 margin to the Annot System, there is no documented evidence that it provided  
518 detritus into the sand fairway, until the emplacement olistostromes of the  
519 Schists à Blocs from the advancing Embrunais-Ubaye thrust sheets.

520

#### 521 Structuring of the Eastern Champsaur Basin and the evolution of crustal 522 shortening

523 The onlap of Champsaur Sandstone onto a tilted succession of Blue  
524 Marls and Nummulitic Limestone indicate that the floor to the Eastern  
525 Champsaur basin was actively deforming during deposition (Vinnels *et al.*  
526 2010). Our studies confirm our earlier work and extend it – stratigraphic  
527 thickness variations across the Eastern Champsaur Basin (Fig. 6) strongly  
528 suggest that open folding continued to influence sand accumulation. However,  
529 sedimentation generally swamped the folds, with turbidites overstepping the  
530 anticline crests. These folds are cored by crystalline basement which outcrops  
531 in the Dourmillouse and Fournel valleys (Fig. 4) and it is this deep-rooting  
532 deformation style that is illustrated on Fig. 9. Basement-rooting deformation is  
533 also shown to provide the structural barrier that separated the distinct turbidite  
534 systems that characterise the Eastern and Western Champsaur Basins. Folds  
535 and thrusts in the adjacent crystalline basement now forming the Ecrins  
536 massif (located on Fig. 1a) are generally interpreted as resulting from tectonic  
537 inversion of half-graben inherited from Jurassic-aged continental rifting (e.g.  
538 Gillcrist *et al.* 1987; Dumont *et al.* 2008).

539 In Provencal basins (Fig. 9), active deformation during deposition of the  
540 Annot turbidites has been characterised as “wedge-top” (Salles *et al.* 2012  
541 and references therein). The term harks back to idealised conceptualisations  
542 of orogenic margins that are divided into a foreland basin and an adjacent  
543 thin-skinned thrust system (e.g. De Celles & Giles 1996). In these models,

544 deposition can occur in the yet-to-deform foreland basin and in small basins  
545 perched on the advancing thrust system (or wedge). However, such  
546 distinctions appear inappropriate to us for the Champsaur district as  
547 deformation of the basin floor during the turbidite deposition was thick-  
548 skinned. Concepts of thrust wedges are unlikely to be directly relevant to  
549 inversion tectonic systems such as the Ecrins or the initial deformation of the  
550 Briançonnais as illustrated on Fig. 9.

551 In Provence, two distinct phases of folding are recognised – one  
552 inferred to result from north-south contraction and traditionally related to  
553 “Pyrenean-Provençal” orogeny (early Tertiary), and a subsequent SW-  
554 directed thrust system (e.g. Siddans 1979; de Graciansky *et al.* 2011).  
555 Likewise, deformation in the crystalline basement and the structural evolution  
556 of the Ecrins basement massif has long been considered result from  
557 punctuated tectonic activity - i.e., two distinct episodes of Alpine deformation,  
558 separated by the unconformity that underlies the transgressive Nummulitic  
559 Limestone (e.g. Gidon 1979). Ford (1996) correlates these with the two  
560 tectonic episodes of Provence. Gupta & Allen (2000) argue that the southern  
561 Ecrins was structured by folds and uplifted fault blocks to modulate the  
562 Nummulitic transgression. It is a deduction that is consistent with the  
563 punctuated model for tectonics in this part of the Alps. However, it seems  
564 unlikely that initiation of a second episode of tectonic activity simply coincided  
565 with the transgression of the Nummulitic megasequence across the  
566 basement.

567 Recent radiometric dating of thrust zones in the Ecrins massif  
568 (Bellanger *et al.* 2015) implies that deformation in the basement straddled  
569 deposition of the post-Nummulitic successions of the Eastern Champsaur  
570 Basin. Using these data, Butler (2017) proposes that crustal shortening in the  
571 Ecrins and, by inference, beneath the Eastern Champsaur Basin was  
572 continuous in time. Where the instantaneous syn-orogenic surface is sub-  
573 areal, deformation is accompanied by erosion, marked by exhumation and  
574 cooling of the basement. Where the instantaneous syn-orogenic surface  
575 moves to below sea-level, deformation is accompanied by deposition. The  
576 transition between these states is marked by transgression and its  
577 diachroneity reflected by the sub-Nummulitic unconformity.

578 Dividing deformation episodes on the basis of an unconformity has a  
579 long tradition in orogenic geology (see discussion in Gray *et al.* 1997).  
580 However, is the interplay between uplift due to crustal shortening, with  
581 concomitant erosion, and subsidence due to long-wavelength orogenic  
582 loading of the lithosphere, with concomitant deposition, that defines the  
583 stratigraphic record of orogenesis. Focussing on unconformities, with their  
584 implications for missing time-sections, can create the false impression that  
585 deformation was punctuated. It is our proposal that the crust beneath the  
586 Champsaur district experienced deformation that was essentially continuous  
587 over geological time, through the late Eocene and into the Oligocene.

588 The arrival of the Embrunais-Ubaye Thrust Sheets in the late Rupelian  
589 (Dumont *et al.* 2008) marked a transition in the tectonic style to thin-skinned  
590 shearing. Deformation of the turbidites of the Eastern Champsaur Basin,  
591 dominantly by folding, happened beneath this over-riding allochthon.  
592 Shortening within the basin was approximately 4 km with differential  
593 shortening through the Champsaur Sandstone stratigraphy implying  
594 significant simple shear distributed penetratively through the multilayer  
595 (Bürgisser & Ford 1998). It is likely that thick-skinned deformation in the  
596 underlying crust continued, eventually arching the Embrunais-Ubaye thrust  
597 sheets and uplifting the adjacent Ecrins massif. The challenge facing further  
598 investigations of syn-tectonic sedimentation lies in unravelling deformed  
599 sedimentary basins to reveal their stratigraphy. These are endeavours that we  
600 hope will be encouraged by our study of the eastern Champsaur Basin.

601

## 602 **Conclusions**

603

604 Linked stratigraphic, sedimentological and structural studies in the Eastern  
605 Champsaur Basin, SE France, reveal:

606 1 – The Champsaur Sandstones of the basin represent a continuation of the  
607 late Eocene-early Oligocene Annot system. Therefore, the well-known type  
608 locations in Provence represent only a small part of this system and their  
609 causative turbidity currents continued up to, and presumably over and beyond  
610 what is now the Ecrins basement massif. There is no evidence of flow  
611 ponding in the Eastern Champsaur Basin, indeed bed character suggests

612 substantial sediment bypass continuing into more distal settings. These  
613 locations await further study.

614 2 – The continuity of sand fairways around the orogen can only arise where  
615 foredeep axial slopes and hence palaeobathymetry dipped monotonically in  
616 the direction of sediment dispersal. Therefore, during the early Oligocene, the  
617 Alpine foredeep became progressively deeper clockwise around the western  
618 Alpine arc in SE France. This implies greater tectonic subsidence, perhaps  
619 coupled with the building of depositional gradients in this direction, during the  
620 late Palaeogene. The proto-Ecrins basement massif did not lie at shallow  
621 bathymetric levels as suggested in some Alpine syntheses but, during the  
622 early Oligocene, lay under water depths significantly deeper than the  
623 Provençal sector of the foredeep.

624 3 – The floor to the foredeep basin within which the Annot turbidites were  
625 deposited was actively deforming. This has been inferred elsewhere for the  
626 type localities of the Annot system to the south of Champsaur where the  
627 foredeep megasequence rests on a gently folded substrate of Mesozoic  
628 strata. For the Eastern Champsaur, Mesozoic strata have been largely eroded  
629 so that the foredeep megasequence unconformably overlies basement. Small  
630 pockets of Triassic and Lower Jurassic strata are preserved, folded and thrust  
631 into basement beneath this unconformity, testifying to significant crustal  
632 deformation predating the deposition of the Nummulitic Limestone. We  
633 interpret the pre-Nummulitic deformation to form simply an early part of crustal  
634 shortening that progressed steadily through the late Eocene and early  
635 Oligocene.

636 4 – The Champsaur Sandstone succession is capped by the Schistes à Blocs,  
637 an olistostrome encased in dark shales, a harbinger of the advancing  
638 Embrunais-Ubaye thrust sheets. Notwithstanding the likely run-out distance of  
639 the causative debris flows ahead of the advancing thrust sheet, this  
640 olistostromal formation is not interbedded with the Champsaur Sandstone.  
641 Presumably these flows arrived on a basin floor that had largely been starved  
642 of sand supply and were not themselves the cause of this starvation.

643 5 - Deformation continued after the deposition of the sedimentary fill to the  
644 Eastern Champsaur Basin, with c. 4 km of shortening accommodated in the

645 footwall to the over-riding the tectonic allochthon represented by the  
646 Embrunais-Ubaye thrust sheets.

647

648 Our study demonstrates that even in deformed parts of the Alpine orogen it is  
649 still possible to establish detailed stratigraphies in turbidites, thereby opening  
650 opportunities to extend tectono-stratigraphic investigations elsewhere in this  
651 and other orogens. Such work may improve understanding not only of the  
652 turbidite systems and their use as analogues for modern deep-water systems  
653 but also of the tectonic relationships between orogens and their “forelands”.

654

655

656 Funding: This study was sponsored as part of the Turbidites Research Group  
657 (2007-2010), funded by Anadarko, BG, BP, ConocoPhillips, Devon Energy,  
658 Marathon, Maersk, Nexen, Petronas, StatoilHydro and Woodside. JV was  
659 supported by BHP-Billiton.

660

#### 661 Acknowledgements

662 We thank Mary Ford, Nigel Woodcock and an anonymous referee for  
663 providing insightful comments on drafts of this paper. RWHB thanks Thierry  
664 Dumont for discussions on Alpine geology and timings – although of course  
665 the authors take sole responsibility for the information and interpretations  
666 presented here.

667

#### 668 **References**

669

- 670 Bellanger, M., Augier, R., Bellahsen, N., Jolivet, L., Monié, P., Baudin, T. &  
671 Beyssax, O. 2015. Shortening of the European Dauphinois margin  
672 (Oisans Massif, Western Alps): New insights from RSCM maximum  
673 temperature estimates and  $^{40}\text{Ar}/^{39}\text{Ar}$  in situ dating. *Journal of*  
674 *Geodynamics*, **83**, 37-64.
- 675 Bouma, A. H. 1959. Some data on turbidites from the Alpes Maritimes  
676 (France). *Geologie en Mijnbouw*, **21**, 223–227.
- 677 Brunt, R.L., McCaffrey, W.D. & Butler, R.W.H. 2007. Setting and architectural  
678 elements of the Champsaur Sandstones, France. In: Nilsen, T.H., Shew,

679 R.D., Steffens, G.S. & Studlick, J.R.J. (eds), *Atlas of deep-water outcrops*  
680 (American Association of Petroleum Geologists, Studies in Geology, **56**,  
681 188-191

682 Bürgisser, J. & Ford, M. 1998. Overthrust shear deformation of a foreland  
683 basin: structural studies south-east of the Pelvoux massif, SE France.  
684 *Journal of Structural Geology*, **20**, 1455-1475.

685 Butler, R.W.H. 1992. Evolution of Alpine fold-thrust complexes: a linked  
686 kinematic approach. In: Mitra, S. & Fisher, G. (eds) *Structural geology of fold*  
687 *and thrust belts*. Johns Hopkins University Press, Baltimore, 29-44.

688 Butler, R.W.H. 2017. Basement-cover tectonics, structural inheritance and  
689 deformation migration in the outer parts of orogenic belts: a view from the  
690 western Alps. In: Law, R.D., Thigpen, R., Merschat, A., Stowell, H. &  
691 Bailey, C. (eds) *Linkages and Feedbacks in Orogenic Process: A volume*  
692 *in Honor of Robert D Hatcher Jr.* Memoir of the Geological Society of  
693 America, **213**, 55-74.

694 Butler, R.W.H. & McCaffrey, W.D. 2004. Nature of thrust zones in deep water  
695 sand-shale sequences: outcrop examples from the Champsaur sandstones  
696 of SE France. *Marine and Petroleum Geology*, **21**, 911-921.

697 Cotton, J.T. & Koyi, H.A. 2000. Modeling of thrust fronts above ductile and  
698 frictional detachments: Application to structures in the Salt Range and Potwar  
699 Plateau, Pakistan. *Geological Society of America Bulletin*, **112**, 351-363.

700 Cunha, R.S., Tinterri, R. & Magalhaes, P.M. 2017. Annot Sandstone in the Peïra  
701 Cava basin: An example of an asymmetric facies distribution in a confined  
702 turbidite system (SE France). *Marine and Petroleum Geology*, **87**, 60-79.

703 Dahlstrom, C.D.A. 1969. Balanced cross-sections. *Canadian Journal of Earth*  
704 *Sciences*, **6**, 743-757.

705 De Celles, P.G. & Giles, K.A. 1996. Foreland basin systems. *Basin Research*, **8**,  
706 105-123.

707 Debelmas, J. 1989. On some key features of the evolution of the western Alps.  
708 In: Sengor, A.M.C. (ed). *Tectonic Evolution of the Tethyan Region*. Springer,  
709 Dordrecht. 23-42.

710 Debelmas, J., Durozoy, G., Kerckhove, C. Monjuvent, G., Mouterde, R. &  
711 Pêcher, A. 1980. *Carte géologique de la France à 1:50,000: feuille*  
712 *Orcières*. Bureau de Recherches Géologiques et Minières, Orleans.

- 713 Dumont, T., Champagnac, J-D., Crouzet, C. & Rochat, P. 2008. Multistage  
714 shortening in the Dauphiné zone (French Alps): the record of Alpine  
715 collision and implications for pre-Alpine restoration. *Swiss Journal of*  
716 *Geosciences*, **101**, S89-S110.
- 717 Dumont, T., Schwartz, S., Guillot, S., Simon-Labric, T., Tricart, P. & Jourdan,  
718 S. 2012. Structural and sedimentary records of the Oligocene revolution in  
719 the Western Alpine arc. *Journal of Geodynamics*, **56-57**, 18-38.
- 720 Etienne, S., Mulder, T., Bez, M., Desaubliaux, G., Kwasniewski, A., Parize,  
721 O., Dujoncquoy, E. & Salles, T. 2012. Multiple scale characterization of  
722 sand-rich distal lobe deposit variability: examples from the Annot  
723 Sandstones Formation, Eocene–Oligocene, SE France. *Sedimentary*  
724 *Geology*, **273**, 1-18.
- 725 Ford, M. 1996. Kinematics and geometry of early Alpine, basement-involved  
726 folds, SW Pelvoux Massif, SE France. *Eclogae Geologicae Helvetica*, **89**,  
727 269-295.
- 728 Ford, M. & Lickorish, W.H. 2004. Foreland basin evolution around the western  
729 Alpine Arc. In: Joseph, P. & Lomas, S. (eds) *Deep-water sedimentation in*  
730 *the Alpine basin of SE France*. Special Publications of the Geological  
731 Society, London, **221**, 39-63.
- 732 Ford, M., Lickorish, W.H. & Kuszniir, N.J. 1999. Tertiary foreland sedimentation  
733 in the Southern Subalpine Chains, SE France: a geodynamic appraisal.  
734 *Basin Research*, **11**, 315-336.
- 735 Ford, M., Duchêne, S., Gasquet, D. & Vanderhaeghe, O. 2006. Two-phase  
736 orogenic convergence in the external and internal SW Alps. *Journal of the*  
737 *Geological Society*, **163**, 815-826.
- 738 Gidon, M. 1979. Le rôle des étapes successives de déformation dans la  
739 tectonique alpine du massif du Pelvoux (Alpes occidentales). *Comptes*  
740 *rendus de l'Académie des Sciences, Paris*, **288**, 803-806.
- 741 Gillcrist, R., Coward, M.P., & Mugnier, J-L. 1987. Structural inversion: examples  
742 from the Alpine foreland and French Alps. *Geodinamica Acta*, **1**, 5-34.
- 743 de Graciansky, P-C., Roberts, D.G., and Tricart, P., 2011, *The Western Alps,*  
744 *from rift to passive margin to orogenic belt*. Developments in Earth Surface  
745 Processes, Elsevier, **14**, pp. 1-398.

- 746 Gray, D.R., Foster, D.A. & Bucher, M. 1997. Recognition and definition of  
747 orogenic events in the Lachlan Fold Belt. *Australian Journal of Earth*  
748 *Sciences*, **44**, 489-501.
- 749 Gupta, S. 1997. Tectonic control on paleovalley incision at the distal margin of  
750 the early Tertiary Alpine foreland basin, southeastern France. *Journal of*  
751 *Sedimentary Research*, **67**, 1030-1043.
- 752 Gupta, S. & Allen, P.A. 2000. Implications of foreland paleotopography for  
753 stratigraphic development in the Eocene distal Alpine foreland basin.  
754 *Geological Society of America Bulletin*, **112**, 515-530.
- 755 Joseph, P. & Lomas, S.A. 2004. Deep-water sedimentation in the Alpine  
756 foreland basin of SE France: New perspectives on the Grès d'Annot and  
757 related systems — An introduction. In: Joseph, P. & Lomas, S. (eds).  
758 *Deep-water sedimentation in the Alpine basin of SE France*. Special  
759 Publications of the Geological Society, London, **221**, 1-16.
- 760 Kerckhove, C. 1969. La 'zone du flysch' dans les nappes de l'Embrunais-  
761 Ubaye (Alpes Occidentales). *Géologie Alpine, Grenoble*, **45**, 5-204.
- 762 Kerckhove, C., Cochonat, P. & Debelmas, J. 1978. Tectonique du  
763 soubassement parautochtone des nappes de l'Embrunais-Ubaye sur leur  
764 bordure occidentale, du Drac au Verdon. *Géologie Alpine, Grenoble*, **54**,  
765 67-82.
- 766 Michard, A., Avigad, D., Goffé, B. and Chopin, C., 2004. The high-pressure  
767 metamorphic front of the south Western Alps (Ubaye-Maira transect,  
768 France, Italy). *Schweizerische Mineralogische und Petrographische*  
769 *Mitteilungen*, **84**, 215-235.
- 770 Mulder, T., Callec, Y., Parize, O., Joseph, P., Schneider, J.L., Robin, C.,  
771 Dujoncquoy, E., Salles, T., Allard, J., Bonnel, C. & Ducassou, E. 2010.  
772 High-resolution analysis of submarine lobes deposits: Seismic-scale  
773 outcrops of the Lauzanier area (SE Alps, France). *Sedimentary Geology*,  
774 **229**, 160-191.
- 775 Pffifner, O.A. 1985. Displacements along thrust faults. *Eclogae Geologicae*  
776 *Helvetica*, **78**, 313-333.
- 777 Price, N.J. & Cosgrove, J.W. 1990. *Analysis of Geological Structures*.  
778 Cambridge University Press, Cambridge.

- 779 Puigdefabregas, C., Gjelberg, J. & Vaksdal, M. 2004. The Gres d'Annot in the  
780 Annot syncline: outer basin-margin onlap and associated soft-sediment  
781 deformation. In: Joseph, P. and Lomas, S. A. (eds). *Deep-Water*  
782 *Sedimentation in the Alpine Basin of SE France: New Perspectives on the*  
783 *Gres D'Annot and Related Systems*. Geological Society of London Special  
784 Publication **221**, 367-388.
- 785 Ramsay, J.G. 1974. Development of chevron folds. *Geological Society of*  
786 *America Bulletin*, **85**, 1741-1754.
- 787 Salles, L., Ford, M. & Joseph, P. 2014. Characteristics of axially-sourced  
788 turbidite sedimentation on an active wedge-top basin (Annot Sandstone, SE  
789 France): *Marine and Petroleum Geology*, **56**, 305-323.
- 790 Schmid, S.M., Fügenschuh, B., Kissling, E. & Schuster, R. 2004. Tectonic map  
791 and overall architecture of the Alpine orogeny. *Eclogae geologicae*  
792 *Helveticae*, **97**, 93-117.
- 793 Siddans, A.W.B. 1979. Arcuate fold and thrust patterns in the Subalpine chains  
794 of Southeast France. *Journal of Structural Geology*, **1**, 117-126.
- 795 Sinclair, H.D. 1997. Tectonostratigraphic model for underfilled peripheral  
796 foreland basins: An Alpine perspective: *Geological Society of America*  
797 *Bulletin*, **109**, 324-346.
- 798 Smith, R. 2004. Silled sub-basins to connected tortuous corridors: sediment  
799 distribution systems on topographically complex sub-aqueous slopes. In:  
800 Lomas, S.A. & Joseph, P. (eds) *Confined turbidite systems*. Special  
801 Publications of the Geological Society, London, **222**, 23-43.
- 802 Stanbrook, D.A. and Clark, J.D., 2004. The Marnes Brunès Inférieures in the  
803 Grand Coyer remnant: characteristics, structure and relationship to the Grès  
804 d'Annot. . In: Joseph, P. & Lomas, S. (eds). *Deep-water sedimentation in*  
805 *the Alpine basin of SE France*. Special Publications of the Geological  
806 Society, London, **221**, 285-300.
- 807 Stanley, D. J. & Bouma, A. H. 1964. Methodology and paleogeographic  
808 interpretation of Flysch formations: a summary of studies in the Maritime  
809 Alps. In: A. H. Bouma and A. Brouwer (eds.), *Turbidites*. Developments in  
810 Sedimentology **3**. Elsevier, Amsterdam, 34–64.

811 Tricart, P. 1981. Les marqueurs des mouvements verticaux du socle aux  
812 abords du massif du Pelvoux (Alpes occidentales). *Revue de géologie*  
813 *dynamique et de géographie physique Paris*, **23**, 289-300.

814 Vinnels, J.S., Butler, R.W.H., McCaffrey, W.D. & Lickorish, W.H. 2010.  
815 Sediment distribution and architecture around a bathymetrically complex  
816 basin: Eastern Champsaur Basin, SE France. *Journal of Sedimentary*  
817 *Research*, **80**, 216-235.

818

819

820 **Figures**

821

822 Fig. 1. a) Location map of the Eastern Champsaur Basin (boxed area is Fig 4)  
823 in the context of the western Alps (modified after Schmid *et al.* 2004). b) the  
824 continuity of the Annot sand system in SE France, in the context of other  
825 Eocene-Oligocene turbidite units in the external Western Alps (modified after  
826 Joseph & Lomas 2004). The grey area represents thrust sheets, including the  
827 Embrunais-Ubaye and lateral equivalents that have over-riden the Annot  
828 sand system, which are shown as inferred at depth. SF – Selle Fault.

829

830 Fig. 2. Idealised Nummulitic stratigraphy (modified after Joseph & Lomas  
831 2004)

832

833 Fig. 3. Fold geometries, all of which face west, in various parts of the  
834 Champsaur Sandstone (see Fig. 4 for place names). a) looking NNW onto Pic  
835 Felix Neff, to illustrate the broadly harmonic and relatively long-wavelength  
836 folding of the upper stratigraphic packages (unit A6 on Figs 4 and 6). b)  
837 looking SE onto polyharmonic folding in sandstone units (unit A3 on Figs 4  
838 and 6) directly above the regional detachment zone of the Blue Marls. The  
839 visible cliff height is 100m. c) looking SSW onto chevron-style folding in the  
840 lower units of the thinly bedded Champsaur sandstone (Unit A1 on Figs 4 and  
841 6). The bottom of the image is at an altitude of c 2200m, the skyline reaches  
842 up to 3085m.

843

844 Fig. 4. Geological map of the Eastern Champsaur Basin showing the  
845 distribution and structure of the mappable units within the Champsaur  
846 Sandstone as defined by this study (Fig. 4). The location of lines of section in  
847 Fig 5a, b are annotated (X-X' and Y-Y' respectively), along with logged  
848 stratigraphic sections (circled numbers, 1-9; Fig. 6) and field photographs of  
849 Fig. 5.

850

851 Fig. 5. Selected key panoramas that serve to trace the bed-set stratigraphy of  
852 the Champsaur Sandstone, with varying accessibility. The viewpoints are  
853 shown on Fig. 4. Stratigraphic boundaries are shown as yellow lines while

854 fold axial traces are shown by red dashed lines. a) illustrates the west face of  
855 Le Piquet, a strike-section, together with the location and content of two  
856 sedimentary logs (1 and 2 on Fig. 4) that characterise the upper part of the  
857 Champsaur Sandstone. b) Is an oblique view of the north side of the  
858 Dourmillouse Valley, which provides a dip-section through the fold belt. This  
859 hillside is partly accessible permitting ground-truthing of units and completion  
860 of measured sections. c) illustrates part of the steep south side of the Fournel  
861 Valley. This provides exceptional exposures for the stratigraphy and folds but  
862 has limited accessibility.

863

864 Fig. 6. A selection of lithostratigraphic logs with defined mappable units (A1 –  
865 A6-A7), correlated through the basin. The locations are shown on Fig. 4.  
866 These sections were constructed with direct bed-measurement and field  
867 observations – their location being restricted by accessibility in the terrain.

868

869 Fig. 7. Cross-sections through the Eastern Champsaur Basin. The lines of  
870 section are shown in Fig. 2. a) the northern basin; b) the Dourmillouse  
871 transect.

872


873 Fig. 8. A line-length restoration of the Dourmillouse cross-section through the  
874 basin. The thicknesses of bed-sets are shown (intra-formation rulings), and  
875 retained between the final state section (a) and its restored version (b).  
876 Selected bed-set levels are illustrated (in colour) together with the top  
877 basement (in red).

878



879 Fig. 9. A schematic representation of the extent of the Annot turbidite system,  
880 connecting its well-studied proximal areas in Provence to the almost un-  
881 studied, more distal regions that include the Aiguilles d'Arves basin, via the  
882 Eastern Champsaur Basin described here. The diagram is inspired by an  
883 equivalent perspective provided by Joseph & Lomas (2004). Evidence for  
884 deformation and its timing beneath the sand fairway within the Ecrins and  
885 beneath the Aiguilles d'Arves sector is discussed by Butler (2017).

886


887




a





b









b)



c)

