



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/164186/>

Version: Accepted Version

Article:

Wailoo, Allan, Hernandez Alava, Monica, Manca, Andrea et al. (2017) Use of Mapping to Estimate Utility Values from Non-Preference-Based Outcome Measures for Cost per QALY Economic Analysis: Good Research Practices Task Force. *Value in Health*. pp. 18-27.
ISSN: 1524-4733

<https://doi.org/10.1016/j.jval.2016.11.006>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 **Use of Mapping to Estimate Utility Values from Non-Preference-Based**
2 **Outcome Measures for Cost per QALY Economic Analysis: Good**
3 **Research Practices Task Force**

4 Allan Wailoo¹, Monica Hernandez Alava¹, Andrea Manca², Aurelio Mejia³, Joshua Ray⁴,
5 Bruce Crawford⁵, Marc Botteman⁶, Jan Busschbach⁷.

6 ¹ *School of Health and Related Research, University of Sheffield, UK*

7 ² *Centre for Health Economics, University of York, York, England, UK*

8 ³ *Instituto de Evaluación Tecnológica en Salud, Bogotá, Colombia.*

9 ⁴ *F. Hoffmann-La Roche, Basel, Switzerland*

10 ⁵ *IMS health, Tokyo, Japan.*

11 ⁶ *Pharmerit International, Bethesda, US*

12 ⁷ *Section Medical Psychology, Erasmus Medical Center, Rotterdam, The Netherlands.*

13 Draft 19th May 2016

14 Total words: Current 8189

15 **Abstract:**

16 **1. BACKGROUND. THE ISPOR TASK FORCE PROCESS**

17 ISPOR to insert details here

18 **2. INTRODUCTION**

19 The assessment of health-related quality of life is critical in the evaluation of health care
20 technologies and services, and in regulatory and reimbursement decisions. “Preference-based
21 measures” (PBMs) play a central role in these evaluations. They allow patients to describe
22 the impact of ill health and have an associated “utility” score (or tariff) for each of those
23 health state descriptions where a value of 1 represents full health, 0 represents the value of
24 dead, and negative values (if defined by the PBM) represent states worse than death. These
25 utility scores can then be used for the calculation of Quality Adjusted Life Years (QALYs),
26 which are an outcome metric for health benefit used in many health economic evaluations.

27 The most widely-used PBMs are *generic*: applicable to a wide range of diseases, patients and
28 interventions. Examples include the EQ-5D¹, SF-6D², a derivative of the SF-36 instrument,
29 and the Health Utilities Index (HUI)³. Many national guidelines for economic evaluation
30 suggest or require the use of these generic instruments, such as England and Wales⁴, Spain⁵,
31 France⁶, Thailand, Finland, Sweden, Poland, New Zealand, Canada, Colombia and The
32 Netherlands. Some recommend the use of a particular instrument, usually the EQ-5D⁷.

33 In many situations, clinical studies do not include a PBM. Often they will include one or
34 more of the many patient-reported outcome measures (PROMs) which are not full PBMs
35 because they do not have an associated, preference-based scoring system. Thus they do not
36 permit construction of a QALY measure. Studies typically will also include physical
37 outcomes (not patient-reported) which are measured “objectively”, that is, without the
38 interpretation of or report by the patient. In the absence of a PBM outcome, researchers will
39 need to derive the “missing” PBM in order to estimate QALYs from these studies. In these
40 circumstances the question is whether it is possible, and how, to predict the value that a PBM
41 would have taken had this been collected, given what we know about the observed clinical
42 outcome(s) and allowing for the mediating effect of the individual characteristics of study
43 participants. “Mapping” attempts to answer this question and, in so doing, bridges the gap
44 that often exists between available evidence on the effect of a health technology in one metric
45 and the requirement for decision makers to express it in a different one (QALYs). It can also
46 be used to provide a means of converting outcomes in one PBM to a different PBM.

47 “Mapping” makes use of another dataset, which may be observational rather than
48 experimental. This dataset must have the same outcomes that are measured in the relevant
49 clinical study/studies, and the patients’ responses to a standard PBM instrument. This
50 external dataset is used to estimate a statistical relationship between the two types of outcome
51 measure. Combining the estimated statistical relationship together with the outcome data
52 from the trial allows an estimate of the effect of the treatment in health utility terms and
53 subsequently may be used to calculate QALYs. The practice of fitting a statistical model to
54 health utility data has variously been referred to as “mapping,” “cross-walking” and

55 “transfer to utility”⁸. “Mapping” has entered into common usage so is used throughout this
56 report.

57 In the context of economic evaluation, the evidence gap which gives rise to the need for
58 mapping is commonly encountered. For example, Kearns *et al* (2013)⁹ reviewed 79 recent
59 NICE Technology Appraisals and found that mapping models were used in almost a quarter
60 of cases. These included mapping from the Psoriasis Area Severity Index (PASI) in patients
61 with psoriasis, from the Functional Assessment of Cancer Therapy – General (FACT-G) in
62 patients with cervical cancer, and from the Patient Assessment of Constipation – Symptoms
63 (PAC-SYM) and Patient Assessment of Constipation – Quality of Life (PAC-QOL) in
64 women with chronic constipation, *inter alia*. The need for mapping may arise because of a
65 failure to include a PBM in the relevant clinical studies (as described above), or because
66 those studies are not sufficient alone to provide the utility information to estimate cost-
67 effectiveness. There could be a requirement for extrapolation beyond the range of health
68 states observed in clinical studies or a requirement to synthesise evidence from several
69 clinical studies, not all of which include evidence on PBMs. Thus, mapping is an issue both
70 for economic evaluation alongside trial data analysis without PBMs as well as for many
71 economic modelling studies. And because studies that have been conducted historically will
72 remain part of the evidence base as comparators for the evaluation of new technologies,
73 mapping is likely to remain a requirement for some time, even when good practices for utility
74 estimation are followed in contemporary clinical studies¹⁰.

75 The current practice of mapping includes substantial variation in methods which are known
76 to lead to differences in cost-effectiveness estimates^{11,12}. The purpose of this Task Force
77 report is to set out Good Research Practices that are relevant for the conduct of mapping
78 studies for use in all types of QALY-based economic evaluation. The recommendations also
79 have broader relevance to all situations where analysts wish to estimate preference-based
80 outcomes as a function of any other variables, for example, where utilities are used as
81 measures of provider performance¹³. Recommendations cover all areas of mapping practice:
82 the selection of datasets for the mapping estimation, model selection and performance
83 assessment, reporting standards, and the use of results including the appropriate reflection of
84 variability and uncertainty. Such recommendations are critical in the face of inconsistent
85 current practices, substantial variation in results between approaches and the risk of bias in
86 several methods. Whilst other recommendations have been made^{14,15}, this document is unique
87 because it takes an international perspective, is comprehensive in its coverage of the aspects
88 of mapping practice, and reflects the current state of the art.

89 **3. PRE-MODELLING CONSIDERATIONS**

90 Prior to undertaking a statistical analysis for the purpose of mapping, the analyst must
91 consider a number of different factors relating to the proposed and potential uses of the
92 mapping itself. These uses create requirements for the dataset(s) in which the statistical
93 analyses will be undertaken and tested.

94 Mapping is almost always undertaken with some pre-defined purpose and in many of those
95 cases this is to inform a specific cost-effectiveness analysis (CEA). Clear understanding of
96 the evidence gap to be addressed requires an understanding of relevant existing utility
97 evidence, the requirements of the decision-making body that will assess the results of the
98 analysis and the CEA in which the results are to be used. These factors help to inform the
99 analytical choices which ensure unbiased estimates in the cost-effectiveness study. There will
100 be requirements to appropriately reflect uncertainty and, additionally in some situations, the
101 variability of estimates (for example, if simulating individual patients in a cost-effectiveness
102 model).

103 The needs of the CEA will help guide the analyst's choice of methods and datasets that can
104 be expected to perform appropriately for these specific needs. Where the analysis is to be
105 used to populate a decision analytic model, one needs to consider what health states are
106 reflected in that model – how are they defined and how do those definitions relate to both the
107 clinical outcome measure or measures of relevance and the target PBM? If there is little
108 overlap between the clinical outcomes and the PBM then mapping is unlikely to be
109 successful. A descriptive comparison of the content of the different outcome measures,
110 including the suggested PBM, is a useful starting point. This will highlight the specific facets
111 of health each instrument measures. It is not a requirement for the PBM and clinical
112 outcomes to address the same symptoms or functional (dis)abilities in order for mapping to
113 be an appropriate approach but they do need to measure the same underlying concepts.

114 Many models, such as transition state models, will typically define a relatively small number
115 of discrete health states. Other situations may require a combination of health states that can
116 be derived in part from a mapping study and in part from other evidence. For example, the
117 model may differentiate health states based on a disease outcome measure and the therapy
118 patients are receiving, or the adverse events they experience, or their comorbidities. Mapping
119 and other existing evidence can provide a range of options for addressing these evidence
120 gaps.

121 Mapping outcomes to the utilities of a PBM is usually done with regression analyses. At one
122 end of the spectrum, there are rare occasions where regression models can be avoided entirely
123 simply by taking the mean and variance of the utility value for patients with the relevant
124 health criteria. This simple approach is entirely legitimate if there is a single summary
125 measure of disease to explain utility with no additional covariates that are considered
126 important and there are sufficient observations of patients within each category. However, it
127 should be noted that this may limit the generalisability of the mapping to other CEAs where
128 these conditions do not hold.

129 Regression type analyses do become a requirement once additional covariate and/or
130 extrapolation outside the range of the observed data are required, as is often the case. This
131 might be because there are multiple disease specific outcome measures that reflect different
132 dimensions of disease that collectively are used to estimate health utility. Or it could be
133 because the analyst wishes to incorporate the effect of socio-demographics on health utility.
134 For instance, age is likely to be a relevant variable in many situations as it will be related to

135 health and quality of life. Another reason to consider regression models for mapping is the
136 possibility of the need to extrapolate beyond the range of disease severity observed in the
137 data. Whilst extrapolation beyond the range of the data is best avoided in any situation, this is
138 not always feasible. Mapping studies are frequently based on datasets that do not include the
139 full range of patient disease severity, particularly when these datasets are from randomised
140 controlled trials with exclusion criteria for comorbidities and other aspects of severity. This
141 contrasts with the needs of decision models, particularly those for patients with chronic
142 conditions, which may model patients' lifetimes and thus span the entire feasible spectrum of
143 disease.

144 It is well established that some methods for such regression analyses exhibit bias, the extent
145 of which is in part dependent on the target utility measure. More details are provided in
146 section 4, but it can be noted at this point that bias is typically greatest at the extremes of
147 disease severity – for patients in severe ill- health these approaches overestimate their true
148 health utility and for those in good health they underestimate health utility¹⁶. With this in
149 mind the analyst must assess the requirements of the CEA. For instance, what is the range of
150 disease to be addressed by the decision model? This judgment should not only be made
151 against the characteristics of candidate patients at the point in the patient pathway where the
152 technology of interest is being assessed (model baseline), but should be informed by the
153 range of future health states to be covered in the model. Since this may cover a long term
154 extrapolation encompassing patients experiencing diverse pathways including disease
155 progression, therapy response and disease remission, a very wide range of disease severity
156 can sometimes be covered.

157 Similar considerations influence the requirements for datasets in which the mapping function
158 is to be estimated. Additional requirements are that, obviously, candidate datasets must come
159 from studies of individuals completing both the relevant clinical outcome measure(s) and the
160 target PBM simultaneously. There is no reason why randomised studies would be more
161 desirable for mapping studies. Indeed, as alluded to above, randomised studies often have less
162 diverse patients than other study types in terms of disease severity because of strict inclusion
163 and exclusion criteria and limited follow up. Observational studies may be more likely to be
164 drawn from representative patient groups, have larger sample sizes and can be conducted at
165 relatively low cost. Where there is more than one candidate dataset then consideration should
166 be given to the additional data fields the different studies include which may facilitate more
167 precise estimates of the target PBM as well as the sample size, generalisability of the patient
168 population and any potential biases in the study designs. However, this needs to be balanced
169 with the use of those values in subsequent CEAs. The availability of information on
170 respondents' age, for example, is likely to improve model fit and ought to be incorporated
171 into a CEA. Datasets may be combined where common covariates exist and differences
172 between patients and study designs are not expected to influence the relationship between
173 covariates and PBM.

174 Uncertainty in the estimates should be minimized. This is facilitated in part by the use of
175 datasets with larger numbers of observations and by avoiding extrapolation beyond the range
176 of the data when feasible. Matching the range of disease severity in the dataset with the

177 population of the CEA is important, but the range of other patient characteristics used as
178 covariates in the mapping model are also relevant here.

179 Finally, the analyst needs to be aware of any potential biases in the dataset. Biases in this
180 situation refers to those factors which influence a patient’s reported health utility other than
181 through an impact on the clinical outcome measure(s) used as explanatory variables. For
182 instance, in some situations the types of therapies patients are receiving may exert some bias,
183 for example, where those therapies are associated with adverse events unrelated to the clinical
184 outcome being measured in the mapping dataset.

185 Summary of pre-modelling recommendations

1. Consider the use or potential uses of the mapping:
 - a. Is it for use in a cohort decision model, patient level model or trial-based cost-effectiveness analysis?
 - b. What are the health states that require utility estimates from the mapping and how do they relate to the PBM?
 - c. What is the range of disease severity for which utility values are required?
2. Provide a descriptive account of the clinical explanatory variable, the dependent PBM and the extent to which they overlap.
3. Assess if a regression-based mapping is required.
 - a. How many health states require estimates of utility?
 - b. Are there additional covariates of importance?
 - c. Are there sufficient observations within each category?
4. Identify if more than one dataset is potentially available for estimation. Compare the characteristics of candidate datasets.
5. To what extent does the distribution of patient characteristics in the sample datasets reflect those that are the subject of the cost effectiveness analysis? In particular, are all extremes of disease

186 **4. MODELLING AND DATA ANALYSIS**


187 *Selection of the statistical model*

188 Utility measures tend to exhibit a number of non-normal distributional characteristics. These
189 measures can be considered a type of limited dependent variable at both the top and bottom
190 of their ranges: by definition a value of 1 is the maximum value that can be achieved and is
191 considered equal to “full health”. There is a lower limit which varies by instrument,
192 sometimes referred to as the “pits” state. Note that these limits in utilities are not the same as
193 “censoring”.

194 Additional aspects of the distribution of utilities that influence the statistical model choice are
195 the presence of large spikes in the distribution (typically at the “full health” upper bound),
196 skewness, multimodality and gaps in the range of feasible values. Figure 1 displays examples
197 of the distribution of EQ5D-3L from a range of different disease areas. The extent to which
198 these features are present varies according to the instrument and scoring algorithm of the
199 PBM that is the target for the mapping study, and the nature of the patient group. The
200 presence of any of these features makes the application of simple statistical regression

201 methods challenging and this is compounded when several of these features are
 202 simultaneously present.

203 Figure 1. The distribution of EQ5D-3L across different disease areas



204 There is considerable evidence that these distributional features result in systematic bias
 205 when linear regression methods are used to analyze the EQ-5D-3L instrument, the most
 206 commonly studied patient reported outcome in the mapping literature^{17,18,19}. Similar findings
 207 have been shown to apply to models like the Tobit¹⁹ (designed to deal with limited dependent
 208 variables), two-part models²⁰ (which attempt to address the mass of observations seen at full
 209 health) and censored least absolute deviations models^{21,22}. A common finding in those reports
 210 is that expected health utility associated with mild health states is underestimated whilst

211 utility for more severe health states is overestimated. When mapping studies with these biases
212 are used in economic evaluations, clinically effective therapies appear less cost-effective than
213 they truly are. Studies have shown that the magnitude of this bias is not trivial^{11,12}.

214 Recent work compares the performance of different statistical methods for mapping. One set
215 of methods estimate the summary utility score directly. Amongst these direct methods, there
216 is some empirical evidence to support the performance of two approaches: the limited
217 dependent variable mixture model approach^{19,12} and the beta-based regression approaches^{17,23}.

218 Alternatively, indirect methods estimate utilities as part of a two-stage procedure²⁴. These
219 methods have also demonstrated improvements over standard methods in some
220 settings^{12,25,26,27}. In the first stage, a so-called “response mapping” model uses a series of
221 (either dependent or independent) separate regression functions to estimate the level on each
222 separate domain of the descriptive system of the target PBM. Models suitable for ordered
223 categorical data should be used for this first stage and the correlation between dimension
224 responses incorporated²⁷. It is then straightforward to calculate the expected utility score as
225 stage 2 of the procedure based on the probabilities assigned to each of the health states in the
226 descriptive system and their associated utilities. This separation allows the analyst to apply
227 any utility tariff to the models estimated in stage 1, according to their requirements. However,
228 it should be noted that the appropriateness of the model and its fit is specific to the tariff in
229 which it has been tested. Furthermore, response mapping models require sufficient
230 observations in each of the levels of the descriptive system. Without this, the model(s) cannot
231 be estimated.

232 We do not advocate any specific set of methods as the performance of different methods will
233 vary according to the characteristics of the target utility measure, the disease and patient
234 population in question, the nature of the explanatory clinical variables and the form of
235 intended use in the CEA. We therefore suggest that it is wise to use a model type for which
236 there is existing empirical evidence of good performance, and that respects the key features
237 of the target utility measure, particularly the limited range of feasible utility values that can
238 be taken in order to avoid problems in implementing results in a cost-effectiveness model.

239 Obviously, mapping does require analysts to adhere to good practice for statistical analysis in
240 general. Below, we highlight some aspects of good practice that relate in particular to
241 mapping. For instance, a plot of the distribution of the target utility measure provides a
242 starting point for considering potentially appropriate modelling methods for direct analysis of
243 the utility index. Analysts should use models that have theoretical plausibility, whose key
244 assumptions hold, and that have a body of existing empirical evidence supporting their
245 validity in the mapping literature. The use of models that do not meet these criteria requires
246 additional justification and the results should be subject to additional scrutiny. This additional
247 justification can be in the form of evidence that demonstrates that the mapping does not
248 suffer from bias in the particular application, or that the nature of that bias is not an issue
249 given the use of the mapping in CEA. For example, if the analyst intends to populate a cohort
250 decision model where only a small number of health states are defined and these health states
251 are not located at the extremes of poor/good health, then bias from the mapping may have a

252 negligible effect on estimated cost-effectiveness. However, it is difficult to assess the impact
253 of any potential bias *a priori*.

254 In most situations it will be extremely important to utilize mapping methods that meet the
255 criteria set out above. This is because the extent and impact of biased estimates on cost-
256 effectiveness will be significant and predictions outside the feasible utility range could be
257 made. For example, model-based CEAs where health states are at the extremes of disease
258 severity, individual patient simulation models, or analyses based on individual level data such
259 as CEAs conducted alongside a single clinical trial will all be at risk of substantially biased
260 cost-effectiveness estimates if inappropriate mapping methods are applied.

261 We note that some model types will require iterative estimation methods. It is imperative that
262 the analyst ensures proper convergence of the estimation algorithm, whether undertaken in a
263 classical²⁸ or Bayesian²⁹ framework.

264 It is also typical for candidate datasets to comprise multiple observations from the same
265 individuals over time. In general one should seek to make use of all observations. Multilevel
266 models can be used to reflect the correlations between these observations. At a minimum,
267 clustered standard errors should be calculated. Where there are reasons to believe that there
268 has been a break in the relationship between the covariates and the PBM then separate
269 models should be estimated and the stability of the parameters tested.

270 *The selection of covariates*

271 In most situations the dataset in which the mapping is to be performed will contain
272 information on a range of potential explanatory variables. The primary decision for the
273 analyst concerns the choice of non-preference-based measure that will serve as the key link
274 between the clinical effectiveness data and the preference-based one. In many situations the
275 non-preference-based measure will be obvious because it will be the primary outcome
276 measure used in clinical studies, or the sole quality of life instrument amongst the secondary
277 outcomes. However, often those measures are formed of individual questions, which in turn
278 can be reported either as dimension scores or a single summary score. Typically, there will be
279 greater explanatory power from a regression model that uses disaggregated information from
280 an outcome measure as explanatory variables. However, not only does this increase the
281 number of explanatory variables but it may not provide the link to clinical evidence in a form
282 that is widely usable (see, for example, Longworth et al³⁰ who modelled the 36 individual
283 question responses to the EORTC instrument). This can be illustrated using the example of
284 Rheumatoid Arthritis (RA). Typically, cost-effectiveness studies make use of the Health
285 Assessment Questionnaire (HAQ) mapped to a preference-based instrument³¹. The HAQ is a
286 summary score of functional impairment that ranges from 0-3 derived from 8 sub-sections
287 each of which is comprised of 2 or 3 individual questions. Whilst the analyst may find a
288 better performing model if using the individual item or dimension scores as explanatory
289 variables, as opposed to the single 0-3 summary score, this should not be the sole criteria for
290 covariate choice (see, for example, Bansback et al³²). Where the mapping function is to be
291 used to estimate health utility from individual questions or component scores, as might be the

292 case in an economic evaluation conducted alongside a clinical trial, such an approach will be
293 useful. However, decision models that synthesize data from several clinical studies will
294 typically rely on the published results which will report only the summary score.

295 In other settings the analyst may have a choice of one or more disease specific outcomes. In
296 Ankylosing Spondylitis (AS) for example, clinical studies typically report both BASDAI and
297 BASFI outcomes measures of disease activity and functional impairment. The conceptual
298 overlap with a preference-based instrument may be improved by the inclusion of multiple
299 instruments and, hence, model fit.

300 Covariates can also be sociodemographic, disease characteristics and treatments. It is good
301 practice to include covariates in order to avoid mis-specification of the model (resulting in the
302 effects of the omitted variable being allocated to the error term and biased estimates for the
303 coefficients). This remains the case even though the economic evaluation may not be
304 designed to directly use each of these explanatory variables. The analyst can still use the
305 mapping and simply set the value of the explanatory variable to that appropriate to their
306 setting. This is preferable to omitting the explanatory variable. Of course, judgment is
307 required here in order to avoid the inclusion of covariates that are highly correlated in the
308 interest of developing a parsimonious mapping model.

309 Covariates should be theoretically justified *a priori* and reported in a manner that permits
310 analysts to use results whether the covariate in question is used directly in their specific CEA
311 or not. For instance, for most uses of mapping functions in CEA, the inclusion of age as a
312 covariate is required and should be retained in preferred models even if not statistically
313 significant. This allows any effect of ageing, independent of that which is captured as part of
314 the clinical outcome measure(s), to be properly reflected. Where the mapping is intended for
315 use in a CEA alongside a trial, covariates common to both the mapping dataset and the trial
316 can be used to improve the generalizability of one to the other.

317

Summary of statistical modelling recommendations

1. Consider whether the cost-effectiveness analysis requires a formal regression based mapping model approach, or if it is suitable to take the mean value for sub-samples of patients.
2. If regression is required then model selection should be based on:
 - a. Consideration of the most straightforward statistical model type whose assumptions are compatible with the target utility instrument. Use a plot of the distribution of the utility data to help inform that choice.
 - b. Existing empirical evidence of the performance of different methods. There is no reason for this to be restricted to evidence from any specific disease area.
 - c. The type of cost-effectiveness analysis where the mapping will be used and the extent to which biased estimates will affect the results.
3. For response mapping, models should be selected that respect the ordered nature of the categorical data in the descriptive system. Expected values should be calculated analytically.
4. Selection of the preferred mapping model is an iterative process that should conform to good practice common to all regression analyses.
5. Covariates should be theoretically justified *a priori*. Exclusion of covariates, even if they are not to be used in the cost-effectiveness model, risks mis-specification.

318 **5. REPORTING OF MAPPING STUDIES**

319 Mapping studies often form an important element of evidence submitted to decision-making
320 Health Technology Assessment (HTA), pricing or reimbursement authorities. The findings
321 must, therefore, be reported in a manner that allows a full assessment of the quality and
322 relevance of the mapping by those that do not have access to the individual level data. In
323 addition to this transparency requirement, it will be helpful to other analysts that sufficient
324 information is reported to use the results in their own CEAs.

325 *The dataset*

326 Where more than one dataset could feasibly be used for mapping, provide a qualitative
327 account of the selection rationale, at a minimum. The characteristics of the sample used in the
328 estimation dataset must be provided fully. All variables should be described in terms of a
329 measure of central tendency and distribution. Special attention should be given to the full
330 distribution of patient observations at the extremes of disease severity, as described by the
331 disease specific measures to be used as explanatory variables. This gives an indication of the
332 extent to which the sample overlaps with the patients that are the focus of any CEA and,
333 therefore, the extent of extrapolation required beyond the observed data.

334 Full information must be provided about the methods for sampling patients, both in the study
335 as a whole and those sub-samples selected for use in the mapping study.

336 Many studies will include multiple observations from the same individuals over time. In this
337 situation, it is important to report the pattern of those multiple, longitudinal observations and
338 any features of the patients that change over those observations. For instance, if the follow-up
339 period is substantial, then age is an important variable that will vary substantially from
340 baseline. The number of available observations will differ according to the combination of
341 covariates selected and this can lead to substantial differences between any final analysis and
342 the description of the entire study sample. This also has implications for the ability to
343 compare between models using measures of fit or penalised likelihood statistics.

344 *Justification of statistical model type*

345 As outlined above, there are numerous statistical challenges inherent in the analysis of utility
346 data arising from its distributional features. The analyst should seek to select and justify their
347 choice of method(s) *a priori* with reference to existing literature that has tested alternative
348 methods using the target preference-based measure in question, examination of the
349 distributional features in the estimation dataset, and the proposed use of the mapping function
350 in any future cost-effectiveness study.

351 An algebraic description of the model is transparent, concise, unambiguous and ensures
352 results can be used correctly by any competent analyst. Non-standard models, that have not
353 been described elsewhere, must always contain such a description. An example of a predicted
354 value from the mapping regression for some set of covariates should be reported. In some
355 publications, additional software that calculates predictions for user defined inputs has been
356 provided^{25,33}.

357 *Justification for covariates used and how specified*

358 Datasets used for mapping will typically offer the analysts a broad range of potential
359 explanatory variables. These cover disease specific outcome measures, which often may be
360 scored either as multiple components or summary index scores, of which there may be more
361 than one, clinical measures, symptom specific information and demographics *inter alia*. A
362 theoretical justification should be given for the inclusion of all variables within the set to be
363 examined in the statistical analyses. It is instructive to provide an account of the dimensions
364 of quality of life covered in the disease specific outcome(s) and contrast them with those
365 covered by the target utility-based measure.

366 The methods used to move from a potentially large set of explanatory variables to a preferred
367 model that is likely to include a smaller number, and in a particular form, must be detailed.
368 There are many ways in which such regression models can be determined⁹.

369 *Model selection and performance*

370 Theoretical justification for the selection of model type(s) should be provided drawing on
371 previous literature and the specific features of the mapping to be performed, with a particular
372 focus on the target utility measure. Regression models make assumptions which should be
373 explicitly acknowledged and tested or assessed for plausibility. The proposed use of the
374 mapping, if known, should also be discussed. Relevant aspects include the range of disease
375 for which the results will be used, the manner in which uncertainty is to be considered and
376 whether the analysis requires only expected utility values conditional on covariates (as is
377 typically the case in a cohort decision model) or if simulated data is required (as in a trial-
378 based analysis or patient level simulation model).

379 Results must be reported in a manner that provides transparency: readers of the results must
380 be made aware of the process of selecting a preferred model(s) from the set of feasible ones
381 and they must be provided with sufficient information to judge the validity of that process.
382 This means that they need to be able to fully assess the performance of the preferred model(s)
383 (and will require details on at least some aspects of performance of the less preferred
384 models). Judgements are required at each stage of the model building process: reporting
385 needs to highlight these judgements and their rationale. Sufficient information should be
386 supplied to allow readers to be able to use the results of the mapping model in future cost-
387 effectiveness studies.

388 One aspect of performance that is particularly important is model fit – the extent to which
389 modelled values coincide with those observed in the data. Movement to a preferred model
390 should not mechanically follow some rule-based on overall fit. Specific judgement will be
391 required and this will be context specific; for example, whether or not to include a particular
392 covariate. Detailed information on model fit is required, however, for the final preferred
393 model(s). Summary measures of fit like the R^2 are of very limited value here, particularly
394 when presented in isolation, and provide little information of the validity of the mapping for
395 use in subsequent CEA. The degree of between patient variability is inherently high in quality
396 of life data, given the (warranted) subjective nature of quality of life. This results in relatively

397 low R^2 statistics. Penalised likelihood statistics, such as the Akaike Information Criteria and
398 Bayesian Information Criteria (AIC/BIC), provide a more appropriate means for comparisons
399 of specifications within model types. Other summary measures of fit such as the Mean
400 Absolute Error (MAE) and Root Mean Squared Error (RMSE) have typically been applied in
401 the mapping literature. These measures have their origins in the field of forecasting. It should,
402 therefore, be recognised that these measures can appear very insensitive when applied in the
403 mapping field because of the limited range of the dependent utility variable and the degree of
404 variability inherent in patient outcome data. Any measure of model fit should be reported
405 both for entire sample and for specific data ranges, defined in terms of the clinical
406 explanatory variable(s). A plot of mean predicted and mean observed utility values
407 conditional on the clinical variable helps to identify the existence and location of any
408 systematic bias (see, for example, Wailoo et al.³³) and where that bias occurs.

409 The fit of a model should not be assessed solely by reference to the point estimates of the
410 predicted values compared to the data. It should also consider the uncertainty around those
411 predictions and the model outputs once patient variability is included, as described below.

412 *Reporting of results*

413 All coefficient values must be reported to a sufficient number of decimal places to permit
414 accurate estimation. Rescaling and centering covariates around their sample mean can
415 facilitate this. Uncertainty in the estimated coefficients and associated correlation is
416 imperative to allow the reflection of parameter uncertainty in the CEA – the covariance
417 matrix should therefore be routinely reported³⁴ to allow probabilistic sensitivity analysis
418 (PSA) to be undertaken. In addition to parameter uncertainty, the use of a mapping function
419 to impute data at the individual level (for example, when conducting an analysis alongside a
420 clinical trial) requires that the individual level variation is also reflected. In real world data, it
421 is obvious that individuals with identical observable characteristics do not report identical
422 health utility values. If mapping regression models are used simply to impute the same
423 conditional expected value for these individuals, that individual level unexplained variability
424 has been ignored and misrepresents both the clinical study and the results of the mapping.
425 Information on the assumed degree and form of this variability is contained in the mapping
426 regression error term(s) distribution and can be used as the basis for simulation methods that
427 reflect this. Therefore, it is also essential that details of the error terms are reported routinely.
428 With the availability of on-line materials, published mapping studies have no reason not to
429 include these important items of information.

430 The guidance above relating to model selection suggests that one ought not select a model
431 that is capable of producing estimates that lie outside the feasible range for the utility scale.
432 But if such a model has been selected then when sampling from the mapping function, either
433 for uncertainty or variability analysis, the frequency with which these samples lie outside the
434 feasible range must be reported. It must also be reported how such unfeasible values were
435 subsequently used or amended in the CEA. When a mapping is produced without any specific
436 CEA in mind, it can still be useful to report the results of a simulated dataset from the model.
437 This can help inform future CEAs and also forms a means of comparing the distribution of

438 the data simulated from the model to the distribution of the original data (and can thus be
439 used as part of the model selection process).

440 *Empirical Validation*

441 As with other statistical models, validation of the mapping model is relevant. Much of the
442 guidance reported here is based on this requirement. The description of the dataset and the
443 decision problem in which it is to be used, the process of model building and the performance
444 of the final preferred model – each of these elements provides information on validation. To
445 what extent can we have confidence that the model’s predictions are accurate within the
446 relevant patient group and to what extent might they be relevant in other similar patient
447 groups? Existing UK guidelines on mapping recommend empirical validation¹⁴ in this
448 respect, described as estimation of the model in two datasets, either from two separate studies
449 (external validation) or from splitting a single dataset (internal validation), though numerous
450 other methods can be used for internal validation (for example, using bootstrapping-based
451 approaches). In many situations, these empirical validation techniques will simply not be an
452 option because there is only one candidate dataset of insufficient sample size to contemplate
453 splitting.

454 Where any of these validation methods could feasibly be undertaken, there remains
455 uncertainty about which of the available range of methods are most appropriate in the
456 mapping setting and the additional value of the information these analyses provide. Sample
457 splitting imposes the additional penalty of reduced sample size for estimation. For these
458 reasons, we believe it would be premature to recommend empirical validation be conducted
459 for all mapping studies. This is consistent with approaches undertaken for other regression-
460 derived inputs to CEA.

461 Validation of alternative *methodological* approaches to the analysis of utility data can be
462 achieved through repeated head-to-head testing in real-world and simulated datasets from
463 different disease areas. However, routine multi-sample validation methods are not required
464 for standard applied mapping studies because of the limitations noted above.

Summary of reporting standards recommendations

1. Describe relevant differences between datasets that are candidates for mapping estimation
2. Give full details of the selected dataset. Describe how the study was run and patients were sampled. Provide baseline and follow-up characteristics including the distribution of patients' disease severity. Missingness in the longitudinal pattern of responses should be described.
3. Plot the distribution of the utility data.
4. Justify the type of model(s) selected with reference to the characteristics of the target utility distribution and the proposed use of the mapping function.
5. Compare the dimensions of health covered by the target utility instrument and those covered by the explanatory clinical measure(s).
6. Describe the approach to determining the final model. Include tests conducted and judgements made.
7. Summary measures of fit are of limited value for the total sample. Provide information on fit conditional on disease severity as measured by the clinical outcome measure(s). A plot of mean predicted versus mean observed utility conditional on the clinical variable(s) should be included.
8. Coefficient values, error term(s) distributions(s), variances and covariances are required.
9. Provide an example predicted value for some set of covariates. Consider providing a program that calculates predictions for user defined inputs.
10. Parameter uncertainty in a mapping regression should be reflected using standard methods for Probabilistic Sensitivity Analysis (PSA). Assessment of model suitability for use in cost-effectiveness analysis should also consider the distribution of utility values for PSA, with particular focus on whether these lie outside the feasible utility range for the PBM.
11. When imputing data from a mapping function individual level variability should be incorporated using simulation methods and information about the distribution of the error term(s). These simulated data can be compared to the raw observed data, including an assessment of the range of values compared to the feasible range for the PBM.
12. Re-estimation of mapping results in a separate dataset is not routinely required.

466 **6. THE USE OF RESULTS FROM MAPPING MODELS.**

467 *Selection of a mapping model for a cost-effectiveness study*

468 Analysts may often need to select an existing mapping, perhaps from the published literature,
 469 to populate their cost-effectiveness model. In some situations, there may be no existing
 470 mapping that matches the population of interest. This might be that the precise characteristics
 471 of the patients do not match in terms of demographics, stage or severity of disease. In other
 472 situations it may be a more fundamental disparity such as the mapping being based on
 473 patients with a different disease. For example, the EORTC QLQ30 is a PROM used with
 474 patients with any type of tumour. Mappings have been estimated based on samples of patients
 475 with breast cancer³⁵. Judgements about the suitability of a mapping study in a CEA should be
 476 based on an assessment of the differences between the patients or diseases in question. Are
 477 these differences likely to make the relationship between the mapping covariates and the
 478 target PBM non-generalizable?

479 *Predicted values*

480 The primary use of mapping for economic evaluation is to predict the mean health state
481 utility value for a set of explanatory variables: in other words, the expected value conditional
482 on covariates. If the guidance presented here has been followed, then a full understanding of
483 the model specification and the estimated coefficients will have been provided and it will be
484 obvious how to derive the required expected values. It may also be helpful for the mapping
485 study to report the expected utility value and standard error for a given set of covariates for
486 future reference. Some published studies go further and provide pre-programmed spreadsheet
487 calculators as supplementary files^{25,33}.

488 *Variability*

489 A full specification of the statistical model and its estimated results, including error term(s)
490 distribution(s), provides the required information to allow an analyst to reflect individual
491 level variability. At its simplest, this may comprise a single normally distributed error term
492 with mean zero and variance as reported. It is, therefore, straightforward to sample from the
493 relevant conditional distribution to reflect variability around any required health state/patient
494 characteristics.

495 *Uncertainty*

496 PSA is the standard accepted method for reflecting parameter uncertainty in health economic
497 models. Monte Carlo simulation can be used to sample from the relevant joint distribution for
498 regression model inputs, including mapping studies, provided the model specification,
499 coefficient estimates and variance-covariances are reported.

500 **7. CONCLUSIONS**

501 Whilst the inclusion of appropriate preference-based measures in clinical studies is always
502 recommended (see ISPOR Good Practice Guide Wolowacz et al¹⁰ for guidance on this issue),
503 this will not always be feasible or sufficient for the needs of economic evaluation. Mapping
504 is, therefore, needed to allow analysts to bridge the gap between clinical evidence and the
505 evidence required for economic evaluation. Provided that mapping analyses are undertaken
506 appropriately, reported transparently and their results used appropriately, decision makers can
507 be confident in the validity of estimates obtained in this manner.

508 **8. REFERENCES**

¹ The EuroQol Group (1990). EuroQol-a new facility for the measurement of health-related quality of life. Health Policy 16(3):199-208.

² Brazier, JE, Roberts, JR,. The estimation of a preference-based index from the SF-12. Medical Care, 2004;42(9):851-859.

-
- ³ Horsman, J., Furlong, W., Feeny, D., Torrance, D. The health utilities index (HUI®): concepts, measurement properties, and applications, *Health Qual Life Outcomes*. 2003; 1: 54.
- ⁴ NICE Guide to the Methods of Technology Appraisal 2013
- ⁵ Lopez-Bastida J, Oliva J, Antoñanzas F, Garcia-Altes A, Gisbert R, Mar J, Puig-Junoy J. Spanish recommendations on economic evaluation of health technologies. *Eur J Health Econ*. (2010);11:513–20.
- ⁶ Haute Autorité de santé (HAS) http://www.has-sante.fr/portail/upload/docs/application/pdf/2012-10/choices_in_methods_for_economic_evaluation.pdf 2013.
- ⁷ ISPOR. Pharmacoeconomic Guidelines Around The World . Available at: <http://www.ispor.org/PEguidelines/index.asp> (accessed 1st Feb 2016)
- ⁸ Mortimer D, Segal L, Sturm J. Can we derive an “exchange rate” between descriptive and preference-based outcome measures for stroke? Results from the transfer to utility (TTU) technique. *Health Qual Life Outcomes*. 2009;7:33.
- ⁹ Kearns, B. Ara, R., Wailoo, A., Manca, A., Hernandez Alava, M., Abrams, K., Campbell, M. (2013) “Good Practice Guidelines for the use of Statistical Regression Models in Economic Evaluations”, *Pharmacoeconomics*, DOI 10.1007/s40273-013-0069-y
- ¹⁰ Wolowacz, S. et al Good Research Practices for Collecting Health-State Utility Values for Economic Models in Clinical Studies. forthcoming
- ¹¹ Pennington, B., Davis, S. (2014) “Mapping from the Health Assessment Questionnaire to the EQ-5D: The Impact of Different Algorithms on Cost-Effectiveness Results”, *Value in Health*, Vol.17(8):762-71.
- ¹² Hernandez Alava, M., Wailoo, A., Wolfe, F., Michaud, K. (2014) “A comparison of direct and indirect methods for the estimation of health utilities from clinical outcomes”, *Medical Decision Making*, Vol: 34:919–930.
- ¹³ Devlin, N., Parkin, D. & Browne, J. (2009). Using the EQ-5D as a performance measurement tool in the NHS (Report No. 09/03). London, UK: Department of Economics, City University London.
- ¹⁴ Longworth, L., Rowen, D. NICE DSU Technical Support Document 10: The use of mapping methods to estimate health state utility values. 2011.
- ¹⁵ Petrou S, Rivero-Arias O, Dakin H, Longworth L, Oppe M, Froud R, Gray A. Preferred reporting items for studies mapping onto preference-based outcome measures: The MAPS statement. *Health and Quality of Life Outcomes* (2015) 13:106
- ¹⁶ Brazier JE, Yang Y, Tsuchiya A, Rowen DL. A review of studies mapping (or cross walking) non-preference based measures of health to generic preference-based measures. *Eur J Health Econ* 2009;11:215–25.
- ¹⁷ Basu, A. and A. Manca (2012). "Regression estimators for generic health-related quality of life and quality-adjusted life years." *Med Decis Making* 32(1): 56-69.
- ¹⁸ Khan, Kamran A., et al. "Mapping between the Roland Morris Questionnaire and generic preference-based measures." *Value in Health* 17.6 (2014): 686-695.
- ¹⁹ Hernández Alava M, Wailoo AJ, Ara R. Tails from the Peak District: Adjusted Limited Dependent Variable Mixture Models of EQ-5D Health State Utility Values. *Value Health* 2012; 15: 550-561.
- ²⁰ Acaster S, Pinder B, Mukuria C, et al. Mapping the EQ-5D index from the cystic fibrosis questionnaire-revised using multiple modelling approaches, *Health and Quality of Life Outcomes* 2015;13:33.
- ²¹ Pullenayegum EM, Tarride J, Xie F, et al. Analysis of health utility data when some subjects attain the upper bound of 1: are Tobit and CLAD models appropriate? *Value Health* 2010;13:487–94.
- ²² Huang I, Frangakis C, Atinson MJ, et al. Addressing ceiling effects in health status measures: a comparison of techniques applied to measures for people with HIV disease. *Health Serv Res* 2008;43:327–39.
- ²³ Hunger M, Baumert J, Holle R. Analysis of SF-6D index data: is beta regression appropriate? *Value in Health* 2011;14(5) : 759-67.
- ²⁴ Tsuchiya, A., Brazier, J., McColl, E., Parkin, D. (2002) “Deriving preference-based single indices from non-preference-based condition specific instruments: Coverting AQLQ into EQ5D indices”, University of Sheffield, Health Economics and Decision Science Discussion paper 02/01.
- ²⁵ Wailoo, A., Hernández, M., Philips, C., Brophy, S., Siebert, S. (2015) “Modelling health state utility values in Ankylosing Spondylitis: comparisons of direct and indirect methods”, *Value in Health*, Vol.18(4):425-431.
- ²⁶ Gray A, Rivero-Arias O, Clarke P. Estimating the association between SF-12 responses and EQ-5D utility values by response mapping. *Med Decis Making*. 2006;26(1):18–29
- ²⁷ Conigliani, C., Manca, A., Tancredi, A. (2015) “Prediction of patient-reported outcome measures via multivariate ordered probit models”, *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, Vol.

²⁸ Cramer, J.S., 1986, *Econometric applications of maximum likelihood methods* (Cambridge University Press, New York, NY).

²⁹ Gelman, Andrew, and Kenneth Shirley. "Inference from simulations and monitoring convergence." *Handbook of Markov Chain Monte Carlo* (2011): 163-174.

³⁰ Longworth L, Yang Y, Young T, Mulhern B, Hernández Alava M, Mukuria C, et al. Use of generic and condition-specific measures of health-related quality of life in NICE decision-making: a systematic review, statistical modelling and survey. *Health Technol Assess* 2014;18(9)

³¹ Tosh J, Brennan A, Wailoo A, Bansback N. The Sheffield rheumatoid arthritis health economic model. *Rheumatology* (2011) 50 Suppl 4:iv26-iv31

³² Bansback, N., Marra, C., Tsuchiya, A., et al. (2007) "Using the health Assessment Questionnaire to Estimate Preference-based single indices in patients with Rheumatoid Arthritis", *Arthritis and Rheumatism*, Vol.57(6):963-71.

³³ Wailoo, A., Hernandez Alava, M. Escobar Martinez A. (2014) Modelling the relationship between the WOMAC Osteoarthritis Index and EQ-5D, *Health and Quality of Life Outcomes*, 12:37, DOI: 10.1186/1477-7525-12-37

³⁴ Ara, R., Wailoo, A.J. NICE DSU Technical Support Document 12: The use of health state utility values in decision models. 2011.

³⁵ Crott R, Briggs A. Mapping the QLQ-C30 quality of life cancer questionnaire to EQ-5D patient preferences. *Eur J Health Econ* (2010) 11:427–434