

This is a repository copy of *Coherency and constructions for monoids*.

White Rose Research Online URL for this paper:  
<https://eprints.whiterose.ac.uk/164764/>

Version: Accepted Version

---

**Article:**

Gould, Victoria [orcid.org/0000-0001-8753-693X](https://orcid.org/0000-0001-8753-693X), Hartmann, Miklos, Ruskuc, Nik et al. (2 more authors) (2020) *Coherency and constructions for monoids*. *The Quarterly Journal of Mathematics*. pp. 1461-1488. ISSN 1464-3847

<https://doi.org/10.1093/qmath/haaa040>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# COHERENCY AND CONSTRUCTIONS FOR MONOIDS

YANG DANDAN, VICTORIA GOULD, MIKLÓS HARTMANN, NIK RUŠKUC, AND RIDA-E ZENAB

ABSTRACT. A monoid  $S$  is *right coherent* if every finitely generated subact of every finitely presented right  $S$ -act is finitely presented. This is a finiteness condition, and we investigate whether or not it is preserved under some standard algebraic and semigroup theoretic constructions: subsemigroups, homomorphic images, direct products, Rees matrix semigroups, including Brandt semigroups, and Bruck–Reilly extensions. We also investigate the relationship with the property of being weakly right noetherian, which requires all right ideals of  $S$  to be finitely generated.

## 1. INTRODUCTION

A *finitary property* for a class of algebraic structures  $\mathcal{A}$  is a property that is certainly satisfied by all finite algebras in  $\mathcal{A}$ . Of course, one hopes that the property will also be satisfied by some infinite members of  $\mathcal{A}$ , in such a way that it forces finite-like behaviour. Studying algebras via their finitary properties is a classical tool, introduced by Noether and Artin in the early part of the last century in the context of descending and ascending chain conditions for ‘classical’ algebras such as rings. It has very broad implications today in areas ranging from structure theory to decidability problems.

In this paper we work with the class of monoids, and the finitary property is that of right coherency. We arrive at this property by considering representations of monoids by mappings of sets, as we now describe. Let  $S$  be a monoid with identity 1. A *right  $S$ -act* is a set  $A$  together with a map  $A \times S \rightarrow A$  where  $(a, s) \mapsto as$ , such that for all  $a \in A$  and  $s, t \in S$  we have  $a1 = a$  and  $a(st) = (as)t$ . A right action of  $S$  on  $A$  may also be viewed as a morphism from  $S$  to the monoid of all mappings of  $A$  to itself (with left-to-right composition). We also have the dual notion of a *left  $S$ -act*. Right  $S$ -acts over a monoid  $S$  may be regarded as the non-additive analogue of right  $R$ -modules over a (unital) ring  $R$ .

---

*Date:* August 22, 2020.

*2010 Mathematics Subject Classification.* Primary: 20M10, 20M30, Secondary: 20M12, 20M17.

*Key words and phrases.* monoid,  $S$ -act, coherency, regular, finitary properties.

The authors acknowledge the support of the Engineering and Physical Sciences Research Council (EPSRC) grant EP/I032312/1. Research also partially supported by the Hungarian Scientific Research Fund (OTKA) grant PD115705. The research of Yang Dandan was supported by grant 20170604 of the Young Talents Project of Shaanxi Association for Science and Technology, by grants 20103176174 and JB180714 of the Fundamental Research Funds for the Central Universities, and by grant 2020JM-178 of Shaanxi Province Basic Research Program of Natural Science.

Although the study of the two notions diverges considerably once technicalities set in, one can often begin by forming analogous notions and asking corresponding questions.

A monoid  $S$  is said to be *right coherent* if every finitely generated subact of every finitely presented right  $S$ -act is finitely presented. *Left coherency* is defined dually;  $S$  is *coherent* if it is both right and left coherent. These notions are analogous to their name-sakes for a ring  $R$  [3] (where, of course,  $S$ -acts are replaced by  $R$ -modules). Coherency is a finitary condition for both rings and monoids. As demonstrated by Wheeler [19], it is intimately related to the model theory of  $S$ -acts and  $R$ -modules. Indeed, the first order theory of right  $S$ -acts (respectively, right  $R$ -modules) has a *model companion* if and only if  $S$  (respectively,  $R$ ), is right coherent. Having a model companion ensures that the first order theory is well behaved, in particular, it is amenable to the application of concepts of stability [4, 5, 13].

This paper is the third in a recent series [9, 8] investigating coherency of monoids. Its predecessors were concerned with free monoids in certain varieties of monoids and unary monoids. In particular, [9] showed that all free monoids are coherent, building upon the earlier observation in [6] that free commutative monoids are coherent, and resolving an open question from that paper. This theme is continued in [8], where it was shown that any free left ample monoid is coherent, while free inverse monoids and free ample monoids of rank  $> 1$  are not. In this paper we change tack somewhat, and note that little is known concerning the preservation of right coherency of monoids under standard algebraic constructions. Our primary aim is to start investigations exploring this strand. This also entails establishing connections between right coherency and finitary properties relating to the lattice of right ideals of a monoid.

In what follows we outline the organisation and content of the paper. In Section 2 we introduce the background material needed for the rest of the paper. This in particular includes the relationship between coherency and right congruences. In the case of rings, coherency can be characterised by properties of certain right ideals. Although monoids have right ideals, where a right ideal  $I$  of  $S$  is a subset such that  $IS \subseteq I$ , they do not play the same role as in the case for rings, because we do not obtain *all* single generator right  $S$ -acts by factoring out by a right ideal. In this regard the natural notion corresponding to a right ideal in a ring is that of a *right congruence* of a monoid, that is, an equivalence relation  $\rho$  on  $S$  such that  $ac\rho bc$  for all  $a, b, c \in S$  with  $a\rho b$ . Right ideals give rise to right congruences, but not all right congruences are obtained in this way. In Section 2 we will review the standard conditions expressing coherency in terms of properties of congruences, as well as all other background material needed for the rest of the paper.

In Section 3 we explore further the link between congruences and coherency, and relate the latter with another finiteness condition, that of being *weakly right noetherian*. A monoid  $S$  has this property if every right ideal is finitely generated, and is *right noetherian* if every right congruence is finitely generated. From Normak [15], a right noetherian monoid is right coherent. In the parallel situation for rings, it is well known that the notions of being right noetherian and of being weakly right noetherian coincide, and that they imply right coherency (see, for example, [17]). On the other hand, for a monoid, the notion of being

weakly right noetherian is easily seen to be weaker than that of being right noetherian. For example, any group is weakly right noetherian, but, since (finitely generated) right congruences correspond to (finitely generated) subgroups, it follows that not all groups are right noetherian. Moreover, [6, Example 3.1] tells us that, in general, a weakly right noetherian monoid need not be right coherent. However, if we restrict our attention to *regular* monoids  $S$ , meaning that for all  $a \in S$  there exists  $b \in S$  such that  $a = aba$ , then we can demonstrate a stronger link between right ideals and coherency (Theorem 3.2), from which we are able to deduce:

- Any weakly right noetherian regular monoid is right coherent (Corollary 3.3).

It is known that there are right coherent monoids that are not weakly right noetherian, for example, any free monoid of rank greater than one [9]. We exhibit two further examples at the end of Section 3, both of them regular, in order to demonstrate the independence of various conditions we consider.

In Section 4 we show that certain natural submonoids of right coherent monoids are right coherent. In particular:

- For any right coherent monoid  $S$ , any monoid ideal and any monoid  $\mathcal{J}$ -class are right coherent (Corollary 4.2 and Theorem 4.8).

In the final part of Section 4 and the entire Section 5 we focus on Brandt monoids  $\mathcal{B}(M, I)^1$ , for which we can present a complete picture:

- A monoid  $M$  is right coherent if and only if any Brandt monoid  $\mathcal{B}(M, I)^1$  is right coherent (Corollary 4.10 and Theorem 5.1).

We also consider two related constructions – Rees matrix semigroups and Bruck–Reilly extensions – but for them are only able to present partial results in Proposition 3.5 and Corollaries 3.4, 4.11.

Section 6 is devoted to right coherency of direct products. We exhibit an example which shows that direct products do not preserve coherency in general, and then prove:

- If  $S$  is a right coherent monoid and  $T$  is a finite monoid, then  $S \times T$  is right coherent (Theorem 6.4).

We regard the work in this paper as an opener in the discussion of the behaviour of coherency with regard to natural semigroup constructions, and its relation to other finitary properties. As we proceed we suggest a number of directions for future work. As far as possible we have attempted to make the material self contained, but refer the reader to [11] and [12] for more details of the terminology.

## 2. PRELIMINARIES

Let  $S$  be a monoid with identity 1. We allow  $\emptyset$  to be a subact of any  $S$ -act. The monoid  $S$  may be regarded as a right  $S$ -act over itself, and as such its subacts are exactly its right ideals, so that, in particular, we allow  $\emptyset$  to be a right ideal. In the category of right  $S$ -acts, a morphism between right  $S$ -acts  $A$  and  $B$  is a map  $\theta : A \rightarrow B$  such that  $(as)\theta = (a\theta)s$  for all  $a \in A$  and  $s \in S$ ; we refer to  $\theta$  as an  $S$ -morphism. Since  $S$ -acts form a variety of algebras, the free  $S$ -act  $F_S(X)$  over any set  $X$  exists. The following describes its very transparent structure.

**Proposition 2.1** ([12, Construction 1.5.14]). *Let  $X$  be a set and  $S$  a monoid. Then  $F_S(X) = X \times S$  with the action of  $S$  defined as  $(x, s) \cdot t = (x, st)$ .*

For simplicity in the above we abbreviate  $(x, s)$  to  $xs$  and  $x1$  to  $x$ .

For a right  $S$ -act  $A$  and  $a \in A$  define  $aS := \{as : s \in S\}$ . Then  $A$  is *finitely generated* if there exist  $a_1, \dots, a_n \in A$  such that

$$A = a_1S \cup a_2S \cup \dots \cup a_nS.$$

This is equivalent to  $A$  being an image of a finitely generated free  $S$ -act  $F_S(X)$  where  $|X| = n$ , under an  $S$ -morphism. Thus  $A$  is  $S$ -isomorphic to a quotient of  $F_S(X)$ , as we now explain.

**Definition 2.2.** Let  $A$  be a right  $S$ -act. An equivalence relation  $\rho$  on  $A$  is a *right  $S$ -act congruence on  $A$*  (*congruence* for short, when there is no danger of confusion) if  $a \rho b$  implies  $as \rho bs$  for all  $a, b \in A$  and  $s \in S$ .

If  $\rho$  is a congruence on a right  $S$ -act  $A$ , then the quotient

$$A/\rho = \{a\rho : a \in A\}$$

is a right  $S$ -act under  $(a\rho)s = (as)\rho$ , for any  $a \in A$  and  $s \in S$ . The congruences of  $S$  considered as a right  $S$ -act are precisely the right congruences of  $S$  considered as a monoid. For any right  $S$ -act  $A$ , and any  $a \in A$ , the annihilator of  $a$  is

$$\mathbf{r}(a) := \{(u, v) \in S \times S : au = av\};$$

this is a right congruence on  $S$  and  $S/\mathbf{r}(a) \cong aS$ .

Generating sets for congruences, and, in particular, congruences that can be generated by finite sets, play a key role in this paper. For a set  $H$  consisting of ordered pairs we let  $H^{-1} := \{(x, y) : (y, x) \in H\}$ . We then let  $\overline{H} = H \cup H^{-1}$ , which is the symmetric closure of  $H$ . This notation will be used throughout the paper without further comment. If  $H \subseteq A \times A$ , where  $A$  is an  $S$ -act, we denote by  $\langle H \rangle$  the congruence on  $A$  *generated* by  $H$ ; that is the least congruence on  $A$  that contains  $H$ . Note that  $\langle H \rangle = \langle \overline{H} \rangle$ , which enables us to replace a (finite) generating set by a symmetric (finite) generating set for any congruence. When we want to emphasise the act within which we are generating a congruence, we will write  $\langle H \rangle_A$  for  $\langle H \rangle$ . The following is a standard description of  $\langle H \rangle$ :

**Lemma 2.3** ([12, Lemma 1.4.37]). *Let  $S$  be a monoid,  $A$  an  $S$ -act, and  $H \subseteq S \times S$ . Then for  $a, b \in A$  we have  $(a, b) \in \langle H \rangle$  if and only if  $a = b$  or there exist  $(c_1, d_1), \dots, (c_n, d_n) \in \overline{H}$  and  $t_1, \dots, t_n \in S$  such that*

$$(1) \quad a = c_1 t_1, d_1 t_1 = c_2 t_2, \dots, d_n t_n = b.$$

**Definition 2.4.** A sequence of the form (1) is called an  $\overline{H}$ -sequence connecting  $a$  and  $b$ .

**Definition 2.5.** A right  $S$ -act  $A$  over a monoid  $S$  is *finitely presented* if it is isomorphic to  $F_S(X)/\rho$  for some finite set  $X$  and finitely generated congruence  $\rho$ .

It is well-known from universal algebra that the notion of finite presentability does not depend on the choice of finite generating set; see [2, Section 1.5 and Theorem 2.5.5]. In particular, any finitely presented monogenic (single-generator)  $S$ -act is of the form  $S/\rho$  where  $\rho$  is a finitely generated right congruence on  $S$ . Having set up all the necessary notations, we now give some equivalent characterisations of right coherent monoids that will be used interchangeably throughout the paper:

**Theorem 2.6** ([5, 6]). *The following are equivalent for a monoid  $S$ :*

- (1)  $S$  is right coherent;
- (2) for any finitely generated right congruence  $\rho$  on  $S$  and any elements  $a, b \in S$ :
  - (i) the subact  $(a\rho)S \cap (b\rho)S$  of the right  $S$ -act  $S/\rho$  is finitely generated;
  - (ii) the annihilator

$$\mathbf{r}(a\rho) = \{(u, v) : au \rho av\}$$

*is a finitely generated right congruence on  $S$ ;*

- (3) for any finite  $X$  and finitely generated right congruence  $\rho$  on  $F_S(X)$  and any  $a, b \in F_S(X)$ :
  - (i) the subact  $(a\rho)S \cap (b\rho)S$  of the right  $S$ -act  $F_S(X)/\rho$  is finitely generated;
  - (ii) the annihilator

$$\mathbf{r}(a\rho) = \{(u, v) : au \rho av\}$$

*is a finitely generated right congruence on  $S$ .*

The equivalence of (1) and (2) in Theorem 2.6 is [6, Corollary 3.4]; that of (2) and (3) follows from [5, Theorem 6] and [6, Proposition 3.2].

Let us briefly review some basic concepts of semigroup theory which will be used throughout. For a monoid  $S$  we will denote by  $E(S)$  the set of all idempotents of  $S$ . There is a natural partial order on  $E(S)$  defined by  $e \leq f$  if and only if  $ef = fe = e$ . Green's relation  $\mathcal{R}$  is defined as follows: for  $a, b \in S$  we have  $a \mathcal{R} b$  if and only if  $aS = bS$ , or, equivalently,  $a = bs$ ,  $b = at$  for some  $s, t \in S$ . The relation  $\mathcal{L}$  is defined dually,  $\mathcal{H} = \mathcal{R} \cap \mathcal{L}$  and  $\mathcal{D} = \mathcal{R} \circ \mathcal{L} = \mathcal{L} \circ \mathcal{R}$ . Finally,  $a \mathcal{J} b$  if and only if  $SaS = SbS$ . A monoid is regular if and only if every  $\mathcal{R}$ -class contains an idempotent, or equivalently, every  $\mathcal{L}$ -class contains an idempotent. If there is a *unique* idempotent in each  $\mathcal{R}$ -class and in each  $\mathcal{L}$ -class then the monoid is *inverse*.

Associated to  $\mathcal{J}$  is a preorder  $\leq_{\mathcal{J}}$  defined by  $a \leq_{\mathcal{J}} b$  if and only if  $SaS \subseteq SbS$ . Clearly,  $a \mathcal{J} b$  if and only if  $a \leq_{\mathcal{J}} b$  and  $b \leq_{\mathcal{J}} a$ . More details about Green's relations and of other semigroup notions may be found in [11].

In this paper we will consider some standard semigroup constructions, built over a monoid  $M$ . For convenience we now briefly introduce them; further details may be found in [11].

Let  $M$  be a monoid,  $I, \Lambda$  non-empty sets, and  $P = (p_{\lambda i})$  a  $\Lambda \times I$  matrix with entries from  $M$ . Then the *Rees matrix semigroup*  $\mathcal{M}(M; I, \Lambda; P)$  over  $M$  is the set  $I \times M \times \Lambda$  with multiplication

$$(i, a, \lambda)(j, b, \mu) = (i, ap_{\lambda j}b, \mu).$$

When  $M$  is a group  $\mathcal{M}(M; I, \Lambda; P)$  is a completely simple semigroup (i.e. a semigroup with no proper two-sided ideals possessing minimal left and right ideals), and all such semigroups arise in this way; see [11, Theorem 3.3.1].

This construction can be generalised to a *Rees matrix semigroup with zero*  $\mathcal{M}^0(M; I, \Lambda; P)$  as follows: this time the entries of  $P$  come from  $M \cup \{0\}$ , where  $0$  is a new symbol not in  $P$ . The set of elements of  $\mathcal{M}^0(M; I, \Lambda; P)$  is  $(I \times M \times \Lambda) \cup \{0\}$ , and the multiplication is given by:

$$(i, a, \lambda)(j, b, \mu) = \begin{cases} (i, ap_{\lambda j}b, \mu) & \text{if } p_{\lambda j} \in M, \\ 0 & \text{if } p_{\lambda j} = 0, \end{cases}$$

$$0(i, a, \lambda) = (i, a, \lambda)0 = 0 \cdot 0 = 0.$$

Rees matrix semigroups with  $0$  over a group with the additional condition that every row and every column of  $P$  contain at least one entry over  $M$  are precisely completely  $0$ -simple semigroups, i.e. semigroups with  $0$ , with no proper non-zero ideals, and possessing minimal non-zero left and right ideals.

A special case of the latter construction is obtained by taking  $I = \Lambda$ , and  $P$  the  $I \times I$  identity matrix. This yields a *Brandt semigroup*  $\mathcal{B}(M; I)$ . Here the set of elements is  $(I \times M \times I) \cup \{0\}$  and the multiplication is given by

$$(i, a, j)(j, b, \ell) = \begin{cases} (i, ab, \ell) & \text{if } j = k, \\ 0 & \text{if } j \neq k, \end{cases}$$

$$0(i, a, j) = (i, a, j)0 = 0 \cdot 0 = 0.$$

Brandt semigroups over groups are precisely completely  $0$ -simple inverse semigroups.

Rees matrix semigroups, or Brandt semigroups, are not (except in degenerate cases) monoids; if we adjoin an identity we refer to them as *Rees matrix monoids* or *Brandt monoids*, respectively.

The final construction we introduce here is that of a *Bruck–Reilly extension*  $\text{BR}(M, \theta)$  of a monoid  $M$ . Here,  $M$  is any monoid,  $\theta : M \rightarrow M$  an endomorphism, the set of elements is  $\mathbb{N}_0 \times M \times \mathbb{N}_0$ , where  $\mathbb{N}_0$  denotes the natural numbers *including* zero, and multiplication

is given by

$$(a, g, b)(c, h, d) = (a - b + t, (g\theta^{t-b})(h\theta^{t-c}), d - c + t) \quad \text{where } t = \max\{b, c\}.$$

Special instances of this construction were introduced by Bruck [1] and Reilly [16], after whom it is now named. In the case where the image of  $\theta$  is contained in the group of units of  $M$ , properties such as regularity pass between  $M$  and  $\text{BR}(M, \theta)$ ; making such a restriction here does not advance our results and so we do not impose it.

Bruck–Reilly extensions of groups are precisely bisimple inverse  $\omega$ -semigroups, i.e. inverse semigroups with a single  $\mathcal{D}$ -class in which the natural order on idempotents is isomorphic to the inversely ordered chain  $\omega$ , i.e.  $0 > 1 > 2 > \dots$ .

### 3. COHERENCY AND NOETHERIAN PROPERTIES

We have commented that, in general, a weakly right noetherian monoid need not be right coherent. However, this implication *does* hold for regular monoids. In fact, we can prove a more general result, building on [7, Theorem 5.2], where Gould showed that any regular monoid in which every right ideal is principal is right coherent.

We make repeated use of the following straightforward observation.

**Lemma 3.1.** [7, Lemma 5.1] *Let  $I$  be a right ideal of a monoid  $S$ , let  $\rho$  be a right congruence on  $S$  and let  $x, y \in S$ .*

(i) *The set*

$$I\rho = \{s \in S : s\rho a \text{ for some } a \in I\} = \bigcup\{a\rho : a \in I\}$$

*is a right ideal of  $S$  containing  $I$ .*

(ii) *The set*

$$(I, x) = \{t \in S : xt \in I\}$$

*is a right ideal of  $S$ .*

(iii) *If  $x\rho y$  then  $(I\rho, x) = (I\rho, y)$ .*

The right ideal  $I\rho$  defined in (i) above is known as the  $\rho$ -closure of  $I$ . In the following result, the reader should keep in mind the following notational subtlety: for an element  $a \in S$ ,  $(aS)\rho$  is a right ideal of  $S$ , whereas  $(a\rho)S$  is the monogenic subact of  $S/\rho$  generated by  $a\rho$ .

**Theorem 3.2.** *If  $S$  is a regular monoid in which for every finitely generated right congruence  $\rho$  and every  $a, b, x, y \in S$  the right ideals  $(aS)\rho \cap (bS)\rho$  and  $(aS, x) \cap (bS, y)$  are finitely generated, then  $S$  is right coherent.*

*Proof.* Let  $\rho$  be a finitely generated right congruence on  $S$ , where  $S$  satisfies the conditions of the hypothesis. We may assume that  $\rho = \langle X \rangle$  for some finite symmetric subset  $X \subseteq S \times S$ .



We show that the conditions of Theorem 2.6(2) hold. To see that (2)(i) holds, let  $a, b \in S$ . Note that the map  $\iota: (aS)\rho \cap (bS)\rho \rightarrow (a\rho)S \cap (b\rho)S$  such that  $x \mapsto x\rho$  is a surjective morphism between right  $S$ -acts. Since the right ideal  $(aS)\rho \cap (bS)\rho$  is finitely generated by assumption, so also is its image  $(a\rho)S \cap (b\rho)S$ .

Still with  $a \in S$  we now show that the right congruence  $\mathbf{r}(a\rho)$  is finitely generated, so that the condition from Theorem 2.6(2)(ii) holds. We proceed by identifying a finite subset  $Y$  of  $\mathbf{r}(a\rho)$ , and then by showing that  $Y$  is indeed a generating set.

By assumption the right ideal  $(aS)\rho$  is finitely generated, so that, as  $S$  is regular,

$$(aS)\rho = \bigcup_{e \in K} eS$$

for some finite non-empty set  $K \subseteq E(S)$ . For each  $e \in K$  choose  $z_e \in S$  with  $e\rho az_e$ . Since  $a \in (aS)\rho$  so we may choose and fix  $f \in K$  such that  $a \in fS$ , so that  $a = fa\rho az_fa$ , giving that  $(1, z_fa) \in \mathbf{r}(a\rho)$ .

For  $p, q \in K$  and  $x, y \in S$  we have by assumption that  $(pS, x) \cap (qS, y)$  is a finitely generated right ideal. Hence there exists a finite (possibly empty) set  $L(p, q, x, y) \subseteq E(S)$  with

$$(pS, x) \cap (qS, y) = \bigcup_{h \in L(p, q, x, y)} hS.$$

Now let

$$Y := \{(1, z_fa)\} \cup \{(z_phx, z_qyh) \in \mathbf{r}(a\rho) : p, q \in K, (x, y) \in X, \\ h \in L(p, p, x, x) \cup L(p, q, x, y)\}.$$

The set  $Y$  is finite and  $Y \subseteq \mathbf{r}(a\rho)$ . Let  $\tau = \langle Y \rangle$  so that  $\tau \subseteq \mathbf{r}(a\rho)$ ; we need to prove the reverse inclusion. To this end we state and verify two claims.

**Claim 1.** *Suppose  $(c, d) \in X$ ,  $p \in K$  and  $t \in S$  with  $pct = ct$ . Then there exists  $q \in K$  such that  $(z_pct, z_qdt) \in \tau$  and  $qdt = dt$ .*

*Proof.* From  $pct = ct$  we have  $t \in (pS, c)$ , and hence there exists  $h \in L(p, p, c, c)$  such that  $ht = t$ . Also we have  $pch = ch$ , and so

$$(2) \quad az_pch \rho pch = ch \rho dh.$$

It follows that  $dh \in (aS)\rho$ , and so there exists  $q \in K$  with  $qdh = dh$ . Now

$$(3) \quad dh = qdh \rho az_qdh.$$

Together with (2) this yields  $(az_pch, az_qdh) \in \rho$  and so  $(z_pch, z_qdh) \in \mathbf{r}(a\rho)$ , giving  $(z_pch, z_qdh) \in Y$  by definition. Now, to verify the claim, we have

$$z_pct = z_pcht \tau z_qdht = z_qdt \quad \text{and} \quad qdt = qdht = dht = dt,$$

as required. □

**Claim 2.** *Suppose  $(c, d) \in X$ ,  $p \in K$  and  $t, w \in S$ . If  $pct = ct$  and  $dt = aw$  then  $(z_pct, w) \in \tau$ .*

*Proof.* Note that  $fdt = faw = aw = dt$ , implying  $t \in (pS, c) \cap (fS, d)$ , and hence  $ht = t$  for some  $h \in L(p, f, c, d)$ . Now,

$$az_pch \rho pch = ch \rho dh = fdh \rho az_fdh,$$

implying  $(z_pch, z_fdh) \in \mathbf{r}(a\rho)$ , and hence  $(z_pch, z_fdh) \in Y \subseteq \tau$ . But then

$$z_pct = z_pcht \tau z_fdh = z_fdt = z_faw \tau w,$$

as claimed.  $\square$

Returning to the main proof, suppose  $(u, v) \in \mathbf{r}(a\rho)$ , i.e.  $(au, av) \in \rho$ . Then either  $au = av$ , or else there is a sequence

$$(4) \quad au = c_1t_1, d_1t_1 = c_2t_2, \dots, d_nt_n = av,$$

where for  $1 \leq j \leq n$  we have  $(c_j, d_j) \in X$  and  $t_j \in S$ . In the first case we have

$$u = 1u \tau z_fau = z_fav \tau 1v = v.$$

Consider now the second case. First note that  $fc_1t_1 = faw = au = c_1t_1$ . Thus, repeatedly applying Claim 1 along the sequence (4), yields idempotents  $f = p_0, p_1, \dots, p_n \in K$  such that

$$(5) \quad z_{p_0}c_1t_1 \tau z_{p_1}d_1t_1 = z_{p_1}c_2t_2 \tau \dots \tau z_{p_n}d_nt_n$$

and

$$p_id_it_i = d_it_i \quad \text{for all } 1 \leq i \leq n.$$

Furthermore, using Claim 2 with  $(c, d) = (c_n, d_n)$ ,  $p = p_{n-1}$ ,  $t = t_n$  and  $w = v$ , yields  $v \tau z_{p_{n-1}}c_nt_n \tau z_{p_n}d_nt_n$ . Similarly, Claim 2 with  $(c, d) = (d_1, c_1)$ ,  $p = p_1$ ,  $t = t_1$  and  $w = u$ , gives  $u \tau z_{p_1}d_1t_1 \tau z_{p_0}c_1t_1$ . Combining this with (5) yields  $(u, v) \in \tau$ , as required.  $\square$

**Corollary 3.3.** *Every weakly right noetherian regular monoid is right coherent.*

Examples of weakly right noetherian regular monoids include completely regular monoids which can be expressed as a weakly (right) noetherian semilattice  $Y$  of completely simple semigroups  $S_\alpha, \alpha \in Y$ , where, additionally, for each  $\alpha \in Y$  we have that  $S_\alpha$  has finitely many  $\mathcal{R}$ -classes. They also include any regular  $\omega$ -semigroup, that is, any regular semigroup whose semilattice of idempotents is isomorphic to an inversely well-ordered chain  $\omega$ , i.e.  $0 > 1 > 2 > \dots$ . If  $S$  is of this kind, then it is well known that any right (and, indeed, left) ideal of  $S$  is principal, so that, by Theorem 3.2,  $S$  must be coherent. As we mentioned in Section 2, bisimple inverse  $\omega$ -semigroups are precisely the Bruck–Reilly extensions of groups, and so we have:

**Corollary 3.4.** *Every Bruck–Reilly extension  $\text{BR}(G, \theta)$  of a group  $G$  is coherent.*

The conditions of Theorem 3.2 are in fact weaker than the condition of being weakly right noetherian, as we now show via two examples. The first concerns a Rees matrix semigroup  $\mathcal{M} = \mathcal{M}(G; I, \Lambda; P)$  over a group  $G$  (i.e. a completely simple semigroup). It is easy to see that for any  $i \in I$ , the set  $R_i = \{(i, g, \lambda) : g \in G, \lambda \in \Lambda\}$  is a right ideal, and is generated

by any of its elements. Thus any right ideal  $K$  of  $\mathcal{M}$  is of the form  $K = \bigcup_{j \in J} R_j$  for some  $J \subseteq I$ .

**Proposition 3.5.** *Let  $S = \mathcal{M}(G; I, \Lambda; P)^1$  be a Rees matrix monoid over a group  $G$ . Then  $S$  satisfies the conditions of Theorem 3.2 and hence is right coherent. However,  $S$  is weakly right noetherian if and only if  $I$  is finite.*

*Proof.* We begin by noting that the proper right ideals of  $S$  are exactly the right ideals of  $\mathcal{M}$ , and thus of the form  $K = \bigcup_{j \in J} R_j$  where  $J \subseteq I$ . Moreover,  $K$  is finitely generated if and only if  $J$  is finite. The final statement of the proposition follows immediately. We also observe that if  $K$  and  $L$  are finitely generated right ideals of  $S$ , then so is  $K \cap L$ . Thus to show that the conditions of Theorem 3.2 hold, it is sufficient to show that for any finitely generated right congruence  $\rho$  on  $S$  and any  $\mathbf{a}, \mathbf{b} \in S$  the right ideals  $(\mathbf{a}S)\rho$  and  $(\mathbf{a}S, \mathbf{b})$  are finitely generated.

Let  $X$  be a finite symmetric set of generators for  $\rho$ . Suppose first that  $1\rho = \{1\}$ , so that  $X$  consists entirely of pairs of the form  $((i, g, \lambda), (j, h, \mu))$ . If  $\mathbf{a} = 1$  then  $(\mathbf{a}S)\rho = S = 1S$ . Otherwise, let  $\mathbf{a} = (\ell, k, \nu)$  so that  $\mathbf{a}S = \{\ell\} \times G \times \Lambda$ . If  $(o, x, \kappa) \rho (\ell, y, \tau)$  for some  $(\ell, y, \tau)$  then either  $(o, x, \kappa) = (\ell, y, \tau)$  or  $(o, x, \kappa) = (i, g, \lambda)\mathbf{t}$  for some  $(i, g, \lambda)$  with  $((i, g, \lambda), (j, h, \mu)) \in X$  and  $\mathbf{t} \in S$ . In the first case  $o = \ell$  and in the second  $o = i$ . Since  $X$  is finite we deduce that  $(\mathbf{a}S)\rho$  is finitely generated.

Suppose now that  $1\rho \neq \{1\}$ , so that  $X$  contains at least one pair of the form  $((i, g, \lambda), 1)$ . Again, if  $\mathbf{a} = 1$  then  $(\mathbf{a}S)\rho = S$ . On the other hand, if  $\mathbf{a} = (\ell, k, \nu)$  then

$$1 \rho (i, g, \lambda) = (i, g, \lambda)(\ell, p_{\lambda\ell}^{-1}, \lambda) \rho 1(\ell, p_{\lambda\ell}^{-1}, \lambda) = (\ell, p_{\lambda\ell}^{-1}, \lambda),$$

so that  $1 \in ((\ell, p_{\lambda\ell}^{-1}, \lambda)S)\rho = (\mathbf{a}S)\rho$ , giving that  $(\mathbf{a}S)\rho = S$ .

Now we have:

- if  $\mathbf{a} = 1$  then  $(\mathbf{a}S, \mathbf{b}) = S$ ;
- if  $\mathbf{b} = 1$  then  $(\mathbf{a}S, \mathbf{b}) = \mathbf{a}S$ ;
- if  $\mathbf{a}, \mathbf{b} \neq 1$  and  $\mathbf{a}, \mathbf{b} \in R_i$  for some  $i \in I$ , then  $(\mathbf{a}S, \mathbf{b}) = S$ ;
- if  $\mathbf{a}, \mathbf{b} \neq 1$  with  $\mathbf{a} \in R_i, \mathbf{b} \in R_j$  where  $i \neq j$ , then  $(\mathbf{a}S, \mathbf{b}) = \emptyset$ .

In any case  $(\mathbf{a}S, \mathbf{b})$  is finitely generated, which concludes the proof.  $\square$

We end this section by giving two examples of regular monoids that are right coherent but are not weakly right noetherian. The first of them is a Brandt monoid over a group, and it does not even satisfy the conditions of Theorem 3.2.

**Proposition 3.6.** *Let  $S = \mathcal{B}(G; I)^1$ , where  $G$  is a group and  $I$  is infinite, and let  $\rho = \langle \langle (i, g, i), 1 \rangle \rangle$  for fixed  $i \in I, g \in G$ . Then  $S$  is right coherent, but the ideal  $(0S)\rho$  is not finitely generated.*

*Proof.* That  $S$  is right coherent will be proved in Theorem 5.1. To prove the second assertion, let  $J = \{0\} \cup \bigcup_{j \in I, j \neq i} R_j$ , and we claim that  $J = (0S)\rho$ . For any  $j \in I$  with  $j \neq i$  we have

$$(j, h, j) = 1(j, h, j) \rho (i, g, i)(j, h, j) = 0,$$

and it follows that the right ideal  $J \subseteq (0S)\rho = 0\rho$ . We claim that 1, or equivalently  $(i, g, i)$ , cannot be in  $(0S)\rho$ . Recall that since  $R_i$  is a right ideal, we have  $(i, g, i) \in (0S)\rho$  if and only if  $(i, h, \lambda) \in (0S)\rho$  for any  $(i, h, \lambda) \in R_i$ . But, the latter is impossible, as we now show. Suppose that  $(i, h, \lambda) \in R_i$  and  $(i, h, \lambda)$  is related to  $\mathbf{s} \in S$  via an  $X$ -sequence of length 1. Then  $(i, h, \lambda) = \mathbf{c}\mathbf{t}$  and  $\mathbf{d}\mathbf{t} = \mathbf{s}$ , where  $(\mathbf{c}, \mathbf{d})$  or  $(\mathbf{d}, \mathbf{c})$  is  $((i, g, i), 1)$  and  $\mathbf{t} \in S$ . If  $(\mathbf{c}, \mathbf{d}) = ((i, g, i), 1)$ , then either  $\mathbf{t} = 1$  and then  $\mathbf{s} = 1$ , or  $\mathbf{t} = (i, g^{-1}h, \lambda)$  and then also  $\mathbf{s} = (i, g^{-1}h, \lambda)$ . Similarly, if  $(\mathbf{d}, \mathbf{c}) = ((i, g, i), 1)$ , then  $\mathbf{s} = (i, gh, \lambda)$ . Since a single step in an  $X$ -sequence starting from 1 can only take us to  $(i, g, i)$ , our claim follows by induction on the length of  $X$ -sequences. Consequently,  $(0S)\rho = J$ . As  $I$  is infinite,  $(0S)\rho$  is not finitely generated as a right ideal of  $S$ .  $\square$

For our second example, we return to a regular semigroup  $S$  with  $\omega$ -chain of idempotents. If  $S$  is bisimple, then, as commented in Section 2,  $S$  is isomorphic to a Bruck–Reilly extension  $\text{BR}(G, \theta)$ . This construction was extended by Warne [18], who noticed that one could replace  $\mathbb{N}_0$  by  $\mathbb{Z}$ , and, using the same rule for multiplication, obtain an inverse monoid with chain of idempotents isomorphic to  $\mathbb{Z}$  (where the natural correspondence inverts the order). This construction yields an *extended Bruck–Reilly* extension  $\text{EBR}(G, \theta)$ . Note that  $\text{EBR}(G, \theta)$  is inverse, and is neither a monoid nor weakly right noetherian.

**Proposition 3.7.** *Let  $G$  be a group,  $\theta: G \rightarrow G$  a homomorphism, and  $S = \text{EBR}(G, \theta)^1$ . Then  $S$  is right coherent, but not weakly right noetherian.*

*Proof.* Denote the identity of  $G$  by  $e$ . It is entirely routine to check that the right ideals of  $S$  are precisely  $S, S \setminus \{1\}, \emptyset$  and  $R_i = \{(j, g, k) : j \geq i\}$  for every  $i \in \mathbb{Z}$ , and that  $S \setminus \{1\}$  is the only right ideal that is not finitely generated.

To prove that  $S$  is right coherent we make use of Theorem 3.2. To this end, let  $\rho$  be a right congruence on  $S$  generated by the finite symmetric set  $X \subseteq S \times S$ . Since the right ideals of  $S$  are linearly ordered, it is enough to show that for any  $\mathbf{a} \in S$  and  $\mathbf{e} \in E(S)$  we can have neither  $(\mathbf{a}S)\rho$  nor  $(\mathbf{e}S, \mathbf{a})$  being equal to the non-finitely generated right ideal  $S \setminus \{1\}$ .

We first consider  $(\mathbf{a}S)\rho$ . If  $\{1\}$  is not a  $\rho$ -class then as  $(\mathbf{a}S)\rho$  is a union of  $\rho$ -classes, we have  $(\mathbf{a}S)\rho \neq S \setminus \{1\}$ . On the other hand, if  $\{1\}$  is a  $\rho$ -class then let  $n$  be an integer smaller than any number appearing in any of the first coordinates of elements of  $X$ . Then  $(n, e, n)\rho = \{(n, e, n)\}$ , because  $(n, e, n)$  cannot be written as  $\mathbf{c}\mathbf{t}$  where  $(\mathbf{c}, \mathbf{d}) \in X$  and  $\mathbf{t} \in S$ . As a consequence,  $(\mathbf{a}S)\rho \neq S \setminus \{1\}$  for any  $\mathbf{a} \in S$ .

Consider now the right ideal  $(\mathbf{e}S, \mathbf{a})$ . If  $\mathbf{a} \in \mathbf{e}S$  then  $(\mathbf{e}S, \mathbf{a}) = S$  and if  $\mathbf{a} = 1$ , then  $(\mathbf{e}S, \mathbf{a}) = \mathbf{e}S$ . Otherwise,  $\mathbf{e} = (i, e, i)$  and  $\mathbf{a} = (p, c, q)$  where  $p < i$ , and then an easy calculation yields  $(\mathbf{e}S, \mathbf{a}) = (q + i - p, e, q + i - p)S$ .  $\square$

## 4. SUBMONOIDS

In this section we begin the examination of when right coherency of a monoid  $S$  is inherited by certain monoid subsemigroups. Recall that if  $T$  is a subsemigroup of a monoid  $S$  then  $T$  is a *retract* of  $S$  if there is a morphism (called *retraction*)  $\theta : S \rightarrow S$  such that  $\text{Im } \theta = T$  and  $\theta|_T : T \rightarrow T$  is the identity map on  $T$ . It is clear that if  $T$  is a retract of  $S$  then it must be a monoid subsemigroup.

**Proposition 4.1.** [8, Theorem 6.3] *Let  $S$  be a right coherent monoid and let  $T$  be a retract of  $S$ . Then  $T$  is right coherent.*

**Corollary 4.2.** *Let  $I$  be an ideal of a right coherent monoid  $S$  such that  $I$  has an identity. Then  $I$  is right coherent.*

*Proof.* If  $e$  is the identity of  $I$ , then it is easy to see that  $\theta : S \rightarrow I$  given by  $a\theta = ea$  is a retraction onto  $I$ . The result now follows by Proposition 4.1.  $\square$

For a semigroup  $M$ , we denote by  $M^\perp$  the monoid obtained by adjoining an identity  $\perp$  whether or not  $M$  already possesses one. As another application of the above result, we obtain one half of the following:

**Proposition 4.3.** *Let  $M$  be a monoid. Then  $M$  is right coherent if and only if  $M^\perp$  is right coherent.*

*Proof.* In what follows we denote the identity of  $M$  by  $1_M$ , and that of  $M^\perp$  by  $\perp$ . If  $M^\perp$  is right coherent, then so is  $M$  by Corollary 4.2, since it is a monoid ideal.

For the converse, let  $\rho$  be a finitely generated right congruence on  $M^\perp$ , and let  $\nu = \rho \cap (M \times M)$  be its restriction to  $M$ . We claim that  $\nu$  is finitely generated. Specifically, we prove that if  $\rho$  is generated by a finite set  $H$  then

$$\nu = \langle K \rangle_M \quad \text{where} \quad K = (H \cap (M \times M)) \cup \{(1_M, a) : a \in M, (\perp, a) \in \overline{H}\}.$$

That  $K \subseteq \nu$  and hence  $\langle K \rangle_M \subseteq \nu$  follows from the definition, and the observation that  $(\perp, a) \in \overline{H}$  implies  $(1_M, a) = (\perp 1_M, a 1_M) \in \rho$ . For the reverse inclusion, let  $a, b \in M$  with  $a \neq b$  and  $(a, b) \in \rho$ . Then there is an  $H$ -sequence

$$a = c_1 t_1, d_1 t_1 = c_2 t_2, \dots, d_n t_n = b,$$

where  $(c_i, d_i) \in \overline{H}$  and  $t_i \in M^\perp$  for  $1 \leq i \leq n$ . Multiplying this sequence on both sides by  $1_M$  yields a  $K$ -sequence over  $M$  from  $a$  to  $b$ , as required.

We now want to prove that  $\mathbf{r}(a\rho)$  is finitely generated for any  $a \in M^\perp$ . If  $a = \perp$  then  $\mathbf{r}(a\rho) = \rho$  and there is nothing to prove. So suppose  $a \in M$ . We claim that:

$$(6) \quad \mathbf{r}(a\rho) = \kappa \quad \text{where} \quad \kappa = \langle \mathbf{r}(a\nu) \cup \{(1_M, \perp)\} \rangle_{M^\perp}.$$

Since  $\mathbf{r}(a\nu) \cup \{(1_M, \perp)\} \subseteq \mathbf{r}(a\rho)$ , so  $\kappa \subseteq \mathbf{r}(a\rho)$ . For the reverse inclusion, consider  $(u, v) \in \mathbf{r}(a\rho)$ . If  $(u, v) \in M \times M$  then  $au\rho av$ , so  $au\nu av$ , and hence  $(u, v) \in \mathbf{r}(a\nu) \subseteq \kappa$ . Clearly  $(\perp, \perp)$  belongs to both  $\mathbf{r}(a\rho)$  and to  $\kappa$ . Finally, if  $u = \perp$  and  $v \in M$ , then we have

$a\mathbf{1} = a = a\mathbf{1}_M \rho av$ , so that  $(\mathbf{1}_M, v) \in \mathbf{r}(a\nu) \subseteq \kappa$ ; together with  $(\mathbf{1}, \mathbf{1}_M) \in \kappa$  this implies  $(\mathbf{1}, v) \in \kappa$ , as required.

Having established (6) it immediately follows that  $\mathbf{r}(a\rho)$  is finitely generated. Indeed, since  $M$  is right coherent by assumption, and  $\nu$  is finitely generated, it follows that  $\mathbf{r}(a\nu)$  is finitely generated as an  $M$ -act. Thus, any finite generating set for  $\mathbf{r}(a\nu)$  together with  $(\mathbf{1}_M, \mathbf{1})$  will generate  $\mathbf{r}(a\rho)$  as an  $M^\perp$ -act.

Now let  $a, b \in M^\perp$ ; we need to show that  $(a\rho)M^\perp \cap (b\rho)M^\perp$  is finitely generated. If  $(a\rho)M^\perp = (\mathbf{1}\rho)M^\perp$  then  $(a\rho)M^\perp \cap (b\rho)M^\perp = (b\rho)M^\perp$ ; likewise if  $(b\rho)M^\perp = (\mathbf{1}\rho)M^\perp$ . So suppose neither  $(a\rho)M^\perp$  nor  $(b\rho)M^\perp$  equals  $(\mathbf{1}\rho)M^\perp$ . Then  $a, b \in M$  and

$$(a\rho)M^\perp \cap (b\rho)M^\perp = (a\nu)M \cap (b\nu)M.$$

The right hand side is finitely generated as an  $M$ -act, because  $M$  is right coherent by assumption and  $\nu$  is finitely generated. Hence it is also finitely generated as an  $M^\perp$ -act, completing the proof of the claim and the proposition.  $\square$

There is an analogue of Proposition 4.3 for  $M^0$ , the monoid obtained by adjoining a new zero element to  $M$ , but we will postpone it until the next section (Corollary 5.6).

Corollary 4.2 is about principal ideals possessing an identity. With some further work, we obtain a corresponding result for monoid  $\mathcal{J}$ -classes. We first present a general result on extendability of right congruences on a submonoid.

**Definition 4.4.** Let  $T$  be a subsemigroup of a monoid  $S$ , and suppose that  $T$  has identity  $e$ . We say that  $T$  is:

- (i) *left unitary* if  $a, ab \in T$  implies that  $b \in T$ ;
- (ii) *weakly left unitary* if  $a, ab \in T$  implies that  $eb \in T$ .

The notions of being right unitary and weakly right unitary are dual.

If  $T$  is left unitary, then it is weakly left unitary. For an immediate counterexample to the converse, consider the bicyclic monoid  $B$ ; the subsemigroup  $\{(1, 1)\}$  is weakly left unitary but not left unitary.

**Definition 4.5.** A subsemigroup  $T$  of a monoid  $S$  has the *strong right congruence extension property SRCEP* if, for any right congruence  $\rho$  on  $T$ , we have

$$\rho = \rho^S \cap (T \times T)$$

and  $T$  is a union of  $\rho^S$ -classes, where  $\rho^S$  is the right congruence on  $S$  generated by  $\rho$ .

We note that  $T$  has SRCEP if and only if for any right congruence  $\rho$  on  $T$  we have  $\rho = \tau \cap (T \times T)$ , and  $T$  is a union of  $\tau$ -classes, for *some* right congruence  $\tau$  on  $S$ .

**Lemma 4.6.** *Let  $S$  be a monoid and let  $T$  be a monoid subsemigroup with identity  $e$ . If  $T$  is weakly left unitary then  $T$  has SRCEP.*

*Proof.* Let  $\rho$  be a right congruence on  $T$ . Clearly,  $\rho \subseteq \rho^S \cap (T \times T)$ . Conversely, if  $a \in T$  and  $a \rho^S b$  where  $b \in S$  and  $b \neq a$ , then there exists a  $\rho$ -sequence

$$a = c_1 t_1, d_1 t_1 = c_2 t_2, \dots, d_n t_n = b$$

where  $(c_i, d_i) \in \rho$  and  $t_i \in S$  for  $1 \leq i \leq n$ . From  $a = c_1 t_1 = (c_1 e) t_1 = c_1 (e t_1)$  and the fact  $T$  is weakly left unitary, we have  $e t_1 \in T$ , so that  $d_1 t_1 = (d_1 e) t_1 = d_1 (e t_1) \in T$ . In the same way we obtain  $e t_2, \dots, e t_n \in T$ , and it follows that  $b \in T$  and  $a \rho b$  as required.  $\square$

**Lemma 4.7.** *Let  $J$  be a monoid  $\mathcal{J}$ -class of  $S$  with identity  $e$ . Then  $J$  is weakly left and right unitary.*

*Proof.* Let  $a, ab \in J$ . Then  $ab = aeb \leq_{\mathcal{J}} eb \leq_{\mathcal{J}} e \mathcal{J} ab$  implies that  $eb \mathcal{J} ab$ , that is,  $eb \in J$ , showing that  $J$  is weakly left unitary. Dually,  $J$  is weakly right unitary.  $\square$

**Theorem 4.8.** *Let  $S$  be a right coherent monoid and let  $J$  be a monoid  $\mathcal{J}$ -class of  $S$ . Then  $J$  is right coherent.*

*Proof.* From Lemmas 4.6 and 4.7, we have that  $J$  is weakly unitary and has SRCEP. Let  $e$  be the identity of  $J$  and let  $\rho$  be a finitely generated right congruence on  $J$ .

We will check that the conditions of Theorem 2.6(2) hold. First we show that  $\mathbf{r}(a\rho)$  is finitely generated for any  $a \in J$ , so that Condition (2)(ii) holds. Since  $S$  is right coherent and  $\rho^S$  is finitely generated,  $\mathbf{r}(a\rho^S)$  is generated by a finite set  $H \subseteq S \times S$ . We claim that  $\mathbf{r}(a\rho)$  is generated by the finite set

$$H' = \{(ec, ed) : (c, d) \in H \text{ and } ec, ed \in J\}.$$

To see this, note first that for any  $(c, d) \in H$  we have

$$a(ec) = (ae)c = ac \rho^S ad = (ae)d = a(ed),$$

and it follows from the fact that  $J$  has SRCEP that  $H' \subseteq \mathbf{r}(a\rho)$ . For the converse, suppose that  $(u, v) \in \mathbf{r}(a\rho)$  where  $u \neq v$ , so that  $u, v \in J$  and  $au \rho av$ . Hence  $au \rho^S av$  so that  $(u, v) \in \mathbf{r}(a\rho^S)$  and there is an  $H$ -sequence

$$(7) \quad u = c_1 t_1, d_1 t_1 = c_2 t_2, \dots, d_n t_n = v$$

where  $(c_i, d_i) \in \overline{H}$  and  $t_i \in S$ , for  $1 \leq i \leq n$ . Multiplying the sequence (7) on the left by  $a$  we have

$$au = ac_1 t_1, ad_1 t_1 = ac_2 t_2, \dots, ad_n t_n = av$$

and it follows that  $au \rho^S ac_i t_i \rho^S ad_i t_i \rho^S av$ , for  $1 \leq i \leq n$ . Since  $J$  has SRCEP, we have  $ac_i t_i, ad_i t_i \in J$  and then  $ec_i, ed_i, t_i e \in J$  for  $1 \leq i \leq n$ , giving  $(ec_i, ed_i) \in H'$ . Returning to sequence (7), multiplying through on both left and right by  $e$ , we obtain an  $H'$ -sequence connecting  $u$  to  $v$ . Hence  $\mathbf{r}(a\rho)$  is generated by  $H'$ .

We now turn our attention to the Condition (2)(ii) in Theorem 2.6. At this stage it is helpful to notice that if  $s, t \in S$  and  $s \rho^S t$ , then, since  $\rho^S$  is generated by pairs of elements from  $J$ , we have  $es \rho^S et$  and, if  $s \neq t$ , then  $es = s$  and  $et = t$ .

Let  $a, b \in J$  be such that  $(a\rho)J \cap (b\rho)J \neq \emptyset$ . Consequently,  $au\rho av$  for some  $u, v \in J$  so that  $au\rho^S bv$  and  $(a\rho^S)S \cap (b\rho^S)S \neq \emptyset$ . Since  $S$  is right coherent we have that

$$(a\rho^S)S \cap (b\rho^S)S = YS$$

for some finite  $Y \subseteq S/\rho^S$ . Let  $X$  be a transversal of the  $\rho^S$ -classes in  $Y$ , let  $X' = eX \cap J$  and put  $Y' = \{x'\rho : x' \in X'\}$ . We claim that

$$(a\rho)J \cap (b\rho)J = Y'J.$$

Let  $x'\rho \in Y'$  where  $x' = ex \in J$  and  $x \in X$ . Then  $x\rho^S \in Y$  so there are  $p, q \in S$  with  $ap\rho^S bq\rho^S x$ . It follows by an earlier remark that  $ap\rho^S bq\rho^S ex = x' (= x) \in J$  and by SRCEP,  $ap, bq \in J$  and  $a(ep) = ap\rho bq = b(eq)\rho x'$ . Further,  $ep, eq \in J$  since  $J$  is weakly left unitary. This shows that  $Y' \subseteq (a\rho)J \cap (b\rho)J$ .

Conversely, suppose that  $c\rho \in (a\rho)J \cap (b\rho)J$  so that  $c\rho ah\rho bk$  for some  $h, k \in J$ . Then  $c\rho^S \in (a\rho^S)S \cap (b\rho^S)S$  so that  $c\rho^S = (x\rho^S)w$  for some  $x\rho^S \in Y$ , where we can take  $x \in X$ , and  $w \in S$ . It follows that  $ah\rho^S bk\rho^S xw$ , giving that  $xw \in J$  and  $ah\rho bk\rho xw$ , since  $J$  satisfies the SRCEP. Now  $xw = exw \leq_{\mathcal{J}} ex \leq_{\mathcal{J}} e$ , so that  $ex \in J$  and hence  $ex \in X'$  and  $(ex)\rho \in Y'$ . Finally,  $xw = exw$  so that as  $J$  is weakly left unitary and  $ex \in J$ , we have  $ew \in J$  and  $xw = e(xw) = (ex)w = (ex)(ew)$ , and we conclude  $c\rho = ((ex)\rho)(ew) \in Y'J$ . Thus  $(a\rho)J \cap (b\rho)J = Y'J$ , as required.  $\square$

Let  $M$  be a monoid. It is easy to see that  $M$  is a monoid subsemigroup of any Brandt monoid  $\mathcal{B}(M, I)^1$  or Bruck–Reilly extension  $\text{BR}(M, \theta)$  built over  $M$ . In the remainder of this section we set up machinery to show that if either  $\mathcal{B}(M, I)^1$  or  $\text{BR}(M, \theta)$  is right coherent, then so is  $M$ . Though our approach is quite technical, it allows us to treat these two cases, and others, together.

We will make use of the relations  $\tilde{\mathcal{R}}_E, \tilde{\mathcal{L}}_E$  and  $\tilde{\mathcal{H}}_E$  on a monoid  $S$ , introduced in [14], where  $E \subseteq E(S)$  is a set of idempotents. For any  $a, b \in S$  we have that  $a \tilde{\mathcal{R}}_E b$  if and only if for all  $e \in E$ ,

$$ea = a \Leftrightarrow eb = b.$$

The relation  $\tilde{\mathcal{L}}_E$  is defined dually, and we put  $\tilde{\mathcal{H}}_E = \tilde{\mathcal{R}}_E \cap \tilde{\mathcal{L}}_E$ . Clearly,  $\tilde{\mathcal{R}}_E, \tilde{\mathcal{L}}_E$  and  $\tilde{\mathcal{H}}_E$  are equivalence relations on  $S$ . It is easy to see that  $\mathcal{R} \subseteq \tilde{\mathcal{R}}_E, \mathcal{L} \subseteq \tilde{\mathcal{L}}_E$  and  $\mathcal{H} \subseteq \tilde{\mathcal{H}}_E$ , and if  $S$  is regular and  $E = E(S)$ , then  $\mathcal{R} = \tilde{\mathcal{R}}_E, \mathcal{L} = \tilde{\mathcal{L}}_E$  and so  $\mathcal{H} = \tilde{\mathcal{H}}_E$ . In this vein, if  $u \tilde{\mathcal{R}}_E v$  and  $uS \subseteq wS \subseteq vS$  for some  $w \in S$ , then it is easy to see that  $u \tilde{\mathcal{R}}_E w \tilde{\mathcal{R}}_E v$ .

Suppose now that  $M$  is a monoid subsemigroup of a monoid  $S$ , with identity  $e$ , and that the following conditions hold:

- (a)  $e \in E$  for some  $E \subseteq E(S)$ ;
- (b)  $\tilde{\mathcal{H}}_E$  is a right congruence;
- (c)  $M$  is the  $\tilde{\mathcal{H}}_E$ -class of  $e$ ;



- (d) for any  $u, v \in S$ , if  $e\tilde{\mathcal{R}}_E u \tilde{\mathcal{H}}_E v$ , then  $u = u(pq), v = v(pq)$  for some  $p, q \in S$  with  $up, vp \in M$ .

Note that (d) would certainly hold if  $S$  were regular and  $E = E(S)$ , from [11, Lemmas 2.2.1 and 2.2.2]. In this case  $M$  would be a group, which we know to be right (and left) coherent.

**Proposition 4.9.** *Let  $M$  be a monoid subsemigroup of a right coherent monoid  $S$ , such that  $M$  has identity  $e$  and (a)–(d) above hold. Then  $M$  is right coherent.*

*Proof.* First note that if  $a, ab \in M$ , then as  $e\tilde{\mathcal{H}}_E a$  and  $\tilde{\mathcal{H}}_E$  is a right congruence on  $S$ , we have  $eb\tilde{\mathcal{H}}_E ab \in M$ , so that  $M$  is weakly left unitary. By Lemma 4.6  $M$  has SRCEP.

Suppose now that  $\rho$  is a finitely generated right congruence on  $M$ ; say  $\rho = \langle H \rangle_M$ , where  $H \subseteq M \times M$  is finite. Then  $\rho^S := \langle H \rangle_S$  is finitely generated as a right congruence on  $S$ . Notice that as  $\tilde{\mathcal{H}}_E$  is also a right congruence on  $S$ , and  $H \subseteq \tilde{\mathcal{H}}_E$ , we have that  $\rho^S \subseteq \tilde{\mathcal{H}}_E$ .

Let  $a \in M$ ; we show that  $\mathbf{r}(a\rho)$  is finitely generated. By assumption that  $S$  is right coherent, we have  $\mathbf{r}(a\rho^S) = \langle K \rangle_S$  for some finite  $K \subseteq S \times S$ .

For any  $(u, v) \in K$  we have by definition that  $au\rho^S av$ , so that as  $\rho^S \subseteq \tilde{\mathcal{H}}_E$  and  $\tilde{\mathcal{H}}_E$  is a right congruence, it follows that  $eu\tilde{\mathcal{H}}_E ev$ . For any pair  $(u, v) \in K$  with  $e\tilde{\mathcal{R}}_E eu \tilde{\mathcal{H}}_E ev$ , choose elements  $p_{(u,v)}, q_{(u,v)} \in S$  guaranteed by (d) such that

$$eu = eu(p_{(u,v)}q_{(u,v)}), \quad ev = ev(p_{(u,v)}q_{(u,v)}) \quad \text{and} \quad eup_{(u,v)}, evp_{(u,v)} \in M.$$

Notice that

$$a eup_{(u,v)} = aup_{(u,v)} \rho^S avp_{(u,v)} = a evp_{(u,v)},$$

so that as  $M$  has SRCEP we have

$$K' = \{(eup_{(u,v)}, evp_{(u,v)}) : (u, v) \in K, e\tilde{\mathcal{R}}_E eu \tilde{\mathcal{H}}_E ev\}$$

is a finite subset of  $\mathbf{r}(a\rho)$ .

We claim that  $K'$  generates  $\mathbf{r}(a\rho)$ . To see this, suppose that  $(h, k) \in \mathbf{r}(a\rho)$ , where  $h \neq k$ , so that  $ah\rho^S ak$  and there exists a  $K$ -sequence

$$h = u_1 t_1, v_1 t_1 = u_2 t_2, \dots, v_n t_n = k$$

where  $(u_i, v_i) \in \overline{K}$  and  $t_i \in S$  for  $1 \leq i \leq n$ . Consequently,

$$h = eh = eu_1 t_1, ev_1 t_1 = eu_2 t_2, \dots, ev_n t_n = ek = k.$$

We have  $hS \subseteq eu_1 S \subseteq eS$ , so by an earlier remark,  $e\tilde{\mathcal{R}}_E eu_1$ , and we have already noted that  $eu_1 \tilde{\mathcal{H}}_E ev_1$ . Consequently, there exist  $p_{(u_1, v_1)}, q_{(u_1, v_1)}$  with

$$(eu_1 p_{(u_1, v_1)}, ev_1 p_{(u_1, v_1)}) \in K', \quad eu_1 p_{(u_1, v_1)} q_{(u_1, v_1)} = eu_1 \quad \text{and} \quad ev_1 p_{(u_1, v_1)} q_{(u_1, v_1)} = ev_1.$$

Thus

$$h = eu_1 t_1 = eu_1 p_{(u_1, v_1)} q_{(u_1, v_1)} t_1 = (eu_1 p_{(u_1, v_1)}) (eq_{(u_1, v_1)} t_1).$$

Since  $M$  is weakly left unitary and  $eu_1p_{(u_1, v_1)} \in M$ , we deduce that  $eq_{(u_1, v_1)}t_1$  lies in  $M$  and

$$h = (eu_1p_{(u_1, v_1)})(eq_{(u_1, v_1)}t_1) \langle K' \rangle_M (ev_1p_{(u_1, v_1)})(eq_{(u_1, v_1)}t_1) = ev_1t_1 = eu_2t_2 \in M.$$

Continuing in this manner we obtain  $h \langle K' \rangle_M ev_n t_n = k$ . Thus  $\langle K' \rangle_M = \mathbf{r}(a\rho)$ , as required.

Next we show that  $(a\rho)M \cap (b\rho)M$  is finitely generated for any  $a, b \in M$ . Since  $S$  is right coherent there is a finite set  $C \subseteq S$  such that

$$(a\rho^S)S \cap (b\rho^S)S = \bigcup_{c \in C} (c\rho^S)S.$$

For any  $c \in C$  we have  $c\rho^S au\rho^S bv$  for some  $u, v \in S$ , whence  $c\tilde{\mathcal{H}}_E au\tilde{\mathcal{H}}_E bv$ . Since  $ea = a$  we have  $ec = c$ . If  $c\tilde{\mathcal{R}}_E e$ , then we choose  $p_c, q_c$  such that  $c = cp_cq_c$  and  $cp_c \in M$ . Notice that  $cp_c\rho^S aup_c\rho^S bvp_c$  so that by Lemma 4.6 as  $M$  has SRCEP,  $cp_c\rho a(eup_c)\rho b(evpc)$ , and  $eup_c, evpc \in M$  as  $M$  is weakly left unitary. Thus

$$T \subseteq (a\rho)M \cap (b\rho)M,$$

where

$$T = \{(cp_c)\rho : c \in C, c\tilde{\mathcal{R}}_E e\}.$$

Conversely, if  $w \in M$  and  $w\rho ah\rho bk$  for some  $h, k \in M$ , then we have  $w\rho^S cl$  for some  $c \in C$  and  $\ell \in S$ . As  $M$  has SRCEP,  $cl \in M$  and from  $clS \subseteq cS \subseteq eS$  we have that  $c\tilde{\mathcal{R}}_E e$ . Choosing  $p_c, q_c$  as above we have  $(cp_c)\rho \in T$  and

$$w\rho cl = cp_cq_c\ell = (cp_c)(eq_c\ell).$$

Weak left unitariness of  $M$  gives that  $eq_c\ell \in M$ . Thus  $w\rho \in ((cp_c)\rho)M$ . It follows that  $T$  generates  $(a\rho)M \cap (b\rho)M$ , as required.  $\square$

**Corollary 4.10.** *Let  $S = \mathcal{B}(M; I)^1$  be a Brandt monoid over a monoid  $M$ . If  $S$  is right coherent, then so is  $M$ .*

*Proof.* Let  $1_M$  be the identity of  $M$  and put  $E = \{(i, 1_M, i) : i \in I\} \cup \{0\}$ . It is easy to see that  $\{1\}$  and  $\{0\}$  are  $\tilde{\mathcal{R}}_E$  and  $\tilde{\mathcal{L}}_E$ -classes, and for any  $(i, a, j), (k, b, \ell)$  we have that

$$(i, a, j) \tilde{\mathcal{R}}_E (k, b, \ell) \Leftrightarrow i = k$$

and

$$(i, a, j) \tilde{\mathcal{L}}_E (k, b, \ell) \Leftrightarrow j = \ell.$$

The relation  $\tilde{\mathcal{H}}_E$  is a congruence on  $S$ . Fixing  $i \in I$  and putting  $M_i = \{i\} \times M \times \{i\}$ , we see that (a), (b) and (c) hold with  $e$  being  $(i, 1_M, i)$ . Further, if  $(i, a, j) \tilde{\mathcal{H}}_E (i, b, j)$ , then with  $p = (j, 1_M, i), q = (i, 1_M, j)$  we see that (d) also holds. Thus if  $S$  is right coherent, then by Proposition 4.9, so is  $M_i$ . But, since  $M_i$  is isomorphic to  $M$ , we conclude that  $M$  is right coherent.  $\square$

The next proof, although the calculations are different, follows the exact pattern of the previous, and so we omit it.

**Corollary 4.11.** *Let  $S = BR(M, \theta)$  be a Bruck–Reilly extension of a monoid  $M$ . If  $S$  is right coherent, then so is  $M$ .*

**Remark 4.12.** An easy adjustment to Corollary 4.11 allows us to apply Proposition 4.9 to  $EBR(M, \theta)^1$ .

The monoids in Corollaries 4.10, 4.11 and Remark 4.12 all possess an *inverse skeleton* [10]; the techniques of [10] would provide an alternative unified approach to these results.

**Question 4.13.** Let  $S$  be a right coherent monoid. Theorem 4.8 tells us that if  $J$  is a monoid  $\mathcal{J}$ -class, then  $J$  inherits right coherency from  $S$ . Is the same true for (i)  $J^1$ , any semigroup  $\mathcal{J}$ -class with an identity adjoined? (ii) any monoid principal factor  $I$ ? or (iii)  $I^1$  where  $I$  is a semigroup principal factor with an identity adjoined?

## 5. BRANDT SEMIGROUPS

We saw in Proposition 3.5 that Rees matrix monoids over groups are right coherent. We do not know whether the analogue of this result for Rees matrix monoids with zero holds. We are, however, able to prove that it does for Brandt monoids, not only over groups but in fact over arbitrary right coherent monoids. This complements Corollary 4.10, and is the main result of this section:

**Theorem 5.1.** *Let  $M$  be a right coherent monoid and  $I$  a non-empty set. Then the Brandt monoid  $\mathcal{B}(M; I)^1$  over  $M$  is also right coherent.*

The rest of this section will be devoted to the proof of Theorem 5.1. Throughout  $M$  will denote an arbitrary right coherent monoid,  $S = \mathcal{B}(M; I)^1$  a Brandt monoid over  $M$ , and  $\rho$  a right congruence on  $S$ . The identity of  $M$  will be denoted by  $1_M$ , and that of  $S$  simply by  $1$ . The proof proceeds along the following lines:

- In Lemma 5.2 we establish a classification of congruences on  $S$ , along with certain specific forms of their generating sets.
- In Lemma 5.3 we establish a connection between a congruence on  $S$  and a certain congruence on a free  $M$ -act.
- We verify the usual properties that ensure right coherency – namely finite generation of annihilators and intersections – in Lemmas 5.4 and 5.5 respectively.

We single out one element of the index set  $I$ , denote it by  $\bullet$ , and note that the following holds:

$$(8) \quad (i, b, j) \rho (k, c, j) \Leftrightarrow (i, b, \bullet) \rho (k, c, \bullet) \quad \text{for all } i, j, k \in I, b, c \in M.$$

**Lemma 5.2.** *The right congruence  $\rho$  satisfies precisely one of the following:*

- (i)  $\rho = S \times S = \langle (1, 0) \rangle_S$ ; or

(ii)  $1\rho = \{1\}$  and  $\rho$  has a set of generators

$$H = \{((i, b, \bullet), (k, c, \bullet)) : (i, b, k, c) \in A\} \cup \{((j, d, \bullet), 0) : (j, d) \in B\},$$

for some suitable index sets  $A, B$ ; or

(iii)  $\{1\} \neq 1\rho \neq S$  and  $\rho$  has a set of generators

$$H = \{((i, 1_M, i), 1)\} \cup \{((i, b, \bullet), (i, c, \bullet)) : (i, b, i, c) \in A\} \\ \cup \{((j, d, \bullet), 0) : (j, d) \in B\},$$

for some  $i \in I$  and index sets  $A \neq \emptyset$  and  $B$ .

Further, in (ii) and (iii), if  $\rho$  is finitely generated, we can take  $A$  and  $B$  to be finite.

*Proof.* Note that  $\rho = S \times S$  if and only if  $(1, 0) \in \rho$ , and then  $\rho$  is generated by  $\{(1, 0)\}$ . We suppose therefore that  $(1, 0) \notin \rho$ .

Choose a set of generators  $K$  for  $\rho$ . If  $((i, b, j), (k, c, \ell)) \in K$  and  $j \neq \ell$ , then

$$(i, b, j) = (i, b, j)(j, 1_M, j) \rho (k, c, \ell)(j, 1_M, j) = 0,$$

so that  $((i, b, j), 0), ((k, c, \ell), 0) \in \rho$ . On the other hand, if we are given that  $((i, b, j), 0), ((k, c, \ell), 0) \in K$ , then  $((i, b, j), (k, c, \ell)) \in \rho$ . Thus we may replace any pair in  $K$  with differing third coordinates by two pairs in which the second coordinate is 0. Thus we may take those pairs to be of the form  $((j, d, \bullet), 0)$ .

By (8) we may assume that all the remaining pairs in  $K$ , not involving 0 or 1, are of the form  $((i, b, \bullet), (k, c, \bullet))$ . If there are no pairs involving 1, we may obtain a set of generators as in (ii).

Finally we suppose that there is a pair in  $K$  of the form  $(1, (i, b, j))$  or  $((i, b, j), 1)$ . Since  $1\rho$  is a submonoid of  $S$  not containing 0, we see that  $i = j$  and if  $(1, (k, c, k))$  or  $((k, c, k), 1) \in K$ , then  $i = k$ . We now observe that

$$1\rho(i, b, i) = (i, b, i)(i, 1_M, i) \rho 1(i, 1_M, i) = (i, 1_M, i).$$

We can therefore add  $((i, 1_M, i), 1)$  to  $K$  and replace any pair  $(1, (i, b, j))$  or  $((i, b, j), 1)$  by  $((i, 1_M, \bullet), (i, b, \bullet))$ . By the preceding discussion and (8), the congruence generated by the set will be unchanged, i.e. still equal to  $\rho$ . Further, if we have a pair  $((j, b, \bullet), (k, c, \bullet)) \in K$ , then since

$$(j, b, \bullet) = 1(j, b, \bullet) \rho (i, 1_M, i)(j, b, \bullet)$$

we deduce that either  $(j, b, \bullet) \rho 0 \rho (k, c, \bullet)$ , so that we may replace  $((j, b, \bullet), (k, c, \bullet))$  in  $K$  by  $((j, b, \bullet), 0), ((k, c, \bullet), 0)$ , or else  $j = i$ . We can perform the same manoeuvre for  $k$ . Finally, if necessary to make the set  $A$  non-empty, we can add a pair from the diagonal, and we obtain a set of generators as in (iii).

It follows from the above argument that if the original generating set  $K$  was finite, then  $H$  would be finite as well, and the lemma is proved.  $\square$

From now on we assume that we fix a set of generators  $H$  for  $\rho$  in the form guaranteed by Lemma 5.2. We define the *support of  $A$* , denoted by  $\text{supp } A$ , in cases (ii) and (iii) above, to be the set of elements of  $I$  appearing as first or third coordinates in  $A$ . Let  $\rho_A \subseteq \rho$  be the right congruence on  $S$  generated by

$$H_A = \{((i, b, \bullet), (k, c, \bullet)) : (i, b, k, c) \in A\}.$$

It is easy to see that  $\{1\}$  and  $\{0\}$  are  $\rho_A$ -classes, and if two triples in  $S$  are  $\rho_A$ -related, they have the same third coordinate.

**Lemma 5.3.** *Suppose that  $\rho \neq S \times S$ . Let  $F_A$  be the free right  $M$ -act on  $\{x_h : h \in \text{supp } A\}$ , and let  $\tau_A$  be the congruence on  $F_A$  generated by*

$$G_A = \{(x_i b, x_k c) : (i, b, k, c) \in A\}.$$

Then for any  $(p, d), (q, e) \in I \times M$  we have

$$(p, d, \bullet) \rho_A (q, e, \bullet) \Leftrightarrow x_p d \tau_A x_q e.$$

*Proof.* Suppose first that  $x_p d \tau_A x_q e$ . If  $x_p d = x_q e$ , then  $(p, d, \bullet) = (q, e, \bullet)$ . We suppose therefore that there is a  $G_A$ -sequence

$$x_p d = u_1 t_1, v_1 t_1 = u_2 t_2, \dots, v_n t_n = x_q e$$

where  $n \in \mathbb{N}$ ,  $(u_j, v_j) \in \overline{G}_A$  and  $t_j \in M$  for  $1 \leq j \leq n$ ; write  $u_j = x_{i_j} b_j, v_j = x_{k_j} c_j$ .

Then

$$\begin{aligned} (p, d, \bullet) &= (i_1, b_1, \bullet)(\bullet, t_1, \bullet) \rho_A (k_1, c_1, \bullet)(\bullet, t_1, \bullet) = (i_2, b_2, \bullet)(\bullet, t_2, \bullet) \\ &\quad \rho_A (k_2, c_2, \bullet)(\bullet, t_2, \bullet) \rho_A \dots \rho_A (k_n, c_n, \bullet)(\bullet, t_n, \bullet) = (q, e, \bullet). \end{aligned}$$

For the reverse implication, suppose  $(p, d, \bullet) \rho_A (q, e, \bullet)$ . If  $(p, d, \bullet) = (q, e, \bullet)$  then  $x_p d \tau_A x_q e$ . Otherwise, we have a  $\rho_A$ -sequence

$$(p, d, \bullet) = \mathbf{b}_1 \mathbf{m}_1, \mathbf{c}_1 \mathbf{m}_1 = \mathbf{b}_2 \mathbf{m}_2, \dots, \mathbf{c}_n \mathbf{m}_n = (q, e, \bullet)$$

where  $n \in \mathbb{N}$ ,  $(\mathbf{b}_j, \mathbf{c}_j) \in \overline{H}_A$  and  $\mathbf{m}_j \in S$  for  $1 \leq j \leq n$ ; write  $\mathbf{b}_j = (i_j, b_j, \bullet)$ ,  $\mathbf{c}_j = (k_j, c_j, \bullet)$ . Since each  $\mathbf{b}_j, \mathbf{c}_j$  have common third coordinate, the element 0 does not appear in the above sequence. By replacing any  $\mathbf{m}_j = 1$  by  $(\bullet, 1_M, \bullet)$ , we can assume that  $\mathbf{m}_j = (\bullet, m_j, \bullet)$  for  $1 \leq j \leq n$ . Then

$$x_p d = x_{i_1} b_1 m_1 \tau_A x_{k_1} c_1 m_1 = x_{i_2} b_2 m_2 \tau_A \dots \tau_A x_{k_n} c_n m_n = x_q e,$$

thus completing the proof of the lemma.  $\square$

We now turn to right annihilators.

**Lemma 5.4.** *If  $\rho$  is finitely generated then for any  $\mathbf{a} \in S$  the right annihilator  $\mathbf{r}(\mathbf{a}\rho)$  is finitely generated.*

*Proof.* Note that  $\mathbf{r}(1\rho) = \rho$  and  $\mathbf{r}(0\rho) = S \times S$ , which are both finitely generated. So we assume that  $\mathbf{a} = (u, a, v)$  and that  $\mathbf{r}(\mathbf{a}\rho) \neq S \times S$ ; in particular  $(\mathbf{a}, 0) \notin \rho$ . If there is a  $((i, 1_M, i), 1) \in H$  and  $u \neq i$  then  $\mathbf{a}\rho 0$ , a contradiction. Thus, we assume that if we are in case (iii), then  $u = i$ .

To construct a finite set of generators of  $\mathbf{r}(\mathbf{a}\rho)$ , first choose a finite set  $T$  of generators of  $\mathbf{r}(a)$  and let

$$R_1 = \{((v, x, v), (v, y, v)) : (x, y) \in T\} \cup \{(1, (v, 1_M, v))\};$$

clearly,  $R_1 \subseteq \mathbf{r}(\mathbf{a}\rho)$ .

If  $u$  is in the support of  $A$ , then, by Theorem 2.6, there is a finite set of generators  $U$  of  $\mathbf{r}((x_u a)\tau_A)$ . For any  $(s, t) \in U$  we have  $x_u a s \tau_A x_u a t$ , and it follows from Lemma 5.3 that  $R_2 \subseteq \mathbf{r}(\mathbf{a}\rho)$ , where

$$R_2 = \{((v, s, v), (v, t, v)) : (s, t) \in U\}.$$

Still supposing that  $u$  is in the support of  $A$ , let us consider a pair  $\alpha = (j, d) \in B$ . If  $(x_u a)\tau_A M \cap (x_j d)\tau_A M \neq \emptyset$  (which can only happen if  $j$  is also in the support of  $A$ ), then, again by Theorem 2.6, we have that

$$(x_u a)\tau_A M \cap (x_j d)\tau_A M = \bigcup_{p \in N(\alpha)} (x_{\alpha, p} w_{\alpha, p})\tau_A M$$

for some finite set  $N(\alpha)$ . For any  $p \in N(\alpha)$ , there are elements  $h_{\alpha, p}, k_{\alpha, p} \in M$  such that

$$x_u a h_{\alpha, p} \tau_A x_{\alpha, p} w_{\alpha, p} \tau_A x_j d k_{\alpha, p}.$$

By Lemma 5.3 we have that

$$(u, a h_{\alpha, p}, \bullet) \rho_A (j, d k_{\alpha, p}, \bullet)$$

whence

$$(u, a, v)(v, h_{\alpha, p}, \bullet) = (u, a h_{\alpha, p}, \bullet) \rho (j, d, \bullet)(\bullet, k_{\alpha, p}, \bullet) \rho 0$$

and so

$$((v, h_{\alpha, p}, v), 0) \in \mathbf{r}(\mathbf{a}\rho).$$

Thus, letting

$$R_3 = \{((v, h_{\alpha, p}, v), 0) : \alpha \in B, p \in N(\alpha)\},$$

we have that  $R_3 \subseteq \mathbf{r}(\mathbf{a}\rho)$ .

We now show that  $R = R_1 \cup R_2 \cup R_3$  generates  $\mathbf{r}(\mathbf{a}\rho)$ . We let  $\nu = \langle R \rangle$ , the right congruence of  $S$  generated by  $R$ .

We have already observed that  $R_1 \cup R_2 \cup R_3 \subseteq \mathbf{r}(\mathbf{a}\rho)$ , and so  $\nu \subseteq \mathbf{r}(\mathbf{a}\rho)$ . For the reverse inclusion suppose  $(\mathbf{r}, \mathbf{s}) \in \mathbf{r}(\mathbf{a}\rho)$ ; we aim to show that  $(\mathbf{r}, \mathbf{s}) \in \nu$ .

First consider the case where  $\mathbf{a}\mathbf{r} = \mathbf{a}\mathbf{s} = 0$ , recalling  $\mathbf{a} = (u, a, v)$ . Then we must have either  $\mathbf{r} = 0$ , or else  $\mathbf{r} = (k, r, \ell)$  with  $v \neq k$ . In the latter case we have

$$(k, r, \ell) = 1(k, r, \ell) \nu (v, 1_M, v)(k, r, \ell) = 0.$$

Hence  $\mathbf{r} \nu 0$ ; by symmetry  $\mathbf{s} \nu 0$ , and therefore, by transitivity,  $(\mathbf{r}, \mathbf{s}) \in \nu$ .

Next consider  $\mathbf{ar} = \mathbf{as} \neq 0$ . If both  $\mathbf{r}$  and  $\mathbf{s}$  are triples, say  $\mathbf{r} = (k, r, \ell)$ ,  $\mathbf{s} = (h, s, m)$ , then  $v = k = h$ ,  $\ell = m$  and using  $R_1$  we have that  $(\mathbf{r}, \mathbf{s}) \in \nu$ . If only one of them is a triple, say  $\mathbf{r} = (k, r, \ell)$  and  $\mathbf{s} = 1$ , then from  $(u, a, v)(k, r, \ell) = (u, a, v)1$ , we have  $k = \ell = v$ , and as  $(u, a, v)(v, 1_M, v) = (u, a, v)(v, r, v)$ , we have that  $1 \nu (v, 1_M, v) \nu (v, r, v) = \mathbf{r}$ , i.e. again  $(\mathbf{r}, \mathbf{s}) \in \nu$ . Finally if  $\mathbf{r} = \mathbf{s} = 1$  then it is obvious that  $(\mathbf{r}, \mathbf{s}) \in \nu$ .

Finally, consider the case where  $\mathbf{ar} \neq \mathbf{as}$ . Since  $(\mathbf{ar}, \mathbf{as}) \in \rho$  there must be an  $H$ -sequence

$$(9) \quad \mathbf{ar} = (u, d, \ell) = \mathbf{c}_1 \mathbf{t}_1, \mathbf{d}_1 \mathbf{t}_1 = \mathbf{c}_2 \mathbf{t}_2, \dots, \mathbf{d}_n \mathbf{t}_n = (u, e, m) = \mathbf{as},$$

where  $(\mathbf{c}_j, \mathbf{d}_j) \in \overline{H}$ ,  $\mathbf{t}_j \in S$  for  $1 \leq j \leq n \in \mathbb{N}$ .

Let us first assume that 0 does not occur in the sequence (9). Immediately, we deduce that  $u \in \text{supp } A$ . If we are in case (iii), where a generator  $((i, 1_M, i), 1)$  lies in  $H$ , then we note that  $\text{supp } A = \{i\}$  and  $u = i$ . Then by multiplying every element of the above sequence *on the left* by  $(i, 1_M, i)$ , we may obtain a (possibly shorter) sequence connecting  $\mathbf{ar}$  and  $\mathbf{as}$ , in which 0 does not appear and the generating pairs from  $\overline{H}$  occurring are all of the form  $((i, b, \bullet), (i, c, \bullet)) \in H$ , where  $(i, b, i, c) \in A$ . Thus, we can assume, in both cases (ii) and (iii), that  $(\mathbf{c}_j, \mathbf{d}_j) \in \overline{H}_A$ . It is then easy to see that the third coordinate of every element of the sequence (9) is equal to  $\ell$  (in particular  $m = \ell$ ), so that multiplying *on the right* by  $(\ell, 1_M, \ell)$ , we can assume that all the  $\mathbf{t}_j$ s are triples.

As 0 is not in the sequence (9), and as  $(u, a, v)1 = (u, a, v)(v, 1_M, v)$  and  $(1, (v, 1_M, v)) \in R$ , we can assume without loss of generality that both  $\mathbf{r}$  and  $\mathbf{s}$  are triples, say  $\mathbf{r} = (v, r, \ell)$ ,  $\mathbf{s} = (v, s, \ell)$ . From  $(u, ar, \ell) \rho_A (u, as, \ell)$  we have  $(u, ar, \bullet) \rho_A (u, as, \bullet)$  by (8), and then by Lemma 5.3 we have  $x_u ar \tau_A x_u as$ . Thus  $(r, s) \in \mathbf{r}((x_u a) \tau_A)$ , which is generated by  $U$ . Therefore it is easy to see (using the generators in  $R_2$ ) that  $(\mathbf{r}, \mathbf{s}) \in \nu$ .

Finally, we consider the case where 0 appears somewhere in the sequence (9). We claim that

$$(10) \quad (\mathbf{r}, 0) \in \nu.$$

The assertion is obvious if  $\mathbf{r} = 0$ . Also  $\mathbf{r} \neq 1$  because  $\mathbf{r}(\mathbf{a}\rho) \neq S \times S$ . We can therefore assume that  $\mathbf{r} = (w, r, \ell)$ . If  $w \neq v$ , then

$$(w, r, \ell) = 1(w, r, \ell) \nu (v, 1_M, v)(w, r, \ell) = 0,$$

proving (10) in this case.

Suppose now that  $w = v$ , and consider the first occurrence of 0 in (9), say  $\mathbf{c}_1 \mathbf{t}_1, \dots, \mathbf{c}_i \mathbf{t}_i \neq 0$  and  $\mathbf{d}_i \mathbf{t}_i = 0$ . Noting that  $\mathbf{c}_i \mathbf{t}_i \neq 1$ , we can assume as above that

$$(u, a, v)(v, r, \ell) \rho_A \mathbf{c}_i \mathbf{t}_i.$$

Since  $\mathbf{d}_i \mathbf{t}_i = 0$ , we are forced to have

$$\mathbf{c}_i = (j, d, \bullet), \mathbf{d}_i = 0 \quad \text{for some } \alpha = (j, d) \in B.$$

We can therefore take  $\mathbf{t}_i = (\bullet, x, \ell)$  and notice that

$$(u, ar, \bullet) \rho_A (j, dx, \bullet),$$

whence  $x_u a r \tau_A x_j d x$ . Thus

$$(x_u a) \tau_A M \cap (x_j d) \tau_A M \neq \emptyset$$

so that

$$x_u a r \tau_A x_j d x \tau_A x_{\alpha, p} w_{\alpha, p} y$$

for some  $p \in N(\alpha)$  and  $y \in M$ . Notice that

$$x_u a r \tau_A x_{\alpha, p} w_{\alpha, p} y \tau_A x_u a h_{\alpha, p} y,$$

so that using  $R_2$  we have

$$((v, r, \ell), (v, h_{\alpha, p} y, \ell)) \in \nu.$$

Then using  $R_3$ ,

$$\mathbf{r} = (v, r, \ell) \nu (v, h_{\alpha, p} y, v) = (v, h_{\alpha, p}, v) (v, y, \ell) \nu 0,$$

completing the proof of (10).

Analogously to (10) we have  $(\mathbf{s}, 0) \in \nu$ , and hence  $(\mathbf{r}, \mathbf{s}) \in \nu$  by transitivity. This completes the proof that  $\mathbf{r}(\mathbf{a}\rho) = \nu$ , and hence of the lemma.  $\square$

The following lemma deals with intersections:

**Lemma 5.5.** *If  $\rho$  is finitely generated then for any  $\mathbf{a}, \mathbf{b} \in S$  we have that  $(\mathbf{a}\rho)S \cap (\mathbf{b}\rho)S$  is finitely generated.*

*Proof.* If  $\mathbf{a}$  or  $\mathbf{b}$  is  $\rho$ -related to 1 or 0, or if  $\mathbf{a} = \mathbf{b}$ , then  $(\mathbf{a}\rho)S \cap (\mathbf{b}\rho)S$  is monogenic. Suppose therefore that  $\mathbf{a} \neq \mathbf{b}$  and that both  $\mathbf{a}$  and  $\mathbf{b}$  are triples, say  $\mathbf{a} = (u, a, v)$ ,  $\mathbf{b} = (w, b, z)$ .

If  $(u, a, v)S \cap (w, b, z)S \neq \{0\}$ , then  $u = w$  and  $aM \cap bM \neq \emptyset$ . Since  $M$  is right coherent,  $aM \cap bM = \bigcup_{c \in C} cM$  for some finite set  $C$ , and then

$$U = \bigcup_{c \in C} (u, c, u)\rho \subseteq (\mathbf{a}\rho)S \cap (\mathbf{b}\rho)S.$$

Further, provided that  $u, w \in \text{supp } A$ , with the  $M$ -act  $F_A/\tau_A$  we consider the intersection  $((x_u a)\tau_A)M \cap ((x_w b)\tau_A)M \neq \emptyset$ . Since  $M$  is right coherent, this intersection is finitely generated by Theorem 2.6, i.e. there exists a finite set  $D \subseteq M$  such that

$$(11) \quad ((x_u a)\tau_A)M \cap ((x_w b)\tau_A)M = \bigcup_{d \in D} ((x_u a d)\tau_A)M.$$

If we let

$$V = \{(u, ad, \bullet)\rho : d \in D\}$$

then it follows from (11) and Lemma 5.3 that

$$V \subseteq (\mathbf{a}\rho)S \cap (\mathbf{b}\rho)S.$$

Let  $W$  be the subact of  $S/\rho$  generated by  $U \cup V$ ; we claim that  $W = (\mathbf{a}\rho)S \cap (\mathbf{b}\rho)S$ . By the preceding discussion, we have  $W \subseteq (\mathbf{a}\rho)S \cap (\mathbf{b}\rho)S$ . For the reverse inclusion, first note that  $0\rho \in (\mathbf{a}\rho)S \cap (\mathbf{b}\rho)S$  always, and if  $(\mathbf{a}\rho)S \cap (\mathbf{b}\rho)S = \{0\rho\}$  there is nothing to prove.



So suppose now that there exist  $\mathbf{h}, \mathbf{g} \in S$  such that  $(\mathbf{ah})\rho = (\mathbf{bg})\rho \neq 0\rho$ ; we want to show that  $(\mathbf{ah})\rho \in W$ .

Now, from  $(\mathbf{ah}, \mathbf{bg}) \in \rho$  we have that either  $\mathbf{ah} = \mathbf{bg} \neq 0$  or else there exists an  $H$ -sequence from one to the other. In the former case  $u = w$ ,  $aM \cap bM \neq \emptyset$ , and  $(\mathbf{ah})\rho \in US$ .

Now suppose that there exists an  $H$ -sequence

$$(12) \quad (u, a, v)\mathbf{h} = \mathbf{c}_1\mathbf{t}_1, \mathbf{d}_1\mathbf{t}_1 = \mathbf{c}_2\mathbf{t}_2, \dots, \mathbf{d}_n\mathbf{t}_n = (w, b, z)\mathbf{g},$$

where  $(\mathbf{c}_i, \mathbf{d}_i) \in \overline{H}$ ,  $\mathbf{t}_i \in S$  for  $1 \leq i \leq n \in \mathbb{N}$ . Since  $(\mathbf{ah})\rho \neq 0\rho$ , no element in the sequence equals 0. We can assume as in Lemma 5.4 that every pair  $(\mathbf{c}_i, \mathbf{d}_i) \in \overline{H}_A$  and every  $\mathbf{t}_i$  is a triple. Note that  $u, w \in \text{supp } A$ . We know that  $\mathbf{h}, \mathbf{g} \neq 0$ . If either of them were 1, then  $(\mathbf{a}\rho)S \cap (\mathbf{b}\rho)S$  would be principal, and our job would be done. We can therefore assume that  $\mathbf{h} = (v, h, y)$ ,  $\mathbf{g} = (z, g, y)$ . Then the sequence (12) implies

$$(u, a, v)(v, h, y) \rho_A (w, b, z)(z, g, y)$$

giving that

$$(u, ah, \bullet) \rho_A (w, bg, \bullet)$$

using (8), and consequently  $x_u ah \tau_A x_w bg$  by Lemma 5.3. Therefore

$$(x_u ah) \tau_A = (x_w bg) \tau_A \in (x_u a) \tau_A M \cap (x_w b) \tau_A M.$$

By the definition of the set  $D$  it follows that there exists  $d \in D$  and  $m \in M$  such that  $(x_u ah) \tau_A = (x_u adm) \tau_A$ . By Lemma 5.3 and (8) we now have  $(u, ah, y) \rho (u, adm, y)$ , and hence

$$(\mathbf{ah})\rho = (u, ah, y)\rho = ((u, ad, \bullet)\rho)(\bullet, m, y) \in VS.$$

This completes the proof of  $(\mathbf{ah})\rho \in W$ , and of the entire lemma.  $\square$

*Proof of Theorem 5.1.* The theorem follows immediately from Lemmas 5.4 and 5.5, using Theorem 2.6.  $\square$

As a consequence we can prove that right coherency is preserved by adjoining or removing a zero element from a monoid, paralleling Proposition 4.3 for identity elements.

**Corollary 5.6.** *A monoid  $M$  is right coherent if and only if  $M^0$  is right coherent.*

*Proof.* ( $\Rightarrow$ ) Note that  $(M^0)^1$  is isomorphic to the  $1 \times 1$  Brandt monoid  $\mathcal{B}(M; \{1\})$ . Hence if  $M$  is right coherent then so is  $(M^0)^1$  by Theorem 5.1. But then it follows that  $M^0$  is right coherent by Proposition 4.3.

( $\Leftarrow$ ) Suppose that  $M^0$  is right coherent, and let  $\rho$  be a finitely generated right congruence on  $M$ . Then  $\rho^0 = \rho \cup \{(0, 0)\}$  is a finitely generated right congruence on  $M^0$ . Let  $a \in M$ . It is easy to see that  $\mathbf{r}(a\rho^0) = \mathbf{r}(a\rho) \cup \{(0, 0)\}$ . Let  $H \subseteq M \times M$  be a finite set such that  $H \cup \{(0, 0)\}$  generates  $\mathbf{r}(a\rho^0)$ . Since  $M^0$  has no zero divisors, a standard argument gives that  $H$  generates  $\mathbf{r}(a\rho)$ .

Now let  $a, b \in M$  and consider  $(a\rho^0)M^0 \cap (b\rho^0)M^0$ . For any  $u, v \in M^0$  we have  $au\rho^0bv$  if and only if  $au = bv = 0$  or  $au, bv \in M$  and  $au\rho^0bv$ . By assumption,  $(a\rho^0)M^0 \cap (b\rho^0)M^0$  is finitely generated, say by  $\{x\rho^0 : x \in X\} \cup \{0\rho^0\}$ , where  $X$  is a finite subset of  $M$ . It is then easy to show, bearing in mind that  $0\rho^0 = \{0\}$ , that  $\{x\rho : x \in X\}$  generates  $(a\rho)M \cap (b\rho)M$ .  $\square$

Unfortunately, we do not know whether the analogues of the above result hold for arbitrary Rees matrix semigroups with or without zero, or for Bruck–Reilly extensions.

**Question 5.7.** If  $M$  is right coherent monoid, are the following necessarily right coherent: (a) an arbitrary Rees matrix monoid  $\mathcal{M}(M; I, \Lambda; P)^1$ ; (b) an arbitrary Rees matrix monoid with zero  $\mathcal{M}^0(M; I, \Lambda; P)^1$ ; (c) an arbitrary Bruck–Reilly extension  $\text{BR}(M, \theta)$ ?

## 6. DIRECT PRODUCTS

In this section we initiate the investigation into when the direct product of two monoids is right coherent. We begin with the following easy observation:

**Proposition 6.1.** *If the direct product  $S \times T$  of two monoids is right coherent then so are both  $S$  and  $T$ .*

*Proof.* The mapping  $S \times T \rightarrow S \times \{1_T\} \cong S$ ,  $(s, t) \mapsto (s, 1_T)$  is a retraction, and hence, by Proposition 4.1,  $S$  is right coherent. The proof for  $T$  is analogous.  $\square$

Next we observe that the converse does not hold: it is possible for the direct product of two right coherent monoids not to be right coherent.

**Example 6.2.** Let  $F$  be the free monoid generated by  $\{a, x, b\}$ . By [9],  $F$  is right coherent, and we claim that  $F \times F$  is not right coherent.

To this end, let  $\rho$  be the right congruence on  $F \times F$  generated by

$$H = \{((ax, 1), (a, x)), ((ab, 1), (aa, 1))\}.$$

We show that the annihilator  $\nu = \mathbf{r}((a, b)\rho)$  is not finitely generated. To see this note that for every  $n \geq 1$  we have

$$(13) \quad (ax^n, b) \rho (ax^{n-1}, xb) \rho \dots \rho (a, x^n b).$$

What is more, the elements appearing in (13) form a complete  $\rho$ -class, because no further rules from  $H$  can be applied to any of them. It follows that  $(x^n, 1)\nu = \{(x^n, 1)\}$  for all  $n \geq 1$ , and it is easy to see this is also true for  $n = 0$ .

In a similar fashion, we have

$$(ax^n b, b) \rho (ax^{n-1} b, xb) \rho \dots \rho (ab, x^n b) \rho (aa, x^n b) \rho (axa, x^{n-1} b) \rho \dots \rho (ax^n a, b),$$

and it follows that  $(x^n b, 1)\nu (x^n a, 1)$ .

Now, let  $K$  be an arbitrary finite set of pairs from  $F \times F$ , and let  $n$  be a natural number greater than the length of any words appearing in any component of any element of  $K$ . It follows that no element of  $K$  has  $x^n a$  as a component. On the other hand, if  $(x^n a, 1)$  is written as  $(u, v)(w, z)$ , with  $(u, v)$  a component of a pair in  $K$ , then  $(u, v) = (x^i, 1)$  for some  $i \geq 0$ . It follows from the discussion above that  $(u, v)$  is not  $\nu$ -related to any element of  $F \times F$  other than itself and so there does not exist a  $K$ -sequence starting from  $(x^n a, 1)$  and ending in  $(x^n b, 1)$ . Therefore  $K$  cannot generate  $\nu$ , and we deduce that  $\nu$  is not finitely generated. Hence  $F \times F$  is not right coherent.

In Example 6.2, certainly  $F \times F$  is a submonoid of  $G \times G$ , where  $G$  is the free group on  $\{a, b, x\}$ . By [6], groups are coherent.

**Corollary 6.3.** *The class of right coherent monoids is not closed under submonoids.*

By way of contrast to Example 6.2, direct products with finite monoids do preserve right coherency:

**Theorem 6.4.** *If  $S$  is a right coherent monoid and  $T$  is a finite monoid then  $S \times T$  is right coherent.*

We will prove this by reference to the original definition of coherency. The main work in doing so is contained in the following result, which may be of independent interest. In what follows, we denote the identities of  $S$  and  $T$  by  $1_S$  and  $1_T$ , respectively.

**Proposition 6.5.** *Let  $S$  be a monoid, let  $T$  be a finite monoid and let  $A$  be an  $S \times T$ -act. Then  $A$  is a right  $S$ -act with the action  $as = a(s, 1_T)$  for all  $a \in A$  and  $s \in S$ . Further:*

- (i)  *$A$  is finitely generated as an  $S$ -act if and only if it is finitely generated as an  $S \times T$ -act;*
- (ii)  *$A$  is finitely presented as an  $S$ -act if and only if it is finitely presented as an  $S \times T$ -act.*

*Proof.* The first statement is straightforward to verify.

(i) ( $\Rightarrow$ ) If  $A$  is finitely generated as an  $S$ -act by a set  $X$ , then  $X$  also generates  $A$  as an  $S \times T$ -act.

( $\Leftarrow$ ) If  $A$  is finitely generated as an  $S \times T$ -act by  $X$ , then for every  $a \in A$  there exist  $(s, t) \in S \times T$  and  $x \in X$  such that  $a = x(s, t) = x(1_S, t)(s, 1_T)$ , which shows that  $A$  is finitely generated as an  $S$ -act by the set  $\{x(1_S, t) : x \in X, t \in T\}$ .

(ii) ( $\Rightarrow$ ) Suppose that  $A$  is finitely presented as an  $S$ -act. Let  $U$  be a finite generating set for  $A$  as an  $S$ -act such that  $u(1_S, t) \in U$  for all  $u \in U$  and  $t \in T$ . Let  $X = \{x_u : u \in U\}$ . Since finite presentability does not depend on the choice of the generating set, there is an  $S$ -morphism  $\theta : F_S(X) \rightarrow A$ , extending the map  $x_u \mapsto u$ , such that  $\ker \theta$  is generated by a finite set of pairs  $H \subseteq F_S(X) \times F_S(X)$ .

Let  $\psi : F_{S \times T}(X) \rightarrow A$  be the  $S \times T$ -morphism extending the map  $x_u \mapsto u$  and let  $H' \subseteq F_{S \times T}(X) \times F_{S \times T}(X)$  be defined by

$$H' = \{(x_u(s, 1_T), x_v(t, 1_T)) : (x_u s, x_v t) \in H\} \cup \{(x_u(1_S, t), x_{u(1_S, t)}) : x_u \in X, t \in T\}.$$

By construction,  $H' \subseteq \ker \psi$ ; we claim that  $H'$  generates  $\ker \psi$ . To this end, suppose that  $(x_u(s, t))\psi = (x_v(h, k))\psi$ , so that  $u(s, t) = v(h, k)$  in  $A$ . Then  $u(1_S, t)(s, 1_T) = v(1_S, k)(h, 1_T)$  so that, regarding  $A$  as an  $S$ -act, we have  $(u(1_S, t))s = (v(1_S, k))h$ . It follows that  $(x_{u(1_S, t)}s)\theta = (x_{v(1_S, k)}h)\theta$  so that, as  $H$  generates  $\ker \theta$ , we have an  $\overline{H}$ -sequence

$$x_{u(1_S, t)}s = (x_{p_1}c_1)t_1, (x_{q_1}d_1)t_1 = (x_{p_2}c_1)t_1, \dots, (x_{q_n}d_n)t_n = x_{v(1_S, k)}h$$

where  $(x_{p_i}c_i, x_{q_i}d_i) \in \overline{H}$  and  $t_i \in S$ ,  $1 \leq i \leq n$ . We therefore have an  $\overline{H'}$ -sequence

$$x_u(s, t) = x_u(1_S, t)(s, 1_T), x_{u(1_S, t)}(s, 1_T) = x_{p_1}(c_1, 1_T)(t_1, 1_T), x_{q_1}(d_1, 1_T)(t_1, 1_T) =$$

$$x_{p_2}(c_2, 1_T)(t_2, 1_T), \dots, x_{q_n}(d_n, 1_T)(t_n, 1_T) = x_{v(1_S, k)}(h, 1_T), x_v(1_S, k)(h, 1_T) = x_v(h, k)$$

connecting  $x_u(s, t)$  and  $x_v(h, k)$ . Thus  $\ker \psi$  is generated by  $H'$ , giving that  $A$  is finitely presented as an  $S \times T$ -act.

( $\Leftarrow$ ) Suppose that  $A$  is finitely presented as an  $S \times T$ -act. Let  $U$  be a finite generating set for  $A$  as an  $S \times T$ -act, let  $X = \{x_u : u \in U\}$  and let  $\theta : F_{S \times T}(X) \rightarrow A$  be the  $S \times T$ -morphism extending the map  $x_u \mapsto u$ . By assumption,  $\ker \theta$  is finitely generated, say by  $H \subseteq F_{S \times T}(X) \times F_{S \times T}(X)$ , where  $H$  is finite. Let

$$X' = \{x_{u(1_S, t)} : u \in U, t \in T\}$$

and put

$$H' = \{(x_{u(1_S, c't)}c, x_{v(1_S, d't)}d) : (x_u(c, c'), x_v(d, d')) \in H, t \in T\}.$$

Let  $\psi : F_S(X') \rightarrow A$  be the  $S$ -morphism extending the map  $x_{u(1_S, t)} \mapsto u(1_S, t)$ . It is easy to check that  $\psi$  is onto and  $H' \subseteq \ker \psi$ .

To show that  $\ker \psi$  is generated by  $H'$ , suppose that  $(x_{u(1_S, t)}s)\psi = (x_{v(1_S, k)}h)\psi$ , so that  $u(s, t) = v(h, k)$  in  $A$  and hence  $(x_u(s, t))\theta = (x_v(h, k))\theta$ . Since  $\ker \theta$  is generated by  $H$ , we have an  $\overline{H}$ -sequence

$$x_u(s, t) = x_{p_1}(c_1, c'_1)(s_1, t_1), x_{q_1}(d_1, d'_1)(s_1, t_1) = x_{p_2}(c_2, c'_2)(s_2, t_2), \\ \dots, x_{q_n}(d_n, d'_n)(s_n, t_n) = x_v(h, k),$$

where  $(x_{p_i}(c_i, c'_i), x_{q_i}(d_i, d'_i)) \in \overline{H}$  and  $(s_i, t_i) \in S \times T$ , for  $1 \leq i \leq n$ . We therefore have an  $\overline{H'}$ -sequence

$$x_{u(1_S, t)}s = (x_{p_1(1_S, c'_1 t_1)}c_1)s_1, x_{q_1(1_S, d'_1 t_1)}d_1)s_1 = (x_{p_2(1_S, c'_2 t_2)}c_2)s_2, \\ \dots, (x_{q_n(1_S, d'_n t_n)}d_n)s_n = x_{v(1_S, k)}h,$$

where  $(x_{p_i(1_S, c'_i t_i)}c_i, x_{q_i(1_S, d'_i t_i)}d_i) \in \overline{H'}$  and  $s_i \in S$  for  $1 \leq i \leq n$ , connecting  $x_{u(1_S, t)}s$  and  $x_{v(1_S, k)}h$ . It follows that  $\ker \psi$  is finitely generated and hence  $A$  is finitely presented as an  $S$ -act.  $\square$

*Proof of Theorem 6.4.* Let  $A$  be a finitely presented  $S \times T$ -act and  $B$  a finitely generated  $S \times T$ -subact of  $A$ . Then, by Theorem 6.5,  $A$  is also a finitely presented  $S$ -act, and  $B$  is finitely generated as an  $S$ -subact of  $A$ . Since  $S$  is right coherent we have that  $B$  is finitely presented as an  $S$ -act, and then by Theorem 6.5 we conclude that  $B$  is also finitely presented as an  $S \times T$ -act, concluding that  $S \times T$  is indeed right coherent.  $\square$

**Question 6.6.** Let  $S$  be a right coherent monoid. For which classes of monoids  $\mathcal{C}$  is it the case that  $S \times T$  is right coherent for all  $T \in \mathcal{C}$ ?

Theorem 6.5 tells us that the class of finite monoids is one example of a class  $\mathcal{C}$  as above.

## REFERENCES

- [1] R.H. Bruck, *A Survey of Binary Systems*, Springer, Berlin, 1958.
- [2] S. Burris and H. Sankapanavar, *A Course on Universal Algebra*, Springer, New York, 1981.
- [3] S.U. Chase, ‘Direct product of modules’, *Trans. Amer. Math. Soc.* **97** (1960), 457–473.
- [4] P. Eklof and G. Sabbagh, ‘Model completions and modules’, *Ann. Math. Logic* **2** (1971), 251–295.
- [5] V. Gould, ‘Model companions of  $S$ -systems’, *Quart. J. Math., Oxford* **38** (1987), 189–211.
- [6] V. Gould, ‘Coherent monoids’, *J. Australian Math. Soc.* **53** (1992), 166–182.
- [7] V. Gould, ‘A rank for right congruences on inverse semigroups’, *Bull. Australian Math. Soc.* **76** (2007), 55–68.
- [8] V. Gould and M. Hartmann, ‘Coherency, free inverse monoids and free left ample monoids’, *Math. Proc. Cambridge Phil. Soc.* **163** (2017), 23–45.
- [9] V. Gould, M. Hartmann and N. Ruškuc, ‘Free monoids are coherent’, *Proc. Edin. Math. Soc. (2)* **60** (2017), 127–131.
- [10] V. Gould and R.-E.Zenab, ‘Semigroups with skeletons and Zappa-Szep products’, *C.G.A.S.A.* **1** (2013), 59–89.
- [11] J.M. Howie, *Fundamentals of Semigroup Theory*, Oxford University Press, Oxford, 1995.
- [12] M. Kilp, U. Knauer and A.V. Mikhalev, *Monoids, Acts, and Categories*, de Gruyter, Berlin, 2000.
- [13] A. Ivanov, ‘Structural problems for model companions of varieties of polygons’, *Siberian Math. J.* **33** (1992), 259–265.
- [14] M.V. Lawson, ‘Semigroups and ordered categories. I. the reduced case’, *J. Algebra* **141** (1991), 422–462.
- [15] P. Normak, ‘On noetherian and finitely presented polygons’, *Uc. Zap. Tartu Gos. Univ.* **431** (1977), 37–46.
- [16] N.R. Reilly, ‘Bisimple  $\omega$ -semigroups’, *Proc. Glasgow Math. Association* **7** (1966), 160–167.
- [17] J. Rotman, *An Introduction to Homological Algebra*, Academic Press, New York, 1979.
- [18] R.J. Warne, ‘ $I$ -bisimple semigroups’, *Trans. American Math. Soc.* **130** (1968), 367–386.
- [19] W.H. Wheeler, ‘Model companions and definability in existentially complete structures’, *Israel J. Math.* **25** (1976), 305–330.

SCHOOL OF MATHEMATICS AND STATISTICS, XIDIAN UNIVERSITY, XI'AN 710071, P. R. CHINA

*Email address:* ddyang@xidian.edu.cn

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, HESLINGTON, YORK, YO10 5DD, UK

*Email address:* victoria.gould@york.ac.uk

*Email address:* hmikklos@gmail.com

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF ST ANDREWS, ST ANDREWS, KY16 9SS,  
UK

*Email address:* nik.ruskuc@st-andrews.ac.uk

DEPARTMENT OF MATHEMATICS SUKKUR IBA UNIVERSITY, PAKISTAN

*Email address:* ridaezenab@iba-suk.edu.pk