

This is a repository copy of *Unravelling the mental health productivity puzzle:where do we start?*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/164930/>

Version: Published Version

---

**Article:**

Maddison, Paula, Castelli, Adriana [orcid.org/0000-0002-2546-419X](https://orcid.org/0000-0002-2546-419X) and Tiffin, Paul A [orcid.org/0000-0003-1770-5034](https://orcid.org/0000-0003-1770-5034) (2020) Unravelling the mental health productivity puzzle:where do we start? The British journal of psychiatry : the journal of mental science. ISSN: 1472-1465

<https://doi.org/10.1192/bjp.2020.142>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

## Editorial

# Unravelling the mental health productivity puzzle: where do we start?

Paula Maddison, Adriana Castelli and Paul A. Tiffin



## Summary

In this editorial we define 'productivity' and 'efficiency' in a mental health service context, outlining the key challenges to measuring these in practice. We attempt to bring clarity of thought to this often debated, but rarely researched area.

## Keywords

Productivity; efficiency; mental health; economics; mental health outcomes.

## Copyright and usage

© The Authors 2020. Published by Cambridge University Press on behalf of the Royal College of Psychiatrists.

**Paula Maddison** (pictured) is a PhD student in the Department of Health Sciences at the University of York and Research Lead for Currency Development and Clinical Outcomes with Tees Esk & Wear Valleys NHS Foundation Trust. **Adriana Castelli** is a Senior Research Fellow at the Centre for Health Economics, University of York. **Paul A. Tiffin** is a Reader in Psychometric Epidemiology in the Department of Health Sciences at the University of York and at Hull York Medical School, a National Institute for Health Research Career Development Fellow and an honorary consultant psychiatrist at Tees, Esk and Wear Valleys NHS Foundation Trust.

Thus, NHS Improvement's report highlighted the degree of variation across services, often built to identical specifications, serving similar populations. However, the sources of this variability remain obscure. Without an understanding of the reasons underlying such differences how can we discern between 'warranted' and 'unwarranted' variation? And unless we are able to accurately define and measure the productivity and efficiency of mental health services, the potential sources of such variation will remain unknown.

'Productivity isn't everything, but in the long-run it is almost everything', stated the Nobel laureate Paul Krugman. Indeed, talk of 'productivity' abounds as the UK National Health Service (NHS), and health services internationally, face unprecedented financial pressures. Mental health services have been especially challenged by the pincer movement of growing demand and workforce shortages, with an urgent need to 'do more with less'. The global economic burden of poor mental health is considerable. For example, in England, it is estimated to cost £105 billion/year. Thus, the effectiveness of these services has implications for the wider economy. In the UK the government has responded by setting out potential strategies for improving the clinical and cost-effectiveness of care.<sup>1</sup> Indeed, an extra £20 billion of NHS funding has been pledged over a 5-year period in England alone (equivalent to a 3.4% annual increase), of which approximately £2.3 billion is expected to be protected for mental health services. The intention is that 380 000 more people with depression and anxiety will receive treatment by 2023–2024, delivered by an enhanced workforce. However, services will need to be as efficient as possible if such aspirations are to be realised.

## Is there any evidence that the NHS is not operating productively or efficiently?


In 2018, NHS Improvement published findings from a large-scale project assessing variations in a broad range of factors across community health services and mental healthcare providers.<sup>2</sup> This highlighted substantial differences across services in a number of areas. For example, the average cost of in-patient treatment between the most and least expensive was found to vary by up to 20%. Variations in sickness absence rates were as high as 116%, with staff engagement, retention, direct care time and frequency all varying markedly between organisations. The report concluded that reducing such variation could save £1 billion/year, if best practice was consistently followed.

## What are 'productivity' and efficiency?

The terms 'productivity' and 'efficiency' are often used in debates regarding healthcare service performance, though rarely in their correct technical sense. Indeed, they are often used as shorthand for clinical activity levels. But does 'looking busy' equate to high levels of either productivity or efficiency? David Maguire, a senior policy analyst at the King's Fund, recently commented on this issue in a blog perceptively entitled: 'The NHS needs to be more productive – or is it more efficient?' (<https://www.kingsfund.org.uk/blog/2019/03/nhs-productive-or-efficient>). So, what do these terms mean, and how should they be applied to mental health services?

'Productivity' measures the quantity of output produced from given inputs, expressed as the ratio of the two. Thus, in healthcare, productivity should relate to the amount of 'care' produced, given the resources available (e.g. funds). However, the ultimate aim is not just to provide care, but to improve health. Therefore, particularly in the context of mental health, productivity must also encompass something greater. Patients seek help not to experience the care itself, but to feel better. Moreover, people who require support from mental health services do not all require the same type of care in order to recover, even if they have similar conditions. Therefore, it is inadequate to simply measure the quantity of patients who have had care delivered without considering its quality. The nature of care should be such that it supports recovery, allowing people to live fulfilling and meaningful lives. Thus, productivity measures must encapsulate both the quantity and quality of the care delivered to each patient.

In contrast to productivity, 'efficiency' is concerned with whether the process is completed using the optimal combination of inputs to produce the maximum outputs, with the least cost and amount of waste. The difference between productivity and efficiency is illustrated with an example in Fig. 1. The solid curve in Fig. 1 depicts a production function, which captures all combinations of inputs and outputs that are 'technically efficient' given



**Fig. 1** The solid curve depicts a 'technically efficient' production function. The degree of 'technical inefficiency' is thus defined by the vertical distance, denoted by the dashed line between  $P_1$  and  $P^*_1$ . This illustrates that a high level of productivity can be achieved while operating at a suboptimal level of efficiency.

the current technological process. Further, the illustrated curve suggests diminishing marginal productivity, i.e. each additional unit of input produces progressively less output. This is the same as saying that the production function exhibits decreasing returns to scale, or the more inputs used, the lower the return in terms of outputs. Assume now that we have two organisations ( $P_1$  and  $P_2$ ) producing one single output ( $O$ ) each using a single input ( $I$ ), given the production function depicted. In terms of the production function, the more inputs used, the lower the return in terms of outputs. Organisation  $P_1$  has a higher level of productivity compared with  $P_2$ , as measured by the ratio of outputs to inputs, which for  $P_1$  ( $O_1/I_1$ ) is larger than that for  $P_2$  ( $O_2/I_2$ ). However,  $P_1$  is not technically efficient as it is not operating on the production function. In fact, it would be feasible for  $P_1$  to produce even more output by producing on the production function at point  $P^*_1$  in Fig. 1, yielding an output of  $O^*_1$  ( $>O_1$ ). The degree of 'technical inefficiency' is thus defined by the vertical distance, denoted by the dashed line between  $P_1$  and  $P^*_1$ . This example shows that it is possible for an organisation to achieve a high level of productivity while operating at a suboptimal level of efficiency.

### **Towards the measurement of productivity and efficiency in mental health services**

One major hurdle in working towards measuring productivity and efficiency in a mental health context is the challenge in identifying the key components of the production process. Inputs are the easiest part of the process to define, consisting of labour (medical, nursing and other healthcare staff), intermediate goods and services (clinical and pharmaceutical supplies, utilities, hotel services, etc.) and capital (non-labour input such as land and building with an asset life greater than 1 year). Ideally, outputs (in terms of care delivered)

should capture the whole course of treatment provided to patients across different care settings. In practice this is often impossible to achieve, given the inability of healthcare systems to track patients across primary, secondary and tertiary care settings. Thus, one often measures the number of patients treated (discharged) and/or the amount of care provided (interventions/activities), usually adjusted to allow for patient case-mix (e.g. the nature, severity and complexity of presentation).

Outcomes, which ideally should reflect the effectiveness of the outputs (care) delivered, are the most challenging element of the process to measure. In fact, selecting and measuring relevant outcomes is not straightforward. Mental health providers have a contractual obligation to collect and report routine outcome measures (ROMs), although completion rates are often low in practice. The potential for ROMs to contribute to a mental health service productivity index has been explored.<sup>3</sup> It was found that multiple barriers existed in relation to the use of ROMs. These included a lack of IT support as well as the perceptions among clinicians as to their usefulness (or otherwise). Given these factors, the report concluded, at the time, that it would be 'premature' to utilise these measures to evaluate service productivity.

It is also the case that different stakeholders may not agree on which outcomes matter the most. We need a lexicon that defines what outcomes are important from the differing perspectives of commissioners, clinicians and, most crucially, patients and carers. For example, the 'recovery' movement emphasises the use of the subjective, individual CHIME factors (connectedness, hope, identity, meaning and empowerment).<sup>4</sup> This contrasts with the currently widely adopted symptom-based questionnaires and rating scales. Indeed, it is well recognised that symptom severity does not always correlate well with functioning and is also at odds with the philosophy of the recovery movement. Moreover, it is unknown as to whether the scores from such symptom measures are




associated with 'hard outcomes', such as the risk of compulsory readmission to hospital.

In a recent King's Fund report, Collins sets out the challenges of conceptualising mental health outcomes, writing that many physical health traumas have a clear path to recovery and a reasonably clear shared sense of priorities. 'In the care of people with mental health conditions, and perhaps people with other chronic conditions, there is a much broader range of sometimes contested objectives.'<sup>5</sup>

We need to ensure that these different perspectives are accommodated within the process of evaluating productivity and efficiency. Inevitably this means that our optimising problem becomes multidimensional in nature. It is therefore quite possible that allocating resources in particular ways may improve outcomes in one domain while degrading those in other areas. However, there may also be situations characterised by 'pareto-dominance', whereby outcomes can be improved in one area without a negative impact elsewhere.

### The way forward

To assess whether the limited funds available are being put to best use, we need a clear framework with which to define and measure productivity and efficiency in the context of mental health services. Moreover, increasingly, workforce availability, as much as monetary resources, are constraining how care can be delivered. There are also important issues related to the optimum (most efficient) stage within the evolution of a mental disorder at which to intervene. Indeed, 'an ounce of prevention is often better than a pound of cure', although identifying such 'sensitive' or 'critical periods' to target for early intervention is not straightforward. We also need an agreed basket of meaningful outcome measures with which to evaluate the effectiveness of care. If the mental health productivity puzzle is to be solved, we need to begin by identifying each piece clearly and put them together, one by one.

**Paula Maddison** , BSc, Department of Health Sciences, University of York; and Currency and Tariff Development Unit, Tees Esk & Wear Valleys NHS Foundation Trust, UK; **Adriana Castelli** , PhD, Centre for Health Economics, University of York, UK; **Paul A. Tiffin** , MD, PhD, Department of Health Sciences, University of York; Hull York Medical School; and Tees Esk & Wear Valleys NHS Foundation Trust, UK

**Correspondence:** Paula Maddison. Email: [paulamaddison1@nhs.net](mailto:paulamaddison1@nhs.net)

First received 19 Jun 2019, final revision 25 Jun 2020, accepted 5 Jul 2020

### Author contributions

All authors actively contributed to the following: substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; AND drafting the work or revising it critically for important intellectual content; AND final approval of the version to be published; AND agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

### Declaration of interest

None.

ICMJE forms are in the supplementary material, available online at <https://doi.org/10.1192/bjp.2020.142>.

### References

- 1 NHS England. *Five Year Forward View for Mental Health: One Year On*. NHS England, 2017.
- 2 NHS Improvement. *NHS Operational Productivity: Unwarranted Variations. Mental Health Services and Community Health Services*. NHS Improvement, 2018.
- 3 Jacobs R. *Investigating Patient Outcome Measures in Mental Health (Working Paper)*. University of York Centre for Health Economics, 2009.
- 4 Leamy M, Bird V, Boutillier C, Williams J, Slade M. Conceptual framework for personal recovery in mental health: systematic review and narrative synthesis. *Br J Psychiatry* 2011; **199**: 445–52.
- 5 Collins B. *Outcomes for Mental Health Services: What Really Matters?* King's Fund, 2019: p 6.