UNIVERSITY of York

This is a repository copy of *Developments in mannose-based treatments for Uropathogenic Escherichia coli induced urinary tract infections*.

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/165058/</u>

Version: Accepted Version

Article:

Hatton, Natasha, Baumann, Christoph George orcid.org/0000-0002-8818-972X and Fascione, Martin Anthony orcid.org/0000-0002-0066-4419 (2020) Developments in mannose-based treatments for Uropathogenic Escherichia coli induced urinary tract infections. Chembiochem. ISSN 1439-7633

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/

Developments in mannose-based treatments for Uropathogenic *Escherichia coli* induced urinary tract infections

Natasha E. Hatton^[a], Christoph G. Baumann*^[b] and Martin A. Fascione*^[a]

Abstract: During their lifetime almost half of women will experience a symptomatic urinary tract infection (UTI) with a further half experiencing a relapse within six months. Currently UTIs are treated with antibiotics, but increasing antibiotic resistance rates highlight the need for new treatments. Uropathogenic *Escherichia coli* (UPEC) is responsible for the majority of symptomatic UTI cases and thus has become a key pathological target. Adhesion of type one pilus subunit FimH at the surface of UPEC strains to mannose-saturated oligosaccharides located on the urothelium is critical to pathogenesis. Since the identification of FimH as a therapeutic target in the late 1980s, a substantial body of research has been generated focusing on the development of FimH-targeting mannose-based antiadhesion therapies. In this review we will discuss the design of different classes of these mannose-based compounds and their utility and potential as UPEC therapeutics.

1. Introduction

1.1. Structure of the urinary tract, urothelium and uroplakins The primary function of the urinary tract is to collect, transport, store and remove urine from the body, eliminating toxic waste products and metabolites generated by the kidneys.^[3] The urinary tract can be anatomically categorized into two subsections: the upper urinary tract (e.g. the kidneys and the ureters)^[4] and the lower urinary tract (e.g. the bladder and the urethra).^[4] The surface of the urinary tract is lined with a specialized epithelium known as urothelium.^[3] The biochemical and morphological features of the urothelium vary depending on its location within the urinary tract.^[3]

Generally, the urothelium is composed of three different layers; a basal cell layer attached to the basement membrane, an intermediate layer and an apical layer consisting of large hexagonal cells known as umbrella cells (Figure 1a).^[5] Umbrella cells are multinucleated, highly differentiated and accumulate a large amount of uroplakin proteins on their surface. This accumulation leads to the forming of a two-dimensional plaque,^{[3,} ^{6]} which acts as a barrier to water and other toxic materials in the urine.^[7] In humans there are four different uroplakin (UP) proteins: UPIa, UPIb, UPII and UPIIIa.^[8] These units come together to form a heterodimer (Figure 1b) with six of these heterodimers combining to form the uroplakin plaque (Figure 1c).^[5] UPIa and UPIb belong to the tetraspanin family,^[5] consisting structurally of four rod-like transmembrane domains. The first and second transmembrane domains are connected through a small extracellular loop, with a second large extracellular loop

[a]	Miss N. E. Hatton, Dr M. A. Fascione			
	York Structural Biology Lab, Department of Chemistry			
	University of York			
	York, YO10 5DD, UK			
	E-mail: martin.fascione@york.ac.uk			
[b]	Dr C. G. Baumann			
	Department of Biology, University of York			
	York, YO10 5DD, UK			
	Email: christoph.baumann@york.ac.uk			

Figure 1. a) Structure of the urothelium with basal cells attached to a basement membrane, an intermediate layer and a layer of umbrella cells. **b)** Structure of the uroplakin (UP) plaque consisting of six heterodimeric units with each unit composed of two dimers UPIa/II and UPIb/IIIa. **c)** Structure of the heterodimer units where the green ellipse represents a high mannose containing *N*-glycan and the grey ellipses represent complex *N*-glycans. **d)** Structure of UPIa and **e)** structure of UPIb. Both structures consist of four transmembrane domains and have an approximate molecular weight of 30 kDa.^[9] The green ellipses represent glucosamine residues and the yellow ellipses represent glactose residues.

connecting the third and fourth transmembrane domains. The main difference between the two uroplakins is UPIa contains a high mannose glycan attached to the second extracellular domain at residue Asn169 (Figure 1d). In comparison, UPIb contains a tetraantennary fucosylated complex glycan attached to the second extracellular domain at the Asn131 residue (Figure 1e).^[5] UPII and UPIIIa are each structurally composed of a single transmembrane domain.

1.2 Urinary tract infections

Urinary tract infection (UTI) is the third most common type of infection experienced by humans after respiratory and gastrointestinal infections.^[10] Most UTI cases affect the lower urinary tract^[11] and can be either symptomatic or asymptomatic,^[12] with symptoms including increased urination frequency, pain during urination and blood in the urine.^[13] If left untreated UTIs can result in kidney damage, allowing bacteria to enter the blood stream resulting in urosepsis. Urosepsis accounts for 5-7% of severe sepsis cases reported, with a mortality rate of between 25% and 60%.^[14]

Women are significantly more likely to experience a UTI than men^[15] due to the female urethrae being significantly shorter than a males (4 cm vs 20 cm).^[16] Approximately 40-50% of women will experience a symptomatic UTI within their lifetime, with over half of these women suffering a relapse within six months.^[17] Age is another well-recognized risk factor, with UTIs being the second most common form of infection in the non-institutionalized elderly population, accounting for 25% of

infection cases.^[15] Due to their weakened immune systems approximately 10% of males and 20% of females over the age of 80 suffer from an asymptomatic UTI.^[18] Other risk factors for UTIs include urinary catheterization and diabetes.^[19]

UTIs are caused by the invasion of foreign pathogens, with *Uropathogenic Escherichia coli* (UPEC) being responsible for 80% of cases. *Staphylococcus saprophytics* accounts for a further 10-15%, and the remaining cases are caused by *Klebsiella, Enterobacter*, and *Proteus* species.^[20] UTIs can be classed as uncomplicated or complicated. For a UTI to be classed as complicated the patient must also suffer from either an underlying illness such as diabetes, a structural malformation of the urinary tract, or an obstruction of urine flow.^[21] Complicated UTIs are generally more difficult to treat,^[20] meaning the infections are often chronic with several different Gram-positive and Gram-negative bacteria present.

Currently UTIs are treated with a course of antibiotic such as Nitrofurantoin or Trimethoprim.^[18] However, an increasing problem observed in the treatment of UTIs is antibiotic resistance - studies demonstrate UPEC strains contain over 30 different resistance genes to trimethoprim, with clinical resistance occurring in 16.7% of cases.^[22] Nitrofurantoin is still active against *E. coli*, with fewer cases of resistance being reported than with Trimethoprim. However, Nitrofurantoin has a higher incidence of significant side effects like pulmonary fibrosis,^[23] and it is predicted that resistance to both antibiotics will increase. The enduring challenge of antibiotic resistance means that researching new effective treatments for UPECinduced UTIs is a clinical priority.^[22]

1.3 Uropathogenic Escherichia coli pathogenesis pathway

UPEC is responsible for the majority of reported uncomplicated UTI cases,^[17] thus identifying new targets within UPEC could serve as the basis for developing new treatments for both acute and recurrent UTIs.

The six stages of UPEC pathogenesis are summarized in Figure 2.^[24] The bacteria initially colonize the periurethral areas and the urethra, travelling up the urethra while growing as planktonic cells in the urine. While in the urinary tract, UPEC interact with and adhere to the urothelium. Once adhered, UPEC grows on the surface of the umbrella cells of the urothelium forming a biofilm, facilitating invasion of the epithelial cells. Once within the umbrella cells UPEC can begin multiplying, forming an intracellular bacterial population (IBC); this allows for further formation of a quiescent intracellular reservoir (QIR).^[25] UPEC can then invade the intermediate layers of the urothelium and lay dormant. These bacteria are protected from antibiotic treatment, making them extremely difficult to eliminate and thus



Figure 2. The pathogenesis cycle for UPEC consists of six stages: stage 1) colonization of the periurethral areas and the urethra, stage 2) movement of UPEC up the urethra, stage 3) UPEC adherence, stage 4) biofilm formation, stage 5) epithelial cell invasion and formation of intracellular bacterial population, and stage 6) colonization of the urinary tract and kidneys by UPEC followed by entry into the blood stream.

the source of many recurrent infections.^[26] If untreated, UPEC will continue to colonize up the urinary tract, progressing to the kidneys.^[25] This colonization can result in kidney tissue damage

and provides UPEC access to the blood stream, resulting in urosepsis.

Invasion of the urothelium by UPEC occurs via a membrane zippering mechanism.^[27] This mechanism is stimulated by UPEC binding to the urothelium, which activates a complex signaling cascade, resulting in localized rearrangement of the urothelium actin cytoskeleton.^[27b] The cytoskeleton rearrangement leads to the envelopment and internalization of the bound UPEC (Figure 3). This complex signaling cascade has been shown to be reliant on many factors, such as focal adhesions; e.g. Src,^[28] phosphoinositide 3-kinase,^[27b] Rho-family GTPases; actin bundling and adaptor proteins, e.g. α -actinin and vinculin;^[27b, 29] lipid raft components, e.g. caveolin-1;^[30] and microtubules. Treatment of a host cell with a microtubule-disrupting agent, such as nocodazole or vinblastine, has been shown to inhibit host cell invasion by UPEC.^[31]





Adhesion of UPEC to the urothelium is mediated by UPEC binding to terminal D-mannose units on UPIa. Without adhesion to the sugar UPEC would remain free in the urine and be removed from the bladder during urination, preventing the initial UPEC infection from progressing into a symptomatic UTI. To bind to terminal mannose units UPEC produce multiple 3-micron long rod-like structures on their surface known as type 1 pili (Figure 4).^[32] The type 1 pilus consists of multiple different subunits, including repeating units of the FimA protein, which form a 7 nm-thick right-handed helical rod. This rod is joined to a 3 nm-thick distal tip fibrillum, which itself is composed of three further subunits: two adapter proteins; FimF and FimG; and an adhesion protein, FimH.^[32]



Figure 4. Structural organisation of the type 1 pilus which includes the following subunits: FimA (blue), FimC (gray), FimD (green rectangle), FimF (purple), FimG (green oval), and FimH (red). The position of the N-terminal domain (N, orange) and two C-terminal domains (C1, pink circle; C2, gray circle) relative to the transmembrane pore (green rectangle) in FimD are also indicated.

The exact process by which the type 1 pilus adheres to the urinary tract was unknown for a long time. One of the original theories proposed was that adhesion was mediated by the repeating FimA units which comprised most of the type one pilus.^[33] However, it is now recognized that the role of the repeating FimA units is to provide a structural scaffold.^[34] A wealth of research conducted in the late 1980s discovered that the minor subunits, FimF, FimG and FimH, were involved in adhesion.^{[35],[36]} Christiansen and co-workers found that recombinant bacteria lacking either all three subunits or just FimH displayed no ability to bind to erythrocytes, indicating that FimH plays a critical role in adhesion.^[37] Klemm and co-workers^[38] confirmed this observation, providing direct evidence that FimH is responsible for mannose-mediated adhesion.^[38]

1.4 FimH structure and catch bond mechanism

FimH is comprised of two domains (Figure 5). The first is a *C*-terminal pilin domain (FimH_{PD}), which attaches FimH to the pilus rod^[39] through the neighboring subunit FimG. This attachment occurs by donor strand complementation,^[40] and is an effective way of providing strong inter-molecular linkages through the donation of one β -strand from one subunit (e.g. FimG) into the β -sandwich of the neighboring subunit (e.g. FimH_P).^[41] The second FimH domain is the *N*-terminal lectin domain (FimH_{LD}), which contains a mannose-binding pocket^[39] which can bind to mannose sugars on the urothelium, mediating adhesion of UPEC to the urinary tract.^[32]



Figure 5. Crystal structure of FimH (red) with a heptylmannoside (green) ligand bound in the N-terminal lectin domain (FimH_{LD}) (PDB ID 4LOV).^[2] Residues with side chains shown as sticks (blue) are involved in non-covalent interactions with heptylmannoside, sodium ions (purple spheres) or water molecules (red spheres). The dashed lines (purple) denote short-range, non-covalent interactions.

The two-domain structure of FimH allows the type-1 pilus to form a catch-bond when bound.^[40] A catch-bond is a type of non-covalent interaction in which the dissociation lifetime of the bond increases when force is applied^[42] (Figure 6). Wirtz and coworkers provided the first unequivocal evidence for the existence of catch-bonds.^[43] Prior to this, catch-bonds were treated as one potential explanation for why the adhesion affinity of some bacteria (e.g. UPEC) increases in moderate shear flow. An alternative theory was the transport limiting binding model. This model, which has fallen out of favor, suggests that increased shear flow causes an increase in both dissociation and association rates, thus an enhancement in bacterial adhesion would have to be caused by an increase in bacterial association not a decrease in bacterial dissociation.^[44]

The catch-bond in FimH is biphasic, meaning under moderate force (such as that experienced by the sides of the urinary tract during urination) FimH binds to ligands with higher affinity, whereas exposure to higher shear flow leads to a decrease in binding affinity due to an increase in the dissociation rate for the FimH-ligand interaction.^[42] Moderate mechanical force induces the separation of the FimH_{LD} and FimH_P subunits (Figure 6). This separation switches the lectin domain from a low affinity state to a high affinity state^[45] reducing the rate of spontaneous ligand release, resulting in a 1000-fold higher affinity of FimH for mannose sugars under moderate flow



Figure 6. Catch bond mechanism for FimH binding to mannose in urethral lumen during a) no shear flow conditions where unbound and bound forms exist at equilibrium, and b) moderate shear flow conditions where the bound state is favored. Under moderate shear flow conditions, the hydrodynamic drag on the micron-sized UPEC bacterium (not to scale) results in a physical force on the tether which activates the catch bond mechanism in FimH. The higher affinity binding to mannose under these conditions prevents the UPEC bacterium from being flushed out of the urinary tract during urination.

conditions compared with static conditions.^[40] The relatively weaker affinity of FimH under static conditions favors invasion of UPEC along the urinary tract during static conditions, while the high affinity of FimH under moderate flow conditions enables UPEC to be retained in the urinary tract during urination. Glockshuber and co-workers^[40] investigated how the different flow conformation states affect the binding of FimH to *N*-glycans. A mixture of synthetic α -linked mono- and dimannosides were used to represent the natural terminal α -D-mannoside moieties present on FimH targeted glycoproteins in the bladder.^[40] Using these ligands, kinetic and structural characterization of the binding properties of FimH under both static and flow conditions could be investigated. It was demonstrated that the increased affinity of FimH under flow conditions compared to static conditions was ligand independent.^[40] Glockshuber and coworkers further found that dimannosides bound with higher affinity compared to monosaccharides, with the difference in affinity determined by the rate of spontaneous ligand dissociation.^[40] Under static conditions FimH binds to all natural terminal α-D-mannoside structures with medium affinity, while under flow conditions FimH binds to all D-mannosides at a 2000fold higher affinity with a 70,000-fold decrease in ligand dissociation rate and 30-fold increase in ligand association rate.^[40] Glockshuber and co-workers additionally found that even though FimH favors monovalent N-glycan binding when Nglycans are in short supply, each N-glycan can bind up to three FimH units at once.^[40]

1.5 Extended FimH binding pocket



Figure 7. A depiction of the main interactions that occur between mannose based pentasaccharide 1 and the extended FimH binding site.^[46] Red indicates interactions mediated by Van der Waals, aromatic stacking and hydrophobic interactions, blue indicates interactions mediated by hydrogen bonding.

The FimH lectin domain contains 157 amino acids, which assemble to form an 11-stranded β-barrel structure.^[39] Encompassed within the β -barrel structure is a polar binding pocket^[40] (residues Asn46, Asp47, Asp54, Gln133, Asn135 and Asp140) to which terminal mannose units can engage in a complex network of hydrogen bonding and electrostatic interactions.^[39] The polar binding pocket is surrounded by a hydrophobic region, (residues Phe1, Ile13 and Phe142)^[39] which contains a tyrosine gate (residues Tyr48, Ile52, Thr51 and Tyr137).[39] This region provides support for the binding site through electrostatic interactions with the tyrosine gate, which is shown to be influential in the ability of ligands to enter the binding site.^[46] Further interactive features of the FimH lectin domain are a small hydrophobic pocket^[47] adjacent to the sugar binding pocket (residues IIe52, Tyr137 and Asn138),^[39] a salt bridge (residues Arg98 and Glu50) which facilitates further hydrogen bonding,^[47] and the Tyr48 and Tyr137 residues, which partake in hydrophobic and ring stacking interactions^[47-48] as well as forming direct and water-mediated hydrogen bonds to ligands.^[48] A summary of the main interactions that occur between the FimH lectin domain and mannose-based pentasaccharide 1 is presented in Figure 7.

As depicted in Figure 7, the binding of a mannose-based ligand to the extended FimH binding site is the result of a complex network of interactions and hydrogen bonding. As such, small changes to the structure of the ligand can disrupt this binding network, resulting in either a decrease or complete loss of ligand affinity. Epimers of mannose (e.g. glucose and galactose) exhibit no inhibitory effects on FimH, [39, 49] due to the lack of an axial C-2 hydroxyl group preventing them from partaking in critical interactions with the polar binding pocket. The five-membered sugar fructose, which like mannose contains an axial C-2 hydroxyl group, has been shown to have weak activity for FimH. $^{\rm [49a]~[39,~49b]}$ A further stereochemical requirement exhibited by the FimH lectin domain is that the terminal mannose unit must be in the alpha orientation.[50] Terminal amannosides can engage in water-mediated hydrogen bonding within the FimH lectin domain;^[39] these interactions would not be possible if the mannoside was in the β -configuration. A further characteristic of the extended FimH binding site that facilitates critical interactions with ligands are its hydrophobic residues. These residues surround the mannose binding pocket and consists of the hydrophobic support and tyrosine gate residues as well as residue Thr51, which extends out of the hydrophobic

WILEY-VCH

ridge along a hydrophobic groove.[39] Unlike the polar binding pocket, the tyrosine gate structure is flexible, interacting with ligands by a mixture of m-stacking and van der Waals interactions.^[39] The flexibility of the tyrosine gate is the result of the Tyr48 residue being able to rotate, allowing for three different tyrosine gate conformations; open, closed and half open.^[51] In its unbound resting state FimH adopts an open conformation; here the side chain of the Tyr48 residue is positioned inwards facing residues Asp47 and Arg98 (Figure 8a).^[51] In the closed gate conformation the Tyr48 residue shifts alignment towards the Thr51 residue (Figure 8b).[52] If the side chain of the Tyr48 residue is aligned somewhere between the open and closed position, then the gate is classified as half open.^[51] When bound to a ligand the tyrosine gate can either remain in its open conformation or shift to the half open or closed conformation.^[51] A conformational change often accompanies ligand binding in order to minimize Gibbs free energy, thus different ligands will bind to different conformational states.^[51-52] When a ligand is flexible and can extend out of the binding site like some oligomannosides,^[46] the open conformation is adopted as this allows the second mannose residue in the oligosaccharide to interact with the Tyr48 residue of the binding site.^[47] When a ligand contains a sterically rigid aglycone group (e.g. 1,4-biaryl group) the tyrosine gate will shift to the closed conformation, positioning the aglycone residue outside of the tyrosine gate. This conformational shift is favored by the formation of strong π -stacking interactions between the aglycon and the outer side of the aromatic ring on the Tyr48 residue.[47, 53]



Figure 8. a) Structure of FimH lectin domain when the tyrosine gate as open PDB 4AV0^[51] b) Structure of the FimH lectin domain when the tyrosine gate is closed PDB 4AT1^[52]. Note the movement of the Tyr48 residue (blue) from facing the Asp47 residue (yellow) in the open conformer to facing the Thr51 residue (pink) in the closed conformer. The Arg98 residue (green) plays a role in stabilizing the surface loop conformation on which the Thr51 residue is located.

2. Mannose based FimH inhibitors

This review will survey and critically assess recent progress in the development of FimH targeting mannose-based analogs with potential applications against UPEC-induced UTIs. The review will focus on three categories of mannose-based FimH inhibitors: oligosaccharide-based FimH inhibitors (section 2.1), α -D-mannopyranoside-based inhibitors (section 2.2.) and polyvalent mannose-based inhibitors (section 2.3). Note that informative reviews on this subject have previously been published. $^{[39,\ 54]}$

2.1 Oligosaccharide based FimH inhibitors

FimH primarily targets large branched mannose-capped oligosaccharides found on *N*-glycosylated proteins which line the urinary tract^[39] such as UPIa.^[55] The preference of FimH for mannose-capped oligosaccharides can be seen in ligand affinity assays, with α -D-mannose-capped oligosaccharides displaying 100-200 times greater affinity for FimH than α -D-mannose monomers.^[39] This increase in affinity is due to mannose-based oligosaccharides being able to interact with the extended FimH binding site.

Due to their role as natural FimH ligands and their ability to act as FimH inhibitors, much research has been performed into using mannose-capped oligosaccharides as a curative treatment for UPEC-induced UTIs. Sharon and co-workers^[56] investigated the ability of a large range of mannose glycosides and mannosecapped oligosaccharides to inhibit E. coli-induced yeast aggregation.^[56] The three most potent inhibitors were oligosaccharides 2, 3 and trisaccharide 4 (Figure 9),^[56] with research suggesting that trisaccharide 4 is the optimal size for binding to the extended FimH binding site.^[56] Further investigation found that the high binding affinity displayed by trisaccharide 4 was due to a few key structural components. Firstly, trisaccharide 4 contains a Man-a1,3-Man linkage at the non-reducing terminus; this linkage has been shown to be highly preferential in FimH binding.^[57] A second beneficial structural feature is the presence of a N-acetyl glucosamine sugar, as neither mannotriose 5 nor mannopentaose 6 (Figure 9) exhibited significantly higher FimH binding affinity than Man-α1,3-Man 7 (Figure 9).^[57] Furthermore, docking studies suggest that the structure of trisaccharide 4 allows both the central mannose unit and the GlcNAc unit to interact with the tyrosine gate, accounting for the high affinity displayed by trisaccharide 4.[®]



Figure 9. Compares the structures of potent oligosaccharide based FimH inhibitors (oligosaccharides 2, 3, and trisaccharide 4) to weak oligosaccharide based FimH inhibitors (mannotriose 5, mannopentaose 6 and Manα-1,3Man 7).

A further oligosaccharide with high affinity to FimH is oligosaccharide **1** (Figure 10, K_d = 20 nM)^[57]), displaying a 10-fold increase in affinity compared to Man- α 1,3-Man.^[46, 57] Oligosaccharide **1** contains a terminal chitobiose unit; these units are ubiquitous in nature, providing a bridge between mannosides and asparagine residues in the Asn-X-Ser/Thr motif found in *N*-linked glycoproteins.^[46] Chitobiose units have been shown to interact with the extended FimH binding site. A crystal structure of oligomannoside **1** bound to the extended FimH

receptor binding domain identified some of the key binding interactions.^[46] Firstly, Man4 at the non-reducing end was shown to be anchored into the polar binding pocket. Further interactions were mediated through the tyrosine gate interacting with the Man- α 1,3-Man- β 1,4-GlcNAc backbone via the α 1,3 and the first *β*1,4 glycosidic linkages. The second mannose unit at the non-reducing end, Man5, was not directly recognized by FimH due to the unit partially extending out of the binding site. GlcNAc1 was also shown to partially exit the binding site, folding over residue Thr51. Man5 and GlcNAc1 displayed the most flexibility of all the saccharide units,^[46] likely due to being the least closely bound to the extended FimH binding site. This crystal structure provides further evidence that trisaccharide 4 may construct the optimal motif for FimH binding, containing the critical Man-a1,3-Man-B1,4-GlcNAc backbone needed for interactions with the extended FimH binding site.



Figure 10. Structure of the potent FimH inhibitor oligomannoside 1.

The oligomannoside structures discussed here were obtained by a number of different methods including chemical synthesis, isolation from yeast extract and isolation from the urine of patients suffering from mannosidosis and GM1 gangliosidosis.⁶⁶

Chemical synthesis is one of the most common methods used to obtain mannose-capped oligosaccharides. As is common in carbohydrate synthesis, the synthesis of these oligosaccharides relies on a complex scheme of protection, glycosylation and deprotection steps. To further add to this complexity, potent FimH inhibitors such as trisaccharide 4 and oligosaccharide 1 contain a critical β-mannosidic bond between the central mannose unit and the GlcNAc unit. β-Mannosidic bonds have long been recognized as the most difficult linkages to construct for carbohydrate chemists.^[58] Over the last 25 vears multiple groups have conducted a wealth of research into the development of β -mannosidic bonds, developing synthetic methods to afford β-mannosides in a high yielding and selective manner.^[58-59] Despite this research, the requirement for preactivation and need for specific mannoside donor protection groups (e.g. O-4,6 benzylidene protection)^[60] still makes β mannoside synthesis challenging, and consequently many opt to prepare simpler *a*-mannoside counterparts. A further factor restricting the use of mannose capped oligosaccharides as therapeutics is their size which makes them unlikely to be orally absorbed. Lack of oral absorption coupled with synthetic complexity, has therefore restricted the application of mannosecapped oligosaccharides, and as such many research groups have chosen to focus on the more easily synthesized α -Dmannopyranoside based inhibitors as FimH antagonists.

2.2 α -D-Mannopyranoside based inhibitors

One of the first classes of D-mannopyranoside based FimH inhibitors were alkyl mannosides (Figure 11, Table 1 entry 1). These were discovered serendipitously when De Greve and coworkers found butyl- α -D-mannoside occupied the mannose binding site.^[61] Investigations found butyl- α -D-mannoside (K_d =

151 nM) to bind to FimH with 15-30 times greater affinity than α-D-mannose (K_d = 2.3 μM).^[61] Further investigation into the binding affinity of alkyl mannosides found that in general binding affinity increases as alkyl chain length increases, with the binding affinity peaking for hept-D-mannoside (K_d = 5 nM).^[61] One theory explaining this trend is that increasing the chain length increases the van der Waals interactions with the hydrophobic groove and tyrosine gate, but once the alkyl chain is longer than a heptyl chain it extends beyond the hydrophobic region into solvent-exposed areas,^[39] increasing the free energy of binding and thus lowering binding affinity.



Figure 11. Structure of the alkyl mannoside scaffold where n = 0-7.

The alkyl mannosides used by De Greve and coworkers^[61] were synthesized using a procedure reported by Tidén and co-workers^[62] for the synthesis of octyl and tetradecyl mannosides. Here alkyl alcohols underwent silver triflatepromoted glycosylations to 2,3,4,6-tetra-*O*-benzoyl- α -Dmannopyranosyl bromide (Scheme 1), followed by Zemplén deacylation to afford the deprotected alkyl mannoside ligand.



Scheme 1. Reaction scheme for the synthesis of alkyl mannosides, by glycosylation with aliphatic alcohols using an AgOTf activation system.

Sharon and co-workers^[63] reported aromatic α -mannosides (Table 1 entry 2) to be powerful inhibitors of the adherence of type 1 fimbriated E. coli to both yeast and intestinal epithelial cells. With p-nitrophenyl- α -mannoside 8 (Figure 12) shown to be approximately 70 times more effective than methyl a-mannoside for both inhibiting yeast agglutination by E. coli 025 and inhibiting the adherence of E. coli 0128 to guinea pig ileal epithelial cells.^[56a, 63] The two most potent aromatic α-mannoside inhibitors reported by Sharon and co-workers^[56a, 63] were *p*-nitro-O-chlorophenyl-α-mannoside (pNoClPαMan) 9 and 4methylumbelliferyl-α-mannoside (MeUmbαMan) 10 (Figure 12). These analogs were shown to increase the inhibition of yeast agglutination by E. coli 025 by a factor of 717 (pNoCIPaMan 9) and 600 (MeUmbaMan 10) compared to methyl a-mannoside. A significant increase in inhibition of E. coli 0128 adherence to guinea pig ileal epithelial cells was also observed [470 times for (pNoCIPaMan 9) and 1015 times for (MeUmbaMan 10) with respect to methyl a-mannoside]. This increase in affinity displayed by FimH for aromatic a-mannosides compared with methyl a-mannoside is likely due to the aromatic side chain being able to establish favorable π - π stacking interactions with the tyrosine gate.[64]



Figure 12. Structure of p-nitrophenyl-α-mannoside **8**, p-nitro-O-chlorophenylα-mannoside (pNoClPαMan) **9** and 4-methylumbelliferyl-α-mannoside (MeUmbαMan) **10**.

Building upon the high potency displayed by *p*-nitrophenylα-mannoside, Lindhorst and co-workers synthesized a series of *para*-substituted aryl α-D-mannosides, with one analog achieving significantly higher potency than *p*-nitrophenyl-α-mannoside [relative inhibitory potential (RIP) of 1.6 vs 1].^[65] Further work into the development of an aromatic α-mannoside based inhibitor was performed by Han and co-workers.^[47] Here they synthesized a series of aryl-substituted α-D-phenylmannosides, evaluating their biological activity using a guinea pig red blood cell based hemagglutination (HA) assay. This assay measures the concentration of inhibitor needed for >90% HA inhibition, and this concentration is known as the hemagglutination inhibition (HAI) titre. The HAI titre of new compounds can be compared to that of other compounds in order to assess the potency of new compounds.

Han and co-workers synthesized an extensive series of aryl substituted α -D-phenylmannosides functionalized with a variety of groups (Cl, NO₂, CN, OMe, CO₂Me, NHAc, CO₂H, CONH₂, CH₂CO₃Me) at either the *ortho*, *meta* or *para* position.^[47] Generally, it was noted that phenylmannosides functionalized at the *ortho* and *meta* positions gave better potency than phenylmannosides functionalized at the *para* position. The most potent phenylmannoside contained a methyl ester group at both *meta* positions, being three times more potent than the phenylmannoside containing a single *meta* methyl ester.^[47]

Another well-investigated group of FimH-targeting analogs are biphenyl mannosides (Table 1 entry 3).^[47, 53, 66] Han and coworkers hypothesized that the potency of aryl mannosides could be increased by the addition of a second aryl ring. This hypothesis was explored using analogs containing an additional aryl ring system in conjugation with the parent ring of the aryl mannoside. The most potent analog was biphenyl mannoside **11** (HAI = 1 μ M, IC₅₀ = 0.94 μ M)^[47] which contained a methyl ester at the *meta* position of the second ring system (Figure 13). A high resolution X-ray crystal structure of biphenyl mannoside **11**, showed that both aromatic rings were able to partake in hydrophobic and π - π stacking interactions with the closed Tyr48 residue, with the methyl ester further able to hydrogen bond with the salt bridge (residues Arg98 and Glu50).^[47]



Figure 13. Structure of biphenyl mannoside 11.

Further SAR evaluation of the terminal biphenyl mannoside aryl ring identified two potent biphenyl mannosides: biphenyl mannoside **12** (HAI = 0.15 μ M) and biphenyl mannoside **13** (HAI = 0.37 μ M, IC₅₀ = 0.74 μ M)^[66a] (Figure 14).^[47] Biphenyl mannoside **12** displayed good potency in the HAI assay, yet due to its insolubility could not be used in a biofilm assay.^[66b] However, biphenyl mannoside **13** was used in a biofilm assay and showed impressive activity, with an IC₅₀ of 0.74 μ M.^[66b] Initial dosing in mice established biphenyl mannoside **13** to be orally bioavailable and stable to metabolism, and the only metabolic degradation pathway detected was hydrolysis of the glycosidic bond. Metabolic degradation of biphenyl mannoside **13** should be minimal as >95% was excreted in the urine unchanged. Moreover, no apparent toxicity was observed up to a dose of 200 mg/kg, using physiological

changes and survival as assessment parameters. Further investigation using an adapted preclinical murine model suggested that biphenyl mannoside **13** was effective at preventing infections, treating established UTIs and increasing the activity of antibiotic treatments.^[66b]



Figure 14. Structure of biphenyl mannoside 12 and biphenyl mannoside 13.

co-workers Han and previous work on α-Dphenylmannosides showed that substitution at the ortho position with chloro and cyano groups could significantly increase potency.^[47] Using widely available biphenyl mannoside 11 as a scaffold (Figure 15), Han and co-workers also synthesized a number of biphenyl mannosides substituted at the ortho position of the top ring, and observed the general potency trend $CF_3 > CI$ = Me > OMe > F.^[66a] One explanation for the increase in potency upon ortho substitution is that the presence of ortho substituents increases the hydrophobic contact with the tyrosine gate at residues Ile52 or Ile13. This explanation is supported by the potency trend seen above, as addition of a CF₃ group would result in the largest increase in hydrophobic contact and thus the largest increase in potency. Further investigation was performed using biphenyl mannoside **14** (IC₅₀ = 1.35 μ M)^[66a] as a scaffold, with a similar potency trend observed: CF₃ > Me > Cl.^[66a] Two final analogs were synthesized; biphenyl mannoside 15 (HAI = 0.01 μ M, IC₅₀ = 0.043 μ M)^[66a] and biphenyl mannoside **16** (HAI = 0.02 μ M, IC₅₀ = 0.073 μ M),^[66a], with both displaying increased potency to biphenyl mannoside **13** (HAI = 0.37 μ M).^[66a]



Figure 15. Structure of biphenyl mannoside 11, biphenyl mannoside 14, biphenyl mannoside 15 and biphenyl mannoside 16 where R= CF₃, Cl, Me, OMe, F and R' = CF₃, Me, Cl.

Biphenyl mannosides 15 and 16 displayed good potency, but their low log(p) values of -6.27 and -8.46, respectively, suggest they would have poor oral absorption.[66a] Biphenyl mannoside 17 (Figure 16) had a significantly higher log(p) of -3.89, and absorption.[66a] therefore should have better oral Pharmacokinetic studies in mice showed biphenyl mannoside 17 $(IC_{50} = 0.16 \ \mu M)^{[66a]}$ to be a promising lead candidate, maintaining a concentration in the urine and plasma well above the predicted minimum effective concentration for over 6 h. Further in vivo tests used a chronic infection mouse model; this model uses C3H/HeN mice with chronic cystitis at two-weeks post-infection. The efficacy of compounds can be monitored by the number of colony forming units (CFU) present at selective time periods after compound administration.[66b] Mice treated with biphenyl mannoside 17 showed a significant reduction of

chronic cystitis six hours post-treatment; however, by 24 hours post-treatment the number of CFU had started to increase, but remained lower than in control mice.^[66b] This CFU increase was shown to be prevented by administering three doses of biphenyl mannoside **17** every eight hours. These *in vivo* experiments suggest that biphenyl mannoside **17** provided a promising lead compound which could be used in preclinical trials.^[66b]



Figure 16. Structure of biphenyl mannoside 17.

A limitation of the antagonistic studies performed on previously designed biphenylmannosides and other α -Dmannopyranoside based inhibitors are the methods used; e.g. a fluorescence polarization assay, an isothermal titration calorimetry (ITC) based assay, and a surface plasmon resonance (SPR)-based assay. These methods rely on the use of $\mathsf{Fim}\mathsf{H}_{\mathsf{LD}},^{[1a]}$ which is locked in the high-affinity $\mathsf{Fim}\mathsf{H}$ conformation.^[2] As discussed previously, the FimH lectin can adopt three conformational states; an unbound low-affinity conformation, a bound medium-affinity conformation which occurs under static conditions, and a bound high-affinity conformation which occurs under shear force.^[1a] Therefore, only measuring an antagonist affinity against one FimH conformation could limit the accuracy of the results. Antagonistic studies performed using most target-based assays and cell based assays (e.g. hemagglutinin assays or a flow cytometry based assay) are not affected by these inaccuracies, as they use E. coli cells producing $FimH_{FL}$, which can adopt all three conformers.^[1a] Monomeric FimH_{FL} is inherently unstable^[1a, 67] and thus a native model must be used instead. This model is



Figure 17. Structure of some substituted biphenyl mannoside analogs with promising FimH activity and pharmacokinetic properties,^[1] such as dissociation constants (K_D) and logP_e values assessed using a parallel artificial membrane permeability (PAMP) assay.

PAMPA LogP_e [cm/s] = -5.1 ± 0.1

PAMPA LogP_e [cm/s] = -5.0 ± 0.1

composed of FimH expressed as a biomolecular complex, where the incomplete fold of FimH_{PD} is complemented by a synthetic donor-strand peptide. Subsequently, comparison of the binding affinities of a series of biphenyl analogs showed an approximate 100-fold decrease in affinity from FimH_{LD} to FimH_{FL}. Recent investigations into biphenyl mannoside affinity for FimH_{LD} and FimH_{FL} has led to the design of some promising substituted biphenyl mannoside analogs (Figure 17).^[11] These analogs display nanomolar potency against FimH_{FL} and subnanomolar potencies against FimH_{LD}, alongside promising pharmacokinetic properties.^[11]

The ortho-substituted biphenyl mannosides were synthesized via two routes (Scheme 2). Most analogs were synthesized using Pathway A. The first step of this pathway is Lewis acid mediated glycosylation of mannose pentaacetate with a 2-substituted 4-bromophenol analog (step a). This is followed by a Suzuki cross-coupling with a commercially available 3-substituted phenylboronic acid derivative to give a protected ortho-substituted 4'-biphenyl mannoside (step b). The final step is a Zemplén deacetylation, affording the deprotected ortho-substituted 4'-biphenyl mannoside (step c). Synthesis of the diamide analogs was performed using Pathway B, following the same initial and final steps as Pathway A (step a and d), but using a 3,5-di-(N-methylaminocarbonyl)-phenylboronic acid pinacol ester synthesized in-house (Pathway C) in the Suzuki cross-coupling reaction (step d).[66a]



A further class of D-mannopyranoside based inhibitors are squarate mannosides (Table 1 entry 4). Lindhorst and coworkers used computer-based docking methods to predict the binding affinities of a number of FimH inhibitors; e.g. methyl mannose, p-nitro-phenyl-a-mannoside and two squarate mannosides (squarate mannoside 18 and squarate mannoside 19, Figure 18). An enzyme-linked immunosorbent assay (ELISA) was performed to test the inhibitors ability to inhibit type 1 fimbriae-mediated bacterial adhesion, with methyl mannose used as a standard (RIP = 1). Squarate mannosides 18 and 19 were shown to be the most potent inhibitors in the ELISA, with RIPs of 1600 and 6900, respectively.^[64] For the inhibitors analyzed the expected inverse trend between docking score and RIP was observed. One notable exception was squarate mannoside 18, which had a predicted FimH affinity higher than that of squarate mannoside 19. However, the ELISA showed squarate mannoside 19 to be 4 times more potent that squarate mannoside 18.[64]



Figure 18. Structure of squarate mannoside 18 and squarate mannoside 19.

Following the discovery of squarate mannosides as potent FimH inhibitors^[64] it was suggested that the high potency observed with these ligands could be the result of covalent bond formation between the squarate mannoside **18** and the FimH binding site (Scheme 3). To see if this type of reaction was possible, squarate mannoside **18** was reacted with a L-phenylalanine ester under physiological conditions. The reaction was shown to be successful.^[68]



Scheme 3. Reaction scheme for proposed covalent bond formation between squarate mannoside 18 and the *N*-terminus of the FimH lectin (residue Phe1).

However, diamide squarate mannoside **20** (Figure 19) was also synthesized and its structure prevents covalent cross-linking to FimH, yet was shown to be a potent inhibitor of FimH (IC₅₀ = 6.38 μ M ± 3.7 compared to nitrophenol IC₅₀ =274 μ M ± 110). This suggests that the potency of squarate mannosides cannot be attributed to covalent cross-linking with the FimH binding site.^[68]



Figure 19. Structure of diamide squarate mannoside 20.

Septanoses (Table 1 entry 5) were introduced by Ernst and co-workers^[69] following the observation that methyl βseptanosides bound to the jackbean lectin concanavalin A (ConA) (another mannose-selective lectin).^[69] Ernst and coworkers^[69] used a competitive binding assay and ITC experiments to assess the potency of multiple septanose analogs on FimH_{LD}. The results of these experiments showed 2-O-n-heptyl-1,6-anhydro-D-glycero-D-galactitol 21 (Figure 20) displayed a ten-fold lower potency for FimH than n-heptyl α -Dmannopyranoside 22 (Figure 20) (IC₅₀ = $1.37 \pm 0.3 \,\mu$ M vs 0.064 \pm 0.02 µM, K_D = 0.26 µM vs 0.029 µM).^[69] Further investigations showed that while 2-O-n-heptyl-1,6-anhydro-Dglycero-D-galactitol 21 (Figure 20) establishes the same hydrogen bonding network at the FimH binding site as *n*-heptyl α -D-mannopyranoside **22** (Figure 20),^[69] the formation of this network results in a loss of conformational flexibility, causing a loss of entropy.[69]



Figure 20. Structure, IC₅₀ values and K_D values of 2-O-*n*-heptyl-1,6-anhydro-D-glycero-D-galactitol 21 and n-heptyl α -D-mannopyranoside 22⁽⁶⁹⁾

Gouin and co-workers developed a further class of Dmannopyranoside based inhibitors: thiazolylaminomannosides (TazMan) and neothiazolylaminomannosides (NeoTazMan) (Table 1 entry 6).^[70] A series of TazMan analogs were originally developed based on scaffold 1 (Figure 21); with the most potent analog, thiazolylaminomannoside 23, being 100-times more potent at preventing adherent-invasive Escherichia coli (AIEC) attaching to the intestinal cells compared to n-heptyl a-D-21).^[70b] mannopyranoside 22 (Figure Despite thiazolylaminomannoside 23 displaying high in vitro potency, it showed limited in vivo efficiency, likely due to having low pH stability and low water solubility; this restricts the applications of thiazolylaminomannoside 23.[70b] To combat these poor in vivo results, Gouin and co-workers synthesized a second neothiazolylaminomannoside (NeoTazMan) series based on scaffold **2** (Figure 21).^[70a] Thiazolylaminomannoside **23** NeoTazMans counterpart neothiazolylaminomannoside 24 (Figure 21) was shown to have improved in vivo properties, being stable to both enzymatic and acid hydrolysis. However, neothiazolvlaminomannoside 24 was shown to be 2.8 times less potent than thiazolylaminomannoside 23 with an IC_{50} of 194 nM vs 70 nM.^[70a] These initial investigations show that both TazMan and NeoTazMan are effective FimH inhibitors. However, it is unknown how applicable these analogs will be as therapeutics in the treatment of UPEC-induced UTIs, as existing studies have only explored their applications in the treatment of Crohn's Disease.^[70]



Figure 21. Structure of thiazolylaminomannosides scaffold 1 and neothiazolylaminomannoside scaffold 2 and structure and IC_{50} values of thiazolylaminomannosides 23 and neothiazolylaminomannoside 24.^[70a]

The final class of α -D-mannopyranoside based inhibitors discussed in this review are indolynylmannosides (Table 1 entry 7), which were first investigated by Ernst and co-workers.^[71] Ernst and co-workers synthesized a series of indolylphenyl and indolinylphenyl α -D-mannosides and then investigated their inhibitory and pharmacokinetic properties. The most promising analog was indolinylphenyl **25** (Figure 22) achieving an IC₅₀ of 20 nM.^[71] Administering a low dose of indolinylphenyl **25** (1 mg / kg) in a mouse model achieved a minimum inhibitory concentration in the bladder for > 8 h.^[71] Furthermore, a 1 mg / kg dose of indolinylphenyl **25** was shown to reduce colony-forming units in the bladder by a factor of 3.7 compared to untreated mice; this result is in line with the reduction seen in mice treated with ciprofloxacin (8 mg / kg).^[71]



Figure 22. Structure and IC₅₀ value of indolinylphenyl 25.^[71]

A factor that must be considered when designing mannose-based inhibitors is selectivity, as humans possess other mannose binding lectins (e.g. human mannose binding proteins) meaning a lack of selectivity can result in off-target reactions. Ernst and co-workers^[72] have previously investigated the selectivity of five potent FimH antagonists (all with nM IC₅₀ values (Figure 23) against eight human mannose receptors. If an analogue displayed 10⁵ times greater potency for FimH over the mannose binding proteins it was classified as selective, and assumed to not cause adverse effects due to non-selective binding.^[72] All inhibitors tested were shown to have a 10⁵ fold lower affinity for the human mannose binding proteins than FimH, confirming their selectivity.^[72] The FimH inhibitors have been optimized to contain hydrophobic substituents at their reducing end; these substituents can interact with the tyrosine gate

located at the entrance of the FimH binding site. The tyrosine gate is a feature unique to FimH, which likely explains the selectivity displayed by these inhibitors.^[72] Furthermore, multivalent ligand presentation is known to be hugely important in nature, and as the inhibitors investigated here are only capable of monovalent binding they likely only display low affinity for human mannose receptors.^[72]



Figure 23. Structures of the α -D-mannopyranoside based inhibitors, n-heptyl α -D-mannopyranoside 22, biphenyl α -D-mannopyranoside derivatives 26 and 27, indolylphenyl mannoside derivative 28 and squarate mannosides derivative 29.

Table 1. A comparison of the $\alpha\mbox{-}\mbox{-}\mbox{-}\mbox{mannopyranoside}$ based inhibitors discussed.





2.3. Use of polyvalent mannose scaffolds in FimH

In nature carbohydrate ligand presentation is mainly multivalent,^[72] suggesting that the potency of α -D-mannopyranoside based inhibitors may be limited due to only achieving monovalent targeting. The phenomenon describing how many multivalent ligands display greater affinity than their



Figure 24. Diagram showing two potential mechanisms whereby multivalent ligands can increase apparent binding affinity. a) Clustering effect where a multivalent ligand binds to one receptor initially and then captures additional receptors as they diffuse into close proximity resulting in clustering of the ligand -bound receptors. b) Induced aggregation whereby multivalent ligands bind to multiple lectins on different bacterial units cross linking them together.

WILEY-VCH

monovalent counterparts has been termed the "cluster glycoside effect".^[73] Multivalent ligand presentation can increase lectin binding via multiple factors, including receptor clustering and cross-linking.^[74] Receptor clustering occurs when monovalent lectins / ligands are anchored to the cell membrane. The presence of a multivalent binding species can capture receptors freely diffusing in the membrane, thus inducing receptor clustering (Figure 24a). However, this mechanism is unlikely to occur in the bacterial outer membrane (OM) due to the restricted lateral diffusion of OM proteins.^[75] Receptor clustering may occur by default in the bacterial OM near sites of beta-barrel transmembrane protein insertion (i.e. BAM complex will mediate insertion of FimD) due to the restricted lateral diffusion. As a consequence, the transmembrane FimD usher protein may already be present at the extracellular surface in clusters.^[74] Multivalent ligand presentation can further increase lectin binding through cross-linking. Here, individual ligands on a multivalent species bind to lectins on separate target units, cross-linking them and aggregating bacteria together (Figure 24b).^[74] Due to the complexity of multivalent carbohydrate interactions, it is often an accumulation of multiple factors that cause the "cluster glycoside effect". In the case of FimH it is likely that multivalent mannose-based ligands bind to multiple different FimH units, resulting in the formation of E. coli clusters. Multivalent glycomaterials attached to filters have been used to aggregate E. coli and remove them from solution.[76]

There has been substantial research into the development of FimH-targeting polyvalent mannose scaffolds, with applications in the treatment of UPEC-induced UTIs. Multiple different polyvalent mannose scaffolds have been investigated with three of the major ones being multimeric heptylmannosides,^[77] glycoclusters^[78] and dendrimers.^[79]

One of the most researched polyvalent scaffolds are multimeric heptylmannosides (Table 2); these aim to build upon the high-inhibitory potency displayed by heptyl α-D-mannoside (HM) (Figure 5). Gouin and co-workers synthesized an initial series of multivalent glycoconjugates (Table 2 entry 1) based on a 1,1,1-tris(hydroxymethyl)ethane or pentaerythritol core and measured their inhibitory potencies using a HAI-assay and a bladder binding assay (BBA) with human bladder cell line 5637.^[77a] Both the HAI titre and the BBA data showed a positive correlation between potency and valency.^[77a] A second multivalent HM glycoconjugates series (Figure 25, Table 2 entry 2) was designed based on a carbohydrate core, with one heptamannoside analog tethered to a ring-opened βcvclodextrin.^[77b] Binding affinity of this series was examined using HAI titre and ITC measurements. As with the previous series, HAI titre measurements showed a positive correlation between valency and potency, with the heptavalent opened β cyclodextrin glycoconjugate achieving a titre in the nanomolar region (60 nM).^[77b] Though initial ITC measurements supported this trend, latter measurements varied, likely due to increased calorimetry noise reducing the accuracy of the K_d measurements.^[77b] Two further heptavalent β-cyclodextrinlinked HM glycoconjugates (Table 2 entry 3) were synthesized using two different spacer lengths. Their biophysical properties were assessed using ITC, extracting K_d values and molar ratios. Both heptavalent β-cyclodextrin-linked glycoconjugates were significantly more potent than their monovalent counterparts, with K_d values in the nanomolar range.^[77c] Reverse titration measurements showed that the use of a shorter spacer unit led to superior potency (K_d = $2.9 \text{ nM} \pm 0.03$)

compared to the use of a longer linker (K_d = 33.0 nM ± 6.6), despite only achieving partial binding occupancy (molar ratios = 3.01 ± 0.03 for the short spacer and = 7.7 ± 0.06 for the longer spacer).^[77c] This suggests potency is not solely dependent on valency. The *in vivo* properties of both heptavalent β-cyclodextrin-linked glycoconjugates were assessed using a murine cystitis model with C3H/HeN mice. Both heptavalent glycoconjugates were shown to be 100 times more potent than their monovalent counterparts.^[77c]



Figure 25. Structure of multivalent HM glycoconjugate series designed using a carbohydrate core where n= 0, 1,2 and 6 (when using β -cyclodextrin core).

Further work into development of a multivalent heptylmannoside has investigated the design and use of glyconanoparticles ^[77e] (Table 2 entry 4) and multimeric heptylmannosides^[77d] (Table 2 entry 5) in targeting adherent invasive *Escherichia coli* (AIEC), a bacterial strain present in the ileal lesion of Crohn's disease patients.^[77d]

Table 2. Summary of the multivalent heptamannoside structure discussed. Entry Core unit Reference [77a] 1) Backbone Link n =1 - 4 [77b] 2) n = 0, 1, 2 and 6 [77c] 3) -Cyclodextr core n = 1, 6 or 7 [77e] 4)

5) [77]

Core structure prior to erization-induced self assembly

Mannose-based glycoclusters are a further polyvalent mannose scaffold that has been investigated. Multiple different glycocluster series have been synthesized using a variety of WILEY-VCH

different backbone scaffolds (Table 3); e.g. cysteine residues (Table 3 entry 1),^[78a] functionalized pentaerythritol (Table 3 , entry 2),^[78b] thiourea-bridged clusters (Table 3, entry 4),^[80] and carbohydrate centered clusters (Table 3, entry 5).^[78d] The *in vitro* efficacy of different glycocluster series have been assessed using a variety of methods; e.g. IC₅₀ measurements, inhibition of baker's yeast agglutination assay, HAI titre, and ELISA. Generally, a positive correlation between potency and valency was observed. However, other factors such as scaffolds structure (e.g. inclusion of a phenyl unit in the scaffold) were shown to significantly contribute to potency.

 Table 3. Summary of the organic scaffold used in the synthesis of mannosebased glycoclusters.

Entry	Organic scaffold	Structure	Reference
1	Cysteine	H ₂ N OH	[78a]
2	Azide or alkyne bearing pentaerythritols	N_3	(786)
3	Thiourea- bridged clusters	$\overset{S}{\overset{R_1 \searrow \overset{N}{}}{}_{R_2} \overset{R_3}{}_{R_4}}$	[78c, 80]
4	Peptide-bridged clusters	O ₂ N OH OH	[80]
5	Carbohydrate centred cluster		[78d]

Investigations have also focused on the synthesis of a mannose-bearing dendrimer.^[79] Roy and co-workers synthesized multiple series of mannose-bearing dendrimers using different scaffolds. One scaffold of particular note was the α -amino-L-lysine scaffold, from which they synthesized a dendrimer containing 16 mannose units with a potency 500 times greater than methyl mannose (HAI titre = 1.0 μ M vs 500 μ M).^[79b]

Diamond nanoparticles can also serve as a polyvalent mannose scaffold. Mannose-functionalized diamond nanoparticles where originally applied to the detection and removal of *E. coli* from bacteria-polluted water.^[76b] More recently mannose-functionalized diamond nanoparticles have been demonstrated to be potent *E. coli* anti-adhesives, displaying impressive potency in a bladder cell adhesion assay (RIP = 9259 vs 1 for methyl mannose)^[81] as well as reducing *E. coli* biofilm formation.^[81]

A final scaffold discussed here is fullerenes. Coppercatalysed click chemistry was used to functionalize fullerenes with mannose moieties,^[82] yielding three fullerene analogs bearing 12 mannoside units.^[83] ITC and SPR showed these polyvalent analogs to be capable of multivalent binding and found that both polyvalent and monovalent analogs displayed a K_d in the nanomolar range.^[83] A further HAI- assay was performed which found the polymeric fullerenes to be between 2.8-30 times more potent than their monomeric counterparts.

Other polyvalent mannose scaffolds reported include glycopeptides,^[79d, 84] polyvalent nanoparticles^[85] and pillar[5]arene derivative.^[86] Research into these scaffolds is in its infancy but initial *in vitro* investigations appear promising.

Oral administration is the preferred route for drug administration, and as such the applications of any analogue which cannot be orally administered are limited. Due to their large structure and high hydrogen bond donor and acceptor content, polyvalent mannosides are unlikely to be orally absorbed. This could severely limit their applications in the treatment of UTIs but also makes them potentially powerful candidates in the treatment of AIEC-induced Crohn's disease. ^{[77e] [77d]} Indeed clinical trials are currently being performed on the use of multivalent inhibitor Sibofimloc (TAK-018/EB8018) in the treatment of Crohn's disease.

In this review we have focused on discussing the advances being made towards the development of mannosebased UPEC targeting UTI treatments. It is worth noting that research has also been done to develop galactose analogs capable of inhibiting the F9 pilus, a homolog to the type one pilus, which has an important role in the maintenance of UPEC-induced UTIs.^[89] While it is not the aim of this review to cover advances in the development of these galactose-based inhibitors, interested readers can find information on this topic published elsewhere ^{[89] [90]}.

3.0 Summary and outlook

We have summarized the main areas of interest regarding research into mannose-based FimH inhibitors. In the current climate, where antibiotic resistance is becoming ever more prevalent, there is a need to develop effective targeted antibacterial treatments. Overall the studies discussed illustrate the great potential of mannose-based inhibitors as targeted treatments against UPEC-induced UTIs. Analogs from each class of mannose-based inhibitor have shown impressive potency against FimH, and warrant further investigation. However, there are disadvantages associated with the use of each inhibitor class. Mannose-capped oligosaccharides most accurately reflect the structure of natural FimH ligands, yet their likely lack of oral bioavailability and complex chemical synthesis widely restricts their use. Simpler α -D-mannopyranoside based inhibitors are currently the most explored class of mannosebased FimH inhibitors and potentially the most promising. The main advantages of *α*-D-mannopyranoside inhibitors are their simpler and smaller structures, making them easier to synthesize and providing a better chance of oral absorption. However, the development of analogs with both high in vivo efficacy and high oral bioavailability is challenging. A further limitation to α-D-mannopyranoside based inhibitors is that their structure prevents polyvalent FimH targeting, potentially limiting their potency. Research into polyvalent mannose scaffolds is still largely in its infancy, and it is not known as yet how effective these inhibitors will be. The potential applications of a highaffinity polyvalent α -D-mannopyranoside based inhibitor are vast, yet their high molecular weight and hydrophilic properties could impact their oral bioavailability, preventing them from functioning as an effective UPEC-induced UTI treatment. These analogs may find alternative applications as a potential treatment for Crohn's disease. It is expected that continued work into both established and novel FimH targeting methods will further advance the field, and take us closer to the development of an effective treatment for UPEC-induced UTIs.

4.0 Acknowledgements

The authors thank The University of York and the Engineering and Physical Sciences Research Council (1944882) for research funding.

14, 749-757.

31. 623-636.

17.

257.

489

528.

15, 247-256.

341-352.

2009, 19, 129-139.

[1] a) K. Mayer, D. Eris, O. Schwardt, C. P. Sager, S. Rabbani, S. Kleeb and B. Ernst, J. Med. Chem. 2017, 60, 5646-5662; b) W. 445 Schonemann, J. Cramer, T. Muhlethaler, B. Fiege, M. Silbermann, S. Rabbani, P. Datwyler, P. Zihlmann, R. P. Jakob, C. P. Sager, M. 58, 1995-1998. Smiesko, O. Schwardt, T. Maier and B. Ernst, ChemMedChem 2019, [2] S. Vanwetswinkel, A. N. Volkov, Y. G. J. Sterckx, A. Garcia-Pino, L. Buts, W. F. Vranken, J. Bouckaert, R. Roy, L. Wyns and N. A. J. van Nuland, J. Med. Chem. 2014, 57, 1416-1427. Soc. 2019, 141, 936-944. [3] D. R. Hickling, T. T. Sun and X. R. Wu, Microbiol. Spectr. 2015, 3, [4] J. Kaufman, M. Temple-Smith and L. Sanci, BMJ Paediatr. open 2019, 3, e000487-e000487. [5] I. Katnik-Prastowska, J. Lis and A. Matejuk, Glycoconj. J. 2014, 13. R611-R613. [6] a) C. C. A. Hu, F. X. Liang, G. Zhou, L. Y. Tu, C. H. A. Tang, J. Zhou, G. Kreibich and T. T. Sun, Mol Biol Cell 2005, 16, 3937-3950; b) B. Kachar, F. X. Liang, U. Lins, M. X. Ding, X. R. Wu, D. Stoffler, U. Aebi and T. T. Sun, J. Mol. Biol. 1999, 285, 595-608. [7] G. Lee, Int Neurourol J 2011, 15, 4-12. [8] M. A. Matuszewski, K. Tupikowski, L. Dolowy, B. Szymanska, J. Dembowski and R. Zdrojowy, Cent European J Urol 2016, 69, 252-[9] a) UniProt in UniProtKB - O00322 (UPK1A_HUMAN), Vol. 2020; 655 b) UniProt in UniProtKB - O75841 (UPK1B_HUMAN), Vol. 2020. [10] M. S. Najar, C. L. Saldanha and K. A. Banday, Ind. J. Nephrol [11] C. W. Tan and M. P. Chlebicki, Singap. Med. J. 2016, 57, 485-[12] L. E. Nicolle, Infect. Dis. Clin. North Am. 1997, 11, 647-&. and J. W. Janetka, J. Med. Chem. 2010, 53, 4779-4792. [13] T. L. Greenhow, Y. Y. Hung, A. M. Herz, E. Losada and R. H. Pantell, Pediatr. Infect. Dis. J. 2014, 33, 595-599. [14] F. M. E. Wagenlehner, A. Pilatz, K. G. Naber and W. Weidner, Eur. J. Clin. Invest. 2008, 38, 45-49. [49] a) D. C. Old, Journal of Gen. Microbiol. 1972, 71, 149-&; b) I. E. [15] B. Foxman, Dis. Mon. 2003, 49, 53-70. [16] J. A. Ryu and B. Kim, Radiographics 2001, 21, 1169-1185. [17] T. Klein, D. Abgottspon, M. Wittwer, S. Rabbani, J. Herold, X. H. Jiang, S. Kleeb, C. Luthi, M. Scharenberg, J. Bezencon, E. Gubler, L. J. Pang, M. Smiesko, B. Cutting, O. Schwardt and B. Ernst, J. Med. Chem. 2010, 53, 8627-8641. [18] T. A. Rowe and M. Juthani-Mehta, Aging Health 2013, 9, 519-4799 [19] B. Foxman, Infect. Dis. Clin. North. Am. 2014, 28, 1-+. [20] A. Ronald, Am. J. Med. 2002, 113, 14S-19S. 1085-1090. [21] M. D. Melekos and K. G. Naber, Int. J. Antimicrob. Agents 2000, 2012, 7, 1404-1422. [22] B. Wiedemann, A. Heisig and P. Heisig, Antibiotics 2014, 3, [23] N. N. T. Goemaere, K. Grijm, P. T. W. van Hal and M. A. den Bakker, J. Med. Case Rep 2008, 2, 169. [24] L. Cegelski, G. R. Marshall, G. R. Eldridge and S. J. Hultgren, 123-168 Mol. Microbiol. 2006, 61, 1556-1568. Lett. 2014, 16, 1116-1119. ChemComm. 2015, 51, 13283-13285. 2003, 68, 8142-8148.

15

Nat. Revi. Microbiol. 2008, 6, 17-27 [25] M. E. Terlizzi, G. Gribaudo and M. E. Maffei, Front. Microbiol.

2017, 8, 23 [26] M. G. Blango, E. M. Ott, A. Erman, P. Veranic and M. A. Mulvey, Plos One 2014, 9, 9.

[27] a) M. A. Mulvey, Y. S. Lopez-Boado, C. L. Wilson, R. Roth, W.

C. Parks, J. Heuser and S. J. Hultgren, Science 1998, 282, 1494-

1497; b) J. J. Martinez, M. A. Mulvey, J. D. Schilling, J. S. Pinkner

and S. J. Hultgren, EMBO J. 2000, 19, 2803-2812.

[28] D. S. Eto, T. A. Jones, J. L. Sundsbak and M. A. Mulvey, PLoS Pathog. 2007, 3, 949-961.

[29] J. J. Martinez and S. J. Hultgren, Cell. Microbiol. 2002, 4, 19-28.

[30] M. J. Duncan, G. J. Li, J. S. Shin, J. L. Carson and S. N.

Abraham, J. Biol. Chem. 2004, 279, 18944-18951.

[31] B. K. Dhakal and M. A. Mulvey, J. Biol. Chem. 2009, 284, 446-454.

[32] J. D. Schilling, M. A. Mulvey and S. J. Hultgren, J. Infect. Dis. 2001, 183, S36-S40.

[33] M. S. Hanson and C. C. Brinton, Nature 1988, 332, 265-268.

[34] C. Spaulding, H. L. Schreiber, W. L. Zheng, K. W. Dodson, J. E.

Hazen, M. Conover, F. B. Wang, P. Svenmarker, A. Luna-Rico, O Francetic, M. Andersson, S. Hultgren and E. H. Egelman, Elife 2018,

7.25. [35] M. S. Hanson, J. Hempel and C. C. Brinton, J. Bacteriol. 1988,

170, 3350-3358.

[36] K. A. Krogfelt and P. Klemm, Microb. Pathog. 1988, 4, 231-238.

[37] P. Klemm and G. Christiansen, Mol. Gen. Gen. 1987, 208, 439-

[38] K. A. Krogfelt, H. Bergmans and P. Klemm, Infect. Immun. 1990,

[39] L. K. Mydock-McGrane, T. J. Hannan and J. W. Janetka, Expert Opin. Drug Discov. 2017, 12, 711-731.

[40] M. M. Sauer, R. P. Jakob, T. Luber, F. Canonica, G. Navarra, B. Ernst, C. Unverzagt, T. Maier and R. Glockshuber, J. Am. Chem.

[41] I. Le Trong, P. Aprikian, B. A. Kidd, W. E. Thomas, E. V. Sokurenko and R. E. Stenkamp, J. Struct. Biol. 2010, 172, 380-388.

[42] M. Forero, W. E. Thomas, C. Bland, L. M. Nilsson, E. V.

Sokurenko and V. Vogel, Nano Lett. 2004, 4, 1593-1597.

[43] K. Konstantopoulos, W. D. Hanley and D. Wirtz, Curr. Biol. 2003,

- [44] a) R. R. Isberg and P. Barnes, Cell 2002, 110, 1-4; b) K. C.
- Chang and D. A. Hammer, Biophys. J. 1999, 76, 1280-1292

[45] a) P. Aprikian, V. Tchesnokova, B. Kidd, O. Yakovenko, V.

Yarov-Yarovoy, E. Trinchina, V. Vogel, W. Thomas and E.

Sokurenko, J. Biol. Chem. 2007, 282, 23437-23446; b) I. Le Trong,

P. Aprikian, B. A. Kidd, M. Forero-Shelton, V. Tchesnokova, P.

Rajagopal, V. Rodriguez, G. Interlandi, R. Klevit, V. Vogel, R. E. Stenkamp, E. V. Sokurenko and W. E. Thomas, Cell 2010, 141, 645-

[46] A. Wellens, C. Garofalo, H. Nguyen, N. Van Gerven, R. Slattegard, J. P. Hernalsteens, L. Wyns, S. Oscarson, H. De Greve,

S. Hultgren and J. Bouckaert, PLoS One 2008, 3, 13.

[47] Z. F. Han, J. S. Pinkner, B. Ford, R. Obermann, W. Nolan, S. A. Wildman, D. Hobbs, T. Ellenberger, C. K. Cusumano, S. J. Hultgren

[48] C. Jarvis, Z. F. Han, V. Kalas, R. Klein, J. S. Pinkner, B. Ford, J. Binkley, C. K. Cusumano, Z. Cusumano, L. Mydock-McGrane, S. J.

Hultgren and J. W. Janetka, ChemMedChem 2016, 11, 367-373.

Salit and E. C. Gotschlich, J. Exp. Med. 1977, 146, 1169-1181.

[50] C. S. Hung, J. Bouckaert, D. Hung, J. Pinkner, C. Widberg, A.

DeFusco, C. G. Auguste, R. Strouse, S. Langermann, G. Waksman

and S. J. Hultgren, Mol. Microbiol. 2002, 44, 903-915.

[51] A. Wellens, M. Lahmann, M. Touaibia, J. Vaucher, S. Oscarson, R. Roy, H. Remaut and J. Bouckaert, Biochemistry 2012, 51, 4790-

[52] G. Roos, A. Wellens, M. Touaibia, N. Yamakawa, P. Geerlings, R. Roy, L. Wyns and J. Bouckaert, ACS Med. Chem. Lett. 2013, 4,

[53] L. J. Pang, S. Kleeb, K. Lemme, S. Rabbani, M. Scharenberg, A. Zalewski, F. Schadler, O. Schwardt and B. Ernst, ChemMedChem

[54] a) M. Hartmann and T. K. Lindhorst, Chem. Eur. J. 2011, 3583-3609; b) L. Mousavifar, M. Touaibia and R. Roy, Acc. Chem. Res. 2018, 51, 2937-2948; c) S. G. Gouin, Future Med. Chem. 2016, 8, 1903-1906; d) S. G. Gouin, G. Roos and J. Bouckaert in Discovery and Application of FimH Antagonists, Eds.: P. H. Seeberger and C. Rademacher), Springer International Publishing, Cham, 2014, pp.

[55] B. Xie, G. Zhou, S. Y. Chan, E. Shapiro, X. P. Kong, X. R. Wu, T. T. Sun and C. E. Costello, J. Biol. Chem. 2006, 281, 14644-14653.

[56] a) N. Sharon, FEBS Lett. 1987, 217, 145-157; b) N. Firon, I.

Ofek and N. Sharon, Carbohydr. Res. 1983, 120, 235-249.

[57] J. Bouckaert, J. Mackenzie, J. L. de Paz, B. Chipwaza, D.

Choudhury, A. Zavialov, K. Mannerstedt, J. Anderson, D. Pierard, L. Wyns, P. H. Seeberger, S. Oscarson, H. De Greve and S. D. Knight,

[58] M. Heuckendorff, J. Bendix, C. M. Pedersen and M. Bols, Org.

[59] a) D. Crich and S. X. Sun, J. Org. Chem. 1996, 61, 4506-4507; b) D. Crich and S. X. Sun, Tetrahedron 1998, 54, 8321-8348; c) D. Crich and H. M. Li, J. Org. Chem. 2000, 65, 801-805; d) D. Crich and S. X. Sun, J. Org. Chem. 1997, 62, 1198-1199; e) M. Heuckendorff, P. S. Bols, C. B. Barry, T. G. Frihed, C. M. Pedersen and M. Bols,

[60] a) D. Crich and O. Vinogradova, J. Org. Chem. 2006, 71, 8473-8480; b) D. Crich, M. de la Mora and A. U. Vinod, J. Org. Chem.

[61] J. Bouckaert, J. Berglund, M. Schembri, E. De Genst, L. Cools, M. Wuhrer, C. S. Hung, J. Pinkner, R. Slattegard, A. Zavialov, D. Choudhury, S. Langermann, S. J. Hultgren, L. Wyns, P. Klemm, S.

Oscarson, S. D. Knight and H. De Greve, Mol. Microbiol. 2005, 55, 441-455 [62] S. Oscarson and A. K. Tiden. Carbohvdr. Res. 1993, 247, 323-328. [63] N. Firon, S. Ashkenazi, D. Mirelman, I. Ofek and N. Sharon, Infect. Immun. 1987, 55, 472-476. [64] O. Sperling, A. Fuchs and T. K. Lindhorst, Org. Biomol. Chem. 2006, 4, 3913-3922. [65] T. K. Lindhorst, S. Kotter, J. Kubisch, U. Krallmann-Wenzel, S. Ehlers and V. Kren, Eur. J. Org. Chem. 1998, 1998, 1669-1674. [66] a) Z. F. Han, J. S. Pinkner, B. Ford, E. Chorell, J. M. Crowley, C. K. Cusumano, S. Campbell, J. P. Henderson, S. J. Hultgren and J. W. Janetka, J. Med. Chem. 2012, 55, 3945-3959; b) C. K. Cusumano, J. S. Pinkner, Z. Han, S. E. Greene, B. A. Ford, J. R. Crowley, J. P. Henderson, J. W. Janetka and S. J. Hultgren, Sci. Trans. Med. 2011, 3, 109ra115-109ra115. [67] M. Vetsch, P. Sebbel and R. Glockshuber, J. Mol. Biol. 2002, 322, 827-840. [68] C. Grabosch, M. Hartmann, J. Schmidt-Lassen and T. K. Lindhorst, ChemBioChem 2011, 12, 1066-1074. [69] C. P. Sager, B. Fiege, P. Zihlmann, R. Vannam, S. Rabbani, R. P. Jakob, R. C. Preston, A. Zalewski, T. Maier, M. W. Peczuh and B. Ernst, Chem. Sci. 2018, 9, 646-654. [70] a) D. A. Dorta, T. Chalopin, A. Sivignon, J. de Ruyck, T. I. Dumych, R. O. Bilyy, D. Deniaud, N. Barnich, J. Bouckaert and S. G. Gouin, ChemMedChem 2017, 12, 986-998; b) S. Brument, A. Sivignon, T. I. Dumych, N. Moreau, G. Roos, Y. Guerardel, T. Chalopin, D. Deniaud, R. O. Bilyy, A. Darfeuille-Michaud, J. Bouckaert and S. G. Gouin, J. Med. Chem. 2013, 56, 5395-5406. [71] X. H. Jiang, D. Abgottspon, S. Kleeb, S. Rabbani, M. Scharenberg, M. Wittwer, M. Haug, O. Schwardt and B. Ernst, J. Med. Chem. 2012, 55, 4700-4713. [72] M. Scharenberg, O. Schwardt, S. Rabbani and B. Ernst, J. Med. Chem. 2012, 55, 9810-9816. [73] a) J. J. Lundguist and E. J. Toone, Chem. Rev. 2002, 102, 555-578; b) Y. C. Lee and R. T. Lee, Acc. Chem. Res. 1995, 28, 321-327. [74] S. Cecioni, A. Imberty and S. Vidal, Chem. Rev. 2015, 115, 525-561. [75] P. Rassam, N. A. Copeland, O. Birkholz, C. Tóth, M. Chavent, A. L. Duncan, S. J. Cross, N. G. Housden, R. Kaminska, U. Seger, D. M. Quinn, T. J. Garrod, M. S. P. Sansom, J. Piehler, C. G. Baumann and C. Kleanthous, Nature 2015, 523, 333-336. [76] a) A. Bernardi, J. Jimenez-Barbero, A. Casnati, C. De Castro, T. Darbre, F. Fieschi, J. Finne, H. Funken, K. E. Jaeger, M. Lahmann, T. K. Lindhorst, M. Marradi, P. Messner, A. Molinaro, P. V. Murphy, C. Nativi, S. Oscarson, S. Penades, F. Peri, R. J. Pieters, O. Renaudet, J. L. Reymond, B. Richichi, J. Rojo, F. Sansone, C. Schaffer, W. B. Turnbull, T. Velasco-Torrijos, S. Vidal, S. Vincent, T. Wennekes, H. Zuilhof and A. Imberty, Chem. Soc. Rev. 2013, 42, 4709-4727; b) M. Hartmann, P. Betz, Y. C. Sun, S. N. Gorb, T. K. Lindhorst and A. Krueger, Chem. Eur. J. 2012, 18, 6485-6492. [77] a) S. G. Gouin, A. Wellens, J. Bouckaert and J. Kovensky, ChemMedChem 2009, 4, 749-755; b) M. Almant, V. Moreau, J. Kovensky, J. Bouckaert and S. G. Gouin, Chem. Eur. J. 2011, 17, 10029-10038; c) J. Bouckaert, Z. L. Li, C. Xavier, M. Almant, V. Caveliers, T. Lahoutte, S. D. Weeks, J. Kovensky and S. G. Gouin, Chem. Eur. J. 2013, 19, 7847-7855; d) A. Sivignon, X. B. Yan, D. A. Dorta, R. Bonnet, J. Bouckaert, E. Fleury, J. Bernard, S. G. Gouin, A. Darfeuille-Michaud and N. Barnich, Mbio 2015, 6, 9; e) X. B. Yan, V. La Padula, S. Favre-Bonte and J. Bernard, Eur. Polym. J. 2019, 112, 170-175 [78] a) A. Schierholt, M. Hartmann, K. Schwekendiek and T. K. Lindhorst, Eur. J. Org. Chem. 2010, 2010, 3120-3128; b) M. Touaibia, T. C. Shiao, A. Papadopoulos, J. Vaucher, Q. G. Wang, K. Benhamioud and R. Roy, Chem. Commun. 2007, 380-382; c) T. K. Lindhorst, C. Kieburg and U. Krallmann-Wenzel, Glycoconjug. J. 1998, 15, 605-613; d) M. Dubber, O. Sperling and T. K. Lindhorst, Org. Biomol. Chem. 2006, 4, 3901-3912. [79] a) M. Touaibia, A. Wellens, T. C. Shiao, Q. Wang, S. Sirois, J. Bouckaert and R. Roy, ChemMedChem 2007, 2, 1190-1201; b) A. Papadopoulos, T. C. Shiao and R. Roy, Mol. Pharmaceut. 2012, 9, 394-403; c) J. D. K. Twibanire, N. K. Paul and T. B. Grindley, New J. Chem. 2015, 39, 4115-4127; d) N. Nagahori, R. T. Lee, S. Nishimura, D. Page, R. Roy and Y. C. Lee, ChemBioChem 2002, 3, 836-844. [80] S. Kötter, U. Krallmann-Wenzel, S. Ehlers and T. K. Lindhorst, J. Chem. Soc. Perkin Trans. 1 1998, 2193-2200.

[81] A. Barras, F. A. Martin, O. Bande, J. S. Baumann, J. M. Ghigo, R. Boukherroub, C. Beloin, A. Siriwardena and S. Szunerits, *Nanoscale* **2013**, *5*, 2307-2316.

[82] J. F. Nierengarten, J. Iehl, V. Oerthel, M. Holler, B. M. Illescas, A. Munoz, N. Martin, J. Rojo, M. Sanchez-Navarro, S. Cecioni, S. Vidal, K. Buffet, M. Durka and S. P. Vincent, *Chem. Commun.* **2010**, *46*, 3860-3862.

[83] M. Durka, K. Buffet, J. lehl, M. Holler, J. F. Nierengarten, J. Taganna, J. Bouckaert and S. P. Vincent, *Chem. Commun.* **2011**, 47, 1321-1323.

[84] A. Schierholt, M. Hartmann and T. K. Lindhorst, *Carbohydr. Res* **2011**, *346*, 1519-1526.

[85] B. Schmidt, S. Sankaran, L. Stegemann, C. A. Strassert, P. Jonkheijm and J. Voskuhl, *J. Mat. Chem. B.* **2016**, *4*, 4732-4738.

[86] I. Nierengarten, K. Buffet, M. Holler, S. P. Vincent and J. F.

Nierengarten, Tetrahedron Lett. 2013, 54, 2398-2402.

[87] W. Reinisch, X. Hebuterne, A. Buisson, S. Schreiber, P.

Desreumaux, J. M. Paillarse and C. Bonny, *J Crohns Colitis* **2020**, *14*, S479-S480.

[88] W. Reinisch, X. Hebuterne, A. Buisson, S. Schreiber, P.

Desreumaux, C. Primas, J. M. Paillarse and C. Bonny,

Gastroenterology 2020, 158, S1211-S1212.

[89] V. Kalas, M. E. Hibbing, A. R. Maddirala, R. Chugani, J. S. Pinkner, L. K. Mydock-McGrane, M. S. Conover, J. W. Janetka and

S. J. Hultgren, PNAS 2018, 115, E2819-E2828.

[90] A. Salminen, V. Loimaranta, J. A. F. Joosten, A. S. Khan, J. Hacker, R. J. Pieters and J. Finne, *J. Antimicrob. Chemother.* **2007**, *60*, 495-501.