



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/165368/>

Version: Published Version

Article:

de Oliveira, Claire, Iwajomo, Tomisin, Gomes, Tara et al. (2020) Le rôle de la formation à la faculté de médecine à l'égard des pratiques de prescription d'opioïdes des médecins: données probantes d'Ontario, Canada. Canadian Journal of Psychiatry. ISSN: 0706-7437

<https://doi.org/10.1177/0706743720931240>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC) licence. This licence allows you to remix, tweak, and build upon this work non-commercially, and any new works must also acknowledge the authors and be non-commercial. You don't have to license any derivative works on the same terms. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

The Role of the Medical School Training on Physician Opioid Prescribing Practices: Evidence from Ontario, Canada

The Canadian Journal of Psychiatry /
La Revue Canadienne de Psychiatrie

1-11

© The Author(s) 2020



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/0706743720931240

TheCJP.ca | LaRCP.ca



Le rôle de la formation à la faculté de médecine à l'égard des pratiques de prescription d'opioïdes des médecins: données probantes d'Ontario, Canada

**Claire de Oliveira, MA, PhD^{1,2,3,4} , Tomisin Iwajomo, MPH²,
Tara Gomes, MHSc, PhD^{3,4,5,6}, and Paul Kurdyak, MD, MSc, PhD, FRCP(C)^{2,3,4,7} **

Abstract

Background: Recent research found that physicians who completed medical school training at top-ranked U.S. medical schools prescribed fewer opioids than those trained at lower ranked schools, suggesting that physician training may play a role in the opioid epidemic. We replicated this analysis to understand whether this finding holds for Ontario, Canada.

Methods: We used data on all opioid prescriptions written by Ontario physicians between 2013 and 2017 from the Narcotics Monitoring System. Using the Corporate Provider Database and ICES Physician Database, which contain medical school of training, we linked patients who filled opioid prescriptions with their respective prescribing physician. Available data on Canadian medical school rankings were obtained from *Maclean's* news magazine. We used regression analysis to assess the relationship between number of opioid prescriptions and medical school ranking.

Results: Compared to the United States, average annual number of opioid prescriptions per physician was lower in Ontario (236 vs. 78). Unlike the United States, we found little evidence that physicians trained at lower ranked medical schools prescribed more than their top-ranked school counterparts after controlling for specialty and location of practice. However, primary care physicians trained at non-English-speaking foreign schools prescribed the most opioids even after excluding opioid maintenance therapy-related prescriptions.

Conclusion: The role of medical school training on opioid prescribing patterns among Ontario physicians differs from that in the United States likely due to greater homogeneity of curricula among Canadian schools. Ensuring physicians trained abroad receive additional pain management/addiction training may help address part of the opioid epidemic in Ontario.

Abrégé

Contexte : Une recherche récente a observé que les médecins qui suivaient leur formation médicale dans les écoles de médecine américaines les mieux cotées prescrivaient moins d'opioïdes que ceux formés dans des écoles moins bien cotées, ce

¹ Centre for Health Economics/Hull York Medical School, University of York, Heslington, United Kingdom

² Institute for Mental Health Policy Research (IMHPR), Centre for Addiction and Mental Health (CAMH), Toronto, Ontario, Canada

³ ICES, Toronto, Ontario, Canada

⁴ Institute of Health Policy, Management and Evaluation (IHPE), University of Toronto, Ontario, Canada

⁵ Li Ka Shing Knowledge Institute, St. Michael's Hospital, Toronto, Ontario, Canada

⁶ Leslie Dan Faculty of Pharmacy, University of Toronto, Ontario, Canada

⁷ Department of Psychiatry, University of Toronto, Ontario, Canada

Corresponding Author:

Claire de Oliveira, MA, PhD, Hull York Medical School/Centre for Health Economics, Alcuin A Block, University of York, Heslington, York YO10 5DD, United Kingdom.

Email: claire.deoliveira@york.ac.uk

qui suggère que la formation des médecins peut jouer un rôle dans l'épidémie d'opioïdes. Nous avons reproduit cette analyse pour comprendre si ce résultat tient pour l'Ontario, Canada.

Méthodes : Nous avons utilisé les données de toutes les prescriptions d'opioïdes écrites par des médecins ontariens entre 2013 et 2017, tirées du Système de surveillance des stupéfiants et des substances contrôlées. À l'aide de la Base de données centrale sur les fournisseurs de services de santé et de la Base de données des médecins de l'IRSS, qui contient la formation à l'école de médecine, nous avons lié les patients qui faisaient remplir des prescriptions d'opioïdes à leur médecin prescripteur respectif. Les données disponibles sur les classements des facultés de médecine canadiennes ont été obtenues du magazine d'actualité *Maclean's*. Nous avons utilisé l'analyse de régression pour évaluer la relation entre le nombre de prescriptions d'opioïdes et le classement des facultés de médecine.

Résultats : Comparé aux États-Unis, le nombre annuel moyen de prescriptions d'opioïdes par médecin était plus faible en Ontario (236 contre 78). Contrairement aux États-Unis, nous avons trouvé peu de données probantes indiquant que les médecins formés à des facultés de médecine moins bien cotées prescrivaient davantage que leurs homologues des écoles au sommet du classement, après contrôle de la spécialité et de l'emplacement de la pratique. Toutefois, les médecins des soins de première ligne formés dans des écoles étrangères non anglophones prescrivaient le plus d'opioïdes, même après l'exclusion des prescriptions liées au traitement d'entretien aux opioïdes.

Conclusion : Le rôle de la formation à la faculté de médecine dans les modèles de prescription d'opioïdes chez les médecins de l'Ontario diffère de celui des États-Unis, probablement en raison de la plus grande homogénéité des programmes d'études dans les écoles canadiennes. Assigner aux médecins formés à l'étranger une formation additionnelle sur la gestion de la douleur/dépendance pourrait contribuer à résoudre en partie l'épidémie d'opioïdes en Ontario.

Keywords

opioids, physician prescribing, administrative data

Introduction

Opioid overdose is a leading public health issue in North America. Over the last two decades, there has been an increase in opioid use in Canada and in the United States; however, this pattern has changed in recent years.^{1,2} In 2015, both Canada and the United States had the highest opioid consumption per capita worldwide.³ Furthermore, opioid-related deaths have increased dramatically in both countries.⁴⁻⁶ In Ontario, more than half of all opioid-related deaths in 2016 involved prescription drugs (dispensed or diverted), with fentanyl being the most commonly nonprescribed opioid.⁷ Given this, policy makers have sought ways to address this epidemic.⁸

Evidence suggests that increased use of opioids may be driven partly by physician prescribing practices. Among the suggested key drivers behind the increased opioid use in the last 20 years is the aggressive lobbying of physicians to prescribe opioids liberally to manage chronic pain.⁹ Using U.S. Medicare emergency department data, Barnett et al. (2017) found that long-term opioid use was significantly higher among patients treated by physicians with a higher propensity to prescribe opioids.¹⁰ Other U.S. research found that physicians who completed medical school training at top medical schools wrote significantly fewer opioid prescriptions than those trained in lower ranked schools even after controlling for specialty and location of practice.² The authors concluded this was due to physician education rather than patient selection across physicians or physician selection across medical schools. These findings suggest there may be a role for physician education in addressing the opioid epidemic. However, there is no evidence as to

whether this relationship holds for other jurisdictions such as Canada. We hypothesize that, given the smaller number of Canadian medical schools, and potentially greater homogeneity across curricula, the U.S. findings do not apply to Canada. Our aim was to replicate the U.S. study using data from Ontario, Canada's most populous province, by examining the relationship between physician opioid prescribing patterns and medical school training and compare our results to those reported in the United States.

Methods

Data Sources

We accessed administrative health care data through ICES, an independent, nonprofit research institute, which maintains provincial health care records (www.ices.on.ca). We employed data on all opioid prescriptions written by physicians practicing in Ontario. These data were obtained from the Narcotics Monitoring System database that includes information on all opioid analgesic prescriptions dispensed from community retail pharmacies across the province and prescribers. Patients in the Narcotics Monitoring System who received and filled an opioid prescription were linked to their respective prescribing physician using unique physician identifiers. We used the Corporate Provider Database to identify all physicians actively practicing in Ontario from 2013 to 2017. These physicians were linked with the ICES Physician Database that contains information on physician demographics, reported specialty, postal code of practice, and year and school of graduation. Data on annual medical school rankings from 2013 to 2017 were obtained from

Maclean's,¹¹ a Canadian news magazine, which publishes university rankings. Rankings incorporate information on students (e.g., success of students at winning academic awards), faculty (e.g., number who have won major awards), resources (e.g., amount of money available for expenses), student support (e.g., assistance available to students), and reputation (e.g., how ready graduates are to embark on successful careers); information on *Maclean's* ranking methodology can be found elsewhere.¹² In Canada, there are 17 medical schools in which 6 are located in Ontario.

Physician Cohort. We examined opioid prescribing practices among all active physicians practicing in Ontario who graduated up until 2011 (school of graduation data were not available thereafter), with a specialty that required the degree of MD, and who were active during the analysis period (i.e., physicians who had a status of either “unrestricted,” “special case license,” or “group practice only” in the Corporate Provider Database). We excluded all physicians with missing data on graduation year, medical school, and specialty and location of practice and those who retired or died during the analysis period. Our final sample included 30,844 physicians (of 39,919 active physicians) with an observation for each year of the analysis period for a total of 154,220 physician-years.

We used the Statistics Canada's Postal Code Conversion File¹³ and Canada Census data to link physician postal code of practice to the respective census division. To characterize physician census division of practice, we obtained information on population density (number of people per square kilometers), percentage of population with a high school diploma or less, percentage of population unemployed, percentage of individuals in the lowest neighbourhood income quintile, percentage of individuals living in a rural setting, and percentage of individuals living in Northern Ontario (i.e., North East and North West Local Health Integration Networks).

Medical School Rankings. Based on *Maclean's* ranking data, and in line with previous research,² we created a composite medical school rank variable based on the average of the medical school's rankings across all years of available data to deal with changes in rankings over time (see Table A1 in the Online Appendix). These data include rankings for 15 of the 17 medical schools in Canada (from highest to lowest ranked, where the highest was assigned 1 and the lowest 15): McGill University, University of Toronto, University of British Columbia, Queen's University, University of Alberta, McMaster University, Dalhousie University, University of Ottawa, University of Western Ontario, University of Calgary, Université de Montréal, Université Laval, University of Saskatchewan, University of Manitoba, and Université de Sherbrooke (Figure A1 in the Online Appendix depicts medical school rankings; rankings among top schools have been quite consistent over time). There was no ranking for Memorial University of Newfoundland or the Northern Ontario School of Medicine; therefore, these schools were

considered separately. We grouped foreign medical schools into three categories: United States; United Kingdom, Ireland, Australia, and New Zealand (UKIANZ, i.e., other English-speaking countries); and Other. (Figure A2 in the Online Appendix provides number of physicians by medical school of training.)

Analysis

Descriptive analysis. We produced summary statistics of annual prescription-related outcomes such as total number of opioid prescriptions written in Ontario, average number of opioid prescriptions written per physician (with and without physician-years with no opioid prescriptions, i.e., zeroes), and the percentage of physician-years with zero prescriptions. We examined the relationship between number of opioid prescriptions and medical school ranking. Primary care physicians (PCPs) accounted for most opioid prescriptions (80%); thus, we examined all physicians and PCPs separately (where PCPs included three subspecialty categories—general practice, family practice, and internal medicine; Currie and Schnell use the term GP instead of PCP; however, the definition is the same).

In line with previous work, we tried to understand whether patient sorting across physicians (i.e., whether patients were more likely to choose physicians based on their prescribing patterns) or physician sorting across medical schools (i.e., whether physicians chose medical schools whose training, e.g., aligned with their approach toward opioids) played a role in our findings.² Given our data, we could not ascertain whether physicians self-selected into certain medical schools. Nonetheless, we could investigate whether physicians who trained at lower ranked medical schools were more likely to practice in specialties and/or locations where patient use of opioids might be greater. Therefore, we examined the average number of opioid prescriptions by medical school ranking for the top 8 opioid prescribing medical specialties (general practice, orthopedic surgery, general surgery, obstetrics and gynecology, plastic surgery, emergency medicine, urology, and otolaryngology) and hematology/oncology (a composite medical specialty that included hematology, medical oncology, and radiation oncology). We also produced these numbers by characteristics of census division of practice.

Regression analysis. Next, we undertook regression analyses to assess the validity of our descriptive findings, while accounting for relevant variables, such as physician specialty and census division of practice, in line with Schnell and Currie (2017). Pooling data from all years, we estimated a linear regression model through ordinary least squares, where the dependent variable was the number of opioid prescriptions written by each physician by year and census division and the independent variables included indicator variables for each ranked school, where McGill was the reference case; indicator variables for each unranked school (the Northern

Ontario School of Medicine and Memorial University); indicator variables for U.S. schools, United Kingdom, Ireland, Australia, and New Zealand schools, and other foreign schools, respectively; and indicator variables for physician specialty, census division, and year. We estimated this equation including physician-years with and without zero opioid prescriptions for all physicians and PCPs separately. Robust standard errors were estimated and clustered by physician.

We also estimated the equation for each of the top 8 opioid prescribing specialties and hematology/oncology to understand whether the relationship between prescribing and medical school differed by specialty (thus, we dropped the indicator variables for specialty but included indicator variables for subspecialty for PCPs to account for differences across the three subspecialties). If medical training plays a role, the prescribing relationship will likely be weaker among specialties that receive subsequent training in pain management and addiction compared to those that do not.

Finally, we examined whether opioid prescribing differed across graduation cohorts, as more recent graduates may have received different training on pain management and addiction than older cohorts, by estimating the equation for each graduation cohort: before 1982, 1982 to 1991, 1992 to 2001, and 2002 to 2011.

Sensitivity analysis. Prior research was not able to examine the number of pills included in each prescription; this may impact our findings as the quantity of medication dispensed can vary considerably between prescriptions. Therefore, we also estimated regression models using the number of opioids dispensed as the dependent variable.

Previous work included *all* opioids prescribed regardless of purpose.² However, some opioids (methadone and buprenorphine/naloxone) are typically prescribed to treat opioid dependence and thus have a different purpose than those prescribed to treat pain. Furthermore, given the nature of their dispensing (typically daily dispensing), their inclusion could affect our primary outcome.¹⁴ Thus, we conducted a sensitivity analysis excluding all opioid maintenance therapy (OMT)-related prescriptions.

Results

Descriptive Analysis

From 2013 to 2017, there was an overall decrease in the number of opioid prescriptions in Ontario (Table 1). Physicians accounted for most opioid prescriptions (87%). The average annual number of opioid prescriptions per physician across all years was 77.79 (standard deviation [*SD*] = 138.31) including physician-years with no opioid prescriptions (i.e., zeroes) and 103.58 (*SD* = 151.01) excluding physician-years with no opioid prescriptions. These numbers were higher for PCPs (124.67, *SD* = 164.85 and 140.06, *SD* = 168.45, respectively). Among physicians excluded from the analysis ($n = 9,075$, 22.7% of all active physicians in the

Corporate Provider Database between 2012 and 2017 inclusive), most were missing data on medical school of training (99.6%); many were also missing data on specialty (63%). Just under half (47%) did not prescribe at any point during the analysis period. The average annual number of prescriptions per physician among excluded physicians was 20.45 and 52.50 with and without physician-years with no opioid prescriptions, respectively (results not shown).

We found a negative relationship between medical school ranking and average number of opioid prescriptions; in other words, physicians from higher ranked schools (where higher ranked schools were assigned a lower number) were more likely to write opioid prescriptions (Figure 1A; Table A2 in the Online Appendix). For example, physicians trained at Université de Sherbrooke prescribed less than those trained at the University of Toronto (39.08, *SD* = 68.81 vs. 82.96, *SD* = 142.35, respectively). Furthermore, physicians trained at the Northern Ontario School of Medicine prescribed more (121.74, *SD* = 85.91) than those trained at both ranked and unranked schools. Findings were similar for PCPs only, except PCPs trained at the University of Western Ontario (137.05, *SD* = 171.28), the University of Manitoba (141.31, *SD* = 230.41), and other foreign schools (153.29, *SD* = 192.53), who prescribed more opioids than other PCPs (Figure 1B; Table A3 in the Online Appendix).

Results were generally similar when we excluded physician-years with no opioid prescriptions. Again, on average, physicians from higher ranked schools wrote more opioid prescriptions (Online Appendix Table A2). In particular, physicians trained at the University of Western Ontario had an average number of prescriptions (122.37, *SD* = 162.37) close to the Northern Ontario School of Medicine value (125.83, *SD* = 83.34). Findings differed slightly when we examined PCPs only (Online Appendix Table A3). While the negative relationship between medical school rank and opioid prescriptions held, PCPs trained at the University of Manitoba (160.12, *SD* = 239.05) and the University of Western Ontario (153.17, *SD* = 174.14) were identified as the top prescribers among ranked schools (Online Appendix Table A3). However, PCPs trained at other foreign schools (166.43) were the top prescribers among all physicians.

We also examined the number of opioid prescriptions by specialty and location of practice (Table 2). General practice was by far the top opioid prescribing specialty followed by orthopedic surgery, general surgery, obstetrics and gynecology, plastic surgery, emergency medicine, urology, and otolaryngology. Combined, these specialties accounted for 82% of all opioid prescriptions (the top 8 specialties in the United States, which accounted for 84% of all opioid prescriptions, were general practice, orthopedic surgery, emergency medicine, pain medicine, physical medicine and rehabilitation, obstetrics and gynecology, anesthesiology, and general surgery; see Online Appendix Table A4 for average number of opioid prescriptions for the top 20 specialties). Typically, physicians trained at top ranked Canadian schools tended

Table 1. Summary Statistics for Annual Opioid Prescription Measures.

	2013	2014	2015	2016	2017	All
Total number of all opioid prescriptions ^a	2,743,809	2,862,487	2,799,863	2,754,514	2,591,261	13,751,934
All physicians						
Number of physicians	30,844	30,844	30,844	30,844	30,844	30,844
Number of prescribing physicians	24,440	24,085	23,294	22,512	21,481	—
Number of physician-years	—	—	—	—	—	154,220
Number of opioid prescriptions	2,543,112	2,611,391	2,449,569	2,309,765	2,082,376	11,996,213
Percentage of total prescribed	92.69	91.23	87.49	83.85	80.36	87.23
Average opioid prescriptions per year including zeroes (std. deviation)	82.45 (139.55)	84.66 (150.57)	79.42 (137.84)	74.89 (134.87)	67.51 (127.02)	77.79 (138.31)
Average opioid prescriptions per year excluding zeroes (std. deviation)	104.05 (149.43)	108.42 (162.66)	105.16 (149.83)	102.6 (148.59)	96.94 (142.53)	103.58 (151.01)
Zeroes (%)	20.76	21.91	24.48	27.01	30.36	—
Primary care physicians						
Number of physicians	15,166	15,166	15,166	15,166	15,166	15,166
Number of prescribing physicians	14,017	13,882	13,574	13,234	12,794	—
Number of physician-years	—	—	—	—	—	75,830
Number of opioid prescriptions	2,005,360	2,070,284	1,933,355	1,813,976	1,631,074	9,454,049
Percentage of total prescribed	73.09	72.32	69.05	65.85	62.95	68.75
Average opioid prescriptions per year including zeroes (std. deviation)	132.22 (164.98)	136.51 (180.82)	127.48 (163.16)	119.61 (160.32)	107.55 (152.07)	124.67 (164.85)
Average opioid prescriptions per year excluding zeroes (std. deviation)	143.06 (167.03)	149.13 (183.95)	142.43 (166.18)	137.06 (164.50)	127.49 (157.70)	140.06 (168.45)
Zeroes (%)	7.58	8.47	10.50	12.74	15.64	—

Note. Standard deviations are displayed in parentheses and are clustered by physician.

^aObtained from the Narcotics Monitoring System and includes all opioid prescriptions regardless of prescriber.




Figure 1. Average number of opioid prescriptions by medical school rank. (A) All physicians. (B) Primary care physicians. See Tables A2 and A3 in the Online Appendix for values for all physicians and primary care physicians, respectively.

to practice in more densely populated, urban areas located in southern Ontario.

Regression Analysis

After controlling for physician specialty and location of practice, the negative relationship between opioid prescriptions and medical school ranking remained (Figure 2A); however, most school rank coefficients were not statistically significant (see Table A5 in the Online Appendix for coefficients). Compared to McGill University, physicians trained at the University of Toronto (11.81, P value < 0.001), the University of Western Ontario (14.48, P value < 0.001), and other foreign schools (17.16, P value < 0.001) prescribed slightly more opioids, while those trained at the Université de Montréal (−13.61, P value = 0.04) and the University of Saskatchewan (−12.49, P value = 0.03) prescribed slightly less.

We found similar results for PCPs only, albeit a less steep gradient. Few coefficients were statistically significant; however, those for the University of Manitoba (31.04, P value = 0.02) and other foreign schools (40.73, P value < 0.001) were much higher compared to McGill University (although smaller in value—16.97, P value < 0.001—the coefficient for the University of Toronto was also statistically significant; Figure 2B; see Table A6 in the Online Appendix for full list of coefficients). When we excluded physician-years with no opioid prescriptions, few coefficients were statistically significant, the largest of which were for other foreign schools (17.59, P value < 0.001), the University of Western Ontario (14.31, P value < 0.001), and the University of Toronto (12.24, P value < 0.001). We found similar results when we restricted our analysis to PCPs only; the largest coefficients were for other foreign schools (39.16, P value < 0.001), the University of Manitoba (33.07, P value = 0.02), and the University of Toronto (16.85, P value = 0.02; see Tables A7 and A8 in the Online Appendix for full

Table 2. Opioid Prescriptions among Physicians in Ontario by Specialty and Location of Practice Based On Medical School of Graduation.

	Full Sample	Ranked									
		McGill	UofT	UBC	Queen's	Alberta	MAC	DAL	Ottawa	Western	Calgary
Number of Physicians	30,844	1,380	6,976	388	1,987	453	2,668	725	2,235	3,060	429
Specialties (top 8)											
General practice	14,609	496	3,420	177	901	182	1,434	329	1,155	1,559	177
Orthopedic surgery	592	23	150	12	48	7	42	17	44	68	15
General surgery	734	38	162	7	65	17	51	14	45	99	8
Obstetrics and gynecology	825	32	168	10	59	7	82	14	47	114	9
Plastic surgery	232	11	70	5	22	5	18	^a	11	43	10
Emergency medicine	319	24	40	5	49	6	27	12	30	43	9
Urology	287	18	82	^a	34	^a	11	12	21	30	8
Otolaryngology	276	12	92	0	24	5	15	8	17	35	^a
Census division of practice											
Population density (people/km ²)	1,465.8	1,706.0	2,179.0	1,637.3	969.0	1,552.8	1,193.9	1,107.5	805.1	984.0	1,431.7
Percentage high school or less	44.4	43.1	43.7	43.8	44.7	43.9	45.5	44.1	42.9	46.0	44.2
Percentage unemployed	8.4	8.3	8.6	8.4	8.2	8.4	8.3	8.3	7.9	8.3	8.5
Percentage low-income quintile	22.1	23.0	23.7	23.1	21.8	22.7	22.0	21.6	20.5	21.0	23.2
Percentage rural	8.1	7.3	5.4	8.6	11.7	9.1	8.7	8.7	10.9	11.3	9.3
Percentage Northern Ontario ^b	5.7	5.0	3.8	7.5	6.0	7.3	7.1	6.6	9.5	4.8	9.3
	Full Sample	Ranked					Unranked			Foreign	
		U. Mon.	Laval	U. Sask.	U. Man	U. Sher.	MUN	NOSM	US	UKIANZ	Other
Number of Physicians	30,844	234	124	262	647	152	445	80	391	1,788	6,420
Specialties (top 8)											
General practice	14,609	106	42	94	307	52	180	79	128	804	2,987
Orthopedic surgery	592	4	^a	7	14		17	0	6	33	80
General surgery	734	7	^a	5	9	5	9	0	5	43	142
Obstetrics and gynecology	825	9	^a	^a	11	10	12	0	10	38	187
Plastic surgery	232	^a	^a	^a	^a	^a	^a	0	^a	8	12
Emergency medicine	319	^a	^a	^a	^a	^a	10	0	18	6	26
Urology	287	^a	^a	^a	5	^a	^a	0	^a	12	39
Otolaryngology	276	^a	^a	^a	^a	7	5	0	^a	17	26
Census division of practice											
Population density (people/km ²)	1,465.8	704.8	1,019.6	1,219.3	1,589.9	858.1	927.5	86.2	2,126.5	1,433.7	1,463.3
Percentage high school or less	44.4	44.0	42.2	44.1	44.8	42.4	44.9	49.1	44.3	44.9	44.5
Percentage unemployed	8.4	7.7	7.9	8.4	8.6	7.6	8.2	8.6	8.7	8.5	8.5
Percentage low-income quintile	22.1	20.2	21.4	22.3	23.1	19.6	21.2	23.2	24.7	22.4	21.2
Percentage rural	8.1	13.7	8.7	7.4	12.0	9.6	10.9	32.6	6.4	8.9	5.7
Percentage Northern Ontario ^a	5.7	10.7	8.9	9.2	14.2	5.3	6.5	70.0	3.6	6.7	3.8

Note. UofT = University of Toronto; UBC = University of British Columbia; MAC = McMaster University; DAL = Dalhousie University; U. Mon. = Université de Montréal; U. Sask. = University of Saskatchewan; U. Man. = University of Manitoba; U. Sher. = Université de Sherbrooke; NOSM = Northern Ontario School of Medicine; UKIANZ = United Kingdom, Ireland, Australia, and New Zealand; MUN = Memorial University.

^aSuppressed due to cell count less than 5.

^bIncludes North West and North East Local Health Integration Networks.

list of coefficients). We also estimated regressions by specialty and graduation cohort but found no gradient in either case; few coefficients were statistically significant (results are available upon request).

Sensitivity Analysis

We found similar results using number of opioids dispensed as the outcome. We found a weak negative relationship and few statistically significant coefficients in the adjusted analysis. Findings were qualitatively the same for PCPs only and excluding physician-years with no opioid prescriptions and by specialty and graduation cohort (results are available upon request). Moreover, our main results did not change when we excluded all OMT-related prescriptions. The largest coefficients for the “all physicians” model were for other foreign schools (21.01, P value < 0.001), followed by the University of Western Ontario (15.90, P value < 0.001), and the University of Toronto (12.95, P value < 0.001) including zeroes; this was largely the case for the “PCPs only” model (other foreign schools, 50.65, P value < 0.001; University of Toronto, 21.48, P value < 0.001; University of Western Ontario, 18.83, P value < 0.001). When we excluded physician-years with no opioid prescriptions, the largest coefficients for the “all physicians” model were for other foreign schools (22.98, P value < 0.001), the University of Western Ontario (16.82, P value < 0.001), and the University of Toronto (14.71, P value < 0.001); this was also the case for the “PCPs only” model (other foreign schools, 48.69, P value < 0.001; the University of Toronto, 22.47, P value < 0.001; and the University of Western Ontario, 18.93, P value < 0.001; see Tables A9 to A12 in the Online Appendix for full list of coefficients).

Discussion

Many developed countries are currently facing an opioid epidemic. It has been hypothesized that medical schools may have different approaches to pain management and addiction training, which may influence appropriate opioid prescribing among their trainees. This analysis sought to understand whether physician training played a role within this context in Ontario as found in the United States. The results suggest the role of medical school training on opioid prescribing among physicians practicing in Ontario is weaker than in the United States likely due to greater homogeneity of curricula among Canadian schools. We found a weak negative relationship between annual opioid prescriptions and medical school rank, where higher ranked schools prescribed more opioids. This finding held when we stratified the analysis by specialty and graduation cohort and when we examined the number of opioids dispensed and excluded OMT-related prescriptions. Nonetheless, we found some outliers—physicians and PCPs trained at non-English-speaking foreign schools and PCPs trained at the University of Manitoba tended to prescribe slightly

more opioids. This was also the case when we considered non-OMT prescriptions only, with the exception of the University of Manitoba.

Schnell and Currie (2018) found a positive relationship between the number of opioids prescribed and medical school rank in the United States, where physicians trained at top medical schools, such as Harvard, prescribed less than those trained at lower ranked schools.² While Harvard-trained physicians wrote an average of 95 opioid prescriptions per year (including zeroes), physicians from the lowest ranked U.S. medical schools wrote over 3 times more (299). Moreover, the authors found that foreign doctors wrote fewer opioid prescriptions than U.S.-trained physicians (albeit with differences by world region), while physicians from unranked U.S. schools were more similar to physicians from the lowest ranked schools. Our results differ from theirs. First, we found a weak negative relationship between opioid prescriptions and medical school rank even after controlling for specialty and location of practice. This may be due, in part, to there being fewer medical schools in Canada (17 vs. 147 in the United States) and less heterogeneity among medical school curricula. Second, the average number of opioid prescriptions written by Ontario physicians was similar to (or lower than) that of top-ranked U.S. medical schools. Third, physicians trained at non-English-speaking foreign schools prescribed more opioids than those trained at Canadian- or English-speaking foreign schools. This highlights the need to ensure physicians trained abroad undergo additional training around pain management and addiction. Also, worth noting, Ontario physicians trained at Québec medical schools were among the lowest opioid prescribers. Other work has found that Québec is one of the provinces with the lowest rates of opioid use.¹⁵ Thus, there is likely a different culture toward opioid prescribing in Québec, which may be worth exploring further. Residency training, fellowships, and continuing medical education should also be considered as they are likely to play a larger role than initial medical training (although addiction fellowship are relatively new). Unfortunately, this information was not available. In an effort to address this, we examined whether physicians trained at lower ranked medical schools were systematically more likely to practice in specialties where patient need for opioids might be higher. We did not find this to be the case. The potential influence of pharmaceutical firms on medical school training pain and prescription curriculum may have also played a role, but again, our analysis was limited by the existing data.

There have been recent efforts on behalf of Canadian medical schools and physician training programs to improve training content in pain management, opioid prescribing, and addiction/substance use disorders for early career practitioners and to provide continuing education to all practicing physicians¹⁶; this may explain the decline in opioid prescriptions in recent years. In November 2016, Health Canada and the then Federal Minister of Health, Dr Jane Philpott, held the “Summit on Problematic Opioid Use,” which included




Figure 2. Estimated mean difference (relative to McGill University) in the number of opioid prescriptions by medical school, controlling for specialty and census division of practice. (A) All physicians. (B) Primary care physicians. McGill University is the reference case. See Tables A5 and A6 in the Online Appendix for values for all physicians and primary care physicians, respectively.

the Association of Faculties of Medicine of Canada. Since then, the Association of Faculties of Medicine of Canada, which includes all 17 faculties of medicine, has committed to improving foundational core competencies in medical education around opioid prescribing and pain management. Furthermore, in 2017, guidelines were developed to inform opioid prescribing for adults with chronic noncancer pain.¹⁷ In Ontario, the Narcotics Monitoring System was introduced in May 2012 to collect and monitor data on all narcotics dispensed.¹⁸ The introduction of a fentanyl patch-for-patch program,¹⁹ which requires patients prescribed fentanyl return used patches to pharmacies before receiving more patches, and the delisting of high-strength opioids from the Ontario formulary (except for palliative patients)²⁰ have also helped curb opioid prescribing.

To our knowledge, this is the first study to examine this issue outside the United States and thus contributes to the

emerging literature around policies/interventions to address the opioid epidemic. We used a population-based sample of all physicians practicing in Ontario and examined all opioids prescribed, regardless of form, strength, and/or duration. Furthermore, we examined the number of opioids dispensed, which was not available in previous work.² Our work has some limitations. We were unable to examine recent graduates (i.e., those who qualified since 2011), and many physicians had missing data on medical school of training. Furthermore, among excluded physicians, the average number of prescriptions was lower than that of included physicians. Thus, there may be some sample selection bias, which we did not control for; however, this would have been more problematic had we found an effect. Data on medical school rankings were only available for more recent years; furthermore, despite research suggesting that the general procedure used by *Maclean's* is sound, these

rankings are subject to criticism.²¹ We were only able to observe where each physician completed their medical school training and not where they obtained specialty training, which limited our analysis. We were not able to adjust for physician volume, clinical need of patients, or in-hospital opioid administration; the inclusion of these variables may have reduced model coefficients. Moreover, we did not control for duration of prescriptions; however, the proportion of short-term prescriptions (5 days or less) did not differ much among schools. We examined Ontario data that only include 36% of all physicians in practicing in Canada.²² Results may have differed had we used data for the entire country. Finally, we did not examine other health providers who prescribe opioids such as dentists, nurses/nurse practitioners, midwives, and chiropodists. Although physicians make up most prescribers, dental prescribers in Ontario are responsible for about 20% to 25% of opioid prescriptions (although these are typically of short duration and low dose).²³

In sum, our results suggest that the role of medical school training on opioid prescribing patterns among physicians in Ontario is not as strong as in the United States. Moreover, the average number of opioids prescribed by Ontario physicians is similar to that of top-ranked U.S. medical schools. Nonetheless, ensuring that physicians trained abroad receive additional pain management and addiction training may help address part of the current opioid epidemic in Ontario and potentially elsewhere in Canada. Future work should seek to explore prescribing patterns among other health professionals and replicate this analysis in other jurisdictions where opioid use is also problematic, such as Alberta²⁴ and Australia,²⁵ for example.

Authors' Note

This study was supported by ICES which is funded by an annual grant from the Ontario Ministry of Health. In 2018, the institute formerly known as the Institute for Clinical Evaluative Sciences formally adopted the initialism ICES as its official name. This change acknowledges the growth and evolution of the organization's research since its inception in 1992 while retaining the familiarity of the former acronym within the scientific community and beyond. The opinions, results, and conclusions reported in this article are those of the authors and are independent from the funding sources. No endorsement by ICES or the Ontario Ministry of Health is intended or should be inferred. Parts of this material are based on data and/or information compiled and provided by the Canadian Institute for Health Information (CIHI). However, the analyses, conclusions, opinions, and statements expressed in the material are those of the author(s) and not necessarily those of CIHI.

Data Access

The data set from this study is held securely in coded form at ICES. While data sharing agreements prohibit ICES from making the data set publicly available, access may be granted to those who meet prespecified criteria for confidential access available at www.ices.on.ca/DAS. The full data set creation plan and underlying analytic

code are available from the authors upon request, understanding that the computer programs may rely upon coding templates or macros that are unique to ICES and are therefore either inaccessible or may require modification.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iDs

Claire de Oliveira, MA, PhD  <https://orcid.org/0000-0003-3961-6008>

Paul Kurdyak, MD, MSc, PhD, FRCP(C)  <https://orcid.org/0000-0001-8115-7437>

Supplemental Material

Supplemental material for this article is available online.

References

- Gomes T, Juurlink DN, Dhalla IA, Mailis-Gagnon A, Paterson JM, Mamdani MM. Trends in opioid use and dosing among socio-economically disadvantaged patients. *Open Med.* 2011; 5(1):e13-e22.
- Schnell M, Currie J. Addressing the opioid epidemic: is there a role for physician education? *Am J Health Econ.* 2018;4(3): 383-410.
- Pain and Policy Studies Group. Opioid consumption motion chart; 2016 [accessed 2018 Jul 31]. <https://ppsg.medicine.wisc.edu/chart>.
- Rudd RA, Seth P, David F, Scholl L. Increases in drug and opioid-involved overdose deaths—United States, 2010-2015. *Morb Mortal Wkly Rep (MMWR).* 2016; 65(5051):1445-1452.
- Gomes T, Greaves S, Martins D, et al. Latest trends in opioid-related deaths in Ontario: 1991 to 2015. Toronto (ON): Ontario Drug Policy Research Network; 2017.
- British Columbia Coroners Service. Illicit drug overdose deaths in BC. 2007 January 1–2017 March 31. Burnaby (BC): British Columbia Coroners Service; 2017.
- Gomes T, Khuu W, Martins D, et al. Contributions of prescribed and non-prescribed opioids to opioid related deaths: population based cohort study in Ontario, Canada. *BMJ.* 2018;362(4):k3207.
- Government of Canada. Responding to Canada's opioid crisis; 2018 [accessed 2018 Jul 31]. <https://www.canada.ca/en/health-canada/services/substance-abuse/prescription-drug-abuse/opioids/responding-canada-opioid-crisis.html> Date modified: 2018-05-18.
- Gomes T, Juurlink DN. Opioid use and overdose: what we've learned in Ontario. *Healthc Q.* 2016;18(4):8-11.
- Barnett M, Olenski A, Jena A. Opioid-prescribing patterns of emergency physicians and risk of long-term use. *N Engl J Med.* 2017;376(2):663-673.

11. Maclean's University Rankings 2018: Canada's top Medical/ Doctoral schools; 2018. [accessed 2018 Jul 31]. <https://www.macleans.ca/education/university-rankings/medical-doctoral-universities/>.
12. Maclean's University Rankings. 2017: How Maclean's selects Canada's top schools; 2017. [accessed 2018 Jul 31]. <http://www.macleans.ca/education/university-rankings-2017-how-macleans-selects-canadas-top-schools/>.
13. Statistics Canada. Postal Code^{OM} Conversion File (PCCF), Reference Guide; 2015 [accessed 2018 Jul 31]. <http://www.statcan.gc.ca/pub/92-154-g/92-154-g2015001-eng.htm> Date modified: 2015-11-30.
14. Gomes T, Pasricha S, Martins D, et al. Behind the prescriptions: A snapshot of opioid use across all Ontarians. Toronto (ON): Ontario Drug Policy Research Network; 2017 [accessed 2018 July 31]. <http://www.stmichaelshospital.com/pdf/ODPRN-report-opioids.pdf>.
15. Public Health Agency of Canada. Special Advisory Committee on the Epidemic of Opioid Overdoses. National report: Apparent opioid-related deaths in Canada (January 2016 to September 2017) web-based report. Ottawa (ON): Public Health Agency of Canada; 2018.
16. The Association of Faculties of Medicine of Canada. Report on the AFMC response to the Canadian opioid crisis; 2017 [accessed 2018 Jul 31]. https://afmc.ca/sites/default/files/documents/2017-11-AFMC-HealthCanadaOpioidReport_en.pdf.
17. Busse JW, Craigie S, Juurlink DN, et al. Guideline for opioid therapy and chronic noncancer pain. *CMAJ*. 2017;189(18):E659-E666.
18. Ministry of Health and Long-Term Care Narcotic Monitoring System (NMS) Pharmacy Reference Manual; 2012. [accessed 2020 Feb 20]. http://www.health.gov.on.ca/en/pro/programs/drugs/resources/narcotics_manual.pdf.
19. Tadrous M, Greaves S, Martins D, et al. Evaluation of the fentanyl patch-for-patch program in Ontario, Canada. *Int J Drug Policy*. 2019;66(1):82-86.
20. Martins D, Khuu W, Tadrous M, et al. Impact of delisting high-strength opioid formulations from a public drug benefit formulary on opioid utilization in Ontario, Canada. *Pharmacoepidemiol Drug Saf*. 2019;28(5):726-733.
21. Nancy H, Rick W, Steven W, Hubert W. Maclean's ranking of Canadian universities, UBC Statistics Department Technical Report 189; 1999 [accessed 2019 Aug 22]. <https://www.stat.ubc.ca/technical-reports-archive/doc/189.pdf>.
22. Canadian Medical Association. Number of physicians by province/territory and specialty, Canada, 2018; 2018 [accessed 2018 Jul 31]. <https://www.cma.ca/Assets/assets-library/document/en/advocacy/01-physicians-by-specialty-province-e.pdf>.
23. Health Quality Ontario. Starting on opioids; 2018 [accessed 2018 Jul 31]. <https://www.hqontario.ca/Portals/0/Documents/system-performance/starting-on-opioids-en.pdf>.
24. Gomes T, Paterson JM, Caetano P, Sketris I, Henry D. CNODES analysis: Safety of oral opioid use in Canada. Part 1: Changes in the dispensing of oral opioid drugs in Canadian provinces between 2008 and the end of 2013. *Canadian Network for Observational Drug Effect Studies (CNODES)*; 2015 [accessed 2020 Feb 20]. https://www.researchgate.net/publication/279204495_CNODES_analysis_Safety_of_oral_Opioid_use_in_Canada_Part_1_Changes_in_the_dispensing_of_oral_opioid_drugs_in_Canadian_provinces_between_2008_and_the_end_of_2013.
25. Rintoul AC, Dobbin MD, Drummer OH, Ozanne-Smith J. Increasing deaths involving oxycodone, Victoria, Australia, 2000-09. *Injury Prevent*. 2011;17(4):254-259.