



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/165440/>

Version: Published Version

Article:

Kelly, C., Morgan, G., Bannard, C. et al. (2020) Early pragmatics in deaf and hard of hearing infants. *Pediatrics*, 146 (Supplement 3). S262-S269. ISSN: 0031-4005

<https://doi.org/10.1542/peds.2020-0242E>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Early Pragmatics in Deaf and Hard of Hearing Infants

Ciara Kelly, PhD,^a Gary Morgan, PhD,^b Colin Bannard, PhD,^c Danielle Matthews, PhD^a

abstract

BACKGROUND AND OBJECTIVES: A set of important pragmatic skills emerge during infancy and pave the way for later language learning. It is thought these early social communication skills develop through infant–caregiver interaction. In a microanalysis, we tested whether deaf and hard of hearing (DHH) infants (typically at high risk of reduced access to rich communicative interaction in infancy) are less likely to engage in gestural and vocal pragmatic behaviors.

METHODS: We coded the naturalistic communication of 8 DHH infants who had no additional needs, who were not preterm or low birth weight, whose parents were hearing, monolingual English speakers, and who had spoken English as their primary target language. The frequency of use of 5 types of infant communication known to positively predict later language development (show gestures, give gestures, index-finger pointing, communicative vocalizations, and early word use) was compared with that of 8 typically hearing infants matched for age, sex, and socioeconomic status.

RESULTS: Hearing loss had a significant negative effect on the frequency with which infants engaged in all types of early communication that predict later language development.

CONCLUSIONS: DHH infants are at high risk of delay in the gestural and vocal communicative skills that lay the foundations for later language. Delay in the gestural domain suggests this is not simply a consequence of difficulties in imitating auditory stimuli. There is significant potential to lift DHH infants onto a positive developmental trajectory by supporting caregivers to nurture interaction from the first year.

WHAT'S KNOWN ON THIS SUBJECT: How often typically hearing infants engage in pragmatic behaviors (like pointing) predicts language development. These early skills are thought to develop through interaction, but evidence is mixed regarding whether deaf and hard of hearing infants show early pragmatic delay.

WHAT THIS STUDY ADDS: Deaf and hard of hearing infants aged 12 to 18 months engage less frequently than matched, typically hearing peers in early gestural and vocal communicative behaviors that predict later spoken language. Caregivers need support from infancy to nurture pragmatic development.

^aDepartment of Psychology, University of Sheffield, Sheffield, United Kingdom; ^bDivision of Language & Communication Science, City, University of London, London, United Kingdom; and ^cDepartment of Psychology, University of Liverpool, Liverpool, United Kingdom

Dr Kelly designed the study, reviewed the literature, collected the data, adapted the coding scheme, led coding, and cowrote and revised the initial manuscript; Dr Bannard conducted the statistical analyses and critically reviewed and revised the manuscript; Prof Morgan critically reviewed and revised the manuscript; Dr Matthews conceptualized and designed the study, supervised data collection and analysis, acquired funding, and cowrote and revised the initial manuscript; and all authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

DOI: <https://doi.org/10.1542/peds.2020-0242E>

Accepted for publication Aug 31, 2020

Address correspondence to Colin Bannard, PhD, Department of Psychology, University of Liverpool, 2nd Floor, Eleanor Rathbone Building, Bedford St S, Liverpool L69 7ZA, United Kingdom. E-mail: colin.bannard@liverpool.ac.uk

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2020 by the American Academy of Pediatrics

To cite: Kelly C, Morgan G, Bannard C, et al. Early Pragmatics in Deaf and Hard of Hearing Infants. *Pediatrics*. 2020;146(Supplement_3):e20200242E

The development of early pragmatic (ie, social communicative) skills during infancy paves the way for later language.^{1,2} These skills are thought to emerge in the context of infant-caregiver interaction, during which caregivers intuitively scaffold communicative development.³ For example, hearing parents intuitively respond to their infants using speech and vocal cues (eg, gasps and exclamations to indicate interest), and, by 5 months, typically hearing infants come to expect a vocal response to their babble.⁴ Furthermore, fluent signing deaf parents intuitively use visual-tactile strategies during interaction to indicate interest and support visual perception of language, and, by 24 months, deaf and hard of hearing (DHH) infants of fluent signing deaf parents have learned to look to their caregivers' faces more often than infants from hearing dyads do.^{5,6} However, ~95% of DHH children are born to hearing parents,⁷ who typically are not fluent sign language users, have little to no experience communicating with a DHH infant, and, for the majority of families, have spoken language as the goal for their child (typically, providing an oral, sign-supported, or sign-spoken bilingual environment⁸).

Hearing parents with a DHH infant often find scaffolding communicative development challenging, whether targeting a spoken language, a signed language, or both. When learning a signed language, hearing parents can struggle with fluency and the adaptations necessary to support their DHH infant's visual perception of language.⁹⁻¹² When supporting development of a spoken language (with or without signed support), ensuring access to communication is often not immediately intuitive.^{9,13} Professional support for families to meet these challenges is highly variable,¹⁴ as is social support^{15,16} and parents' belief in their ability to support their child's communicative

development (ie, feelings of self-efficacy).^{17,18} Because of reduced access to language as well as the impact hearing loss can have on the quality of communicative interactions, delays in communicative development and later language learning are found in the majority of DHH infants.^{8,19,20} To date, however, it is unclear whether delays emerge in social communication during infancy; current evidence regarding delays in the key early communicative behaviors that pave the way for language development is sparse and equivocal.^{21,22}

THE RELATIONSHIP BETWEEN EARLY PRAGMATIC SKILLS AND LATER LANGUAGE DEVELOPMENT

In recent research with typically developing, typically hearing infants, researchers have identified specific early communicative behaviors as being theoretically important indicators of readiness for language learning and good empirical predictors of later language development.^{2,23-27} Positive predictors include the frequency with which infants produce give gestures, show gestures, index-finger pointing, and communicative vocalizations (operationalized as vocalizations made within 1 second of looking to the caregiver's face). Alongside first words, these behaviors are thought to be positive predictors because they indicate social-cognitive readiness for language development, and because they elicit timely and informative caregiver responses, which are known to scaffold further development.²⁸⁻³² The positive predictive value of these early communicative acts seems to be specific to these behaviors. Indeed, the frequency with which infants rely on some other gestures (eg, open-hand pointing) is a negative predictor of language development.^{25,27} Thus, when assessing early pragmatic development in DHH infants, it has become apparent that investigating

precise frequencies of specific infant naturalistic behaviors is important.

THE COMMUNICATION OF DHH INFANTS: CURRENT KNOWLEDGE

Relatively little is known about how hearing loss affects social communication in the first 2 years of life. There is evidence that DHH 22 month olds have difficulty in maintaining joint attention (a state arising when caregiver and infant are mutually aware of sharing attention to the same thing).^{19,33-35} However, in research with younger infants to date, it has been suggested that DHH infants communicate with their hearing parents at a broadly similar rate to their typically hearing peers,^{21,36,37} albeit with some evidence of a reduced frequency of gesture use.²² A limitation of these studies is that researchers have collapsed across a broad range of behaviors (likely including both positive and negative predictors of spoken language in one count) or have employed binary measures of whether a behavior is in repertoire, rather than measuring the frequency with which that behavior is used for communication. It is the latter that is known to predict outcomes.²⁷ Moreover, although DHH infants' vocalizations have been explored in terms of their phonology (with evidence of delay^{38,39}), there has been little study of the communicative use of vocalizations (regardless of phonological properties). It has recently been found that one of the most valuable predictors of typically hearing infants' transition to speech is the frequency with which they produce communicative vocalizations that are then responded to by the caregiver with language that is relevant to the infant's focus of attention. An infant who benefits from such interactions with a frequency 1 SD above the mean at 12 months is predicted to produce ~28 more words than the average infant by 19 months.²⁷

THE CURRENT STUDY

Motivated by the evidence for the predictive value of specific early pragmatic indicators for later language development and the gap in knowledge about the development of these skills in DHH infants, we used a fine-grained coding scheme to measure the early naturalistic communication of DHH infants. We hypothesized that DHH infants who are at risk of reduced access to communicative interaction (in this case, those with hearing parents whose target primary language was spoken English) would engage less frequently than closely matched, typically hearing infants in the communicative behaviors that are positive predictors of spoken language development. If such differences are observed, this would (1) provide health care professionals with a more complete picture of the developmental trajectory of DHH infants' early pragmatic skills and (2) suggest a need to start targeted support of pragmatic development during the first year of life.

METHOD

Participants

Participants were 8 DHH infants, 8 typically hearing infants, and their primary caregivers. The DHH infants were recruited across England and Scotland through the United Kingdom's National Deaf Children's Society database of families with DHH children. The typically hearing infants were participants in the control condition of a longitudinal randomized controlled trial,⁴⁰ in which researchers collected data using the same procedure as the current study. All caregivers gave informed consent for their data to be used for further research. Participation was subject to the following inclusion criteria: infants were term (born no more than 3 weeks before the due date) and with a birth weight >2.5 kg and no

other known disabilities or developmental delays. All caregivers had no known physical, mental, or learning disabilities. See Table 1.

Written informed consent was obtained from each caregiver. The Psychology Ethics Committee at the University of Sheffield approved this study.

Procedure

Infant-caregiver dyads were video recorded in free play together for 25 minutes in their home, from 2 different camera angles and without the researcher present.

Coding

Following the coding scheme reported in Donnellan et al,²⁷ video recordings were coded for infant gestures, vocalizations, gaze to caregiver's face, and recognizable British Sign Language (BSL) signs by using EUDICO Linguistic Annotator software (ELAN)⁴¹ by the first author. Gestures were categorized as either give, show, or index-finger point. All nonvegetative vocalizations (ie, vocalizations containing speech sounds as opposed to coughs, sneezes, hiccups, etc.^{42,43}) were categorized as either infant vocalization (ie, vocalizations with or without a consonant that were not

recognizable words) or words (ie, a recognizable word). Gaze-coordinated vocalizations were automatically identified as any infant vocalization occurring within 1 second of a gaze to the caregiver. Words were orthographically transcribed. BSL signs were translated to English and orthographically transcribed. Because only 2 infants produced signs (2 and 35 signs, respectively), these were not included in statistical analyses. A new coding pass of each video was made for each behavior type (ie, each video was coded 4 times, once each for gestures, vocalizations, gaze, and signs. See Fig 1). Any periods of time in which the infant was not within view of the cameras were identified and not coded. The observable coding time was <25 minutes (but not <21 minutes and 11 seconds) for 4 DHH infants and 6 typically hearing infants. On-screen time was controlled for in analyses.

Reliabilities

Twenty-five percent of video recordings (8 play sessions) were coded for reliability by a second coder. Counts of gestures, infant vocalizations, gaze to caregiver's face, and words were highly correlated ($r = 0.99$, $r = 1.00$, $r = 0.99$, and $r = 0.98$, respectively). Cohen's κ revealed that

TABLE 1 Characteristics of the DHH Infants

Participant	Sex	Age, mo; d	Left Ear	Right Ear	Age HAS Received, wk	Family Mode of Communication	IMD ^a Decile
1	Male	12; 21	Moderate-severe	Moderate	5	Spoken English	6
3	Male	13; 07	Moderate-severe	Profound	8	Mostly SSE	5
6	Female	12; 25	Profound	Severe	12	Spoken English and occasionally SSE	6
9	Female	12; 30	Severe-profound	Moderate-severe	9	Spoken English and SSE	7
4	Male	19; 09	Moderate	Moderate	6	Spoken English and rarely SSE	6
5	Male	18; 17	Severe	Severe	12	Spoken English	4
7	Male	18; 12	Moderate	Moderate	15	Spoken English	2
8	Male	18; 20	Moderate	Severe	8	Spoken English	7

IMD, Indices of Multiple Deprivation; SSE, sign-supported English.

^a Areas considered to be within the most deprived 10% of England and Scotland = 1; areas considered to be within the least deprived 10% of England and Scotland = 10.




FIGURE 1
Example of video coding in EUDICO Linguistic Annotator software (ELAN).

coders categorized gestures as either gives, shows, or index-finger points with high reliability ($k = 0.97$; $P < .001$).

Statistical Analysis

To assess whether the DHH infants raised with spoken English as their primary target language differed from typically hearing infants in their production of communicative acts known to predict later spoken language development, we fitted 2 multilevel negative binomial regression models to the data (using lme4 version 1.1-23). This model type was chosen because the outcome variable is a count measure but does not meet the assumption for a Poisson regression (equality of variance and mean). A set of counts for each of the communicative acts (give and show gestures, index-finger pointing, gaze-coordinated vocalizations, and early word use) was entered as the dependent variable, hearing status as a fixed effect (DHH or typical-hearing), participant and age as random effects on the intercept, and communication

type as random effects on the intercept and slope. In model 1, we included all prelinguistic acts alongside early word use because this provided the best overall picture of infants' readiness for spoken language. In model 2, given that DHH infants have been observed to show a delay in the onset of spoken word use at 18 months,^{21,44} we then restricted the analysis by removing word production from the set of communicative behaviors, thereby focusing on prelinguistic communicative acts. In both models, we included a scaled and centered on-screen time variable as a control because infants varied in how much on-screen coding time they had (4 DHH and 6 typically hearing infants <25 minutes). Whether we included on-screen time in this way or as an exposure offset with a coefficient of 1 did not affect outcomes.

RESULTS

The demographic characteristics of the 8 DHH infants are shown in Table 1. All were born with bilateral

permanent hearing loss identified within 3 weeks of birth through newborn hearing screening and received bilateral hearing aids (HAs) by 15 weeks. All were continuing to use bilateral HAs at the time of testing. Hearing loss ranged from moderate (41–70 dB) to profound (>95 dB). Information on specific hearing thresholds (aided or unaided) was not available. Four infants were aged between 12 and 13 months (2 boys), and 4 were aged between 18 and 19 months (all boys). The 8 typically hearing infants were closely matched to the DHH infants for age, sex, and socioeconomic status (determined by using the English and Scottish Indices of Multiple Deprivation: a government measure based on neighborhood income, employment, health provision, and housing). All caregivers were female, typically hearing, and monolingual English-speaking. Four parents were learning BSL to provide sign support for spoken English.

In Table 2, we present descriptive statistics by group for the specific

communicative acts under study. Relative to the typically hearing group, the DHH group produced all communicative acts with a lower median frequency.

Hearing loss had a significant negative effect on the frequency of infant communication (model 1: $b = -0.853$; 95% confidence interval: -1.515 to -0.192 ; $P = .01$). According to model estimates, DHH infants produced 5 fewer communicative behaviors in 25 minutes of play than their matched typically hearing peers. This effect held when the data set was restricted to prelinguistic communicative acts only (model 2 [removing words]: $b = -0.770$; 95% confidence interval: -1.516 to -0.024 ; $P = .04$). According to model estimates, DHH infants produced 4 fewer prelinguistic communicative behaviors in 25 minutes of play than their matched typically hearing peers.

DISCUSSION

We analyzed the early communicative behaviors of infants with moderate-to-profound hearing loss who had no additional needs, who were not preterm or low birth weight, whose parents were hearing, monolingual English speakers, and who had spoken English as their primary target language. These infants were closely matched demographically with typically hearing infants. Fine-grained, naturalistic observation of the specific communicative behaviors that positively predict later spoken language development suggested that

hearing loss puts infants at significant risk of delay. Analyses restricted only to prelinguistic communication suggested that the risk extends to those early gestural and vocal communicative developments that pave the way for language. By producing fewer communicative behaviors during interaction, DHH infants are likely to experience fewer learning opportunities, which cumulatively would affect later language learning.

Where do these early gestural and vocal communicative skills come from? Their development is thought to depend on frequent, finely timed interactions between caregiver and infant during earlier stages of development⁴⁵⁻⁴⁷ (see Mood et al, this supplement). Because both vocal and gestural prelinguistic communication was affected, we assume that reduced communication was not simply the product of reduced imitation of verbal stimuli: this would not explain gestural delay. Rather, communicative tools, including pointing gestures, are likely not being used as frequently by DHH infants because of (1) reduced access to timely caregiver reinforcement of their early communication and (2) reduced access to cues that support joint attention and, thereby, provide insights into how people use gestures and vocalizations to communicate.⁴⁸ Hearing parents typically use nonlinguistic vocal cues (eg, gasps and exclamations) to regulate interaction, alert the infant to a topic of interest, initiate joint attention

with their infant, and respond to them. These nonlinguistic vocal cues are used alongside gestures. By using sound and gesture in predictable, synchronous ways,⁴⁹ hearing parents intuitively make it easier for infants to understand how communicative acts can direct others' attention. However, DHH infants are more likely to miss some of these cues, and, so, early joint attentional episodes are more likely to break down before learning occurs.^{33,50,51} Furthermore, hearing parents tend not to use visual cues as frequently or as proficiently in comparison with deaf fluent signing parents, which again places joint attention at risk.^{10,34,52} The critical point, here, is that this study strongly suggests that access to sound is important not just for learning spoken language through imitation but for the nonlinguistic uses of sound that hearing parents intuitively use in synchrony with other cues to regulate the back-and-forth of the earliest interactions.

The implications of these findings are that families need support to nurture specific pragmatic developments during the first 2 years of life (namely the appropriate communicative interaction around infant give gestures, show gestures, index-finger pointing, communicative vocalizations, and, eventually, words and signs). A major challenge in this work will be engaging sensitively with parents at what, for many, is an emotionally turbulent time,⁵³⁻⁵⁵ when not all families are ready to participate in interventions or research.

TABLE 2 Descriptive Statistics for Infant Production of Communicative Acts as a Function of Hearing Status

Variable	DHH Group (N = 8)					Typically Hearing Group (N = 8)				
	Mean	SD	Median	Minimum to Maximum	% scoring >0	Mean	SD	Median	Minimum to Maximum	% scoring >0
Give gestures	3.88	4.05	2.50	0-12	75 (n = 6)	9.75	9.59	8.50	0-26	88 (n = 7)
Show gestures	1.13	2.10	0.00	0-6	38 (n = 3)	4.75	5.39	3.50	0-16	75 (n = 6)
Index-finger points	3.00	5.81	0.50	0-17	50 (n = 4)	6.50	8.07	2.00	0-19	63 (n = 5)
Gaze-coordinated vocalizations	15.75	11.31	15.00	6-42	100 (n = 8)	25.00	20.56	19.50	5-61	100 (n = 8)
Word production	6.63	11.56	0.50	0-29	50 (n = 4)	22.00	30.92	1.50	0-74	63 (n = 5)

Descriptive statistics were prorated for the amount of on-screen coding time from a 25 min recording.

In this study, we took the approach of exploring a fine-grained set of behaviors in a hard-to-reach group of infants that was as homogenous as possible (eg, no known language development risk factors other than hearing loss). Although this allowed us to control for potential confounding variables, this had the effect of limiting the sample size and generalizability beyond infants with these characteristics. A complementary approach would be to include a broader range of infants (including those with cochlear implants and those primarily learning to sign) and then control for additional characteristics. Further research will allow us to fully understand the source of the current observed differences with (1) longitudinal studies used to predict

DHH child outcomes from both early infant communication and caregiver practice and (2) trials of early intensive intervention to assess how support programs change parenting behaviors and child outcomes. These interventions would need to support all aspects of early communication, including pragmatic aspects of language development.⁵⁶

In sum, the current study used a fine-grained analysis of infant pragmatic development and found DHH infants are at significant risk of delay in precisely the types of early gestural and vocal communication that are important for future language development. The observation that DHH children are at risk for reduced use of gestures from infancy is novel and striking.

The implications for health care providers and allied health professionals are clear. By responding to the call to action⁵⁷ from the first year of a DHH child's life (ie, screening for risks to infant-caregiver interaction and supporting caregivers to nurture early social communication), there is real potential to lift children onto a positive learning trajectory and open the way to the cumulative learning experiences that build from these early moments.

ABBREVIATIONS

BSL: British Sign Language
DHH: deaf and hard of hearing
HA: hearing aid

This article is licensed under the Creative Commons Attribution 4.0 International License (CC-BY 4.0). The named authors alone are responsible for the views expressed in this publication.

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: The first phase of this work was supported by a University of Sheffield scholarship and Funds for Women Graduates Foundation Main Grant awarded to Ciara Kelly. The second phase was supported by an Alan Kelly Legacy to the University of Sheffield. Danielle Matthews was supported by British Academy grant MD\170025.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

REFERENCES

1. Adamson LB. *Communication Development During Infancy*. Madison, WI: Brown & Benchmark; 1995
2. Tomasello M. *Origins of Human Communication*. Cambridge, MA: The MIT Press; 2008
3. Papoušek H, Bornstein MH. Didactic Interactions: Intuitive Parental Support of Vocal and Verbal Development in Human Infants. In: Papoušek H, Jürgens U, Papoušek M, eds. *Nonverbal Vocal Communication: Comparative and Developmental Approaches*. New York, NY: Cambridge University Press; 1992:209–229
4. Goldstein MH, Schwade JA, Bornstein MH. The value of vocalizing: five-month-old infants associate their own noncry vocalizations with responses from caregivers. *Child Dev*. 2009;80(3):636–644
5. Lieberman AM, Hatrak M, Mayberry RI. Learning to look for language: development of joint attention in young deaf children. *Lang Learn Dev*. 2014; 10(1)
6. Brooks R, Singleton JL, Meltzoff AN. Enhanced gaze-following behavior in deaf infants of deaf parents. *Dev Sci*. 2020;23(2):e12900
7. Mitchell RE, Karchmer MA. Chasing the mythical ten percent: parental hearing status of deaf and hard of hearing students in the United States. *Sign Lang Stud*. 2004;4(2):138–163
8. Lederberg AR, Schick B, Spencer PE. Language and literacy development of deaf and hard-of-hearing children: successes and challenges. *Dev Psychol*. 2013;49(1):15–30
9. Koester LS. Intuitive parenting as a model for understanding parent-infant interactions when one partner is deaf. *Am Ann Deaf*. 1992;137(4):362–369
10. Waxman R, Spencer P. What mothers do to support infant visual attention: sensitivities to age and hearing status. *J Deaf Stud Deaf Educ*. 1997;2(2): 104–114
11. Arnesen K, Enerstvedt RT, Engen EA, Engen T, Høie G, Vonen AM. The linguistic milieu of Norwegian children with hearing loss. *Am Ann Deaf*. 2008; 153(1):65–77
12. DeLana M, Gentry MA, Andrews J. The efficacy of ASL/English bilingual education: considering public schools. *Am Ann Deaf*. 2007;152(1):73–87

13. Beatrijs W, Kristiane VL, Mieke VH. Parental strategies used in communication with their deaf infants. *Child Lang Teach The*. 2019;35(2): 165–183
14. Rees R, Mahon M, Herman R, Newton C, Craig G, Marriage J. Communication interventions for families of pre-school deaf children in the UK. *Deafness Educ Int*. 2015;17(2):88–100
15. MacTurk RH, Meadow-Orlans KP, Koester LS, Spencer PE. Social support, motivation, language, and interaction. A longitudinal study of mothers and deaf infants. [published correction appears in *Am Ann Deaf*: 1993;138(3):308]. *Am Ann Deaf*. 1993;138(1):19–25
16. Meadow-Orlans KP, Steinberg AG. Effects of infant hearing loss and maternal support on mother-infant interactions at 18 months. *J Appl Dev Psychol*. 1993; 14(3):407–426
17. DesJardin JL. *Maternal Self-Efficacy and Involvement: Supporting Language Development in Young Deaf Children With Cochlear Implants [PhD thesis]*. Los Angeles, CA: University of California; 2004
18. DesJardin JL, Eisenberg LS. Maternal contributions: supporting language development in young children with cochlear implants. *Ear Hear*. 2007;28(4): 456–469
19. Lederberg AR, Everhart VS. Conversations between deaf children and their hearing mothers: pragmatic and dialogic characteristics. *J Deaf Stud Deaf Educ*. 2000;5(4):303–322
20. Levine D, Strother-Garcia K, Golinkoff RM, Hirsh-Pasek K. Language development in the first year of life: what deaf children might be missing before cochlear implantation. *Otol Neurotol*. 2016;37(2):e56–e62
21. Spencer PE. Communication behaviors of infants with hearing loss and their hearing mothers. *J Speech Hear Res*. 1993;36(2):311–321
22. Robinshaw HM. The pattern of development from non-communicative behaviour to language by hearing impaired and hearing infants. *Br J Audiol*. 1996;30(3):177–198
23. Beuker KT, Rommelse NNJ, Donders R, Buitelaar JK. Development of early communication skills in the first two years of life. *Infant Behav Dev*. 2013; 36(1):71–83
24. Boundy L, Cameron-Faulkner T, Theakston A. Exploring early communicative behaviours: a fine-grained analysis of infant shows and gives. *Infant Behav Dev*. 2016;44:86–97
25. Lüke C, Grimminger A, Rohlfing KJ, Liskowski U, Ritterfeld U. In infants' hands: identification of preverbal infants at risk for primary language delay. *Child Dev*. 2017;88(2):484–492
26. McKean C, Law J, Mensah F, et al. Predicting meaningful differences in school-entry language skills from child and family factors measured at 12 months of age. *Int J Early Child*. 2016;48(3):329–351
27. Donnellan E, Bannard C, McGillion ML, Slocombe KE, Matthews D. Infants' intentionally communicative vocalizations elicit responses from caregivers and are the best predictors of the transition to language: a longitudinal investigation of infants' vocalizations, gestures and word production. *Dev Sci*. 2020;23(1):e12843
28. McGillion ML, Herbert JS, Pine JM, Keren-Portnoy T, Vihman MM, Matthews D. Supporting early vocabulary development: what sort of responsiveness matters? *IEEE Trans Auton Ment Dev*. 2013;5(3):240–248
29. Tamis-LeMonda CS, Kuchirko Y, Song L. Why is infant language learning facilitated by parental responsiveness? *Curr Dir Psychol Sci*. 2014;23(2): 121–126
30. Yoder PJ, Warren SF. Can Developmentally Delayed Children's Language Development Be Enhanced through Prelinguistic Intervention? In: Kaiser AP, Gray DB, eds. *Enhancing Children's Communication: Research Foundations for Intervention*. Baltimore, MD: Brookes Publishing; 1993:35–62
31. Yoder PJ, Warren SF. Maternal responsivity mediates the relationship between prelinguistic intentional communication and later language. *J Early Interv*. 1999;22(2):126–136
32. Bruner JS. From communication to language – a psychological perspective. *Cognition*. 1976;3(3):255–287
33. Prezbindowski AK, Adamson LB, Lederberg AR. Joint attention in deaf and hearing 22 month-old children and their hearing mothers. *J Appl Dev Psychol*. 1998;19(3):377–387
34. Lederberg AR, Everhart VS. Communication between deaf children and their hearing mothers: the role of language, gesture, and vocalizations. *J Speech Lang Hear Res*. 1998;41(4):887–899
35. Rinaldi P, Baruffaldi F, Burdo S, Caselli MC. Linguistic and pragmatic skills in toddlers with cochlear implant. *Int J Lang Commun Disord*. 2013;48(6): 715–725
36. Yoshinaga-Itano C, Stredler-Brown A. Learning to communicate: babies with hearing impairments make their needs known. *Volta Review*. 1992;94(2): 107–129
37. Zaidman-Zait A, Dromi E. Analogous and distinctive patterns of prelinguistic communication in toddlers with and without hearing loss. *J Speech Lang Hear Res*. 2007;50(5):1166–1180
38. Moeller MP, Hoover B, Putman C, et al. Vocalizations of infants with hearing loss compared with infants with normal hearing: Part I-phonetic development. *Ear Hear*. 2007;28(5):605–627
39. Iyer SN, Oller DK. Prelinguistic vocal development in infants with typical hearing and infants with severe-to-profound hearing loss. *Volta Rev*. 2008; 108(2):115–138
40. McGillion M, Pine JM, Herbert JS, Matthews D. A randomised controlled trial to test the effect of promoting caregiver contingent talk on language development in infants from diverse socioeconomic status backgrounds. *J Child Psychol Psychiatry*. 2017;58(10): 1122–1131
41. Sloetjes H, Wittenburg P. Annotation by category: ELAN and ISO DCR. In: *Proceedings of the Sixth International Conference on Language Resources and Evaluation (LREC'08)*. May 26 to June 1, 2008; Marrakech, Morocco
42. D'Odorico L, Cassibba R. Cross-sectional study of coordination between infants' gaze and vocalizations towards their mothers. *Early Dev Parent*. 1995; 4(1):11–19
43. Murillo E, Belinchón M. Gestural-vocal coordination: longitudinal changes and predictive value on early lexical development. *Gesture*. 2012;12(1):16–39

44. Moeller MP, Hoover B, Putman C, et al. Vocalizations of infants with hearing loss compared with infants with normal hearing: Part II—transition to words. *Ear Hear*. 2007;28(5):628–642
45. Bruner JS. *Child's Talk: Learning to Use Language*. New York, NY: W. W. Norton & Company; 1983
46. Locke JL. *The Child's Path to Spoken Language*. Cambridge, MA: Harvard University Press; 1995
47. Matthews D. Learning How to Communicate in Infancy. In: Rowland CF, Theakston AL, Ambridge B, Twomey KE, eds. *Current Perspectives on Child Language Acquisition: How Children Use Their Environment to Learn*. Amsterdam, The Netherlands: John Benjamins Publishing Company; 2020: 11–38
48. Matthews D, Behne T, Lieven E, Tomasello M. Origins of the human pointing gesture: a training study. *Dev Sci*. 2012;15(6):817–829
49. Gogate LJ, Bahrick LE, Watson JD. A study of multimodal motherese: the role of temporal synchrony between verbal labels and gestures. *Child Dev*. 2000;71(4):878–894
50. Spencer PE, Meadow-Orlans KP, Koester LS, Ludwig JL. Relationships across Developmental Domains and over Time. In: Meadow-Orlans KP, Spencer PE, Koester LS, eds. *The World of Deaf Infants: A Longitudinal Study*. New York, NY: Oxford University Press; 2004: 205–217
51. Gale E, Schick B. Symbol-infused joint attention and language use in mothers with deaf and hearing toddlers. *Am Ann Deaf*. 2009;153(5):484–503
52. Depowski N, Abaya H, Oghalai J, Bortfeld H. Modality use in joint attention between hearing parents and deaf children. *Front Psychol*. 2015;6: 1556
53. Kurtzer-White E, Luterman D. Families and children with hearing loss: grief and coping. *Ment Retard Dev Disabil Res Rev*. 2003;9(4):232–235
54. Lederberg AR, Golbach T. Parenting stress and social support in hearing mothers of deaf and hearing children: a longitudinal study. *J Deaf Stud Deaf Educ*. 2002;7(4):330–345
55. Meadow-Orlans KP, Koester LS, Spencer PE, MacTurk RH. Theoretical Rationale for the Longitudinal Study. In: Meadow-Orlans KP, Spencer PE, Koester LS, eds. *The World of Deaf Infants: A Longitudinal Study*. New York, NY: Oxford University Press; 2004:11–23
56. Roberts MY. Parent-implemented communication treatment for infants and toddlers with hearing loss: a randomized pilot trial. *J Speech Lang Hear Res*. 2019;62(1):143–152
57. Szarkowski A, Young A, Matthews D, Meinzen-Derr J. Pragmatics development in deaf and hard of hearing children: a call to action. *Pediatrics*. 2020;146(suppl 3): e20200242L

Early Pragmatics in Deaf and Hard of Hearing Infants
Ciara Kelly, Gary Morgan, Colin Bannard and Danielle Matthews
Pediatrics 2020;146;S262
DOI: 10.1542/peds.2020-0242E

Updated Information & Services

including high resolution figures, can be found at:
http://pediatrics.aappublications.org/content/146/Supplement_3/S262

References

This article cites 45 articles, 0 of which you can access for free at:
http://pediatrics.aappublications.org/content/146/Supplement_3/S262#BIBL

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



PEDIATRICS[®]

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Early Pragmatics in Deaf and Hard of Hearing Infants
Ciara Kelly, Gary Morgan, Colin Bannard and Danielle Matthews
Pediatrics 2020;146;S262
DOI: 10.1542/peds.2020-0242E

The online version of this article, along with updated information and services, is located on the World Wide Web at:
http://pediatrics.aappublications.org/content/146/Supplement_3/S262

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 345 Park Avenue, Itasca, Illinois, 60143. Copyright © 2020 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN[®]

