

This is a repository copy of *Leishmania differentiation requires ubiquitin conjugation mediated by a UBC2-UEV1 E2 complex*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/165466/>

Version: Accepted Version

Article:

Burge, Rebecca J., Damianou, Andreas, Wilkinson, Anthony J. orcid.org/0000-0003-4577-9479 et al. (2 more authors) (2020) *Leishmania differentiation requires ubiquitin conjugation mediated by a UBC2-UEV1 E2 complex*. PLOS PATHOGENS. e1008784. ISSN 1553-7366

<https://doi.org/10.1371/journal.ppat.1008784>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

***Leishmania* differentiation requires ubiquitin conjugation mediated by a UBC2-UEV1 E2 complex**

Rebecca J. Burge¹, Andreas Damianou^{1,2}, Anthony J. Wilkinson³, Boris Rodenko⁴,
Jeremy C. Mottram¹

¹York Biomedical Research Institute and Department of Biology, University of York, UK.

²Wellcome Centre for Integrative Parasitology, Institute of Infection, Immunity and Inflammation, College of Medical Veterinary and Life Sciences, University of Glasgow, UK

³York Biomedical Research Institute and York Structural Biology Laboratory, Department of Chemistry, University of York, UK.

⁴UbiQ Bio BV, Amsterdam Science Park, The Netherlands.

Running title: E2 ubiquitin-conjugating complex of *Leishmania*

Abstract

Post-translational modifications such as ubiquitination are important for orchestrating the cellular transformations that occur as the *Leishmania* parasite differentiates between its main morphological forms, the promastigote and amastigote. 2 E1 ubiquitin-activating (E1), 13 E2 ubiquitin-conjugating (E2), 79 E3 ubiquitin ligase (E3) and 20 deubiquitinating cysteine peptidase (DUB) genes can be identified in the *Leishmania mexicana* genome but, currently, little is known about the role of E1, E2 and E3 enzymes in this parasite. Bar-seq analysis of 23 E1, E2 and HECT/RBR E3 null mutants generated in promastigotes using CRISPR-Cas9 revealed numerous loss-of-fitness phenotypes in promastigote to amastigote differentiation and mammalian infection. The E2s UBC1/CDC34, UBC2 and UEV1 and the HECT E3 ligase HECT2 are required for the successful transformation from promastigote to amastigote and UBA1b, UBC9, UBC14, HECT7 and HECT11 are required for normal proliferation during mouse infection, with $\Delta ubc2$ and $\Delta uev1$ exhibiting the most extreme loss-of-fitness phenotypes. Null mutants could not be generated for the E1 UBA1a or the E2s UBC3, UBC7, UBC12 and UBC13, suggesting these genes are essential in promastigotes. X-ray crystal structure analysis of UBC2 and UEV1, orthologues of human UBE2N and UBE2V1/UBE2V2 respectively, reveal a heterodimer with a highly conserved structure and interface. Furthermore, recombinant *L. mexicana* UBA1a can load ubiquitin onto UBC2, allowing UBC2-UEV1 to form K63-linked di-ubiquitin chains *in vitro*. Notably, UBC2 can cooperate *in vitro* with human E3s RNF8 and BIRC2 to form non-K63-linked polyubiquitin chains, showing that UBC2 can facilitate ubiquitination independent of UEV1, but association of UBC2 with UEV1 inhibits this ability. Our study demonstrates the dual essentiality of UBC2 and UEV1 in the differentiation and intracellular survival of *L. mexicana* and shows that the interaction between these two proteins is crucial for regulation of their ubiquitination activity and function.

Author summary

The post-translational modification of proteins is key for allowing *Leishmania* parasites to transition between the different life cycle stages that exist in its insect vector and mammalian host. In particular, components of the ubiquitin system are important for the transformation of *Leishmania* from its insect (promastigote) to mammalian (amastigote) stage and normal infection in mice. However, little is known about the role of the enzymes that generate ubiquitin modifications in *Leishmania*. Here we characterise 28 enzymes of the ubiquitination pathway and show that many are required for life cycle progression or mouse infection by this parasite. Two proteins, UBC2 and UEV1, were selected for further study based on their importance in the promastigote to amastigote transition. We demonstrate that UBC2 and UEV1 form a heterodimer capable of carrying out ubiquitination and that the structural basis for this activity is conserved between *Leishmania*, *Saccharomyces cerevisiae* and humans. We also show that the interaction of UBC2 with UEV1 alters the nature of the ubiquitination activity performed by UBC2. Overall, we demonstrate the important role that ubiquitination enzymes play in the life cycle and infection process of *Leishmania* and explore the biochemistry underlying UBC2 and UEV1 function.

Introduction

Leishmaniasis is a neglected tropical disease caused by parasites of the genus *Leishmania*. This disease, which has a tropical and sub-tropical distribution, is transmitted by the bite of a sandfly and causes around 70,000 deaths annually (1). During their complex, digenetic life cycle, *Leishmania* differentiate between two main morphological forms: the motile, extracellular promastigote form in the sand fly vector and the non-motile, intracellular amastigote form in the mammalian host. Of the numerous promastigote forms that exist in the sandfly, procyclic promastigotes are the most actively dividing and metacyclic promastigotes are the transmissible form (2). In order for *Leishmania* cells to transition between the disparate environments of the sandfly mouthparts and mammalian phagolysosomes, substantial changes in gene expression, particularly at the post-transcriptional level, are required (3, 4).

The post-translational modifications phosphorylation and ubiquitination, for example, are thought to contribute significantly to the differentiation process (5-8).

Ubiquitination regulates numerous cellular processes including proteasomal degradation, endocytic trafficking and DNA repair (9). Addition of the 8.5 kDa protein ubiquitin (Ub) to proteins is carried out by the sequential actions of E1 ubiquitin-activating (E1), E2 ubiquitin-conjugating (E2) and E3 ubiquitin ligase (E3) enzymes. Typically, an E1 activates ubiquitin in an ATP-dependent manner by adenylating its C-terminus, allowing a thioester bond to form between the E1 active site and ubiquitin. Subsequently, ubiquitin is transferred to the active site of an E2 via trans-thioesterification and then onto the substrate with the help of an E3 ligase. Most commonly, ubiquitination occurs on a lysine residue, although modification of cysteine, serine, threonine and the N-termini of proteins are also possible (10). E3 ligases can be grouped into two categories based on their mechanism of action. Cys-dependent E3s such as the HECT (Homologous to the E6-AP Carboxyl Terminus) and RBR (Ring-Between-Ring) E3s contain a cysteine residue that forms a thioester bond with ubiquitin prior to transfer to the substrate. Conversely, Cys-independent E3s such as the RING (really interesting new gene) and U-box E3s facilitate the direct transfer of ubiquitin between E2 and substrate by providing a scaffold that orients the ubiquitin-charged E2 relative to the substrate (11). Ubiquitin-like modifiers (Ubls) such as SUMO and Nedd8 have a similar, but distinct, E1-E2-E3 conjugation pathway to that of ubiquitin (12). The removal of ubiquitin modifications is carried out by deubiquitinating enzymes (DUBs) (9).

A huge diversity of ubiquitin modifications exists, due in part to the ability of ubiquitin itself to be ubiquitinated on any of the epsilon amino groups of its 7 lysine side chains or on the alpha amino group of its N-terminus. This allows for the formation of linear or branched polyubiquitin chains. Acetylation, phosphorylation and the modification of ubiquitin with other Ubls (including SUMO and Nedd8) are also possible (13). The huge array of potential modifications permits a range of signalling outcomes. For example, K48-linked chains, the most common ubiquitin chain type, target proteins for proteasomal degradation. In contrast, K63-linked chains

typically provide a non-degradative signal, for example in promoting the recruitment of proteins to sites of DNA damage (9).

To date, two ubiquitin-activating enzymes have been described in *Leishmania major* (14), and at least 20 cysteine peptidase DUBs exist in *Leishmania mexicana*, many of which are essential for the promastigote to amastigote transition (8). The importance of the ubiquitination system in these species is further exemplified by the finding that the *Leishmania* proteasome is essential for parasite survival, since many forms of ubiquitin modification target proteins for proteasomal degradation (15). The *L. major* Atg8 and Atg12 UbIs have been shown to play a role in parasite autophagy (16, 17) and, although they have yet to be properly described in *Leishmania*, SUMO- and Nedd8-conjugation systems exist in the closely-related kinetoplastid *Trypanosoma brucei* (18-23). The UbIs Ufm1 and Urm1 also exist in *Leishmania* (24-26). However, little is currently known about the role that ubiquitin E1, E2 and E3 enzymes play in *Leishmania* biology. In this study we characterise the E1, E2 and E3 enzymes of *L. mexicana* by analysing the fitness of an E1, E2 and HECT/RBR E3 null mutant library throughout the *Leishmania* life cycle. We show that, amongst others, the E2 ubiquitin-conjugating enzyme UBC2 and the ubiquitin E2 variant UEV1, are essential for the promastigote to amastigote transition and subsequently characterise these proteins in biochemical and structural detail.

Results

Ubiquitination gene families in *L. mexicana*

Initial analysis of the *L. mexicana* genome using Interpro and PFAM domain searches identified 4 ubiquitin-activating E1 (UBA), 15 ubiquitin E2-conjugating (UBC) and 81 E3 ligase genes (S1 Table). The putative E3s included 14 HECT, 1 RBR, 57 RING, 4 RING-CH-type and 5 U-box E3s. Upon more detailed analysis, however, LmxM.08.0220 and LmxM.02.0390 were found to be orthologues of *T. brucei* Uba2 and Ubc9, which have been identified as an E1 catalytic subunit and E2 enzyme for the Ubl SUMO respectively and were named UBA2

and UBC9 based on this orthology (21). Similarly, LmxM.01.0710 (UBA3) was found to share more similarity with HsUBA3 (68% query cover, 35.8% identity, E value: $8e^{-64}$), a Nedd8-activating enzyme catalytic subunit, than the ubiquitin E1 HsUBA1 (46% query cover, 31.7% identity, E value: $3e^{-25}$) and LmxM.24.1710 (UBC12) appears to be an orthologue of *T. brucei* Ubc12, an E2 Nedd8-conjugating enzyme (23). For two of the HECT domain-containing genes, *HECT13* and *HECT14*, only partial HECT domains of 69 and 60 amino acids were identified respectively, suggesting that they do not function as E3s. Following the removal of Ubl E1 and E2s and possible pseudo-HECTs from the list of identified ubiquitination genes, 2 ubiquitin E1s, 13 ubiquitin E2s and 79 E3 ligase genes were proposed to be present in the *L. mexicana* genome. Of these, UBC1, a ubiquitin E2, is related to *T. brucei* CDC34 (99% query cover, 55.2% identity, E value: $2e^{-105}$), required for cytokinesis and infection progression of bloodstream form parasites in mice (27). UBC4, also a ubiquitin E2, is related to *T. brucei* PEX4 (100% query cover, 58.6% identity, E value: $5e^{-72}$), implicated in the ubiquitination of TbPEX5, a cytosolic receptor involved in peroxisome biogenesis (28).

Of the putative ubiquitination genes identified, the predicted molecular weights of encoded proteins range from between 115-127 kDa for E1s, 16-49 kDa for E2s, 147-733 kDa for HECT E3 ligases and 9-288 kDa for RING, RING-CH-type and U-box E3 ligases. Only one RBR-type E3 ligase, with a predicted molecular weight of 58 kDa, was identified. An alignment of UBA1a and UBA1b with human ubiquitin-activating E1 enzymes (S1A Fig) revealed conservation of the catalytic cysteine residue (C596 in UBA1a and C651 in UBA1b, equivalent to C632 in HsUBA1). Similarly, an alignment of *L. mexicana* and human E2-conjugating enzymes showed conservation of the conserved catalytic cysteine in all *L. mexicana* E2s except UEV1 (S1B Fig), suggesting the latter is a non-catalytic ubiquitin E2 variant (29). Furthermore, the HPN (His-Pro-Asn) motif, which is conserved in human E2-conjugating enzymes and which contains an asparagine residue that can be important for catalysis (30-32), was found to be partially or completely missing in UBC3, UBC6, UBC11, UBC14 and UEV1. A HECT domain alignment for *L. mexicana* HECT E3 ligases and 4 human HECT E3

ligases showed that HECTs 1-12 contained the conserved catalytic cysteine residue. Consistent with the identification of a partial HECT domain in HECT13 and HECT14, a putative catalytic cysteine was absent in these sequences (S1C Fig).

A bar-seq screen reveals the importance of ubiquitination genes in the *L. mexicana* life cycle

In order to investigate the role of ubiquitination, SUMOylation and Neddylation genes in the life cycle of *L. mexicana*, the 4 E1, 15 E2 and HECT and RBR subgroups of E3 genes (12 and 1 gene respectively) from our bioinformatics analysis were targeted for deletion in procyclic promastigotes using CRISPR-Cas9 (33). The resistance cassettes used for integration into the targeted gene locus contained unique, 12 nucleotide barcodes to allow for the identification of individual null mutant lines in a pooled library. Details of the primers and plasmids used are provided in S2 Table. Null mutant lines were validated using PCRs to detect removal of the gene of interest and integration of the blasticidin resistance cassette (S2A Fig and S2B Fig) and, where a null mutant was not immediately generated, three rounds of transfection were performed to reduce the likelihood of not obtaining a null mutant line due to technical failure. A similar strategy was used for the generation of a Δ *hect12* mutant but will be described elsewhere. In this way, null mutants were generated for all genes except UBA1a, UBC3, UBC7, UBC12 and UBC13, suggesting that 1 out of 2 ubiquitin E1s and 4 out of 13 ubiquitin E2s may be essential in promastigotes. Although mutant clones were obtained for UBA1a, UBC3, UBC7 and UBC12, they still contained the gene of interest (S2C Fig). This lends further support to the proposition that UBA1a, UBC3, UBC7 and UBC12 are essential in promastigotes, since it suggests that gene duplication events have occurred to allow the parasite to retain the gene. Notably, a null mutant was successfully generated for the Nedd8 E1 UBA3 despite UBC12 (Nedd8 E2) appearing to be essential. This suggests that the essential role of UBC12 may be independent of UBA3. None of the SUMOylation, HECT E3

or RBR E3 genes were essential in promastigotes, as evidenced by the successful generation of null mutants for these genes.

Next, barcode analysis by sequencing (bar-seq) , which involves the parallel phenotyping of a null mutant pool using next-generation sequencing, was used to analyse the null mutants (34). Briefly, 58 promastigote-stage null mutants including the 3 E1, 10 E2 (excluding $\Delta ubc5$) and 13 HECT/RBR E3 null mutants alongside 16 DUB, 13 other peptidase and 4 protein kinase null mutants were pooled in equal proportions and in 6 replicate samples. Pooled promastigote (PRO) cultures were grown for 7 days and then induced to form axenic amastigotes (AXA) or used to purify metacyclic stage promastigotes (META). Purified metacyclic cells were used to infect macrophages (inMAC) in culture or mice using footpad injection (FP). At the time points indicated in Fig 1A, DNA was extracted to allow for amplification of the barcoded regions by PCR and quantitative analysis by next generation sequencing; the raw data are available in S4 Table in Damianou *et al.*, 2020 (8). The heat maps in Fig 1B-D show the proportional representation of each null mutant line at each experimental time point, with a gradient of white to red representing proportions ranging between zero and 0.04 (or above) respectively. Proportional representation was calculated by dividing the number of reads for an individual barcode by the total number of reads for all expected barcodes and averaging over the 6 replicates. Loss-of-fitness phenotypes were inferred from decreases in proportional representation from one time point to the next. For $\Delta uba2$, only 2.6×10^6 , instead of 4×10^6 cells, were added to the pools due to the poor prior growth of this cell line. Furthermore, the associated barcode was not detected in all of the PRO 0 h time points, perhaps due to the lower representation of $\Delta uba2$ in the pool, prompting the decision to exclude this cell line from the analysis.

Between the PRO 0 h and PRO 168 h samples, there was no complete loss of any null mutant line from the populations, although several significant loss-of-fitness phenotypes (decreases in proportional representation within the population) were observed between adjacent timepoints (<0.05 , paired t-test, Holm-Šidák method, S4 Table, Damianou *et al.*, 2020 (8)).

For example, reduced fitness was seen at two or more promastigote time points for *Δubc9*, *Δhect2* and *Δhect12*. No loss-of-fitness phenotypes were observed between the PRO 168 h and META samples or between the PRO 168 h and AXA 24 h samples.

Within both the AXA 24 h-AXA 120 h and META-inMAC 12 h intervals, loss-of-fitness was observed for 10 null mutants: *Δuba1b*, *Δubc2*, *Δubc6*, *Δubc8*, *Δubc9*, *Δuev1*, *Δhect2*, *Δhect4*, *Δhect7* and *Δrbr1*, demonstrating good correlation between these two experiments. Of these, strong defects (>3-fold decrease in fitness between both intervals) were observed for *Δubc1/cdc34*, *Δubc2*, *Δuev1* and *Δhect2*. In addition, *Δubc4/pex4*, *Δubc11*, *Δubc14*, *Δhect5*, *Δhect6*, *Δhect11* and *Δhect12* showed loss-of-fitness between the AXA 24 h and AXA 120 h samples and *Δhect10* showed a reduction in fitness between the META and inMAC 12 h samples. Further loss-of-fitness for *Δubc8* and *Δrbr1* was observed between inMAC 12 h and inMAC 72 h. All null mutant lines that showed loss-of-fitness defects in the axenic amastigote and macrophage samples also had compromised fitness in the mouse, although many of the phenotypes observed were more severe. Specifically, *Δuba1b*, *Δubc1/cdc34*, *Δubc2*, *Δubc9*, *Δubc14*, *Δuev1*, *Δhect2*, *Δhect7* and *Δhect11* had at least 20-fold reductions in fitness between the META and FP 3 samples. Additionally, *Δuba3* and *Δhect1* showed small decreases in fitness during this interval. Notably, *Δubc1/cdc34*, *Δubc2*, *Δuev1* and *Δhect2* showed the most dramatic and consistent loss-of-fitness phenotypes, including >5-fold, >25-fold and >200-fold decreases in fitness across the axenic amastigote, macrophage infection and mouse infection experiments respectively (calculated between the PRO 168 h/META sample and the experimental endpoints). A pBLAST search revealed that HECT2 is related to human UBE3C (28% query cover, 35.3% identity, $1e^{-64}$ E value), a HECT E3 ligase that ubiquitinates proteasome substrates to enhance proteasomal processivity (35). Overall, *Δubc2* and *Δuev1* had the most severe phenotypes with reductions in fitness of 50-fold in the axenic amastigote, 300-fold in the macrophage and a complete loss-of-fitness (to zero) in the mouse. The strong defects observed for *Δubc1/cdc34*, *Δubc2*, *Δuev1* and *Δhect2* in pooled

axenic amastigote differentiation were also observed individually using a cell viability assay (S2D Fig, S3 Table).

UBC2 and UEV1 are orthologues of human UBE2N and UBE2V1/UBE2V2

Since $\Delta ubc2$ and $\Delta uev1$ showed the most severe loss-of-fitness phenotypes in the bar-seq screen, we decided to investigate their function. The most probable orthologues of *L. mexicana* UBC2 (LmxM.04.0680) and UEV1 (LmxM.13.1580) were identified in the *Trypanosoma brucei*, *Saccharomyces cerevisiae* and human genomes using protein BLAST searches and the two groups of sequences aligned. Fig 2A shows that there is a high level of conservation between UBC2 and its predicted orthologues. Specifically, UBC2 shares 77%, 66% and 70% amino acid identity with the equivalent *T. brucei*, *S. cerevisiae* and human proteins respectively. Such a high level of shared identity suggests that UBC2 functions are likely to be well-conserved. In contrast, UEV1 is less well conserved with its orthologues, sharing 70%, 54% and 47% amino acid identity with the equivalent *T. brucei*, *S. cerevisiae* and human proteins respectively (Fig 2B). Consequently, there may be more functional divergence within this gene family. In humans, UBE2N and UBE2V1/V2 form a heterodimeric complex specific for K63-linked ubiquitination and have been linked to inflammatory signalling (36) and DNA damage response pathways (37).

UBC2 and *UEV1* encode proteins of 148 and 138 amino acid residues respectively, corresponding to predicted molecular weights of 17 kDa and 16 kDa. Based on the alignment in Fig 2A, the putative catalytic cysteine of UBC2 is C85, equivalent to C87 in *S. cerevisiae* UBC13 and C87 in human UBE2N (38). UBC2 also contains an HPN motif 10 amino acids N-terminal of C85 that may be involved in catalysis (30-32). All of the proteins in the UEV1 alignment (Fig 2B) lack catalytic cysteine residues and HPN motifs, consistent with UEV1 being part of the noncatalytic UEV family of E2s (29). For the UEV1 alignment, isoform 3 (canonical sequence in UniProtKB) of UBE2V1 was used as this is the most similar isoform to UEV1. However, UBE2V1 is predicted to have numerous isoforms produced by alternative

splicing. For example, a 30 residue N-terminal extension of isoform 2 (UEV-1A) relative to UBE2V2 has previously been shown to account for their differing functions (39). Since UEV1 lacks such an N-terminal extension, it could be predicted to be more similar in function to UBE2V2 than UBE2V1.

***L. mexicana* UBC2 and UEV1 form a heterodimer**

To permit biochemical characterisation of *L. mexicana* UBC2 and UEV1, His-Im9-tagged (40) UBC2 and UEV1 were expressed in *Escherichia coli* and purified following tag removal. In parallel, UBA1a, a putative ubiquitin-activating enzyme and the product of one of our proposed promastigote-essential genes, was similarly expressed and purified. Two putative ubiquitin-activating enzymes exist in *L. mexicana*, and, like *T. brucei* UBA1a and UBA1b, are more closely related to human UBA1 than UBA6 (14). Specifically, *L. mexicana* UBA1a and UBA1b share 36% and 33% amino acid identity with HsUBA1 respectively and 28% identity with each other. Based on its higher shared identity with HsUBA1 and potential essentiality, it was reasoned that UBA1a would be more likely to show a broad E2 specificity (comparable to that of HsUBA1) and therefore be capable of loading ubiquitin onto UBC2 (41, 42). All 3 proteins showed a good level of purity as assessed by InstantBlue staining (Fig 3A). For UBC2 and UEV1 no additional proteins were detected, whereas the UBA1a sample contained 2 extra proteins at around 74 and 100 kDa. These appeared below UBA1a, suggesting the presence of co-purified protein or UBA1a degradation products.

The orthologues of UBC2 and UEV1 in *T. brucei*, *S. cerevisiae* and humans have previously been shown to form a heterodimeric complex (43-46). To test whether this was also true in *L. mexicana*, a size-exclusion chromatography multi-angle laser light scattering (SEC-MALLS) approach was used. The chromatograms in Fig 3B show peaks in the refractive index representing the purified UBC2 and UEV1 samples at 25 and 26 min respectively. Measurement of both the refractive index and multi-angle laser light scattering allowed an estimation of the molecular weights of the recombinant proteins at 18.7 kDa and 18.2 kDa for

UBC2 and UEV1 respectively (dashed lines above peaks in refractive index). These values are close to the predicted molecular weights of 17 kDa for UBC2 and 16 kDa for UEV1. That both of these proteins eluted as a single peak is indicative of them existing in monomeric form while also reflecting the high quality of the protein preparations. When UBC2 and UEV1 were mixed in an equimolar ratio, an elution peak was seen at 23 min, indicating the presence of a higher molecular weight species (estimated at 34.2 kDa). Given that no peaks were seen representing UBC2 and UEV1 monomers, it can be assumed that all protein material in this sample existed in heterodimeric UBC2-UEV1 complexes. Therefore, *L. mexicana* UBC2 and UEV1 readily associate to form a heterodimeric UBC2-UEV1 complex.

Structure of the UBC2-UEV1 heterodimer

To investigate the conservation and interactions of UBC2 and UEV1 at the structural level, we sought crystals of their complex. Crystals, which appeared after two days from polyethylene glycol-containing solutions, were sent for data collection at the Diamond Light Source, with the best crystal yielding a dataset extending to 1.7 Å spacing. The structure was solved by molecular replacement using the coordinates of the human UBE2N-UBE2V2 complex as the search model (PDB ID: 1J7D) (47). There are two UBC2-UEV1 heterodimers (PDB ID: 6ZM3) in the crystallographic asymmetric unit. These were found to have a similar structure and subunit organisation following superposition of chains equivalent to UBC2, UEV1 or the UBC2-UEV1 heterodimer by secondary structure matching (SSM) procedures, with RMSDs of 0.32 Å, 0.74 Å and 1.13 Å for 146, 132 and 276 equivalent atoms respectively. Since residues Gly20 to Asn24 are poorly defined in the electron density maps for one of the UEV1 chains, the alternative UBC2-UEV1 heterodimer was the focus of our analysis.

Like its *S. cerevisiae* and human orthologues, UBC2 has a canonical E2 structure comprising a 4-stranded antiparallel β -sheet flanked by four α -helices (Fig 4A). UEV1 exhibits a similar topology, but its polypeptide chain is shorter and the prominent pair of α -helices at the C-terminus of UBC2 (α 3 and α 4) are missing from UEV1. Relative to human UBE2V1 and

UBE2V2, UEV1 has a shorter segment leading into the first α -helix (Fig 2B), a feature it shares with *S. cerevisiae* Mms2. Superposing UBC2 with human UBE2N, UEV1 with human UBE2V2 or the UBC2-UEV1 and UBE2N-UBE2V2 heterodimers (S3A Fig) gives RMSDs of 0.82, 1.15 and 1.13 Å for 147, 135 and 271 equivalent atoms respectively, demonstrating a high level of structural conservation between these orthologues. The conserved active site residues, His75, Pro76, Asn77 and Cys85, reside on an extended segment of the polypeptide that connects strand β 4 and helix α 2 of UBC2, with the thiol group of Cys85 projecting out of the catalytic cleft (Fig 4B). The high level of sequence conservation around these residues is reflected in the close proximity of their superposed catalytic clefts.

The interface between UBC2 and UEV1 (Fig 4C-D, S3B Fig) involves the β 2, β 3 and β 4 strands and the loops following β 1, β 3 and β 4 of UBC2 and the N-terminus, α 1 helix and loop following β 1 of UEV1. The buried surface area of this interface is 1,466 Å². The core of the interface contains a number of hydrophobic residues (Tyr32, Phe55, Leu70 and Val81 of UBC2 and Pro5, Phe8, Leu11, Leu14 and Phe39 of UEV1) that contribute strongly to the interaction. Hydrophobic residues are highly conserved at these positions (Fig 2), highlighting the importance of the associated hydrophobic interactions for complex formation. These hydrophobic interactions are complemented by a number of polar intermolecular interactions, mostly notably between the side chains of Glu53 of UBC2 and Asn7 of UEV1 and Arg68 of UBC2 and the main chain carbonyl oxygen of Ile37 of UEV1. Both of these interactions are conserved in the human complex although Met is present in place of Ile. Notably, a salt bridge is formed between Arg83 of UBC2 and Glu18 of UEV1. Although these residues are conserved in *S. cerevisiae*, human UBE2V1/2 has Gln in place of Glu. Asp38 and Glu39 residues in human UBE2V2 provide hydrogen bonds for the interaction interface but are replaced by Thr and Ala (at positions 34 and 35 respectively) in UEV1.

Ubiquitin transfer occurs between UBA1a and UBC2 *in vitro*

To assess whether UBA1a, UBC2 and UEV1 are an active E1, active E2 and inactive E2 variant respectively, recombinant proteins were tested in thioester intermediate assays. Fig 5A shows the results of incubating UBA1a, UBC2 and UEV1 in different combinations in the presence of ubiquitin and ATP. In these assays, human ubiquitin, which has 2 amino acid substitutions relative to *L. mexicana* ubiquitin (S4A Fig), was used. When UBA1a was present under non-reducing conditions, the appearance of a UBA1a~Ub thioester intermediate (at around 125 kDa) was observed over time. Under reducing conditions, this intermediate was lost, confirming the presence of thioester-linked UBA1a~Ub. When UBA1a and UBC2 were combined under non-reducing conditions, the appearance of an additional protein band at 26 kDa was observed, suggesting the transfer of ubiquitin between UBA1a and UBC2 to form thioester-linked UBC2~Ub. This protein complex was lost under reducing conditions, confirming its identity as a UBC2~Ub thioester intermediate. In contrast, when UBA1a and UEV1 were combined, no lower mobility species of UEV1 were observed, suggesting that UEV1 is unable to receive ubiquitin from UBA1a. When UBA1a, UBC2 and UEV1 were combined, no UBC2~Ub thioester intermediate was observed. It was reasoned that UEV1 could prevent UBC2 from binding ubiquitin or, alternatively, that the presence of UEV1 facilitates the release of ubiquitin from UBC2 or its transfer onto substrate(s) in solution. Conversely, the alternative *L. mexicana* ubiquitin E1, UBA1b, was unable to transfer ubiquitin to UBC2, despite being capable of forming thioester-linked UBA1b~Ub (S4B Fig). UBA1b was also unable to transfer ubiquitin onto UEV1 (S4B Fig), showing that ubiquitin cannot be loaded onto UEV1 by either of the two *L. mexicana* ubiquitin E1 enzymes. These results support the identities of UBA1a, UBC2 and UEV1 as an active E1, active E2 and inactive E2 variant respectively.

UBC2 and UEV1 conjugate ubiquitin *in vitro*

Previous *in vitro* studies have demonstrated the ability of *S. cerevisiae* UBC13 and MMS2 and their human counterparts to promote the formation of ubiquitin chains in the absence of E3 enzyme (30, 39, 44, 45, 48). To test whether the *L. mexicana* enzymes share this ability,

different combinations of UBC2 and UEV1 were incubated with UBA1a, ubiquitin and ATP and di-ubiquitin formation monitored by immunoblotting for mono- and poly-ubiquitinated conjugates under reducing conditions. This approach was chosen in order to distinguish between UBC2, UEV1 and di-ubiquitin, which all have similar molecular weights. Di-ubiquitin formation was observed after 30 min in the presence of UBC2 and UEV1 and increased by 90 min (Fig 5B). In contrast, no di-ubiquitin formation was observed in the absence of either UBC2 or UEV1, suggesting that both proteins are required for di-ubiquitin formation. An additional protein that may represent tri-ubiquitin was observed above di-ubiquitin at the 90 minute time point, suggesting that higher order chains may be being assembled. Additionally, since the reducing conditions would disrupt any UBC2-Ub thioester intermediates, the protein at around 26 kDa in samples containing UBC2 may represent auto-ubiquitinated UBC2. Comparably, *in vitro* auto-ubiquitination of human UBE2N on K92, equivalent to K90 in UBC2, has been observed (45). Auto-ubiquitinated UBA1a also appears to be present at the top of the blot.

Since *S. cerevisiae* UBC13 and MMS2 and their human orthologues have been shown to specifically form K63-linked ubiquitin chains (44, 45), the reactions described above were additionally probed with a K63 linkage-specific antibody. The right-hand panel in Fig 5B shows that K63-linked ubiquitin conjugates are formed by UBC2-UEV1. Furthermore, UBC2 and UEV1 were unable to conjugate K63R mutant ubiquitin, demonstrating the essential requirement for K63 of ubiquitin in the formation of free ubiquitin chains by UBC2-UEV1 (Fig 5C). Pre-incubation of UBC2 with the covalent UBE2N inhibitor NSC697923 reduced di-ubiquitin formation in a concentration-dependent manner, showing that K63-linked ubiquitin chain formation is dependent upon UBC2 catalytic activity (Fig 5D). That NSC697923 was likely to inhibit UBC2 was rationalised based on the fact that all 4 of the residues that were mutated to make UBE2N resistant to NSC697923 are also found in UBC2 (49). Complete inhibition of di-ubiquitin formation was not achieved, perhaps due to an insufficiently long incubation time for UBC2 with NSC697923. These experiments demonstrate that UBC2 and UEV1 can form free K63-linked ubiquitin chains *in vitro* (Fig 5E).

The crystal structure of *S. cerevisiae* Mms2-Ubc13 covalently linked to a donor ubiquitin molecule revealed the structural basis of K63 linkage specificity in ubiquitin chain formation. In this structure, Mms2 directs the K63 residue of a putative acceptor ubiquitin into the Ubc13 active site, where it can attack Gly76 of the donor ubiquitin bound to the Ubc13 active site cysteine to form an isopeptide bond (50). An overlay of this structure (PDB ID: 2GMI) with chains A and B of our UBC2-UEV1 structure allowed the positions of the donor and acceptor ubiquitins to be revealed in the context of the *L. mexicana* complex (S3C Fig). This produces a plausible model of the quaternary complex without significant steric clashes and suggests that the same strategy is used to confer Lys63-linkage specificity in *L. mexicana* and *S. cerevisiae*. In support of this, two Mms2 residues shown to be required for acceptor ubiquitin binding in the *S. cerevisiae* Mms2-Ubc13-Ub structure, Ser27 and Thr44, are conserved in *L. mexicana* UEV1 (Ser28 and Thr45).

UBC2 can cooperate with human E3s to allow polyubiquitination *in vitro*

As the cognate E3s for UBC2 and UEV1 have not yet been identified, human E3s known to catalyse ubiquitination in coordination with UBE2N were tested for their ability to similarly cooperate with UBC2 and/or UEV1. The reasons for doing this were threefold. Firstly, to investigate whether UBC2 could carry out the typical E2 role of facilitating ubiquitin transfer to substrates via E3 enzymes. Secondly, in the hope of making inferences about *L. mexicana* E3s that could be part of the UBC2 ubiquitination cascade and, lastly, to explore the conservation of E2-E3 interactions between *L. mexicana* and humans. For this purpose, two RING E3 ligases, BIRC2 and RNF8, were selected for testing on the basis that they are known to interact with human UBE2N (51-53). In humans, BIRC2 has wide-ranging roles including in regulating apoptosis and cell proliferation (51, 54) and RNF8 has well-characterised roles in DNA damage signalling (55-57). HUWE1, a HECT E3 ligase that is not known or predicted to interact with UBE2N, was also chosen for comparison (58). All recombinant E3s used were GST-tagged and, with the exception of HUWE1 which was N-terminally truncated, full-length.

Fig 6 shows the ubiquitination profiles observed for BIRC2, RNF8 or HUWE1 incubated with different combinations of UBC2 and UEV1 in reactions containing UBA1a, ubiquitin and ATP. When BIRC2 or RNF8 were incubated with UBC2 in the absence of UEV1, a prominent pattern of polyubiquitination was seen (upper panel). Based on the K63-linked ubiquitin and GST blots (middle and lower panels respectively), the polyubiquitination observed was not K63-linked and occurred both as a result of E3 auto-ubiquitination and of free chain formation and/or ubiquitination of other proteins (such as UBA1a, UBC2 or UEV1) in solution. When BIRC2 or RNF8 were incubated with UEV1 in the absence of UBC2, polyubiquitination did not occur. However, a single protein was present in the ubiquitin conjugate blot at around 130 kDa, likely representing ubiquitinated UBA1a (as observed in Fig 5B-D). Similarly, polyubiquitination was not observed when BIRC2 or RNF8 were incubated with both UBC2 and UEV1. In these samples, however, di-ubiquitin formation was notably increased, suggesting that UEV1 is able to regulate UBC2 such that it switches its activity between facilitating polyubiquitination by E3s and forming K63-linked ubiquitin chains in complex with UEV1 (Fig 6B). In contrast, polyubiquitination was observed when either UBC2, UEV1 or UBC2 and UEV1 were combined with HUWE1, suggesting that physical interaction between E2s and HUWE1 may be required for HUWE1 activity. Alternatively, the truncated nature of the recombinant HUWE1 protein, which lacks its UBA and WWE protein-protein interaction domains, may have encouraged non-specific interaction with and ubiquitin transfer from UBA1a. No ubiquitin conjugates were observed in the absence of both UBC2 and UEV1.

High functional conservation of human and *Leishmania* enzymes

In order to further interrogate the functional conservation between *Leishmania* and human enzymes, the ability of UBC2 to extend ubiquitin chains on the human U-box E3 ligase CHIP was investigated. *In vitro* monoubiquitination of CHIP by UBE2W can be followed by the extension of ubiquitin chains by UBE2N-UBE2V1 and provides an example of a pair of cooperating E2s with distinct chain initiation and elongation functions (59-61). To simplify the experimental setup and interpretation of results, it was decided to select a single E1 enzyme

to facilitate ubiquitin transfer to both human UBE2W and *L. mexicana* UBC2. In a thioester assay, *L. mexicana* UBA1a was shown to be equally competent at transferring ubiquitin to UBE2W as human UBA1 (Fig 7A) and was therefore selected for use in subsequent experiments. In addition to the reducible UBE2W-Ub thioester bands observed, an additional, non-reducible band was observed following incubation of E1 and E2 enzymes. This is likely to be N-terminally ubiquitinated UBE2W (62, 63).

When *L. mexicana* UBC2 and UEV1 were incubated with human UBE2W and CHIP in the presence of E1, ubiquitin and ATP, polyubiquitinated CHIP was observed (Fig 7B). When UBE2W was absent from this reaction, no CHIP ubiquitination or free chain formation was seen. Alternatively, when UBC2 and UEV1 were absent, only monoubiquitinated CHIP was observed, suggesting that UBE2W is priming CHIP with a single ubiquitin modification that can then be extended by UBC2 and UEV1 (Fig 7C). In this respect, UBC2 and UEV1 behave in a similar manner to human UBE2N-UBE2V1 (59). The absence of free chain formation in the presence of UBC2, UEV1 and CHIP, however, is in contrast to what was reported for human UBE2N-UBE2V1 and CHIP (59, 60). In reactions where UBE2W was present but CHIP was absent, a strong band at around 34 kDa was observed, likely corresponding to N-terminally ubiquitinated UBE2W as in Fig 7A (62, 63).

Discussion

Trypanosomatids are amongst the most ancient of eukaryotes and possess some highly divergent biochemistry, for example compartmentalisation of the glycolytic pathway or mRNA trans-splicing (64, 65). Despite this, our bioinformatic analysis of the *L. mexicana* genome revealed numerous ubiquitin conjugation system components: 2 ubiquitin E1, 13 ubiquitin E2 and 79 E3 ligase genes, including 12 HECT E3s, 1 RBR E3, 5 U-box RING E3s, 57 RING and 4 RING-CH-type E3s. These numbers are similar to those of another single-celled eukaryote, *S. cerevisiae*, which has 1 E1, 11 E2s and 60-100 E3s, of which 5 are HECT E3s, 2 are RBR E3s, 2 are U-box E3s, one is a RING CH-type E3 and the rest are RING E3s (66, 67). In

contrast, humans have 2 E1s, 40 E2s and over 600 E3s, of which 28 are HECT E3s, around 15 are RBR E3s, 9 are U-box E3s, 11 are RING CH-type E3s and the rest are RING E3s (67-71). We also identified a putative SUMO E1 catalytic subunit (UBA2), a Nedd8 E1 catalytic subunit (UBA3), a SUMO E2 (UBC9) and a Nedd8 E2 (UBC12), although E1, E2 and E3 genes for UbIs were not extensively searched for. Despite this, previous characterisation of the *T. brucei* orthologues of UBA2 and UBC9 by *in vitro* SUMOylation assays and UBA3 and UBC12 by affinity purification of Nedd8 lend further weight to their proposed functions (21, 23). Of the putative ubiquitin E2s identified, 5 were missing the conserved asparagine residue thought to be important for E2 catalysis (30-32). UEV1, which also lacks the catalytic cysteine residue, is part of the non-catalytic ubiquitin E2 variant protein family. For UBC3, UBC6, UBC11 and UBC14, in contrast, the missing asparagine residue may indicate non-canonical methods of E2 catalysis.

Previous work has underscored the importance of the ubiquitination system in *Leishmania* by demonstrating both the essentiality of the parasite proteasome and a key role for DUBs in the life cycle of this parasite (8, 15). Our generation of a select ubiquitination gene null mutant library revealed that 1 out of 2 ubiquitin E1s and 4 out of 13 ubiquitin E2s could not be deleted and therefore may be essential in promastigotes. A smaller proportion (4 out of 20) cysteine peptidase DUBs were shown to be essential in promastigotes, perhaps due to a greater degree of redundancy in DUB function (8). UBC12, a putative Nedd8 E2, also appears to be essential in promastigotes, suggesting it functions independently of the Nedd8 E1 UBA3, which is non-essential in promastigotes. This could be due to the presence of an additional Nedd8 E1 or a Neddylation-independent function of UBC12. In contrast, none of the HECT or RBR E3 ligases were essential in promastigotes, likely attributable to the considerable functional redundancy that characterises E3 ligases (72).

When interpreting our bar-seq data, we reasoned that since null mutants were more likely to exhibit decreases in proportional representation as the experiments progressed, our analysis should be limited to the identification of loss-of-fitness phenotypes. This is because decreases

in the relative abundance of a subset of null mutant lines in the population would lead to an increase in the proportional representation of the remaining null mutant lines, potentially mimicking gain-of-fitness phenotypes. Despite this limitation, the bar-seq approach allowed us to identify numerous fitness phenotypes associated with the promastigote and amastigote life cycle stages. In particular, loss-of-fitness was identified for more than one interval between promastigote time points for $\Delta ubc9$, hinting at a role for SUMOylation (UBC9), and $\Delta hect2$ and $\Delta hect12$, for HECT E3-mediated ubiquitination, in promastigote growth and/or survival. These genes were not absolutely required for survival, however, given our ability to detect $\Delta ubc9$, $\Delta hect2$ and $\Delta hect12$ in the metacyclic promastigote samples. Also of interest was the high degree of correlation between the data from the axenic amastigote, macrophage and mouse infection experiments, supporting previous findings that only small transcriptomic differences exist between axenic and intracellular amastigotes (3). The strong phenotypes observed for $\Delta ubc1/cdc34$, $\Delta ubc2$, $\Delta uev1$ and $\Delta hect2$ in the amastigote stages suggest an important role for these genes in the successful transformation from promastigote to amastigote. Since $\Delta hect2$ also showed loss-of-fitness in the promastigote stage, the effect of HECT2 deletion on cell survival/proliferation may be a more general one. For example, if the function of human UBE3C is shared with HECT2, then the build-up of harmful, incompletely-degraded proteasome substrates during the differentiation process could explain the requirement for HECT2 in amastigotes and the more subtle effect of HECT2 deletion on promastigotes (35). Notably, the observed requirement for UBC1/CDC34 during *L. mexicana* mouse infection mirrors the finding that TbCDC34 is required for infection of mice with bloodstream form *T. brucei* (27). Additionally, the human orthologues of both UBC2 and UEV1 (UBE2N and UBE2V1 respectively) have been implicated in the differentiation of various human cell types, perhaps pointing to a general role for these protein families in differentiation processes (29, 73-75). Since both ubiquitination and deubiquitination enzymes have been found to be essential in the promastigote to amastigote transition, interplay between the activities of E2/E3s (UBC1/CDC34, UBC2, UEV1 and HECT2) and DUBs (DUB4, DUB7 and DUB13) could be crucial for maintaining an optimal abundance and/or state of modification of protein

targets that are required for differentiation. $\Delta uba1b$, $\Delta ubc14$, $\Delta hect7$ and $\Delta hect11$ were lost at later stages and $\Delta ubc9$ showed cumulative loss throughout the experiments, suggesting that the deleted genes are required for normal amastigote proliferation, including during mouse infection.

Our finding that UBC2 and UEV1 are highly conserved at both the sequence and structural level suggests that their function may be shared between distantly related species. For example, that UEV1 is more similar in sequence to HsUBE2V2 and ScMMS2 than to HsUBE2V1 hints at a possible function in DNA damage repair (39). Our observation that UBC2 and UEV1 form a heterodimer was not unexpected and is consistent with the similar phenotypes observed for $\Delta ubc2$ and $\Delta uev1$ in promastigote to amastigote differentiation. We also found that UBC2 is a monomer *in vitro*. This property is similar to HsUBE2N but contrasts with ScUbc13, which is a homodimer *in vitro* (44). Given that dimerisation does not affect ubiquitin transfer by another human E2, UBE2W, this difference may not be physiologically relevant (76). Our X-ray crystal structure of the UBC2-UEV1 heterodimer revealed high conservation of the UBC2 active site and UBC2-UEV1 interface, highlighting the importance of stable UBC2-UEV1 interaction in the function of this complex across diverse eukaryotes.

Subsequent biochemical characterisation of UBC2 and UEV1 revealed that UBA1a and UBC2 are a functional ubiquitin E1-E2 pair. This demonstrates a level of specificity for E1-E2 interactions in *L. mexicana*, despite both UBA1a and UBA1b being more closely related to HsUBA1 than HsUBA6. That UBA1a and UBA1b appear to have unique functions is interesting since there is no obvious requirement for two ubiquitin E1s in other single-celled eukaryotes. *S. cerevisiae*, for example, has only one ubiquitin E1. A differential requirement for UBA1a and UBA1b is further supported by the likelihood that UBA1a, but not UBA1b, is an essential gene in promastigotes.

The UBC2-UEV1 heterodimer, like its *S. cerevisiae* and human orthologues, is able to form K63-linked ubiquitin chains *in vitro*, suggesting a role for non-degradative ubiquitin

modifications in *Leishmania* differentiation (30, 39, 44, 45, 48). Most of the chains observed in our *in vitro* assays were di-ubiquitin, supporting our suggestion that UEV1 is more similar to human UBE2V2, which forms di-ubiquitin *in vitro*, than UEV-1A (isoform 2 of UBE2V1), which forms polyubiquitin chains (39). Furthermore, our structural modelling of UBC2-UEV1 in complex with donor and acceptor ubiquitins, together with the conservation of key UEV1 residues that are required for acceptor ubiquitin interaction, are consistent with a role for UEV1 in dictating K63-linked chain specificity by correctly orienting the acceptor ubiquitin (50). In contrast, the donor ubiquitin is thought to exhibit flexible positioning around the covalent Cys85 linkage (50, 77). The physiological role of ubiquitin chains generated by *Leishmania* UBC2-UEV1 is currently unknown and an important area for further investigation. However, previous research has shown that free ubiquitin chains are present in both *S. cerevisiae* and human cells and that their levels (in *S. cerevisiae*) increase following heat shock, DNA damage or oxidative stress (78). Furthermore, unanchored K63-linked chains generated by the human E3 ligases TRAF6 and TRIM32 have been shown to interact with and activate protein kinases (79, 80) and unanchored K48-linked chains can inhibit the proteasome (81), demonstrating that free ubiquitin chains can perform regulatory functions. Curiously, *S. cerevisiae* HUL5, the E3 ligase partly responsible for free chain formation upon stress induction (78), is related to *L. mexicana* HECT2 (45% query cover, 32.6% identity, E value: $5e^{-47}$). This raises the possibility that HECT2 is involved in the response to environmental stresses that trigger the promastigote to amastigote transition and may explain the requirement for HECT2 in amastigotes. Exploring the potential interaction between UBC2-UEV1 or HECT2 and protein kinases involved in stress responses is an interesting avenue for further study. Additionally, the identification of UBC2 and UEV1 in an interactome of *L. mexicana* DUB2, which is able to cleave K63-linked diubiquitin *in vitro*, suggests a possible interplay between these proteins in the regulation of K63-linked ubiquitin chains (8).

For our *in vitro* experiments, human BIRC2 and RNF8 provided useful tools for examining the activity of UBC2 in the absence of available *Leishmania* E3s. Both BIRC2 and RNF8 were

able to form polyubiquitin chains (non-K63-linked) in a UBC2-dependent manner. Intriguingly, UEV1 effectively inhibited this reaction by switching UBC2 activity towards unanchored K63-linked di-ubiquitin formation. This ability of UEV1 could allow UBC2 to flip between direct (covalent attachment of ubiquitin to substrates) and indirect (unanchored ubiquitin chain binding to regulated proteins) mechanisms of regulating other proteins as well as between sets of protein targets, depending on the availability of UEV1. The proposed role for UEV1 in specifying K63-linked chain formation is further supported by the observation that polyubiquitin chains produced in the presence of UBC2 and BIRC2 or RNF8 are not K63-linked. In support of our finding that UBC2 acts alone with RNF8, human UBE2V1 and UBE2V2 have been shown to be dispensable for the function of UBE2N and RNF8 in DNA damage signalling (82). Conversely, the formation of polyubiquitin chains on human CHIP was seen in the presence of both UBC2 and UEV1. In this example, K63-linked ubiquitin chains could be formed in a RING-dependent manner similar to that described for rat RNF4, where the RING domain stabilises the positions of human UBE2N holds the donor ubiquitin in a 'folded-back' conformation poised for nucleophilic attack by K63 of the acceptor ubiquitin bound to UBE2V2 (77). The ability of UBC2-UEV1 to extend ubiquitin chains on CHIP illustrates the potential for UBC2-UEV1 to act in coordination with other E2s in *L. mexicana*, although an E2 with such a role has yet to be identified.

Previous studies have investigated the importance of DUBs and the parasite proteasome at various stages of the *Leishmania* life cycle (8, 15). Supplementing this, our study explores the requirement for selected E1, E2 and E3 enzymes across the life cycle of *Leishmania*. Our detailed investigation of UBC2 and UEV1, identified as essential in amastigotes, demonstrates high levels of conservation at both the structural and functional level. Consequently, our finding that the nature of UBC2 activity can be regulated by UEV1 has implications for orthologous proteins in other species.

Materials and Methods

Bioinformatic identification of ubiquitination genes

Ubiquitin E1-activating enzyme, E2-conjugating enzyme and E3 ligase genes were identified in the *L. mexicana* genome by performing Interpro and PFAM domain searches in TriTrypDB (<https://tritrypdb.org/tritrypdb/>). The following Interpro and PFAM identification codes were used: IPR018075, PF10585, IPR019572, IPR028077, IPR000608, IPR000569, PF00632, IPR002867, PF01485, IPR001841, IPR011016, IPR003613 and PF04564. UniProt (<https://www.uniprot.org/>) was also used to search for genes annotated with the terms “HECT” or “RBR”. Protein BLAST searches were used to determine the percentage query cover, percentage identity and E value for two gene sequences.

Leishmania culture

L. mexicana (MNYC/BZ/62/M379) promastigotes were grown in HOMEM (Gibco) supplemented with 10% v/v heat-inactivated Fetal Bovine Serum (FBS) (Gibco) and 1% v/v Penicillin/Streptomycin (Sigma-Aldrich) at 25°C. Typically, cells were split around twice a week. Selection drugs were added to the medium as appropriate: 10 µg mL⁻¹ blasticidin (InvivoGen), 40 µg mL⁻¹ puromycin (InvivoGen), 10-15 µg mL⁻¹ G418 (InvivoGen), 50 µg mL⁻¹ hygromycin (InvivoGen) and 50 µg mL⁻¹ nourseothricin (Jena Bioscience).

Null mutant library generation

Null mutants were generated using the CRISPR-Cas9-based approach as previously described (8, 33). Primer sequences to allow amplification of the single guide DNAs (sgDNAs) and repair cassettes for gene deletion were designed by entering the relevant gene identifiers into an automated web tool (<http://www.leishgedit.net/Home.html>). Primers for amplification of the repair cassettes contained primer binding sites for pTBlast_v1, pTPuro_v1 or pTNeo_v1 plasmids (sequences available from leishgedit.net) and 30 nt homology arms to allow recombination (83). Additionally, a T7 polymerase promoter sequence, 10 nt linker and 12 nt unique barcode were inserted into the 5' end of the upstream forward primer in the following

order: 5'-TAATACGACTCACTATAAACTGGAAGXXXXXXXXXXXXX-3', where X represents the barcoded region.

PCR reactions for cassette amplification contained 30 ng of plasmid (pTBlast_v1, pTPuro_v1 or pTNeo_v1), 0.2 mM dNTPs, 2 μ M each of forward and reverse primer, 1U Q5 DNA Polymerase (NEB), 1x Q5 reaction buffer (NEB) and distilled water to make the volume up to 40 μ L. The reverse primer used for generating all guides was aaaagcaccgactcgggtgccacttttcaagttgataacggactagccttattttaacttgctatttctagctctaaaac. The PCR was run with the following settings: 94°C for 5 min, 45 cycles of 94°C for 30 s, 65°C for 30 s and 72°C for 2 min 15 s and 72°C for 7 min. For the sgRNAs, PCR reactions were set up in a similar manner but with a total volume of 20 μ L. The PCR program used was 98°C for 30 s, 35 cycles of 98°C for 10 s, 60°C for 30 s and 72°C for 15 s and 72°C for 10 min.

Mid-log phase *L. mexicana* Cas9 T7 procyclic promastigotes were transfected either with whole PCR reactions (120 μ L total volume) or 2.5 μ g of DNA purified from PCR reactions using the QIAquick PCR Purification Kit (Qiagen). 8×10^6 log phase cells were prepared by spinning down (1,000 x g for 10 min), washing with 1 x PBS and resuspending in 1 x Cytomix (66.7 mM Na₂HPO₄, 23.3 mM NaH₂PO₄, 5 mM KCl, 50 mM HEPES and 150 μ M CaCl₂, pH 7.3) or P3 solution from the P3 Primary Cell 4D-Nucleofector X Kit (Lonza). Next, cells were pulsed twice using the Nucleofector 2B device (Lonza) and the X-001 program or using the Amaxa 4D-Nucleofector (Lonza) and the FI-115 programme (for the Cytomix and P3 solutions respectively) and then placed into 5 mL of HOMEM media with 20% FBS and 1% Penicillin/Streptomycin. As a negative control, the parental cell line was transfected with water in place of DNA.

Following recovery of the cells at 25°C, appropriate antibiotics were added (10 μ g mL⁻¹ blasticidin, 40 μ g mL⁻¹ puromycin, 15 μ g mL⁻¹ G418) to select for transfectants. Selection was performed either on a population level (10 mL containing the total transfected population)

or along with cloning into 96-well plates at 1:6, 1:66 and 1:726 dilutions. Dilutions were carried out in HOMEM supplemented with 20% v/v FBS and 1% v/v Penicillin/Streptomycin.

To extract genomic DNA for analysis, 500 μ L-5 mL of mid-log phase promastigotes were centrifuged (1,000 x g for 10 min), washed once in 1 x PBS and processed using the Qiagen DNeasy Blood and Tissue Kit and the manufacturer's protocol. For genotype analysis, PCR reactions were set up with 1 μ L of genomic DNA and either Q5 DNA Polymerase (NEB) or LongAmp Taq (NEB) with the manufacturer's protocols.

Bar-seq screen

The bar-seq screen was performed as described previously in Damianou *et al.*, 2020 (8); the raw data are available in S4 Table. Statistical analyses were performed between adjacent experimental time points using paired t-tests and the Holm-Šídák method in GraphPad Prism 8.

Cell viability assay

L. mexicana cultures were grown to stationary phase ($>1 \times 10^7$ mL⁻¹) and resuspended at 1×10^6 cells per mL in amastigote medium (Schneider's Drosophila medium [Gibco], 20% FBS [Gibco] and 15 μ g mL⁻¹ HEMIN [Sigma], adjusted to pH 5.5). 200 μ L cell samples were prepared in sextuplicate in 96-well plates and included the parental cell line (Cas9 T7) as a positive control. Also included were media-only, negative control samples. At 0 h, 48 h and 120 h, 20 μ L of 125 μ g mL⁻¹ resazurin (in 1 x PBS) was added to sample wells and the plate incubated at 37°C for 8 h. The fluorescence at 590 nm was then read using the POLARstar Omega Plate Reader (BMG Labtech). Relative viability was calculated for each sample by averaging fluorescence readings across the 6 replicates, subtracting the average for the negative control and then dividing by the value obtained for the Cas9 T7 positive control.

Sequence alignments

S. cerevisiae sequences were obtained from UniProtKB (84), human sequences from UniProtKB or NCBI and *L. mexicana* and *T. brucei* sequences from TriTrypDB (85). Sequence alignments were performed using T-Coffee (86) and structural annotation using ESPript 3.0 (87).

Protein expression and purification

UBA1a (LmxM.23.0550), *UBC2* (LmxM.04.0680) and *UEV1* (LmxM.13.1580) genes were codon-optimised for *E. coli* expression and synthesised by DC Biosciences. Genes were amplified by PCR and inserted into an expression vector (Protein Production Facility, University of York) in frame with an N-terminal His tag and Im9 solubility tag (40) using In-Fusion cloning (Takara). Primers used for gene amplification were 5'-TCCAGGGACCAGCAATGCTTTCTGAGGAAGAGCAAAAAC-3' and 5'-TGAGGAGAAGGCGCGTTAAAAGCGATAGCGGTAGCGGATG-3' for *UBA1a*, 5'-TCCAGGGACCAGCAATGTTGACCACTCGTATCATTAAGG-3' and 5'-TGAGGAGAAGGCGCGTCATGGTTTGGCGTACTTACGAG-3' for *UBC2* and 5'-TCCAGGGACCAGCAATGGTCGAGGTTCCGCGC-3' and 5'-TGAGGAGAAGGCGCGTCAGTAGGTACTACCCTCC-3' for *UEV1*. Insert integration and sequence were confirmed by DNA sequencing (Eurofins). Plasmids were then transformed into BL21-Gold (DE3) cells (Agilent Technologies). Cells were grown overnight in 5 mL of LB medium with 25 µg mL⁻¹ kanamycin and used to seed 500 mL cultures for growth at 37°C. When an optical density of around 0.5 at 600 nm was reached, cultures were equilibrated to 20°C and 1 mM of isopropyl 1-thio-β-D-galactopyranoside was added to induce recombinant protein production. Cultures were then grown for a further 24 h at 20°C.

The standard purification procedure was as follows: bacterial pellets from 500 mL cultures were resuspended in 25 mL buffer A (20 mM NaH₂PO₄·2H₂O, 20 mM Na₂HPO₄·12H₂O, 0.3 M NaCl, 30 mM imidazole and 5 mM β-mercaptoethanol, pH 7.4), DNase-treated and homogenised using a cell disruptor (Constant Systems Ltd). The lysate was centrifuged

(35,000 x g, 10 min, 4°C), filtered and loaded onto a HisTrap Fast Flow Crude column (GE Healthcare). After column washes with buffer A, a gradient of buffer B (20 mM NaH₂PO₄·2H₂O, 20 mM Na₂HPO₄·12H₂O, 0.3 M NaCl, 0.5 M imidazole and 5 mM β-mercaptoethanol, pH 7.4) was applied to elute bound protein. His-Im9 tag cleavage was then carried out using at least one tenth HRV 3C protease (Protein Production Facility, University of York) to sample protein (mg). This was followed by imidazole removal using either overnight dialysis at 4°C in buffer C (20 mM NaH₂PO₄·2H₂O, 20 mM Na₂HPO₄·12H₂O, 0.3 M NaCl and 5 mM β-mercaptoethanol, pH 7.4) or a desalt column. Removal of the His-Im9 tag leaves an additional 3 amino acids (Gly, Pro, Ala) at the N-terminus. Following cleavage, the sample was reappplied to the HisTrap column and application of buffer A and B used to separate the tag from the protein of interest. Sample fractions were concentrated and size-exclusion chromatography (SEC) carried out using HiLoad 16/600 Superdex 200 preparation grade (pg) or 75 pg columns (GE Healthcare) for UBA1a and UBC2 or UEV1 respectively. Buffer used for SEC contained 50 mM HEPES and 150 mM NaCl with either 2 mM DTT (UBC2 and UEV1) or 1 mM TCEP (UBA1a). Fractions containing purified protein were identified by SDS-PAGE analysis, pooled, concentrated and stored at -80°C.

For the UBC2 and UEV1 samples used for SEC-MALLS, purification was carried out as described above but with the following minor changes: buffers A-C did not contain β-mercaptoethanol, UBC2 was not purified by SEC and the buffer used for SEC of UEV1 was 25 mM Tris-HCl, 150 mM NaCl, pH 8. Storage was in the final purification buffers plus 1 mM DTT.

Size-exclusion chromatography multi-angle laser light scattering (SEC-MALLS)

Prior to loading, mixtures of UBC2 and UEV1 were prepared as required and incubated on ice for 30 min. 120 µL of each sample was then loaded onto a Superdex 200 HR10/300 gel filtration column (Sigma-Aldrich) and run in 25 mM Tris-HCl, 150 mM NaCl and 1 mM DTT, pH 8. Light scattering and refractive index measurements were taken using a Dawn Heleos II

and Optilab rEx detector (Wyatt) respectively. Concentrations of samples loaded were between 1.2 and 2 mg mL⁻¹.

Crystallisation and structure determination

For crystallisation experiments, UBC2 and UEV1 were mixed in a 1:1 molar ratio to a final concentration of 6.6 mg mL⁻¹ in buffer containing 50 mM HEPES, 150 mM NaCl and 2 mM DTT and incubated on ice for 30 min. Crystallisation conditions were screened (PACT premier HT-96 screen, Molecular Diagnostics) in a 96-well sitting drop format. Drops consisting of 150 nL of protein and 150 nL of reservoir solution were mixed and incubated above 100 µL of reservoir solution. Crystals appeared after two days at 25°C in drops prepared with a reservoir solution consisting of 0.1 M Bis-Tris propane, pH 7.5, 0.2 M sodium formate and 20% PEG. A single crystal was captured in a fine nylon loop, cryo-cooled in liquid nitrogen, and sent for data collection at the Diamond Light Source (Beamline I03). The diffraction data, extending to a nominal resolution of 1.7 Å, were processed using the 3dii pipeline in *xia2* (88). The crystals belonged to space group *P*2₁ with two UBC2-UEV1 heterodimers in the asymmetric unit.

The structure was solved by molecular replacement in the program MOLREP (89) implemented in the CCP4i2 interface (90). The search model used was the coordinate set for the human UBE2N-UBE2V2 complex (PDB ID: 1J7D). Model rebuilding and refinement were carried out using iterations of the programs Buccaneer (91) Refmac5 (92, 93) and Coot (94) in CCP4i2 (90). The electron density maps were of good quality allowing the confident tracing of the protein chains in the two heterodimers (AB and CD) with the exception of residues in the α1-β1 loop of UEV1 chains (B and D) where the maps were of poorer quality such that residues Gly22 and Ser23 could not be built in Chain D. It is assumed that this region of the structure has higher mobility. Data collection and refinement statistics are given in S4 Table. The coordinates and structure factors for the *L. mexicana* UBC2-UEV1 complex are available in the Protein Data Bank (PDB ID:6ZM3).

Structure analysis

Superposition of structures and RMSD determination were performed using “superpose structures” in CCP4mg (95). UBC2-UEV1 interface analysis was performed using PISA (version 1.52) (96).

Thioester formation assay

Reactions contained 300 nM UBA1a, 2.5 μ M E2, 20 μ M human ubiquitin (Boston Biochem) and 5 mM ATP as indicated in 40 μ L ubiquitination assay buffer (50 mM HEPES, pH 7.5, 100 mM NaCl, 10 mM MgCl₂ and 2 mM DTT). Reactions were incubated for 0-10 min at room temperature and then quenched with sample buffer with or without reducing agent. Samples containing reducing agent were heated at 90°C for 5 min. Samples were run on an SDS-PAGE gel and stained with InstantBlue Coomassie Protein Stain (Expedeon). For the UBE2W thioester assay, bovine ubiquitin (Ubiquigent) was used in place of human ubiquitin.

Di-ubiquitin formation assay

Reactions contained 100 nM UBA1a, 2.5 μ M of UBC2 and UEV1, 100 μ M of human wild-type (Boston Biochem) or K63R ubiquitin (2B Scientific) and 5 mM ATP as indicated in 40 μ L of ubiquitination assay buffer. Samples were incubated at 37°C for between 0 and 90 min as indicated. For the inhibitor assay, 0-50 μ M of NSC697923 (Abcam) was pre-incubated with UBC2 and UEV1 in reaction buffer for 15 min at room temperature prior to addition of UBA1a and ATP. The reaction was then incubated at 37°C for 90 min. The final DMSO concentration in these reactions was 0.5%. Following incubation, reducing sample buffer was added and the samples heated at 90°C for 5 min. Samples were separated by SDS-PAGE and Western blotting carried out using a mouse mono- and polyubiquitinated ubiquitin conjugate (Ubiquigent) or mouse Ub-K63 (ThermoFisher) antibody with HRP-conjugated anti-mouse secondary antibody (GE Healthcare or Promega). In addition to the experimental samples, 100 ng of K63 di-ubiquitin positive control (Ubiquigent) was loaded where indicated.

E3 cooperation assay

Reactions were prepared with 100 nM UBA1a, 2.5 μ M E2 (UBC2 or UEV1), 1 μ M human E3 (BIRC2, RNF8 or HUWE1), 100 μ M ubiquitin (Boston Biochem) and 5 mM ATP as indicated in ubiquitination assay buffer in 40 μ L total reaction volume. BIRC2, RNF8 and (N-terminally truncated) HUWE1 were all GST-tagged and sourced from Ubiquigent. Samples were then incubated at 30°C for 1 h prior to SDS-PAGE and Western blotting with a mouse mono- and polyubiquitinated ubiquitin conjugate (Ubiquigent), mouse Ub-K63 (ThermoFisher) or rabbit anti-GST (Abcam) antibody with HRP-conjugated anti-mouse (GE Healthcare) or HRP-conjugated anti-rabbit (GE Healthcare) secondary antibody as appropriate.

CHIP priming and extension assay

Reactions were prepared with 0.1 μ M UBA1a, 2.5 μ M 6His-tagged UBE2W (Ubiquigent), 1 μ M CHIP (Ubiquigent), 0.1 mM human ubiquitin (Boston Biochem) and 2 mM ATP (Sigma-Aldrich) in ubiquitination assay buffer in 50 μ L total reaction volume. Samples were incubated at 30°C for 1 h prior to SDS-PAGE and Western blotting with a mouse mono- and polyubiquitinated conjugate (Ubiquigent) or rabbit CHIP antibody (Calbiochem) with HRP-conjugated anti-mouse (GE Healthcare) or HRP-conjugated anti-rabbit (GE Healthcare) secondary antibody as appropriate.

Supplementary methods

Thioester formation assay

Reactions contained 6 pmol E1 (UBA1a or UBA1b), 2.5 μ M E2 (UBC2 or UEV1), 2 nmol human ubiquitin (Boston Biochem) and 2 mM ATP as indicated in 20 μ L ubiquitination assay buffer. Reactions were incubated for 30 min at 30°C. Sample buffer with or without reducing agent was then added and the samples containing reducing agent heated at 90°C for 5 min. Samples were run on an SDS-PAGE gel and stained with InstantBlue Coomassie Protein

Stain (Expedeon). Recombinant *L. mexicana* UBA1b was donated by Daniel Harris, University of Glasgow.

Acknowledgements

We would like to thank Ubiquigent for providing a 3-month internship for RB where the biochemical characterisation of UBC2-UEV1 was carried out. Specifically, we would like to thank Jane Hilton and Sheelagh Frame for technical support and Jason Brown for reviewing the manuscript. Additionally, we thank Daniel Harris (University of Glasgow) for donating the recombinant *L. mexicana* UBA1b, Jared Cartwright (Protein Production Facility, University of York) for providing plasmid backbone, H3C protease and valuable advice for protein expression and purification and Farid El Oualid (UbiQ) for valuable discussions. We also thank Diamond Light Source for beamline access to I03 (Proposal No MX-18598-30) that contributed to the results presented here, Marek Brzozowski for crystal-harvesting and valuable discussions, Johan Turkenburg for help with data processing and Eleanor Dodson for advice on structure solution and refinement.

References

1. Torres-Guerrero E, Quintanilla-Cedillo MR, Ruiz-Esmenjaud J, Arenas R. Leishmaniasis: a review. *F1000Res*. 2017;6:750.
2. Gossage SM, Rogers ME, Bates PA. Two separate growth phases during the development of *Leishmania* in sand flies: implications for understanding the life cycle. *Int J Parasitol*. 2003;33(10):1027-34.
3. Fiebig M, Kelly S, Gluenz E. Comparative Life Cycle Transcriptomics Revises *Leishmania mexicana* Genome Annotation and Links a Chromosome Duplication with Parasitism of Vertebrates. *PLoS Pathog*. 2015;11(10):e1005186.
4. De Pablos LM, Ferreira TR, Walrad PB. Developmental differentiation in *Leishmania* lifecycle progression: post-transcriptional control conducts the orchestra. *Curr Opin Microbiol*. 2016;34:82-9.
5. Morales MA, Watanabe R, Dacher M, Chafey P, Osorio y Fortéa J, Scott DA, et al. Phosphoproteome dynamics reveal heat-shock protein complexes specific to the *Leishmania donovani* infectious stage. *Proc Natl Acad Sci U S A*. 2010;107(18):8381-6.
6. Tsigankov P, Gherardini PF, Helmer-Citterich M, Späth GF, Myler PJ, Zilberstein D. Regulation dynamics of *Leishmania* differentiation: deconvoluting signals and identifying phosphorylation trends. *Mol Cell Proteomics*. 2014;13(7):1787-99.
7. Cayla M, Rachidi N, Leclercq O, Schmidt-Arras D, Rosenqvist H, Wiese M, et al. Transgenic analysis of the *Leishmania* MAP kinase MPK10 reveals an auto-inhibitory

- mechanism crucial for stage-regulated activity and parasite viability. *PLoS Pathog.* 2014;10(9):e1004347.
8. Damianou A, Burge RJ, Catta-Preta CMC, Geoghegan V, Nievas YR, Newling K, et al. Essential roles for deubiquitination in *Leishmania* life cycle progression. *PLoS Pathog.* 2020;16(6):e1008455.
 9. Komander D, Rape M. The ubiquitin code. *Annu Rev Biochem.* 2012;81:203-29.
 10. McClellan AJ, Laugesen SH, Ellgaard L. Cellular functions and molecular mechanisms of non-lysine ubiquitination. *Open Biol.* 2019;9(9):190147.
 11. Deshaies RJ, Joazeiro CA. RING domain E3 ubiquitin ligases. *Annu Rev Biochem.* 2009;78:399-434.
 12. Taherbhoy AM, Schulman BA, Kaiser SE. Ubiquitin-like modifiers. *Essays Biochem.* 2012;52:51-63.
 13. Swatek KN, Komander D. Ubiquitin modifications. *Cell Res.* 2016;26(4):399-422.
 14. Boer DR, Bijlmakers MJ. Differential Inhibition of Human and Trypanosome Ubiquitin E1S by TAK-243 Offers Possibilities for Parasite Selective Inhibitors. *Sci Rep.* 2019;9(1):16195.
 15. Khare S, Nagle AS, Biggart A, Lai YH, Liang F, Davis LC, et al. Proteasome inhibition for treatment of leishmaniasis, Chagas disease and sleeping sickness. *Nature.* 2016;537(7619):229-33.
 16. Williams RA, Woods KL, Juliano L, Mottram JC, Coombs GH. Characterization of unusual families of ATG8-like proteins and ATG12 in the protozoan parasite *Leishmania major*. *Autophagy.* 2009;5(2):159-72.
 17. Williams RA, Smith TK, Cull B, Mottram JC, Coombs GH. ATG5 is essential for ATG8-dependent autophagy and mitochondrial homeostasis in *Leishmania major*. *PLoS Pathog.* 2012;8(5):e1002695.
 18. Liao S, Wang T, Fan K, Tu X. The small ubiquitin-like modifier (SUMO) is essential in cell cycle regulation in *Trypanosoma brucei*. *Exp Cell Res.* 2010;316(5):704-15.
 19. Klein CA, Droll D, Clayton C. SUMOylation in *Trypanosoma brucei*. *PeerJ.* 2013;1:e180.
 20. López-Farfán D, Bart JM, Rojas-Barros DI, Navarro M. SUMOylation by the E3 ligase TbSIZ1/PIAS1 positively regulates VSG expression in *Trypanosoma brucei*. *PLoS Pathog.* 2014;10(12):e1004545.
 21. Ye K, Zhang X, Ni J, Liao S, Tu X. Identification of enzymes involved in SUMOylation in *Trypanosoma brucei*. *Sci Rep.* 2015;5:10097.
 22. Iribarren PA, Di Marzio LA, Berazategui MA, De Gaudenzi JG, Alvarez VE. SUMO polymeric chains are involved in nuclear foci formation and chromatin organization in *Trypanosoma brucei* procyclic forms. *PLoS One.* 2018;13(2):e0193528.
 23. Liao S, Hu H, Wang T, Tu X, Li Z. The Protein Neddylation Pathway in *Trypanosoma brucei*: FUNCTIONAL CHARACTERIZATION AND SUBSTRATE IDENTIFICATION. *J Biol Chem.* 2017;292(3):1081-91.
 24. Gannavaram S, Sharma P, Duncan RC, Salotra P, Nakhasi HL. Mitochondrial associated ubiquitin fold modifier-1 mediated protein conjugation in *Leishmania donovani*. *PLoS One.* 2011;6(1):e16156.
 25. Gannavaram S, Connelly PS, Daniels MP, Duncan R, Salotra P, Nakhasi HL. Deletion of mitochondrial associated ubiquitin fold modifier protein Ufm1 in *Leishmania donovani* results in loss of β -oxidation of fatty acids and blocks cell division in the amastigote stage. *Mol Microbiol.* 2012;86(1):187-98.
 26. Sharma V, Sharma P, Selvapandiyan A, Salotra P. *Leishmania donovani*-specific Ub-related modifier-1: an early endosome-associated ubiquitin-like conjugation in *Leishmania donovani*. *Mol Microbiol.* 2016;99(3):597-610.
 27. Rojas F, Koszela J, Búa J, Llorente B, Burchmore R, Auer M, et al. The ubiquitin-conjugating enzyme CDC34 is essential for cytokinesis in contrast to putative subunits of a SCF complex in *Trypanosoma brucei*. *PLoS Negl Trop Dis.* 2017;11(6):e0005626.
 28. Gualdrón-López M, Chevalier N, Van Der Smissen P, Courtoy PJ, Rigden DJ, Michels PAM. Ubiquitination of the glycosomal matrix protein receptor PEX5 in

Trypanosoma brucei by PEX4 displays novel features. *Biochim Biophys Acta*. 2013;1833(12):3076-92.

29. Sancho E, Vilá MR, Sánchez-Pulido L, Lozano JJ, Paciucci R, Nadal M, et al. Role of UEV-1, an inactive variant of the E2 ubiquitin-conjugating enzymes, in in vitro differentiation and cell cycle behavior of HT-29-M6 intestinal mucosecretory cells. *Mol Cell Biol*. 1998;18(1):576-89.

30. Wu PY, Hanlon M, Eddins M, Tsui C, Rogers RS, Jensen JP, et al. A conserved catalytic residue in the ubiquitin-conjugating enzyme family. *EMBO J*. 2003;22(19):5241-50.

31. Jones WM, Davis AG, Wilson RH, Elliott KL, Sumner I. A conserved asparagine in a ubiquitin-conjugating enzyme positions the substrate for nucleophilic attack. *J Comput Chem*. 2019.

32. Berndsen CE, Wiener R, Yu IW, Ringel AE, Wolberger C. A conserved asparagine has a structural role in ubiquitin-conjugating enzymes. *Nat Chem Biol*. 2013;9(3):154-6.

33. Beneke T, Madden R, Makin L, Valli J, Sunter J, Gluenz E. A CRISPR Cas9 high-throughput genome editing toolkit for kinetoplastids. *R Soc Open Sci*. 2017;4(5):170095.

34. Smith AM, Heisler LE, Mellor J, Kaper F, Thompson MJ, Chee M, et al. Quantitative phenotyping via deep barcode sequencing. *Genome Res*. 2009;19(10):1836-42.

35. Chu BW, Kovary KM, Guillaume J, Chen LC, Teruel MN, Wandless TJ. The E3 ubiquitin ligase UBE3C enhances proteasome processivity by ubiquitinating partially proteolyzed substrates. *J Biol Chem*. 2013;288(48):34575-87.

36. Wu X, Karin M. Emerging roles of Lys63-linked polyubiquitylation in immune responses. *Immunol Rev*. 2015;266(1):161-74.

37. Lee BL, Singh A, Mark Glover JN, Hendzel MJ, Spyropoulos L. Molecular Basis for K63-Linked Ubiquitination Processes in Double-Strand DNA Break Repair: A Focus on Kinetics and Dynamics. *J Mol Biol*. 2017;429(22):3409-29.

38. Motegi A, Sood R, Moinova H, Markowitz SD, Liu PP, Myung K. Human SHPRH suppresses genomic instability through proliferating cell nuclear antigen polyubiquitination. *J Cell Biol*. 2006;175(5):703-8.

39. Andersen PL, Zhou H, Pastushok L, Moraes T, McKenna S, Ziola B, et al. Distinct regulation of Ubc13 functions by the two ubiquitin-conjugating enzyme variants Mms2 and Uev1A. *J Cell Biol*. 2005;170(5):745-55.

40. Shevket SH, Gonzalez D, Cartwright JL, Kleanthous C, Ferguson SJ, Redfield C, et al. The CcmC-CcmE interaction during cytochrome. *J Biol Chem*. 2018;293(43):16778-90.

41. Jin J, Li X, Gygi SP, Harper JW. Dual E1 activation systems for ubiquitin differentially regulate E2 enzyme charging. *Nature*. 2007;447(7148):1135-8.

42. David Y, Ziv T, Admon A, Navon A. The E2 ubiquitin-conjugating enzymes direct polyubiquitination to preferred lysines. *J Biol Chem*. 2010;285(12):8595-604.

43. Crozier TWM, Tinti M, Larance M, Lamond AI, Ferguson MAJ. Prediction of Protein Complexes in. *Mol Cell Proteomics*. 2017;16(12):2254-67.

44. Hofmann RM, Pickart CM. Noncanonical MMS2-encoded ubiquitin-conjugating enzyme functions in assembly of novel polyubiquitin chains for DNA repair. *Cell*. 1999;96(5):645-53.

45. McKenna S, Spyropoulos L, Moraes T, Pastushok L, Ptak C, Xiao W, et al. Noncovalent interaction between ubiquitin and the human DNA repair protein Mms2 is required for Ubc13-mediated polyubiquitination. *J Biol Chem*. 2001;276(43):40120-6.

46. Ulrich HD, Jentsch S. Two RING finger proteins mediate cooperation between ubiquitin-conjugating enzymes in DNA repair. *EMBO J*. 2000;19(13):3388-97.

47. Moraes TF, Edwards RA, McKenna S, Pastushok L, Xiao W, Glover JN, et al. Crystal structure of the human ubiquitin conjugating enzyme complex, hMms2-hUbc13. *Nat Struct Biol*. 2001;8(8):669-73.

48. Pastushok L, Moraes TF, Ellison MJ, Xiao W. A single Mms2 "key" residue insertion into a Ubc13 pocket determines the interface specificity of a human Lys63 ubiquitin conjugation complex. *J Biol Chem*. 2005;280(18):17891-900.

49. Hodge CD, Edwards RA, Markin CJ, McDonald D, Pulvino M, Huen MS, et al. Covalent Inhibition of Ubc13 Affects Ubiquitin Signaling and Reveals Active Site Elements Important for Targeting. *ACS Chem Biol*. 2015;10(7):1718-28.
50. Eddins MJ, Carlile CM, Gomez KM, Pickart CM, Wolberger C. Mms2-Ubc13 covalently bound to ubiquitin reveals the structural basis of linkage-specific polyubiquitin chain formation. *Nat Struct Mol Biol*. 2006;13(10):915-20.
51. Bertrand MJ, Milutinovic S, Dickson KM, Ho WC, Boudreault A, Durkin J, et al. cIAP1 and cIAP2 facilitate cancer cell survival by functioning as E3 ligases that promote RIP1 ubiquitination. *Mol Cell*. 2008;30(6):689-700.
52. Plans V, Scheper J, Soler M, Loukili N, Okano Y, Thomson TM. The RING finger protein RNF8 recruits UBC13 for lysine 63-based self polyubiquitylation. *J Cell Biochem*. 2006;97(3):572-82.
53. Lok GT, Sy SM, Dong SS, Ching YP, Tsao SW, Thomson TM, et al. Differential regulation of RNF8-mediated Lys48- and Lys63-based poly-ubiquitylation. *Nucleic Acids Res*. 2012;40(1):196-205.
54. Samuel T, Okada K, Hyer M, Welsh K, Zapata JM, Reed JC. cIAP1 Localizes to the nuclear compartment and modulates the cell cycle. *Cancer Res*. 2005;65(1):210-8.
55. Kolas NK, Chapman JR, Nakada S, Ylanko J, Chahwan R, Sweeney FD, et al. Orchestration of the DNA-damage response by the RNF8 ubiquitin ligase. *Science*. 2007;318(5856):1637-40.
56. Mailand N, Bekker-Jensen S, Faustrup H, Melander F, Bartek J, Lukas C, et al. RNF8 ubiquitylates histones at DNA double-strand breaks and promotes assembly of repair proteins. *Cell*. 2007;131(5):887-900.
57. Huen MS, Grant R, Manke I, Minn K, Yu X, Yaffe MB, et al. RNF8 transduces the DNA-damage signal via histone ubiquitylation and checkpoint protein assembly. *Cell*. 2007;131(5):901-14.
58. Kar G, Keskin O, Nussinov R, Gursoy A. Human proteome-scale structural modeling of E2-E3 interactions exploiting interface motifs. *J Proteome Res*. 2012;11(2):1196-207.
59. Soss SE, Yue Y, Dhe-Paganon S, Chazin WJ. E2 conjugating enzyme selectivity and requirements for function of the E3 ubiquitin ligase CHIP. *J Biol Chem*. 2011;286(24):21277-86.
60. Zhang M, Windheim M, Roe SM, Pegg M, Cohen P, Prodromou C, et al. Chaperoned ubiquitylation--crystal structures of the CHIP U box E3 ubiquitin ligase and a CHIP-Ubc13-Uev1a complex. *Mol Cell*. 2005;20(4):525-38.
61. Ye Y, Rape M. Building ubiquitin chains: E2 enzymes at work. *Nat Rev Mol Cell Biol*. 2009;10(11):755-64.
62. Tatham MH, Plechanovová A, Jaffray EG, Salmen H, Hay RT. Ube2W conjugates ubiquitin to α -amino groups of protein N-termini. *Biochem J*. 2013;453(1):137-45.
63. Scaglione KM, Basrur V, Ashraf NS, Konen JR, Elenitoba-Johnson KS, Todi SV, et al. The ubiquitin-conjugating enzyme (E2) Ube2w ubiquitinates the N terminus of substrates. *J Biol Chem*. 2013;288(26):18784-8.
64. Hannaert V, Bringaud F, Opperdoes FR, Michels PA. Evolution of energy metabolism and its compartmentation in Kinetoplastida. *Kinetoplastid Biol Dis*. 2003;2(1):11.
65. Donelson JE, Gardner MJ, El-Sayed NM. More surprises from Kinetoplastida. *Proc Natl Acad Sci U S A*. 1999;96(6):2579-81.
66. Finley D, Ulrich HD, Sommer T, Kaiser P. The ubiquitin-proteasome system of *Saccharomyces cerevisiae*. *Genetics*. 2012;192(2):319-60.
67. Lin H, Li S, Shu HB. The Membrane-Associated MARCH E3 Ligase Family: Emerging Roles in Immune Regulation. *Front Immunol*. 2019;10:1751.
68. Lorenz S, Cantor AJ, Rape M, Kuriyan J. Macromolecular juggling by ubiquitylation enzymes. *BMC Biol*. 2013;11:65.
69. Scheffner M, Kumar S. Mammalian HECT ubiquitin-protein ligases: biological and pathophysiological aspects. *Biochim Biophys Acta*. 2014;1843(1):61-74.
70. Eisenhaber B, Chumak N, Eisenhaber F, Hauser MT. The ring between ring fingers (RBR) protein family. *Genome Biol*. 2007;8(3):209.

71. Li W, Bengtson MH, Ulbrich A, Matsuda A, Reddy VA, Orth A, et al. Genome-wide and functional annotation of human E3 ubiquitin ligases identifies MULAN, a mitochondrial E3 that regulates the organelle's dynamics and signaling. *PLoS One*. 2008;3(1):e1487.
72. Iconomou M, Saunders DN. Systematic approaches to identify E3 ligase substrates. *Biochem J*. 2016;473(22):4083-101.
73. Wu X, Yamamoto M, Akira S, Sun SC. Regulation of hematopoiesis by the K63-specific ubiquitin-conjugating enzyme Ubc13. *Proc Natl Acad Sci U S A*. 2009;106(49):20836-41.
74. Zhang W, Zhuang Y, Zhang Y, Yang X, Zhang H, Wang G, et al. Uev1A facilitates osteosarcoma differentiation by promoting Smurf1-mediated Smad1 ubiquitination and degradation. *Cell Death Dis*. 2017;8(8):e2974.
75. Lentucci C, Belkina AC, Cederquist CT, Chan M, Johnson HE, Prasad S, et al. Inhibition of Ubc13-mediated Ubiquitination by GPS2 Regulates Multiple Stages of B Cell Development. *J Biol Chem*. 2017;292(7):2754-72.
76. Vittal V, Wenzel DM, Brzovic PS, Klevit RE. Biochemical and structural characterization of the ubiquitin-conjugating enzyme UBE2W reveals the formation of a noncovalent homodimer. *Cell Biochem Biophys*. 2013;67(1):103-10.
77. Branigan E, Plechanovová A, Jaffray EG, Naismith JH, Hay RT. Structural basis for the RING-catalyzed synthesis of K63-linked ubiquitin chains. *Nat Struct Mol Biol*. 2015;22(8):597-602.
78. Braten O, Shabek N, Kravtsova-Ivantsiv Y, Ciechanover A. Generation of free ubiquitin chains is up-regulated in stress and facilitated by the HECT domain ubiquitin ligases UFD4 and HUL5. *Biochem J*. 2012;444(3):611-7.
79. Di Rienzo M, Antonioli M, Fusco C, Liu Y, Mari M, Orhon I, et al. Autophagy induction in atrophic muscle cells requires ULK1 activation by TRIM32 through unanchored K63-linked polyubiquitin chains. *Sci Adv*. 2019;5(5):eaau8857.
80. Xia ZP, Sun L, Chen X, Pineda G, Jiang X, Adhikari A, et al. Direct activation of protein kinases by unanchored polyubiquitin chains. *Nature*. 2009;461(7260):114-9.
81. Piotrowski J, Beal R, Hoffman L, Wilkinson KD, Cohen RE, Pickart CM. Inhibition of the 26 S proteasome by polyubiquitin chains synthesized to have defined lengths. *J Biol Chem*. 1997;272(38):23712-21.
82. Huen MS, Huang J, Yuan J, Yamamoto M, Akira S, Ashley C, et al. Noncanonical E2 variant-independent function of UBC13 in promoting checkpoint protein assembly. *Mol Cell Biol*. 2008;28(19):6104-12.
83. Beneke T, Gluenz E. LeishGEdit: A Method for Rapid Gene Knockout and Tagging Using CRISPR-Cas9. *Methods Mol Biol*. 2019;1971:189-210.
84. Consortium U. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res*. 2019;47(D1):D506-D15.
85. Aslett M, Aurrecochea C, Berriman M, Brestelli J, Brunk BP, Carrington M, et al. TriTrypDB: a functional genomic resource for the Trypanosomatidae. *Nucleic Acids Res*. 2010;38(Database issue):D457-62.
86. Notredame C, Higgins DG, Heringa J. T-Coffee: A novel method for fast and accurate multiple sequence alignment. *J Mol Biol*. 2000;302(1):205-17.
87. Robert X, Gouet P. Deciphering key features in protein structures with the new ENDscript server. *Nucleic Acids Res*. 2014;42(Web Server issue):W320-4.
88. Winter G, Lobley CM, Prince SM. Decision making in xia2. *Acta Crystallogr D Biol Crystallogr*. 2013;69(Pt 7):1260-73.
89. Vagin A, Teplyakov A. MOLREP: an Automated Program for Molecular Replacement. *Journal of Applied Crystallography*. 1997;30(6):1022-5.
90. Potterton L, Agirre J, Ballard C, Cowtan K, Dodson E, Evans PR, et al. CCP4i2: the new graphical user interface to the CCP4 program suite. *Acta Crystallogr D Struct Biol*. 2018;74(Pt 2):68-84.
91. Cowtan K. The Buccaneer software for automated model building. 1. Tracing protein chains. *Acta Crystallogr D Biol Crystallogr*. 2006;62(Pt 9):1002-11.

92. Murshudov GN, Vagin AA, Dodson EJ. Refinement of macromolecular structures by the maximum-likelihood method. *Acta Crystallogr D Biol Crystallogr.* 1997;53(Pt 3):240-55.
93. Murshudov GN, Skubák P, Lebedev AA, Pannu NS, Steiner RA, Nicholls RA, et al. REFMAC5 for the refinement of macromolecular crystal structures. *Acta Crystallogr D Biol Crystallogr.* 2011;67(Pt 4):355-67.
94. Emsley P, Lohkamp B, Scott WG, Cowtan K. Features and development of Coot. *Acta Crystallogr D Biol Crystallogr.* 2010;66(Pt 4):486-501.
95. McNicholas S, Potterton E, Wilson KS, Noble ME. Presenting your structures: the CCP4mg molecular-graphics software. *Acta Crystallogr D Biol Crystallogr.* 2011;67(Pt 4):386-94.
96. Krissinel E, Henrick K. Inference of macromolecular assemblies from crystalline state. *J Mol Biol.* 2007;372(3):774-97.




Figure 1. Life cycle phenotyping of ubiquitination gene null mutants. Fifty-eight null mutant lines were pooled ($n = 6$) as procyclic promastigotes and grown to stationary phase. Cells were then induced to differentiate into axenic amastigotes *in vitro* or metacyclic promastigotes purified and used to infect macrophages or mice. **A** Experimental workflow showing the time points at which DNA was extracted for barcode amplification and next generation sequencing.

The heat maps for promastigote to **B** axenic amastigote, **C** macrophage infection or **D** mouse infection experiments show the average proportional representation of each null mutant at each experimental time point, calculated by dividing the number of reads for null mutant-specific barcodes by the total number of reads for expected barcodes. Samples included represent promastigote time-point zero (PRO 0 h), early-log phase (PRO 24 h), mid-log phase (PRO 48 h), late-log phase (PRO 72 h), stationary phase (PRO 168 h), early axenic amastigote differentiation (AXA 24 h), post-axenic amastigote differentiation (AXA 120 h), purified metacyclic promastigotes (META), early macrophage infection (inMAC 12 h), late macrophage infection (inMAC 72 h), 3 week footpad mouse infection (FP 3) and 6 week footpad mouse infection (FP 6).




Figure 2. Alignments of *LmUBC2* and *LmUEV1* with selected orthologues. Sequence alignment and structural annotation were performed using T-Coffee and ESPript 3.0 respectively for **A** *LmUBC2* and **B** *LmUEV1*. Red boxes indicate amino acid identity, red characters show similarity within the highlighted group and blue frames highlight similarity across groups. Positions of the UBC2 HPN motif and catalytic cysteine are shown by a black

line and triangle respectively. Secondary structures derived from the crystallised *L. mexicana* UBC2-UEV1 heterodimer (Fig 4) are represented above the sequence alignment with helices represented by spirals and beta sheets by arrows. Asterisks denote important interface residues in the *L. mexicana* complex. *Lm*, *Leishmania mexicana*; *Tb*, *Trypanosoma brucei*; *Sc*, *Saccharomyces cerevisiae*; *Hs*, *Homo sapiens*.




Figure 3. *UBC2* and *UEV1* form a stable heterodimer *in vitro*. **A** SDS-PAGE gel showing recombinant UBA1a, UBC2 and UEV1 stained with InstantBlue stain. Proteins were expressed in *E. coli* and purified by nickel affinity and size-exclusion chromatography. 1 µg of each protein was loaded onto the gel. **B** Elution profiles of UBC2, UEV1 and a 1:1 molar mix of UBC2-UEV1 presented as changes in refractive index over time. Curved lines show changes in refractive index for UBC2-UEV1 (black), UBC2 (red) and UEV1 (blue). The expected mass (as estimated from light scattering data) in kDa is indicated by a dashed line above the peak corresponding to each protein sample.




Figure 4. Structure of the UBC2-UEV1 heterodimer. **A** Ribbon diagram showing the crystal structure of UBC2 (green) and UEV1 (blue) in complex. The location of the N- and C-termini are highlighted along with the numbering of alpha helices (α) and beta strands (β). **B** Zoom-in of the conserved catalytic residues in UBC2. The HPN motif and proposed catalytic cysteine are shown as cylinders coloured by atom (red, oxygen; blue, nitrogen; yellow, sulfur). **C-D** Zoom-in of interface between UBC2 and UEV1 with UBC2 or UEV1 as a surface fill model respectively. Amino acid residues are shown as cylinders coloured by atom (red, oxygen; blue, nitrogen). Hydrogen bonds are denoted by dashed lines.




Figure 5. *UBA1a* and *UBC2* cooperate in ubiquitin transfer *in vitro*. **A** Thioester assay demonstrating the ability of *UBA1a* and *UBC2* to form thioester bonds with ubiquitin. *UBA1a*, ubiquitin and ATP were incubated with *UBC2* and *UEV1* as indicated in ubiquitination assay buffer for up to 10 min at room temperature. Samples were treated with either reducing or non-reducing sample buffer and visualised by SDS-PAGE with InstantBlue stain. **B** Di-ubiquitin formation assay. *UBA1a*, ubiquitin and ATP were incubated with *UBC2* and *UEV1* as indicated in ubiquitination assay buffer for up to 90 min at 37°C. Reactions were then treated with reducing sample buffer and visualised by immunoblotting with either ubiquitin conjugate or K63-linked ubiquitin antibodies. **C** Di-ubiquitin formation assay performed as for **B** but with

wild-type ubiquitin substituted for K63R ubiquitin where shown. **D** UBC2 and UEV1 were pre-incubated with UBE2N inhibitor NSC697923 as indicated for 15 min prior to setting up a di-ubiquitin formation assay as in **B**. For **C** and **D**, samples were treated with reducing sample buffer prior to SDS-PAGE and immunoblotting with ubiquitin conjugate antibody. Data shown in **A-D** are representative. No 2/V1, no UBC2 or UEV1. **E** Schematic summarising the findings of **A-D**. UBA1a activates ubiquitin in an ATP-dependent manner, allowing ubiquitin to bind to the active site of UBA1a via a thioester linkage. Ubiquitin is then transferred to UBC2 (green) which, when present in complex with UEV1 (yellow), can generate free K63-linked ubiquitin chains.




Figure 6. Cooperation of UBC2 with human E3s in in vitro polyubiquitination. **A** UBA1a, ubiquitin and ATP were incubated with UBC2, UEV1 and human E3s (BIRC2, RNF8 and HUWE1) as indicated in ubiquitination assay buffer for 1 h at 30°C. Reactions were visualised by immunoblotting with either ubiquitin conjugate, K63-linked ubiquitin or GST antibodies as

Schematic summarising **A-B**. UBA1a activates ubiquitin in an ATP-dependent manner, allowing ubiquitin to bind to the active site of UBA1a via a thioester linkage. Ubiquitin is then transferred to HsUBE2W which monoubiquitinates HsCHIP. Alternatively, HsUBE2W can ubiquitinate its own N-terminus. Once primed with monoubiquitin, UBC2-UEV1 can extend ubiquitin chains on HsCHIP. Lm, *Leishmania mexicana*; Hs, *Homo sapiens*. Where species is not indicated, proteins are from *L. mexicana*.

S1 Table. *Summary of L. mexicana ubiquitination genes.*

Gene name	Type	<i>L. mexicana</i> gene ID	Size (kDa)	<i>T. brucei</i> orthologue
UBA1a	Ubiquitin E1	LmxM.23.0550	115.1	Tb927.8.2640
UBA1b	Ubiquitin E1	LmxM.34.3060	126.7	Tb11.v5.0675, Tb927.9.12650
UBA2	SUMO E1	LmxM.08.0220	109.7	Tb927.5.3430
UBA3	Nedd8 E1	LmxM.01.0710	56.1	Tb927.9.4620
UBC1/CDC34	Ubiquitin E2	LmxM.31.0960	30	Tb11.01.5790
UBC2	Ubiquitin E2	LmxM.04.0680	17.1	Tb927.9.8000
UBC3	Ubiquitin E2	LmxM.05.0930	48.8	Tb927.7.6960
UBC4/PEX4	Ubiquitin E2	LmxM.07.0850	20	Tb927.8.920
UBC5	Ubiquitin E2	LmxM.22.0610	28.2	Tb927.7.2540
UBC6	Ubiquitin E2	LmxM.33.1555	18.4	Tb927.4.2710
UBC7	Ubiquitin E2	LmxM.24.2130	17	Tb927.8.6090
UBC8	Ubiquitin E2	LmxM.31.0700	26.3	Tb927.11.13940
UBC9	SUMO E2	LmxM.02.0390	17.1	Tb927.2.2460
UBC10	Ubiquitin E2	LmxM.32.2770	26.1	Tb927.2.3720
UBC11	Ubiquitin E2	LmxM.33.0900	31.7	Tb927.4.3190, Tb927.4.3460
UBC12	Nedd8 E2	LmxM.24.1710	21.6	Tb927.8.6510
UBC13	Ubiquitin E2	LmxM.34.1300	16.7	Tb927.5.1000
UBC14	Ubiquitin E2	LmxM.24.2140	36.4	Not identified
UEV1	Ubiquitin E2 variant	LmxM.13.1580	16	Tb927.11.3310
HECT1	HECT E3 ligase	LmxM.36.6340	455.4	Tb927.10.8390
HECT2	HECT E3 ligase	LmxM.29.0910	153	Tb927.6.2370
HECT3	HECT E3 ligase	LmxM.07.0280	666.2	Tb927.8.1590
HECT4	HECT E3 ligase	LmxM.13.1470	592.1	Tb927.11.3390
HECT5	HECT E3 ligase	LmxM.36.4370	276.8	Tb11.v5.0817, Tb927.10.9750
HECT6	HECT E3 ligase	LmxM.26.2370	421.9	Tb927.9.1770
HECT7	HECT E3 ligase	LmxM.31.1090	178.7	Tb927.11.14330

HECT8	HECT E3 ligase	LmxM.31.3930	453.9	Tb927.11.16260
HECT9	HECT E3 ligase	LmxM.33.3960	149.3	Tb927.4.770
HECT10	HECT E3 ligase	LmxM.34.2450	250.9	Tb927.9.13360
HECT11	HECT E3 ligase	LmxM.34.5390	732.6	Tb927.4.310
HECT12	HECT E3 ligase	LmxM.34.4000	147.1	Not identified
HECT13	HECT E3 ligase	LmxM.33.3400	36.3	Not identified
HECT14	HECT E3 ligase	LmxM.26.2650	63.5	Tb927.9.1350
RBR1	RBR E3 ligase	LmxM.29.2170	58.1	Tb927.6.3780
Not given	RING-type zinc finger-containing	LmxM.15.1410	52.8	Tb927.9.5260
Not given	RING-type zinc finger-containing	LmxM.21.0023	12.2	Tb927.10.1810
Not given	RING-type zinc finger-containing	LmxM.24.1610	13.1	Not identified
Not given	RING-type zinc finger-containing	LmxM.34.0660	133.9	Tb927.10.3800
Not given	RING-type zinc finger-containing	LmxM.34.4500	9.4	Tb927.9.10170
Not given	RING-type zinc finger-containing	LmxM.01.0430	81.2	Tb927.9.4050
Not given	RING-type zinc finger-containing	LmxM.02.0140	94.1	Tb927.2.2360
Not given	RING-type zinc finger-containing	LmxM.03.0010	104.9	Not identified
Not given	RING-type zinc finger-containing	LmxM.04.0540	63.7	Not identified
Not given	RING-type zinc finger-containing	LmxM.04.0640	35.7	Not identified
Not given	RING-type zinc finger-containing	LmxM.07.0370	96.5	Not identified
Not given	RING-type zinc finger-containing	LmxM.08.1091	41.1	Tb927.5.2920
Not given	RING-type zinc finger-containing	LmxM.08.1215	56.3	Tb927.5.3070
Not given	RING-type zinc finger-containing	LmxM.09.0140	57.3	Tb927.11.12190
Not given	RING-type zinc finger-containing	LmxM.09.0350	88.4	Not identified
Not given	RING-type zinc finger-containing	LmxM.09.0430	51.9	Tb927.11.12440
Not given	RING-type zinc finger-containing	LmxM.09.1210	44.9	Not identified
Not given	RING-type zinc finger-containing	LmxM.10.0440	17.6	Not identified
Not given	RING-type zinc finger-containing	LmxM.10.1140	39.3	Tb927.8.4570
Not given	RING-type zinc finger-containing	LmxM.12.0100	32	Tb927.6.1190
Not given	RING-type zinc finger-containing	LmxM.12.1240	96	Tb927.1.4520

Not given	RING-type zinc finger-containing	LmxM.25.0740	122.7	Tb927.11.750
Not given	RING-type zinc finger-containing	LmxM.25.2230	38	Tb11.v5.0818, Tb927.3.2340
Not given	RING-type zinc finger-containing	LmxM.25.2290	33.2	Tb927.3.2410
Not given	RING-type zinc finger-containing	LmxM.26.1170	55.2	Tb927.7.770
Not given	RING-type zinc finger-containing	LmxM.26.1560	204.2	Tb927.7.1090
Not given	RING-type zinc finger-containing	LmxM.27.0570	59.9	Tb11.v5.0388, Tb927.11.1190
Not given	RING-type zinc finger-containing	LmxM.27.1200	44.1	Tb11.1410, Tb927.11.1810
Not given	RING-type zinc finger-containing	LmxM.28.0760	83	Tb927.11.8010
Not given	RING-type zinc finger-containing	LmxM.28.1300	45.9	Tb927.5.4280
Not given	RING-type zinc finger-containing	LmxM.28.1830	99.6	Tb927.11.8010
Not given	RING-type zinc finger-containing	LmxM.29.1230	78.8	Tb927.6.2710
Not given	RING-type zinc finger-containing	LmxM.30.2890	135.3	Tb927.8.7200
Not given	RING-type zinc finger-containing	LmxM.31.0560	19.8	Not identified
Not given	RING-type zinc finger-containing	LmxM.13.0230	44.9	Tb927.11.4560
Not given	RING-type zinc finger-containing	LmxM.16.0650	11.6	Tb927.5.3745
Not given	RING-type zinc finger-containing	LmxM.17.0030	21	Tb927.7.6190
Not given	RING-type zinc finger-containing	LmxM.17.0290	76.2	Not identified
Not given	RING-type zinc finger-containing	LmxM.17.0400	113.6	Not identified
Not given	RING-type zinc finger-containing	LmxM.18.1150	39.4	Tb927.10.12940
Not given	RING-type zinc finger-containing	LmxM.18.1570	16.9	Tb927.10.12460
Not given	RING-type zinc finger-containing	LmxM.19.1150	288.2	Tb11.v5.0394, Tb927.10.15750
Not given	RING-type zinc finger-containing	LmxM.36.1930	32.6	Not identified
Not given	RING-type zinc finger-containing	LmxM.36.2930	59.4	Not identified
Not given	RING-type zinc finger-containing	LmxM.36.3830	34	Tb927.11.9660
Not given	RING-type zinc finger-containing	LmxM.21.1295	28.2	Not identified
Not given	RING-type zinc finger-containing	LmxM.22.0060	84.8	Tb927.3.5640, Tb927.7.2050
Not given	RING-type zinc finger-containing	LmxM.22.0190	50.8	Not identified
Not given	RING-type zinc finger-containing	LmxM.23.0280	117.9	Not identified

Not given	RING-type zinc finger-containing	LmxM.23.1730	42.9	Tb927.8.3460
Not given	RING-type zinc finger-containing	LmxM.24.0080	55.3	Tb927.11.4860
Not given	RING-type zinc finger-containing	LmxM.24.0300	193.2	Tb927.11.5020
Not given	RING-type zinc finger-containing	LmxM.24.1380	44.1	Tb927.8.6850
Not given	RING-type zinc finger-containing	LmxM.24.1390	37.3	Tb927.8.6850
Not given	RING-type zinc finger-containing	LmxM.25.0340	72	Tb927.3.5390
Not given	RING-type zinc finger-containing	LmxM.34.2520	42.8	Tb927.9.13300
Not given	RING-type zinc finger-containing	LmxM.29.2360	60.8	Tb927.6.3640
Not given	RING-CH-type zinc finger-containing	LmxM.08_29.1630	135	Tb927.3.4220
Not given	RING-CH-type zinc finger-containing	LmxM.27.0120	70.1	Tb927.3.710
Not given	RING-CH-type zinc finger-containing	LmxM.30.0030	115.9	Not identified
Not given	RING-CH-type zinc finger-containing	LmxM.30.1920	140.1	Tb927.4.4880, Tb927.8.7580
Not given	U-box E3 ligase	LmxM.06.0510	71.2	Not identified
Not given	U-box E3 ligase	LmxM.13.0660	76	Tb927.11.4210
Not given	U-box E3 ligase	LmxM.26.1600	72.8	Not identified
Not given	U-box E3 ligase	LmxM.27.2480	53	Tb927.2.5240
Not given	U-box E3 ligase	LmxM.08_29.0780	66.8	Tb927.3.3560

S2 Table. *Summary of primers and plasmids.*

Oligo number	Target/associated gene	Sequence	Direction	Notes
OL5674	BSD (blasticidin resistance gene)	GACGATACAAGTCAGGTTG	Reverse	Blasticidin resistance cassette for diagnostic PCRs 1
OL6137	gDNA	aaaagcaccgactcgggtgccacttttcaagttgataacggactagccttattttaactgctatttctagctctaaac	Reverse	Standard primer for T7RPol expression of sgRNA (contains Cas9 scaffold for generation of sgRNA PCR)
OL6367	LmxM.36.6340	gaaattaatacgactcactataggACCATTGCTCTCAAAACAAGgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6370	LmxM.29.0910	gaaattaatacgactcactataggAGTCCTTCGCACAAACACTGgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6373	LmxM.07.0280	gaaattaatacgactcactataggAAGGGTTCCAATAGTTGGGGgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6376	LmxM.13.1470	gaaattaatacgactcactataggACACACACACGCCAATAAAAgtttagagctagaaatagc	Forward	5' sgRNA primer
OL6379	LmxM.36.4370	gaaattaatacgactcactataggTCCATTACACGGCAGCACTGgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6382	LmxM.26.2370	gaaattaatacgactcactataggAGGACGCAACCAGAGGCAAAgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6385	LmxM.31.1090	gaaattaatacgactcactataggACTAAGGTGAGGACGTCACAgtttagagctagaaatagc	Forward	5' sgRNA primer

OL6388	LmxM.31.39 30	gaaattaatacgactcactataggCTTTGGTGCGGTTTCCATCTgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6391	LmxM.33.39 60	gaaattaatacgactcactataggTTCACAAAGTACTCTGAGGCgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6394	LmxM.34.24 50	gaaattaatacgactcactataggTACCAAGAGCGAGTACTGTGgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6397	LmxM.34.53 90	gaaattaatacgactcactataggTAACAATAGCGTTTCTACGGgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6400	LmxM.29.21 70	gaaattaatacgactcactataggGCCATTACGACATGGGAAAgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6838	LmxM.02.03 90	gaaattaatacgactcactataggACCCTCTCTGTCGCTCCCAAgtttagagctagaaatagc	Forward	5' sgRNA primer
OL6841	LmxM.04.06 80	gaaattaatacgactcactataggAGAGTGGGGAAGGGTGGATAgtttagagctagaaatagc	Forward	5' sgRNA primer
OL6844	LmxM.05.09 30	gaaattaatacgactcactataggGAGAACAGGGGTGGGGGTGGgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6847	LmxM.07.08 50	gaaattaatacgactcactataggTGCGTGTGTGTATGTGTGTAgtttagagctagaaatagc	Forward	5' sgRNA primer
OL6850	LmxM.13.15 80	gaaattaatacgactcactataggGTGCAAGTGAATATGTTGTCgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6853	LmxM.22.06 10	gaaattaatacgactcactataggCTCTTGCTTCCTTCACAAGCgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6856	LmxM.24.17 10	gaaattaatacgactcactataggTGTACAGTGTGTCTTCAAATgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6859	LmxM.24.21 30	gaaattaatacgactcactataggCAACGCGCTTCTCTGGTCTTgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6862	LmxM.31.07 00	gaaattaatacgactcactataggGTGGTGGCGGCTGGGATCTGgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6865	LmxM.31.09 60	gaaattaatacgactcactataggTGATGATGATGATGCCACTTgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6868	LmxM.32.27 70	gaaattaatacgactcactataggCTTGTAGCGAGGAGTAGGAAgtttagagctagaaatagc	Forward	5' sgRNA primer
OL6871	LmxM.33.09 00	gaaattaatacgactcactataggACAGGTGGGAAGGCGAGTTAgtttagagctagaaatagc	Forward	5' sgRNA primer

OL6874	LmxM.33.15 55	gaaattaatacgactcactataggCGACGGCTCGTGTCGTTTTgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6877	LmxM.34.13 00	gaaattaatacgactcactataggTGTGTTTTCTTTCCTTTCgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6880	LmxM.24.21 40	gaaattaatacgactcactataggGTCGATGAGGATGATACACGgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6883	LmxM.23.05 50	gaaattaatacgactcactataggCGCGCCTGAGTGTGAGTGCGgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6886	LmxM.34.30 60	gaaattaatacgactcactataggTTGTGCTTTGCGCTGTGTTTgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6889	LmxM.08.02 20	gaaattaatacgactcactataggCTACGCTACTCGTTACTTTGgttttagagctagaaatagc	Forward	5' sgRNA primer
OL6892	LmxM.01.07 10	gaaattaatacgactcactataggACGACGGCGTAGATGCGCGAggttttagagctagaaatagc	Forward	5' sgRNA primer
OL7153	LmxM.36.63 40	ACCCCCCTCCCCTGAGCATCGCAGGTGATTAATACGACTCACTATAAAAC TGGAAGAAATTAACCTTgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL7154	LmxM.36.63 40	GAAGGGGAGGCGGCGCAAACGAAACGACCCccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL7155	LmxM.36.63 40	gaaattaatacgactcactataggTAGCAGGGCGTATCGAACGTgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL7156	LmxM.29.09 10	GGTCGACGTCGGCTTGCCTCGCAGCTCCCTTAATACGACTCACTATAAAAC TGGAAGTTGCATGTGGCCgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL7157	LmxM.29.09 10	CTCGTTCTGACTGCTCCACGAAGCCCACCGccaatttgagagacctgtgc	Reverse	Downstream reverse primer for

				null mutant generation
OL7158	LmxM.29.09 10	gaaattaatacgactcactataggCGTGGCGCTACAACGCGTCTgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL7159	LmxM.07.02 80	CCCCCTCTCTCTGTTTCTCGGCGAGACCCTTAATACGACTCACTATAAACT GGAAGCCCGGAACCCTTgtataatgcagacctgctgc	Forward	Upstream forward primer for null mutant generation
OL7160	LmxM.07.02 80	CGCACACGCCTCTGCCCCCTCCCCCTCCCCccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL7161	LmxM.07.02 80	gaaattaatacgactcactataggAGTAGTAGAGTGGGGGAGGgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL7162	LmxM.13.14 70	CCTCGTCCGGCAACACACACACATACAGACTAATACGACTCACTATAAAAC TGGAAGAACGTTGTGTCCgtataatgcagacctgctgc	Forward	Upstream forward primer for null mutant generation
OL7163	LmxM.13.14 70	GAGCAAACATGCACCCCCTCCCCATCCCCTccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL7164	LmxM.13.14 70	gaaattaatacgactcactataggATCAGCTGCCACGTATGTGGgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL7165	LmxM.36.43 70	AACGCTGATATAGAGCCCACCACTCTCTTCTAATACGACTCACTATAAACT GGAAGGGGCCAACCCTTgtataatgcagacctgctgc	Forward	Upstream forward primer for null mutant generation
OL7166	LmxM.36.43 70	CCCCTCCAACAAGCAAGCATCCCACGCCCGccaatttgagagacctgtgc	Reverse	Downstream reverse

				primer for null mutant generation
OL7167	LmxM.36.43 70	gaaattaatacgcactcactataggGCACGTGGAGCTCTTTGACGgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL7168	LmxM.26.23 70	CGTGACCAGCTCAGGCACCTAGGGCAGCGGTAATACGACTCACTATAAAA CTGGAAGCACACCATGTCCgtataatgcagacctgctgc	Forward	Upstream forward primer for null mutant generation
OL7169	LmxM.26.23 70	CACGTCAGGAGCAACGCTGCATGGGCTCCGccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL7170	LmxM.26.23 70	gaaattaatacgcactcactataggTCACCGCCGTAGACGGAAAGgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL7171	LmxM.31.10 90	TGGAGCACACCTCAGCCACGGTACATTCCGTAATACGACTCACTATAAAAC TGGAAGTTTGGTTCCCGGgtataatgcagacctgctgc	Forward	Upstream forward primer for null mutant generation
OL7172	LmxM.31.10 90	ACTCCGCAACAAACGTCTATACGGTCACATccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL7173	LmxM.31.10 90	gaaattaatacgcactcactataggATAGCTGCTTTGACAGCCTGgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL7174	LmxM.31.39 30	GATCGTCCCGCGTGTGCGGGTCGTTGGTGATAATACGACTCACTATAAAAC TGGAAGGTGTGCATGTCCgtataatgcagacctgctgc	Forward	Upstream forward primer for null mutant generation

OL7175	LmxM.31.39 30	CTTCGCAGATACGTACGCATAGAAAGACGCccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL7176	LmxM.31.39 30	gaaattaatacgactcactataggGTCGGTGACATGCTCGAGAgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL7177	LmxM.33.39 60	GTGATTGCTCTCGTTGTGGGGCCGACTTATTAATACGACTCACTATAAACT GGAAGCCCAATTCCCGGgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL7178	LmxM.33.39 60	GTGAGCCCTTCGCAACGGCGGGGCGCGGCTccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL7179	LmxM.33.39 60	gaaattaatacgactcactataggAAACAACCAAGCATGCCGCGgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL7180	LmxM.34.24 50	CCTTCAACCCTCCTCCACCCTTCCCTCCCTTAATACGACTCACTATAAACT GGAAGTGTGTCATGTCCgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL7181	LmxM.34.24 50	GATGCAACAGGCATAAATACAAAAGAGAAccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL7182	LmxM.34.24 50	gaaattaatacgactcactataggCGAAAAGAGCGCGAGATGAAgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL7183	LmxM.34.53 90	AGAAAGATAGTTATCATCACGCGGCGGTGCTAATACGACTCACTATAAAAC TGGAAGAAACCTTCCCGGgtataatgcagacctgtgc	Forward	Upstream forward primer for

				null mutant generation
OL7184	LmxM.34.53 90	CAGATGAGCAACAGCACCAAAAGAAAAGGCccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL7185	LmxM.34.53 90	gaaattaatacgactcactataggGACAACGAAAGGAATGACGCgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL7186	LmxM.29.21 70	GACACTGACGCGGCCGCGAATATATCTATTTAATACGACTCACTATAAACT GGAAGATGACGTTGTCCgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL7187	LmxM.29.21 70	CCTCTCTCCCGTTTCCTGTACTGCTGCCAccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL7188	LmxM.29.21 70	gaaattaatacgactcactataggTTCCTACCAGTCAGTGCTGTgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL7994	LmxM.36.63 40	AGTTTGGGTGGTTTCCGTGT	Forward	Within gene forward (for diagnostic PCR)
OL7995	LmxM.36.63 40	GTATTTGCCGCGTTCTCGAC	Reverse	Within gene reverse (for diagnostic PCR)
OL7996	LmxM.36.63 40	GAGTTTCTCGCGTCACCTCA	Forward	Forward upstream of 5' HR region (for diagnostic PCR)

OL7997	LmxM.29.09 10	CCTCCTTGCCCACCAAAGAT	Forward	Within gene forward (for diagnostic PCR)
OL7998	LmxM.29.09 10	GAGCGCACATACATTGCCAG	Reverse	Within gene reverse (for diagnostic PCR)
OL7999	LmxM.29.09 10	TCTTTCGCTTTCAGGCTGTCT	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL8001	LmxM.07.02 80	AGCTGCAGAAGGTCAAGTCC	Forward	Within gene forward (for diagnostic PCR)
OL8002	LmxM.07.02 80	CTTACCGTTCGGCTCCAACT	Reverse	Within gene reverse (for diagnostic PCR)
OL8003	LmxM.07.02 80	AGGGTGGGTACATCAGTGCT	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL8004	LmxM.13.14 70	CAACGCTGCAACAGCTTCTT	Forward	Within gene forward (for diagnostic PCR)
OL8005	LmxM.13.14 70	AGTCACGACTGATCTCCCGA	Reverse	Within gene reverse (for

				diagnostic PCR)
OL8006	LmxM.13.14 70	GGGCGCGCACTCATTTTTAT	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL8007	LmxM.36.43 70	TCCTCGTTGGACTCAAGCAC	Forward	Within gene forward (for diagnostic PCR)
OL8008	LmxM.36.43 70	GATCGGCGAACGATCTGACT	Reverse	Within gene reverse (for diagnostic PCR)
OL8009	LmxM.36.43 70	GCCACTCTCGTGGTTACTCA	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL8011	LmxM.26.23 70	CGTAATGGCACTCAATGGCG	Forward	Within gene forward (for diagnostic PCR)
OL8012	LmxM.26.23 70	GATACATGGCTCCCTCGTCG	Reverse	Within gene reverse (for diagnostic PCR)
OL8013	LmxM.26.23 70	CGGTCTCTACGCGATTCGTT	Forward	Forward upstream of 5' HR region (for

				diagnostic PCR)
OL8349	LmxM.31.10 90	ACCTTCGAGACGTTCCCAAC	Forward	Within gene forward (for diagnostic PCR)
OL8350	LmxM.31.10 90	CGCTGGTTGCAGTAAGGAGA	Reverse	Within gene reverse (for diagnostic PCR)
OL8351	LmxM.31.10 90	AACTGCTTCCGTCTGACGTT	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL8353	LmxM.31.39 30	TGACGACAACGGTGGAAGAG	Forward	Within gene forward (for diagnostic PCR)
OL8354	LmxM.31.39 30	CCAAGCGCTTCGTAATGCTC	Reverse	Within gene reverse (for diagnostic PCR)
OL8355	LmxM.31.39 30	TCACGTTGGTGGCGTTTTTC	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL8356	LmxM.33.39 60	CAAGGTGAGGACCTGCTGAG	Forward	Within gene forward (for diagnostic PCR)

OL8357	LmxM.33.39 60	AGAACAGACTGCCCGTAAGC	Reverse	Within gene reverse (for diagnostic PCR)
OL8358	LmxM.33.39 60	GGGATCCGGAGAAACAACCA	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL8360	LmxM.34.24 50	GATCTAGAGCGGTGGTTCCG	Forward	Within gene forward (for diagnostic PCR)
OL8361	LmxM.34.24 50	TACCGCATGAGGTGCGATAC	Reverse	Within gene reverse (for diagnostic PCR)
OL8362	LmxM.34.24 50	TGGAGGGCGAGAGAAGAACT	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL8364	LmxM.34.53 90	CACGGTTGTGAAGGGCAAAG	Forward	Within gene forward (for diagnostic PCR)
OL8365	LmxM.34.53 90	CAATGATTTTCGCAGGAGGCG	Reverse	Within gene reverse (for diagnostic PCR)
OL8366	LmxM.34.53 90	TCGTTCATCCTTCGGTGCAA	Forward	Forward upstream of 5' HR region

				(for diagnostic PCR)
OL8367	LmxM.29.21 70	TCGCAGTGGAACGCTACTTT	Forward	Within gene forward (for diagnostic PCR)
OL8368	LmxM.29.21 70	GACCTGTTGCACAGGAGACA	Reverse	Within gene reverse (for diagnostic PCR)
OL8369	LmxM.29.21 70	GCGATCTTTACTGCCAACGC	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL8640	LmxM.32.27 70	GCACAGACACACATACACACACACAGACTAATACGACTCACTATAAAAC TGGAAGGCAATGCGCTGGgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL8641	LmxM.32.27 70	TACATCATCTCAGCAGACAGACATAACCAAccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL8642	LmxM.32.27 70	gaaattaatacgactcactataggAACCTGGAACCAAGTGCGTTgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL8643	LmxM.24.21 40	ATCACGTCAACCCTAACCGCATCCCCACCGTAATACGACTCACTATAAAAC TGGAAGATTTCTAGCAAgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation

OL8644	LmxM.24.21 40	GACGCAACAACAAGAATACATCGACCTCCTccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL8645	LmxM.24.21 40	gaaattaatacgactcactataggGCGCCGGCGATGATGACAAGgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL9327	BSD (blasticidin resistance gene)	CACAAGGTCCCCCAGTAAAA	Reverse	Blasticidin resistance cassette for diagnostic PCRs 2
OL9371	LmxM.23.05 50	TGCTGGCACTGGCGCACGTCAAGGTAACCATAATACGACTCACTATAAAAC TGGAAGGGCTTCCGGTGGgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL9372	LmxM.23.05 50	CAAATCACCATTAGTCATCCTCTGTGGCCGccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL9373	LmxM.23.05 50	gaaattaatacgactcactataggCCGCGTGGTGTGGCCATCTAgtttagagctagaaatagc	Reverse	3' sgRNA primer
OL9374	LmxM.34.30 60	CAGACACGAAAACATAGCTTCAAACCTACCGTAATACGACTCACTATAAACT GGAAGCCTCAGTACCAAggtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL9375	LmxM.34.30 60	CCGGGTGGGGTGGGCATGGCCGGGGGGCGTccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation

OL9376	LmxM.34.30 60	gaaattaatacgactcactataggGCAGCCAGCAGCTCCGGATTgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL9377	LmxM.08.02 20	TGTGGTGCGCCAAGAACGCCACACGCCCATATACTACTATAAAAC TGGAAGTTAGGCCGGTGGgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL9378	LmxM.08.02 20	TGATTCATGGCAAGTCGTCACCGCGGTCCAccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL9379	LmxM.08.02 20	gaaattaatacgactcactataggGTGCTTGCGTCTGGAGCACGgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL9380	LmxM.01.07 10	GTTTGAGCAACAAGCACCGGAGCAGTGCCTTAATACGACTCACTATAAAAC TGGAAGAAGACGTACCAAgataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL9381	LmxM.01.07 10	TGACCAGCGGTAGAACCTCCGTTGCGCCCGccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL9382	LmxM.01.07 10	gaaattaatacgactcactataggACGACGGCGTAGATGCGCGAgtttagagctagaaatagc	Reverse	3' sgRNA primer
OL9383	LmxM.02.03 90	CGCTTACAGGACGCAGGGTGTGTGGGGAATACTACTACTATAAAA CTGGAAGGCTATATCGTGGgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL9384	LmxM.02.03 90	TCCCCAAGCCCCACACGCACACAAAAGCCTccaatttgagagacctgtgc	Reverse	Downstream reverse primer for

				null mutant generation
OL9385	LmxM.02.03 90	gaaattaatacgactcactataggACGCGAGAGAAAGGTTGCGAgtttagagctagaaatagc	Reverse	3' sgRNA primer
OL9386	LmxM.04.06 80	CGTTGTACGCCTCCCTTCGCTCCCCGTCCTTAATACGACTCACTATAAAAC TGGAAGATATCTCTCCAAgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL9387	LmxM.04.06 80	TGCACACGCACGCACGCACATGTCAGGCATccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL9388	LmxM.04.06 80	gaaattaatacgactcactataggAGGCATACACACATGTCACAgtttagagctagaaatagc	Reverse	3' sgRNA primer
OL9389	LmxM.05.09 30	TCCCCGTGTGTCTTTGGACTCCTTCTGCCCTAATACGACTCACTATAAACT GGAAGCGATAATCGTGGgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL9390	LmxM.05.09 30	TACGTCGCGGCGCTACCGCCACTCCCCCCCccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL9391	LmxM.05.09 30	gaaattaatacgactcactataggTGTGTGTGTGTGTGTGTGGAgtttagagctagaaatagc	Reverse	3' sgRNA primer
OL9392	LmxM.07.08 50	AAGTCAGGGGACCGACTCTCCCCTGTGCCGTAATACGACTCACTATAAAAC TGGAAGTATAGTCTCCAAgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL9393	LmxM.07.08 50	GCAGTCGCCCACCCCCACCCCCCACTGCCAccaatttgagagacctgtgc	Reverse	Downstream reverse

				primer for null mutant generation
OL9394	LmxM.07.08 50	gaaattaatacgactcactataggCTGGAGAGGAACCAAAACGAgtttagagctagaaatagc	Reverse	3' sgRNA primer
OL9395	LmxM.13.15 80	ATCTCTCCTCCCTCGGTGGTCTTTTCCCCATAATACGACTCACTATAAACT GGAAGTAGCGATCGTGGgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL9396	LmxM.13.15 80	AAAGCACCGAGACAATCAACAGTGGCTCCAccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL9397	LmxM.13.15 80	gaaattaatacgactcactataggTTTCGCAGTATCACACCCGCgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL9398	LmxM.22.06 10	CTTTTTTAATTGGTGTGAGCGCCGTCTTTCTAATACGACTCACTATAAACT GGAAGCGCGATCTCCAAgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL9399	LmxM.22.06 10	AACCAGGGAACTAGCCACGCACATGCCGACccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL9400	LmxM.22.06 10	gaaattaatacgactcactataggACACCGCACCTGTAGCAGTGgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL9401	LmxM.24.17 10	TTTTTCTCCTCGCTCACTTCATTTTCAGATCTAATACGACTCACTATAAACTG GAAGATCGCATCGTGGgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation

OL9402	LmxM.24.17 10	ACGACACGGCGAGGGCCCGCCAAGTCGTGCccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL9403	LmxM.24.17 10	gaaattaatacgactcactataggGGTGGAGCGCGTGAAGGCGGgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL9404	LmxM.24.21 30	CGTGCCTGCTCTTGTGCGCTTGCCTCGGTGTAATACGACTCACTATAAAAC TGGAAGGCGCTTCTCCAgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL9405	LmxM.24.21 30	ATCTCTAGGACCCAACAAACCACCGACTCAccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL9406	LmxM.24.21 30	gaaattaatacgactcactataggGTCACACACGCAACACGTAGgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL9407	LmxM.31.07 00	TCTTCTCCGTTGCTGTTTCTTGCTCCACCATAATACGACTCACTATAAACT GGAAGTGAATTACGCGGgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL9408	LmxM.31.07 00	ACACCTTCACGAGGAATGCCTCAAGCGCACccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL9409	LmxM.31.07 00	gaaattaatacgactcactataggGAGTAAGCGGTGGGTATAAGgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL9410	LmxM.31.09 60	AACGGAAGAAGAATTAAGGCGCCCTCCTTAATACGACTCACTATAAAAC TGGAAGCATTCACTCTAAgtataatgcagacctgtgc	Forward	Upstream forward primer for

				null mutant generation
OL9411	LmxM.31.09 60	TCGTACGTCGCGCTGTTTTGGCTGCTACCGccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL9412	LmxM.31.09 60	gaaattaatacgactcactataggTCGACGTCACGGCCAGACCTgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL9413	LmxM.33.09 00	ACCAAAGAGTTCCAGTCCAGACCTTCCCCATAATACGACTCACTATAAACT GGAAGGGAAGAGGACGGgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL9414	LmxM.33.09 00	CTTCGTAGTTTATCAGAGGGAGGTTCCGCCccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL9415	LmxM.33.09 00	gaaattaatacgactcactataggAGTGAACACAGATCCCTAGGgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL9416	LmxM.33.15 5	TCCTTCATGTTTGCTGGGGTTTTTTTTTCATAATACGACTCACTATAAACTG GAAGACGGCGCATTCCgtataatgcagacctgtgc	Forward	Upstream forward primer for null mutant generation
OL9417	LmxM.33.15 55	TTGACACGAAGTGGTTGCACAAGCGAGTATccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL9418	LmxM.33.15 55	gaaattaatacgactcactataggGACCGGGTACTAGTTCCGTAgtttagagctagaaatagc	Reverse	3' sgRNA primer
OL9419	LmxM.34.13 00	CCTATCTGTTCTGTTGCTCTCATCTCACCTAATACGACTCACTATAAACT GGAAGCCTTCAGGAGGGgtataatgcagacctgtgc	Forward	Upstream forward

				primer for null mutant generation
OL9420	LmxM.34.1300	TCTTCGGTTCTCTCCTTGGTTTCACTTCCccaatttgagagacctgtgc	Reverse	Downstream reverse primer for null mutant generation
OL9421	LmxM.34.1300	gaaattaatacgcactcactataggAAAGTGAATCCGGGAGCGGTgttttagagctagaaatagc	Reverse	3' sgRNA primer
OL9479	LmxM.32.2770	CCAAAGGGTTTCTTCCGCAC	Forward	Within gene forward (for diagnostic PCR)
OL9480	LmxM.32.2770	TGTTTGCCATGGACGATGCT	Reverse	Within gene reverse (for diagnostic PCR)
OL9481	LmxM.32.2770	CATTTCCGGCACACGCTTTCA	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL9482	LmxM.24.2140	GCATGGATTCTGCGTTGACC	Forward	Within gene forward (for diagnostic PCR)
OL9483	LmxM.24.2140	GGTACAAAGACGCGCTTCAC	Reverse	Within gene reverse (for diagnostic PCR)
OL9484	LmxM.24.2140	AATGTACAAGCGCACCCACT	Forward	Forward upstream of 5' HR region

				(for diagnostic PCR)
OL9545	LmxM.23.05 50	ATTCTGGAGCGGTGCAAAGA	Forward	Within gene forward (for diagnostic PCR)
OL9546	LmxM.23.05 50	ACGAGAGACGTGGTTGTCAC	Reverse	Within gene reverse (for diagnostic PCR)
OL9547	LmxM.23.05 50	GAGGTCGGCGAATGAAAGGT	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL9548	LmxM.34.30 60	GACTTTATCGCTGCGGCAAG	Forward	Within gene forward (for diagnostic PCR)
OL9549	LmxM.34.30 60	CCGTTCCACAGCATACCCTT	Reverse	Within gene reverse (for diagnostic PCR)
OL9550	LmxM.34.30 60	TTTCATCTCGTACCCGCCTG	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL9551	LmxM.01.07 10	CACCGTCGTCGACATGGATT	Forward	Within gene forward (for diagnostic PCR)

OL9552	LmxM.01.07 10	TGGGCCATGTGAATGACTCG	Reverse	Within gene reverse (for diagnostic PCR)
OL9553	LmxM.01.07 10	CTGTCGACATCGCTCCTAGTC	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL9554	LmxM.02.03 90	TCCGTCTCTCCCTCAAGTACAA	Forward	Within gene forward (for diagnostic PCR)
OL9555	LmxM.02.03 90	TAGCCCTTCCAATGATGCCT	Reverse	Within gene reverse (for diagnostic PCR)
OL9556	LmxM.02.03 90	ATCGTGAGAGGGAGACGGG	Forward	Forward upstream of 5' HR region (for diagnostic PCR) 1
OL9557	LmxM.04.06 80	ACAACGCGCATTATCAAGGAG	Forward	Within gene forward (for diagnostic PCR)
OL9558	LmxM.04.06 80	GGCTCGACATGAGAATCTGGA	Reverse	Within gene reverse (for diagnostic PCR)
OL9559	LmxM.04.06 80	TGCCTCTTACCCATCCAACAC	Forward	Forward upstream of 5' HR region

				(for diagnostic PCR)
OL9560	LmxM.05.09 30	ACGCGTCTTCCTCCATCAAG	Forward	Within gene forward (for diagnostic PCR)
OL9561	LmxM.05.09 30	GCCTCCTTTGTTTCGAGTGC	Reverse	Within gene reverse (for diagnostic PCR)
OL9562	LmxM.05.09 30	GACTCCGATCATCCCGTGTG	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL9563	LmxM.07.08 50	CCGACCCGGACATTCATCTG	Forward	Within gene forward (for diagnostic PCR)
OL9564	LmxM.07.08 50	AGACTCGCGTAACCCTCCAT	Reverse	Within gene reverse (for diagnostic PCR)
OL9565	LmxM.07.08 50	AATCGAGATCCGCTTACCGC	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL9754	LmxM.07.02 80	TCGGCACGCGCGCTTGCGCAGGAGTCGATGccaatttgagagacctgtgc	Reverse	Downstream reverse primer for

				null mutant generation 2
OL9755	LmxM.07.02 80	gaaattaatacgactcactataggGGGAAGGGTTCCAATAGTTGGGGgttttagagctagaaa tagc	Forward	5' sgRNA primer 2
OL9756	LmxM.07.02 80	gaaattaatacgactcactataggTGGACACGCTGAGCGAGACCGGGgttttagagctagaa atagc	Reverse	3' sgRNA primer 2
OL10120	LmxM.02.03 90	ACGCGCACGAATTCTTTTCC	Forward	Forward upstream of 5' HR region (for diagnostic PCR) (2)
OL10121	LmxM.08.02 20	TACCTGGGGCTCTCGATTCA	Forward	Within gene forward (for diagnostic PCR)
OL10122	LmxM.08.02 20	TTGAGTTCCCGACATCTGCC	Reverse	Within gene reverse (for diagnostic PCR)
OL10123	LmxM.08.02 20	GTACGCACCTCCCAACACTT	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL10124	LmxM.13.15 80	GAAACTTTCGCCTGCTGGA	Forward	Within gene forward (for diagnostic PCR)
OL10125	LmxM.13.15 80	TCAGTACGTAGACCCCTCCG	Reverse	Within gene reverse (for diagnostic PCR)

OL10126	LmxM.13.15 80	TAAGCGCAATTCCATCCCGT	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL10127	LmxM.22.06 10	GATCAACCTCAGCAGGGTGT	Forward	Within gene forward (for diagnostic PCR)
OL10128	LmxM.22.06 10	GCGCCTCTAGGTTAAGCGAT	Reverse	Within gene reverse (for diagnostic PCR)
OL10129	LmxM.22.06 10	CGTGAAGAGGCCGCAAAAAT	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL10132	LmxM.24.17 10	CTCCCTTCCTTCTGCCAGTG	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL10166	LmxM.31.07 00	TCCAGCTGCCATCCGATTAC	Forward	Within gene forward (for diagnostic PCR)
OL10167	LmxM.31.07 00	GTCCGTCAAACCCTTCGAGT	Reverse	Within gene reverse (for diagnostic PCR)

OL10168	LmxM.31.07 00	TCTGGCTGATACCGACACAC	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL10169	LmxM.31.09 60	TGGACGAGTGTGCATTTCCA	Forward	Within gene forward (for diagnostic PCR)
OL10170	LmxM.31.09 60	CGCCACGATGCTTGATTAGC	Reverse	Within gene reverse (for diagnostic PCR)
OL10171	LmxM.31.09 60	CGCAGACTTGGCATCGAGTA	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL10172	LmxM.33.09 00	AGCTCTTACCACCCCGAGAA	Forward	Within gene forward (for diagnostic PCR)
OL10173	LmxM.33.09 00	CGGGCTACAGCACCAGTAAT	Reverse	Within gene reverse (for diagnostic PCR)
OL10174	LmxM.33.09 00	GGAAGAGGTTGCGTCCTTG	Forward	Forward upstream of 5' HR region (for diagnostic PCR)

OL10175	LmxM.33.15 55	TCTTAAGGAGCTGGCCAACAG	Forward	Within gene forward (for diagnostic PCR)
OL10176	LmxM.33.15 55	CTTGAACCTCTTCAGGCGTTGC	Reverse	Within gene reverse (for diagnostic PCR)
OL10177	LmxM.33.15 55	TAGTCGCAGTTCGCAGAGTG	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL10283	LmxM.24.21 30	CGGCTGCAAAAGGAACTCGTA	Forward	Within gene forward (for diagnostic PCR)
OL10284	LmxM.24.21 30	GTACAGCGATGGTGAAGTCT	Reverse	Within gene reverse (for diagnostic PCR)
OL10285	LmxM.24.21 30	GATTCCTGCGGCTGTTCTTA	Forward	Forward upstream of 5' HR region (for diagnostic PCR)
OL10329	LmxM.24.17 10	CATGCAGAAGCTGGATGGAGT	Forward	Within gene forward (for diagnostic PCR)
OL10330	LmxM.24.17 10	GATGATGTAGTTGCCGCCCAT	Reverse	Within gene reverse (for

				diagnostic PCR)
OL10540	Recoded LmxM.13.15 80	TCCAGGGACCAGCAATGGTCGAGGTTCCGCGC	Forward	Forward primer for integration into pETFPP vectors
OL10541	Recoded LmxM.13.15 80	TGAGGAGAAGGCGCGTCAGTAGGTACTACCCTCC	Reverse	Reverse primer for integration into pETFPP vectors
OL10602	pETFPP vectors	TAATACGACTCACTATAGGG	Forward	T7 sequencing primer
OL10603	pETFPP vectors	TATGCTAGTTATTGCTCAGCGGT	Reverse	T7 term sequencing primer
OL10633	pGL2813 (recoded UBA1a)	CACGGTGACCTGTGTCGTAA	Forward	Sequencing primer
OL10634	pGL2813 (recoded UBA1a)	GTTCTTCGGCTGCCGTTTTT	Forward	Sequencing primer
OL10635	pGL2813 (recoded UBA1a)	ATGGGACCAGAAACGGAAGG	Forward	Sequencing primer
RB1	Recoded LmxM.23.05 50	TCCAGGGACCAGCAATGCTTTCTGAGGAAGAGCAAAAAC	Forward	Forward primer for integration into pETFPP vectors
RB2	Recoded LmxM.23.05 50	TGAGGAGAAGGCGCGTTAAAAGCGATAGCGGTAGCGGATG	Reverse	Reverse primer for integration

				into pETFPP vectors
RB3	Recoded LmxM.04.06 80	TCCAGGGACCAGCAATGTTGACCACTCGTATCATTAAGG	Forward	Forward primer for integration into pETFPP vectors
RB4	Recoded LmxM.04.06 80	TGAGGAGAAGGCGCGTCATGGTTTGGCGTACTTACGAG	Reverse	Reverse primer for integration into pETFPP vectors

Plasmid name	Description	Backbone	Resistance	Source
pGL2662	Template vector for PCR amplification of blasticidin resistance cassette for gene knockout	NA	Ampicillin	pTBlast_v1, Gluenz lab, University of Oxford
pGL2663	Template vector for PCR amplification of G418 resistance cassette for gene knockout	NA	Ampicillin	pTNeo_v1, Gluenz lab, University of Oxford
pGL2667	Template vector for PCR amplification of puromycin resistance cassette for gene knockout	NA	Ampicillin	pTPuro_v1, Gluenz lab, University of Oxford
pETFPP_4	HIS-Im9-3cProtease-ORF	pETYSBLIC3c	Kanamycin	York Technology Facility

pGL2813	UBA1a in pETFPP_4	pETFPP_4	Kanamycin	Self-generated
pGL2817	UBC2 in pETFPP_4	pETFPP_4	Kanamycin	Self-generated
pGL2837	UEV1 in pETFPP_4	pETFPP_4	Kanamycin	Self-generated

S3 Table. Individual viability assay of $\Delta ubc1$, $\Delta ubc2$, $\Delta uev1$ and $\Delta hect2$ during axenic amastigote differentiation. Raw fluorescence (590 nm) readings and relative viabilities compared to the Cas9 T7 cell line are included.

Experiment 1														
0 h					48 h					120 h				
Nega tive	T7 Cas9 pare ntal	Δubc 1	Δubc 2	$\Delta uev1$	Nega tive	T7 Cas9 pare ntal	Δubc 1	Δubc 2	$\Delta uev1$	Nega tive	T7 Cas9 pare ntal	Δubc 1	Δubc 2	$\Delta uev1$
8452	3731 3	3856 4	5501 8	36498	1627 9	4883 2	4655 7	5116 9	46109	1086 2	5069 6	2429 6	1691 4	15591
8594	3718 1	3795 7	5535 9	36876	1668 4	4775 0	4329 5	5109 8	46276	1130 8	4523 7	1926 6	1664 2	15236
8725	3808 1	3690 2	5881 8	38100	1712 0	4767 3	3736 5	4962 3	41846	1142 2	4377 9	1722 2	1667 2	15481
8620	3673 7	3735 8	5531 9	36852	1704 4	4506 0	3730 4	4156 9	43443	1137 2	4148 0	1750 0	1640 1	14850
8659	3668 2	3960 6	5875 1	37500	1678 5	4682 6	4364 7	4489 5	45494	1140 0	4614 3	1808 0	1641 4	14917
8371	3627 0	4048 9	5850 1	37343	1627 4	5781 7	3848 4	5653 5	50919	1092 5	5632 1	2335 6	1660 8	15192
Experiment 2														

0 h					48 h					120 h				
Nega tive	T7 Cas9 pare ntal	Δubc 1	Δubc 2	$\Delta uev1$	Nega tive	T7 Cas9 pare ntal	Δubc 1	Δubc 2	$\Delta uev1$	Nega tive	T7 Cas9 pare ntal	Δubc 1	Δubc 2	$\Delta uev1$
9856	3810 2	3792 6	5608 2	41017	1756 5	5938 7	4722 2	5082 2	47819	1418 3	4264 8	2259 0	1908 0	18250
1001 6	3845 3	3877 0	5694 6	41503	1827 5	5338 3	4265 3	4883 7	49687	1494 5	3893 0	2042 9	1949 3	17989
9990	3857 2	3805 9	5646 6	41668	1864 5	5610 0	4530 6	5072 8	56363	1495 6	3836 5	2008 7	2000 0	18326
9980	3804 5	3734 9	5846 2	40373	1830 6	5656 2	4651 9	4747 8	50766	1493 8	3871 3	2069 1	1989 8	18012
9974	3768 8	3746 4	5657 7	41282	1842 9	5026 6	4352 6	4994 1	48353	1466 5	3958 3	2101 9	2038 0	18160
9714	3768 7	3783 7	5583 3	40204	1759 0	5011 8	4643 0	4276 1	48598	1349 8	3664 0	2314 3	1985 7	18807

Experiment 1								
0 h			48 h			120 h		
Negative	T7 Cas9 parental	$\Delta hec2$	Negative	T7 Cas9 parental	$\Delta hec2$	Negative	T7 Cas9 parental	$\Delta hec2$
21143	54662	54366	17452	52061	36335	13387	48159	15157
21585	55822	54093	18396	53755	40325	13972	45988	15441
21692	55745	54605	18383	53359	42512	13946	43715	15538
21441	55740	54675	18420	52141	41370	13849	42746	15542
21401	55370	55008	18401	52583	38966	13797	43026	15526
21238	56019	54677	17856	52287	34640	13235	46959	15239

Experiment 2								
0 h			48 h			120 h		
Negative	T7 Cas9 parental	Δ hect2	Negative	T7 Cas9 parental	Δ hect2	Negative	T7 Cas9 parental	Δ hect2
26946	45285	57003	26946	45107	29833	22632	30187	14212
28348	45175	40858	28348	45503	32500	35053	40382	14506
28254	47848	57771	28254	46726	34150	28293	40316	14696
28114	42693	57020	28114	47753	34330	26289	39611	14533
27386	46252	53986	27386	48782	30720	26922	40196	13968
27880	46024	43447	27880	45568	25247	29833	39931	13100

Experiment 1				Experiment 2				Average			
Time into differentiation (h)				Time into differentiation (h)				Time into differentiation (h)			
Cell line	0	48	120	Cell line	0	48	120	Cell line	0	48	120
Cas9 T7	1	1	1	Cas9 T7	1	1	1	Cas9 T7	1	1	1
Δ ubc1	1.05185	0.75587	0.24232	Δ ubc1	0.99324	0.75042	0.27607	Δ ubc1	1.02254	0.75315	0.2592
Δ ubc2	1.71697	1.00481	0.14957	Δ ubc2	1.66158	0.83757	0.21344	Δ ubc2	1.68928	0.92119	0.1815
Δ uev1	1.01211	0.89745	0.11082	Δ uev1	1.10354	0.88834	0.15139	Δ uev1	1.05783	0.8929	0.1311
Δ hect2	0.97103	0.60421	0.05444	Δ hect2	1.34611	0.17646	0	Δ hect2	1.15857	0.39034	0.02722

S4 Table. Data collection and refinement statistics for UBC2-UEV1 crystal structure.

Data collection	
Space group	P2 ₁
Cell dimensions	
a, b, c (Å)	32.52, 72.56, 120.10
α, β, γ (°)	90, 91.84, 90
Resolution (Å) ^a	46.25-1.70 (1.73 -1.70)
Total No. of reflections ^a	260,600 (14,037)
No. of unique reflections ^a	61,439 (3,267)
Completeness (%) ^a	100 (100)
Multiplicity ^a	4.2 (4.3)
I/σ(I) ^a	7.3 (1.2)
R _{merge} ^b	0.096 (1.449)
R _{pim} ^c	0.050 (0.747)
CC1/2	0.9967 (0.529)
Overall B factor from Wilson plot (Å ²)	26.32
Refinement	
Resolution (Å)	46.29-1.70
R _{work} ^d / R _{free} ^e	0.222 / 0.260
No reflections (free)	61,440 / 3,018
Completeness (%)	100 (99.7)
No. of non-H atoms	
Protein	4,652
Water	266

Ion	5
Total	4,923
B-factors	
Protein Main Chain (\AA^2)	29.4
Protein Side Chain (\AA^2)	33.8
Water (\AA^2)	32.4
R.m.s. deviations ^f	
Bonds (\AA)	0.0087
Angles ($^\circ$)	1.597
Ramachandran plot ^g	96.41 / 2.51 / 1.08

^a Values in parentheses correspond to the outer resolution shell

^b $R_{\text{merge}} = \frac{\sum_{hkl} \sum_i |I_i - \langle I \rangle|}{\sum_{hkl} \sum_i I_i}$ where I_i is the intensity of the i th measurement of a reflection with indexes hkl and $\langle I \rangle$ is the statistically weighted average reflection intensity.

^c $R_{\text{pim}} = \frac{\sum_{hkl} [1/(n-1)]^{1/2} \sum_i |I_i - \langle I \rangle|}{\sum_{hkl} \sum_i I_i}$

^d $R_{\text{factor}} = \frac{\sum ||F_o| - |F_c||}{\sum |F_o|}$ where F_o and F_c are the observed and calculated structure factor amplitudes, respectively.

^e R_{free} is the R -factor calculated with 5 % of the reflections chosen at random and omitted from refinement

^f Root-mean-square deviation of bond lengths and bond angles from ideal geometry.

^g Percentage of residues in preferred/allowed/outlier regions.

A

	590	600	610	620	630
LmUBA1a	YDPP	EKG	IPLC	TLK	NFPN
LmUBA1b	GAAD	QTGG	IPMC	TLR	NFPY
HsUBA1	QDPP	EKS	IPIC	TLK	NFPN
HsUBA6	RDPF	EEE	IPFC	TLK	SFPA

B

	80	90	100
UBC1	IS..S..F...	WHPN.VY.A.D....	GRVCI
UBC2	LT..R..I...	YHPN.VD.K.V....	GRICL
UBC3	FT..R..S...	GRFA.VR.E.....	KICST
UBC4	IT..K..V...	FHPN.VEFH.T....	GNVCL
UBC5	RT..K.D.V...	FHPN.VY.V.D....	GNICM
UBC6	LT..R..I...	FNPV.VS.E.D....	GAVCE
UBC7	TT..P..C...	FHPN.VD.T.R....	GAICL
UBC8	CN..R..I...	LHPN.VD.ERS....	GSVCL
UBC9	TP..V..L...	FHPN.VY.P.S....	GTVCL
UBC10	RT..K..I...	FHPN.IS.E.R....	GDICV
UBC11	CT..P..S...	GR..LK.T.G....	MQICL
UBC12	TGPRR..I...	FHPN.IE.G.DDGKEDWG	VCLGM
UBC13	TT..K..I...	YHPN.IN.N.N....	GGICL
UBC14	TG..P..F...	RSHP.LVQ.E.....	RCLQH
UEV1	LS..K..V...	NLPC.VD.S.D....	GTVNRE
HsUBE2A	VS..K..M...	FHPN.VY.A.D....	GSICL
HsUBE2B	LS..K..M...	FHPN.VY.A.D....	GSICL
HsUBE2C	LT..P..C...	YHPN.VD.T.Q....	GNICL
HsUBE2D1	TT..K..I...	YHPN.IN.S.N....	GSICL
HsUBE2D2	TT..R..I...	YHPN.IN.S.N....	GSICL
HsUBE2D3	TT..R..I...	YHPN.IN.S.N....	GSICL
HsUBE2D4	TT..K..I...	YHPN.IN.S.N....	GSICL
HsUBE2E1	RT..R..I...	YHCN.IN.S.Q....	GVICL
HsUBE2E2	RT..R..I...	YHCN.IN.S.Q....	GVICL
HsUBE2E3	RT..R..I...	YHCN.IN.S.Q....	GVICL
HsUBE2G1	IT..E..I...	WHPN.VD.K.N....	GDVCI
HsUBE2G2	TC..E..M...	FHPN.IY.P.D....	GRVCI
HsUBE2H	MN..A..KI...	FHPN.ID.E.AS....	GTVCL
HsUBE2J1	LT..A..N...	GRFE.VG.K.....	KICLS
HsUBE2J2	IT..P..N...	GRFK..C.N....	TRLC
HsUBE2K	IT..K..I...	WHPN.IS.SVT....	GAICL
HsUBE2L3	KT..K..I...	YHPN.ID.E.K....	GQVCL
HsUBE2N	MT..K..I...	YHPN.VD.K.L....	GRICL
HsUBE2N-like	MT..K..I...	YHPN.VD.K.L....	ERISL
HsUBE2O	LS..Q..C...	SGRLNPN.LY.D.N...	GKVCV
HsUBE2Q1	VS..P..V...	LSGGYVL.G.G....	GAICM
HsUBE2Q2
HsUBE2R1	LT..K..M...	WHPN.IY.E.T....	GDVCI
HsUBE2R2	LT..K..M...	WHPN.IY.E.N....	GDVCI
HsUBE2S	LT..K..I...	FHLN.VG.A.N....	GEICV
HsUBE2T	LT..P..I...	YHPN.ID.S.A....	GRICL
HsUBE2U	IT..I..P...	FHPN.VD.PHT....	GQPC
HsUBE2V1	VT..K..I...	NMNG.VN.SSN....	GVVDP
HsUBE2V2	VT..K..I...	NMNG.IN.NSS....	GMVD
HsUBE2W	TG..ENIP...	VHPH.VY.S.N....	GHICL


C

	280	290	300	310
HECT1	..TQADADTV	LPRAATCF	FFA.IEVPDY	STLEV
HECT2	LAMNASGPDH	LPSAATCF	S.TLKLPPY	RDYAT
HECT3	..RWAVTKEDL	LPQAHTCF	N.KIDLPVY	TSAAV
HECT4	VDSSSCDDTR	LPQASTCF	Y.WLRLPCY	SSSTAV
HECT5	..GGVNHDS	LPTVSTCF	FL.YLKLPRY	SSKAV
HECT6	LVGCGDEPGR	LPTAHTCF	SPIS.LLLQPC	TTTADL
HECT7	EDASIKEEDM	LPSAMTC	Q.N.YLKLQ	PYNTKE
HECT8	..ENNESVDY	LPRAQAC	GS.LITIPQY	TSLEMF
HECT9	KDMESKKEQT	LPSCNTCF	FL.YLKLPPY	TTREIM
HECT10	ALHLPESSEWA	LPSVNTCF	FR.YLKLPPY	PSSEEL
HECT11	..THNGNKDT	LPTSSTCF	FFA.LRLPSY	SSSAEK
HECT12	..LRCADEGR	LPVGHGCT	N.QLDLPD	YNDKQI
HECT13
HECT14
HsHWE1	..RDDRSTDR	LPSAHTCF	N.QLDLPAY	ESFEKL
HsITCH	..K.VGKENW	LPRTSHTCF	N.RLLDPYK	SSYEQL
HsUBE3A	..KNGPDTER	LPTSHTCF	N.VLLLPEY	SSSKEKL
HsUBE3C	IHNHGSGLER	LPTASTCF	MN.LLKLPEY	DETLL

Supplementary figure 1. Alignments of *L. mexicana* and *H. sapiens* protein sequences.


Sequences obtained from TriTrypDB (*L. mexicana*) and NCBI or UniProt (*H. sapiens*) were

aligned using T-Coffee. Residues in the local vicinity of the putative active site cysteine of **A** ubiquitin E1-activating enzymes, **B** E2-conjugating enzymes and **C** E3 ligase HECT domains are shown. Red boxes indicate amino acid identity, red characters show similarity within the highlighted group and blue frames highlight similarity across groups. Black bar indicates the position of the conserved HPN motif in E2 genes and the black triangles highlight the conserved catalytic cysteine residues in all classes of protein. *L. mexicana* genes are indicated with the prefix Lm and *H. sapiens* genes with the prefix Hs. Where no prefix is given, sequences belong to *L. mexicana*.





Supplementary figure 2. Confirmation of the null mutant library. **A** Schematic of PCRs performed to identify null mutants from heterozygotes or mutants with additional gene copies. Primers were designed to amplify regions within the gene of interest (red primers) and between the 5'-UTR of the gene and the blasticidin resistance cassette of the edited DNA (black primers). **B** PCRs performed on genomic DNA from parental (P) and mutant (M) cells are shown for genes that were deleted successfully. Due to the large size of *HECT3* (18.6

kb), only the first 8 kbp of the gene was targeted for editing. The *HECT3* gene PCR was designed to amplify within the targeted region. **C** PCRs performed on genomic DNA from parental (P) and mutant (M) cells are shown for genes that could not be deleted (no mutant clones were obtained for UBC13). **D** Relative viability of $\Delta ubc1$, $\Delta ubc2$, $\Delta uev1$ and $\Delta hect2$ compared to the parental Cas9 T7 line during promastigote to amastigote differentiation. Time elapsed since the initiation of differentiation is marked on the x-axis. Data are an average of two independent experiments, each with 6 biological replicates.



Supplementary figure 3. Structural analysis of the UBC2-UEV1 heterodimer. **A** Superposition of chains A and B for UBC2-UEV1 (dark green and dark blue, PDB ID: 6ZM3), HsUBE2N-UBE2V2 (darker green and darker blue, PDB ID: 1J7D) and ScUbc13-Mms2 (light green and light blue, PDB ID: 1JAT). Locations of the N- and C-termini are shown. **B** Zoom-in

of the interface between UBC2 and UEV1 showing residues thought to contribute most significantly to complex formation according to analysis in the program PISA (96). Residues are labelled in black for UBC2 and blue for UEV1. Residues are coloured by atom (red, oxygen; blue, nitrogen). Hydrogen bonds are denoted by dashed lines. **C** Superposition of UBC2-UEV1 onto the structure of the UBE2N-UBE2V2-Ub complex (PDB ID: 2GMI) (50) showing UBC2 (green) and UEV1 (blue) as space fill models and the positions of acceptor and donor ubiquitins (orange ribbons) obtained from the UBE2N-UBE2V2-Ub structure. C85 is represented by black (carbon) and yellow (sulfur) spheres. K63 of ubiquitin is highlighted as cylinders coloured by atom (blue, nitrogen).



Supplementary figure 4. *UBA1b cannot transfer ubiquitin to UBC2 in vitro.* **A** Alignment of *L. mexicana* and *H. sapiens* ubiquitin protein sequences. Red boxes indicate amino acid identity, red characters show similarity within the highlighted group and blue frames highlight similarity across groups. **B** UBA1b and ubiquitin were incubated with UBC2, UEV1 and ATP as indicated in ubiquitination assay buffer for 30 min at 30°C. Samples were treated with either reducing or non-reducing sample buffer and visualised by SDS-PAGE with InstantBlue stain.