

This is a repository copy of *Prevalence and Predictors of Reincarceration after Correctional Center Release: A Population-based Comparison of Individuals with and without Schizophrenia in Ontario, Canada: Prévalence et prédicteurs de la réincarcération après la libération d'un centre correctionnel : une comparaison dans la population-de personnes souffrant ou non de schizophrénie en Ontario, Canada.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/165974/>

Version: Accepted Version

Article:

Mahentharan, Mayuri, Kouyoumdjian, Fiona G., de Oliveira, Claire orcid.org/0000-0003-3961-6008 et al. (4 more authors) (2020) Prevalence and Predictors of Reincarceration after Correctional Center Release: A Population-based Comparison of Individuals with and without Schizophrenia in Ontario, Canada: Prévalence et prédicteurs de la réincarcération après la libération d'un centre correctionnel : une comparaison dans la population-de personnes souffrant ou non de schizophrénie en Ontario, Canada. Canadian Journal of Psychiatry. ISSN: 0706-7437

<https://doi.org/10.1177/0706743720953018>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Prevalence and predictors of re-incarceration after correctional centre release: a population-based comparison of individuals with and without schizophrenia in Ontario, Canada.

Journal:	<i>The Canadian Journal of Psychiatry/La Revue canadienne de psychiatrie</i>
Manuscript ID	CJP-2020-140-OR.R1
Manuscript Type:	Original Research
Date Submitted by the Author:	21-Jun-2020
Complete List of Authors:	Mahentharan, Mayuri; Ontario Renal Network Kouyoumdjian, Fiona; McMaster University, Department of Family Medicine de Oliveira, Claire; University of York, Hull York Medical School/Centre for Health Economics; Centre for Addiction and Mental Health, Institute for Mental Health Policy Research Iwajomo, Tomi; Centre for Addiction and Mental Health, General Psychiatry Simpson, Alexander; Centre for Addiction and Mental Health (CAMH), Chief of Forensic Psychiatry Jones, Roland; Centre for Addiction and Mental Health, ; University of Toronto, Psychiatry Kurdyak, Paul; Centre for Addiction and Mental Health, General Psychiatry
Key Words:	Schizophrenia, Incarceration, Healthcare policy
Abstract:	<p>Objectives: Individuals with schizophrenia are over-represented in correctional facilities relative to their population-based prevalence. The purpose of this study was to determine the rate and predictors of reincarceration of individuals with schizophrenia after release from correctional facilities.</p> <p>Methods: This was a retrospective cohort study that included all people released from Ontario's provincial correctional facilities from January 1 to December 31, 2010. Individuals with schizophrenia were identified using a population-based algorithm. The primary outcome was time to reincarceration. Covariates included sociodemographic characteristics (age, sex, neighbourhood income quintile, urban/rural residence), health service utilization (primary care physician visits, psychiatrist visits, psychiatric and non-psychiatric hospitalizations, emergency department visits) and other clinical comorbidity. Survival analysis was used to examine the association between schizophrenia and re-incarceration.</p> <p>Results: Among 46,928 individuals, N = 3237 (7%) had a diagnosis of schizophrenia. Approximately 67.5% of these individuals were reincarcerated within five years following their first release in 2010,</p>

	<p>compared to 58.8% of individuals without schizophrenia. Individuals with schizophrenia were 40% (HR: 1.39, 95% CI: 1.33-1.45) more likely to be reincarcerated following release than the control group after adjusting for demographic characteristics. This association reduced to 8% (1.08, 95% CI:1.03-1.14) after adjusting for prior health service utilization, prior correctional involvement, and comorbidities.</p> <p>Conclusion: Individuals with schizophrenia were more likely to experience reincarceration after release from correctional facilities. This risk is partly explained by prior correctional involvement, health service utilization and comorbidities. Future research should focus on risk factors predicting the higher reincarceration rate and interventions to reduce correctional involvement.</p>

SCHOLARONE™
Manuscripts

Prevalence and predictors of re-incarceration after correctional centre release: a population-based comparison of individuals with and without schizophrenia in Ontario, Canada.

Mayuri Mahentharan MSc¹

Fiona G. Kouyoumdjian MD MPH PhD^{2,3}

Claire de Oliveira PhD^{1,3,4}

Tomisin Iwajomo MPH^{1,3}

Alexander I F Simpson MBChB BMedSci⁵

Roland Jones MD PhD⁵

Paul Kurdyak MD PhD^{1,3,4}

Institutional Affiliations

¹Institute for Mental Health Policy Research, CAMH, Toronto, ON

²Department of Family Medicine, McMaster University, Hamilton, ON

³ICES, Toronto, ON

⁴Institute of Health Policy, Management and Evaluation, Faculty of Medicine, Toronto, ON

⁵Division of Forensic Psychiatry, CAMH, Toronto, ON

Corresponding Author:

Paul Kurdyak

Institute for Mental Health Policy Research

CAMH

33 Russell Street, T305

Toronto, ON

M5S 2S1

Email: paul.kurdyak@camh.ca

Abstract

Objectives: Individuals with schizophrenia are over-represented in correctional facilities relative to their population-based prevalence. The purpose of this study was to determine the rate and predictors of reincarceration of individuals with schizophrenia after release from correctional facilities.

Methods: This was a retrospective cohort study that included all people released from Ontario’s provincial correctional facilities from January 1 to December 31, 2010. Individuals with schizophrenia were identified using a population-based algorithm. The primary outcome was time to reincarceration. Covariates included sociodemographic characteristics (age, sex, neighbourhood income quintile, urban/rural residence), health service utilization (primary care physician visits, psychiatrist visits, psychiatric and non-psychiatric hospitalizations, emergency department visits) and other clinical comorbidity. Survival analysis was used to examine the association between schizophrenia and re-incarceration.

Results: Among 46,928 individuals, N = 3237 (7%) had a diagnosis of schizophrenia. Approximately 67.5% of these individuals were reincarcerated within five years following their first release in 2010, compared to 58.8% of individuals without schizophrenia. Individuals with schizophrenia were 40% (HR: 1.39, 95% CI: 1.33-1.45) more likely to be reincarcerated following release than the control group after adjusting for demographic characteristics. This association reduced to 8% (1.08, 95% CI:1.03-1.14) after adjusting for prior health service utilization, prior correctional involvement, and comorbidities.

Conclusion: Individuals with schizophrenia were more likely to experience reincarceration after release from correctional facilities. This risk is partly explained by prior correctional involvement, health service utilization and comorbidities. Future research should focus on risk factors predicting the higher reincarceration rate and interventions to reduce correctional involvement.

Keywords: Schizophrenia, Correctional Involvement, Incarceration, Health Services

Introduction

Individuals with schizophrenia are at higher risk of incarceration compared to individuals without serious mental illness(1, 2). This higher rate of incarceration has translated into a schizophrenia prevalence in correctional facilities that is 3-6 times the population prevalence(3). Estimates from Canadian studies similarly suggest a higher prevalence of schizophrenia and other psychotic disorders in custodial populations(4, 5).

The higher rate of incarceration amongst individuals with schizophrenia may be due in part to higher rates of recidivism and potential re-incarceration, or effects of criminal justice interventions. For instance, reoffending after correctional facility release is increased by 40% amongst individuals with severe mental illnesses(3) compared to those without. In particular, a significant factor for higher risk of reincarceration for individuals with schizophrenia in contrast to those without is technical violation. Technical violations arise from criminal justice sanctions (court appearances, probation and parole requirement). Individuals with schizophrenia may have challenges adhering to probation and parole requirements resulting in a greater likelihood of getting new charges for these violations. The high rate of incarceration and reincarceration amongst individuals with schizophrenia may also be partially due to poor access to treatment(6), and evidence suggests that timely access to care among individuals with schizophrenia who have been released from correctional facilities can reduce the likelihood of reincarceration(7, 8).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The objective of this study was to determine the prevalence of schizophrenia in persons released from Ontario correctional centres in 2010 and to estimate reincarceration rates among these individuals ~~with~~when compared to those without a diagnosis of schizophrenia. We used comprehensive population-based correctional and health administrative data for all Ontario residents who were released from provincial correctional facilities in 2010.

Methods

Study Setting and Design

The Ontario provincial correctional system is responsible for the detention of all persons on remand, and sentenced inmates subject to sentences of less than 2 years. Sentenced people have an average length of stay of approximately 60 days, and people on remand of approximately 40 days, although the median remand length of stay is closer to one week.

Individuals who were released from Ontario’s provincial correctional facilities from January 1, 2010 to December 31, 2010 served as the study population, based on available data(9) (10).

Exclusion criteria included individuals whose age was less than 18 and greater than 105 at the time of the first correctional facility release data in 2010 (the index event); individuals whose data could not be linked to a valid identifier in the administrative data(10), individuals missing information on their place of residence, and individuals who were not residents of Ontario for the three years prior to the admission date of the index incarceration – as health service utilization data would not be available for these individuals.

Data Sources

The Ministry of Community Safety and Correctional Services dataset provided data for individuals within this cohort from 2005 to 2015. ~~Given that the~~ The index release was the first release in 2010, and the maximum period of time for a provincial correctional event is 2 years. Since the data are retrospective to 2005, a three-year lookback is the maximum amount of time for an individual with the longest detention (2 years) given the correctional data goes back to 2005. ~~Therefore,~~ three years prior to correctional centre was the lookback duration that ensured the same correctional and health service utilization observation period for each subject regardless the duration of index correctional event duration. Subjects were followed from the index release in 2010 to a maximum follow up date of December 31, 2015 to capture correctional involvement and health service utilization.

The Ministry of Community Safety and Correctional Services provided socio-demographic data including age, sex, self-reported race, address on correctional entry (data on income quintile and rurality of residence were accessed at ICES, Toronto-Ontario's largest health data repository, using the address provided), and dates of entry and release from provincial custody from 2005 to 2015. These data were linked with health administrative data contained within ICES using valid ICES Key Numbers (IKN), which are unique person identifiers that are encoded Ontario Health Insurance Plan (OHIP) numbers. The linkage rate was approximately 97%.

Data related to health service utilization were obtained using administrative health information datasets including: the Canadian Institute for Health Information Discharge Abstract Database

(CIHI DAD), the Ontario Mental Health Reporting System (OMHRS), the National Ambulatory Care Reporting System (NACRS), and Ontario Health Insurance Plan Claims Database (OHIP). These health databases capture and report administrative, clinical, and demographic data such as hospital admission and discharge, length of stay, emergency department registrations, and ambulatory encounters. The Registered Persons Database (RPDB) was used to obtain population and demographic data. The RPDB is a population-based registry maintained by the Ministry of Health and Long-Term Care in Ontario. The dataset has demographic information on individuals that use the health system including date of birth, sex, address, date of death, and OHIP eligibility and status changes. All these data sources are held at ICES. ICES is an independent, non-profit research institute whose legal status under Ontario’s health information privacy law allows it to collect and analyze health care and demographic data, without consent, for health system evaluation and improvement. These datasets were linked using unique encoded identifiers and analyzed at ICES. The use of data in this project was authorized under section 45 of Ontario’s Personal Health Information Protection Act, which does not require review by a Research Ethics Board.

Exposure Variable

The provincial correctional sample was stratified based on the presence of a diagnosis of schizophrenia prior to the index event. Persons with a schizophrenia diagnosis were identified using a validated non-specific algorithm using OHIP billing codes, emergency department visits and hospitalizations⁽¹¹⁾. Psychosis not otherwise specified (NOS) is often indistinguishable from

1
2
3 schizophrenia and therefore we included this diagnosis. The definition was ≥ 1 hospitalization
4 (DAD/OMHRS) or ≥ 3 OHIP claims in a three-year period with DSM-IV/ICD9 dx 295, 298, ICD10
5
6 F20, F25, F29⁽¹¹⁾.
7
8
9

10 11 **Outcomes**

12
13
14
15 The primary outcome of this study was re-incarceration, measured as time to first readmission
16
17 to a provincial correctional facility after the date of first release in 2010. Secondary outcomes of
18
19 this study were measures of correctional events and health service utilization. Outcomes
20
21 included: the number of admissions to provincial correctional facilities within five years of the
22
23 index event, the total number of days in custody in provincial correctional facilities within five
24
25 years of the index event, and the time to first contact for each of the following health services
26
27 after release: primary care physician (PCP) contact (including total, mental health-related and
28
29 non-mental health-related contact), psychiatrist contact, emergency department visit (for all
30
31 causes and mental health reason), medical hospitalization and psychiatric hospitalization.
32
33
34
35
36
37
38
39
40

41 **Covariates**

42
43
44 Sociodemographic characteristics included age, sex, neighbourhood income quintile, and
45
46 urban/rural residence. Age was divided into six categories (18-29, 30-39, 40-49, 50-64, 65-74
47
48 and 75+ years). Neighbourhood income levels were obtained using Census Canada information
49
50 provided by Statistics Canada (May 2006). Neighbourhood income levels were divided into five
51
52 categories based on distribution, in which income level 1 was defined as the lowest income
53
54
55
56
57
58
59
60

quintile and level 5 as the highest based on dissemination areas⁽¹²⁾. Rurality was defined using the Rurality Index of Ontario (RIO2008) score, areas that scored 0-39 were considered urban and areas that scored 40 or more were considered rural⁽¹³⁾. Self-reported race and variables describing provincial correctional involvement were obtained from the dataset provided by the Ministry of Community Safety and Correctional Services. These categorical variables included: the cumulative length of time in custody from all correctional admissions in three years prior to the index incarceration, length of the index incarceration event, and the number of prior correctional events in three years prior to the index incarceration.

PCP visits were defined as the total number of visits for all reasons for the periods mentioned above. Visits to PCPs were further categorized as mental health related and non-mental health related visits based on a validated algorithm using OHIP billing codes⁽¹⁴⁾. The total number of psychiatry visits were also identified in OHIP billing codes. To differentiate between PCP and psychiatrist visits, the ICES Physician Database (IPDP) was used to identify physician specialties for accuracy. Total hospitalizations and psychiatric hospitalizations were identified in CIHI DAD and OMHRS. The total number of emergency department visits was captured using NACRS. Clinical characteristics of the sample were captured using the ACG® System Collapsed Adjusted Diagnosis Groups (CADGs) from the John Hopkins ACG® System Version 10, which has 12 categories⁽¹⁵⁾. CADGs were used as a proxy measure of morbidity and have shown good prediction of mortality in the general population⁽¹⁶⁾ and in a population of individuals with a diagnosis of schizophrenia⁽¹⁵⁾.

Statistical Analysis

1
2
3 Health service utilization encounters were calculated for each participant in the three year
4
5 period before the entry date of the index incarceration and in the five year period after the
6
7 index release – defined as the first release in 2010. Healthcare utilization was also measured
8
9 during the index incarceration. We examined episodes in custody in the three years prior to the
10
11 entry date of the index incarceration and five years after the index event.
12
13
14
15

16 The descriptive data were summarized using frequencies and proportions for categorical data,
17
18 and defined based on the data distribution. The distribution of continuous variables was
19
20 measured using mean (standard deviation) and median (interquartile range) to measure the
21
22 variations. Cox proportional hazards model was selected as the most appropriate modelling
23
24 method to reflect our interest in the time to event, and to be able to include other covariates in
25
26 the time to event model. Violation of the Cox PH proportionality assumption was tested using
27
28 Schoenfeld residuals and time-dependent covariates in the analysis. All analyses were
29
30 performed in SAS version 9.4 (SAS Institute, Cary, NC)(17).
31
32
33
34
35
36
37
38
39
40

41 Results

42
43 In 2010, there were 51,013 individuals released from Ontario's provincial correctional facilities.
44
45 After excluding individuals who were missing information location on residence (N = 3448),
46
47 missing age variable (N = 5), younger than 18 at the index event (N = 17) or not Ontario
48
49 residents in the three years prior to the admission date of individuals' first incarceration in 2010
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

(N=615), the cohort included 46,928 individuals. Among the 46,928 individuals, 7% (N = 3237) had a diagnosis of schizophrenia prior to the index event date.

The characteristics of individuals released from correctional facilities with and without schizophrenia are outlined in Table 1. Individuals with schizophrenia were older, more likely to be female, more likely to live in a neighbourhood in the lowest neighbourhood income quintile, and less likely to reside in rural regions.

Correctional involvement in the three years prior to the index correctional entry date is outlined in Table 2. Individuals with schizophrenia had almost double the median aggregate length of correctional centre stay compared to individuals without schizophrenia (66 (IQR – 13-176) vs. 38 (IQR – 7-123)). The median length of stay of the index event and the number of prior incarcerations was also higher for individuals with schizophrenia. The prior health service utilization is outlined in Supplementary Table 1, and shows higher rates of use of all prior health services measured.

Correctional involvement following index correctional event release date is outlined in Table 3. Slightly more individuals with schizophrenia had at least one incarceration event in the five years following release than individuals without schizophrenia (67.5% vs. 58.8%). The median time to re-incarceration was shorter for individuals with vs. without schizophrenia (156 days

(IQR 52-410) vs. 219 days (IQR 81-531)) although total length of time in custody did not differ between the groups.

Supplementary Table 2 shows health service utilization in the five years following release from correctional centres. As with health service utilization prior to correctional centre entry, individuals with schizophrenia had higher rates of all health service utilization in comparison to individuals without schizophrenia. Twenty-four percent of people with schizophrenia (N = 776; 24.0%) did not see a psychiatrist within 5 years following correctional centre release, and the median time post-release for a person with schizophrenia to see a psychiatrist was 114 days (IQR: 30-374).

In a multivariable Cox regression model adjusting for age, sex, neighbourhood income quintile and rurality, individuals with schizophrenia had a 40% increased risk of reincarceration (adjusted HR: 1.39; 95% CI 1.33-1.45)(Fig. 1A) compared to those without schizophrenia. Further adjusting for prior correctional involvement, comorbidity, and prior health service utilization, individuals with schizophrenia were only at 8% increased risk of reincarceration compared to individuals without schizophrenia (adjusted HR: 1.08; 95% CI 1.03-1.14)(Fig. 1B).

Discussion

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Approximately 1 in 14 individuals (7%) released from provincial correctional facilities in 2010 had a diagnosis of schizophrenia. People with schizophrenia had a higher number of correctional encounters prior to and following release than individuals without schizophrenia, and were 40% more likely to be reincarcerated within 5 years following correctional release than individuals without schizophrenia. That this relative increase in re-incarceration was substantially attenuated by prior health service utilization, correctional involvement and comorbidity suggests that these factors may play an important role in re-incarceration. That prior correctional involvement is an important contributor is not surprising; however, prior health service utilization may point to opportunities to intervene if individuals with better pre-incarceration access to care have reduced rates of reincarceration.

We found that individuals with schizophrenia are highly overrepresented in Ontario’s provincial correctional facilities compared to the general population, which is consistent with prior research (3, 18-22). The schizophrenia prevalence within this cohort was notably high (7%), compared with estimates of schizophrenia prevalence observed in prior studies(3) and in the general population (approximately 1%)(23, 24). Our findings that a diagnosis of schizophrenia is associated with a higher rate of re-incarceration is also consistent with existing literature(13, 14, 25, 26).

The findings also highlight service fragmentation following the index release in 2010. In the five years following release, approximately 1 in 4 individuals with schizophrenia did not see a psychiatrist. Given the complexities associated with support for individuals who have a

1
2
3 diagnosis of schizophrenia such as the continuation of antipsychotic medication, the need for
4
5 other psychosocial interventions such as cognitive behavioural therapy, the need for regular
6
7 physical health assessments, stable housing, among other socio-economic supports(27), timely
8
9 access to specialty mental health care services, including psychiatrists, quickly following release
10
11 should be an expected standard of care for individuals with schizophrenia who have had
12
13 correctional involvement. Indeed, 44% of individuals with schizophrenia had at least one
14
15 psychiatric hospitalization following index correctional event release. Lack of psychiatric care
16
17 and treatment may contribute to both re-incarceration and psychiatric hospitalizations. Indeed,
18
19 the same factors that create challenges adhering to conditions of release (parole, etc.) likely
20
21 contribute to challenges attending psychiatric follow-up.
22
23
24
25
26
27
28
29

30 There are several limitations that merit discussion. Given the limited availability of socio-
31
32 demographic data, we were unable to include largely cited socio-demographic factors
33
34 including: marital status, education level and housing status(8, 28-32). Our diagnosis of
35
36 schizophrenia relies on clinical codes in administrative health data rather than a comprehensive
37
38 clinical assessment, and there may be misclassification of schizophrenia status. However, the
39
40 algorithm used to identify individuals with a primary diagnosis of schizophrenia was validated
41
42 using population-based health administrative databases in Ontario(11). Finally, the correctional
43
44 data did not include information on the criminal charges or convictions. Distinguishing whether
45
46 individuals with schizophrenia within this cohort were incarcerated for violent or non-violent
47
48 crimes or for administrative offences such as failing to comply with probation/parole or to
49
50 attend court appearances could have contributed to the analysis in understanding the length of
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

correctional episodes and whether individuals were receiving the type of care necessary. In addition, this study focused on Ontario’s provincial correctional facilities and did not include data on federal correctional involvement.

Conclusion

Individuals with schizophrenia are overrepresented in correctional facilities in comparison to the general population. Additionally, individuals with schizophrenia also experience higher rates of re-incarceration. Further work is needed to understand the needs of individuals with schizophrenia and risk factors contributing to re-incarceration. However, these individuals would likely benefit from more coordinated and integrated health service delivery following correctional release.

Data Access

The dataset from this study is held securely in coded form at ICES. While data sharing agreements prohibit ICES from making the dataset publicly available, access may be granted to those who meet pre-specified criteria for confidential access, available at www.ices.on.ca/DAS. The full dataset creation plan and underlying analytic code are available from the authors upon request, understanding that the computer programs may rely upon coding templates or macros that are unique to ICES and are therefore either inaccessible or may require modification.

Acknowledgements

This study was supported by ICES, which is funded by an annual grant from the Ontario Ministry of Health and Long-Term Care (MOHLTC). Parts of this material are based on data and information compiled and provided by: the Ontario Ministry of Health and Long-term Care and the Canadian Institute for Health Information (CIHI). The analyses, conclusions, opinions and statements expressed herein are solely those of the authors and do not reflect those of the funding or data sources; no endorsement is intended or should be inferred.

Conflicts of Interest

No author has a conflict of interest related to this study.

Funding

This study was funded by a grant from the Canadian Institutes for Health Research.

References

1. Ramsay CE, Goulding, S. M., Broussard, B., Cristofaro, S. L., Abedi, G. R., & Compton, M. T. Prevalence and psychosocial correlates of prior incarcerations in an urban, predominantly African-American sample of hospitalized patients with first-episode psychosis. *J Am Acad Psychiatry Law*. 2011;39(1):57-64.

2. Rautanen M, & Lauerma, H. Imprisonment and diagnostic delay among male offenders with schizophrenia. *Crim Behav Ment Health*. 2011;21(4):259-64.

3. Fazel S, & Seewald, K. Severe mental illness in 33,588 prisoners worldwide: systematic review and meta-regression analysis. *Br J Psychiatry*. 2012;200(5):364-73.

4. Bland RC, Newman, S. C., Thompson, A. H., & Dyck, R. J. Psychiatric disorders in the population and in prisoners. *Int J Law Psychiatry*. 1998;21(3):273-9.

5. Corrado RR, Cohen, I., Hart, S., & Roesch, R. Comparative examination of the prevalence of mental disorders among jailed inmates in Canada and the United States. *Int J Law Psychiatry*. 2000;23(5-6):633-47.

6. Fisher WH, Packer, I. K., Simon, L. J., & Smith, D. Community mental health services and the prevalence of severe mental illness in local jails: are they related? *Adm Policy Ment Health*. 2000;27(6):371-82.

7. Hawthorne WB, Folsom, D. P., Sommerfeld, D. H., Lanouette, N. M., Lewis, M., Aarons, G. A., Conklin, R. M., Solorzano, E., Lindamer, L. A., & Jeste DV. Incarceration Among Adults Who Are in the Public Mental Health System: Rates, Risk Factors, and Short-Term Outcomes. *Psychiatric Services*. 2012;63(1):26-32.

- 1
2
3 8. Lamberti JS. Understanding and preventing criminal recidivism among adults with
4
5 psychotic disorders. *Psychiatr Serv.* 2007;58(6):773-81.
6
7
- 8 9. Kouyoumdjian F, Schuler, A., Hwang, S. W. & Matheson, F. Research on the health of
9
10 people who experience detention or incarceration in Canada: a scoping review. *BMC public*
11
12 *health.* 2015;15(419):1-8.
13
14
- 15 10. Kouyoumdjian FG, Cheng S. Y., Fung, K., Orkin, A. M., McIsaac K. E., Kendall, C., Kiefer, L.,
16
17 Matheson, F. I., Green, S. E., and Hwang, S. W. . The health care utilization of people in prison
18
19 and after prison release: A population-based cohort study in Ontario, Canada. *PloS one.* August
20
21 3, 2018;13(8).
22
23
- 24 11. Kurdyak P, Lin, E., Green, D., & Vigod, S. Validation of a Population-Based Algorithm to
25
26 Detect Chronic Psychotic Illness. *Can J Psychiatry.* 2015;60(8):362-8.
27
28
- 29 12. Statistics Canada. Income in Canada 2015 [Available from:
30
31 <https://www.statcan.gc.ca/pub/75-202-x/2010000/analysis-analyses-eng.htm>.
32
33
- 34 13. Krajl B. Measuring Rurality - RIO2008 BASIC: Methodology and Results. Toronto (ON):
35
36 OMA Economics Department; 2009.
37
38
- 39 14. Steele L, Glazier R, Lin E, Evans M. Using administrative data to measure ambulatory
40
41 mental health service provision in primary care. *Med Care.* 2004;42(10):960-5.
42
43
- 44 15. Austin PC, Newman A, Kurdyak PA. Using the Johns Hopkins Aggregated Diagnosis
45
46 Groups (ADGs) to predict mortality in a population-based cohort of adults with schizophrenia in
47
48 Ontario, Canada. *Psychiatry Res.* 2012;196(1):32-7.
49
50
51
52
53
54
55
56
57
58
59
60

16. Austin PC, Walraven, C., Wodchis, W., Newman, A. & Anderson, G. M. Using the Johns Hopkins Aggregated Diagnosis Groups (ADGs) to predict mortality in a general adult population cohort in Ontario, Canada. Canadian Institutes of Health Research. 2011;40(10):932-9.

17. SAS. SAS System, Version 9: Installation and Administration Information. 2007.

18. Brinded PM, Simpson, A. I., Laidlaw, T. M., Fairley, N., & Malcolm, F. Prevalence of psychiatric disorders in New Zealand prisons: a national study. Aust N Z J Psychiatry. 2001;35(2):166-73.

19. Sirdifield C, Gojkovic, D., Brooker, C., & Ferriter, M. A systematic review of research on the epidemiology of mental health disorders in prison populations: a summary of findings. J Forensic Psychi Ps. 2009;20(1):S78-S101.

20. Lafortune D. Prevalence and screening of mental disorders in short-term correctional facilities. Int J Law Psychiatry. 2010;33(2):94-100.

21. Senior J, Birmingham, L., Harty, M. A., Hassan, L., Hayes, A. J., Kendall, K., King, C., Lathlean, J., Lowthian, C., Mills, A., Webb, R., Thornicroft, G., & Shaw, J. Identification and management of prisoners with severe psychiatric illness by specialist mental health services. Psychol Med. 2013;43(7):1511-20.

22. Fazel S, & Danesh, J. Serious mental disorder in 23000 prisoners: a systematic review of 62 surveys. Lancet. 2002;359(9306):545-50.

23. Goldner EM, Jones, W., & Waraich, P. Using administrative data to analyze the prevalence and distribution of schizophrenic disorders. Psychiatric Services. 2003;54(7):1017-21.

24. Khan S, Chiu, M., Simpson, A. I., Guttman, A., Jembere, N., & Kurdyak, P. Use of Mental Health Services by Youths and Young Adults Before and During Correctional Custody: A Population-Based Study. *Psychiatr Serv.* 2016;67(7):790-3.
25. Simpson AI, McMaster, J. J., & Cohen, S. N. Challenges for Canada in meeting the needs of persons with serious mental illness in prison. *J Am Acad Psychiatry Law.* 2013;41(4):501-9.
26. National Institute of Health and Care Excellence. Psychosis and schizophrenia in adults. 2015.
27. Health Quality Ontario. Schizophrenia Care in the Community for Adults Ontario: Queen's Printer for Ontario; 2019 [Available from: <https://www.hqontario.ca/Evidence-to-Improve-Care/Quality-Standards/View-all-Quality-Standards/Schizophrenia-Care-in-the-Community/The-Quality-Standard-In-Brief>].
28. Prince JD. Incarceration and hospital care. *J Nerv Ment Dis.* 2006;194(1):34-9.
29. Harty M, Jarrett, M., Thornicroft, G., & Shaw, J. Unmet needs of male prisoners under the care of prison Mental Health Inreach Services. *J Forensic Psychi Ps.* 2012;23(3):285-96.
30. Prince JD, Akincigil, A., & Bromet, E. Incarceration rates of persons with first-admission psychosis. *Psychiatr Serv.* 2007;58(9):1173-80.
31. Robertson AG, Swanson, J. W., Frisman, L. K., Lin, H., & Swartz, M. S. Patterns of justice involvement among adults with schizophrenia and bipolar disorder: key risk factors. *Psychiatr Serv.* 2014;65(7):931-8.
32. Wallace C, Mullen, P. E., & Burgess, P. Criminal offending in schizophrenia over a 25-year period marked by deinstitutionalization and increasing prevalence of comorbid substance use disorders. *Am J Psychiatry.* 2004;161(4):716-27.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Characteristics of study cohort, by schizophrenia diagnosis

Variable	Individuals without schizophrenia	Individuals with schizophrenia	Standardized Difference
	N (%)	N (%)	
Total	43,691 (93.1)	3,237 (7.0)	
Age			
Median (IQR)	32 (24-43)	36 (28-45)	0.30
18-29	19,007 (43.5)	1,004 (31.0)	0.26
30-39	10,775 (24.7)	936 (28.9)	0.10
40-49	9,175 (21.0)	825 (25.5)	0.11
50-64	4,322 (9.9)	425 (13.1)	0.10
65-74	354 (0.8)	39 (1.2)	0.04
75+	58 (0.1)	8 (0.2)	0.03
Sex			
Female	5,280 (12.1)	542 (16.7)	0.13
Male	38,411 (87.9)	2,695 (83.3)	0.13
Self-reported race			
Missing	2,952 (6.8)	223 (6.9)	0.01
Aboriginal	4,763 (10.9)	271 (8.4)	0.09
Black	5,248 (12.0)	447 (13.8)	0.05
East Asian	715 (1.6)	43 (1.3)	0.03
Hispanic	519 (1.2)	26 (0.8)	0.04
South Asian	908 (2.1)	68 (2.1)	0.00
South East Asian	557 (1.3)	46 (1.4)	0.01
West Asian/Arabic	671 (1.5)	46 (1.4)	0.01
White	25,973 (59.4)	1,958 (60.5)	0.02
Other racial origin	1,000 (2.3)	79 (2.4)	0.01
Declined to specify	130 (0.3)	12 (0.4)	0.01
Racial origin unknown	254 (0.6)	17 (0.5)	0.01
Income Quintile			
Missing	3,207 (7.3)	96 (3.0)	0.20
1	15,359 (35.2)	1,334 (41.2)	0.12
2	9,218 (21.1)	696 (21.5)	0.01
3	6,802 (15.6)	508 (15.7)	0.00
4	5,194 (11.9)	330 (10.2)	0.05
5	3,911 (9.0)	273 (8.4)	0.02
Rurality			
Missing	2,398 (5.5)	25 (0.8)	0.27
Urban	35,533 (81.3)	2,953 (91.2)	0.29
Rural	5,760 (13.2)	259 (8.0)	0.17
Comorbidity (cADGs)			

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Mean ± SD	3.0±2.1	4.2±2.0	0.60
Median (IQR)	3(1-4)	4(3-6)	0.61

Legend: IQR – interquartile range; cADGs – Collapsed Aggregated Diagnostic groups; SD – standard deviation

Table 2: Correctional involvement three years prior to the index* incarceration by schizophrenia diagnosis

Variable	Individuals without schizophrenia	Individuals with schizophrenia	Standardized Difference
	N (%)	N (%)	
Total	43,691 (93.1)	3,237 (7.0)	
Length of stay from all correctional entries in 3 years prior to index release			
Median (IQR)	38 (7-123)	66 (13-176)	0.26
No prior incarcerations	22,904 (52.4)	1,327 (41.0)	0.23
1-6	5,115 (11.7)	323 (10.0)	0.06
7-29	4,464 (10.2)	354 (10.9)	0.02
30-89	4,385 (10.0)	430 (13.3)	0.10
90-179	3,414 (7.8)	335 (10.3)	0.09
180-364	2,570 (5.9)	332 (10.3)	0.16
365+	839 (1.9)	136 (4.2)	0.13
Length of stay of index incarceration event			
Median (IQR)	7 (2-37)	11 (3-48)	0.16
1-6	21,101 (48.3)	1,324 (40.9)	0.15
7-29	10,104 (23.1)	840 (25.9)	0.07
30-89	7,434 (17.0)	638 (19.7)	0.07
90-119	1,813 (4.1)	140 (4.3)	0.01
120-364	2,966 (6.8)	273 (8.4)	0.06
365+	273 (0.6)	22 (0.7)	0.01
Number of prior incarcerations			
Median (IQR)	0 (0-2)	1 (0-3)	0.3
0	22,904 (52.4)	1,327 (41.0)	0.23
1	8,151 (18.7)	566 (17.5)	0.03
2	4,868 (11.1)	357 (11.0)	0.00
3	2,929 (6.7)	266 (8.2)	0.06
4	1,846 (4.2)	207 (6.4)	0.10
5	1,071 (2.5)	139 (4.3)	0.10
6	661 (1.5)	84 (2.6)	0.08
7	450 (1.0)	68 (2.1)	0.09
8	259 (0.6)	57 (1.8)	0.11
9	175 (0.4)	46 (1.4)	0.11
10+	377 (0.9)	120 (3.7)	0.19

*Index refers to the first correctional release for each individual in 2010 in Ontario.

Legend: IQR – interquartile range


Table 3: Correctional involvement five years after index release, by schizophrenia diagnosis

Variable	Individuals without schizophrenia	Individuals with schizophrenia	Standardized Difference
	N (%)	N (%)	
Total	43,691 (93.1)	3,237 (7.0)	
Any re-incarceration within 5 years	25,703 (58.8)	2,185 (67.5)	0.18
Time to re-incarceration			
Median (IQR)	219 (81-531)	156 (52-410)	0.21
Not reincarcerated	17,988 (41.2)	1,052 (32.5)	0.18
1-30 days	2,853 (6.5)	359 (11.1)	0.16
31-60	2,249 (5.1)	259 (8.0)	0.12
61-90	1,908 (4.4)	184 (5.7)	0.06
91-240	6,503 (14.9)	528 (16.3)	0.04
241-365	3,143 (7.2)	240 (7.4)	0.01
366-730	4,554 (10.4)	330 (10.2)	0.01
731-1095	2,079 (4.8)	96 (3.0)	0.09
3 years +	2,414 (5.5)	189 (5.8)	0.01
Length of stay from all correctional events in 5 years after index release			
Median (IQR)	83 (20-204)	88 (23-222)	0.07
Not re-incarcerated	17,988 (41.2)	1,052 (32.5)	0.18
1-6 days	3,563 (8.2)	246 (7.6)	0.02
7-29	4,181 (9.6)	373 (11.5)	0.06
30-89	5,597 (12.8)	483 (14.9)	0.06
90-179	4,999 (11.4)	386 (11.9)	0.02
180-364	4,501 (10.3)	402 (12.4)	0.07
365-729	2,473 (5.7)	232 (7.2)	0.06
730+	389 (0.9)	63 (1.9)	0.09

*Index refers to the first correctional release for each individual in 2010 in Ontario.


Legend: IQR – interquartile range

Figure 1A: Time to re-incarceration for individuals with and without schizophrenia from multivariable Cox regression*



*Adjusted for age, sex, neighbourhood income quintile and rurality.

Figure 1B: Recidivism survival curves between individuals with and without schizophrenia.*



*Adjusted for age, sex, neighbourhood income quintile, rurality, comorbidity, prior correctional involvement and prior health service utilization.

Prevalence and predictors of re-incarceration after correctional centre release: a population-based comparison of individuals with and without schizophrenia in Ontario, Canada.

Mayuri Mahentharan MSc¹

Fiona G. Kouyoumdjian MD MPH PhD^{2,3}

Claire de Oliveira PhD^{1,3,4}

Tomisin Iwajomo MPH^{1,3}

Alexander I F Simpson MBChB BMedSci⁵

Roland Jones MD PhD⁵

Paul Kurdyak MD PhD^{1,3,4}

Institutional Affiliations

¹Institute for Mental Health Policy Research, CAMH, Toronto, ON

²Department of Family Medicine, McMaster University, Hamilton, ON

³ICES, Toronto, ON

⁴Institute of Health Policy, Management and Evaluation, Faculty of Medicine, Toronto, ON

⁵Division of Forensic Psychiatry, CAMH, Toronto, ON

Corresponding Author:

Paul Kurdyak

Institute for Mental Health Policy Research

CAMH

33 Russell Street, T305

Toronto, ON

M5S 2S1

Email: paul.kurdyak@camh.ca

Abstract

Objectives: Individuals with schizophrenia are over-represented in correctional facilities relative to their population-based prevalence. The purpose of this study was to determine the rate and predictors of reincarceration of individuals with schizophrenia after release from correctional facilities.

Methods: This was a retrospective cohort study that included all people released from Ontario’s provincial correctional facilities from January 1 to December 31, 2010. Individuals with schizophrenia were identified using a population-based algorithm. The primary outcome was time to reincarceration. Covariates included sociodemographic characteristics (age, sex, neighbourhood income quintile, urban/rural residence), health service utilization (primary care physician visits, psychiatrist visits, psychiatric and non-psychiatric hospitalizations, emergency department visits) and other clinical comorbidity. Survival analysis was used to examine the association between schizophrenia and re-incarceration.

Results: Among 46,928 individuals, N = 3237 (7%) had a diagnosis of schizophrenia. Approximately 67.5% of these individuals were reincarcerated within five years following their first release in 2010, compared to 58.8% of individuals without schizophrenia. Individuals with schizophrenia were 40% (HR: 1.39, 95% CI: 1.33-1.45) more likely to be reincarcerated following release than the control group after adjusting for demographic characteristics. This association reduced to 8% (1.08, 95% CI:1.03-1.14) after adjusting for prior health service utilization, prior correctional involvement, and comorbidities.

Conclusion: Individuals with schizophrenia were more likely to experience reincarceration after release from correctional facilities. This risk is partly explained by prior correctional involvement, health service utilization and comorbidities. Future research should focus on risk factors predicting the higher reincarceration rate and interventions to reduce correctional involvement.

Keywords: Schizophrenia, Correctional Involvement, Incarceration, Health Services

Introduction

Individuals with schizophrenia are at higher risk of incarceration compared to individuals without serious mental illness(1, 2). This higher rate of incarceration has translated into a schizophrenia prevalence in correctional facilities that is 3-6 times the population prevalence(3). Estimates from Canadian studies similarly suggest a higher prevalence of schizophrenia and other psychotic disorders in custodial populations(4, 5).

The higher rate of incarceration amongst individuals with schizophrenia may be due in part to higher rates of recidivism and potential re-incarceration, or effects of criminal justice interventions. For instance, reoffending after correctional facility release is increased by 40% amongst individuals with severe mental illnesses(3) compared to those without. In particular, a significant factor for higher risk of reincarceration for individuals with schizophrenia in contrast to those without is technical violation. Technical violations arise from criminal justice sanctions (court appearances, probation and parole requirement). Individuals with schizophrenia may have challenges adhering to probation and parole requirements resulting in a greater likelihood of getting new charges for these violations. The high rate of incarceration and reincarceration amongst individuals with schizophrenia may also be partially due to poor access to treatment(6), and evidence suggests that timely access to care among individuals with schizophrenia who have been released from correctional facilities can reduce the likelihood of reincarceration(7, 8).

The objective of this study was to determine the prevalence of schizophrenia in persons released from Ontario correctional centres in 2010 and to estimate reincarceration rates among these individuals with compared to those without a diagnosis of schizophrenia. We used comprehensive population-based correctional and health administrative data for all Ontario residents who were released from provincial correctional facilities in 2010.

Methods

Study Setting and Design

The Ontario provincial correctional system is responsible for the detention of all persons on remand, and sentenced inmates subject to sentences of less than 2 years. Sentenced people have an average length of stay of approximately 60 days, and people on remand of approximately 40 days, although the median remand length of stay is closer to one week.

Individuals who were released from Ontario’s provincial correctional facilities from January 1, 2010 to December 31, 2010 served as the study population, based on available data(9) (10).

Exclusion criteria included individuals whose age was less than 18 and greater than 105 at the time of the first correctional facility release data in 2010 (the index event); individuals whose data could not be linked to a valid identifier in the administrative data(10), individuals missing information on their place of residence, and individuals who were not residents of Ontario for the three years prior to the admission date of the index incarceration – as health service utilization data would not be available for these individuals.

Data Sources

The Ministry of Community Safety and Correctional Services dataset provided data for individuals within this cohort from 2005 to 2015. Given that the index release was the first release in 2010, a three year lookback is the maximum amount of time for an individual with the longest detention (2 years) given the correctional data goes back to 2005. Therefore, three years prior to correctional centre was the lookback duration that ensured the same correctional and health service utilization observation period for each subject regardless the duration of index correctional event duration. Subjects were followed from the index release in 2010 to a maximum follow up date of December 31, 2015 to capture correctional involvement and health service utilization.

The Ministry of Community Safety and Correctional Services provided socio-demographic data including age, sex, self-reported race, address on correctional entry (data on income quintile and rurality of residence were accessed at ICES, Toronto-Ontario's largest health data repository, using the address provided), and dates of entry and release from provincial custody from 2005 to 2015. These data were linked with health administrative data contained within ICES using valid ICES Key Numbers (IKN), which are unique person identifiers that are encoded Ontario Health Insurance Plan (OHIP) numbers. The linkage rate was approximately 97%.

Data related to health service utilization were obtained using administrative health information datasets including: the Canadian Institute for Health Information Discharge Abstract Database (CIHI DAD), the Ontario Mental Health Reporting System (OMHRS), the National Ambulatory

Care Reporting System (NACRS), and Ontario Health Insurance Plan Claims Database (OHIP). These health databases capture and report administrative, clinical, and demographic data such as hospital admission and discharge, length of stay, emergency department registrations, and ambulatory encounters. The Registered Persons Database (RPDB) was used to obtain population and demographic data. The RPDB is a population-based registry maintained by the Ministry of Health and Long-Term Care in Ontario. The dataset has demographic information on individuals that use the health system including date of birth, sex, address, date of death, and OHIP eligibility and status changes. All these data sources are held at ICES. ICES is an independent, non-profit research institute whose legal status under Ontario’s health information privacy law allows it to collect and analyze health care and demographic data, without consent, for health system evaluation and improvement. These datasets were linked using unique encoded identifiers and analyzed at ICES. The use of data in this project was authorized under section 45 of Ontario’s Personal Health Information Protection Act, which does not require review by a Research Ethics Board.

Exposure Variable

The provincial correctional sample was stratified based on the presence of a diagnosis of schizophrenia prior to the index event. Persons with a schizophrenia diagnosis were identified using a validated non-specific algorithm using OHIP billing codes, emergency department visits and hospitalizations⁽¹¹⁾. Psychosis not otherwise specified (NOS) is often indistinguishable from schizophrenia and therefore we included this diagnosis. The definition was ≥1 hospitalization

(DAD/OMHRS) or ≥ 3 OHIP claims in a three-year period with DSM-IV/ICD9 dx 295, 298, ICD10 F20, F25, F29⁽¹¹⁾.

Outcomes

The primary outcome of this study was re-incarceration, measured as time to first readmission to a provincial correctional facility after the date of first release in 2010. Secondary outcomes of this study were measures of correctional events and health service utilization. Outcomes included: the number of admissions to provincial correctional facilities within five years of the index event, the total number of days in custody in provincial correctional facilities within five years of the index event, and the time to first contact for each of the following health services after release: primary care physician (PCP) contact (including total, mental health-related and non-mental health-related contact), psychiatrist contact, emergency department visit (for all causes and mental health reason), medical hospitalization and psychiatric hospitalization.

Covariates

Sociodemographic characteristics included age, sex, neighbourhood income quintile, and urban/rural residence. Age was divided into six categories (18-29, 30-39, 40-49, 50-64, 65-74 and 75+ years). Neighbourhood income levels were obtained using Census Canada information provided by Statistics Canada (May 2006). Neighbourhood income levels were divided into five categories based on distribution, in which income level 1 was defined as the lowest income quintile and level 5 as the highest based on dissemination areas⁽¹²⁾. Rurality was defined using

the Rurality Index of Ontario (RIO2008) score, areas that scored 0-39 were considered urban and areas that scored 40 or more were considered rural⁽¹³⁾. Self-reported race and variables describing provincial correctional involvement were obtained from the dataset provided by the Ministry of Community Safety and Correctional Services. These categorical variables included: the cumulative length of time in custody from all correctional admissions in three years prior to the index incarceration, length of the index incarceration event, and the number of prior correctional events in three years prior to the index incarceration.

PCP visits were defined as the total number of visits for all reasons for the periods mentioned above. Visits to PCPs were further categorized as mental health related and non-mental health related visits based on a validated algorithm using OHIP billing codes⁽¹⁴⁾. The total number of psychiatry visits were also identified in OHIP billing codes. To differentiate between PCP and psychiatrist visits, the ICES Physician Database (IPDP) was used to identify physician specialties for accuracy. Total hospitalizations and psychiatric hospitalizations were identified in CIHI DAD and OMHRS. The total number of emergency department visits was captured using NACRS. Clinical characteristics of the sample were captured using the ACG® System Collapsed Adjusted Diagnosis Groups (CADGs) from the John Hopkins ACG® System Version 10, which has 12 categories⁽¹⁵⁾. CADGs were used as a proxy measure of morbidity and have shown good prediction of mortality in the general population⁽¹⁶⁾ and in a population of individuals with a diagnosis of schizophrenia⁽¹⁷⁾.

Statistical Analysis

Health service utilization encounters were calculated for each participant in the three year period before the entry date of the index incarceration and in the five year period after the index release – defined as the first release in 2010. Healthcare utilization was also measured during the index incarceration. We examined episodes in custody in the three years prior to the entry date of the index incarceration and five years after the index event.

The descriptive data were summarized using frequencies and proportions for categorical data, and defined based on the data distribution. The distribution of continuous variables was measured using mean (standard deviation) and median (interquartile range) to measure the variations. Cox proportional hazards model was selected as the most appropriate modelling method to reflect our interest in the time to event, and to be able to include other covariates in the time to event model. Violation of the Cox PH proportionality assumption was tested using Schoenfeld residuals and time-dependent covariates in the analysis. All analyses were performed in SAS version 9.4 (SAS Institute, Cary, NC)(18).

Results

In 2010, there were 51,013 individuals released from Ontario's provincial correctional facilities. After excluding individuals who were missing information location on residence (N = 3448), missing age variable (N = 5), younger than 18 at the index event (N = 17) or not Ontario residents in the three years prior to the admission date of individuals' first incarceration in 2010

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

(N=615), the cohort included 46,928 individuals. Among the 46,928 individuals, 7% (N = 3237) had a diagnosis of schizophrenia prior to the index event date.

The characteristics of individuals released from correctional facilities with and without schizophrenia are outlined in Table 1. Individuals with schizophrenia were older, more likely to be female, more likely to live in a neighbourhood in the lowest neighbourhood income quintile, and less likely to reside in rural regions.

Correctional involvement in the three years prior to the index correctional entry date is outlined in Table 2. Individuals with schizophrenia had almost double the median aggregate length of correctional centre stay compared to individuals without schizophrenia (66 (IQR – 13-176) vs. 38 (IQR – 7-123)). The median length of stay of the index event and the number of prior incarcerations was also higher for individuals with schizophrenia. The prior health service utilization is outlined in Supplementary Table 1, and shows higher rates of use of all prior health services measured.

Correctional involvement following index correctional event release date is outlined in Table 3. Slightly more individuals with schizophrenia had at least one incarceration event in the five years following release than individuals without schizophrenia (67.5% vs. 58.8%). The median time to re-incarceration was shorter for individuals with vs. without schizophrenia (156 days

(IQR 52-410) vs. 219 days (IQR 81-531)) although total length of time in custody did not differ between the groups.

Supplementary Table 2 shows health service utilization in the five years following release from correctional centres. As with health service utilization prior to correctional centre entry, individuals with schizophrenia had higher rates of all health service utilization in comparison to individuals without schizophrenia. Twenty-four percent of people with schizophrenia (N = 776; 24.0%) did not see a psychiatrist within 5 years following correctional centre release, and the median time post-release for a person with schizophrenia to see a psychiatrist was 114 days (IQR: 30-374).

In a multivariable Cox regression model adjusting for age, sex, neighbourhood income quintile and rurality, individuals with schizophrenia had a 40% increased risk of reincarceration (adjusted HR: 1.39; 95% CI 1.33-1.45)(Fig. 1A) compared to those without schizophrenia. Further adjusting for prior correctional involvement, comorbidity, and prior health service utilization, individuals with schizophrenia were only at 8% increased risk of reincarceration compared to individuals without schizophrenia (adjusted HR: 1.08; 95% CI 1.03-1.14)(Fig. 1B).

Discussion

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Approximately 1 in 14 individuals (7%) released from provincial correctional facilities in 2010 had a diagnosis of schizophrenia. People with schizophrenia had a higher number of correctional encounters prior to and following release than individuals without schizophrenia, and were 40% more likely to be reincarcerated within 5 years following correctional release than individuals without schizophrenia. That this relative increase in re-incarceration was substantially attenuated by prior health service utilization, correctional involvement and comorbidity suggests that these factors may play an important role in re-incarceration. That prior correctional involvement is an important contributor is not surprising; however, prior health service utilization may point to opportunities to intervene if individuals with better pre-incarceration access to care have reduced rates of reincarceration.

We found that individuals with schizophrenia are highly overrepresented in Ontario’s provincial correctional facilities compared to the general population, which is consistent with prior research (3, 19-23). The schizophrenia prevalence within this cohort was notably high (7%), compared with estimates of schizophrenia prevalence observed in prior studies(3) and in the general population (approximately 1%)(24, 25). Our findings that a diagnosis of schizophrenia is associated with a higher rate of re-incarceration is also consistent with existing literature(13, 14, 26, 27).

The findings also highlight service fragmentation following the index release in 2010. In the five years following release, approximately 1 in 4 individuals with schizophrenia did not see a psychiatrist. Given the complexities associated with support for individuals who have a

1
2
3 diagnosis of schizophrenia such as the continuation of antipsychotic medication, the need for
4
5 other psychosocial interventions such as cognitive behavioural therapy, the need for regular
6
7 physical health assessments, stable housing, among other socio-economic supports(28), timely
8
9 access to specialty mental health care services, including psychiatrists, quickly following release
10
11 should be an expected standard of care for individuals with schizophrenia who have had
12
13 correctional involvement. Indeed, 44% of individuals with schizophrenia had at least one
14
15 psychiatric hospitalization following index correctional event release. Lack of psychiatric care
16
17 and treatment may contribute to both re-incarceration and psychiatric hospitalizations. Indeed,
18
19 the same factors that create challenges adhering to conditions of release (parole, etc.) likely
20
21 contribute to challenges attending psychiatric follow-up.
22
23
24
25
26
27
28
29

30 There are several limitations that merit discussion. Given the limited availability of socio-
31
32 demographic data, we were unable to include largely cited socio-demographic factors
33
34 including: marital status, education level and housing status(8, 29-33). Our diagnosis of
35
36 schizophrenia relies on clinical codes in administrative health data rather than a comprehensive
37
38 clinical assessment, and there may be misclassification of schizophrenia status. However, the
39
40 algorithm used to identify individuals with a primary diagnosis of schizophrenia was validated
41
42 using population-based health administrative databases in Ontario(11). Finally, the correctional
43
44 data did not include information on the criminal charges or convictions. Distinguishing whether
45
46 individuals with schizophrenia within this cohort were incarcerated for violent or non-violent
47
48 crimes or for administrative offences such as failing to comply with probation/parole or to
49
50 attend court appearances could have contributed to the analysis in understanding the length of
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

correctional episodes and whether individuals were receiving the type of care necessary. In addition, this study focused on Ontario’s provincial correctional facilities and did not include data on federal correctional involvement.

Conclusion

Individuals with schizophrenia are overrepresented in correctional facilities in comparison to the general population. Additionally, individuals with schizophrenia also experience higher rates of re-incarceration. Further work is needed to understand the needs of individuals with schizophrenia and risk factors contributing to re-incarceration. However, these individuals would likely benefit from more coordinated and integrated health service delivery following correctional release.

Data Access

The dataset from this study is held securely in coded form at ICES. While data sharing agreements prohibit ICES from making the dataset publicly available, access may be granted to those who meet pre-specified criteria for confidential access, available at www.ices.on.ca/DAS. The full dataset creation plan and underlying analytic code are available from the authors upon request, understanding that the computer programs may rely upon coding templates or macros that are unique to ICES and are therefore either inaccessible or may require modification.

Acknowledgements

This study was supported by ICES, which is funded by an annual grant from the Ontario Ministry of Health and Long-Term Care (MOHLTC). Parts of this material are based on data and information compiled and provided by: the Ontario Ministry of Health and Long-term Care and the Canadian Institute for Health Information (CIHI). The analyses, conclusions, opinions and statements expressed herein are solely those of the authors and do not reflect those of the funding or data sources; no endorsement is intended or should be inferred.

Conflicts of Interest

No author has a conflict of interest related to this study.

Funding

This study was funded by a grant from the Canadian Institutes for Health Research.

References

1. Ramsay CE, Goulding, S. M., Broussard, B., Cristofaro, S. L., Abedi, G. R., & Compton, M. T. Prevalence and psychosocial correlates of prior incarcerations in an urban, predominantly African-American sample of hospitalized patients with first-episode psychosis. *J Am Acad Psychiatry Law*. 2011;39(1):57-64.

2. Rautanen M, & Lauerma, H. Imprisonment and diagnostic delay among male offenders with schizophrenia. *Crim Behav Ment Health*. 2011;21(4):259-64.

3. Fazel S, & Seewald, K. Severe mental illness in 33,588 prisoners worldwide: systematic review and meta-regression analysis. *Br J Psychiatry*. 2012;200(5):364-73.

4. Bland RC, Newman, S. C., Thompson, A. H., & Dyck, R. J. Psychiatric disorders in the population and in prisoners. *Int J Law Psychiatry*. 1998;21(3):273-9.

5. Corrado RR, Cohen, I., Hart, S., & Roesch, R. Comparative examination of the prevalence of mental disorders among jailed inmates in Canada and the United States. *Int J Law Psychiatry*. 2000;23(5-6):633-47.

6. Fisher WH, Packer, I. K., Simon, L. J., & Smith, D. Community mental health services and the prevalence of severe mental illness in local jails: are they related? *Adm Policy Ment Health*. 2000;27(6):371-82.

7. Hawthorne WB, Folsom, D. P., Sommerfeld, D. H., Lanouette, N. M., Lewis, M., Aarons, G. A., Conklin, R. M., Solorzano, E., Lindamer, L. A., & Jeste DV. Incarceration Among Adults Who Are in the Public Mental Health System: Rates, Risk Factors, and Short-Term Outcomes. *Psychiatric Services*. 2012;63(1):26-32.

- 1
2
3 8. Lamberti JS. Understanding and preventing criminal recidivism among adults with
4
5 psychotic disorders. *Psychiatr Serv.* 2007;58(6):773-81.
6
7
- 8 9. Kouyoumdjian F, Schuler, A., Hwang, S. W. & Matheson, F. Research on the health of
9
10 people who experience detention or incarceration in Canada: a scoping review. *BMC public*
11
12 *health.* 2015;15(419):1-8.
13
14
- 15 10. Kouyoumdjian FG, Cheng S. Y., Fung, K., Orkin, A. M., McIsaac K. E., Kendall, C., Kiefer, L.,
16
17 Matheson, F. I., Green, S. E., and Hwang, S. W. . The health care utilization of people in prison
18
19 and after prison release: A population-based cohort study in Ontario, Canada. *PLoS One.* August
20
21 3, 2018;13(8).
22
23
- 24 11. Kurdyak P, Lin, E., Green, D., & Vigod, S. Validation of a Population-Based Algorithm to
25
26 Detect Chronic Psychotic Illness. *Can J Psychiatry.* 2015;60(8):362-8.
27
28
- 29 12. Statistics Canada. Income in Canada 2015 [Available from:
30
31 <https://www.statcan.gc.ca/pub/75-202-x/2010000/analysis-analyses-eng.htm>.
32
33
- 34 13. Krajl B. Measuring Rurality - RIO2008 BASIC: Methodology and Results. Toronto (ON):
35
36 OMA Economics Department; 2009.
37
38
- 39 14. Steele LS, Glazier, R. H., Lin, E., & Evans, M. Using administrative data to measure
40
41 ambulatory mental health service provision in primary care. *Medical Care.* 2004;42(10):960-5.
42
43
- 44 15. John Hopkins University. Johns Hopkins ACG® Case-Mix Adjustment System Version 10
45
46 [Available from: <http://www.acg.jhsph.edu>.
47
48
- 49 16. Austin PC, Walraven, C., Wodchis, W., Newman, A. & Anderson, G. M. Using the Johns
50
51 Hopkins Aggregated Diagnosis Groups (ADGs) to predict mortality in a general adult population
52
53 cohort in Ontario, Canada. *Canadian Institutes of Health Research.* 2011;40(10):932-9.
54
55
56
57
58
59
60

17. Austin PC, Newman A, Kurdyak PA. Using the Johns Hopkins Aggregated Diagnosis Groups (ADGs) to predict mortality in a population-based cohort of adults with schizophrenia in Ontario, Canada. *Psychiatry Res.* 2012;196(1):32-7.

18. SAS. SAS System, Version 9: Installation and Administration Information. 2007.

19. Brinded PM, Simpson, A. I., Laidlaw, T. M., Fairley, N., & Malcolm, F. Prevalence of psychiatric disorders in New Zealand prisons: a national study. *Aust N Z J Psychiatry.* 2001;35(2):166-73.

20. Sirdifield C, Gojkovic, D., Brooker, C., & Ferriter, M. A systematic review of research on the epidemiology of mental health disorders in prison populations: a summary of findings. *Journal of Forensic Psychiatry & Psychology.* 2009;20(1):S78-S101.

21. Lafortune D. Prevalence and screening of mental disorders in short-term correctional facilities. *Int J Law Psychiatry.* 2010;33(2):94-100.

22. Senior J, Birmingham, L., Harty, M. A., Hassan, L., Hayes, A. J., Kendall, K., King, C., Lathlean, J., Lowthian, C., Mills, A., Webb, R., Thornicroft, G., & Shaw, J. Identification and management of prisoners with severe psychiatric illness by specialist mental health services. *Psychol Med.* 2013;43(7):1511-20.

23. Fazel S, & Danesh, J. Serious mental disorder in 23000 prisoners: a systematic review of 62 surveys. *Lancet.* 2002;359(9306):545-50.

24. Goldner EM, Jones, W., & Waraich, P. Using administrative data to analyze the prevalence and distribution of schizophrenic disorders. *Psychiatric Services.* 2003;54(7):1017-21.

25. Khan S, Chiu, M., Simpson, A. I., Guttman, A., Jembere, N., & Kurdyak, P. Use of Mental Health Services by Youths and Young Adults Before and During Correctional Custody: A Population-Based Study. *Psychiatr Serv.* 2016;67(7):790-3.
26. Simpson AI, McMaster, J. J., & Cohen, S. N. Challenges for Canada in meeting the needs of persons with serious mental illness in prison. *J Am Acad Psychiatry Law.* 2013;41(4):501-9.
27. National Institute of Health and Care Excellence. Psychosis and schizophrenia in adults. 2015.
28. Health Quality Ontario. Schizophrenia Care in the Community for Adults Ontario: Queen's Printer for Ontario; 2019 [Available from: <https://www.hqontario.ca/Evidence-to-Improve-Care/Quality-Standards/View-all-Quality-Standards/Schizophrenia-Care-in-the-Community/The-Quality-Standard-In-Brief>].
29. Prince JD. Incarceration and hospital care. *J Nerv Ment Dis.* 2006;194(1):34-9.
30. Harty M, Jarrett, M., Thornicroft, G., & Shaw, J. Unmet needs of male prisoners under the care of prison Mental Health Inreach Services. *Journal of Forensic Psychiatry & Psychology.* 2012;23(3):285-96.
31. Prince JD, Akincigil, A., & Bromet, E. Incarceration rates of persons with first-admission psychosis. *Psychiatr Serv.* 2007;58(9):1173-80.
32. Robertson AG, Swanson, J. W., Frisman, L. K., Lin, H., & Swartz, M. S. Patterns of justice involvement among adults with schizophrenia and bipolar disorder: key risk factors. *Psychiatr Serv.* 2014;65(7):931-8.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

33. Wallace C, Mullen, P. E., & Burgess, P. Criminal offending in schizophrenia over a 25-year period marked by deinstitutionalization and increasing prevalence of comorbid substance use disorders. *Am J Psychiatry*. 2004;161(4):716-27.

Table 1: Characteristics of study cohort, by schizophrenia diagnosis

Variable	Individuals without schizophrenia	Individuals with schizophrenia	Standardized Difference
	N (%)	N (%)	
Total	43,691 (93.1)	3,237 (7.0)	
Age			
Median (IQR)	32 (24-43)	36 (28-45)	0.30
18-29	19,007 (43.5)	1,004 (31.0)	0.26
30-39	10,775 (24.7)	936 (28.9)	0.10
40-49	9,175 (21.0)	825 (25.5)	0.11
50-64	4,322 (9.9)	425 (13.1)	0.10
65-74	354 (0.8)	39 (1.2)	0.04
75+	58 (0.1)	8 (0.2)	0.03
Sex			
Female	5,280 (12.1)	542 (16.7)	0.13
Male	38,411 (87.9)	2,695 (83.3)	0.13
Self-reported race			
Missing	2,952 (6.8)	223 (6.9)	0.01
Aboriginal	4,763 (10.9)	271 (8.4)	0.09
Black	5,248 (12.0)	447 (13.8)	0.05
East Asian	715 (1.6)	43 (1.3)	0.03
Hispanic	519 (1.2)	26 (0.8)	0.04
South Asian	908 (2.1)	68 (2.1)	0.00
South East Asian	557 (1.3)	46 (1.4)	0.01
West Asian/Arabic	671 (1.5)	46 (1.4)	0.01
White	25,973 (59.4)	1,958 (60.5)	0.02
Other racial origin	1,000 (2.3)	79 (2.4)	0.01
Declined to specify	130 (0.3)	12 (0.4)	0.01
Racial origin unknown	254 (0.6)	17 (0.5)	0.01
Income Quintile			
Missing	3,207 (7.3)	96 (3.0)	0.20
1	15,359 (35.2)	1,334 (41.2)	0.12
2	9,218 (21.1)	696 (21.5)	0.01
3	6,802 (15.6)	508 (15.7)	0.00
4	5,194 (11.9)	330 (10.2)	0.05
5	3,911 (9.0)	273 (8.4)	0.02
Rurality			
Missing	2,398 (5.5)	25 (0.8)	0.27
Urban	35,533 (81.3)	2,953 (91.2)	0.29
Rural	5,760 (13.2)	259 (8.0)	0.17
Comorbidity (cADGs)			

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Mean ± SD	3.0±2.1	4.2±2.0	0.60
Median (IQR)	3(1-4)	4(3-6)	0.61

Legend: IQR – interquartile range; cADGs – Collapsed Aggregated Diagnostic groups; SD – standard deviation

Table 2: Correctional involvement three years prior to the index* incarceration by schizophrenia diagnosis

Variable	Individuals without schizophrenia	Individuals with schizophrenia	Standardized Difference
	N (%)	N (%)	
Total	43,691 (93.1)	3,237 (7.0)	
Length of stay from all correctional entries in 3 years prior to index release			
Median (IQR)	38 (7-123)	66 (13-176)	0.26
No prior incarcerations	22,904 (52.4)	1,327 (41.0)	0.23
1-6	5,115 (11.7)	323 (10.0)	0.06
7-29	4,464 (10.2)	354 (10.9)	0.02
30-89	4,385 (10.0)	430 (13.3)	0.10
90-179	3,414 (7.8)	335 (10.3)	0.09
180-364	2,570 (5.9)	332 (10.3)	0.16
365+	839 (1.9)	136 (4.2)	0.13
Length of stay of index incarceration event			
Median (IQR)	7 (2-37)	11 (3-48)	0.16
1-6	21,101 (48.3)	1,324 (40.9)	0.15
7-29	10,104 (23.1)	840 (25.9)	0.07
30-89	7,434 (17.0)	638 (19.7)	0.07
90-119	1,813 (4.1)	140 (4.3)	0.01
120-364	2,966 (6.8)	273 (8.4)	0.06
365+	273 (0.6)	22 (0.7)	0.01
Number of prior incarcerations			
Median (IQR)	0 (0-2)	1 (0-3)	0.3
0	22,904 (52.4)	1,327 (41.0)	0.23
1	8,151 (18.7)	566 (17.5)	0.03
2	4,868 (11.1)	357 (11.0)	0.00
3	2,929 (6.7)	266 (8.2)	0.06
4	1,846 (4.2)	207 (6.4)	0.10
5	1,071 (2.5)	139 (4.3)	0.10
6	661 (1.5)	84 (2.6)	0.08
7	450 (1.0)	68 (2.1)	0.09
8	259 (0.6)	57 (1.8)	0.11
9	175 (0.4)	46 (1.4)	0.11
10+	377 (0.9)	120 (3.7)	0.19

*Index refers to the first correctional release for each individual in 2010 in Ontario.

Legend: IQR – interquartile range


Table 3: Correctional involvement five years after index release, by schizophrenia diagnosis

Variable	Individuals without schizophrenia	Individuals with schizophrenia	Standardized Difference
	N (%)	N (%)	
Total	43,691 (93.1)	3,237 (7.0)	
Any re-incarceration within 5 years	25,703 (58.8)	2,185 (67.5)	0.18
Time to re-incarceration			
Median (IQR)	219 (81-531)	156 (52-410)	0.21
Not reincarcerated	17,988 (41.2)	1,052 (32.5)	0.18
1-30 days	2,853 (6.5)	359 (11.1)	0.16
31-60	2,249 (5.1)	259 (8.0)	0.12
61-90	1,908 (4.4)	184 (5.7)	0.06
91-240	6,503 (14.9)	528 (16.3)	0.04
241-365	3,143 (7.2)	240 (7.4)	0.01
366-730	4,554 (10.4)	330 (10.2)	0.01
731-1095	2,079 (4.8)	96 (3.0)	0.09
3 years +	2,414 (5.5)	189 (5.8)	0.01
Length of stay from all correctional events in 5 years after index release			
Median (IQR)	83 (20-204)	88 (23-222)	0.07
Not re-incarcerated	17,988 (41.2)	1,052 (32.5)	0.18
1-6 days	3,563 (8.2)	246 (7.6)	0.02
7-29	4,181 (9.6)	373 (11.5)	0.06
30-89	5,597 (12.8)	483 (14.9)	0.06
90-179	4,999 (11.4)	386 (11.9)	0.02
180-364	4,501 (10.3)	402 (12.4)	0.07
365-729	2,473 (5.7)	232 (7.2)	0.06
730+	389 (0.9)	63 (1.9)	0.09

*Index refers to the first correctional release for each individual in 2010 in Ontario.


Legend: IQR – interquartile range

Figure 1A: Time to re-incarceration for individuals with and without schizophrenia from multivariable Cox regression*



*Adjusted for age, sex, neighbourhood income quintile and rurality.

Figure 1B: Recidivism survival curves between individuals with and without schizophrenia.*



*Adjusted for age, sex, neighbourhood income quintile, rurality, comorbidity, prior correctional involvement and prior health service utilization.

Supplement Table 1

Health Service Utilization Three Years Prior to the Index Incarceration*

Variable	Individuals without schizophrenia	Individuals with schizophrenia	Standardized Difference
	N (%)	N (%)	
Total	43,691 (93.1)	3,237 (7.0)	
Total number visits to PCP			
Median (IQR)	5 (1-14)	13 (5-28)	0.58
0	8,005 (18.3)	210 (6.5)	0.36
1-5	14,135 (32.4)	659 (20.4)	0.27
6-10	7,394 (16.9)	569 (17.6)	0.02
11-15	4,244 (9.7)	401 (12.4)	0.09
16-20	2,460 (5.6)	288 (8.9)	0.13
21+	7,453 (17.1)	1,110 (34.3)	0.40
Total number of MH visits to PCP			
Median (IQR)	0 (0-3)	5 (1-12)	0.92
0	23,706 (54.3)	565 (17.5)	0.83
1-5	12,931 (29.6)	1,187 (36.7)	0.15
6-10	2,564 (5.9)	551 (17.0)	0.36
11-15	951 (2.2)	287 (8.9)	0.30
16-20	552 (1.3)	160 (4.9)	0.21
21+	2,987 (6.8)	487 (15.0)	0.27
Total number of non-MH visits to PCP			
Median (IQR)	4 (1-10)	6 (2-14)	0.30
0	9,143 (20.9)	396 (12.2)	0.24
1-5	16,903 (38.7)	1,180 (36.5)	0.05
6-10	7,750 (17.7)	601 (18.6)	0.02
11-15	3,875 (8.9)	328 (10.1)	0.04
16-20	2,044 (4.7)	222 (6.9)	0.09
21+	3,976 (9.1)	510 (15.8)	0.2
Total number of psychiatrist visits			
Median (IQR)	0 (0-0)	4 (0-12)	1.42
0	36,536 (83.6)	885 (27.3)	1.37
1-3	4,248 (9.7)	710 (21.9)	0.34
4-6	1,180 (2.7)	399 (12.3)	0.37
7-9	533 (1.2)	276 (8.5)	0.34
10+	1,194 (2.7)	967 (29.9)	0.79
Total ED visits			
Median (IQR)	1 (0-3)	3 (1-8)	0.60

1				
2				
3	0	15,189 (34.8)	531 (16.4)	0.43
4	1-5	22,118 (50.6)	1,549 (47.9)	0.06
5	6-10	4,042 (9.3)	528 (16.3)	0.21
6	11-15	1,215 (2.8)	237 (7.3)	0.21
7	16-20	493 (1.1)	121 (3.7)	0.17
8	21+	634 (1.5)	271 (8.4)	0.32
9				
10	Total hospitalizations			
11	Median (IQR)	0 (0-0)	1 (0-2)	0.94
12	0	37,706 (86.3)	1,523 (47.0)	0.92
13	1-3	5,572 (12.8)	1,316 (40.7)	0.66
14	4-6	311 (0.7)	276 (8.5)	0.38
15	7-9	63 (0.1)	68 (2.1)	0.19
16	10+	39 (0.1)	54 (1.7)	0.10
17				
18	Total inpatient days			
19	Median (IQR)	0 (0-0)	2 (0-24)	0.97
20	0	38,048 (87.1)	1,550 (47.9)	0.92
21	1-10	4,147 (9.5)	521 (16.1)	0.20
22	11-89	1,451 (3.3)	921 (28.5)	0.73
23	90-180	32 (0.1)	162 (5.0)	0.32
24	181+	13 (0.0)	83 (2.6)	0.23
25				
26	Total psychiatric hospitalizations			
27	Median (IQR)	0 (0-0)	0 (0-2)	1.08
28	0	42,062 (96.3)	1,814 (56.0)	1.07
29	1-3	1,555 (3.6)	1,130 (34.9)	0.87
30	4-6	61 (0.1)	215 (6.6)	0.37
31	7-9	8 (0.0)	48 (1.5)	0.17
32	10+	**	**	**
33				
34	Total psychiatric hospital days			
35	Median (IQR)	0 (0-0)	0 (0-22)	1.08
36	0	42,134 (96.4)	1,825 (56.4)	1.07
37	1-10	795 (1.8)	337 (10.4)	0.36
38	11-89	734 (1.7)	843 (26.0)	0.75
39	90-180	20 (0.0)	151 (4.7)	0.31
40	181+	8 (0.0)	81 (2.5)	0.22
41				
42				
43				
44				
45				
46				

*Index refers to the first correctional release for each individual in 2010 in Ontario.

Legend: PCP – primary care physician; IQR – interquartile range; MH – mental health

Notes: ** Suppressed values due to n<6

Supplementary Table 2

Health service utilization outcomes five years after the index release*

Schizophrenia Indicator				
Variable	Individuals without schizophrenia	Individuals with schizophrenia	All Persons	Standardized Difference
	N (%)	N (%)	N (%)	
Total	43,691 (93.1)	3,237 (7.0)	46,928 (100)	
Number of PCP visits				
Median (IQR)	10 (3-28)	19 (7-43)	11 (3-29)	0.38
0	5,794 (13.3)	164 (5.1)	5,958 (12.7)	0.29
1-5	9,659 (22.1)	501 (15.5)	10,160 (21.7)	0.17
6-10	6,410 (14.7)	434 (13.4)	6,844 (14.6)	0.04
11-15	4,394 (10.1)	317 (9.8)	4,711 (10.0)	0.01
16-20	3,182 (7.3)	293 (9.1)	3,475 (7.4)	0.06
21+	14,252 (32.6)	1,528 (47.2)	15,780 (33.6)	0.30
Number of MH visits to PCP				
Median (IQR)	1 (0-6)	7 (2-19)	1 (0-7)	0.72
0	18,749 (42.9)	454 (14.0)	19,203 (40.9)	0.68
1-5	13,168 (30.1)	992 (30.6%)	14,160 (30.2)	0.01
6-10	3,217 (7.4)	522 (16.1)	3,739 (8.0)	0.27
11-15	1,523 (3.5)	314 (9.7)	1,837 (3.9)	0.25
16-20	832 (1.9)	207 (6.4)	1,039 (2.2)	0.23
21+	6,202 (14.2)	748 (23.1)	6,950 (14.8)	0.23
Number of non-MH visits to PCP				
Median (IQR)	7 (2-16)	8 (3-21)	7 (2-17)	0.17
0	6,837 (15.6)	313 (9.7)	7,150 (15.2)	0.18
1-5	12,505 (28.6)	934 (28.9)	13,439 (28.6)	0.01
6-10	7,685 (17.6)	562 (17.4)	8,247 (17.6)	0.01
11-15	5,026 (11.5)	383 (11.8)	5,409 (11.5)	0.01
16-20	3,223 (7.4)	230 (7.1)	3,453 (7.4)	0.01
21+	8,415 (19.3)	815 (25.2)	9,230 (19.7)	0.14
Total number of psychiatrist visits				
Median (IQR)	0 (0-1)	6 (1-20)	0 (0-1)	1.2
0	31,830 (72.9)	776 (24.0)	32,606 (69.5)	1.12
1-5	7,163 (16.4)	787 (24.3)	7,950 (16.9)	0.20
6-10	1,806 (4.1)	399 (12.3)	2,205 (4.7)	0.30
11-15	861 (2.0)	288 (8.9)	1,149 (2.4)	0.31
16-20	536 (1.2)	212 (6.5)	748 (1.6)	0.28
21+	1,495 (3.4)	775 (23.9)	2,270 (4.8)	0.63
Total ED visits				
Median (IQR)	2 (0-5)	4 (1-12)	2 (0-5)	0.47
0	11,846 (27.1)	491 (15.2)	12,337 (26.3)	0.30
1-5	21,590 (49.4)	1,351 (41.7)	22,941 (48.9)	0.15
6-10	5,844 (13.4)	510 (15.8)	6,354 (13.5)	0.07
11-15	2,093 (4.8)	303 (9.4)	2,396 (5.1)	0.18

16-20	897 (2.1)	162 (5.0)	1,059 (2.3)	0.18
21+	1,421 (3.3)	420 (13.0)	1,841 (3.9)	0.36
Total Mental Health Related				
ED Visits				
Median (IQR)	0 (0-0)	1 (0-3)	0 (0-0)	0.75
0	34,264 (78.4)	1,478 (45.7)	35,742 (76.2)	0.72
1-5	8,299 (19.0)	1,297 (40.1)	9,596 (20.4)	0.47
6-10	616 (1.4)	189 (5.8)	805 (1.7)	0.24
11-15	186 (0.4)	105 (3.2)	291 (0.6)	0.21
16-20	85 (0.2)	48 (1.5)	133 (0.3)	0.14
21+	241 (0.6)	120 (3.7)	361 (0.8)	0.22
Total hospitalizations				
Median (IQR)	0 (0-0)	1 (0-3)	0 (0-0)	0.84
0	34,475 (78.9)	1,383 (42.7)	35,858 (76.4)	0.80
1-5	8,582 (19.6)	1,529 (47.2)	10,111 (21.5)	0.61
6-10	451 (1.0)	226 (7.0)	677 (1.4)	0.31
11-15	115 (0.3)	56 (1.7)	171 (0.4)	0.15
16-20	39 (0.1)	25 (0.8)	64 (0.1)	0.10
21+	29 (0.1)	18 (0.6)	47 (0.1)	0.09
Total inpatient days				
Median (IQR)	0 (0-0)	4 (0-45)	0 (0-0)	0.87
0	34,946 (80.0)	1,420 (43.9)	36,366 (77.5)	0.80
1-10	5,420 (12.4)	463 (14.3)	5,883 (12.5)	0.06
11-89	2,969 (6.8)	818 (25.3)	3,787 (8.1)	0.52
90-180	231 (0.5)	238 (7.4)	469 (1.0)	0.36
181-365	74 (0.2)	122 (3.8)	196 (0.4)	0.26
366+	51 (0.1)	176 (5.4)	227(0.5)	0.33
Total psychiatric hospitalizations				
Median (IQR)	0 (0-0)	0 (0-2)	0 (0-0)	0.97
0	40,975 (93.8)	1,817 (56.1)	42,792 (91.2)	0.96
1-5	2,583 (5.9)	1,206 (37.3)	3,789 (8.1)	0.82
6-10	107 (0.2)	158 (4.9)	265 (0.6)	0.30
11-15	18 (0.0)	42 (1.3)	60 (0.1)	0.15
16-20	7 (0.0)	10 (0.3)	17 (0.0)	0.07
21+	**	**	**	**
Total number of psychiatric hospital days				
Median (IQR)	0 (0-0)	0 (0-35)	0 (0-0)	0.97
0	41,045 (93.9)	1,838 (56.8)	42,883 (91.4)	0.96
1-10	1,131 (2.6)	224 (6.9)	1,355 (2.9)	0.20
11-89	1,321 (3.0)	687 (21.2)	2,008 (4.3)	0.58
90-180	107 (0.2)	200 (6.2)	307 (0.7)	0.34
181-365	43 (0.1)	112 (3.5)	155 (0.3)	0.26
366+	44 (0.1)	176 (5.4)	220 (0.5)	0.33

*Index refers to the first correctional release for each individual in 2010 in Ontario.

Legend: PCP – primary care physician; IQR – interquartile range; MH – mental health

Notes: ** Suppressed values due to n<6

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Prévalence et prédicteurs de la réincarcération après la libération d’un centre correctionnel : une comparaison dans la population-de personnes souffrant ou non de schizophrénie en Ontario, Canada.

Objectifs : Les personnes souffrant de schizophrénie sont surreprésentées dans les établissements correctionnels relativement à leur prévalence dans la population. La présente étude a pour but de déterminer le taux et les prédicteurs de la réincarcération des personnes souffrant de schizophrénie après leur libération des établissements correctionnels.

Méthodes : Il s’agissait d’une étude de cohorte rétrospective qui incluait toutes les personnes libérées des établissements correctionnels provinciaux de l’Ontario du 1^{er} janvier au 31 décembre 2010. Les personnes souffrant de schizophrénie ont été identifiées à l’aide d’un algorithme dans la population. Le résultat principal était le temps écoulé jusqu’à la réincarcération. Les covariables étaient notamment les caractéristiques sociodémographiques (âge, sexe, quintile de revenu du quartier, résidence urbaine/rurale), l’utilisation des services de santé (visites à des médecins de soins de première ligne, visites à un psychiatre, hospitalisations psychiatriques ou autres, visites au service d’urgence) et d’autre morbidité clinique. L’analyse de survie a servi à examiner l’association entre la schizophrénie et la réincarcération.

Résultats : Parmi les 46 928 personnes, N = 3 237 (7 %) avaient un diagnostic de schizophrénie. Environ 67,5 % de celles-ci ont été réincarcérées dans les 5 années suivant leur première libération en 2010, comparé à 58,8 % des personnes ne souffrant pas de schizophrénie. Les personnes souffrant de schizophrénie étaient 40 % (RR : 1,39; IC à 95 % 1,33 à 1,45) plus susceptibles d’être réincarcérées après une libération que le groupe témoin après ajustement des caractéristiques démographiques. Cette association s’est réduite à 8 % (1,08; IC à 95 % 1,03 à 1,14) après ajustement pour utilisation précédente des services de santé, implication correctionnelle précédente et comorbidités.

Conclusion : Les personnes souffrant de schizophrénie étaient plus susceptibles de connaître une réincarcération après une libération des établissements correctionnels. Ce risque

s'explique en partie par l'implication correctionnelle précédente, l'utilisation des services de santé, et les comorbidités. La recherche future devrait mettre l'accent sur les facteurs de risque qui prédisent le taux de réincarcération plus élevé et sur les interventions pour réduire l'implication correctionnelle.