

This is a repository copy of *Characterizing Family Physicians Who Refer to Telepsychiatry in Ontario*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/166050/>

Version: Accepted Version

Article:

Serhal, Eva, Iwajomo, Tomisin, de Oliveira, Claire orcid.org/0000-0003-3961-6008 et al. (2 more authors) (2020) Characterizing Family Physicians Who Refer to Telepsychiatry in Ontario. *Canadian journal of psychiatry-Revue canadienne de psychiatrie*. p. 706743720954061. ISSN 0706-7437

<https://doi.org/10.1177/0706743720954061>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Characterizing Family Physicians Who Refer to Telepsychiatry in Ontario

Journal:	<i>The Canadian Journal of Psychiatry/La Revue canadienne de psychiatrie</i>
Manuscript ID	CJP-2020-122-OR.R1
Manuscript Type:	Original Research
Date Submitted by the Author:	26-Jun-2020
Complete List of Authors:	Serhal, Eva; Centre for Addiction and Mental Health, ; University of Toronto, Institute of Health Policy, Management and Evaluation Iwajomo, Tomisin; Centre for Addiction and Mental Health de Oliveira, Claire; University of York, Hull York Medical School/Centre for Health Economics; Centre for Addiction and Mental Health, Institute for Mental Health Policy Research Crawford, Allison; Centre for Addiction and Mental Health, Northern Psychiatric Outreach Program and Telepsychiatry; University of Toronto, Psychiatry Kurdyak, Paul; Centre for Addiction and Mental Health, General Psychiatry
Key Words:	Telepsychiatry, Primary care, Telehealth, Family Physician, Adult Psychiatry, Telemedicine, Virtual Care, referral, Telemental Health
Abstract:	<p>Objectives: Telepsychiatry can improve access to psychiatric services for those who otherwise cannot easily access care. Family physicians are gatekeepers to specialized care in Ontario, so it is essential to understand predictors relating to referrals to telepsychiatry to better plan services and increase telepsychiatry adoption.</p> <p>Methods: This study used an annual retrospective cross-sectional study design to compare physicians who referred their patients to telepsychiatry each year from fiscal year (FY) 2008 to FY 2016. A one year (FY 2016) comparison of family physicians who referred to telepsychiatry compared to family physicians who did not refer to telepsychiatry matched (1:2) by region was also conducted. Finally, we used statistical modelling to understand the predictors of referring to telepsychiatry among physicians.</p> <p>Results: Between FY 2008 to FY 2016, telepsychiatry visits increased from 925 visits to 13,825 visits. Thirty-two percent of Ontario primary care physicians referred to telepsychiatry in 2017. Several characteristics were notably different between family physicians who refer to telepsychiatry (FPTs) and family physicians who do not refer to telepsychiatry (FPNTs): FPTs were more likely to be from a residence with less than 10,000 people, to have more nurse practitioners in the</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	<p>practice, and to be from a family health team than FPNTs. Rostered patients of FPTs were more likely to reside in rural areas, have more clinical complexity, and to utilize more mental health services compared to FPNTs.</p> <p>Conclusions: There has been an increase in the use of telepsychiatry by patients and family physicians over the study period and family physicians that are live in a rural area, are part of a FHT, have more NPs, with more patients in rural areas with high needs for mental health services are more likely to refer to telepsychiatry.</p>

SCHOLARONE™
Manuscripts

CHARACTERIZING FAMILY PHYSICIANS WHO REFER TO TELEPSYCHIATRY IN ONTARIO

Authors:

Eva Serhal^{1,2}, Tomisin Iwajomo^{1,3}, Claire de Oliveira¹⁻³, Allison Crawford^{1,4}, Paul

Kurdyak¹⁻³

Affiliations

1. The Centre for Addiction and Mental Health, Toronto, Ontario
2. The Institute for Health Policy, Management and Evaluation, University of Toronto, Toronto, Ontario
3. ICES, Toronto, Ontario
4. Dalla Lana School of Public Health, University of Toronto, Toronto, Ontario

Corresponding Author:

Eva Serhal, MBA, PhD
Director, Virtual Mental Health and Outreach & ECHO Ontario
Centre for Addiction and Mental Health
821-250 College Street, Toronto, Ontario
M5T1R8

Abstract

Introduction

Telepsychiatry can improve access to psychiatric services for those who otherwise cannot easily access care. Family physicians are gatekeepers to specialized care in Ontario, so it is essential to understand predictors relating to referrals to telepsychiatry to better plan services and increase telepsychiatry adoption.

Methods

This study used an annual retrospective cross-sectional study design to compare physicians who referred their patients to telepsychiatry each year from fiscal year (FY) 2008 to FY 2016. A one year (FY 2016) comparison of family physicians who referred to telepsychiatry compared to family physicians who did not refer to telepsychiatry matched (1:2) by region was also conducted. Finally, we used statistical modelling to understand the predictors of referring to telepsychiatry among physicians.

Results

Between FY 2008 to FY 2016, telepsychiatry visits increased from 925 visits to 13,825 visits. Thirty-two percent of Ontario primary care physicians referred to telepsychiatry in 2017. Several characteristics were notably different between family physicians who refer to telepsychiatry (FPTs) and family physicians who do not refer to telepsychiatry (FPNTs): FPTs were more likely to be from a residence with less than 10,000 people, to have more nurse practitioners in the practice, and to be from a family health team than

1
2
3 FPNTs. Rostered patients of FPTs were more likely to reside in rural areas, have more
4
5 clinical complexity, and to utilize more mental health services compared to FPNTs.
6
7

8 ***Conclusion***

9
10
11
12 There has been an increase in the use of telepsychiatry by patients and family
13
14 physicians over the study period, although there remains opportunity for significant
15
16 growth. Family physicians that live in rural areas, are part of a FHT, have more NPs,
17
18 with more rural and complex patients were more likely to refer to telepsychiatry. As
19
20 recent pro-telemedicine policies support the growth of telepsychiatry, this study will
21
22 serve as an important baseline.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

Mental illness is among the world's leading causes of sickness and disability; one in five Canadians (two million Ontarians) and one in four people internationally will experience mental illness or addiction.¹⁻³ However, only one-third of Ontario patients reported having their mental health needs met or partially met within primary care.¹ Patients from lower-income countries and socially disadvantaged groups had lower rates of mental health service utilization.⁴⁻⁶ Access to quality mental health services is often limited due to rurality, physician shortages and variation in the availability of mental health services.^{1,5,7-10}

Telepsychiatry, the delivery of psychiatry via secure videoconferencing, can reduce access barriers such as distance, cost of travel, weather, and workforce shortages.¹¹⁻¹⁷ In particular, telepsychiatry can provide access to patients who otherwise are not able to easily access services, potentially reducing unnecessary hospitalizations and ED visits.^{18,19} Several studies have deemed telepsychiatry effective for various mental health conditions and populations.²⁰⁻²⁴ Telemedicine has been shown to reduce isolation and increase the morale of rural primary care practitioners.^{25,26} While there is significant potential for telepsychiatry to connect patients to psychiatrists throughout Ontario, recent evidence has shown minimal adoption by patients.²⁷ In Ontario during the time of this study, in order to deliver and bill for telepsychiatry, psychiatrists were typically required to use a centralized government funded telemedicine platform and patients were typically required to access their telepsychiatry visit at a patient host site

1
2
3 (a clinical site that connects to the centralized provincial network). It is important to note
4
5 that at the time this paper was submitted temporary policy changes to virtual care as a
6
7 result of Covid-19 (introduced March 2020) have created more options for home-based
8
9 care for patients, and have allowed physicians to deliver virtual care via a variety of
10
11 virtual platforms. These recent changes will be discussed in relation to study findings
12
13 in the discussion section.
14
15
16
17
18

19 Family physicians are the gatekeepers to specialist care in Canada and other parts of
20
21 the world.^{28,29} Common reasons for family physician referral to psychiatry include
22
23 obtaining medication recommendations, obtaining second opinions, or non-
24
25 pharmacological treatment recommendations.³⁰ Evidence shows providers who are
26
27 more isolated had more positive feelings about telemedicine.²⁵ Positive perceptions of
28
29 telemedicine included convenience, perceived patient benefits, reduced wait time, cost-
30
31 effectiveness, and access to medical expertise, while negative aspects included cost,
32
33 reduced funding for in-person services, inconvenience, inadequate reimbursement and
34
35 training, medical liability, and physician fear of practice change.^{25,31-34} Previous studies
36
37 suggest nurses may be more enthusiastic about telemedicine than physicians, referring
38
39 physicians tend to be more satisfied than specialists, satisfaction increases relative to
40
41 use and time, and that providers do not recommend telemedicine in all
42
43 circumstances.^{29,35,36} The objective of this study was to examine the characteristics of
44
45 family physicians in Ontario who have referred patients to telepsychiatry (FPTs)
46
47 compared to family physicians who have not referred to telepsychiatry (FPNTs) from FY
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 2008 to FY 2016, in order to understand if there are specific factors that make certain
4
5 family physicians more likely to refer than others.
6
7
8
9

10 **Methods**

13 ***Setting and Study Design***

16
17 We conducted an annual retrospective cross-sectional study assessing descriptive
18 characteristics of FPTs for FYs (2008 to 2016); this time horizon was used because it
19 includes data from when the Ontario Health Insurance Plan (OHIP) physician fee
20 schedule telemedicine codes were introduced until the most recent data available from
21 ICES. Referring physician and patient information were ascertained using the OHIP
22 claims database; if referral information was not available, we looked back within the
23 past 12 months to identify the most frequently seen family physician and, if not
24 available, the most recently seen family physician. These data were linked to
25 population-level health administrative databases held at ICES. Additionally, a one year
26 (FY2016) comparison of FPTs vs FPNTs matched (1:2) by the 14 Local Health
27 Integration Networks (LHIN). LHINs are artificially divided regions of Ontario which were
28 designed to support local approaches to strategic planning, funding and regional
29 coordination of clinical care, and are used in administrative data to classify provincial
30 regions. This study was approved by the Research Ethics Board of Sunnybrook Health
31 Sciences Centre, Toronto.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Data Sources

This study used province-wide health administrative databases that were linked using unique encoded identifiers and analyzed at ICES. OHIP data were used to collect family physician details, including physician number, fee code, service date, and location of service provided, and patient information, including patient ID, diagnosis, service date, and location. The National Ambulatory Care Reporting System was used to collect data on emergency department visits. The Ontario Mental Health Reporting System was used to obtain information on psychiatric hospitalizations. The ICES Physician Database and the Primary Care Population database were used to analyze data on primary care providers and teams, including the number of years since medical school graduation, years of practice since graduation, provider type, interprofessional team details, and total physicians per practice. The Client Agency Program Enrolment database was used to analyze data on physician practice type (defined as patient enrollment type). The Ontario Census Area Profiles and the Registered Persons Database were used to analyze patient-level demographic variables (such as age, sex, the region of residence), as well as geographic details (rurality, income quintiles). Primary care providers who had missing age or sex data, and patients with missing health card numbers, missing age or missing sex were excluded from the analysis.

Provider Characteristics

We selected primary care provider characteristics based on two things: availability within the ICES dataset, and previous literature within the field that assessed predictors

1
2
3 using these measures.³⁴ The primary care provider characteristics used to compare
4
5 FPTs vs FPNTs included age group, sex, number of years since medical school
6
7 graduation, years of practice since graduation, rurality of residence, physician practice
8
9 type (Family Health Team¹ or other), average number of nurse practitioners within the
10
11 practice, and average number of social workers within the practice.
12
13
14

15 ***Patient Characteristics***

16
17
18 We examined patient demographic and service utilization characteristics for patients
19
20 rostered within each physician group (FPT vs FPNT). Patient demographic
21
22 characteristics included age, sex, neighbourhood income quintile (NIQ) (NIQ 1= 0-20%,
23
24 NIQ 2= 20-40%, NIQ 3= 40-60%, NIQ 4= 60-80%, NIQ 5= 80%+), and whether they live
25
26 in a rural or urban area. Patient utilization and medical complexity were considered
27
28 within each annual cross-section by assessing: number of primary care provider visits
29
30 (total, mental health and non-mental health); number of emergency department visits
31
32 (total, mental health and non-mental health-related); number of hospitalizations (total,
33
34 mental health and non-mental health-related); and comorbidities using the Johns
35
36 Hopkins' Aggregated Diagnosis Groups (ADG). Mental health primary care visits were
37
38 defined using a validated algorithm to identify adults accessing care for a mental health
39
40 disorder, which included adult mental health diagnostic codes 295-298, 300-304, 306,
41
42 309, 311, 897-902, 904, 905 (which represents a broad range of adult mental health
43
44
45
46
47
48
49

50
51 ¹ Family health teams are different from many primary care practices because they are funded to include an
52 interprofessional team within the practice, and are also paid via a blended capitation model (where physicians are
53 paid primarily through capitation fees based on a defined basket of services delivered to enrolled patients) or a
54 salary model).¹⁶³
55

1
2
3 and addiction diagnoses as defined by the International Classification of Diseases 10
4 (ICD-10-CA).³⁷ Hospitalizations and ED visits included ICD-10-CA codes F04-F99
5
6 (which excludes dementia). The Johns Hopkins ADG's are categorized based on five
7
8 clinical dimensions, including the duration, the severity, the etiology, the diagnostic
9
10 certainty and the specialty care involvement of the condition.³⁸
11
12
13
14
15

16 ***Analysis***

17 *Descriptive Analysis*

18 Annual Outputs and Trends over Time

19 Annual panels containing physician and patient characteristics were described,
20
21 including counts, column percentages, and means with standard deviations.
22
23

24 Additionally, changes over time were evaluated for each of the physician characteristics.
25
26
27
28
29
30
31

32 Matched Physician Comparison

33 Matched comparisons between FTPs and FTNPs were presented as counts, column
34
35 percentages, and standardized mean differences (SMD). We compared between-group
36
37 differences using the standard mean difference (SMD). Cohen (1988) suggests a small
38
39 effect size is equal to 0.2 SMD, but recent studies suggest that a more prudent
40
41 threshold for assessing effect size is 0.1 SMD, especially with large data sets.^{39,40} Using
42
43 this threshold (0.1 SMD), this study identifies significant differences between groups.
44
45
46
47
48
49
50

51 The model covariates included physician age (categorical; <34, 35-44, 45-54, 55-64,
52
53 64+), sex (Male/Female), number of years after graduation (categorical; 1-10, 11-20,
54
55
56
57
58
59
60

9

1
2
3 21-30, 31-40, 41+ years), rural residence (living in community with less than 10,000
4 people- Yes/No), total nurse practitioner on the team (continuous), total social workers
5 on the team (continuous), if the family physician is a Canadian Medical Graduate
6 (Yes/No), the physician practice type (FHT/Other), patient age (categorical; 0-10, 11-20,
7 21-30, 31-40, 41-50, 51-60, 61-70, 71-80, 81+ years), patient sex (Male/Female),
8 patient NIQ (categorical; 1-5), if the FP has more patients in low NIQ areas (Yes/No),
9 patient residence (Urban/Rural), if the FP has more urban patients than rural (Yes/No),
10 if the FP has more patients with at least one psychosocial ADG (Yes/No), patient region
11 (categorical; LHIN1-14), total number of ED visits (categorical; 0-59, 60-156, 157-257,
12 258-414, 415+), total number of mental health ED visits (categorical;0-3, 4-9, 10-19,
13 20+ visits), total hospitalizations (categorical; 0, 1-2, 3-5, 6-10, 11+), total mental health
14 hospitalizations (categorical; 0, 1-2, 3-5, 6-10, 11+), total primary care visits
15 (categorical; 0-59, 60-156, 157-257, 258-414, 415+) and total mental health primary
16 care provider visits (categorical; 0-59, 60-156, 157-257, 258-414, 415+).

36 Statistical Analysis

37
38 We used statistical modeling to understand the predictors of referring to telepsychiatry
39 among physicians. We used GEE for the statistical model, as it accounts for patient
40 clustering at the provider level.⁴¹ All analyses were carried out using SAS version 9.4
41 (SAS Institute, Cary, NC). All results were presented as adjusted odds ratios (ORs) with
42 corresponding 95% confidence intervals (CIs).
43
44
45
46
47
48
49
50
51

52 **Results**

Descriptive analysis of FTPs

In FY 2016, there were 13,290 family physicians in Ontario; this analysis includes a subset of those family physicians who have referred to telepsychiatry.⁴²

Increase in Patients Accessing and Family Physicians Referring to Telepsychiatry

The number of patients receiving telepsychiatry increased from 925 patients in FY 2008 to 13,825 patients in FY 2016, which represents approximately a fifteen-fold increase (see Table 1). There was substantial growth in the number of FPTs from 474 to 4,224, which represents a nine-fold increase from FY 2008 to FY 2016 (see Table 1). Of all physicians in Ontario, 32% referred to telepsychiatry in FY 2016 (4224/13290). Less than 1% of total patients rostered to FPTs had a telepsychiatry visit in FY 2018 (n= 12,449/3,513,638). [Insert Figure 1]

Physician Rurality

In FY 2008, the percentage of total telepsychiatry visits was highest in the North East LHIN (20%, n = 94), followed by the North West LHIN (16%, n = 74) (Ontario's most northern regions), followed by the South West LHIN (15%, n = 70). In FY 2016, those numbers shifted so the highest percentage of telepsychiatry referrals came from family physicians in regions such as Hamilton Niagara Haldimand Brant (10%, n = 433), Central East (10%, n = 421), and Toronto Central (10%, n = 420) LHINs, with the Central (n = 393) and North East (n = 379) LHIN's family physicians referring 9% each. All LHINs ranged between 4-10% of total telepsychiatry delivery in FY 2016, compared

1
2
3 to FY 2008, where the range spread from 0% (in Mississauga Halton LHIN) to 20% (in
4 the North East LHIN). This shows a more evenly distributed adoption of telepsychiatry
5 from northern regions to all regions of Ontario, many of which still have highly rural and
6 remote communities. [Insert Figure 2]
7
8
9
10
11
12
13
14
15

16 **Comparative Analysis**

17 *Family Physician Characteristics*

18
19
20
21
22 A larger percentage of FPTs were male, compared to FPNTs (59.6% (n = 2427) vs.
23 53.1% (n = 4075), SD = 0.13). A greater percentage of FPTs were from communities
24 with a population that had less than 10,000 people (14.9% (n = 608) vs. 8.2% (n = 633),
25 SMD = 0.21). FPTs had a larger mean number of nurse practitioners and FPNTs (1.71
26 (+/- 4.02) vs. 1.10 (+/- 3.70), SMD = 0.16), finally, a greater percentage of FPTs
27 belonged to FHTs than FPNTs (28.1% (n = 1145) vs. 18% (n = 1381), SMD = 0.24).
28
29
30

31 [Insert Table 1]
32
33
34
35
36
37
38

39 *Comparing Patient Roster Characteristics*

40
41
42 There were a few notable differences in patient roster characteristics between FPTs and
43 FPNTs. FPTs had a larger mean number of attached patients than FPNTs (862.45 (+/-
44 668.36) vs. 788.97 (+/- 605.70), SMD = 0.15). A greater percentage of FPTs had
45 patients on their roster from lower NIQ's than FPNTs (1-3) (77.7% (n = 3167) vs. 69.8%
46 (n = 5353), SMD = 0.18). FPTs had a larger mean number of patients from a rural
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 residence compared to FPNTs (139.28 (+/- 256.05) vs. 54.93 (+/- 137.27), SMD = 0.41).

4
5 A greater percentage of FPTs had more patients from rural areas than urban areas
6
7 (14.6% (N = 594) vs. 6.8% (n = 522), SMD = 0.25). FPNTs had a higher patient mean in
8
9 the Champlain LHIN (which includes Ottawa, a large urban region in Ontario) (FPT =
10
11 61.94 (+/- 236.36), FPTN = 99.43 (+/-287.94), SMD = 0.14). FPTs had higher patient
12
13 means in four LHINs (mostly northern regions): South East (SE) LHIN (FPT = 47.03 (+/-
14
15 209), FPTN = 24.27 (+/-132.92), SMD = 0.13), North Simcoe Muskoka (NSM) LHIN
16
17 (FPT = 59.28 (+/- 230.38), FPNT = 15.12 (+/- 97.56), SMD =0.25), North East (NE)
18
19 LHIN (FPT = 53.59 (+/- 228.57), FPNT = 19.86 (+/- 124.20), SMD =0.18), North West
20
21 (NW) LHIN (FPT = 24.64 (+/- 149.32), FPNT = 6.06 (+/- 101.30), SMD = 0.16).
22
23
24
25
26
27

28
29 The mean number of patients with psychosocial ADG's was higher for FPTs compared
30
31 to FPNTs (258.24 (+/- 204.4) vs. 217.56 (+/- 179.29), SMD = 0.21).The mean mental
32
33 health primary care provider (MHPCP) visits was higher in FPTs compared to FPNTs
34
35 (707.60 (+/- 2863.95) vs. 276.86 (+/- 532.56), SMD = 0.21). Due to the large standard
36
37 deviation, we also examined the median for MHPCP; FPTs had a higher median
38
39 MHPCP visits than FPNTs (229 (IQR=106-415) vs. 189 (IQR=110-350), SMD = 0.24).
40
41
42 FPTs have a higher percentage of patients with 415+ MHPCP visits compared to
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
FPNTs (25.2% (n = 1026) vs. 16.7% (n = 1285), SMD 0.21).

59
60
The mean number of ED visits was higher for FPTs compared to FPTNs (440.82 (+/-
357.7) vs. 291.96 (+/- 240.01), SMD = 0.49). Table 1 describes the difference between
all categories, but the most notable differences are for patients that had 60-156, 157-

1
2
3 257, and 415+ visits. FPTs had a lower percentage of patients with 60-156 ED visits
4 than FPNTs (9.5% (n = 387) vs. 17.2% (n = 1320), SMD = 0.23) and patients with 157-
5
6 257 visits (12.0% (n = 489) vs. 19.5% (n = 1495), SMD = 0.21). FPTs has a greater
7
8 percentage of patients with 415+ ED visits compared to FPNTs (45.6% (n = 1857) vs.
9
10 16.7% (n = 1285), SMD = 0.48).
11
12
13

14
15
16
17 FPTs had a larger mean number of MHED visits for rostered patients compared to
18
19 FPNTs (18.99 (+/- 29.07) vs. 10.91 (+/- 13.07), SMD = 0.36). Full details within each
20
21 category are in Table 1, but the most notable difference was the group that had 0-3 and
22
23 20+ MHED visits. FPTs had a lower percentage of patients with 0-3 MHED visits than
24
25 FPNTs (16.7% (n = 681) vs. 27.5% (n = 2108), SMD = 0.26) and a higher percentage of
26
27 patients with 20+ MHED visits than FPNTs (30.9% (n = 1258) vs. 14.1% (n = 1084),
28
29 SMD = 0.41).
30
31
32

33
34
35 FPTs had a greater mean of hospitalizations compared to FPNTs (57.05 (+/-47.43) vs.
36
37 10.91 (+/-13.07), SMD = 0.36). Full details within each category are in Table 1, but the
38
39 most notable difference was that FPTs had a larger percentage of patients with 11+
40
41 hospitalizations, compared to FPNTs (85.7% (n = 3491) vs. 78.8% (n = 6044), SMD =
42
43 0.18). Of those hospitalizations, mental health specific hospitalizations (MHH) were
44
45 measured. FPTs had a greater mean of MHH visits compared to FPNTs (7.01 (+/-13.61)
46
47 vs. 3.86 (+/-5.43), SMD = 0.36). The most striking differences were the groups that had
48
49 zero visits and those with 11+ visits. FPTs had a lower percentage of patients with zero
50
51 MHH visits compared to FPNTs (13.4% (n = 545) vs. 21.9% (n = 1679), SMD = 0.22).
52
53
54

1
2
3 FPTs had a higher percentage of patients with 11+ MHH visits compared to FPNTs
4
5 (17.8% (n = 725) vs. 6.8% (n = 518), SMD = 0.34).
6
7

8 **Statistical Modeling**

9 *Regression using Generalized Estimating Equations*

10
11
12 A full description of the findings is listed in Table 2. Within this model, a number of
13
14
15 significant correlates of FPTs were identified, including physician age group of 55-64
16
17 (OR=0.85, CI=0.74-0.97), physician age group of 64+ (OR=0.85, CI=0.62-0.85), and
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
FPTs were more likely to be located in rural areas (OR=1.38, CI=1.17-1.62). Compared
to patient rosters of FPNTs, FPT rostered patients were: almost half as likely to be
urban (OR=0.55, CI=0.43-.70), more likely to have psychiatric ADGs (OR=1.2, CI=0.86-
1.66), and more likely to have 4-9 mental health ED (MHED) visits (OR=1.33, CI=1.03-
1.72), patients with 10-19 MHED visits (OR=1.54 CI=1.14-2.07), patients with 20+
MHED visits (OR=2.17, CI=1.60-2.94), patients with 3-5 MHH (OR=1.21, CI=1.08-1.35),
patients with 6-10 MHH (OR=1.32, CI=1.15-1.51), patients with 11+ MHH (OR=1.77,
CI=1.28-2.43), patients with 157-257 mental health primary care (MHPC) visits
(OR=1.24, CI=1.00-1.54), patients with 258-414 MHPC visits (OR=1.38 CI=1.09-1.76),
patients with 415+ MHPC visits (OR=1.61, CI=01.33-1.94). [Insert Table 2].

50 **Discussion**

51
52
53
54
55
56
57
58
59
60
To our knowledge, this is the first study that investigates trends and predictors of family
physicians referring to telepsychiatry at a system-level. Between FY 2008 and FY 2016,

1
2
3 there was a 9-fold increase in the number of primary care physicians referring to
4
5 telepsychiatry and a nearly 15-fold increase in the number of primary care patients
6
7 referred to telepsychiatry. 32% of family physicians referred to telepsychiatry, while less
8
9 than 1% of patients rostered to FPTs utilized telepsychiatry. While patient adoption of
10
11 telepsychiatry continued to increase in Ontario during the study period, it still represents
12
13 a very small percentage of patients in Ontario. These visits represent, for the most part,
14
15 patients that were required to go to a healthcare site to access care and psychiatrists
16
17 that are required to use the provincial telemedicine platform. It is evident that there
18
19 remains a large opportunity to increase patient adoption. As discussed in the
20
21 introduction, it is likely that the recent Covid-19 related temporary policy changes to
22
23 virtual care will present an opportunity for significant growth and adoption of
24
25 telepsychiatry. This study provides an essential benchmark for future analyses,
26
27 comparing how these policy changes have shifted adoption patterns with patients and
28
29 family physician referrals.
30
31
32
33
34
35
36
37

38 Physicians who referred to telepsychiatry in FY 2008 were mainly from northern
39
40 regions, whereas in FY 2016, more referrals from southern regions emerged, many of
41
42 which still have rural areas within their catchments. Referral to telepsychiatry was more
43
44 evenly distributed throughout the province in FY 2016. While historically telepsychiatry
45
46 was used to support rural populations, in later years of the study, there were increasing
47
48 provincially funded telepsychiatry projects that aimed to increase adoption of
49
50 telepsychiatry within more populous regions. While it is important to create access and
51
52 choice for patients in both rural and urban areas, policy-makers and funders should
53
54
55
56
57
58
59
60

1
2
3 consider aspects of digital health equity⁴³, allowing the use of diverse secure virtual
4
5 platforms, funding home-based visits and establishing mechanisms to support the
6
7 appropriate distribution of virtual care throughout the province.
8
9

10
11
12 Past studies have suggested that physicians who were more isolated and were part of an integrated
13
14 health system had positive feelings about telemedicine, and that nurses were more enthusiastic about
15
16 telemedicine than physicians.^{25,36} As such, unsurprisingly, this study found that predictors of family
17
18 physician referrals to telepsychiatry include rurality of residence (for both provider and patient), team-
19
20 based practice models, nurse practitioners on the team, and clinical complexity of patient rosters.
21
22

23
24 This study has numerous strengths; it utilizes a large, multi-year, linked dataset of family
25
26 physician and patient data to help understand important characteristics that are
27
28 common in physicians who refer to telepsychiatry. Additionally, since this study uses
29
30 administrative data, the study was conducted rapidly using a retrospective analysis.
31
32 However, the study also has several limitations. The identification of telepsychiatry visits
33
34 is contingent upon psychiatrists using telepsychiatry billing codes and billing fee for
35
36 service, so it is possible that there is missing data, or an underrepresentation of visits
37
38 completed by physicians paid by salary or alternate funding. Another potential
39
40 limitation, is that while this study includes nurse practitioners as a covariate in the
41
42 analysis, it does not take into account actual nurse practitioner referrals to
43
44 telepsychiatry, which could lead to an underestimate of our total telepsychiatry visits.
45
46 We believe, however, that this will have minimal impact on our study because it is likely
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 that nurse practitioner referrals will be captured within total telepsychiatry visits, but
4
5 rostered (as defined in our methods) to a family physician instead.
6
7
8
9

10 **Conclusion**

11
12 While telepsychiatry has seen continued growth over recent years, there remains
13
14 significant opportunities for increased adoption. This study shows that certain factors,
15
16 including rurality, provider practice type, the number of nurse practitioners on the
17
18 provider's team, as well as patient healthcare and hospital utilization characteristics are
19
20 more likely in FPTs than FPNTs. This study provides a baseline from which to assess
21
22 recent rapid mobilization of telepsychiatry, and a means of doing so using administrative
23
24 health data.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Tables and Figures

Figure 1. Telepsychiatry Adoption by Year (FY2008 and FY2016)




Figure 2. Percentage of Referring Physicians by Region (2008 and 2016)




Table 1. Comparative Analysis of Family Physicians that Refer to Telepsychiatry compared to (2016)

VARIABLE	VALUE	FPT	FPNT	TOTAL	SMD
		N = 4,074	N = 7,674	N = 11,748	
PHYSICIAN CHARACTERISTICS					
Physician Sex	F	1,647 (40.4%)	3,599 (46.9%)	5,246 (44.7%)	0.13
	M	2,427 (59.6%)	4,075 (53.1%)	6,502 (55.3%)	0.13
< 10,000 population in CMA	N	3,466 (85.1%)	7,041 (91.8%)	10,507 (89.4%)	0.21
	Y	608 (14.9%)	633 (8.2%)	1,241 (10.6%)	0.21
Nurse Practitioners on Team	Mean ± SD	1.71 ± 4.02	1.10 ± 3.70	1.31 ± 3.83	0.16
Physician Practice Type	FHT	1,145 (28.1%)	1,381 (18.0%)	2,526 (21.5%)	0.24
	Others	2,929 (71.9%)	6,293 (82.0%)	9,222 (78.5%)	0.24
PATIENT ROSTER CHARACTERISTICS					
Total # of attached patients	Mean ± SD	862.45 ± 668.36	766.97 ± 605.70	800.64 ± 630.13	0.15
# patients in NIQ 1,2&3 > # patients in NIQ 4&5	No	907 (22.3%)	2,127 (27.7%)	3,034 (25.8%)	0.13
	Yes	3,167 (77.7%)	5,353 (69.8%)	8,520 (72.5%)	0.18
Rural Residence	Mean ± SD	139.28 ± 256.05	54.93 ± 137.27	84.67 ± 192.19	0.41
# of patients in urban residence > # of patients in rural residence	No	594 (14.6%)	522 (6.8%)	1,116 (9.5%)	0.25
	Yes	3,480 (85.4%)	6,958 (90.7%)	10,438 (88.8%)	0.16
South East LHIN	Mean ± SD	47.03 ± 209.09	24.27 ± 132.92	32.29 ± 164.22	0.13
Champlain LHIN	Mean ± SD	61.94 ± 263.36	99.43 ± 287.94	86.21 ± 280.09	0.14
North Simcoe Muskoka LHIN	Mean ± SD	59.28 ± 230.38	15.12 ± 97.56	30.69 ± 159.11	0.25
North East LHIN	Mean ± SD	53.59 ± 228.57	19.86 ± 124.20	31.75 ± 169.31	0.18
North West LHIN	Mean ± SD	24.64 ± 149.32	6.06 ± 59.90	12.61 ± 101.30	0.16
# of patients with a psychosocial ADG	Mean ± SD	258.24 ± 204.40	217.56 ± 179.29	231.91 ± 189.52	0.21
Total MH PCP visits	Mean ± SD	707.60 ± 2,863.95	276.86 ± 532.56	428.74 ± 1,765.68	0.21
	Median (IQR)	239 (106-415)	188 (73-335)	204 (82-364)	0.24
	00-59	686 (16.8%)	1,626 (21.2%)	2,312 (19.7%)	0.11
	60-156	702 (17.2%)	1,612 (21.0%)	2,314 (19.7%)	0.1
157-257	795 (19.5%)	1,512 (19.7%)	2,307 (19.6%)	0	

	258-414	865 (21.2%)	1,445 (18.8%)	2,310 (19.7%)	0.06
	415-high	1,026 (25.2%)	1,285 (16.7%)	2,311 (19.7%)	0.21
VARIABLE	VALUE	FPT	FPNT	TOTAL	SMD
Total ED visits	Mean ± SD	440.82 ± 357.70	291.96 ± 240.01	344.45 ± 295.74	0.49
	00-59	493 (12.1%)	1,090 (14.2%)	1,583 (13.5%)	0.06
	60-156	387 (9.5%)	1,320 (17.2%)	1,707 (14.5%)	0.23
	157-257	489 (12.0%)	1,495 (19.5%)	1,984 (16.9%)	0.21
	258-414	848 (20.8%)	1,782 (23.2%)	2,630 (22.4%)	0.06
	415-high	1,857 (45.6%)	1,793 (23.4%)	3,650 (31.1%)	0.48
Total MH ED visits	Mean ± SD	18.99 ± 29.07	10.91 ± 13.07	13.75 ± 20.58	0.36
	00-03	681 (16.7%)	2,108 (27.5%)	2,789 (23.7%)	0.26
	04-09	922 (22.6%)	2,275 (29.6%)	3,197 (27.2%)	0.16
	10-19	1,213 (29.8%)	2,013 (26.2%)	3,226 (27.5%)	0.08
	20-high	1,258 (30.9%)	1,084 (14.1%)	2,342 (19.9%)	0.41
Total hospitalizations	Mean ± SD	57.05 ± 47.43	41.75 ± 36.01	47.15 ± 41.06	0.36
	0 visits	116 (2.8%)	286 (3.7%)	402 (3.4%)	0.05
	01-02 visit	133 (3.3%)	287 (3.7%)	420 (3.6%)	0.03
	03-05 visits	131 (3.2%)	374 (4.9%)	505 (4.3%)	0.08
	06-10 visits	203 (5.0%)	489 (6.4%)	692 (5.9%)	0.06
	11+ visits	3,491 (85.7%)	6,044 (78.8%)	9,535 (81.2%)	0.18
Total MH hospitalizations	Mean ± SD	7.01 ± 13.61	3.86 ± 5.43	4.97 ± 9.31	0.3
	0 visits	545 (13.4%)	1,679 (21.9%)	2,224 (18.9%)	0.22
	01-02 visit	768 (18.9%)	2,078 (27.1%)	2,846 (24.2%)	0.2
	03-05 visits	1,072 (26.3%)	1,959 (25.5%)	3,031 (25.8%)	0.02
	06-10 visits	964 (23.7%)	1,246 (16.2%)	2,210 (18.8%)	0.19
	11+ visits	725 (17.8%)	518 (6.8%)	1,243 (10.6%)	0.34

NIQ= Neighbourhood Income Quintiles
 LHIN=Local Health Integration Networks
 ADG= Aggregated Diagnosis Groupings
 MHPCP= Mental Health Primary Care Practitioner
 ED= Emergency Department
 MH ED= Mental Health Emergency Department
 MH= Mental Health

Table 2. Model using Generalized Estimating Equations comparing FPTs vs FPNTs (2016)

Analysis Of GEE Parameter Estimates			
	Estimate	95% Confidence Limits	
Intercept	-0.9162	-1.1901	-0.6423
Physician Age 35-44 vs <=34	0.8895	0.7855	1.0072
Physician Age 45-54 vs <=34	0.9082	0.7983	1.0333
Physician Age 55-64 vs <=34	0.849	0.7419	0.9717
Physician Age 64+ vs <=34	0.7255	0.6223	0.8458
Sex (Male vs Female)	1.0144	0.9368	1.0983
Physician Rural (Yes vs No)	1.3791	1.174	1.6199
Canadian Medical Graduate (Yes vs No)	1.0333	0.9493	1.1247
Canadian Medical Graduate (Unknown vs No)	1.0231	0.8617	1.2148
More patients in low/med income quintile (Yes vs No)	1.0719	0.969	1.1856
More urban patients (Yes vs No)	0.5491	0.4338	0.6952
Mental Health ED Visits (04-09 vs 00-03)	1.3289	1.0252	1.7226
Mental Health ED Visits (10-19 vs 00-03)	1.5373	1.1442	2.0654
Mental Health ED Visits (20-high vs 00-03)	2.1678	1.601	2.9352
Mental Health Hospitalizations (0 visits vs 01-02 visit)	0.9072	0.7363	1.1179
Mental Health Hospitalizations 03-05 visits vs 01-02 visit	1.2048	1.0749	1.3504
Mental Health Hospitalizations 06-10 visits vs 01-02 visit	1.316	1.1496	1.5064
Mental Health Hospitalizations 11+ visits vs 01-02 visit	1.7662	1.2837	2.4301
Mental Health Primary Care Visits (60-156 vs 00-59)	0.965	0.8134	1.1449
Mental Health Primary Care Visits (157-257 vs 00-59)	1.237	0.9943	1.539
Mental Health Primary Care Visits (258-414 vs 00-59)	1.384	1.0859	1.7638
Mental Health Primary Care Visits (415-high vs 00-59)	1.6089	1.3322	1.943
More patients with psychiatric ADGs (Yes vs No)	1.1979	0.8627	1.6634

ED= Emergency Department

ADG= Aggregated Diagnosis Groups

References

1. Brien S, Grenier L, Kapral M. *Taking stock*. Health Quality Ontario; 2015.
2. Bishop TF, Press MJ, Keyhani S, Pincus HA. Acceptance of insurance by psychiatrists and the implications for access to mental health care. *JAMA psychiatry*. 2014;71(2):176-181.
3. World Health Organization. Mental disorders affect one in four people. *World Health Report*. 2001.
4. Alegría M, Nakash O, NeMoyer A. Increasing equity in access to mental health care: A critical first step in improving service quality. *World Psychiatry*. 2018;17(1):43.
5. Wang PS, Aguilar-Gaxiola S, Alonso J, et al. Use of mental health services for anxiety, mood, and substance disorders in 17 countries in the WHO world mental health surveys. *The Lancet*. 2007;370(9590):841-850.
6. Goldner EM, Jones W, Fang ML. Access to and waiting time for psychiatrist services in a canadian urban area: A study in real time. *The Canadian Journal of Psychiatry*. 2011;56(8):474-480.
7. Meadows GN, Enticott JC, Inder B, Russell GM, Gurr R. Better access to mental health care and the failure of the medicare principle of universality. *Med J Aust*. 2015;202(4):190-194.

- 1
2
3 8. Cunningham PJ. Beyond parity: Primary care physicians' perspectives on access to
4 mental health care: More PCPs have trouble obtaining mental health services for their
5 patients than have problems getting other specialty services. *Health Aff.*
6
7
8 2009;28(Suppl1):w490-w501.
9
- 10
11
12
13 9. Kurdyak P, Zaheer J, Cheng J, Rudoler D, Mulsant BH. Changes in characteristics
14 and practice patterns of ontario psychiatrists: Implications for access to psychiatrists.
15
16
17
18 *The Canadian Journal of Psychiatry.* 2017;62(1):40-47.
19
- 20
21
22 10. Kurdyak P, Stukel TA, Goldbloom D, Kopp A, Zagorski BM, Mulsant BH. Universal
23 coverage without universal access: A study of psychiatrist supply and practice patterns
24 in ontario. *Open Medicine.* 2014;8(3):e87.
25
26
27
28
- 29
30 11. Myers KM, Valentine JM, Melzer SM. Child and adolescent telepsychiatry:
31 Utilization and satisfaction. *Telemedicine and e-Health.* 2008;14(2):131-137.
32
33
34
- 35
36 12. Antonacci DJ, Bloch RM, Saeed SA, Yildirim Y, Talley J. Empirical evidence on the
37 use and effectiveness of telepsychiatry via videoconferencing: Implications for forensic
38 and correctional psychiatry. *Behav Sci Law.* 2008;26(3):253-269.
39
40
41
42
- 43
44 13. Campbell R, O'Gorman J, Cernovsky ZZ. Reactions of psychiatric patients to
45 telepsychiatry. *Mental illness.* 2015;7(2).
46
47
48
- 49
50 14. Deslich S, Stec B, Tomblin S, Coustasse A. Telepsychiatry in the 21st century:
51 Transforming healthcare with technology. *Perspectives in Health Information*
52
53
54
55

1
2
3 *Management/AHIMA, American Health Information Management Association.*

4
5 2013;10(Summer).

6
7
8
9 15. Fleming DA, Edison KE, Pak H. Telehealth ethics. *Telemedicine and e-Health.*

10
11 2009;15(8):797-803.

12
13
14 16. Saeed SA, Diamond J, Bloch RM. Use of telepsychiatry to improve care for people
15 with mental illness in rural North Carolina. *NC Med J.* 2011;72(3):219-222.

16
17
18 17. Ulzen T, Williamson L, Foster PP, Parris-Barnes K. The evolution of a community-
19 based telepsychiatry program in rural alabama: Lessons learned—a brief report.

20
21
22 *Community Ment Health J.* 2013;49(1):101-105.

23
24
25 18. Lyketsos CG, Roques C, Hovanec L, Jones III BN. Telemedicine use and the
26 reduction of psychiatric admissions from a long-term care facility. *J Geriatr Psychiatry*
27 *Neurol.* 2001;14(2):76-79.

28
29
30 19. Hilty DM, Ferrer DC, Parish MB, Johnston B, Callahan EJ, Yellowlees PM. The
31 effectiveness of telemental health: A 2013 review. *Telemedicine and e-Health.*

32
33 2013;19(6):444-454.

34
35
36 20. Chakrabarti S. Usefulness of telepsychiatry: A critical evaluation of

37
38
39 videoconferencing-based approaches. *World journal of psychiatry.* 2015;5(3):286.

- 1
2
3 21. Bishop JE, O'Reilly RL, Maddox K, Hutchinson LJ. Client satisfaction in a feasibility
4 study comparing face-to-face interviews with telepsychiatry. *J Telemed Telecare*.
5
6 2002;8(4):217-221.
7
8
9
10
11 22. Frueh BC, Monnier J, Yim E, Grubaugh AL, Hamner MB, Knapp RG. A randomized
12 trial of telepsychiatry for post-traumatic stress disorder. *J Telemed Telecare*.
13
14 2007;13(3):142-147.
15
16
17
18
19 23. Farabee D, Calhoun S, Veliz R. An experimental comparison of telepsychiatry and
20 conventional psychiatry for parolees. *Psychiatric Services*. 2016;67(5):562-565.
21
22
23
24
25 24. Bashshur RL, Shannon GW, Bashshur N, Yellowlees PM. The empirical evidence
26 for telemedicine interventions in mental disorders. *Telemedicine and e-Health*.
27
28 2016;22(2):87-113.
29
30
31
32
33 25. Sargeant J, Allen M, Langille D. Physician perceptions of the effect of telemedicine
34 on rural retention and recruitment. *J Telemed Telecare*. 2004;10(2):89-93.
35
36
37
38
39 26. Swanson B. Information technology and under-served communities. *J Telemed*
40 *Telecare*. 1999;5(2_suppl):3-10.
41
42
43
44
45 27. Serhal E, Crawford A, Cheng J, Kurdyak P. Implementation and utilisation of
46 telepsychiatry in Ontario: A population-based study. *The Canadian Journal of*
47 *Psychiatry*. 2017;62(10):716-725.
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 28. Grumbach K, Selby JV, Damberg C, et al. Resolving the gatekeeper conundrum:
4
5 What patients value in primary care and referrals to specialists. *JAMA*.
6
7 1999;282(3):261-266.
8
9
10
11 29. Norton PG, Dunn EV, Bestvater D. Referrals in primary care: Is the family physician
12
13 a "Gatekeeper"? *Canadian Family Physician*. 1989;35:1776.
14
15
16
17 30. Steele M, Zayed R, Davidson B, et al. Referral patterns and training needs in
18
19 psychiatry among primary care physicians in canadian rural/remote areas. *Journal of*
20
21 *the Canadian Academy of Child and Adolescent Psychiatry*. 2012;21(2):111.
22
23
24
25 31. Spaulding RJ, Russo T, Cook DJ, Doolittle GC. Diffusion theory and telemedicine
26
27 adoption by kansas health-care providers: Critical factors in telemedicine adoption for
28
29 improved patient access. *J Telemed Telecare*. 2005;11(1_suppl):107-109.
30
31
32
33 32. Whitten P, Franken EA. Telemedicine for patient consultation: Factors affecting use
34
35 by rural primary-care physicians in kansas. *J Telemed Telecare*. 1995;1(3):139-144.
36
37
38
39 33. LeRouge C, Garfield M. Crossing the telemedicine chasm: Have the US barriers to
40
41 widespread adoption of telemedicine been significantly reduced? *International journal of*
42
43 *environmental research and public health*. 2013;10(12):6472-6484.
44
45
46
47 34. Moore MA, Coffman M, Jetty A, Klink K, Petterson S, Bazemore A. Family
48
49 physicians report considerable interest in, but limited use of, telehealth services. *The*
50
51 *Journal of the American Board of Family Medicine*. 2017;30(3):320-330.
52
53
54
55
56
57
58
59
60

- 1
2
3 35. Spaulding R, Cain S, Sonnenschein K. Urban telepsychiatry: Uncommon service for
4 a common need. *Child and Adolescent Psychiatric Clinics*. 2011;20(1):29-39.
5
6
7
8
9 36. Whitten PS, Mackert MS. Addressing telehealth's foremost barrier: Provider as initial
10 gatekeeper. *Int J Technol Assess Health Care*. 2005;21(4):517-521.
11
12
13
14 37. Steele LS, Glazier RH, Lin E, Evans M. Using administrative data to measure
15 ambulatory mental health service provision in primary care. *Med Care*. 2004:960-965.
16
17
18
19 38. Austin PC, van Walraven C, Wodchis WP, Newman A, Anderson GM. Using the
20 Johns Hopkins aggregated diagnosis groups (ADGs) to predict mortality in a general
21 adult population cohort in Ontario, Canada. *Med Care*. 2011;49(10):932.
22
23
24
25
26
27
28 39. Cohen J. *Statistical power analysis for the behavioral sciences*. Routledge; 2013.
29
30
31
32 40. Austin PC. Using the standardized difference to compare the prevalence of a binary
33 variable between two groups in observational research. *Communications in Statistics-
34 Simulation and Computation*. 2009;38(6):1228-1234.
35
36
37
38
39
40 41. Wang M. Generalized estimating equations in longitudinal data analysis: A review
41 and recent developments. *Advances in Statistics*. 2014;2014.
42
43
44
45
46 42. The Ontario Physician Human Resources Data Centre. . (2018, 10). Physicians in
47 Ontario 2017. Retrieved from the Ontario Physician Human Resources data centre:
48 [Http://Www.ophrdc.org/wp-content/uploads/2018/10/PIO-2017-report-with-](http://www.ophrdc.org/wp-content/uploads/2018/10/PIO-2017-report-with-hyperlinks.pdf)
49 [hyperlinks.pdf](http://www.ophrdc.org/wp-content/uploads/2018/10/PIO-2017-report-with-hyperlinks.pdf). *McMaster University*. 2018.
50
51
52
53
54
55
56
57
58
59
60

1
2
3 43. Crawford, A., & Serhal, E. (2020). Digital Health Equity and COVID-19: The
4
5 Innovation Curve Cannot Reinforce the Social Gradient of Health. *Journal of Medical*
6
7 *Internet Research*, 22(6), e19361.
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


Characterizing Family Physicians Who Refer to Telepsychiatry in Ontario

Figure Legends

Figure 1. Telepsychiatry Adoption by Year (FY2008 and FY2016)


Figure 2. Percentage of Referring Physicians by Region (FY 2008 and FY2016)

Figure 1. Telepsychiatry Adoption by Year (FY2008 and FY2016)



Characterizing Family Physicians Who Refer to Telepsychiatry in Ontario

Figure 2. Percentage of Referring Physicians by Region (2008 and 2016)



CARACTÉRISATION DES MÉDECINS DE FAMILLE QUI RÉFÈRENT À LA TÉLÉPSYCHIATRIE EN ONTARIO

Introduction : La télépsychiatrie peut améliorer l'accès aux services psychiatriques pour ceux qui autrement ne peuvent accéder aux soins facilement. Les médecins de famille sont les gardiens des soins spécialisés en Ontario, alors il est essentiel de comprendre les prédicteurs en rapport avec les références à la télépsychiatrie pour mieux planifier les services et stimuler l'adoption de la télépsychiatrie.

Méthodes : La présente étude a utilisé une méthodologie d'étude transversale rétrospective annuelle pour comparer les médecins qui référaient leurs patients en télépsychiatrie chaque année de l'exercice financier (EF) de 2008 à l'EF de 2016. Une comparaison d'un an (EF 2016) des médecins de famille qui référaient en télépsychiatrie avec les médecins de famille qui ne référaient pas en télépsychiatrie, appariés (1:2) par région, a aussi été menée. Enfin, nous avons utilisé une modélisation statistique pour comprendre les prédicteurs de la référence en télépsychiatrie chez les médecins.

1
2
3 **Résultats** : Entre l'EF 2008 et l'EF 2016, les visites en télépsychiatrie sont passées de
4
5 925 à 13 825 visites. Trente-deux pour cent des médecins de première ligne en Ontario
6
7 ont référé à la télépsychiatrie en 2017. Plusieurs caractéristiques étaient notablement
8
9 différentes entre les médecins de famille qui réfèrent en télépsychiatrie (MFT) et les
10
11 médecins de famille qui ne réfèrent pas en télépsychiatrie (MFNT) : les MFT étaient
12
13 plus susceptibles de provenir d'une résidence de moins de 10 000 personnes, d'avoir
14
15 plus d'infirmières praticiennes (IP) dans leur pratique, et de faire partie d'une équipe de
16
17 santé familiale (ESF) que les MFNT. Les patients inscrits étaient plus susceptibles de
18
19 résider en région rurale, de présenter plus de complexité clinique, et d'utiliser plus de
20
21 services de santé mentale comparativement aux MFNT.
22
23
24
25

26
27 **Conclusion** : Il y a eu une augmentation de l'utilisation de la télépsychiatrie par les
28
29 patients et les médecins de famille durant la période de l'étude, bien qu'il demeure une
30
31 possibilité de croissance significative. Les médecins de famille qui habitent en milieu
32
33 rural, qui font partie d'une ESF, qui ont plus d'IP, dont les patients sont davantage
34
35 complexes et en milieu rural étaient plus susceptibles de référer en télépsychiatrie.
36
37 Comme de récentes politiques en faveur de la télépsychiatrie en soutiennent la
38
39 croissance, la présente étude servira de base de départ importante.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


Characterizing Family Physicians Who Refer to Telepsychiatry in Ontario

Figure Legends

Figure 1. Telepsychiatry Adoption by Year (FY2008 and FY2016)


Figure 2. Percentage of Referring Physicians by Region (FY 2008 and FY2016)

Figure 1. Telepsychiatry Adoption by Year (FY2008 and FY2016)



Characterizing Family Physicians Who Refer to Telepsychiatry in Ontario

Figure 2. Percentage of Referring Physicians by Region (2008 and 2016)



CHARACTERIZING FAMILY PHYSICIANS WHO REFER TO TELEPSYCHIATRY IN ONTARIO

Authors:

Eva Serhal^{1,2}, Tomisin Iwajomo^{1,3}, Claire de Oliveira¹⁻³, Allison Crawford^{1,4}, Paul

Kurdyak¹⁻³

Affiliations

1. The Centre for Addiction and Mental Health, Toronto, Ontario
2. The Institute for Health Policy, Management and Evaluation, University of Toronto, Toronto, Ontario
3. ICES, Toronto, Ontario
4. Dalla Lana School of Public Health, University of Toronto, Toronto, Ontario

Corresponding Author:

Eva Serhal, MBA, PhD
Director, Virtual Mental Health and Outreach & ECHO Ontario
Centre for Addiction and Mental Health
821-250 College Street, Toronto, Ontario
M5T1R8

Abstract

Introduction

Telepsychiatry can improve access to psychiatric services for those who otherwise cannot easily access care. Family physicians are gatekeepers to specialized care in Ontario, so it is essential to understand predictors relating to referrals to telepsychiatry to better plan services and increase telepsychiatry adoption.

Methods

This study used an annual retrospective cross-sectional study design to compare physicians who referred their patients to telepsychiatry each year from fiscal year (FY) 2008 to FY 2016. A one year (FY 2016) comparison of family physicians who referred to telepsychiatry compared to family physicians who did not refer to telepsychiatry matched (1:2) by region was also conducted. Finally, we used statistical modelling to understand the predictors of referring to telepsychiatry among physicians.

Results

Between FY 2008 to FY 2016, telepsychiatry visits increased from 925 visits to 13,825 visits. Thirty-two percent of Ontario primary care physicians referred to telepsychiatry in 2017. Several characteristics were notably different between family physicians who refer to telepsychiatry (FPTs) and family physicians who do not refer to telepsychiatry (FPNTs): FPTs were more likely to be from a residence with less than 10,000 people, to have more nurse practitioners in the practice, and to be from a family health team than

1
2
3 FPNTs. Rostered patients of FPTs were more likely to reside in rural areas, have more
4
5 clinical complexity, and to utilize more mental health services compared to FPNTs.
6
7

8 **Conclusion**

9
10
11
12 There has been an increase in the use of telepsychiatry by patients and family
13
14 physicians over the study period, although there remains opportunity for significant
15
16 growth. and familyFamily physicians that ~~are~~ live in ~~a~~ rural areas, are part of a FHT,
17
18 have more NPs, with more rural and complex patients ~~patients in rural areas with high~~
19
20 ~~needs for mental health services are~~ were more likely to refer to telepsychiatry. As
21
22 recent pro-telemedicine policies support the growth of telepsychiatry, this study will
23
24 serve as an important baseline.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

Introduction

Mental illness is among the world's leading causes of sickness and disability; one in five Canadians (two million Ontarians) and one in four people internationally will experience mental illness or addiction ~~at some point in their life, and one in four people internationally will be affected by a mental or neurological disorder.~~¹⁻³ However, only one-third of Ontario patients ~~requiring mental health and addiction services~~ reported having their mental health needs met or partially met within primary care.¹ Patients ~~Similar international studies found that at least two thirds of people with common mental disorders do not receive treatment and patients~~ from lower-income countries and socially disadvantaged groups had lower rates of mental health service utilization.⁴⁻⁶ Access to quality mental health services is ~~limited in many parts of the world, including Canada often limited,~~ due to rurality, physician shortages and variation in the availability of mental health services ~~from primary care providers and outpatient psychiatrists.~~^{1,5,7-10}

Telepsychiatry, ~~which is~~ the delivery of psychiatry via secure videoconferencing, can reduce access barriers such as distance, cost of travel, weather, and workforce shortages.¹¹⁻¹⁷ In particular, telepsychiatry can provide access to outpatient psychiatric care for patients who otherwise are not able to easily access services, potentially reducing unnecessary hospitalizations and ED visits.^{18,19} Several studies have deemed telepsychiatry effective for various mental health conditions and populations.²⁰⁻²⁴ Telemedicine has been shown to reduce isolation and increase the morale of rural primary care practitioners.^{25,26} While there is significant potential for telepsychiatry to

1
2
3 connect patients to psychiatrists throughout Ontario, recent evidence has shown
4 minimal adoption by patients.²⁷ In Ontario during the time of this study, in order to
5 deliver and bill for telepsychiatry, psychiatrists were typically required to use a
6 centralized government funded telemedicine platform ~~the Ontario Telemedicine~~
7 Network (OTN) platform (a centralized, government funded platform) to deliver care.
8 Additionally, other than small pilot tests, and patients were typically required to access
9 their telepsychiatry visit at a patient host site (a clinical site that ~~could connect to the~~
10 OTN-connects to the centralized provincial network). It is important to note that at the
11 time this paper was submitted ~~there were significant~~ temporary policy changes to virtual
12 care as a result of Covid-19 (introduced March 2020), ~~which~~ have created more options
13 for home-based care for patients, and ~~have allowed physicians to deliver virtual care via~~
14 a variety of virtual platforms. ~~have removed the requirement that physicians must be~~
15 members of the OTN network, and use the OTN platforms to deliver and bill for virtual
16 care. These recent changes will be discussed in relation to the study findings in the
17 discussion section.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 Family physicians are the gatekeepers to specialist care in Canada and other parts of
41 the world.^{28,29} Common reasons for family physician referral to psychiatry include
42 obtaining medication recommendations, obtaining second opinions, or ~~for patients who~~
43 are non-responsive to medications, getting a second opinion on a diagnosis, and
44 obtaining non-pharmacological treatment recommendations.³⁰ Evidence shows
45 providers who are more isolated had more positive feelings about telemedicine.²⁵
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Positive perceptions of telemedicine included convenience, perceived patient benefits,
4 reduced wait time, cost-effectiveness, and access to medical expertise,²⁵ while negative
5 aspects included cost, reduced funding for in-person services, inconvenience ~~to~~
6 ~~doctors~~, inadequate reimbursement, ~~inadequate~~ and training, medical liability, ~~issues~~
7 ~~with referral patterns~~, and physician fear of practice change.^{25,31-34} Previous studies
8 suggest nurses ~~are more~~ may be more enthusiastic about telemedicine than physicians,
9 referring physicians ~~are more~~ tend to be more satisfied than ~~the referred~~ specialists,
10 satisfaction increases relative to use and time, and that providers do not recommend
11 telemedicine in all circumstances.^{29,35,36} ~~Understanding if there are differences in family~~
12 ~~physicians who refer to telepsychiatry, compared to those who do not, is an important~~
13 ~~step to understanding how telepsychiatry is currently being used, and how it can be~~
14 ~~deployed more effectively throughout the province.~~ The objective of this study was to
15 examine the characteristics of family physicians in Ontario who have referred patients to
16 telepsychiatry (FPTs) compared to family physicians who have not referred to
17 telepsychiatry (FPNTs) from FY 2008 to FY 2016, in order to understand if there are
18 specific factors that make certain family physicians more likely to refer than others.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

42 **Methods**

43 ***Setting and Study Design***

44
45
46 We conducted an annual retrospective cross-sectional study assessing descriptive
47 characteristics of FPTs for FYs (2008 to 2016); this time horizon was used because it
48
49
50
51
52
53
54

1
2
3 includes data from when the Ontario Health Insurance Plan (OHIP) physician fee
4 schedule telemedicine codes were introduced until the most recent data available from
5 ICES. Referring physician and patient information were ascertained using the OHIP
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

includes data from when the Ontario Health Insurance Plan (OHIP) physician fee schedule telemedicine codes were introduced until the most recent data available from ICES. Referring physician and patient information were ascertained using the OHIP claims database; if referral information was not available, we looked back within the past 12 months to identify the most frequently seen family physician and, if not available, the most recently seen family physician. These data were linked to population-level health administrative databases held at ICES. Additionally, a one year (FY2016) comparison of FPTs vs FPNTs matched (1:2) by the 14 Local Health Integration Networks ~~(the region in which they work) was conducted~~(LHIN). [LHINs are artificially divided regions of Ontario which were designed to support local approaches to strategic planning, funding and regional coordination of clinical care, and are used in administrative data to classify provincial regions.](#) ~~We matched by these 14 LHINs only, and not age or sex because we wanted to ensure we were able to explore differences in age and sex between the groups.~~ This study was approved by the Research Ethics Board of Sunnybrook Health Sciences Centre, Toronto.

Data Sources

This study used province-wide health administrative databases that were linked using unique encoded identifiers and analyzed at ICES. OHIP data were used to collect family physician details, including physician number, fee code, service date, and location of service provided, and patient information, including patient ID, diagnosis, service date, and location. The National Ambulatory Care Reporting System was used to collect data on emergency department visits. The Ontario Mental Health Reporting System was

1
2
3 used to obtain information on psychiatric hospitalizations. The ICES Physician
4
5 Database and the Primary Care Population database were used to analyze data on
6
7 primary care providers and teams, including the number of years since medical school
8
9 graduation, years of practice since graduation, provider type, interprofessional team
10
11 details, and total physicians per practice. The Client Agency Program Enrolment
12
13 database was used to analyze data on physician practice type (defined as patient
14
15 enrollment type). The Ontario Census Area Profiles and the Registered Persons
16
17 Database were used to analyze patient-level demographic variables (such as age, sex,
18
19 the region of residence), as well as geographic details (rurality, income quintiles).
20
21 Primary care providers who had missing age or sex data, and patients with missing
22
23 health card numbers, missing age or missing sex were excluded from the analysis.
24
25
26
27
28

29 ***Provider Characteristics***

30
31
32 We selected primary care provider characteristics based on two things: availability
33
34 within the ICES dataset, and previous literature within the field that assessed predictors
35
36 using these measures.³⁴ The primary care provider characteristics used to compare
37
38 FPTs vs FPNTs included age group, sex, number of years since medical school
39
40 graduation, years of practice since graduation, rurality of residence, physician practice
41
42 type (Family Health Team¹ or other), average number of nurse practitioners within the
43
44 practice, and average number of social workers within the practice.
45
46
47
48
49

50
51 ¹ Family health teams are different from many primary care practices because they are funded to include an
52
53 interprofessional team within the practice, and are also paid via a blended capitation model (where physicians are
54
55 paid primarily through capitation fees based on a defined basket of services delivered to enrolled patients) or a
56
57 salary model).¹⁶³
58
59
60

Patient Characteristics

We examined patient demographic and service utilization characteristics for patients rostered within each physician group (FPT vs FPNT). Patient demographic characteristics included age, sex, neighbourhood income quintile (NIQ) (NIQ 1= 0-20%, NIQ 2= 20-40%, NIQ 3= 40-60%, NIQ 4= 60-80%, NIQ 5= 80%+), and whether they live in a rural or urban area. Patient utilization and medical complexity were considered within each annual cross-section by assessing: number of primary care provider visits (total, mental health and non-mental health); number of emergency department visits (total, mental health and non-mental health-related); number of hospitalizations (total, mental health and non-mental health-related); and comorbidities using the Johns Hopkins' Aggregated Diagnosis Groups (ADG). [Mental health primary care visits were defined using a validated algorithm to identify adults accessing care for a mental health disorder, which included adult mental health diagnostic codes 295-298, 300-304, 306, 309, 311, 897-902, 904, 905 \(which represents a broad range of adult mental health and addiction diagnoses as defined by the International Classification of Diseases 10 \(ICD-10-CA\)\).](#) ~~Mental health primary care visits were defined using a validated algorithm, and included mental health diagnostic codes 295-298, 300-304, 306, 309, 311, 897-902, 904, 905.~~³⁷ Hospitalizations and ED visits included [International Classification of Diseases 10 \(ICD-10-CA\)](#) codes F04-F99 (which excludes dementia). The Johns Hopkins ADG's are categorized based on five clinical dimensions, including the duration, the severity, the etiology, the diagnostic certainty and the specialty care

1
2
3 involvement of the condition. ~~Study subjects may have diagnoses ranging between 0-32~~
4
5 ~~ADGs.~~³⁸
6
7

8 9 **Analysis**

10 11 *Descriptive Analysis*

12 13 Annual Outputs and Trends over Time

14
15 Annual panels containing physician and patient characteristics were described,
16
17 including counts, column percentages, and means with standard deviations.
18
19

20
21 Additionally, changes over time were evaluated for each of the physician characteristics
22
23 ~~to assess if there were changing trends in physicians using telepsychiatry throughout~~
24
25 ~~our study period.~~
26
27
28
29

30 31 Matched Physician Comparison

32
33 ~~A one-year (FY 2016) comparison of FPTs vs FPNTs to compare between-group~~
34
35 ~~differences was conducted by matching (1:2) based on the fourteen LHINs (the region~~
36
37 ~~in which they worked).~~ Matched comparisons between FTPs and FTNPs were
38
39 presented as counts, column percentages, and standardized mean differences (SMD).
40
41 We compared between-group differences using the standard mean difference (SMD).
42
43 Cohen (1988) suggests a small effect size is equal to 0.2 SMD, but recent studies
44
45 suggest that a more prudent threshold for assessing effect size is 0.1 SMD, especially
46
47 with large data sets.^{39,40} Using this threshold (0.1 SMD), this study identifies significant
48
49 differences between groups.
50
51
52
53
54
55
56
57
58
59
60

1
2
3 The model covariates included physician age (categorical; <34, 35-44, 45-54, 55-64,
4 64+), sex (Male/Female), number of years after graduation (categorical; 1-10, 11-20,
5 21-30, 31-40, 41+ years), rural residence (living in community with less than 10,000
6 people- Yes/No), total nurse practitioner on the team (continuous), total social workers
7 on the team (continuous), if the family physician is a Canadian Medical Graduate
8 (Yes/No), the physician practice type (FHT/Other), patient age (categorical; 0-10, 11-20,
9 21-30, 31-40, 41-50, 51-60, 61-70, 71-80, 81+ years), patient sex (Male/Female),
10 patient NIQ (categorical; 1-5), if the FP has more patients in low NIQ areas (Yes/No),
11 patient residence (Urban/Rural), if the FP has more urban patients than rural (Yes/No),
12 if the FP has more patients with at least one psychosocial ADG (Yes/No), patient region
13 (categorical; LHIN1-14), total number of ED visits (categorical; 0-59, 60-156, 157-257,
14 258-414, 415+), total number of mental health ED visits (categorical; 0-3, 4-9, 10-19,
15 20+ visits), total hospitalizations (categorical; 0, 1-2, 3-5, 6-10, 11+), total mental health
16 hospitalizations (categorical; 0, 1-2, 3-5, 6-10, 11+), total primary care visits
17 (categorical; 0-59, 60-156, 157-257, 258-414, 415+) and total mental health primary
18 care provider visits (categorical; 0-59, 60-156, 157-257, 258-414, 415+).

39 Statistical Analysis

40
41 We used statistical modeling to understand the predictors of referring to telepsychiatry
42 among physicians. We used GEE for the statistical model, as it accounts for patient
43 clustering at the provider level.⁴¹ ~~The outcome variable was a referral to telepsychiatry~~
44 ~~(Yes/No) and the covariates are the same as those described in the comparison above,~~
45 ~~however, whether they were part of a family health team or not, and number of nurse~~
46
47
48
49
50
51
52
53
54

1
2
3 ~~practitioner variables were removed, as they were enveloped within the cluster and thus~~
4 ~~would have been considered co-linear.~~ All analyses were carried out using SAS version
5
6 9.4 (SAS Institute, Cary, NC). All results were presented as adjusted odds ratios (ORs)
7
8 with corresponding 95% confidence intervals (CIs).
9
10
11
12

13 14 15 **Results**

16 17 18 *Descriptive analysis of FTPs*

19
20
21 ~~Our study compared the difference between FTPs and FPNTs, using descriptive and~~
22 ~~statistical analyses.~~ In FY 2016, there were 13,290 family physicians in Ontario; this
23
24 analysis includes a subset of those family physicians who have referred to
25
26 telepsychiatry.⁴²
27
28
29

30 31 32 *Increase in Patients Accessing and Family Physicians Referring to Telepsychiatry*

33
34
35 The number of patients receiving telepsychiatry increased from 925 patients in FY 2008
36
37 to 13,825 patients in FY 2016, which represents approximately a fifteen-fold increase
38
39 (see Table 1). There was ~~also~~ substantial growth in the number of FPTs from 474 to
40
41 4,224, which represents a nine-fold increase from FY 2008 to FY 2016 (see Table 1). Of
42
43 all physicians in Ontario, 32% referred to telepsychiatry in FY 2016 (4224/13290). Less
44
45 than 1% of total patients rostered to FPTs had a telepsychiatry visit in FY 2018 (n=
46
47 12,449/3,513,638). [Insert Figure 1]
48
49
50
51
52
53
54

Physician Rurality

In FY 2008, the percentage of total telepsychiatry visits was highest in the North East LHIN (20%, n = 94), followed by the North West LHIN (16%, n = 74) (Ontario's most northern regions), followed by the South West LHIN (15%, n = 70). In FY 2016, those numbers shifted so ~~that~~ the highest percentage of telepsychiatry referrals came from family physicians in ~~more-central~~ regions such as Hamilton Niagara Haldimand Brant (10%, n = 433), Central East (10%, n = 421), and Toronto Central (10%, n = 420) LHINs, with the Central (n = 393) and North East (n = 379) LHIN's family physicians referring 9% each. All LHINs ranged between 4-10% of total telepsychiatry delivery in FY 2016, compared to FY 2008, where the range spread from 0% (in Mississauga Halton LHIN) to 20% (in the North East LHIN). This shows a more evenly distributed adoption of telepsychiatry from ~~strictly~~ northern regions to all regions of Ontario, many of which still have highly rural and remote communities. [Insert Figure 2]

Comparative Analysis

Family Physician Characteristics

A larger percentage of FPTs were male, compared to FPNTs (59.6% (n = 2427) vs. 53.1% (n = 4075), SD = 0.13). A greater percentage of FPTs were from communities with a population that had less than 10,000 people (14.9% (n = 608) vs. 8.2% (n = 633), SMD = 0.21). FPTs had a larger mean number of nurse practitioners and FPNTs (1.71 (+/- 4.02) vs. 1.10 (+/- 3.70), SMD = 0.16), finally, a greater percentage of FPTs

1
2
3 belonged to FHTs than FPNTs (28.1% (n = 1145) vs. 18% (n = 1381), SMD = 0.24).

4
5 [Insert Table 1]

6 7 8 *Comparing Patient Roster Characteristics*

9
10
11
12 There were a few notable differences in patient roster characteristics between FPTs and
13
14 FPNTs. FPTs had a larger mean number of attached patients than FPNTs (862.45 (+/-
15
16 668.36) vs. 788.97 (+/- 605.70), SMD = 0.15). A greater percentage of FPTs had
17
18 patients on their roster from lower NIQ's than FPNTs (1-3) (77.7% (n = 3167) vs. 69.8%
19
20 (n = 5353), SMD = 0.18). FPTs had a larger mean number of patients from a rural
21
22 residence compared to FPNTs (139.28 (+/- 256.05) vs. 54.93 (+/- 137.27), SMD = 0.41).
23
24 A greater percentage of FPTs had more patients from rural areas than urban areas
25
26 (14.6% (N = 594) vs. 6.8% (n = 522), SMD = 0.25). FPNTs had a higher patient mean in
27
28 the Champlain LHIN (which includes Ottawa, a large urban region in Ontario) (FPT =
29
30 61.94 (+/- 236.36), FPTN = 99.43 (+/-287.94), SMD = 0.14). FPTs had higher patient
31
32 means in four LHINs (mostly northern regions): South East (SE) LHIN (FPT = 47.03 (+/-
33
34 209), FPTN = 24.27 (+/-132.92), SMD = 0.13), North Simcoe Muskoka (NSM) LHIN
35
36 (FPT = 59.28 (+/- 230.38), FPNT = 15.12 (+/- 97.56), SMD =0.25), North East (NE)
37
38 LHIN (FPT = 53.59 (+/- 228.57), FPNT = 19.86 (+/- 124.20), SMD =0.18), North West
39
40 (NW) LHIN (FPT = 24.64 (+/- 149.32), FPNT = 6.06 (+/- 101.30), SMD = 0.16).
41
42
43
44
45
46
47
48

49 The mean number of patients with psychosocial ADG's was higher for FPTs compared
50
51 to FPNTs (258.24 (+/- 204.4) vs. 217.56 (+/- 179.29), SMD = 0.21).The mean mental
52
53 health primary care provider (MHPCP) visits was higher in FPTs compared to FPNTs
54
55

1
2
3 (707.60 (+/- 2863.95) vs. 276.86 (+/- 532.56), SMD = 0.21). Due to the large standard
4
5 deviation, we also examined the median for MHPCP; FPTs had a higher median
6
7 MHPCP visits than FPNTs (229 (IQR=106-415) vs. 189 (IQR=110-350), SMD = 0.24).
8
9
10 FPTs have a higher percentage of patients with 415+ MHPCP visits compared to
11
12 FPNTs (25.2% (n = 1026) vs. 16.7% (n = 1285), SMD 0.21).
13
14
15
16

17 The mean number of ED visits was **significantly** higher for FPTs compared to FPNTs
18
19 (440.82 (+/- 357.7) vs. 291.96 (+/- 240.01), SMD = 0.49). Table 1 describes the
20
21 difference between all categories, but the most notable differences are for patients that
22
23 had 60-156, 157-257, and 415+ visits. FPTs had a lower percentage of patients with 60-
24
25 156 ED visits than FPNTs (9.5% (n = 387) vs. 17.2% (n = 1320), SMD = 0.23) and
26
27 patients with 157-257 visits (12.0% (n = 489) vs. 19.5% (n = 1495), SMD = 0.21). FPTs
28
29 has a greater percentage of patients with 415+ ED visits compared to FPNTs (45.6% (n
30
31 = 1857) vs. 16.7% (n = 1285), SMD = 0.48).
32
33
34
35
36
37

38 FPTs had a larger mean number of MHED visits for rostered patients compared to
39
40 FPNTs (18.99 (+/- 29.07) vs. 10.91 (+/- 13.07), SMD = 0.36). Full details within each
41
42 category are in Table 1, but the most notable difference was the group that had 0-3 and
43
44 20+ MHED visits. FPTs had a lower percentage of patients with 0-3 MHED visits than
45
46 FPNTs (16.7% (n = 681) vs. 27.5% (n = 2108), SMD = 0.26) and a higher percentage of
47
48 patients with 20+ MHED visits than FPNTs (30.9% (n = 1258) vs. 14.1% (n = 1084),
49
50 SMD = 0.41).
51
52
53
54
55
56
57
58
59
60

1
2
3 FPTs had a greater mean of hospitalizations compared to FPNTs (57.05 (+/-47.43) vs.
4 10.91 (+/-13.07), SMD = 0.36). Full details within each category are in Table 1, but the
5
6 most notable difference was that FPTs had a larger percentage of patients with 11+
7 hospitalizations, compared to FPNTs (85.7% (n = 3491) vs. 78.8% (n = 6044), SMD =
8 0.18). Of those hospitalizations, mental health specific hospitalizations (MHH) were
9 measured. FPTs had a greater mean of MHH visits compared to FPNTs (7.01 (+/-13.61)
10 vs. 3.86 (+/-5.43), SMD = 0.36). The most striking differences were the groups that had
11 zero visits and those with 11+ visits. FPTs had a lower percentage of patients with zero
12 MHH visits compared to FPNTs (13.4% (n = 545) vs. 21.9% (n = 1679), SMD = 0.22).
13
14 FPTs had a higher percentage of patients with 11+ MHH visits compared to FPNTs
15 (17.8% (n = 725) vs. 6.8% (n = 518), SMD = 0.34).
16
17
18
19
20
21
22
23
24
25
26
27
28

29 **Statistical Modeling**

30 *Regression using Generalized Estimating Equations*

31
32
33 A full description of the findings is listed in Table 2. Within this model, a number of
34 significant correlates of FPTs were identified, including physician age group of 55-64
35 (OR=0.85, CI=0.74-0.97), physician age group of 64+ (OR=0.85, CI=0.62-0.85), and
36
37 FPTs were more likely to be located in rural areas (OR=1.38, CI=1.17-1.62). Compared
38 to patient rosters of FPNTs, FPT rostered patients were: almost half as likely to be
39 urban (OR=0.55, CI=0.43-.70), more likely to have psychiatric ADGs (OR=1.2, CI=0.86-
40 1.66), and more likely to have 4-9 mental health ED (MHED) visits (OR=1.33, CI=1.03-
41 1.72), patients with 10-19 MHED visits (OR=1.54 CI=1.14-2.07), patients with 20+
42
43
44
45
46
47
48
49
50
51
52
53
54

1
2
3 MHED visits (OR=2.17, CI=1.60-2.94), patients with 3-5 MHH (OR=1.21, CI=1.08-1.35),
4
5 patients with 6-10 MHH (OR=1.32, CI=1.15-1.51), patients with 11+ MHH (OR=1.77,
6
7 CI=1.28-2.43), patients with 157-257 mental health primary care (MHPC) visits
8
9 (OR=1.24, CI=1.00-1.54), patients with 258-414 MHPC visits (OR=1.38 CI=1.09-1.76),
10
11 patients with 415+ MHPC visits (OR=1.61, CI=01.33-1.94). [Insert Table 2].
12
13
14
15
16

17 Discussion

18
19 To our knowledge, this is the first study that investigates trends and predictors of family
20
21 physicians referring to telepsychiatry at a system-level. Between FY 2008 and FY 2016,
22
23 there was a 9-fold increase in the number of primary care physicians referring to
24
25 telepsychiatry and a nearly 15-fold increase in the number of primary care patients
26
27 referred to telepsychiatry. 32% of family physicians referred to telepsychiatry, while less
28
29 than 1% of patients rostered to FPTs utilized telepsychiatry. [While patient adoption of](#)
30
31 [telepsychiatry continued to increase in Ontario during the study period, it still represents](#)
32
33 [a very small percentage of patients in Ontario. These visits represent, for the most part,](#)
34
35 [patients that ~~are~~ were required to go to a healthcare site to access care and psychiatrists](#)
36
37 [that are required to use the ~~OTN~~ provincial telemedicine platform. It is evident that there](#)
38
39 [remains a large opportunity to increase patient adoption. ~~As reported elsewhere, in~~](#)
40
41 [2012, only about 7% of psychiatrists were providing telepsychiatry visits, so similar to](#)
42
43 [patients, there is still significant adoption potential for psychiatrists.²⁷-As discussed in](#)
44
45 [the introduction, it is likely that the recent Covid-19 related temporary policy changes to](#)
46
47 [virtual care ~~might~~ will present an opportunity for significant growth and adoption of](#)
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 telepsychiatry. This study can provides an essential benchmark for future analyses,
4 comparing how these policy changes have changedshifted adoption patterns with
5 patients and family physician referrals.
6
7
8
9

10
11
12 Physicians who referred to telepsychiatry in FY 2008 were mainly from northern regions
13 (LHINs) of the province, whereas in FY 2016, more referrals from southern regions
14 (LHINs) emerged, many of which still have rural pockets areas within their catchments.
15
16
17
18

19 Referral to telepsychiatry was much more evenly distributed throughout the province in
20 FY 2016. In FY 2016 physicians from all LHINs referred somewhere between 4-10% of
21 total telepsychiatry, compared to FY 2008 where the range spread from 0% to 20%,
22 with six LHINs that delivered more than 9% of total telepsychiatry each and eight LHINs
23 all delivering less than 4% each. While historically telepsychiatry was used to support
24 rural populations, in later years of the study, there were increasing provincially funded
25 'urban' telepsychiatry projects that aimed to increase adoption of telepsychiatry within
26 urbanmore populous regions. While it i's important to create access and choice for
27 patients in both rural and urban areas, and increase access provincially, policy-makers
28 and funders should consider how existing payment and funding models support
29 prioritization of care to underserved populations, and might consider strategies such as
30 a wait-times dashboard to help better distribute care virtually. aspects of digital health
31 equity⁴³, allowing the use of diverse secure virtual platforms, funding home-based visits
32 and establishing mechanisms to support the appropriate distribution of virtual care
33 throughout the province.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Past studies have suggested that physicians who were more isolated and were part of
4 an integrated health system had positive feelings about telemedicine, and that nurses
5 were more enthusiastic about telemedicine than physicians.^{25,36} As such, ~~unperhaps not~~
6 surprisingly, this study found that predictors of family physician referrals to
7 telepsychiatry include rurality of residence (for both provider and patient), ~~team-based~~
8 ~~practice models, having more~~ nurse practitioners on the team, ~~and clinical complexity~~
9 ~~of patient rosters, and if referring the provider was part of a family health team.~~
10 Additionally, ~~FPTs were more likely to have rostered patients who had more clinical~~
11 ~~complexity and higher healthcare utilization rates. While this might be an indicator of~~
12 ~~patients' overall complexity, the number of people accessing mental health care in the~~
13 ~~ED can also be an indicator of a lack of access to outpatient mental health care. As~~
14 ~~access to telepsychiatry continues to increase, studies measuring potential diversion~~
15 ~~from ED-related mental health visits would be helpful.~~

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37 This study has numerous strengths; it utilizes a large, multi-year, linked dataset of family
38 physician and patient data to help understand important characteristics that are
39 common in physicians who refer to telepsychiatry. Additionally, since this study uses
40 administrative data, the study was conducted rapidly using a retrospective analysis.
41
42 ~~However, t~~~~he~~ study also has several limitations. The identification of telepsychiatry
43 visits is contingent upon psychiatrists using telepsychiatry billing codes and billing fee
44 for service, so it is possible that there is missing data, or an underrepresentation of
45 visits completed by physicians paid by salary or alternate funding. Another potential
46
47
48
49
50
51
52
53
54
55

1
2
3 limitation, is that while this study includes nurse practitioners as a covariate in the
4
5 analysis, it does not take into account actual nurse practitioner referrals to
6
7 telepsychiatry, which could lead to an underestimate of our total telepsychiatry visits.
8
9 We believe, however, that this will have minimal impact on our study because it is likely
10
11 that nurse practitioner referrals will be captured within total telepsychiatry visits, but
12
13 rostered (as defined in our methods) to a family physician instead.
14
15
16
17
18

19 Conclusion

20
21 ~~Over the study period, the number of patients receiving and providers referring to~~
22 ~~telepsychiatry has increased by fifteen-fold and nine-fold, While telepsychiatry has seen~~
23 ~~continued growth over recent years, respectively there remains significant opportunities~~
24 ~~for increased adoption.~~ This study shows that certain factors, including rurality, provider
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FPNTs. This study provides a baseline from which to assess recent rapid mobilization of telepsychiatry, and a means of doing so using administrative health data.

Tables and Figures

Figure 1. Telepsychiatry Adoption by Year (FY2008 and FY2016)




Figure 2. Percentage of Referring Physicians by Region (2008 and 2016)




Table 1. Comparative Analysis of Family Physicians that Refer to Telepsychiatry compared to (2016)

VARIABLE	VALUE	FPT	FPNT	TOTAL	SMD
		N = 4,074	N = 7,674	N = 11,748	
PHYSICIAN CHARACTERISTICS					
Physician Sex	F	1,647 (40.4%)	3,599 (46.9%)	5,246 (44.7%)	0.13
	M	2,427 (59.6%)	4,075 (53.1%)	6,502 (55.3%)	0.13
< 10,000 population in CMA	N	3,466 (85.1%)	7,041 (91.8%)	10,507 (89.4%)	0.21
	Y	608 (14.9%)	633 (8.2%)	1,241 (10.6%)	0.21
Nurse Practitioners on Team	Mean ± SD	1.71 ± 4.02	1.10 ± 3.70	1.31 ± 3.83	0.16
Physician Practice Type	FHT	1,145 (28.1%)	1,381 (18.0%)	2,526 (21.5%)	0.24
	Others	2,929 (71.9%)	6,293 (82.0%)	9,222 (78.5%)	0.24
PATIENT ROSTER CHARACTERISTICS					
Total # of attached patients	Mean ± SD	862.45 ± 668.36	766.97 ± 605.70	800.64 ± 630.13	0.15
# patients in NIQ 1,2&3 > # patients in NIQ 4&5	No	907 (22.3%)	2,127 (27.7%)	3,034 (25.8%)	0.13
	Yes	3,167 (77.7%)	5,353 (69.8%)	8,520 (72.5%)	0.18
Rural Residence	Mean ± SD	139.28 ± 256.05	54.93 ± 137.27	84.67 ± 192.19	0.41
# of patients in urban residence > # of patients in rural residence	No	594 (14.6%)	522 (6.8%)	1,116 (9.5%)	0.25
	Yes	3,480 (85.4%)	6,958 (90.7%)	10,438 (88.8%)	0.16
South East LHIN	Mean ± SD	47.03 ± 209.09	24.27 ± 132.92	32.29 ± 164.22	0.13
Champlain LHIN	Mean ± SD	61.94 ± 263.36	99.43 ± 287.94	86.21 ± 280.09	0.14
North Simcoe Muskoka LHIN	Mean ± SD	59.28 ± 230.38	15.12 ± 97.56	30.69 ± 159.11	0.25
North East LHIN	Mean ± SD	53.59 ± 228.57	19.86 ± 124.20	31.75 ± 169.31	0.18
North West LHIN	Mean ± SD	24.64 ± 149.32	6.06 ± 59.90	12.61 ± 101.30	0.16
# of patients with a psychosocial ADG	Mean ± SD	258.24 ± 204.40	217.56 ± 179.29	231.91 ± 189.52	0.21
Total MH PCP visits	Mean ± SD	707.60 ± 2,863.95	276.86 ± 532.56	428.74 ± 1,765.68	0.21
	Median (IQR)	239 (106-415)	188 (73-335)	204 (82-364)	0.24
	00-59	686 (16.8%)	1,626 (21.2%)	2,312 (19.7%)	0.11
	60-156	702 (17.2%)	1,612 (21.0%)	2,314 (19.7%)	0.1
157-257	795 (19.5%)	1,512 (19.7%)	2,307 (19.6%)	0	

	258-414	865 (21.2%)	1,445 (18.8%)	2,310 (19.7%)	0.06
	415-high	1,026 (25.2%)	1,285 (16.7%)	2,311 (19.7%)	0.21
VARIABLE	VALUE	FPT	FPNT	TOTAL	SMD
Total ED visits	Mean ± SD	440.82 ± 357.70	291.96 ± 240.01	344.45 ± 295.74	0.49
	00-59	493 (12.1%)	1,090 (14.2%)	1,583 (13.5%)	0.06
	60-156	387 (9.5%)	1,320 (17.2%)	1,707 (14.5%)	0.23
	157-257	489 (12.0%)	1,495 (19.5%)	1,984 (16.9%)	0.21
	258-414	848 (20.8%)	1,782 (23.2%)	2,630 (22.4%)	0.06
	415-high	1,857 (45.6%)	1,793 (23.4%)	3,650 (31.1%)	0.48
Total MH ED visits	Mean ± SD	18.99 ± 29.07	10.91 ± 13.07	13.75 ± 20.58	0.36
	00-03	681 (16.7%)	2,108 (27.5%)	2,789 (23.7%)	0.26
	04-09	922 (22.6%)	2,275 (29.6%)	3,197 (27.2%)	0.16
	10-19	1,213 (29.8%)	2,013 (26.2%)	3,226 (27.5%)	0.08
	20-high	1,258 (30.9%)	1,084 (14.1%)	2,342 (19.9%)	0.41
Total hospitalizations	Mean ± SD	57.05 ± 47.43	41.75 ± 36.01	47.15 ± 41.06	0.36
	0 visits	116 (2.8%)	286 (3.7%)	402 (3.4%)	0.05
	01-02 visit	133 (3.3%)	287 (3.7%)	420 (3.6%)	0.03
	03-05 visits	131 (3.2%)	374 (4.9%)	505 (4.3%)	0.08
	06-10 visits	203 (5.0%)	489 (6.4%)	692 (5.9%)	0.06
	11+ visits	3,491 (85.7%)	6,044 (78.8%)	9,535 (81.2%)	0.18
Total MH hospitalizations	Mean ± SD	7.01 ± 13.61	3.86 ± 5.43	4.97 ± 9.31	0.3
	0 visits	545 (13.4%)	1,679 (21.9%)	2,224 (18.9%)	0.22
	01-02 visit	768 (18.9%)	2,078 (27.1%)	2,846 (24.2%)	0.2
	03-05 visits	1,072 (26.3%)	1,959 (25.5%)	3,031 (25.8%)	0.02
	06-10 visits	964 (23.7%)	1,246 (16.2%)	2,210 (18.8%)	0.19
	11+ visits	725 (17.8%)	518 (6.8%)	1,243 (10.6%)	0.34

NIQ= Neighbourhood Income Quintiles
 LHIN=Local Health Integration Networks
 ADG= Aggregated Diagnosis Groupings
 MHPCP= Mental Health Primary Care Practitioner
 ED= Emergency Department
 MH ED= Mental Health Emergency Department
 MH= Mental Health

Table 2. Model using Generalized Estimating Equations comparing FPTs vs FPNTs (2016)

Analysis Of GEE Parameter Estimates			
	Estimate	95% Confidence Limits	
Intercept	-0.9162	-1.1901	-0.6423
Physician Age 35-44 vs <=34	0.8895	0.7855	1.0072
Physician Age 45-54 vs <=34	0.9082	0.7983	1.0333
Physician Age 55-64 vs <=34	0.849	0.7419	0.9717
Physician Age 64+ vs <=34	0.7255	0.6223	0.8458
Sex (Male vs Female)	1.0144	0.9368	1.0983
Physician Rural (Yes vs No)	1.3791	1.174	1.6199
Canadian Medical Graduate (Yes vs No)	1.0333	0.9493	1.1247
Canadian Medical Graduate (Unknown vs No)	1.0231	0.8617	1.2148
More patients in low/med income quintile (Yes vs No)	1.0719	0.969	1.1856
More urban patients (Yes vs No)	0.5491	0.4338	0.6952
Mental Health ED Visits (04-09 vs 00-03)	1.3289	1.0252	1.7226
Mental Health ED Visits (10-19 vs 00-03)	1.5373	1.1442	2.0654
Mental Health ED Visits (20-high vs 00-03)	2.1678	1.601	2.9352
Mental Health Hospitalizations (0 visits vs 01-02 visit)	0.9072	0.7363	1.1179
Mental Health Hospitalizations 03-05 visits vs 01-02 visit	1.2048	1.0749	1.3504
Mental Health Hospitalizations 06-10 visits vs 01-02 visit	1.316	1.1496	1.5064
Mental Health Hospitalizations 11+ visits vs 01-02 visit	1.7662	1.2837	2.4301
Mental Health Primary Care Visits (60-156 vs 00-59)	0.965	0.8134	1.1449
Mental Health Primary Care Visits (157-257 vs 00-59)	1.237	0.9943	1.539
Mental Health Primary Care Visits (258-414 vs 00-59)	1.384	1.0859	1.7638
Mental Health Primary Care Visits (415-high vs 00-59)	1.6089	1.3322	1.943
More patients with psychiatric ADGs (Yes vs No)	1.1979	0.8627	1.6634

ED= Emergency Department

ADG= Aggregated Diagnosis Groups

References

1. Brien S, Grenier L, Kapral M. *Taking stock*. Health Quality Ontario; 2015.
2. Bishop TF, Press MJ, Keyhani S, Pincus HA. Acceptance of insurance by psychiatrists and the implications for access to mental health care. *JAMA psychiatry*. 2014;71(2):176-181.
3. World Health Organization. Mental disorders affect one in four people. *World Health Report*. 2001.
4. Alegría M, Nakash O, NeMoyer A. Increasing equity in access to mental health care: A critical first step in improving service quality. *World Psychiatry*. 2018;17(1):43.
5. Wang PS, Aguilar-Gaxiola S, Alonso J, et al. Use of mental health services for anxiety, mood, and substance disorders in 17 countries in the WHO world mental health surveys. *The Lancet*. 2007;370(9590):841-850.
6. Goldner EM, Jones W, Fang ML. Access to and waiting time for psychiatrist services in a canadian urban area: A study in real time. *The Canadian Journal of Psychiatry*. 2011;56(8):474-480.
7. Meadows GN, Enticott JC, Inder B, Russell GM, Gurr R. Better access to mental health care and the failure of the medicare principle of universality. *Med J Aust*. 2015;202(4):190-194.

- 1
2
3 8. Cunningham PJ. Beyond parity: Primary care physicians' perspectives on access to
4 mental health care: More PCPs have trouble obtaining mental health services for their
5 patients than have problems getting other specialty services. *Health Aff.*
6
7
8
9
10 2009;28(Suppl1):w490-w501.
11
12
13 9. Kurdyak P, Zaheer J, Cheng J, Rudoler D, Mulsant BH. Changes in characteristics
14 and practice patterns of ontario psychiatrists: Implications for access to psychiatrists.
15
16
17
18 *The Canadian Journal of Psychiatry.* 2017;62(1):40-47.
19
20
21 10. Kurdyak P, Stukel TA, Goldbloom D, Kopp A, Zagorski BM, Mulsant BH. Universal
22 coverage without universal access: A study of psychiatrist supply and practice patterns
23 in ontario. *Open Medicine.* 2014;8(3):e87.
24
25
26
27
28
29 11. Myers KM, Valentine JM, Melzer SM. Child and adolescent telepsychiatry:
30 Utilization and satisfaction. *Telemedicine and e-Health.* 2008;14(2):131-137.
31
32
33
34
35 12. Antonacci DJ, Bloch RM, Saeed SA, Yildirim Y, Talley J. Empirical evidence on the
36 use and effectiveness of telepsychiatry via videoconferencing: Implications for forensic
37 and correctional psychiatry. *Behav Sci Law.* 2008;26(3):253-269.
38
39
40
41
42
43 13. Campbell R, O'Gorman J, Cernovsky ZZ. Reactions of psychiatric patients to
44 telepsychiatry. *Mental illness.* 2015;7(2).
45
46
47
48
49 14. Deslich S, Stec B, Tomblin S, Coustasse A. Telepsychiatry in the 21st century:
50 Transforming healthcare with technology. *Perspectives in Health Information*
51
52
53
54

1
2
3 *Management/AHIMA, American Health Information Management Association.*

4
5 2013;10(Summer).

6
7
8
9 15. Fleming DA, Edison KE, Pak H. Telehealth ethics. *Telemedicine and e-Health.*

10
11 2009;15(8):797-803.

12
13
14 16. Saeed SA, Diamond J, Bloch RM. Use of telepsychiatry to improve care for people
15 with mental illness in rural North Carolina. *NC Med J.* 2011;72(3):219-222.

16
17
18 17. Ulzen T, Williamson L, Foster PP, Parris-Barnes K. The evolution of a community-
19 based telepsychiatry program in rural alabama: Lessons learned—a brief report.

20
21
22 *Community Ment Health J.* 2013;49(1):101-105.

23
24
25 18. Lyketsos CG, Roques C, Hovanec L, Jones III BN. Telemedicine use and the
26 reduction of psychiatric admissions from a long-term care facility. *J Geriatr Psychiatry*
27 *Neurol.* 2001;14(2):76-79.

28
29
30 19. Hilty DM, Ferrer DC, Parish MB, Johnston B, Callahan EJ, Yellowlees PM. The
31 effectiveness of telemental health: A 2013 review. *Telemedicine and e-Health.*

32
33 2013;19(6):444-454.

34
35
36 20. Chakrabarti S. Usefulness of telepsychiatry: A critical evaluation of

37
38
39 videoconferencing-based approaches. *World journal of psychiatry.* 2015;5(3):286.

- 1
2
3 21. Bishop JE, O'Reilly RL, Maddox K, Hutchinson LJ. Client satisfaction in a feasibility
4 study comparing face-to-face interviews with telepsychiatry. *J Telemed Telecare*.
5
6 2002;8(4):217-221.
7
8
9
10
11 22. Frueh BC, Monnier J, Yim E, Grubaugh AL, Hamner MB, Knapp RG. A randomized
12 trial of telepsychiatry for post-traumatic stress disorder. *J Telemed Telecare*.
13
14 2007;13(3):142-147.
15
16
17
18
19 23. Farabee D, Calhoun S, Veliz R. An experimental comparison of telepsychiatry and
20 conventional psychiatry for parolees. *Psychiatric Services*. 2016;67(5):562-565.
21
22
23
24
25 24. Bashshur RL, Shannon GW, Bashshur N, Yellowlees PM. The empirical evidence
26 for telemedicine interventions in mental disorders. *Telemedicine and e-Health*.
27
28 2016;22(2):87-113.
29
30
31
32
33 25. Sargeant J, Allen M, Langille D. Physician perceptions of the effect of telemedicine
34 on rural retention and recruitment. *J Telemed Telecare*. 2004;10(2):89-93.
35
36
37
38
39 26. Swanson B. Information technology and under-served communities. *J Telemed*
40
41 *Telecare*. 1999;5(2_suppl):3-10.
42
43
44
45 27. Serhal E, Crawford A, Cheng J, Kurdyak P. Implementation and utilisation of
46 telepsychiatry in Ontario: A population-based study. *The Canadian Journal of*
47
48 *Psychiatry*. 2017;62(10):716-725.
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 28. Grumbach K, Selby JV, Damberg C, et al. Resolving the gatekeeper conundrum:
4
5 What patients value in primary care and referrals to specialists. *JAMA*.
6
7 1999;282(3):261-266.
8
9
10
11 29. Norton PG, Dunn EV, Bestvater D. Referrals in primary care: Is the family physician
12
13 a "Gatekeeper"? *Canadian Family Physician*. 1989;35:1776.
14
15
16
17 30. Steele M, Zayed R, Davidson B, et al. Referral patterns and training needs in
18
19 psychiatry among primary care physicians in canadian rural/remote areas. *Journal of*
20
21 *the Canadian Academy of Child and Adolescent Psychiatry*. 2012;21(2):111.
22
23
24
25 31. Spaulding RJ, Russo T, Cook DJ, Doolittle GC. Diffusion theory and telemedicine
26
27 adoption by kansas health-care providers: Critical factors in telemedicine adoption for
28
29 improved patient access. *J Telemed Telecare*. 2005;11(1_suppl):107-109.
30
31
32
33 32. Whitten P, Franken EA. Telemedicine for patient consultation: Factors affecting use
34
35 by rural primary-care physicians in kansas. *J Telemed Telecare*. 1995;1(3):139-144.
36
37
38
39 33. LeRouge C, Garfield M. Crossing the telemedicine chasm: Have the US barriers to
40
41 widespread adoption of telemedicine been significantly reduced? *International journal of*
42
43 *environmental research and public health*. 2013;10(12):6472-6484.
44
45
46
47 34. Moore MA, Coffman M, Jetty A, Klink K, Petterson S, Bazemore A. Family
48
49 physicians report considerable interest in, but limited use of, telehealth services. *The*
50
51 *Journal of the American Board of Family Medicine*. 2017;30(3):320-330.
52
53
54

- 1
2
3 35. Spaulding R, Cain S, Sonnenschein K. Urban telepsychiatry: Uncommon service for
4 a common need. *Child and Adolescent Psychiatric Clinics*. 2011;20(1):29-39.
5
6
7
8
9 36. Whitten PS, Mackert MS. Addressing telehealth's foremost barrier: Provider as initial
10 gatekeeper. *Int J Technol Assess Health Care*. 2005;21(4):517-521.
11
12
13
14 37. Steele LS, Glazier RH, Lin E, Evans M. Using administrative data to measure
15 ambulatory mental health service provision in primary care. *Med Care*. 2004:960-965.
16
17
18
19 38. Austin PC, van Walraven C, Wodchis WP, Newman A, Anderson GM. Using the
20 Johns Hopkins aggregated diagnosis groups (ADGs) to predict mortality in a general
21 adult population cohort in Ontario, Canada. *Med Care*. 2011;49(10):932.
22
23
24
25
26
27
28 39. Cohen J. *Statistical power analysis for the behavioral sciences*. Routledge; 2013.
29
30
31
32 40. Austin PC. Using the standardized difference to compare the prevalence of a binary
33 variable between two groups in observational research. *Communications in Statistics-
34 Simulation and Computation*. 2009;38(6):1228-1234.
35
36
37
38
39
40 41. Wang M. Generalized estimating equations in longitudinal data analysis: A review
41 and recent developments. *Advances in Statistics*. 2014;2014.
42
43
44
45
46 42. The Ontario Physician Human Resources Data Centre. . (2018, 10). Physicians in
47 Ontario 2017. Retrieved from the Ontario Physician Human Resources data centre:
48 [Http://Www.ophrdc.org/wp-content/uploads/2018/10/PIO-2017-report-with-](http://www.ophrdc.org/wp-content/uploads/2018/10/PIO-2017-report-with-hyperlinks.pdf)
49 [hyperlinks.pdf](http://www.ophrdc.org/wp-content/uploads/2018/10/PIO-2017-report-with-hyperlinks.pdf). *McMaster University*. 2018.
50
51
52
53
54

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

43. Crawford, A., & Serhal, E. (2020). Digital Health Equity and COVID-19: The Innovation Curve Cannot Reinforce the Social Gradient of Health. Journal of Medical Internet Research, 22(6), e19361.