This is a repository copy of *Asymmetric 'Clip-Cycle' Synthesis of Pyrrolidines and Spiropyrrolidines*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/166173/>

Version: Accepted Version

Article:

Clarke, Paul Andrew orcid.org/0000-0003-3952-359X, Ermanis, Kristaps and Maddocks, Christopher (2020) Asymmetric 'Clip-Cycle' Synthesis of Pyrrolidines and Spiropyrrolidines. *Organic Letters*. pp. 8116-8121. ISSN 1523-7052

<https://doi.org/10.1021/acs.orglett.0c03090>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Asymmetric 'Clip-Cycle' Synthesis of Pyrrolidines and Spiropyrrolidines.

Christopher J. Maddocks,[†] Kristaps Ermanis*[‡] and Paul A. Clarke*[†]

[†] Department of Chemistry, University of York, Heslington, York, North Yorks, UK, YO10 5DD.

[‡] Centre for Molecular Informatics, Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, UK, CB2 1EW.

Supporting Information Placeholder

ABSTRACT: The development of an asymmetric 'clip-cycle' synthesis of 2,2- and 3,3-disubstituted pyrrolidines and spiropyrrolidines, which are increasingly important scaffolds in drug discovery programs, is reported. Cbz-protected bis-homoallylic amines were activated by 'clipping' them to thioacrylate *via* an alkene metathesis reaction. Enantioselective intramolecular aza-Michael cyclisation onto the activated alkene, catalyzed by a chiral phosphoric acid, formed a pyrrolidine. The reaction accommodated a range of substitution to form 2,2- and 3,3-disubstituted pyrrolidines and spiropyrrolidines with high enantioselectivities. The importance of the thioester activating group was demonstrated by comparison to ketone and oxoester-containing substrates. DFT studies supported the aza-Michael cyclisation as the rate and stereochemical determining step, and correctly predicted the formation of the major enantiomer. The catalytic asymmetric syntheses of *N*-methyl pyrrolidine alkaloids (*R*)-irnidine and (*R*)-bgugaine, which possess DNA binding and antibacterial properties were achieved using the 'clip-cycle' methodology.

Pyrrolidines appear in nearly 20% of FDA-approved drug molecules which contain a saturated cyclic amine unit,¹ making them one of the most important structural features in modern pharmaceuticals (Figure 1). Recently, spirocyclic amines, including pyrrolidines, have become highly desirable scaffolds^{2,3} as they are conformationally well-defined which allows them to be elaborated along specific vectors. Pyrrolidines are also present in many biologically active natural products.⁴

However, despite this prevalence, the methods available to make them asymmetrically often have drawbacks. A common method is asymmetric lithiation of *N*-Boc pyrrolidine, followed by electrophilic quenching.⁵ The drawbacks of cryogenic temperatures, pyrophoric reagents and the regular use of stoichiometric quantities of chiral diamine are well documented.⁵ The alternative strategies of asymmetric azomethine cycloaddition to olefins,⁶ organocatalytic approaches,⁷ and aza-Michael reactions,⁸ including the asymmetric addition to nitroalkenes⁹ tend to generate products which need to undergo further reactions. An alternative strategy is the *intramolecular* aza-Michael cyclisation, and while there are racemic reactions which generate pyrrolidines in good yields,¹⁰ there are limited examples of asymmetric versions of this reaction. Those which have been reported use chiral auxiliaries on nitrogen,¹¹ or are limited to the formation of specific pyrrolidine substitution patterns.¹² Interest in the synthesis of substituted pyrrolidines and spiropyrrolidines has increased over the last few years due to their recently recognized value in drug discovery projects.^{2,3a,13} However, while new methods to synthesize them have been developed, few are general and asymmetric.¹⁴

Figure 1. Pyrrolidine-containing pharmaceuticals.

To address these limitations, we envisaged the development of the ‘clip-cycle’ procedure. ‘Clip-cycle’ is a strategy which ‘clips’ together a *N*-protected *bis*-homoallylic amine with a thioacrylate to generate an activated alkene, with a pendent nucleophile. The *N*-protected amine can then undergo aza-Michael cyclisation catalyzed by a chiral phosphoric acid (CPA), to yield enantioenriched pyrrolidines (Scheme 1).

Scheme 1. Asymmetric ‘Clip-Cycle’ Reaction.

In this paper, we report the successful implementation of a new two-step, programmable, catalytic approach for the asymmetric synthesis of pyrrolidines and spiropyrrolidines in high enantiopurity. The key features of this ‘clip-cycle’ strategy are that it is: (i) modular, enabling a diverse range of pyrrolidines to be assembled quickly and easily from readily accessible *N*-protected amines and acrylates, (ii) catalytic in both the ‘clip’ and ‘cycle’ steps, (iii) straightforward to perform, (iv) yields substituted pyrrolidine and spiropyrrolidine products directly in high enantiomeric excesses and, (v) the thioester function provides a versatile handle for late stage diversification.

Three different activating groups were examined initially, a ketone, an oxoester and a thioester, in order to determine the optimal electrophile for the asymmetric cyclisation (Scheme 2). Hoveyda-Grubbs 2nd generation catalyst (HG-II) was used to clip Cbz-protected *bis*-homoallylic amine **1a** to each of the electrophilic components. When **1a** was treated with HG-II in the presence of enone **2**, cyclisation to the pyrrolidine occurred spontaneously. Coupling **1a** with *p*-tolyl acrylate **4** generated oxoester **5** in 86% yield. Treatment of **5** with (*R*)-TRIP (20 mol%) in cyclohexane at 80 °C, resulted in only 24% conversion (20% yield) to the pyrrolidine **6**, which was formed in 95:5 e.r. Gratifyingly, when *p*-tolyl thioacrylate **7** was used as the metathesis partner α,β -unsaturated thioester **8a** was formed in 75% yield, which was cyclized to pyrrolidine **9a** using (*R*)-TRIP (20 mol%) in cyclohexane at 80 °C with 100% conversion (83% yield) and 98:2 e.r. (Scheme 2). α,β -Unsaturated thioesters were, therefore, chosen as the activating group of choice due to their intermediate electrophilicity compared to α,β -unsaturated aldehydes/ketones¹⁵ and oxoesters.¹⁶

Scheme 2. Effect of the electrophile on the asymmetric ‘clip-cycle’ synthesis of pyrrolidines.

The ‘clip-cycle’ reaction was shown to be tolerant of a range of substitution at the 3-position of the pyrrolidine (Figure 2). Replacing the methyl substituents with cycloalkyl groups¹⁷ (**9b-d**, Figure 2) did not adversely affect the enantioselectivity or yield of the ‘clip-cycle’ reaction, and generated spirocyclic pyrrolidines. The more sterically congested *gem*-diphenyl substitution (**9e**, Figure 2) was also tolerated in a slightly lower yield and enantioselectivity. Unsubstituted amine (*R*=H, **9f**, Figure 2) was also tolerated, and the pyrrolidine **9f** was formed in 73% yield and 90:10 e.r. The enantioselectivity increased to 94:6 e.r. when an alternative CPA, (*R*)-TiPSY (20 mol%) was used. Reduction of the reaction temperature to 50 °C did not significantly affect the enantioselectivity of the cyclisation to **9f** (Figure 2). The absolute sense of the enantioselectivity in the clip-cycle reaction was determined as (*S*) by the conversion of **9f** into the known methyl oxoester.^{17, 18}

Figure 2. Asymmetric ‘Clip-Cycle’ Synthesis of 3,3-Disubstituted Pyrrolidines. Yields are reported over two steps. Enantiomeric ratios were determined by chiral stationary phase HPLC. In the case of **9f** (ii) and (iii) relate to the use of (*R*)-TiPSY (20 mol%) as the catalyst or the reaction being heated to 50 °C respectively.

A similar variety of substituents were shown to be tolerated at the 2-position of the pyrrolidine¹⁷ (Figure 3). 2,2-Dimethyl substitution (**11a**, Figure 3) resulted in a pyrrolidine product with a 96:4 e.r. Surprisingly, the use of (*R*)-TiPSY resulted in a dramatic decrease in the enantioselectivity of the reaction and the yield (20% yield, 76:24 e.r., **11a**, Figure 3). Running the reaction at 50 °C, did not result in an increase in the enantioselectivity, only in a reduced conversion (34% yield, Figure 3). Changing the solvent to either DCE or toluene only led to a reduction in yield of **11a** (Figure 3).

Figure 3. Asymmetric 'Clip-Cycle' Synthesis of 2,2-Disubstituted Pyrrolidines. Yields are reported over two steps. Enantiomeric ratios were determined by chiral stationary phase HPLC. In the case of **11a** (ii) (*R*)-TiPSY (20 mol%) was used as the catalyst. In (iii-v) the reaction was heated to 50 °C. In the case of **11h** the reaction was run for 48 h.

Replacing the *p*-tolyl thioester with a mesityl thioester, also resulted in 92:8 e.r (**11b**, Figure 3). A range of cycloalkyl substituents could be introduced with no detrimental effect on the enantioselectivity of the pyrrolidine product (**11c-f**, Figure 3). However, when 4-*N*-Cbz-piperidyl or 2,2-diphenyl substitution was present the yields and enantioselectivity of the reaction were lower (**11g-h**, Figure 3). This could be due to the increased steric bulk of the diphenyl groups or the additional *N*-Cbz-group interacting with CPA catalyst. Thus, (*R*)-TRIP was still the optimal CPA for the 'clip-cycle' reaction, although the reaction is slightly less tolerant of 2,2-disubstitution than of 3,3-disubstitution.

Intrigued by the high levels of enantioselectivity, the 'clip-cycle' reaction was investigated computationally. Our working mechanistic hypothesis (Scheme 3) was that the carbamate nitrogen in **12a** directly cyclizes on to the α,β -unsaturated thioester, forming the aminoenol **14**, which then tautomerizes to give the pyrrolidine product **11a**. The aminoenol can have either *S* or *R* stereogenic center configuration, and either *E* or *Z* enol geometry depending on whether the thioester adopts an *s-cis* or *s-trans* conformation in **TS-13A**. Investigation of all 4 combinations was necessary to identify the lowest energy

pathway. Both steps could be catalyzed by the CPA. An alternative pathway proceeding *via* an iminol and corresponding cyclization transition state **TS-13B** was also briefly considered, but early model studies found that this pathway is much higher in energy and was not investigated in more detail.

Scheme 3. Mechanistic hypothesis for enantioselective CPA-catalyzed aza-Michael cyclization.

Conformational searches were done on the starting material, intermediates, products and transition-state-like structures. Selected conformers were then reoptimized at DFT level, and free energies calculated (B3LYP¹⁹/6-31G**²⁰/SMD(cyclohexane)²¹//Mo6-2X²²/def2-TZVP²³/SMD(cyclohexane), see SI for further details). Computational results (Figure 4) show that the Brønsted acid acts as a proton shuttle and catalyzes both the cyclization to give the aminoenol **14**, and its subsequent tautomerisation to the thioester **11a**. The cyclization is the rate-limiting and stereochemical-determining step. Computations indicate that the *S,E* pathway is the most favorable, and the next lowest-energy competing pathway is the enantiomeric *R,Z* pathway. The main difference between these competing transition states is the orientation of the thioester α -proton: in the *S,E* TS it is pointing away from the catalyst pocket, while in the *R,Z* TS it is pointing inwards to allow amide to approach from the other side. This steric clash is presumably the main reason for the difference in the energies. The free energies of the remaining *S,Z* and the *R,E* cyclization transition states were 27.9 kcal/mol and 29.4 kcal/mol, respectively, which is much higher than in the *S,E* and the *R,Z* pathways. Our computational study predicted formation of the (*S*)-product **11a** with 93:7 e.r. which fits well the experimentally observed value of 96:4 e.r.

Figure 4. Results from the computational study of CPA catalyzed enantioselective aza-Michael reaction

N-Methyl pyrrolidine-containing alkaloids such as (*R*)-irnidine^{24a} **16** and (*R*)-bgugaine^{24b} **17** have been shown to have a strong binding affinity for DNA,²⁵ and possess antibiotic properties against Gram +ve bacteria,^{24b} (*R*)-bgugaine is also a potent hepatotoxin.²⁶ To date there have been three racemic total syntheses²⁷ and five total syntheses of single enantiomers of these alkaloids,²⁸ where the stereochemistry was either set from the chiral pool or *via* the use of a chiral auxiliary. One report utilized a Sharpless asymmetric dihydroxylation reaction to install the required stereochemistry.^{28f} Clip-cycle pyrrolidine formation enabled the short, catalytic enantioselective total syntheses of these alkaloids (Scheme 4).

The total syntheses of both (*R*)-irnidine and (*R*)-bgugaine were achieved from a common β -homoproline intermediate **9f**. β -Homoproline thioester **9f** was synthesized in 94:6 e.r. and 71% isolated yield over two steps by the 'clip-cycle' reaction using (*R*)-TiPSY. A Liebeskind-Srogl coupling of **9f** with the appropriate alkylborane (**18** or **19**) yielded **20** or **21** in 73% and 77% yields respectively. Wolf-Kischner reduction of **20** and **21** *via* formation of the tosyl hydrazone (88% and 90% yields for the hydrazone formation from **20** and **21** respectively) was followed by reduction with NaBH₃CN and ZnCl₂ to give **22** in 40% yield and **23** in 81% yield over two steps. Finally, LiAlH₄ reduction of the Cbz groups of **22** and **23** delivered the natural products (*R*)-irnidine **16** and (*R*)-bgugaine **17** in 88% and 68% yields in 6 steps from **1f**.

Scheme 4. Total syntheses of (*R*)-irnidine and (*R*)-bgugaine^a

^aReagents and conditions: **a.** Hoveyda-Grubbs II (10 mol%), CuI, DCE, 50 °C, 14 h, 88%; **b.** (*R*)-TiPSY (20 mol%), cyclohexane, 80 °C, 81%; **c.** **18** or **19**, CuTC, Pd(PPh₃)₄ (5 mol%), Cs₂CO₃, THF, 55 °C, 20 h, 73% (**20**) 77% (**21**); **d.** TsNHNH₂, AcOH (cat.), MeOH, 24 h 88% (**20**), 90% (**21**); **e.** NaBH₃CN, ZnCl₂, MeOH, reflux, 16 h, 40% (**22**), 81% (**23**); **f.** LiAlH₄, THF, 15 h, 88% (**16**), 68% (**17**). HG-II = Hoveyda-Grubbs 2nd generation catalyst, CuTC = copper (I) thiophene-2-carboxylate.

In conclusion we have developed a modular and enantioselective Brønsted acid-catalyzed 'clip-cycle' synthesis of 2,2- and 3,3-disubstituted pyrrolidines and spiropyrrolidines, which are key units in drug discovery programs, and we have used this in the catalytic enantioselective synthesis of (*R*)-irnidine and (*R*)-bgugaine in 18% and 33% overall yields respectively. This method compares favorably with other methods for the synthesis of pyrrolidine natural products, including those which utilize aza-Michael reactions as it avoids the need for cryogenic temperatures²⁹ and yields the natural products in only 6 steps. Investigation into substrates with additional double bond substitution and kinetic resolution of racemic substrates are ongoing and will be reported in due course.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website. Full experimental procedures and analysis data for all new compounds, overview of the computational methods and results (PDF). Full set of DFT output files with optimized structures, frequencies and high-level single-point energies can be found at <https://doi.org/10.17863/CAM.50976>. Additional reference spectroscopic and reaction data can be found at DOI: 10.15124/a01fegbe-9b9e-43ac-8a21-20a9075784c5

AUTHOR INFORMATION

Corresponding Authors

paul.clarke@york.ac.uk, ke291@cam.ac.uk

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

We thank the Department of Chemistry, University of York (CJM), the Leverhulme Trust and Isaac Newton Trust (KE) for financial support. The computational work has been performed using resources provided by the Cambridge Tier-2 system operated by the University of Cambridge Research Computing Service (<http://www.hpc.cam.ac.uk>) funded by EPSRC Tier-2 capital grant EP/P020259/1.

REFERENCES

- (1) Vitaku, E.; Smith, D. T.; Njardarson, J. T. Analysis of Structural Diversity, Substitution Patterns, and Frequency of Nitrogen Heterocycles among U.S. FDA Approved Pharmaceuticals. *J. Med. Chem.* **2014**, *57*, 10257–10274.
- (2) (a) Zheng, Y.-J.; Tice, C. M. The Utilization of Spirocyclic Scaffolds in Novel Drug Discovery. *Expert Opinion on Drug Discovery*, **2016**, *11*, 831–834. (b) Zheng, Y.; Tice, C. M.; Singh, S. B. The Use of Spirocyclic Scaffolds in Drug Discovery. *Bioorg. Med. Chem. Lett.* **2014**, *24*, 3673–3682.
- (3) (a) Fremov, I. V.; Vajdos, F. F.; Borzilleri, K. A.; Capetta, S.; Chen, H.; Dorff, P. H.; Dutra, J. K.; Goldstein, S. W.; Mansour, M.; McColl, A.; Noell, S.; Oborski, C. E.; O'Connell, T. N.; O'Sullivan, T. J.; Pandit, J.; Wang, H.; Wei, B. Q.; Withka, J. M. Discovery and Optimization of a Novel Spiropyrrolidine Inhibitor of β -Secretase (BACE1) through Fragment-Based Drug Design. *J. Med. Chem.* **2012**, *55*, 9069–9088. (b) Griggs, S. D.; Thompson, N.; Tape, D. T.; Fabre, M.; Clarke, P. A. A Two-Step Synthesis of 2-Spiropiperidines. *Chem. - Eur. J.* **2017**, *23*, 9262–9265. (c) Fjølbye, K.; Marlgo, M.; Clausen, R. P.; Juhl, K. Enantioselective Fluorination of Spirocyclic β -Prolinals Using Enamine Catalysis. *Synlett*, **2017**, *28*, 425–428. (d) Thierry, T.; Lebarry, C.; Pfund, E.; Lequeux, T. Photoinduced Nonstabilized Azomethine Ylide Formation for the Preparation of Fluorine Containing Pyrrolidines. *J. Org. Chem.* **2019**, *84*, 5877–5885. (e) Carreira, E. M.; Fessard, T. C. Four-Membered Ring-containing Spirocycles: synthetic Strategies and Opportunities. *Chem. Rev.* **2014**, *114*, 8257–8322.
- (4) Bhat, C.; Tilve, S. G. Recent advances in the synthesis of naturally occurring pyrrolidines, pyrrolizidines and indolizidine alkaloids using proline as a unique chiral synthon. *RSC Adv.* **2014**, *4*, 5405–5452.
- (5) For selected examples see: (a) Gallagher, G. J.; Kerrick, S. T.; Beak, P. Enantioselective Deprotonation: The Structure and Reactivity of an Unsymmetrically Complexed Isopropyllithium/Sparteine/Et₂O Dimer. *J. Am. Chem. Soc.* **1992**, *114*, 5872–5873. (b) Beak, P.; Kerrick, S. T.; Wu, S.; Chu, J. Complex Induced Proximity Effects: Enantioselective syntheses Based on Asymmetric Deprotonations of *N*-Boc-pyrrolidines. *J. Am. Chem. Soc.* **1994**, *116*, 3231–3239. (c) Carbone, G.; O'Brien, P.; Hilmersson, G. Asymmetric Deprotonation Using *s*-BuLi or *i*-PrLi and Chiral Diamines in THF: The Diamine Matters. *J. Am. Chem. Soc.* **2010**, *132*, 15445–15450. (d) Coldham, I.; Patel, J. J.; Sanchez-Jimenez, G. Dynamic Kinetic Resolution of *N*-Boc-2-lithiopyrrolidine. *Chem. Commun.* **2005**, 3083–3085.
- (6) (a) For a review see: Adrio, J.; Carretero, J. C. Recent Advances in the Catalytic Asymmetric 1,3-Dipolar Cycloaddition of Azomethine Ylides. *Chem. Commun.*, **2014**, *50*, 12434–12446. (b) For a recent example see: Cayuelas, A.; Larrañaga, O.; Selva, V.; Nájera, C.; Akiyama, T.; Sansano, J. M.; de Cózar, A.; Miranda, J. I.; Cossío, F. P. Cooperative Catalysis with Coupled Chiral Induction in 1,3-Dipolar Cycloadditions of Azomethine Ylides. *Chem. Eur. J.* **2018**, *24*, 8092–8097.
- (7) Han, M.-Y.; Jia, J.-Y.; Wang, W. Recent Advances in Organocatalytic Asymmetric Synthesis of Polysubstituted Pyrrolidines. *Tetrahedron Letts.* **2014**, *55*, 784–794.
- (8) For reviews on general asymmetric inter- and intramolecular aza-Michael reactions see: (a) Amara, Z.; Caron, J.; Joseph, D. Recent Contributions from the Asymmetric Aza-Michael Reaction to Alkaloid Total Synthesis. *Nat. Prod. Rep.* **2013**, *30*, 1211–1225. (b) Wang, J.; Li, P.; Choy, P. Y.; Chan, A. S. C.; Kwong, F. Y. Advances and Applications in Organocatalytic Asymmetric aza-Michael Addition. *ChemCatChem* **2012**, *4*, 917–925.
- (9) Duschmalé, J.; Wennemers, H. Adapting to Substrate Challenges: Peptides as Catalysts for Conjugate Addition Reactions of Aldehydes to α,β -Disubstituted Nitroolefins. *Chem. Eur. J.* **2012**, *18*, 1111–1120.
- (10) (a) Fustero, S.; Jiménez, D.; Sánchez-Roselló, M.; del Pozo, C. Microwave-Assisted Tandem Cross Metathesis Intramolecular Aza-michael Reaction: An Easy Entry to Cyclic β -Amino Carbonyl Derivatives. *J. Am. Chem. Soc.* **2007**, *129*, 6700–6701. (b) Carlson, E. C.; Rathbone, L. K.; Yang, H.; Collett, N. D.; Carter, R. G. Improved Protocol for Asymmetric, Intramolecular Heteroatom Michael Addition Using Organocatalysis: Enantioselective Syntheses of Homoproline, Pelletierine and Homopipercolic Acid. *J. Org. Chem.* **2008**, *73*, 5155–5158.
- (11) Fustero, S.; Monteagudo, S.; Sánchez-Roselló, M.; Flores, S.; Barrio, P.; del Pozo, C. *N*-Sulfinyl Amines as a Nitrogen Source in the Asymmetric Intramolecular Aza-Michael Reaction: Total Synthesis of (-)-Pinidinol. *Chem. Eur. J.* **2010**, *16*, 9835–9845.
- (12) For representative examples see: (a) Enders, D.; Göddertz, D. P.; Beceno, C.; Raabe, G. Asymmetric Synthesis of Polyfunctionalised Pyrrolidines via a Thiourea Catalyzed Domino Mannich/Aza-Michael Reaction. *Adv. Synth. Catal.* **2010**, *352*, 2863–2868. (b) Zhu, Y.-f.; Cai, C. *N*-Heterocyclic Carbene-Catalyzed Synthesis of 3-Hydroxypyrrolidinones via a Domino Aza-Michael/Aldol Reaction. *New. J. Chem.* **2015**, *39*, 5014–5017.
- (13) For recent examples see: (a) Lee, S.; Lei, H.; Rovis, T. A Rh(III)-Catalyzed Formal [4+1] approach to Pyrrolidines from Unactivated Terminal Alkenes and Nitrene Sources. *J. Am. Chem. Soc.* **2019**, *141*, 12536–12540. (b) Jiang, H.; Gao, H.; Liu, B. Wu, W. Palladium-Catalyzed Selective Aminoamidation and Aminocyanation of Alkenes using Isonitrile as Amide and Cyanide sources. *Chem. Commun.* **2014**, *50*, 15348–15351. (c) Farndon, J. J.; Ma, X.; Bower, J. F. Transition Metal Free C–N Bond Forming Dearomatizations and Aryl C–H Aminations by *in situ* Release of a Hydroxylamine-Based Aminating Agent. *J. Am. Chem. Soc.* **2017**, *139*, 14005–14008. (d) Espinosa, M.; Noda, H.; Shibasaki, M. Synthesis of Unprotected Spirocyclic β -Prolines and β -Homoprolines by Rh-Catalyzed C–H Insertion. *Org. Lett.* **2019**, *21*, 9296–9299. (e) Wang, Y.-Y.; Bode, J. W. Olefin Amine (OLA) Reagents for the Synthesis of Bridged Bicyclic and Spirocyclic Saturated *N*-Heterocycles by Catalytic Hydrogen Atom Transfer (HAT) Reactions. *J. Am. Chem. Soc.* **2019**, *141*, 9739–9745. (f) Flodén, N. J.; Trowbridge, A.; Willcox, D.; Walton, S. M.; Kim, Y.; Gaunt, M. J. Streamlined Synthesis of C(sp³)-Rich *N*-Heterospirocycles Enabled by Visible-Light-Mediated Photocatalysis. *J. Am. Chem. Soc.* **2019**, *141*, 8426–8430. (g) Fominova, K.; Diachuk, T.; Sadova, I. V.; Mykhailiuk, P. K. Synthesis of 2,2-Disubstituted Spirocyclic Pyrrolidines by Intramolecular Dieckmann Condensation. *Eur. J. Org. Chem.* **2019**, 3553–3559. (h) Verma, P.; Richter, J. M.; Chekshin, N.; Qiao, J. X.; Yu, J. -Q. Iridium(I)-Catalyzed α -C(sp³)-H Alkylation of Saturated Azacycles. *J. Am. Chem. Soc.* **2020**, *142*, 5117–5125.
- (14) (a) Fang, X.; Wang, C.-J. Catalytic Asymmetric Construction of Spiropyrrolidines via 1,3-Dipolar Cycloaddition of Azomethine Ylides. *Org. Biomol. Chem.* **2018**, *16*, 2591–2601. (b) Ma, X.; Hazelden, I. R.; Langer, T.; Munday, R. H.; Bower, J. F. Enantioselective Aza-Heck Cyclizations of *N*-(Tosyloxy)carbamates: Synthesis of Pyrrolidines and Piperidines. *J. Am. Chem. Soc.* **2019**, *141*, 3356–3360.
- (15) (a) Liu, H.; Zeng, C.; Guo, J.; Zhang, M.; Yu, S. Enantioselective Synthesis of 2-substituted Pyrrolidines via Domino Cross Metathesis/Intramolecular Aza-Michael Addition. *RSC Advances*, **2013**, *3*, 1666–1668. (b) Kim, H.; Kim, Y.; Kim, S.-G. Or-

- ganocatalytic Asymmetric Mannich/Aza-Michael Cascade Reaction of δ -Formyl- α,β -unsaturated Ketones with Cyclic *N*-sulfimines: Enantioselective Synthesis of Benzosulfamidate-fused Pyrrolidines. *J. Org. Chem.* **2017**, *82*, 8179–8185.
- (16) Ermanis, K.; Hsiao, Y.-T.; Kaya, U.; Jeuken, A.; Clarke, P. A. The Stereodivergent Formation of 2,6-*cis* and 2,6-*trans* Tetrahydropyrans: Experimental and Computational Investigation of the Mechanism of a Thioester Oxy-Michael Cyclisation. *Chem. Sci.* **2017**, *8*, 482–490.
- (17) Procedures for the synthesis of the 2,2- and 3,3-disubstituted substrates are included in the supporting information
- (18) Masson, G.; Riota, S.; Pardo, D. G.; Cossy, J. Access to Enantio-enriched Substituted α -Trifluoromethyl Azepanes from L-Proline. *Org. Lett.* **2018**, *20*, 5019–5022.
- (19) (a) Becke, A. D. Density-Functional Exchange-Energy Approximation with Correct Asymptotic Behavior. *Phys. Rev. A*, **1988**, *38*, 3098–3100; (b) Lee, C.; Yang, W.; Parr, R. G. Development of the Colle-Salvetti Correlation-Energy Formula into a Functional of the Electron Density. *Phys. Rev. B*, **1988**, *37*, 785–789 (c) Becke, A. D. Density-Functional Thermochemistry. III. The Role of Exact Exchange *J. Chem. Phys.*, **1993**, *98*, 5648–5652.
- (20) (a) Hehre, W. J.; Ditchfield, R.; Pople, J. A. Self-Consistent Molecular Orbital Methods. XII. Further Extensions of Gaussian-Type Basis Sets for Use in Molecular Orbital Studies of Organic Molecules. *J. Chem. Phys.*, **1972**, *56*, 2257–2261. (b) Hariharan, P. C.; Pople, J. A. The Influence of Polarization Functions on Molecular Orbital Hydrogenation Energies. *Theor. Chem. Acc.*, **1973**, *28*, 213–222.
- (21) Marenich, A. V.; Cramer, C. J.; Truhlar, D. G. Universal Solvation Model Based on Solute Electron Density and a Continuum Model of the Solvent Defined by the Bulk Dielectric Constant and Atomic Surface Tensions. *J. Phys. Chem. B*, **2009**, *113*, 6378–6396.
- (22) Zhao, Y.; Truhlar, D. The Mo6 Suite of Density Functionals for Main Group Thermochemistry, Thermochemical Kinetics, Noncovalent Interactions, Excited States, and Transition Elements: Two New Functionals and Systematic Testing of Four Mo6-Class Functionals and 12 other Functionals. *Theor. Chem. Acc.*, **2008**, *120*, 215–241.
- (23) (a) Weigend, F.; Ahlrichs, R. Balanced Basis Sets of Split Valence, Triple Zeta Valence and Quadruple Zeta Valence Quality for H to Rn: Design and Assessment of Accuracy. *Phys. Chem. Chem. Phys.*, **2005**, *7*, 3297–3305. (b) Weigend, F. Accurate Coulomb-Fitting Basis Sets for H to Rn. *Phys. Chem. Chem. Phys.*, **2006**, *8*, 1057–1065.
- (24) (a) Melhaoui, A. A New Toxic Alkylpyrrolidine Alkaloid from *Arisarum vulgare*. *Planta Med.* **1998**, *64*, 476–477. (b) Melhaoui, A.; Mallea, M.; Jossang, A.; Bodo, B. Antibiotic and Antifungal Pyrrolidine Alkaloids, from *Arisarum vulgare*. *Nat. Prod. Lett.* **1993**, *2*, 237–242.
- (25) Melhaoui, A.; Belouali, H. DNA Affinity of Active Alkaloids from *Arisarum vulgare* Targ. *J. Ethnopharmacology*, **1998**, *62*, 67–71.
- (26) Rakba, N.; Melhaoui, A.; Loyer, P.; Delcros, J.; Morel, I.; Lescoat, G. Bgugaine, a Pyrrolidine Alkaloid from *Arisarum vulgare* is a Strong Hepatotoxin in Rat and Human Liver Cell Culture. *Toxicol. Lett.*, **1994**, *104*, 239–248.
- (27) (a) Miyata, O.; Takahashi, S.; Tamura, A.; Ueda, M.; Naito, T. Novel Synthesis of Substituted Pyrrolidines and Piperidines via Radical Addition-Ionic Cyclization Reaction of Oxime Ethers. *Tetrahedron*, **2008**, *64*, 1270–1284. (b) Xiao, K.-J.; Wang, Y.; Ye, K.-Y.; Huang, P.-Q. Versatile One-Pot Reductive Alkylation of Lactams/Amides via Amide Activation: Application to the Concise Syntheses of Bioactive Alkaloids ($\pm$)-Bgugaine, ($\pm$)-Coniine, (+)-Preussina, and (–)-Cassine. *Chem. Eur. J.*, **2010**, *16*, 12792–12797. (c) Zhang, W.; Werness, J. B.; Tang, W. Base-Catalysed Intramolecular Hydroamination of Conjugated Enynes. *Org. Lett.* **2008**, *10*, 2023–2026. (d) Zhang, W.; Werness, J. B.; Tang, W. Intramolecular Hydroamination of Conjugated Enynes. *Tetrahedron* **2009**, *65*, 3090–3095.
- (28) (a) Jossang, A.; Melhaoui, A.; Bodo, B. Enantioselective Total Synthesis of Irniine and Bgugaine, Bioactive 2-Alkylpyrrolidine Alkaloids. *Heterocycles*, **1996**, *43*, 755–765. (b) Enders, D.; Thiebes, C. Enantioselective Synthesis of Both Enantiomers of Bgugaine Employing the SAMP-Hydrazone Method. *J. Indian Chem. Soc.*, **1999**, *76*, 561–564. (c) Enders, D.; Thiebes, C. Efficient Stereoselective Syntheses of Piperidine, Pyrrolidine, and Indolizidine alkaloids. *Pure Appl. Chem.*, **2001**, *73*, 573–578. (d) Majik, M. S.; Parameswaran, P. S.; Tilve, S. G. A Concise Synthesis of (*R*)-Bgugaine, a Pyrrolidine Alkaloid from *Arisarum vulgare*. *J. Chem. Res.* **2008**, 121–122. (e) Shu, C.; Li, M.-Q.; Wang, S.-S.; Li, L.; Ye, L.-W. Gold-Catalysed Oxidative Cyclisation of Chiral Homopropargyl Amides: Synthesis of Enantioenriched γ -Lactams. *J. Org. Chem.* **2013**, *78*, 3292–3299. (f) Takahata, H.; Ihara, K.; Kubota, M.; Momose, T. New Asymmetric Synthesis of Enantiomeric Pairs of the 2-Substituted Pyrrolidine: Bgugaine and Irniine. *Heterocycles*, **1997**, *46*, 349–356.
- (29) Fustero, S.; Jiménez, D.; Moscardó, J.; Catalán, S.; del Pozo, C. Enantioselective Organocatalytic Intramolecular Aza-Michael Reaction: a Concise Synthesis of (+)-Sedamine, (+)-Allosedamine, and (+)-Coniine. *Org. Lett.* **2007**, *9*, 5283–5286.