

This is a repository copy of *The predictors of cigarette smoking, smokeless tobacco consumption and use of both forms in adolescents in South Asia : a secondary analysis of the Global Youth Tobacco Surveys*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/166412/>

Version: Accepted Version

---

**Article:**

Mishu, Masuma Pervin [orcid.org/0000-0002-6545-9117](https://orcid.org/0000-0002-6545-9117), Siddiqui, Faraz [orcid.org/0000-0002-2253-3911](https://orcid.org/0000-0002-2253-3911), Shukla, Radha et al. (3 more authors) (2021) The predictors of cigarette smoking, smokeless tobacco consumption and use of both forms in adolescents in South Asia : a secondary analysis of the Global Youth Tobacco Surveys. *Nicotine and tobacco research*. pp. 956-965. ISSN 1469-994X

<https://doi.org/10.1093/ntr/ntaa202>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

Figure 1: Number of study participants in GYTS survey from each country and data analysed

