

This is a repository copy of *Insights into the composition and structural chemistry of gallium(I) triflate*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/166429/>

Version: Accepted Version

Article:

Boronski, Josef, Stevens, Matthew, van IJzendoorn, Bono et al. (2 more authors) (2020) Insights into the composition and structural chemistry of gallium(I) triflate. *Angewandte Chemie International Edition*. ISSN: 1433-7851

<https://doi.org/10.1002/anie.202010837>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Insights into the composition and structural chemistry of gallium(I) triflate

Josef T. Boronski,^[a] Matthew Stevens,^[a] Bono van IJzendoorn,^[a] Adrian C. Whitwood^[a] and John M. Slattery^{*[a]}

Abstract: “GaOTf” is a simple, convenient source of low-valent gallium for synthetic chemistry and catalysis. However, little is currently known about its composition or reactivity. In this work, ⁷¹Ga NMR shows the presence of [Ga(arene)_n]⁺ salts on oxidation of Ga metal with AgOTf in arene solvents. However, a more complex picture of speciation is uncovered by X-ray diffraction studies. In all cases, mixed-valence compounds containing Ga-arene and Ga-OTf coordination motifs, in addition to an unusual “naked” [Ga]⁺ ion, are found. Addition of 18-crown-6 allows for the isolation of a discrete Ga(I) crown complex. Evidence of a potential intermediate in the formation of “GaOTf” has been isolated in the form of the bimetallic silver(I)/gallium(I) cluster anion [Ag₄{Ga(OTf)₃}₄(μ-Ga)₆(OTf)₄]²⁻.

Gallium occupies a fascinating position in group 13. Much of its chemistry is dominated by the +3 oxidation state, like the lighter elements, but it also displays an extensive range of low oxidation state compounds. There have been a number of breakthroughs in the development of starting materials for low-valent gallium chemistry, each of which has advantages and disadvantages. True Ga(I)-halides are thermodynamically unstable, and while they can be trapped and utilized, this requires specialist equipment.^[1] Ga₂Cl₄ (formulated as Ga⁺[GaX₄]⁻) has been known for many years, and has an extensive coordination chemistry with arene ligands,^[2] but often undergoes dis- or comproportionation on reaction with other species. “GaI” can be an excellent source of Ga(I), but like Ga₂Cl₄ also has a tendency towards unexpected behavior.^[3] [Ga₂C₅Me₅][B(Ar^F)₄]⁻ salts are capable of delivering Ga⁺ ions as ligands to suitable transition metal complexes.^[4] [Ga(arene)_n]⁺[Al(OC(CF₃)₃)₄]⁻ (n = 2, 3) salts are well-behaved sources of Ga(I), but these anions are still not as widely used as simple weakly coordinating anions (WCAs) such as triflate.^[5] More recently, Ga(I) species have been accessed via reductive elimination of H₂ from Ga(III) dihydrides.^[6] In addition, a family of Ga-based carbenoids, with differing charges and ring sizes, have emerged as important building blocks in low-valent Ga chemistry and have been used as ligands in coordination chemistry.^[7]

There are a growing number of reports of Ga(I) involvement in catalysis and the development of new, more accessible, well-behaved low-valent gallium sources is particularly important to




Figure 1. Low-valent-Ga catalyzed reaction between acetals, ketals or amins and allyl boronic esters reported by Schneider (top)^[8] and this work (below).

allow this area to grow. [Ga(arene)_n]⁺ salts catalyse the polymerization of alkenes,^[5e, 5f, 6] and have been shown to be active catalysts for cycloisomerisation, transfer hydrogenation and reductive hydroarylation reactions.^[9] There is also evidence for Ga(I) involvement in important mechanistic steps that underpin many catalytic reactions, e.g. reductive elimination, suggesting exciting possibilities for the Ga(I)/Ga(III) redox couple in other catalytic reactions.^[3c, 6] In related work, the use of Ga(0) in synthesis and materials chemistry has received attention.^[10]

Schneider *et al.* recently reported the preparation of a new low-valent gallium reagent “GaOTf” (OTf = [O₃SCF₃]⁻) using a similar route to the preparation of [Ga(arene)_n]⁺[Al(OC(CF₃)₃)₄]⁻ salts,^[5a] where gallium metal is oxidized by AgOTf in arene solvents, or dioxane in the presence of 18-crown-6.^[8] “GaOTf” solutions were then used as ambiphilic catalysts for C-C bond forming reactions (figure 1). Control experiments with Ga(III) salts, Ag(0) or other metal triflates showed that the presence of low oxidation-state Ga was important for catalytic activity. ⁷¹Ga NMR spectroscopy suggested the formation of a [Ga(18-crown-6)][OTf]·(dioxane)_n complex under catalytic conditions and other stoichiometric studies allowed a catalytic cycle involving Ga(I)-alkoxide and -allyl intermediates to be proposed. However, speciation in “GaOTf”, particularly in arene solvents, was not explored in detail. Given the potential wider uses of “GaOTf” as one of the most easily accessible sources of Ga(I) yet reported, this paper explores the composition and structural chemistry of the “GaOTf” system in detail for the first time.

Reaction of AgOTf and Ga metal in toluene under ultrasonic activation (see fig. 1 for summary) led to the immediate formation of a pale-yellow solution and after ca. 1.5 hours of ultrasonication a dense black precipitate. After filtration, ⁷¹Ga NMR spectroscopy showed a single, relatively sharp peak at -692 ppm, which is characteristic of gallium(I) arene complexes.^[2] Storage of the solution overnight at -20 °C led to the formation of pale brown crystals, which X-ray crystallography revealed to contain [Ga][Ga(C₆H₅Me)₂]₂[Ga₃(OTf)₈] (1) (figure 2).

[a] Mr Josef T. Boronski, Mr Bono van IJzendoorn, Mr Matthew Stevens, Dr Adrian C. Whitwood and Dr John M Slattery
Department of Chemistry
The University of York
Heslington
York, YO10 5DD, UK
E-mail: john.slattery@york.ac.uk

Supporting information for this article is given via a link at the end of the document.




Figure 2. Crystal structure of $[\text{Ga}][\text{Ga}(\text{C}_6\text{H}_5\text{Me})_2]_2[\text{Ga}_3(\text{OTf})_6]$ (**1**). Monoclinic, $P2_1/c$, 110K, $R_1 = 0.0784$, $wR_2(\text{all}) = 0.2303$. Some toluene molecules and one $[\text{OTf}]^-$ are disordered over two positions. Disordered parts are omitted for clarity. Selected bond lengths (Å) and angles ($^\circ$): Ga(1)–Ga(2) 2.379(1), Ga(2)–Ga(3) 2.377(1), Ga(4)–C₆(cent) 2.837(8) and 2.861(4), Ga(6)–C₆(cent) 2.809(12) and 2.831(5), Ga(1)–Ga(2)–Ga(3) 140.31(6), C₆(cent)–Ga(4)–C₆(cent) 121.30(18), C₆(cent)–Ga(4)–C₆(cent) 126.95(30).

As anticipated, and consistent with the ^{71}Ga NMR spectrum, **1** contains two bent-sandwich gallium(I) toluene cations, $[\text{Ga}(\text{C}_6\text{H}_5\text{Me})_2]^+$, which have very similar parameters to the bis-benzene complex reported by Schmidbaur $[\text{Ga}(\text{C}_6\text{H}_6)_2][\text{GaCl}_4]$ (av. Ga–C₆ centroid in **1** = 2.835 vs. 2.844 Å, av. C₆(cent)–Ga–C₆(cent) in **1** = 124 vs. 124 $^\circ$).^[11] In addition, an unusual “naked” $[\text{Ga}]^+$ ion, which is stabilized by Ga \cdots O and Ga \cdots F contacts with the anions, rather than arene coordination, is found in the asymmetric unit.

The counterions in **1** are not simple $[\text{OTf}]^-$ ions, but the previously unknown trianion $[\text{Ga}_3(\text{OTf})_6]^{3-}$. This is composed of a catenated $[\text{Ga}_3]^{5+}$ core, coordinated by eight $[\text{OTf}]^-$ and is related to the $\text{Ga}_3\text{X}_5\text{L}_3$ motif identified by Schnöckel in $[\text{Ga}_3\text{I}_5(\text{PET}_3)_3]$,^[12] the monoanion $[\text{Ga}_3(\text{OTf})_6(\text{GaCp}^*)_2]^-$ reported by Linti^[13] and Baines' $[\text{Ga}_3\text{Cl}_4(\text{crypt-222})]^+$ cation.^[14] Indeed, these species can all be viewed as $\text{Ga}_3\text{X}_5\text{L}_3$ -type compounds, where X = I, Cl or OTf and L is either a neutral (PET_3 , GaCp^* or crypt-222) or anionic ($[\text{OTf}]^-$) ligand. In Baines' system one X^- has been lost and the $[\text{Ga}_3]^{5+}$ core forms an additional bond to the cryptand to compensate for this. Comparing the structural parameters of $[\text{Ga}_3(\text{OTf})_6]^{3-}$ with other $\text{Ga}_3\text{X}_5\text{L}_3$ -type compounds shows that the average Ga–Ga bond lengths are shorter in $[\text{Ga}_3(\text{OTf})_6]^{3-}$ (2.374 Å, compared to 2.456 Å in $[\text{Ga}_3\text{I}_5(\text{PET}_3)_3]$, 2.426 Å in $[\text{Ga}_3(\text{OTf})_6(\text{GaCp}^*)_2]^-$ and 2.416 Å in $[\text{Ga}_3\text{Cl}_4(\text{crypt-222})]^+$). These species are all mixed-valence Ga compounds and different oxidation state models have been proposed. In $[\text{Ga}_3(\text{OTf})_6]^{3-}$ a Ga(II)–Ga(I)–Ga(II) model, in analogy to $[\text{Ga}_3\text{I}_5(\text{PET}_3)_3]$, would seem appropriate. However, recent XAS studies suggest that assignment of formal oxidation states in such species is challenging and a Ga(I)–Ga(III)–Ga(I) model is also feasible.^[14b] Other examples of catenated Ga and In species, some with quite extended structures, have been reported and are of note.^[7e, 12, 15]

In an attempt to prevent disproportionation and the formation of higher oxidation state species, “GaOTf” was prepared as described above, but with the addition of four equivalents of hexamethylbenzene, as a more π -basic arene.^[16] The ^{71}Ga NMR spectrum after ultrasonication and filtration featured a single peak at –707 ppm. Colourless crystals were grown by storage of the solution at –20 $^\circ\text{C}$ and X-ray diffraction




Figure 3. Crystal structure of $[\text{Ga}(\text{C}_6\text{H}_5\text{Me})(\text{C}_6\text{Me}_6)]_2[\text{Ga}_2(\text{OTf})_6]$ (**2**). Monoclinic, $P2_1/n$, 110K, $R_1 = 0.0861$, $wR_2(\text{all}) = 0.2444$. The asymmetric unit contains half of the molecular formula, with the other half related to this by a centre of inversion. A toluene molecule in the lattice, disordered over two positions, is omitted for clarity. Selected bond lengths (Å) and angles ($^\circ$): Ga(1)–Ga(1') 2.355(6), Ga(2)–C₆Me₆(cent) 2.525(9), Ga(2)–C₆H₅Me(cent) 2.813(11), C₆(cent)–Ga(2)–C₆(cent) 134.97(30).

studies showed these to have the composition $[\text{Ga}(\text{C}_6\text{H}_5\text{Me})(\text{C}_6\text{Me}_6)]_2[\text{Ga}_2(\text{OTf})_6]$ (**2**) (figure 3).

The $[\text{Ga}(\text{C}_6\text{H}_5\text{Me})(\text{C}_6\text{Me}_6)]^+$ units in **2** are rare examples of mixed bis(arene) gallium(I) complexes.^[16] The Ga–(C₆Me₆) C₆ centroid distance is one of the shortest known, at 2.525 Å.^[2, 17] This is 0.288 Å shorter than the Ga–(C₆H₅Me) C₆ centroid distance (2.813 Å) in **2**, suggesting a significantly enhanced interaction between Ga and the more π -basic arene. Shorter Ga–arene distances have been observed in the mono-arene salt $[\text{Ga}(\text{C}_6\text{Me}_6)][\text{Al}(\text{OC}(\text{CF}_3)_3)_4]$ (2.262 Å), but are otherwise rare.^[17a, 18] The effects of the steric bulk of the C₆Me₆ ligand are clearly evident in the arene interplanar angles in **2**, which are considerably smaller (av. 43 $^\circ$) than those in **1** (av. 56 $^\circ$). Bent-sandwich gallium(I) complexes of bulky arenes have previously been seen to display small interplanar angles, as they are forced into a more linear geometry in order to reduce the steric interaction between the two coordinating arenes.^[16]

As in **1**, the counter anion in **2** is not a simple $[\text{OTf}]^-$, but rather the Ga(II) dianion $[\text{Ga}_2(\text{OTf})_6]^{2-}$.^[19] While the formal oxidation states of Ga in this anion are different to those in $[\text{Ga}_3(\text{OTf})_6]^{3-}$ their structural parameters are similar, with average Ga–Ga bond lengths of 2.376 Å and 2.355 Å respectively. Related, halide-containing anions such as $[\text{Ga}_2\text{I}_6]^{2-}$, which is thought to be a major component of Green's “GaI”,^[3] have also been seen in similar systems. Other Ga(II) compounds that include Ga–Ga bonds, as here, have been known for some time.^[12, 20] More recently, monometallic Ga(II) radicals have also been isolated.^[21] The $[\text{Ga}_2(\text{OTf})_6]^{2-}$ and $[\text{Ga}_3(\text{OTf})_6]^{3-}$ polyanions seen here are presumably formed via partial disproportionation of Ga(I) salts formed after oxidation of Ga by Ag^+ . This suggests that $[\text{OTf}]^-$ is not a suitable anion for stabilizing Ga(I) arene complexes. The larger, highly fluorinated polyanionic species can be viewed as *in situ* generated WCAs, which are less nucleophilic than $[\text{OTf}]^-$ and as such are better able to stabilize $[\text{Ga}(\text{arene})_n]^+$ ions, in analogy to the large $[\text{Al}(\text{OC}(\text{CF}_3)_3)_4]^-$ WCA.

It can now be seen that there are several isolable low-valent-gallium-containing species, both cationic and anionic, within “GaOTf” solutions. The composition of the crystalline material is only a snapshot of some of the potential species




Figure 4. Crystal structure of $[\text{Ga}(\text{18-crown-6})(\text{OTf})]$ (**3**). Monoclinic, $P2_1/n$, 110K, $R_1 = 0.0214$, $wR_2(\text{all}) = 0.0564$. Selected bond lengths (Å) and angles ($^\circ$): Ga(1)-O(1) 2.137(1), Ga(1)-O(1)-S(1) 149.71(8).

present in solution. What is clear, however, is that “GaOTf” is more complex than it might initially appear and multiple Ga coordination environments and oxidation states are present. This realization is very important for the interpretation of catalytic studies involving “GaOTf”, as the catalytically active species could involve several possible Ga oxidation states.

Schneider *et al.* showed that addition of 18-crown-6 to “GaOTf” improved the yields in their catalytic C-C bond forming reactions.^[6] In order to investigate the potential for low-valent Ga-crown complex formation in this system, “GaOTf” solutions were prepared in toluene, as described above, filtered and the filtrate added to a solution containing 18-crown-6. In suitably concentrated solutions, this led to the spontaneous formation of colorless, block-like crystals. This was encouraged by low temperature at the time of filtration. X-ray diffraction studies revealed these to contain $[\text{Ga}(\text{18-crown-6})(\text{OTf})]$ (**3**) (figure 4). Compound **3** is only the second structurally authenticated Ga(I)-18-crown-6 complex, the first being $[\text{Ga}(\text{18-crown-6})(\text{C}_6\text{H}_5\text{F})_2][\text{Al}(\text{OC}(\text{CF}_3)_3)_4]$.^[5b] However, these complexes differ significantly, as **3** features tight ion pairing, whereas $[\text{Ga}(\text{18-crown-6})(\text{C}_6\text{H}_5\text{F})_2][\text{Al}(\text{OC}(\text{CF}_3)_3)_4]$ displays no contacts between the gallium cation and the more weakly coordinating alkoxyaluminate anion. Instead, weak Ga-C₆H₅F interactions above and below the plane of the crown complete the coordination sphere around Ga. Coordination of [OTf]⁻ to Ga in **3** appears to lead to a weakening of the Ga-crown interactions, with average Ga-O(crown) distances of 2.840 and 2.800 Å for **3** and $[\text{Ga}(\text{18-crown-6})(\text{C}_6\text{H}_5\text{F})_2][\text{Al}(\text{OC}(\text{CF}_3)_3)_4]$ respectively. Complex **3** can be compared to the isostructural indium salt $[\text{In}(\text{18-crown-6})(\text{OTf})]$.^[22] In this case, the average In-O(crown) distance is longer (2.874 Å) than in **3**, suggesting that the larger In⁺ ion sits slightly above the crown, compared to Ga. The Ga-OTf distance in **3** (2.137 Å) is 0.233 Å shorter than the In-OTf distance in $[\text{In}(\text{18-crown-6})(\text{OTf})]$, closely mirroring the difference between the ionic radii of gallium(I) and indium(I) (0.23 Å).^[23] A related $[\text{Ga}(\text{12-crown-4})]^+$ salt has also been reported in a recent PhD thesis and coordination of Ga⁺ by the tridentate PMDETA ligand, somewhat related to crown ethers, is also noted for comparison.^[24]

Unlike in **1** and **2**, the presence of 18-crown-6 in **3** appears to facilitate the formation of a Ga(I) salt with a discrete [OTf]⁻




Figure 5. Optimised structure of **3** showing the HOMO (-6.24 eV, at the (RI-)PBE0/def2-TZVPP/(RI-)BP86/SV(P) level).

anion, rather than a complex Ga-containing polyanion. This may be because the crown offers additional stability to Ga(I) towards disproportionation, which could explain its positive impact in Schneider's catalysis.^[6] Indeed, in our hands **3** is considerably more stable than solutions from which **1** or **2** are isolated, which readily disproportionate over time or on removal of the solvent.

In order to explore the electronic structure of **3**, DFT studies were performed (see ESI for details). The optimized structure of **3** (at the (RI-)BP86/SV(P) level) is shown in figure 5. Comparisons between **3** and other gallium(I)-species, such as GaCp*, can be drawn from analysis of the DFT data.^[4f] Natural bond orbital (NBO) calculations (at the (RI-)PBE0/def2-TZVPP/(RI-)BP86/SV(P) level) suggest that there is a small amount of *sp* character to the Ga(I) lone pair (3 % *p* character), which gives some directionality, as shown in figure 5. This is very similar to the hybridization in GaCp* (4 % *p* character), which has an established coordination chemistry as a ligand to other metals, and suggests that **3** may prove to be an interesting new low-valent Ga-based ligand, an area under investigation in our lab.

In order to explore speciation during the formation of “GaOTf” at shorter time periods, a toluene solution of AgOTf and a bead of Ga was sonicated for less than one hour. Filtration produced an orange solution, from which block-like orange crystals were formed after storage at -20 °C for four days. X-ray diffraction studies revealed these to be salt (**4**), which is composed of two $[\text{Ga}(\text{C}_6\text{H}_5\text{Me})_2]^+$ ions, the novel and unusual Ga-Ag cluster ion $[\text{Ag}_4\{\text{Ga}(\text{OTf})_3\}_4(\mu\text{-Ga})_6(\text{OTf})_4]^{2-}$ (figure 6) and a toluene molecule of crystallization. In light of the presence of both Ag(I) and low-valent Ga species in **4**, we speculate that this may represent an intermediate species present at short reaction times during the formation of **1** or related systems. It is also a compound of significant fundamental interest and will be discussed below.

Although the overall XRD data quality for **4** are relatively poor, due to extensive disorder in the $[\text{Ga}(\text{C}_6\text{H}_5\text{Me})_2]^+$ ions and toluene of crystallization, they are sufficient to confirm the composition of the salt and some discussion of the cluster anion, which is quite well resolved, is reasonable. The $[\text{Ag}_4\{\text{Ga}(\text{OTf})_3\}_4(\mu\text{-Ga})_6(\text{OTf})_4]^{2-}$ anion in **4** can be viewed as being formed from a tetrahedral $[\text{Ag}_4]^{4+}$ core, coordinated by four $[\text{Ga}(\text{OTf})_3]^{2-}$ ligands - one at each vertex. Bridging Ga⁺ ions coordinate each of the six edges of the tetrahedral $[\text{Ag}_4]^{4+}$ core,




Figure 6. Molecular structure of the cluster anion from $[\text{Ga}(\text{C}_6\text{H}_5\text{Me})_2]_2[\text{Ag}_4\{\text{Ga}(\text{OTf})_3\}_4(\mu\text{-Ga})_6(\text{OTf})_4]\cdot\text{C}_6\text{H}_5\text{Me}$ (**4**). Monoclinic, $\text{C}2/c$, 110K, $R_1 = 0.0625$, $wR_2(\text{all}) = 0.1828$. Selected bond lengths (Å): av. Ag–Ag 2.871, av. Ag–Ga_(vertex) 2.468, av. Ag–Ga_(edge) 2.631.

forming contacts with oxygen atoms from the $[\text{Ga}(\text{OTf})_3]^{2-}$ units. The bridging Ga* are further coordinated by the remaining four triflate ions, which sit roughly above each face of the tetrahedral $[\text{Ag}_4]^{4+}$ core. A simplified illustration is shown in figure 7.




Figure 7. Simplified structure for $[\text{Ag}_4\{\text{Ga}(\text{OTf})_3\}_4(\mu\text{-Ga})_6(\text{OTf})_4]^{2-}$.

Although uncommon, clusters based around a tetrahedral $[\text{Ag}_4]^{4+}$ core, with vertex-capping or bridging ligands are not without precedent (e.g. $[\text{Ag}_4(\mu_3\text{-I})_4]^{4-}$ and $[\text{Ag}_4(\text{TeC}_4\text{H}_3\text{S})_6]^{2-}$).^[25] Most are based around conventional donor ligands such as halides, S- or Te-based ligands, rather than the unusual low-valent Ga ligands seen here. However, low oxidation state main-group ligands based on stanna-closo-dodecaborate dianions have been found to stabilize an $[\text{Ag}_4]^{4+}$ cluster in $[\{\text{Ag}(\mu_3\text{-SnB}_{11}\text{H}_{11})(\text{PMe}_3)\}_4]^{4-}$.^[26] The Ag–Ag bonds within **4** (av. 2.87 Å) are amongst the shortest found for structures of this type on the Cambridge Crystallographic Database (av. for ten structures containing $[\text{Ag}_4]^{4+}$ tetrahedra is 3.110 Å). Compound **4** contains only the fourth example of structurally authenticated Ag–Ga bonds. The average Ag– $[\text{Ga}(\text{OTf})_3]^{2-}$ bond length of 2.47 Å is shorter than the Ag–Ga bond lengths found in $[\text{Ag}(\text{GaCp}^*)_4]^+$ (average 2.519 Å),^[27] but longer than those found within the complexes $[\text{Ag}\{\text{Ga}^{\text{DipDAB}}\}(\text{IMes})]$ (2.416 Å) and $[\text{Ag}\{\text{Ga}^{\text{DipDAB}}\}(\text{IDip})]$ (2.411 Å); where $\text{DipDAB} = [(\text{DipNCH})_2]^{2-}$, $\text{IMes} = \text{:C}(\text{MesNCH})_2$, $\text{IDip} = \text{:C}(\text{DipNCH})_2$, $\text{Dip} = 2,6\text{-diisopropylphenyl}$ and $\text{Mes} =$

mesityl.^[28] The Ga-containing N-heterocyclic carbene analogue $[\text{Ga}^{\text{DipDAB}}]^-$ is known to be a strong donor ligand, GaCp^* less so, and so based on Ag–Ga bond lengths alone it appears that the donor properties of $[\text{Ga}(\text{OTf})_3]^{2-}$ lie in between that of GaCp^* and $[\text{Ga}^{\text{DipDAB}}]^-$.




Figure 8. The structure of $[(\text{Cp}^*\text{Ga})\text{Cu}(\mu\text{-GaCp}^*)_3\text{Cu}\{\text{Ga}(\text{OTf})_3\}]$ (**5**).^[29]

Only one example of a complex that could be considered to involve the $[\text{Ga}(\text{OTf})_3]^{2-}$ motif has previously been reported, $[(\text{Cp}^*\text{Ga})\text{Cu}(\mu\text{-GaCp}^*)_3\text{Cu}\{\text{Ga}(\text{OTf})_3\}]$ (**5**, figure 8).^[29] The authors formulate this as a Lewis acid/base adduct between $[(\text{Cp}^*\text{Ga})\text{Cu}(\mu\text{-GaCp}^*)_3\text{Cu}]$, comprising two Cu(0) centers, and Ga(III) triflate. An alternative view, in light of our analysis of compound **4**, is that **5** could involve two Cu(I) centres, with the $[(\text{Cp}^*\text{Ga})\text{Cu}(\mu\text{-GaCp}^*)_3\text{Cu}]^{2+}$ ion coordinated by $[\text{Ga}(\text{OTf})_3]^{2-}$. As noted above, for $\text{Ga}_3\text{X}_5\text{L}_3$, it is challenging to unambiguously assign oxidation states to such species.^[14b] However, some features of **5** suggest that a Ga(I)/Cu(I) model is a plausible alternative formulation. The Cu– $\{\text{Ga}(\text{OTf})_3\}$ distance (2.2906 Å) is shorter than the terminal Cu–GaCp* distance (2.3268 Å). This is consistent with the shorter Ag–Ga distances in Ag– $[\text{Ga}(\text{OTf})_3]^{2-}$ complexes compared to Ag–GaCp* complexes, as described above. The average Ga–OTf distances for the “ $\text{Ga}(\text{OTf})_3$ ” unit in **5** (1.975 Å) are more similar to those in **4** (1.996 Å) than to the shorter Ga–OTf distances in Ga(III) systems such as $[\text{Ga}(\text{OTf})_3(\text{THF})_3]$ (1.945 Å).^[13] Finally, NBO calculations carried out on **5** and the dianionic cluster in **4** are consistent with a similar Ga(I) donor model for the “ $\text{Ga}(\text{OTf})_3$ ” units in both. In particular, the gallium center contributes 72% of the electron density to the Ga–Cu bond in **5** and 85 % to the Ga–Ag bond in **4**.

In conclusion, a number of low-valent gallium species have been isolated from “GaOTf” solutions in aromatic solvents. These suggest that “GaOTf” in toluene is best considered a mixed-valence compound, even in the presence of more π -basic arenes, such as hexamethylbenzene. However, the reaction of “GaOTf” with [18]-crown-6 leads to the isolation of a relatively stable univalent gallium crown ether complex. This simplification in speciation, compared to the crown-free system, may be linked to improved catalytic behavior seen by Schneider *et al.* Additionally, the bimetallic silver(I)/gallium(I) cluster anion $[\text{Ag}_4\{\text{Ga}(\text{OTf})_3\}_4(\mu\text{-Ga})_6(\text{OTf})_4]^{2-}$ has been identified at short reaction times. Thus, while “GaOTf” solutions are very promising reagents for the emerging field of low-valent Ga catalysis, care is required when considering catalytic mechanisms, as the complex speciation in these systems mean that it may not always be obvious what the catalytically active metal and/or oxidation state will be.

Acknowledgements

We are grateful to the University of York and EPSRC (grants EP/H011455 and EP/K031589/1) for funding.

Keywords: Low valent gallium • main group chemistry • coordination chemistry • X-ray crystallography • density functional theory

- [1] a) H. Schnöckel, *Chem. Rev.* **2010**, *110*, 4125-4163; b) C. Dohmeier, D. Loos, H. Schnöckel, *Angew. Chem. Int. Ed.* **1996**, *35*, 129-149; c) G. Linti, H. Schnöckel, *Coord. Chem. Rev.* **2000**, *206-207*, 285-319; d) H. Schnöckel, A. Schnepf, in *Adv. Organomet. Chem.*, Vol. 47, Academic Press, **2001**, 235-281.
- [2] S. Hubert, *Angew. Chem. Int. Ed.* **1985**, *24*, 893-904.
- [3] a) R. J. Baker, C. Jones, *Dalton Trans.* **2005**, 1341-1348; b) B. J. Malbrecht, J. W. Dube, M. J. Willans, P. J. Ragogna, *Inorg. Chem.* **2014**, *53*, 9644-9656; c) J. K. Schuster, J. H. Muessig, R. D. Dewhurst, H. Braunschweig, *Chem. Eur. J.* **2018**, *24*, 9692-9697.
- [4] a) B. Buchin, C. Gemel, T. Cadenbach, R. Schmid, R. A. Fischer, *Angew. Chem. Int. Ed.* **2006**, *45*, 1074-1076; b) B. Buchin, C. Gemel, T. Cadenbach, I. Fernandez, G. Frenking, R. A. Fischer, *Angew. Chem. Int. Ed.* **2006**, *45*, 5207-5210; c) T. Cadenbach, C. Gemel, T. Bollermann, I. Fernandez, G. Frenking, R. A. Fischer, *Chem. Eur. J.* **2008**, *14*, 10789-10796; d) T. Cadenbach, C. Gemel, R. Schmid, M. Halbherr, K. Yusenko, M. Cokoja, R. A. Fischer, *Angew. Chem. Int. Ed.* **2009**, *48*, 3872-3876; e) M. Halbherr, T. Bollermann, C. Gemel, R. A. Fischer, *Angew. Chem. Int. Ed.* **2010**, *49*, 1878-1881; f) S. Gonzalez-Gallardo, T. Bollermann, R. A. Fischer, R. Murugavel, *Chem. Rev.* **2012**, *112*, 3136-3170.
- [5] a) J. M. Slattery, A. Higelin, T. Bayer, I. Krossing, *Angew. Chem. Int. Ed.* **2010**, *49*, 3228-3231; b) A. Higelin, C. Haber, S. Meier, I. Krossing, *Dalton Trans.* **2012**, *41*, 12011-12015; c) A. Higelin, U. Sachs, S. Keller, I. Krossing, *Chem. Eur. J.* **2012**, *18*, 10029-10034; d) A. Higelin, S. Keller, C. Gohringer, C. Jones, I. Krossing, *Angew. Chem. Int. Ed.* **2013**, *52*, 4941-4944; e) M. R. Lichtenthaler, A. Higelin, A. Kraft, S. Hughes, A. Steffani, D. A. Plattner, J. M. Slattery, I. Krossing, *Organometallics* **2013**, *32*, 6725-6735; f) M. R. Lichtenthaler, S. Maurer, R. J. Mangan, F. Stahl, F. Monkemeyer, J. Hamann, I. Krossing, *Chem. Eur. J.* **2015**, *21*, 157-165; g) M. R. Lichtenthaler, F. Stahl, D. Kratzert, L. Heidinger, E. Schleicher, J. Hamann, D. Himmel, S. Weber, I. Krossing, *Nat. Commun.* **2015**, *6*; h) K. Grootz, D. Kratzert, D. Himmel, A. Kastro, Z. Yassine, T. Findeisen, I. Krossing, *Angew. Chem. Int. Ed.* **2018**, *57*, 14203-14206.
- [6] R. J. Wehmschulte, R. Peverati, D. R. Powell, *Inorg. Chem.* **2019**, *58*, 12441-12445.
- [7] a) M. C. Kuchta, J. B. Bonanno, G. Parkin, *J. Am. Chem. Soc.* **1996**, *118*, 10914-10915; b) E. S. Schmidt, A. Jockisch, H. Schmidbaur, *J. Am. Chem. Soc.* **1999**, *121*, 9758-9759; c) N. J. Hardman, B. E. Eichler, P. P. Power, *Chem. Commun.* **2000**, 1991-1992; d) C. Jones, P. C. Junk, J. A. Platts, A. Stasch, *J. Am. Chem. Soc.* **2006**, *128*, 2206-2207; e) K. Yurkerwich, G. Parkin, *J. Cluster Sci.* **2010**, *21*, 225-234; f) M. Asay, C. Jones, M. Driess, *Chem. Rev.* **2011**, *111*, 354-396; g) J. Hicks, P. Vasko, J. M. Goicoechea, S. Aldridge, *Nature* **2018**, *557*, 92-95.
- [8] B. Qin, U. Schneider, *J. Am. Chem. Soc.* **2016**, *138*, 13119-13122.
- [9] Z. Li, G. Thiery, M. R. Lichtenthaler, R. Guillot, I. Krossing, V. Gandon, C. Bour, *Adv. Synth. Catal.* **2018**, *360*, 544-549.
- [10] a) S. Araki, H. Ito, Y. Butsugan, *Appl. Organomet. Chem.* **1988**, *2*, 475-478; b) P.-H. Lee, *Bull. Korean Chem. Soc.* **2007**, *28*, 17-28; c) D. Goswami, A. Chattopadhyay, A. Sharma, S. Chattopadhyay, *J. Org. Chem.* **2012**, *77*, 11064-11070; d) K. Kalantar-Zadeh, J. Tang, T. Daeneke, A. P. O'Mullane, L. A. Stewart, J. Liu, C. Majidi, R. S. Ruoff, P. S. Weiss, M. D. Dickey, *ACS Nano* **2019**, *13*, 7388-7395; e) J. Ma, Y. Lin, Y.-W. Kim, Y. Ko, J. Kim, K. H. Oh, J.-Y. Sun, C. B. Gorman, M. A. Voinov, A. I. Smirnov, J. Genzer, M. D. Dickey, *ACS Macro Letters* **2019**, *8*, 1522-1527.
- [11] H. Schmidbaur, U. Thewalt, T. Zafiroopoulos, *Organometallics* **1983**, *2*, 1550-1554.
- [12] A. Schnepf, C. Doriati, *Chem. Commun.* **1997**, 2111-2112.
- [13] G. Linti, A. Seifert, *Z. Anorg. Allg. Chem.* **2008**, *634*, 1312-1320.
- [14] a) J. L. Bourque, P. D. Boyle, K. M. Baines, *Chem. Eur. J.* **2015**, *21*, 9790-9796; b) L. Yang, J. L. Bourque, J. A. McLeod, P. Shen, K. M. Baines, L. Liu, *Inorg. Chem.* **2017**, *56*, 2985-2991.
- [15] a) C. U. Doriati, M. Friesen, E. Baum, A. Ecker, H. Schnöckel, *Angew. Chem. Int. Ed.* **1997**, *36*, 1969-1971; b) M. S. Hill, P. B. Hitchcock, R. Pongtavornpinyo, *Science* **2006**, *311*, 1904-1907; c) G. Linti, A. Seifert, *Z. Anorg. Allg. Chem.* **2008**, *634*, 1312-1320.
- [16] H. Schmidbaur, R. Nowak, B. Huber, G. Müller, *Polyhedron* **1990**, *9*, 283-287.
- [17] a) M. Schorpp, S. Rein, S. Weber, H. Scherer, I. Krossing, *Chem. Commun.* **2018**, *54*, 10036-10039; b) H. Schmidbaur, U. Thewalt, T. Zafiroopoulos, *Angew. Chem. Int. Ed.* **1984**, *23*, 76-77; c) U. Thewalt, T. Zafiroopoulos, H. Schmidbaur, *Z. Naturforsch B* **1984**, *39*, 1642-1646.
- [18] H. Schmidbaur, R. Nowak, B. Huber, G. Müller, *Z. Naturforsch B* **1988**, *43*, 1447-1452.
- [19] C. Fiolka, **2011**, Universität zu Köln, Cologne.
- [20] a) J. C. Beamish, M. Wilkinson, I. J. Worrall, *Inorg. Chem.* **1978**, *17*, 2026-2027; b) J. C. Beamish, R. W. H. Small, I. J. Worrall, *Inorg. Chem.* **1979**, *18*, 220-223.
- [21] A. V. Protchenko, D. Dange, J. R. Harmer, C. Y. Tang, A. D. Schwarz, M. J. Kelly, N. Phillips, R. Tirfion, K. H. Birj Kumar, C. Jones, N. Kaltsoyannis, P. Mountford, S. Aldridge, *Nature Chem.* **2014**, *6*, 315-319.
- [22] C. G. Andrews, C. L. B. Macdonald, *Angew. Chem. Int. Ed.* **2005**, *44*, 7453-7456.
- [23] J. E. Huheey, E. A. Keiter, R. L. Keiter, *Inorganic Chemistry: Principles of structure and reactivity*, Harper Collins, New York, **1993**.
- [24] a) J. L. Bourque, **2017**, The University of Western Ontario, Electronic Thesis and Dissertation Repository, 5026; b) K. Grootz, D. Kratzert, I. Krossing, *Z. Anorg. Allg. Chem.* **2020**, *646*, 523-525.
- [25] a) S. Mishra, E. Jeanneau, S. Daniele, G. Ledoux, *Dalton Trans.* **2008**, 6296-6304; b) J. Zhao, D. Adcock, W. T. Pennington, J. W. Kolis, *Inorg. Chem.* **1990**, *29*, 4358-4360.
- [26] S. Hagen, H. Schubert, C. Maichle-Mössmer, I. Pantenburg, F. Weigend, L. Wesemann, *Inorg. Chem.* **2007**, *46*, 6775-6784.
- [27] T. Bollermann, A. Puls, C. Gemel, T. Cadenbach, R. A. Fischer, *Dalton Trans.* **2009**, 1372-1377.
- [28] S. P. Green, C. Jones, D. P. Mills, A. Stasch, *Organometallics* **2007**, *26*, 3424-3430.
- [29] T. Bollermann, G. Prabusankar, C. Gemel, R. W. Seidel, M. Winter, R. A. Fischer, *Chem. Eur. J.* **2010**, *16*, 8846-8853.