UNIVERSITY of York

This is a repository copy of Paleotsunami research along the Nankai Trough and Ryukyu Trench subduction zones–Current achievements and future challenges.

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/166432/</u>

Version: Accepted Version

Article:

Fujiwara, Osamu, Goto, Kazuhisa, Ando, Ryosuke et al. (1 more author) (2020) Paleotsunami research along the Nankai Trough and Ryukyu Trench subduction zones– Current achievements and future challenges. Earth-Science Reviews. 103333. ISSN 0012-8252

https://doi.org/10.1016/j.earscirev.2020.103333

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: https://creativecommons.org/licenses/

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/

1	Paleotsunami research along the Nankai Trough and Ryukyu Trench subduction zones									
2	 Current achievements and future challenges – 									
3										
4										
5	Authors:									
6	Osamu Fujiwara ^{1*} , Kazuhisa Goto ² , Ryosuke Ando ² and Ed Garrett ^{3,4}									
7										
8	Affiliations:									
9	1. Geological Survey of Japan, National Institute of Advanced Industrial Science and									
10	Technology, AIST Central 7, 1-1-1 Higashi, Tsukuba, 305-8567, Japan.									
11	2. Department of Earth and Planetary Science, The University of Tokyo, 7-3-1 Hongo,									
12	Bunkyo-ku, Tokyo, 113-0033, Japan.									
13	3. Department of Environment and Geography, University of York, Wentworth Way,									
14	Heslington, York, YO10 5NG, United Kingdom									
15	4. Geological Survey of Belgium, Royal Belgian Institute of Natural Sciences, Brussels,									
16	Belgium									
17										
18	*Corresponding Author: Osamu Fujiwara; Email: o.fujiwara@aist.go.jp.									
19										
20	Keywords:									
21	tsunami deposit, paleoseismology, earthquake forecasting, Japanese islands, Maximum possible									
22	tsunami									
23										
24										

25 1. Introduction

Facing the convergent plate boundary along the northwestern Pacific margin, the Japanese islands have repeatedly suffered from great subduction zone earthquakes and related tsunamis. As it is a mountainous country, the large population and industrial activity are concentrated on small coastal plains, making them particularly vulnerable to these natural hazards. The most recent example is the 2011 Tōhoku earthquake and tsunami, which occurred along the northern Japan Trench, where the Pacific Plate subducts beneath the Japanese archipelago.

One area highly likely to generate a future magnitude (M) 8.5+ subduction zone earthquake 32 and tsunami is the Nankai Trough-Ryukyu Trench subduction zone, where the Philippine Sea 33 Plate descends beneath the Japanese archipelago (e.g., HERP, 2013). Forecasting of the next 34 35 earthquake and tsunami in these regions is an urgent issue for the Japanese government's national disaster prevention measures. To address this issue, many research projects have sought 36 to identify the locations, timing, and size of past great earthquakes, based on historical records 37 38 along with geological evidence. Documented Nankai Trough earthquakes and related tsunamis date back to the 7th century CE and suggest that M8-class great earthquakes have occurred at an 39 40 interval of 100-200 years in this region (HERP, 2013), with 75 years having passed since the last events, which occurred in 1944 and 1946 CE. 41

On the other hand, documented records of great earthquakes and tsunamis along the Ryukyu Trench are limited mainly to the 17th century CE and later. It was once thought that no great earthquakes occur along the Ryukyu Trench; however, results from seismology and tsunami deposit studies since 2011 CE have overturned this theory and have highlighted the importance of estimating the magnitude and location of possible future great earthquakes in this region (Ando et al., 2009, 2012a, 2018).

Both the historical and geological records are inevitably biased by age and region as 48 archives for paleoseismic events. Furthermore, geological records tend to be biased towards 49 larger events. However, the geological record has the greatest advantage of all: the length of 50 51 time it covers. Geological and archeological studies have been conducted to supplement the imperfect documented earthquake and tsunami history and to extend the record back to earlier 52 periods. The first line of these studies is "Earthquake archaeology", established in the late 1980s, 53 which seeks to identify the past earthquakes based on evidence of strong ground shaking, such 54 55 as liquefaction features, found in archeological sites (Sangawa, 1992). These studies suggested the possibility of subduction zone earthquakes, such as the 887 and 684 CE Tokai earthquakes 56 along the eastern Nankai trough, which had not been confirmed in the documentary record. 57 Several earlier Nankai Trough earthquakes dating back to 1800-2000 years ago have been also 58 59 proposed from the archaeological data (e.g., Sangawa, 2007). However, this method generally

does not provide information on tsunamis, which are a key characteristic of major subductionzone earthquakes.

A second line of research, paleotsunami studies, commenced along the Ryukyu Trench in 62 the 1960s, and full-scale research along the Nankai Trough began in the 1990s. Tsunamis 63 following two M9-class earthquakes that occurred in close succession in the 21st century, the 64 65 2004 CE Indian Ocean tsunami and 2011 CE Tohoku tsunami, have provided a clear incentive to study the record of past tsunamis and incorporate paleotsunami data into forecasts of future 66 events. Results of paleotsunami studies along the Nankai and Sagami Troughs up to 2007 CE 67 are summarized in Komatsubara and Fujiwara (2007). Garrett et al. (2016) provides a 68 comprehensive review of the recurrence history of great Nankai Trough earthquakes in the 69 70 Middle to Late Holocene, including not only tsunami deposits and liquefaction features on land, 71 but also marine and lacustrine turbidites.

The Geological Society of Japan published a special issue summarizing the results of 72 73 paleotsunami research along the Japanese archipelago as one of the commemorative publications celebrating its 125th anniversary (Journal of the Geological Society of Japan 123, 74 75 No. 10). Wallis et al. (2018) describes the changes in the Japanese government's policy 76 concerning tsunami countermeasures following the Tohoku event, and the changes in the paleotsunami community's research direction reflecting the new policy. The policy (CDMC, 77 78 2011) reflected on the overreliance on geodetic observation data (\sim 130 year-long) and "recent and reliable" historical documents (covering the last ~400 years) in the long-term earthquake 79 prediction for Japan, and stipulated that all information from various different sources including 80 81 geological data should be considered in the new earthquake and tsunami countermeasures (see 82 summary in Goto et al., in this issue). As a response to this situation, in Japan, many paleotsunami studies have been conducted and some are in progress under the auspices of the 83 national and local governments. 84

Tsunami deposit studies along the Nankai and Ryukyu subduction zones have been 85 independently reviewed (e.g., Fujiwara and Tanigawa, 2017; Goto, 2017), but there is no 86 comprehensive review comparing the two or summarizing the latest research. In this review, we 87 intend to address this point. For example, differences in the climate and geology of the two 88 regions create differences in the occurrence and composition of typical tsunami deposits: sandy 89 90 tsunami deposits for the Nankai and tsunami boulders (mainly coral blocks) for the Ryukyu subduction zone. Additionally, as the two regions comprise a continuous subduction zone, the 91 possibility of a giant earthquake spanning both regions must carefully be assessed (Furumoto 92 and Ando, 2009). For this reason, synthesizing the earthquake and tsunami histories of both 93 94 subduction zones is particularly important.

Earthquake and tsunami forecasts operate at various timescales ranging from earthquake 95 96 early warnings seconds before shaking is felt to long-term probability predictions over decades where geological data is predominant (e.g., Satake and Atwater, 2007). Every case requires the 97 specification of three factors: where, when, and how big. From this perspective, this paper 98 reviews the results of research on onshore tsunami deposits, their limitations, and existing 99 100 knowledge gaps, and discusses measures to address the remaining issues. When and where 101 earthquakes occurred is reconstructed based mainly on tsunami deposits studies with the help of 102 other geological evidence. A comprehensive earthquake recurrence model is still incomplete for these subduction zones. This paper therefore also addresses the role that tsunami deposit 103 research plays in the development of this recurrence model. The magnitude of earthquakes that 104 105 occur along a subduction zone varies widely (e.g., Satake and Atwater, 2007); this paper 106 discusses the variation in the size of earthquakes inferred from tsunami deposit research.

The possibility of the occurrence of an unforeseen or unexpected large earthquake and 107 tsunami is a major point of reflection following the 2011 Tohoku event. Learning from this, an 108 109 estimation of the "maximum possible earthquake and tsunami" that could affect the Japanese 110 coast has been included in disaster prevention measures in Japan (CDMC, 2011; Cabinet Office, 2012a). The CDMC (2011) stated that future tsunami countermeasures will basically require the 111 assumption of two possible levels of tsunamis. Level one reflects the largest tsunamis known to 112 113 occur on centennial timescales and forms the basis for designing the height of seawalls and other features. Level two is the maximum possible tsunami, which occurs at a much lower 114 115 frequency than the first level but is considered to be more destructive. The hypothetical size of a 116 level two tsunami is based on studies of tsunami deposits and crustal movements. The 2011 117 Tōhoku tsunami is an example of a level two tsunami. Countermeasures against a level two tsunami could include relocating settlements to higher ground and building huge seawalls, but 118 specific measures are difficult to implement. Complicating attempts to mitigate future impacts, 119 it is unknown whether such extremely large earthquakes and tsunamis have occurred in the past 120 121 along the Nankai and Ryukyu subduction zones. This paper refers to the plausibility of this 122 scenario from the standpoint of the geological evidence.

123

124 **2. Tectonic setting**

125 **2.1. Plate subduction and related geomorphology**

The Nankai and Ryukyu subduction zones mark the northwestern margin of the Philippine Sea Plate (Fig. 1), but the nature of plate subduction, geological structures, and occurrence mode of earthquakes are largely different between the two. The Palau-Kyushu Ridge forms the boundary between the two subduction zones. The Nankai Trough extends for 700 km from the southwest of Shikoku Island in the west to Suruga Bay in the east, where it is also called the Suruga Trough. The Ryukyu Islands consist of numerous small islands extending over 1000 km along the Ryukyu Trench, which belong to the subtropical climate zone and hence the depositional environment is very different from other parts of Japan. The convergence rate is generally 40-50 mm a⁻¹ along the Nankai subduction zone and 50-63 mm a⁻¹ along the Ryukyu subduction zone, respectively. The slope of the descending slab, with some local variations, is generally steeper in the Ryukyu subduction zone.

137 The Amami Plateau, Daito Ridge, Oki-Daito Ridge and basins between them extend in E-W or NW-SE directions on the Philippine Sea Plate. Western extensions of these topographic 138 irregularities subduct beneath the Ryukyu arc (e.g., Kato, 1993). The Ryukyu and Nankai 139 subduction zones are represented by old (40-49 Ma), cold slab that originated from the West 140 141 Philippine Basin and young (<20 Ma), hot slab that originated from the Shikoku Basin, 142 respectively (e.g., Ishizuka et al., 2011). A thick accretionary prism characterizes the Nankai Trough, in which high-angle reverse faults (splay faults) branching off the plate boundary 143 megathrust may move together with the main thrust during a great earthquake and enhance the 144 tsunami (Baba et al., 2006; Moore et al., 2007). 145

146 The Nankai Trough fault region has been empirically divided into five fault planes (rupture 147 zones), which denoted as segments A to E from west to east, respectively (Fig. 1) (Ando, 1975a, b; Ishibashi, 1976, 1981). This classification is mainly based on studies of the source 148 mechanisms of historical earthquakes and the forearc submarine topography, which reflects the 149 150 seismotectonic activities along this region. Each of the five segments has a length of 100-150 151 km along the trough axis and almost corresponds to the distribution of forearc basins (e.g., Awata and Sugiyama, 1989; Sugiyama, 1990). In general, plate boundary earthquakes 152 153 (M8-class) that are considered to have source areas in areas A to B and C to E are called Nankai 154 and Tokai earthquakes, respectively.

Allocating the source areas of past earthquakes to any of segments A to E is based on the implicit assumption that a large earthquake recurs on its own fault plane. The necessity to criticize and review this assignment has recently been pointed out (e.g., Seno, 2012). The Hyūga-nada area (Figs. 1, 2), where M7-class earthquakes occur separately from the Nankai and Tōkai earthquakes, may be another seismic zone denoted as Z (e.g., Wells et al., 2003). This empirical segmentation of rupture zones has not been applied to the Ryukyu Trench as historical and seismological information is scarce.

Flight of marine terraces, mainly from mid-Pleistocene to Holocene in age, are distributed on the peninsulas, capes and islands of the Nankai and Ryukyu subduction zones (Koike and Machida, 2001), and there have been many discussions about their formation process in relation to sea-level changes and crustal movements (e.g. Yoshikawa et al., 1964; Webster et al., 1998). They show a wide regional variation in average uplift rates. For example, the terraces of marine 167 oxygen isotope stage (MIS) 5e are 40-50 m high in many areas, but reach nearly 190 m at Cape
168 Muroto, the boundary between segments A and B. Coseismic uplift and subsidence during
169 historical earthquakes characterize the tips of the peninsulas and their inland plains, respectively
170 (e.g., Shikoku; Geographical Survey Institute, 1952; Tokai; Ishibashi, 1981). In these area,
171 reverse sense crustal deformation is observed in the interseismic period; gradual subsidence of
172 the peninsulas and uplift of the lowland, respectively (Geographical Survey Institute, 2008).

173 Kikai Island, located at the northern margin of the Amami Islands on the subducting 174 Amami Plateau, is the closest island to the Ryukyu Trench axis. It shows an extremely high uplift rate, with the height of the MIS 5e marine terrace exceeding 200 m. On this island, 175 176 Holocene marine terraces consisting of coral reefs record several-meter-scale uplift events that 177 recurred with 1500-year intervals (Webster et al., 1998). The other islands along the Ryukyu 178 Trench do not show remarkable crustal deformation, but the coastal notch in Ishigaki Island records a cumulative uplift of ~ 2 m in the last 2000 years (Kawana, 1989). Some of these 179 marine terraces distributed along the Nankai and Ryukyu coasts have provided the theoretical 180 basis for time-predictable recurrence models for subduction zone earthquakes (e.g., Shimazaki 181 182 and Nakata, 1980).

183

184 2.2. Seismicity

Understanding the state of inter-plate coupling and seismogenesis in the subduction zones provides basic information for estimating the region's potential of great earthquake and tsunami. Especially, in the Hyūga-nada region, connecting the Nankai and Ryukyu subduction zones, this information is the key to identifying the geological and seismological differences between the two.

190

191 Geodetic deformation and inferences of inter-plate coupling along the Nankai Trough and192 Ryukyu Trench subduction zones

193 The source areas of large earthquakes are usually expected to present strong inter-plate 194 coupling during inter-seismic periods (e.g., Lay and Kanamori, 1981). The degree of the coupling can vary over fault areas, reflecting differences in the fault's frictional properties. If a 195 fault creeps steadily, it has a low degree of coupling, while high coupling results from a 196 197 seismogenic fault that is stuck during inter-seismic periods. The degree of the coupling is 198 quantified in geodesy (Savage, 1983) by the slip-deficit rate (SDR) (or back-slip rate), which is 199 the rate of upper plate displacement along with the plate interface driven by the subduction 200 motion of the lower plate. The coupling coefficient is also determined as the SDR divided by 201 the long-term average rate of the relative motion between the two plates. When the fault is 202 completely coupled, the SDR is equal to the long-term average relative plate motion rate and the 203 coupling coefficient is unity. SDRs are estimated from geodetic data by applying slip inversion204 analysis (Savage, 1983).

Several geodetic inferences of inter-plate coupling have been recorded along the Nankai 205 206 Trough subduction zone (Miyazaki and Heki, 2001; Ito and Hashimoto, 2004; Nishimura and Hashimoto, 2006; Wallace et al., 2009; Loveless and Meade, 2010; Yokota et al., 2016), while 207 208 only a few studies have been conducted along the Ryukyu Trench (Ando et al., 2009; Tadokoro 209 et al., 2018) due to limited spatial coverage of onshore GNSS observations in this island area. Along the Nankai Trough, spatial distributions of SDR have been inferred and certain 210 heterogeneous structures have been identified (Fig. 1), owing to a dense onshore GNSS network 211 called GEONET and the recent development of seafloor geodetic observation networks by the 212 213 Japan Coast Guard (Yokota et al., 2016). The seafloor observation sites deployed near the 214 trough axis contribute greatly to resolving the previously unknown coupling at shallow depths of the plate interface (Sagiya and Thatcher, 1999), which caused an ambiguity in the estimation 215 of the maximum earthquake size in the area. Lack of knowledge of shallow coupling was clearly 216 217 illustrated along the Japan Trench subduction zone, which hosted the 2011 Tohoku-oki 218 earthquake (Nishimura et al., 2000; Loveless and Meade, 2011).

219 All of the SDR estimates indicate that the Nankai Trough subduction interface is strongly coupled overall, with Yokota et al. (2016) further resolving regional heterogeneity, with three 220 221 areas of relatively stronger coupling with higher SDRs inferred offshore Shikoku Island, the Kii Peninsula and the region from the Atsumi Peninsula to the Omaezaki Peninsula (Fig. 1). The 222 first two areas seen to coincide with the source areas of the 1946 CE Nankai (M8.0) and the 223 224 1944 CE Tonankai (M7.9) earthquakes, and the third area seems to be included in the source area of the 1854 CE Ansei-Tōkai earthquake (Ando, 1975b). Given such correlations, the 225 226 observed heterogeneous distribution of SDR might have played a role in defining the segmentation of historical earthquakes. However, some earthquakes can involve multiple 227 segments, as in the 1707 Hoei (M8.6) earthquake, which is considered to have ruptured all three 228 229 of the high SDR segments (Ando, 1975b), and the earthquake recurrence pattern is not always 230 characteristic (Seno, 2012).

Near the southwestern end of the Nankai Trough subduction zone, the degree of coupling offshore of southern Kyushu Island becomes rather ambiguous due to poorer observational coverage; however, geodetic inferences still tend to suggest weaker coupling in the southern half of the Hyūga-nada region (Nishimura and Hashimoto, 2006; Wallace et al., 2009; Loveless and Meade, 2010) (see Fig.1), reflecting the rotation of the displacement vectors observed over southern Kyushu. The location of this weak coupling is suggested to be correlated with the subducting Kyushu-Palau ridge (Wallace et al., 2009; Yamamoto et al., 2013). To the south, inferences of inter-plate coupling become more difficult due to the much poorer observationalcoverage.

Along the Ryukyu Trench subduction zone, inter-plate coupling has been evaluated 240 241 qualitatively due to the sparseness of onshore observational stations, with generally weak 242 coupling inferred (Nishimura et al., 2004). However, more quantitative geodetic inferences have 243 been conducted with the addition of ocean bottom geodetic observations. Seafloor displacement 244 data has recently been retrieved from the area offshore the Okinawa Islands, in the center of the Ryukyu Trench, and almost complete coupling of the plate interface is inferred by Tadokoro et 245 al. (2018), in contrast to the onshore-based estimation. It is also suggested that this highly 246 coupled area corresponds to the source area of the 1791 CE M 8 inter-plate earthquake. 247

248

249 Seismological inferences of inter-plate coupling using repeating earthquakes

Repeating earthquakes are moderate-sized earthquakes that are considered to occur 250 repeatedly at the same points on faults due to the similarity in their seismic waveforms. They 251 252 are used to infer the degree of inter-plate coupling in a similar fashion to creep meters (Nadeau 253 and Johnson, 1998). While the absence of repeating earthquakes characterizes both the end members of the fully coupled and the fully decoupled fault, once repeating earthquakes are 254 255 observed, inter-plate coupling can be estimated, with higher production rates reflecting weaker 256 coupling and more frequent occurrence of fault slip. The use of repeating earthquakes can supplement geodetic inferences due to its higher detection capability, reflecting the weaker 257 geometrical spreading nature of the seismic waves than the geodetically-observed static 258 259 displacement. Nevertheless, both observations are inevitably limited in island areas.

260 Igarashi (2010) inferred inter-plate coupling by using repeating earthquakes and mapped its 261 spatial variation on the plate interfaces off the Japanese islands. The background seismic activity of the Nankai Trough is generally low reflecting the strong coupling and repeating 262 earthquakes are rarely found except the shallowest portion of the plate interface near the trough 263 264 axis. However, it is evident that inter-plate coupling becomes weaker from the Hyūga-nada 265 region towards the Ryukyu Trench to the south, matching to the inference from geodetic observations. Yamashita et al. (2012) increased the detection of repeating earthquakes, focusing 266 267 on the Hyūga-nada region, and mapped a more detailed spatial distribution of inter-plate 268 coupling (Fig. 1). In this region, the coupling appears to be weakest along the band where the 269 Kyushu-Palau ridge is being subducted (Yamamoto et al., 2013), with the repeating earthquakes 270 accommodating most of the long-term relative plate motion at shallower (<~20 km) and deeper 271 (>40 km) depths. However, to the south, the coupling increases in the area off Tanegashima 272 Island, corresponding to the northern end of the Ryukyu Trench, where repeating earthquakes 273 accommodate approximately half the relative plate motion.

In the Hyūga-nada region, the history of moderate- and large-sized earthquakes appears to be matched with the inferred coupling distribution (Yamashita et al., 2012). In the northern area of the Hyūga-nada, M 6 and 7-class earthquakes have repeatedly occurred, while in the southern part, the 1996 CE M 6.5 earthquake occurred. Between these regions, major earthquakes are not known, while moderate-sized earthquakes (5<M<6.5) have been observed in the relatively strongly coupled area.

Inter-plate coupling along the Ryukyu Trench seems to show a systematic variation where the northern region – from Tanegashima Island to Amami Island – exhibits relatively strong coupling and the southern region around the Sakishima Islands exhibits weaker coupling. However, the coupling state may not be definitively identified solely based on repeating earthquakes in these island areas; in the central region, while repeating earthquakes suggest moderate coupling, seafloor observations suggest almost full coupling (Tadokoro et al., 2018).

286

Spatial distribution of shallow and deep slow earthquakes; implications for the state of inter-plate coupling

Weak inter-plate coupling does not necessarily mean that faults creep steadily. Weak coupling can reflect intermittent slip events on the plate interfaces. Such slip events are generally called slow earthquakes and include slow slip events (SSEs), low-frequency earthquakes and (non-volcanic) tremor. Slow earthquakes are thought to occur along the transition zones of the seismogenic and aseismic zones of plate interfaces (Ando et al., 2012b; Obara and Kato, 2016) both at the up-dip (~5 km) and down-dip ends (~30 km) of the seismogenic zones that generate the huge earthquakes.

296 The family of slow earthquakes has been discovered over the past twenty years along the 297 Nankai Trough by seismological (e.g., Obara, 2002) and geodetic (e.g., Hirose and Obara, 2005) observations, and their correlation with inferred inter-plate coupling states are confirmed in 298 great detail. Along the plate interface at depths from ~30 to ~40 km, tremor and short-term 299 300 SSEs (S-SSEs) are generally observed with recurrence periods of ~ 6 months (Obara, 2002; 301 Hirose and Obara, 2005), while long-term SSEs with M 7 have recurred at ~10-year intervals particularly beneath Lake Hamana in Tōkai region (Suito and Ozawa, 2009 and references 302 therein). For the Ryukyu Trench, seismological observations suggest that slow earthquakes, 303 called very low frequency earthquakes, probably occur on the plate interface over the entire 304 305 Ryukyu Trench (Ando et al., 2012a), and geodetic observations identify seemingly 306 corresponding short-term SSEs (S-SSEs) and long-term SSEs (L-SSEs) (Heki and Kataoka, 307 2008; Nishimura, 2014). Nishimura (2014) suggested that the occurrence of S-SSEs at shallow 308 depths may be a signature of incomplete inter-plate coupling. Tadokoro et al. (2018) suggested the inferred strong coupling area does not overlap with the SSE areas, and heterogeneity in theareas of SSE occurrence should exist along the Ryukyu Trench.

311 The Hyūga-nada region appears to present a transitional character in terms of the spatial 312 distributions of slow earthquakes. In the other areas of the Nankai Trough, shallow slow earthquakes and deep S-SSEs and tremors are both observed, while deep S-SSE/tremor is not 313 314 observed in the Hyūga-nada region (Obara and Kato, 2016). The occurrence of shallow slow 315 earthquakes has been characterized in great detail owing to the recent deployment of ocean bottom seismometers (Yamashita et al., 2015). Tremor was observed to migrate in the shallow 316 depths of the plate interface, and the lateral extent of the migration was delimited by the 317 possible area of the subducting Kyushu-Palau ridge (Yamamoto et al., 2013). The down-dip 318 319 limit of the migration was not reached until ~ 20 km, corresponding to the hypocentral depths of 320 M 7-class earthquakes in this region, suggested that it stopped at the shallower end of the seismogenic zone (Yamashita et al., 2015). This implies that the intermediate depths of the plate 321 interface are still coupled and have the potential to nucleate a major earthquake. The 322 323 geophysical observations clearly show that the inter-plate coupling of this area is relatively low 324 overall, but these observations still leave the possibility that the rupture of this coupled zone 325 could be triggered by neighboring large earthquakes, or the rupture of this area could facilitate interactions between neighboring large earthquakes, if their timings are synchronized. 326

While the above-mentioned geophysical observations have clarified the spatial characteristics of the inter-plate properties, we should still remember that our experience is limited by the short history of geophysical observations. The states of inter-plate coupling can slowly fluctuate even over the course of several years (Uchida et al., 2016), and limited snapshots can fail to reveal critical properties. Integration of paleoseismological evidence is still critically important to capture phenomena spanning long periods of time.

333

334 3. Record of tsunamis and tsunami deposits

335 3.1. Nankai Trough region

336 3.1.1. Historical Tōkai and Nankai earthquakes and tsunamis

The Nankai Trough region, which is close to the political and cultural center of Japan, the 337 Kyoto-Nara-Osaka area, has the longest and most complete documented record for the 338 339 recurrence of subduction zone earthquakes and tsunamis in the world. According to this 340 1300-year long record, earthquakes have occurred in a variety of modes. Some earthquakes 341 have ruptured almost the entire subduction zone at the same time, while at other times, two 342 earthquakes, referred to as Tokai and Nankai earthquakes, have occurred consecutively, separated by short intervals (Fig. 2). Nankai earthquakes that are historically confirmed include 343 344 the 684 CE Hakuho (M8.0-8.5), 887 CE Nin-na (M8.0-8.5), 1361 CE Kōan (M8.2-8.5), 1707

CE Hoei, 1854 CE Ansei-Nankai (M8.4; only 32 hours after the 1854 Ansei-Tokai earthquake), 345 and 1946 CE Showa-Nankai (M8.0) earthquakes, with further unconfirmed events in 1099 and 346 347 1614 CE. Tōkai earthquakes confirmed by documented records include the 1096 CE Eichō (M8.0-8.5), 1498 CE Meiō (M8.2-8.4), 1707 CE Hōei (M8.6), 1854 Ansei-Tōkai (M8.4), and 348 1944 CE Tonankai (M7.9) earthquakes, and a further rupture may have occurred in 1614 CE. 349 350 The 1707 earthquake was a simultaneous rupture of the Tokai and Nankai fault segments 351 (segments A to D, and perhaps also including the southern part of E) and was the largest earthquake in Japanese history until the occurrence of the 2011 Tohoku earthquake. 352

There are two possibilities for a great earthquake that occurred along the Nankai Trough in 353 the early 17th century. The first is the 1605 CE Keichō earthquake, which generated a large 354 355 tsunami from Kanto to Kyushu. This event has been considered as a "tsunami earthquake", 356 because only weak shaking accompanied the large tsunami (e.g., Ishibashi and Satake, 1998). Ando and Nakamura (2013) suggested that the rupture occurred along a shallow portion of the 357 plate interface. An alternative hypothesis is that the 1605 earthquake occurred along the 358 359 Izu-Ogasawara Trench, to the east of Izu-Bonin Arc (Harada et al., 2013). Instead of 1605 CE 360 event, Ishibashi (2014) proposed the 1614 CE earthquake as a candidate for a 17th century 361 Nankai Trough earthquake, based on records of strong ground shaking in Kyoto and tsunamis at the Kii Peninsula. A Nankai earthquake paired with the 1498 CE Meiō Tōkai earthquake has not 362 363 been confirmed; nevertheless, liquefaction features in western Japan suggest the occurrence of a Nankai earthquake around 1498 CE (Sangawa, 2001, 2007). Ishibashi (2014) proposed four 364 earthquakes that occurred between 1498 and 1512 CE as candidates, but there is no solid record 365 366 of tsunami inundation during this period.

Documentary records confirming the occurrence of a Tōkai earthquake paired with the 1361 CE Nankai earthquake have not yet been discovered. As a candidate for this earthquake, Ishibashi (1998, 2014) proposed an earthquake that featured strong shaking in Kyoto, Nara, and Kumano (north of Shiono-misaki; boundary between the segments B and C) on July 24th 1361 (2 days before the Nankai earthquake). Late 14th century coseismic uplift around Cape Omaezaki (Fig. 2, site 27) shown by emerged boring bivalves (Kitamura et al., 2018c) and marine terrace (Fujiwara et al., 2010) support Ishibashi's hypothesis.

The occurrence of the 1099 CE earthquake was originally inferred from documents suggesting strong ground shaking in the Kyoto, Nara, and Osaka region (but less than 1096 CE earthquake) and coastal subsidence in Kochi (around site 10 in Fig. 2); which matches general characteristics of other Nankai earthquakes (e.g., Ishibashi, 1999). However, the occurrence of this earthquake is uncertain due to the unreliability of underlying evidence, especially the lack of reports of strong ground shaking and tsunami inundation in the area facing segments A and B. Ishibashi (2016) suggested that the 1099 CE earthquake did not actually exist and that the 1096 381 CE earthquake was instead a full-length rupture of the Tōkai and Nankai segments. Ishibashi
382 (2014) also suggested the possibility that the 1099 CE Nankai earthquake did occur but was
383 smaller in magnitude than other Nankai earthquakes.

The possibility that a Tokai earthquake occurred at the same time as the Nankai earthquake 384 on 22nd August, 887 CE has been suggested from a description in the Nihon Sandai Jitsuroku 385 (日本三代実録, "The true history of three reigns of Japan"; completed in 901 CE) (e.g., 386 387 Ishibashi, 1999, 2014). The chronicle's record for that day shows that strong ground shaking was felt in wide area including the Tokai region along with the occurrence of an earthquake and 388 tsunami in the Nankai region. The occurrence of the 684 CE Nankai earthquake is documented 389 in the Nihon Shoki (日本書紀, The Chronicles of Japan, completed in 720 CE), with a report of 390 391 tsunami and coastal submergence in Kochi, but there are no records of a corresponding 392 earthquake in the Tōkai region. Only liquefaction features in archaeological sites in the Tōkai region have suggested as possible evidence of a Tokai earthquake corresponding to the 684 CE 393 Nankai earthquake (e.g., Sangawa, 2001, 2007). However, given the lack of evidence for 394 tsunami to determine the occurrence of the Tokai earthquake in either case, the possibility of an 395 396 inland earthquake from an active fault cannot be ruled out. These imperfections in the historical 397 earthquake record have been a hindrance in discussing the occurrence time of Tokai and Nankai 398 earthquakes and their linkage.

399 The 1498 CE Meiō tsunami might have been somewhat higher than 1707 and 1854 CE tsunamis, ~ 8 m along the Enshu-nada coast, which includes the sites 19 to 27 in Fig. 2 (Hatori, 400 1975). The height of the 1605 CE tsunami was generally 4 - 6 m and up to 13 m in Shikoku 401 402 (Murakami et al., 1996). The 1707 CE tsunami generally reached 5 - 8 m-high along the Tokai 403 and Nankai coasts, over 10 m in some locations on the Kii Peninsula and Shikoku, and 2 - 4 m 404 along the Kyushu coast (Murakami et al., 1996, Watanabe, 1998). The 1854 CE Ansei-Tōkai tsunami had a height of 5 - 8 m, with some isolated cases of ~20 m at the tips of promontories 405 (Watanabe, 1998). The 1854 CE Ansei-Nankai tsunami generally reached 5-6 m-high, 406 407 occasionally over 8 m along the Kii Peninsula and Shikoku (Murakami et al., 1996, Watanabe, 408 1998), and 2 - 4 m along the Kyushu coast (Cabinet Office, 2012b). The 1944 CE tsunami was 3 - 7 m-high along the eastern coast of Kii Peninsula and 0.9 - 2 m-high along the Enshu-nada 409 coast (Watanabe, 1998). The height of the 1946 CE Showa-Nankai tsunami was 2 - 4 m in 410 411 general, up to 6 m on the Kii and Sikoku coasts and 1 - 1.6 m on the Kyushu coast (Murakami et 412 al., 1996, Watanabe, 1998). These tsunamis generally caused greater human and economic 413 damages than resulted from ground shaking during historical Nankai Trough earthquakes (e.g., 414 Yata, 2009, 2018; Ishibashi, 2014).

415

416 3.1.2. Tsunami deposits

Nankai Trough coasts have few extensive wetlands suitable for reconstructing the tsunami 417 418 inundation area, which characterize the major paleotsunami study fields in the world, such as Hokkaido (e.g., Nanayama et al., 2003), Cascadia (e.g., Peters et al., 2007) and Chile (e.g. 419 420 Cisternas et al., 2005). Artificial disturbance of surface sediments reduces the chances of finding tsunami deposits from this region, along with the difficulties of distinguishing tsunami 421 deposits from the other washover (mainly storm) deposits and river-flooding deposits. 422 423 Consequently, coastal back-barrier lakes and ponds, and infilled valleys have mainly been targeted as paleotsunami study sites. The barriers between these sites and the sea are mainly 424 425 composed of sand dunes and beach ridges formed during the last seven to six thousand years, after the mid Holocene sea level high stand in Japan (e.g., Sato et al., 2016a). Many of these 426 427 sites have gradually subsided through the Holocene, and stably accumulated fine-grained 428 sediments, sometimes rich in organic matter (e.g., Okamura and Matsuoka, 2012). They have a high potential for preserving marine overwash event deposits. Coring surveys using hand corers, 429 piston corers, and geoslicers (Nakata and Shimazaki, 1997) have been the main methods to 430 431 reveal the tsunami deposits, while their depositional ages have mainly been determined by 432 radiocarbon dating. Due to the very limited distribution of widespread Holocene tephras along 433 the Nankai and Ryukyu subduction zones, it is difficult to estimate the age of tsunami deposits 434 based on volcanic ash stratigraphy and to compare them between regions, as has been done in 435 the Hokkaido (e.g., Nanayama et al., 2003) and Tōhoku (e.g., Sawai et al., 2012) regions.

Tsunami and possible tsunami deposits have been reported from a total of 23 study sites 436 along the Nankai Trough coast (Fig. 2, sites 7 to 20, 22 to 26, 28, 29, 31 and 32). Adjacent 437 438 points that cannot be distinguished on the map, such as some points in the southern part of the 439 Kii Peninsula, are counted as one. Garett et al. (2016) recalibrated all available published 440 radiocarbon ages to take advantage of the latest radiocarbon calibration curves, updated estimates of local marine radiocarbon reservoir effects, and Bayesian age modeling approaches. 441 As a result, some of the originally reported correlations between sedimentary layers and 442 443 historical earthquakes have been revised. We follow Garrett et al. (2016) for the ages of tsunami 444 deposits summarized here.

Most of the reported tsunami and possible deposits are washover sand deposits intercalated 445 in muddy or peaty deposits, with two exceptions of the tsunami boulders found from the 446 447 southern Kii peninsula (Fig. 2, site 14) and the southern Izu peninsula (Fig. 2, site 32). Sandy 448 tsunami deposits generally show a tapering shape with a fining landward trend and sometimes 449 include remains of marine organism. They may be a single layer, but often consist of multiple layers, each of which shows a graded structure. Sedimentary structures indicating deposition 450 451 from a tractive flow, such as basal erosion surface, cross-lamination, and inverse grading also 452 characterize many cases.

454 Historical tsunami deposits

Examples of tsunami deposits that can be accurately correlated to historical earthquakes are still limited in the Nankai Trough region (Fig. 2). Limited numbers of measurements and chronological uncertainties inherent in the radiocarbon approach make it difficult to distinguish individual historical tsunami deposits that formed at short intervals, often of less than 100 years, and compare them between distant sites.

The easternmost examples are reported by Kitamura and Kobayashi (2014a) and Kitamura et al. (2014) from Shimoda, southern Izu Peninsula (Fig. 2, Site 32). The former is a laminated sand bed about 10 cm thick, with rip-up clasts that is estimated to have been deposited between 463 430-290 cal. BP and 1950 CE, possibly due to the 1854 CE Ansei-Tōkai or 1707 CE Hōei tsunami. The latter is a 32-tonne boulder located on a coastal plateau, which is attributed to the 1854 CE Ansei-Tōkai earthquake based on the radiocarbon ages of sessile marine fossils attached to the boulder.

On the Ita Lowlands (Fig. 2, site 31), one possible sandy tsunami deposit – with diatoms 467 suggesting a marine incursion – was found from a sedimentary sequence covering the last 2500 468 years (Sawai et al., 2016). Radiocarbon dating limited the age of the tsunami bed to 1200-1320 469 CE or 1150-1330 CE and Sawai et al. (2016) attributed it to the 1096, 1099, or 1361 CE Nankai 470 471 Trough earthquake or the 1293 CE Kanto earthquake, centered on the Sagami Trough, east of Izu Peninsula. On the Shimizu Plain, Kitamura and Kobayashi (2014b) reported the deposition 472 473 of gravelly sand beds associated with the erosion of the muddy bay bottom between 670-600 474 and 250–50 cal. BP (Fig. 2, Site 29). They inferred that the sand beds may be attributed to the 475 tsunamis from 1498 CE Meiō or 1707 CE Hōei earthquakes.

476 Komatsubara et al. (2008) conducted paleotsunami research in a back-marsh at Shirasuka (Fig. 2, site 20) using a 4 m-long geoslicer and found exotic sand beds within the muddy 477 478 sedimentary sequence. Comparing the radiocarbon ages obtained from the cores and historical 479 documents, five sand beds, each of which is 10 - 20 cm thick, were attributed to tsunamis 480 associated with the 1498, 1605, 1707, 1854 CE earthquakes and the 1680 or 1699 CE typhoon. This hypothesis was verified with new age models developed using radiocarbon ages (Garrett et 481 al., 2018) and optically stimulated luminescence (OSL) ages (Riedesel et al., 2018) from 482 483 subsequently excavated sediment cores from the same site as the original study. The sand unit A of Komatsubara et al., (2008), located beneath the 1498 CE tsunami deposit and originally 484 485 interpreted as a slope failure deposit, was recognized as a possible tsunami deposit based on the occurrence of marine and brackish diatoms (Garrett et al. 2018). Both radiocarbon and OSL age 486 487 models suggest an age for this tsunami deposit of the late 14th century and the layer may be 488 attributed to the 1361 CE Tōkai earthquake (Garrett et al., 2018; Riedesel et al., 2018).

Fujino et al. (2018) conducted an array coring survey on the Shima Lowland (Fig. 2, site 17), a Holocene drowned valley, and found 10 exotic sand beds within the muddy marsh sequence. A tsunami origin for these beds is inferred from sedimentary features common in modern tsunami deposits, including marine bioclast-rich sand, alternating sub-layers of sand and silt, rip-up clasts, and upward-fining structures. The youngest three tsunami beds were tentatively attributed to 1498, 1096, and 684 CE Nankai Trough earthquakes.

495 A group of possible tsunami boulders, consisting of hundreds of angular boulders weighing up to 100 tons (Namegaya et al., 2011), are found on a wave-cut bench near Cape 496 Shiono-misaki (Fig. 2, site 14). These boulders were derived from the Middle Miocene 497 quartz-porphyry dyke penetrating the basement sedimentary rock, which is located seaward of 498 499 the boulder group. The dyke and boulders are famous as the tourist spot of "Hashigui-iwa", 500 which means large rocks (iwa) look like bridge (Hashi) piles (gui or kui) in Japanese. Excluding smaller ones < 1m in diameter, these boulders were not moved by the 1946 CE Showa-Nankai 501 tsunami or subsequent storms (Shishikura, 2013). Shishikura (2013) recognized two major 502 503 boulder transport events based on the radiocarbon ages of sessile marine fossils, the 12-14th 504 century and the 17-18th century CE, and attributed them to the 1361 and 1707 CE tsunamis, 505 respectively.

Tōkai earthquakes corresponding to the 887 and 684 CE Nankai earthquakes have 506 507 remained unconfirmed for a long time, but have recently been substantiated by the discovery of tsunami deposits on the Otagawa Lowland (Fig. 2, site 25) in western Shizuoka Prefecture 508 509 (Fujiwara et al., 2020). Excavation walls exposed by river improvement work in the lowland 510 revealed a \sim 1-km long coast-normal and 4 to 5 m-deep cross section of the strand plain (Fig. 511 3A). Fujiwara et al. (2020) mapped four sandy tsunami deposits along the excavation walls, 512 each of which shows inundation over 2 km inland from the coast at the time of tsunami, with landward thinning and fining trends. Radiocarbon ages link the younger two tsunami deposits to 513 the 1498 and 1096 CE Tōkai earthquakes. The older two deposits indicate the occurrence of 514 515 Tōkai earthquakes near to the end of the 7th and 9th centuries CE. The late 9th century tsunami 516 deposit, integrated with the description of strong ground shaking in the Nihon Sandai Jitsuroku, demonstrates that the 887 CE earthquake was a full-length rupture of the Tokai and Nankai 517 segments (Fujiwara et al., 2020). As the late 7th century tsunami deposit lacks data to identify 518 519 the precise date of its occurrence, it is not known whether a Tokai earthquake is coincident with 520 the 684 CE Nankai earthquake.

521 For Shikoku and northern Kyushu, in addition to the summary provided by Okamura and 522 Matsuoka (2012) from 26 ponds, Baranes et al. (2016) conducted paleotsunami research in three 523 ponds. Among these 29 sites, historical tsunami deposits have been found in four locations. At 524 Ryujin-ike in northern Kyushu (Fig.2, site 7), separated by a 17 m high and 100 m wide sand

dune, a narrow channel is the only access to the Pacific. According to historical documents, the 525 1707 CE Hoei tsunami, over 10 m-high, was much higher than that of 1854 and 1946 CE 526 tsunamis around Ryujin-ike (e.g., Chida and Nakaue, 2007). Among 40 sand beds intercalated 527 within the 5.4 m long cored humic mud sequence, eight prominent beds, each 5 - 20 cm-thick, 528 are interpreted as washover deposits due to their thinning landward trend and the inclusion of 529 530 beach sand and marine bioclasts (Okamura and Matsuoka, 2012). Based on radiocarbon dating, 531 Okamura and Matsuoka (2012) linked the youngest three beds to the 1707, 1361 and 684 CE tsunamis. 532

Baranes et al. (2016) reported a total of nine washover sand beds from the back-barrier 533 lakes facing the Bungo Channel in western Shikoku (Fig. 2, site 8). A 9.3 m-deep sediment core 534 535 from Lake Ryuuō, consisting of mainly clay beds and covering ~2500-year-long record of 536 deposition, shows an environmental change from a brackish lake to freshwater conditions that occurred around 1000 CE (about 6 m below the lake bottom). Nine washover sand beds occur in 537 the freshwater clay between the depths of 595 and 271 cm, accumulated from 1000 to 1800 CE. 538 539 Although with substantial chronological uncertainties, their depth-to-age model suggests the 540 possibility that four washover sand beds represent the 1099, 1361, 1605, and 1707 CE tsunamis. 541 There are two sand layers deposited around 1500 CE, one of which may be due to a Nankai earthquake that is unknown from historical records and paired with the 1498 Meiō Tōkai 542 543 earthquake. Baranes et al. (2016) concluded that the 1707 CE tsunami, which left the thickest (25 cm) coarse-grained sand bed, was one of the most significant floods over the last 544 millennium in the region. 545

546 Fourteen sandy tsunami deposits ranging from 4800 to 1300 cal BP in age are reported 547 from Tadasu-ike (Fig.2, site 9), located 800 m inland from the modern coast and behind a 5 m 548 high beach ridge (Okamura and Matsuoka, 2012). The sand beds range from 5 to 27 cm in thickness. The pond was inundated by the 1707 and 1946 CE tsunamis, however the record of 549 the last 1300 years is missing in this pond due to artificial disturbance; the youngest bed was 550 551 linked to the 684 CE tsunami (Okamura and Matsuoka, 2012). Kaniga-ike in Shikoku (Fig.2, 552 site 10), located about 400 m inland from the modern coast and fringed by 6 m-high coastal dunes, records six sandy tsunami deposits in the last 2000 years (Okamura and Matsuoka, 2012). 553 The tsunami beds range from 5 to 75 cm in thickness. Based on historical documents and 554 555 limited radiocarbon ages (Cabinet Office, 2011a), the youngest two can be correlated with the 556 1854 CE Ansei-Nankai and 1707 CE Hoei earthquakes, while the third and fourth ones from the top may be attributed to the 1361 and 684 CE Nankai earthquakes, respectively. 557

558

559 Prehistoric tsunami deposits

Tsunami deposits before the oldest documented Nankai Trough earthquake, the 684 CE Hakuho earthquake, have been recorded as far back as 6000 years ago. Kitamura and Kobayashi (2014a) reported four possible tsunami sand beds from a muddy shallow bay sequence in the Shimizu Plain (Fig. 2, site 29), which are dated to 6180–6010 to 5700–5580, 5700–5580 to 5520–5320, 4335–4125 to 4250–4067, and 3670–3540 to 3500–3360 cal. BP, respectively. Kitamura et al. (2013) identified two possible tsunami sand beds from a back-marsh sequence in the Shizuoka Plain covering over the past 4000 years (Fig. 2, site 28).

567 Drowned valleys on the Shima Peninsula archive a long and numerous record of possible 568 paleotsunamis. Okahashi et al. (2005) reported up to 12 washover beds deposited from 6,000 to 569 1,600 cal BP at Osatsu (Fig. 2, site 18). At Shima (Fig. 2, site 17), Fujino et al. (2018) reported 570 seven washover sand beds beneath the three historical tsunami beds. Their depth-to-age model, 571 based on radiocarbon dates, indicates that these ten tsunami beds were deposited from about 572 4,500 to 500 cal BP, with an interval of 100 to 600 years.

573 Lake Hamana, a large brackish lagoon on the Enshu-nada coast (Fig. 2, site 23), has also 574 been targeted for paleotsunami studies by several research groups. Tsuji et al. (1998) analyzed 575 cores of up to 2 m in length from the flood-tide delta and interpreted eight exotic coarse-grained 576 beds with gravel and marine shells as possible tsunami deposits. The four older deposits formed between about 4,800 and 3,200 cal BP; storm surges and tidal channel migration also remain 577 578 possible explanations for them. The four youngest deposits are linked with tsunami inundation in 1854 or 1707, 1498 CE, the 13th century and 1096 CE by Tsuji et al. (1998); however, more 579 580 recent recalibration of radiocarbon data highlights difficulties with ascribing particular historical 581 tsunamis to these deposits (Garrett et al., 2016). Sato et al. (2016b) identified a marine incursion 582 event around 4790 – 4420 cal. BP based on fossil diatom analyses of a sediment core obtained from the central Lake Hamana. 583

The western Hamamatsu Plain (Fig. 2, site 24) has a maximum width of about 4 km and 584 six rows of beach ridges (Fig. 4A). Possible tsunami sand beds intercalated with mud and peat 585 586 were reported from two separate swales (Sato et al., 2016a) and an incised valley (Fujiwara et 587 al., 2013a). Sato et al. (2016a) does not state that the sand beds are tsunami deposits, but 588 additional evidence suggests a possible tsunami origin for them. Each sand bed, generally less than 20 cm thick, shows an erosive base and upward fining trend, and records the occurrence of 589 590 sediment flow in the back-marsh. A rapid increase in marine and brackish diatoms recognized 591 within and just after the deposition of some of the sand beds and suggests coastal subsidence 592 coincident with the marine incursion (Fig. 4D). The uppermost tsunami deposits are progressively younger with decreasing distance from the present coastline. In a 593 594 landward swale (Fig. 4B) and in the Rokkengawa lowland (Fig. 4E), 4 km inland from the 595 modern coast, the uppermost deposit dates to just before the fall of the Kawagodaira pumice

(1210-1187 cal BCE, Tani et al., 2013). In a more seaward swale, 2 km from the present coast
(Fig. 4C), the uppermost deposit dates to 775-895 CE.

In Shikoku and Kyushu, five tsunami deposits have been recognized from Ryujin-ike (Fig. 598 2, site 7) between 3300 and 1600 years ago, thirteen from Tadasu-ike (Fig. 1, site 9) between 599 3300 and 1600 years ago, and two from Kaniga-ike (Fig. 2, site 10) between 1000 BCE and 600 600 CE (Okamura and Matsuoka, 2012). Tanigawa et al. (2018) reported four sandy washover 601 602 deposits tapering landward from sediment cores drilled in a small coastal lowland in Nankoku, Kochi Prefecture (Fig. 2, site 11). The lowland is separated from the Pacific by dunes 450 m 603 wide and 13 m high. The sand beds are laterally extensive through marine-brackish clay and 604 overlying freshwater clay and peat. Tanigawa et al. (2018) interpreted the sand beds as having 605 606 been deposited by tsunamis or unusually large storm surges. The possible tsunami sand beds 607 were found in sediments with ages between 5970 and 2440 cal BP. They attributed the lack of marine incursions after 2440 cal BP to the development of beach ridges that protected the site 608 609 from tsunamis and storm surges.

610 Shimada et al. (2019) reported five tsunami inundations between 5581 and 3640 cal BP 611 from a small drowned valley in Mugi (Fig. 2, site 12), eastern Shikoku. Array coring along 612 the valley axis and diatom analyses of the cored sediments revealed the deposition of at least nine exotic sand and gravel beds in the muddy brackish-to-fresh wetland sequence. 613 Environmental changes inferred from diatom assemblages suggest the occurrence of coastal 614 subsidence associated with the deposition of three of the sand beds. Subsidence of 0.3 m or 615 more was reported around the study site during the 1946 CE Showa-Nankai earthquake (Japan 616 617 Coast Guard, 1948). Shimada et al. (2019) tentatively attributed two other sand beds to tsunami 618 events, while acknowledging the possibility of deposition from a storm surge or far-field 619 tsunami. Reported tsunami deposits can be traced up to 240 m or more inland from the current shoreline, and they range in thickness from 5 to 20 cm with some exceptions up to 75 cm. 620

621

622 3.2. Transitional boundary of Nankai Trough and Ryukyu Trench (Hyūga-nada)

623 3.2.1. Historical earthquakes and tsunamis

The Hyūga-nada region covers the western end of Nankai Trough and the northern end of Ryukyu Trench. Seismic activity in this region is relatively high and small tsunamis were detected following the 1941, 1961, 1968, and 1984 CE earthquakes (Hatori, 1985). Historically, large tsunamis occurred in 1662 and 1769. The 1662 CE earthquake was the largest historical event, with a magnitude of 7.6 (HERP, 2004) and a maximum tsunami height of <5 m at Miyazaki Plain (Hatori, 1985). Hatori (1985) estimated that the 1662 CE tsunami could have inundated the lowland of Miyazaki Plain via the large rivers, while tsunamis might not have been able to overtop the coastal sand dunes because of their height (~10 to 15 m, Nagaoka et al.,1991).

633

634 3.2.2. Paleotsunami deposits

Paleotsunami studies, both historical and prehistoric, are very scarce in this region; a major reason is probably the lack of suitable lowlands. Although the Miyazaki Plain is one candidate site for paleotsunami research, the coastal dunes are too high and the rivers and floodplains are large. Therefore, it is not straightforward to locate tsunami deposits in such a depositional environment. Nevertheless, Ichihara et al. (2015) reported a possible tsunami deposit at around 2000 years ago on this plain.

Further south at Kushima City (Fig. 2, site 6), Yamada et al. (2020) conducted paleotsunami
research and found only one tsunami deposit at 4600 years ago during the period between 2000
and 5000 years ago. Based on this result, they suggested that tsunamigenic earthquakes
probably only occur at low frequencies in the Hyūga-nada region.

645

646 **3.3. Paleotsunamis on the Ryukyu Islands**

647 3.3.1. Historical earthquakes and tsunamis

Most islands in the Ryukyu Islands are surrounded by fringing reefs; these are typically 648 649 wider in the south of the island chain (\sim 1.5 km wide) and narrower in the north (a few hundred 650 meters wide). Based on the geographical distribution and paleotsunami records, we divide these islands into three groups: the Amami, Okinawa and Sakishima Islands. The three groups face 651 the northern, central, and southern Ryukyu Trench, respectively (Fig. 1). In between Kyushu 652 653 and the Amami Islands, there are several islands which we group as the Osumi and Tokara 654 Islands. Since no paleotsunami generated by subduction zone earthquake have been reported, we do not discuss this region in this study. The Ryukyu Islands align parallel to the trench and 655 hence each island has an approximately equal chance of being affected by tsunamis if 656 657 earthquakes are generated along the trench.

658 In contrast to Nankai Trough region, historical records from the Ryukyu Islands are only available for the last ~400 years. Historical and instrumental data show that earthquake and 659 tsunami occurrence in this region is remarkably low in frequency. Indeed, the only earthquakes 660 661 with magnitudes >8 during the past ~400 years were in 1911 CE in the northern Ryukyu Trench 662 (Mw=8.0, e.g., Tsuji, 1997), in 1791 CE in the central Ryukyu Trench (Mw=8.0 - 8.3, Nakamura and Kinjou, 2013; Tadokoro et al., 2018), and in 1771 CE in the southern Ryukyu 663 Trench (Mw>8.0, e.g., Nakamura, 2009). However, because there is no record of ground 664 shaking accompanying the 1791 CE tsunami, it is uncertain whether this resulted from a 665 666 near-field tsunamigenic earthquake (Nakamura and Kinjou, 2013; Tadokoro et al. 2018) or was potentially a far-field tsunami generated somewhere in the Pacific Ocean (Matu'ura, in press).
The 1771 CE tsunami was exceedingly large with maximum run-up heights of approx. 30 m
(e.g., Goto et al., 2010a). However, based on detailed historical records, it is also well known
that the tsunami-affected area was spatially limited to a narrow range of the Sakishima Islands
(Goto et al., 2010a).

Lowland plains suitable for paleotsunami research using sandy tsunami deposits are extremely scarce because of the fringing reef environment. On the other hand, numerous coral boulders are scattered on the reefs surrounding many islands and they are considered as useful geological evidence of past extreme wave events such as tsunamis and storm waves (Fig. 3B, e.g., Kawana and Nakata, 1994; Goto et al., 2010a).

In this section, we summarize paleotsunami research on the Ryukyu Islands based on the comprehensive review by Goto (2017) with additional recent works. As a brief summary, the paleotsunami history of the Amami and Okinawa Islands can be characterized by scarce (or even no) evidence of large tsunamis over the last few thousand years, while the Sakishima Islands can be characterized by frequent large tsunamis with intervals of a few hundred years.

682

683 3.3.2. Paleotsunami deposits on the Amami and Okinawa Islands

684 On the Amami Islands, Kawana and Nakata (2003) estimated the possible occurrence of tsunamis in 360-510 cal BP, 1390-1770 cal BP, and 2100-2110 cal BP, based on dating coral 685 boulders on the reef. Similarly, Kawana (2006) and Iwai and Kawana (2008) suggested possible 686 large tsunamis around 3400 years ago on the Okinawa Islands, based on the dating of beach 687 688 rocks and coral boulders. However, following studies by Goto et al. (2009, 2013), field surveys 689 and remote sensing analyses revealed that all of the coral boulders deposited on the reefs 690 fringing the Amami and Okinawa Islands can be explained by transport during storms despite their heavy weight (~160 tons). The boulders are deposited close to the source area (i.e. around 691 692 the reef edge) and have not been transported a long distance. Since the reef was formed and the 693 boulders were emplaced on the reef at least 2300 years ago (Kawana and Nakata, 2003), the 694 present clast size and spatial distribution of boulders reflects the cumulative impacts of many 695 storm waves since at least 2300 years ago. If large tsunamis had occurred during this period, boulders originally deposited on the reef by storm waves should be transported further inland, as 696 697 is observed in the Sakishima Islands (discussed below). Therefore, Goto et al. (2013) concluded 698 that no tsunamis large enough to significantly modify the characteristic distribution of storm 699 wave boulders have affected these islands in the last 2300 years. Indeed, using numerical 700 modeling, Minamidate et al. (2020) confirmed that boulders in the Okinawa Islands can be 701 explained by the realistic size of storm waves generated by past typhoons.

Another important paleoseismological aspect, especially in the Amami Islands, is a rapid 702 uplift of Kikai Island (e.g., Webster et al., 1998). At Kikai Island, the terrace of MIS 5e is at an 703 elevation of 224 m (e.g., Ota and Omura, 2000). At least 4 emerged Holocene coral terraces are 704 705 observed and each terrace is associated with emergence of 1 to 4 m (Webster et al., 1998; Sugihara et al., 2003; Hongo, 2010). Because of these geomorphological features, it is thought 706 707 that Kikai Island was intermittently uplifted by large earthquakes (Nakata et al., 1978). On the 708 other hand, considering the absence of tsunami boulders but presence of storm boulders both on Kikai and Amami-Oshima Islands (Goto et al., 2013), earthquakes that might have caused the 709 710 uplift of the island may not have generated large tsunamis. The uplift might be explained by slip on an intra-plate fault, such as a splay fault, rather than slip on the subduction interface (Goto, 711 712 2017). This idea may better fit with the fact that nearby Amami-Oshima Island (~40 km to the 713 west) has not been uplifted significantly compared with Kikai Island (Ikeda, 1977) and hence these islands are tilted within a narrow area. Alternatively, Shikakura (2014) numerically 714 modeled crustal movements and the formation of the coral terraces at Kikai Island and 715 716 suggested that large earthquakes are not mandatory requirements, but steady uplift with some 717 intermittent earthquakes may alternatively be enough to explain the uplift as well as the 718 formation of multiple coral terraces.

719

720

3.3.3. Paleotsunamis on the Sakishima Islands

721 Because of historical records that describe the movement of huge boulders during the 1771 722 CE tsunami and the presence of numerous boulders deposited on the reef and on land, identification of historically described 1771 CE tsunami boulders has proved an attractive topic. 723 724 With this objective, geological and historical studies of the 1771 CE tsunami and boulders 725 began earlier than tsunami studies in other regions, with the earliest work by Makino (1968). Makino (1968, 1981) interpreted that all boulders deposited on land were transported by the 726 1771 CE tsunami. However, following studies cautioned the need for careful identification 727 728 because many of the boulders are composed of the Pleistocene Ryukyu Limestone that forms 729 the island itself and may have been sourced from higher elevations and moved by gravity (Kato and Kimura, 1983; Kawana and Nakata, 1994). Nevertheless, there are numerous coral boulders 730 with fresh Holocene coral skeletons that must have been transported from the sea possibly by 731 732 large tsunamis during Holocene (Kato and Kimura, 1983; Kawana and Nakata, 1994).

733 Radiocarbon dating of coral boulders has been performed since early 1980's and boulders 734 with ages around 1771 CE have been discovered, suggesting their origin from this tsunami (e.g., 735 Kato and Kimura, 1983; Kawana and Nakata, 1994). However, many studies have revealed that 736 there are boulders with much older ages than 1771 CE, suggesting that these boulders are likely

to have been deposited by repeated tsunamis during the few thousand years before 1771 CE
(Kawana and Nakata, 1994; Omoto, 2012; Araoka et al., 2013).

- 739 Problems associated with pre-2010 CE studies on these boulders include 1) discrimination 740 of tsunami boulders from storm boulders, and 2) dating accuracy and reliability of tsunami age estimation. Regarding the first issue, Goto et al. (2010b, 2013) investigated clast size and the 741 742 spatial distribution of boulders on the Sakishima Islands and suggested that coral and reef 743 boulders can be classified into two groups. One group has a characteristic distribution of storm wave boulders such as a narrow range in spatial distribution from the reef edge and an 744 745 exponentially fining landward trend in clast-size distribution. The boulders of this group can be considered as of storm wave origin (Goto et al., 2010b, 2013). On the other hand, the other 746 747 group can be characterized by boulders that are deposited along the coast and on land. They are 748 deposited far beyond the possible landward limit of storm wave boulders (i.e. ~300 m from the reef edge, Goto et al., 2009). Also, there is no clear clast size distribution indicative of storm 749 deposition. Based on these features, Goto et al. (2010b, 2013) identified the boulders in this 750 group as of tsunami origin. Later, Watanabe et al. (2016) numerically confirmed the tsunami 751 752 origin of the boulders in the latter group, identifying that extremely long wavelengths are 753 required to explain the deposition of boulders along the coast.
- Regarding the second issue, Omoto (2012, 2019) (Fig. 2, site 3) and Araoka et al. (2013) 754 755 (Fig. 2, site 2) carefully selected samples for dating. They limited measurements to the boulders that meet tsunami boulder identification criteria proposed by Goto et al. (2010b, 2013) and that 756 757 were composed of single massive corals (i.e. Porites sp., Fig. 3B). Based on this approach, 758 Araoka et al. (2013) estimated a tsunami recurrence interval of about 150-400 years on the 759 Sakishima Islands. It should be noted, however, that they measured ages of various sizes of 760 boulders so it is uncertain whether the paleotsunamis that cast the boulders ashore were large or small. 761
- There are some studies that focus on sandy tsunami deposits from the Sakishima Islands. 762 763 Possible sandy tsunami deposits, especially relating to 1771 CE, were previously reported by 764 archaeologists (e.g., Yamamoto, 2008; Kugai, 2011; Nakaza, 2017) and coastal engineers (Nakaza et al., 2013) following the work done by Kawana and Nakata (1994) but their detail 765 history had remained uncertain. Recently, Ando et al. (2018) performed a 120 m-long trench 766 767 survey and found that paleotsunamis may have occurred at approximately 600-year intervals 768 over the last 2000 years (Fig. 2, site 1). This observation is consistent with numerical modeling 769 of the movement of a very large boulder at southeastern Ishigaki Island by Hisamatsu et al. 770 (2014) who suggested at least 2 large tsunamis equivalent to or even larger than the 1771 771 tsunami occurred after 2000 years ago but before 1771 CE. Interestingly, the recurrence interval 772 of paleotsunamis estimated from sandy tsunami deposits (Ando et al., 2018; Kitamura et al.,

2018a, b) is longer than that estimated from boulders (Araoka et al., 2013). This discrepancy 773 774 may be explained by coastal boulders being more sensitive in recording tsunamis because even small tsunamis can transport small boulders to the coast. Therefore, evidence of small to 775 776 medium tsunamis are included in the estimation of tsunami recurrence intervals from boulders. 777 Meanwhile, sandy tsunami deposits that extended far inland may only be suitable to recognize 778 large events. Alternatively, this discrepancy might also reflect the small number of studies 779 carried out on sandy deposits in this region so further research is required. Although some 780 researchers have suggested the absence of a predecessor to the 1771 CE tsunami (Nakaza et al., 2013, 2015), geological evidence of both boulders and sandy deposits provides crucial evidence 781 782 that either small or large tsunamis have repeatedly occurred over the last few thousand years 783 around the Sakishima Islands.

784

785 4. When and where: the history of great earthquakes

This section summarizes the spatial and temporal distribution of paleotsunami evidence along the Nankai and Ryukyu subduction zones. Table 1 summarizes when the historical earthquakes occurred in these regions.

789

790 4.1. Nankai subduction zone

791 Great earthquakes occurred in the Tōkai region nine times in total, including the 792 controversial 1614 CE event, from the end of the 7th century until 1944 CE. On the other hand, 793 great earthquakes occurred in the Nankai region eight times from 684 to 1946 CE. The 1099 CE 794 earthquake was recorded with an incorrect date, and may have actually occurred in 1096 CE 795 (Ishibashi, 2016). The Tōkai and Nankai earthquakes always occur in pairs within a few days to 796 several years, except for the 1498 CE Meio earthquake. A Nankai earthquake paired with the 1498 CE Meiō Tōkai earthquake is likely to have occurred, but we have not been able to 797 definitively ascribe any of the multiple candidates from the historical record. The 1707 CE Hoei 798 799 and 887 CE Nin-na earthquakes were full-length ruptures of both the Tokai and Nankai fault 800 segments.

Some questions remain about the recurrence pattern of Nankai Trough earthquakes. First, 801 the recurrence intervals of earthquakes in the Tokai and Nankai regions appears to be different 802 803 before and after the 1361 CE earthquake (Table 1). Even if the 1614 CE earthquake did exist, it 804 seems unsuitable to include it within a series of great Nankai Trough earthquakes because the 805 seismic intensity and tsunami during the 1614 CE event were much smaller than other 806 earthquakes. If this earthquake is included, the recurrence interval from the 1361 CE earthquake 807 to the 1854 CE earthquake were 137 years (1361 - 1498 CE), 116 years (1498 - 1614 CE), 93 808 years (1614 - 1707 CE), and 147 years (1707 - 1854 CE). On the other hand, before 1361CE,

recurrence intervals were much longer; 203 years (684 - 887CE), 209 years (887 - 1096 CE),
and 265 years (1361 - 1096 CE). It is unlikely that the process of plate boundary earthquakes
suddenly changed at some point and, while it is possible that the apparent change is an artifact
of the small sample number, other explanations may also be considered.

It is possible that other great earthquakes may be missing from the documented record due 813 to a range of circumstances (Koyama, 1999). However, if M8-class earthquakes repeated at 814 815 intervals of ~ 100 years before 1361 CE, due to the large Medieval population, it would be surprising if no historical documents recorded their occurrence. Reflecting the sensitivity with 816 817 which historical records from this period document earthquakes, in the period between 1200 and 1260 CE, there are over 100 earthquakes detailed in documents from Kyoto (Ishibashi, 2014). 818 819 Nevertheless, liquefaction features suggest the occurrence of a Tokai earthquake between 1096 820 and 1361 CE and evidence for a Nankai earthquake between 1099 and 1361 CE has been found 821 from archaeological sites in southern Osaka and the southern Kii Peninsula (Sangawa, 2007). Further searches for tsunami deposits are necessary to identify whether there are additional 822 823 historical Nankai Trough earthquakes other than those shown in Table 1.

824

825 4. 2. Ryukyu subduction zone

In contrast to the Nankai Trough region, there seems to be little evidence of large earthquakes (Mw>8) and tsunamis from the Hyūga-nada or the northern and central Ryukyu Trench regions over the last few thousand years. However, more data needs to be collected, especially from the Hyūga-nada region.

Large tsunamis have repeatedly occurred around the southern Ryukyu Islands. If submarine landslides have contributed to enhancing tsunami magnitudes (Miyazawa et al., 2012; Okamura et al., 2018), it is difficult to estimate earthquake magnitudes. However, it is probably reasonable to infer from the paleotsunami record that Mw>8 earthquakes probably occurred, with some potentially triggering submarine landslides (Hisamatsu et al., 2014).

835

836 5. Variation in size and recurrence time

The size of earthquakes and tsunamis occurring on the Nankai and Ryukyu subduction 837 zones shows a wide variation. However, lesser tsunamis may not leave a trace in the geological 838 839 record. Of course, the creation and preservation potential of tsunami deposits can change 840 depending on the geological and geomorphological conditions of the place where the tsunami 841 hit. For example, intervals between 10 tsunami deposits left in the Shima Lowlands (Fig. 2, site 842 17) between 4500 and 500 years ago show variations from 100 to 600 years (Fujino et al., 2018), 843 which are longer than expected from the historical Nankai Trough record (Fig. 2). The lack of 844 intervening tsunami deposits may simply be due to the lesser size of those tsunamis. Only the

larger tsunamis that reached the study site with sufficient stream power would be recorded inthe sediment record.

In that sense, it is interesting that traces of the 1361 CE tsunami were not found between the 1096 and 1498 CE tsunami deposits in the Ota River Lowland, where a careful investigation was conducted by Fujiwara et al. (2020) on a large outcrop over a length of 1 km (see chapter 3.1.2). This probably indicates that the 1361 CE tsunami was too small to inundate the study area, or that, if the tsunami reached the area, its stream power was too weak to form an identifiable tsunami deposit.

To enhance the contribution of the geological record, it is necessary to expand the search 853 for paleotsunami deposits to sediments that have not been targeted so far. For example, the 2011 854 855 Tōhoku tsunami widely left fine-grained deposits, mainly clay beds, inland from the area where 856 sandy deposits are distributed (e.g., Goto et al., 2011; Fujiwara, 2015). If the mud supply is sufficient, such a mud layer may be left behind after the tsunami. However, muddy 857 paleotsunami deposits are likely to have been overlooked in previous studies. Development of 858 859 criteria and techniques for identifying such muddy tsunami deposits in the geological record 860 will help to expand the range of available sites.

861 Analyses of seawater components (Chagué-Goff et al., 2017) and marine biomarkers in the sediment (e.g., Shinozaki et al., 2015) are among several promising new ways to solve this 862 863 problem. While efforts should be made to develop approaches for identifying tsunami deposits, geological methods alone have their limitations. For example, identification of the difference 864 between a tsunami and a storm surge by sediment transport modeling is also expected 865 866 (Watanabe et al., 2018). Although many studies have performed paleotsunami surveys on 867 low-lying plains or coastal lakes, it is important to expand the survey area to the other regions such as narrow valleys (e.g., Fujiwara, 2015; Abe et al., 2020) in order to increase number of 868 potential sites in the region. 869

There are only a few "treasure troves" where many tsunami deposits are recorded. Methods 870 871 for accurately dating the sediments and estimating the tsunami size from such records may have 872 not been sufficiently developed. Recent advances in coring, non-destructive testing technology (Falvard and Paris, 2017, Paris et al., 2020), chemical analysis methods (e.g. Chagué-Goff et al., 873 2017) and geophysical sensing techniques (e.g., Obrocki et al., 2020) will drive the 874 875 development of more complete tsunami records from such key sites. Advances in radiocarbon 876 dating and calibration techniques will enable more accurate estimation of the occurrence timing 877 of great earthquakes and tsunamis (Ishizawa et al., 2020).

Progress in methods for quantifying the tsunami size from deposits are needed to elucidate the variations in earthquake size. Most lowlands along the Nankai Trough coast are small, with mountains and hills close to the coast. In such a situation, tsunamis reach the landward margin of the lowland irrespective of their size, and it is not possible to compare the differences in inundation distances. The tsunami height could, however, be estimated using sediment transport modeling, using parameters such as grain size, the thickness of tsunami deposits, and the height of coastal barrier (cf. Baranes et al., 2016). Tsunami boulders may be a useful tool for estimating local tsunami size especially in the Ryukyu Islands (e.g., Watanabe et al., 2016). However, the future improvement of models and development of methodologies to collect field data about the boulder source and shape are required for further high accuracy modeling.

888

889 **6. How big**?

As the 2004 Sumatra-Andaman and 2011 Tohoku earthquakes demonstrated, an 890 891 earthquake much larger than any in the region's instrumental or documentary history may occur 892 in the future. Paleoseismological evidence shows that these extraordinary events have occurred repeatedly along subduction zones around the world, including the Kuril Trench (e.g., 893 Nanayama et al., 2003), northern Japan Trench (e.g., Sawai et al., 2012; Sawai, in press), 894 Peru-Chile Trench (e.g., Cisternas et al., 2005), and Sunda-Java Trench (e.g., Malik et al., 2011). 895 896 These giant earthquakes have much longer (\sim 300-500 years or longer) recurrence intervals than 897 those of great earthquakes (M8-class) that have occurred in each subduction zone. According to the available paleotsunami evidence, the Nankai and Ryukyu subduction zones may not be 898 899 exceptions.

900

901 6.1. Nankai subduction zone

902 An updated fault model for the 1707 CE Hoei earthquake based on crustal deformation 903 inferred from the GPS network and evidence of tsunami deposits at Ryujin Pond (Fig. 2, site 7) 904 suggest that its source area extended 70 km further to the west of the 1854 earthquake (Furumura et al., 2011). Okamura and Matsuoka (2012) suggested that earthquakes of similar 905 magnitude to the 1707 event could occur every 300-500 years along this subduction zone. The 906 907 number of tsunami deposits varies from site to site and they are generally less frequent than 908 expected from the frequency of historical earthquakes. This may be the result of only larger tsunamis overwashing the coastal barriers and leaving deposits in back-barrier ponds. The 684, 909 1361 and 1707 CE tsunamis selectively recorded at the Ryujin, Ryuuō, and Kaniga-ike Pond 910 (Fig. 2, sites 7, 8 and 10) may therefore be relatively larger than other historical tsunamis 911 912 (Okamura and Matsuoka, 2012; Baranes et al., 2016). If larger tsunamis result from larger 913 earthquakes, this indicates that larger earthquakes with 300 - 500-year recurrence intervals and 914 smaller earthquakes with shorter recurrence intervals coexist in the Nankai Trough region, 915 suggesting the recurrence pattern of this region is not periodic, but may still be time predictable 916 or slip predictable (Fig. 5A).

Okamura and Matsuoka (2012) also pointed out that the historical period of 1300 years 917 918 may not be enough to consider the maximum-size earthquake and tsunami of this region. The 919 discovery of tsunami deposits thicker than those from the tsunami that followed the 1707 CE 920 Hōei earthquake, the largest in the Nankai region's history, from Tadasu-ike (Fig. 2, site 9), 921 Kaniga-ike (Fig. 2, site 10), Kamoda-oike (Fig.2, site 13) and Sugari-oike (Fig. 2, site 15), all of which were aged around 2000 years ago, led Okamura and Matsuoka (2012) to hypothesize the 922 923 occurrence of a giant tsunami. However, it is unreasonable to discuss the size of a tsunami 924 based on the thickness of tsunami deposits alone. The thickness of different tsunami deposits 925 within a site may be influenced by the state of the tide and the time of year, which controls vegetation cover and therefore sediment transport. Furthermore, changes in coastal 926 927 geomorphology over time may change a site's susceptibility to accumulating and preserving 928 tsunami deposits (Garrett et al., 2016). Underdeveloped low dunes and beach ridges can allow large amounts of overflow, even during lesser tsunamis, resulting in the formation of thicker 929 tsunami deposits in back-barrier ponds (Fujiwara, 2015). Additionally, the estimated ages for 930 the thick tsunami bed in each of Okamura and Matsuoka's (2012) ponds has substantial 931 932 uncertainties of hundreds of years. Consequently, it is ambiguous whether the thick tsunami bed 933 in each of the ponds was formed by one extremely large tsunami or series of closely spaced 934 large tsunamis.

935 Deposition of a thick tsunami deposits around 2000 years ago is not seen at Nankoku (Fig. 2, site 11) (Tanigawa et al., 2018) or Ryuuō Pond (Fig. 2, site 8) (Baranes et al., 2016). In 936 937 addition, at Ryūjin Pond in northern Kyushu, a sand bed deposited around 3300 years ago is the 938 thickest, with only a minor sand bed deposited around 2000 years ago. Thick tsunami deposits 939 formed around 2000 years ago have not been found from the Tokai region (Fujiwara, 2013; 940 Fujiwara and Tanigawa, 2017). The late 7th century tsunami deposits in the Otagawa Lowland is thick, over 40 cm more than 2 km inland from the coast at the time of tsunami occurrence, but 941 its distribution is limited to the topographic lows in the lowland (Fujiwara et al., 2020). 942 943 Fujiwara et al. (2020) interpreted that the tsunami was not so large and inundation and sand bed 944 deposition was limited to the topographic depressions.

Kitamura (2016) summarized the spatio-temporal distribution of reported tsunami deposits from the eastern coast of the Nankai Trough, including sites 23, 24, 26, 28, 29, and 32 in Figure 2, and concluded that no tsunami deposits with a wide regional distribution that might suggest the occurrence of "level 2" tsunami over the past 4000 years have been found in this area. However, due to the limitation in the age control of the tsunami deposits, it would be difficult to deny the existence of a giant tsunami on this basis alone.

951

952 6.2. Hyūga-nada and the Ryukyu subduction zone

Historical and geological evidence is too scarce to evaluate whether large tsunamigenic earthquakes have been generated in the Hyūga-nada region. Historical records suggest no large earthquakes with magnitude >8 have occurred over the last 400 years in this region. Geological evidence from the Pacific coast of central and southern Kyushu is probably still insufficient to discuss paleoseismicity in this region, although recent works suggest the rare occurrence of tsunamigenic earthquakes (Ichihara et al., 2015; Yamada et al., 2020).

959 M8-class earthquakes can be generated elsewhere along the Ryukyu Trench based on 960 historical and instrumental records; however, the Ryukyu Trench is generally considered as an aseismic zone in terms of the potential occurrence of giant earthquakes of magnitude ~ 9 , 961 although potential coupling of the plate boundary and the consequent occurrence of extremely 962 963 large earthquakes along the Ryukyu Trench has been suspected by some researchers (e.g., Ando 964 et al., 2009, 2012a; Aydan and Tokashiki, 2018; Hsu et al., 2012). Geological evidence shows that tsunamigenic earthquakes may be rare or even nonexistent in the northern and central 965 Ryukyu Trench (Goto et al., 2013). Even though Kikai Island in the northern Ryukyu Trench 966 has been uplifted quickly and the repeated occurrence of large earthquakes has been suspected 967 (e.g., Nakata et al., 1978), relatively small earthquakes just beneath the island (Goto, 2017) or 968 969 stable uplift without large earthquakes (Shikakura, 2014) may also explain this record.

Nakamura and Sunagawa (2015) studied the distribution of very low frequency 970 971 earthquakes (VLFEs) along entire the Ryukyu Trench. They suggested that occurrence of 972 VLFEs is high in the northern and central Ryukyu Trench but low in the southern Ryukyu 973 Trench. Therefore, it is likely that the southern Ryukyu Trench has potential to generate large earthquakes, unlike the northern and central regions (Nakamura and Sunagawa, 2015). 974 975 Nishimura (2014) also suggested that short-term slow slip events (SSEs) cluster near the areas 976 where large earthquakes have occurred historically (e.g., the 1911 and 1771 CE events). Nevertheless, the sources of these large earthquakes do not overlap with the clusters of 977 short-term SSEs and the reason remains uncertain. 978

979 Arai et al. (2016) and Nakamura (2017) suggested there is a strongly coupled zone at 980 shallower plate boundary depths in the southern Ryukyu Trench. Therefore, localized tsunamigenic earthquakes at the southern Ryukyu Trench inferred from tsunami boulders (Goto 981 et al., 2013) may be explained by the heterogeneity of the coupling along the Ryukyu Trench. 982 983 Nevertheless, the source model of the 1771 CE tsunami is still controversial with proposed 984 models including an intra-plate earthquake plus a submarine landslide (Imamura et al., 2008; 985 Miyazawa et al., 2012), a tsunami earthquake (Nakamura, 2009), splay faulting (Hsu et al., 2013), and a large submarine landslide near the trench axis (Okamura et al., 2018). By assuming 986 987 a large subduction zone earthquake along the Ryukyu Trench, Ando et al. (2018) estimated a 988 seismic coupling coefficient of 20%. Alternatively, if a submarine landslide played an important role in the generation of the 1771 CE tsunami and predecessors, then a special tectonic state capable of generating large earthquakes only at the southern Ryukyu Trench may not need to be considered. Relationships among VLFEs, short-term SSEs, and large earthquakes (Mw \sim 8) are still uncertain, so further seismological research with reference to paleotsunami results is required along the Ryukyu Trench.

- 994
- 995

996

997 7.1. Earthquake recurrence model

7. Knowledge gaps and future work

998 Since Kanamori and McNally (1982) explained a series of different size earthquakes off 999 the Colombian coast as the rupture of different combinations of multiple asperities, similar 1000 phenomena have been reported from subduction zones around the world. Some examples show 1001 that giant (M9-class) and large (M7-8 class) earthquakes may share fault plane asperities, and 1002 the former have much longer recurrence intervals than the latter (Cisternas et al., 2005, 2017). 1003 The Nankai and Ryukyu subduction zones may be no exception. This may indicate that the 1004 recurrence pattern of subduction zone earthquakes is "nonpredictable" (Fig.5A).

1005 The recurrence of Nankai Trough earthquakes has generally been explained as the repeated rupture of all or part of characteristic fault planes (segments) A, B, C, D and E (Fig. 1 and Table 1006 1007 1). Observations that the amount of uplift during the 1707, 1854, and 1946 CE Nankai earthquakes at Cape Muroto was proportional to the length of the interval preceding each 1008 1009 earthquake led to the proposal of the time-predictable recurrence model (Fig. 5A) by Shimazaki 1010 and Nakata (1980). Kumagai (1996) applied this model to Nankai Trough earthquakes using the time series of great earthquakes since 684 CE. However, according to current knowledge, 1011 1012 Kumagai's series of historical earthquakes contains some uncertainties, and the reliability of the recurrence model is ambiguous. For example, the 1230 CE earthquake and the 1498 CE Meiō 1013 earthquake in the Nankai region (as opposed to the undisputed Tokai event) in his series are still 1014 1015 undetermined. Results of paleotsunami research in the Otagawa Lowland (Fujiwara et al., 2020) 1016 does not support the time-predictable recurrence model. According to this study, the 1361 CE 1017 Tōkai earthquake has the longest pre-earthquake interval (265 years following the 1096 CE earthquake), but its tsunami seems to have been less extensive than others, suggesting a smaller 1018 1019 magnitude of earthquake (see chapter 5).

Seno (2012) proposed a new earthquake recurrence model for Nankai Trough earthquakes that challenged the traditional idea of the characteristic earthquake model, which partitions the Nankai Trough seismogenic zone into ruptures A-E and assign the epicenters of historical earthquakes to them. He assumed two complementary earthquakes that have different non-overlapping asperities mainly based on the spatial distribution of seismic intensities, while 1025 also considering tsunami heights and coastal deformation estimated from documented records 1026 and liquefaction features in archaeological sites (Fig. 5B). According to his idea, historical great 1027 Nankai Trough earthquakes are classified into two types; "Hōei-type" earthquakes (887, 1361, 1707 and 1944+1946 CE) and the "Ansei-type" earthquakes (684, 1096+1099, 1489, 1028 1029 1854+1854 CE), which alternately occur with recurrence intervals of around 350 and 400 years, 1030 respectively. Seno's (2012) model seems to explain the variations in the recurrence intervals 1031 and extent of source area of each earthquake in Nankai Trough; however, the reliability of model is reduced by the decrease in the number of reliable documentary records with increasing 1032 age, especially before the 1096 CE earthquake. 1033

In contrast to the Nankai Trough region, the establishment of an earthquake recurrence 1034 1035 model along the Ryukyu Trench is not straightforward because of limited historical and 1036 geological data. Presently available data suggest that earthquakes with magnitudes of around 8 have occurred patchily in space and time from the northern to the central Ryukyu Trench. The 1037 1038 repeated occurrence of these earthquakes is uncertain. On the other hand, in the southern 1039 Ryukyu Trench, the repeated occurrence of large tsunamis is suggested (Omoto, 2012; Araoka 1040 et al., 2013; Ando et al., 2018). Therefore, the future occurrence of large earthquakes in this region is likely. 1041

The amount of reported paleo-tsunami data is still insufficient to update the earthquake recurrence model in these regions (Figure 2). Updating the model also requires improved accuracy in the dating and the identification of tsunami deposits, which is a particularly important issue with sites surveyed in the early stages of the development of the field. Re-examination of these sites with current knowledge and technology, as discussed in chapter 5, may reveal some rules in the recurrence mode of the earthquakes in these regions.

1048 Coastal deformation data will enhance the revision of earthquake recurrence models (e.g., 1049 Garrett et al, 2016). In particular, uplift along the western Suruga Bay coast, including Cape 1050 Omaezaki (representing the rupture of segment E), and the southern Kii Peninsula (representing 1051 the rupture of segment C) is key to proving the hypothesis of Ansei- and Hōei-type earthquakes 1052 (Fig. 5B). Kitamura et al. (2019) suggested a possible seismic uplift occurred around 400 CE 1053 from the Shimizu Plain (Fig. 2, site 29).

1054 Integration of offshore paleoseismic features, such as turbidite sequences, with onshore 1055 tsunami deposits also will help us to consider which recurrence model is plausible. In fact, the 1056 integrated study of coastal deformation and onshore tsunami deposits, and offshore 1057 seismo-turbidite sequences in the Cascadia subduction zone and northern San Andreas fault 1058 systems has revealed the recurrence mode of great earthquakes in this region (e.g., Goldfinger et 1059 al., 2003 a, b). These studies also contribute to regional earthquake and tsunami 1060 countermeasures through underpinning tsunami simulations (e.g., Priest et al., 2017). Although there are some initial studies on seismo-turbidites in the Nankai (Garrett et al., 2016 and
references therein) and the Ryukyu subduction zone (e.g., Ujiie et. al. 1997; Ikehara et al., 2017),
at present, coordination of onshore and offshore paleoseismic research is an area where future
progress is expected.

1065

1066 7.2. Maximum possible earthquake and tsunami

1067 As a response to the occurrence of unexpectedly huge 2011 Tohoku earthquake, the Cabinet Office (2011b) attempted to quantify the maximum possible earthquake and tsunami for 1068 the Nankai Trough subduction zone ("maximum scenario" in Fig. 1). The earthquake source 1069 consists of the strongly coupled zone of 10 to 20 km depth, which has been considered to be the 1070 1071 main seismogenic zone, with the addition of down-dip weakly coupled zones at ~ 30 to 40 km 1072 depth, which are the location of the occurrence of SSE and deep low-frequency tremor. Its western end is delimited by the Kyushu-Palau ridge. The up-dip zone from 10 km to the trough 1073 axis, which may also generate large tsunamis, is assumed as an additional tsunami source area. 1074 According to the worst scenario by the Cabinet Office (2012a), the height of the tsunami 1075 1076 occurring from this M9 earthquake could exceed 10 m over a wide area along the Pacific coast 1077 of western Japan, and reach in excess of 30 m in some places. However, no geological evidence, 1078 such as tsunami deposits, was taken into account for this catastrophic earthquake and tsunami 1079 scenario (e.g., Wallis et al., 2018). Determining when the most recent outsize earthquake and 1080 tsunami occurred and how large it was will help to address this concern.

1081

1082 Largest tsunami in the area estimated from geological data

1083 The "2 ka giant tsunami hypothesis" by Okamura and Matsuoka (2012), derived from the discovery of extremely thick tsunami deposits, played an important role in making people aware 1084 of the possibility of a giant earthquake and tsunami in the Nankai Trough region and raising 1085 1086 awareness of disaster management. Nevertheless, it has not been tested. First, it is necessary to 1087 ensure that multiple seismic events are not confused with a single major event. To solve this 1088 issue, the extremely thick tsunami deposits should be re-examined by increasing the number of 1089 reliable radiocarbon ages and using new age modeling methods (e.g., Bronk Ramsey, 2009; Lougheed and Obrochta, 2016, 2019). 1090

1091 Other causes of the thick tsunami deposit in individual ponds should also be considered. 1092 Earthquake-induced submarine landslides are a candidate source for locally thick tsunami 1093 deposits. They generate much larger, but regionally limited, tsunamis than would be expected 1094 based on the size of concurrent earthquakes (e.g., Tappin et al., 2008; Harbitz et al., 2013; 1095 Kawamura et al., 2017). In fact, many submarine landslides have been found on the submarine 1096 slopes along the Nankai Trough and some of them have areas of $\geq 10 \text{ km}^2$ (Moriki et al., 2017). 1097 The study of the evolutionary history of both coastal and submarine landforms may be the key1098 to solving this problem.

1099 Reconstruction of the inundation area is the most important indicator for determining the 1100 size of a tsunami from its deposit (e.g., Fujiwara, 2015). In the case of eastern Hokkaido, the 1101 discovery of prehistoric tsunami deposits showing inundation areas several times larger than 1102 those of historical M8-class earthquakes revealed that unusually large tsunamis repeatedly 1103 occurred on the Kuril Trench (e.g., Nanayama et al., 2003). Similarly, the 869 CE Jogan tsunami deposit showed much larger inundation areas than those of other historical earthquakes 1104 that occurred in the northern Japan Trench, suggesting an unprecedentedly large earthquake 1105 (Satake et al., 2008; Namegaya et al., 2010, Sawai et al., 2012). 1106

1107 The Hamamatsu Plain (Fig. 2, site 24) is the first candidate site where we can suggest that 1108 no giant tsunami has occurred in the eastern Nankai Trough over the latter half of the Holocene based on the tsunami inundation area shown by the tsunami deposits. The strand plain has 1109 prograded seaward over the last 6000 years following the Holocene sea-level high stand (e.g., 1110 1111 Sato et al., 2011, 2016a), as evidenced by preservation of rows of sandy beach ridges. This 1112 progradation process can be seen in the vertical change of diatom assemblages in swale deposits, which shows a change from brackish-marine to freshwater conditions (Fig. 4B). In harmony 1113 with the progradation of the coast, older washover deposits, probably including tsunami 1114 1115 deposits, are distributed inland and younger ones are found in more seaward locations (Fig. 4 B, C). Additionally, at each site, younger tsunami deposits tend to be thinner (Fig. 4 B, C). 1116 Assuming tsunamis (or other washover events) of similar size over time, the progradation of the 1117 1118 coast would make it less likely that each successive tsunami would reach the inland areas, and 1119 the observed sequence of tsunami deposits would be left on the Hamamatsu Plain (Fig. 6). If an 1120 extremely large tsunami occurred over the past few thousand years, its deposit would have extended from the coast to further inland, as shown at the top of the Fig. 6. However, no such 1121 trace has yet been found (Fujiwara, 2013; Fujiwara and Tanigawa, 2017). 1122

Along the Ryukyu subduction zone, the 1771 CE tsunami at the Sakishima Islands is the largest known event. Based on numerical modeling of boulder transport Hisamatsu et al. (2014) suggested that at least one prehistoric tsunami event was even larger than the 1771 CE tsunami. However, since information is limited, further geological data is required to clarify the size of prehistoric tsunamis.

1128

1129 Crustal movements suggesting outsized earthquakes along the Nankai Trough

1130 In discussing the possibility of a giant earthquake, it is necessary to assess a wide range of 1131 crustal deformation patterns occurring at different temporal and spatial scales. In the case of 1132 eastern Hokkaido, where the recurrence of giant tsunami along the Kuril Trench is shown by tsunami deposits, there is a contradiction between the coastal deformation pattern shown by
geological and geodetic data. In this area, 60 to 80 km above the top of the subducting plate,
marine terraces showing 20 to 50 m of net uplift in the past 125 ka (Okumura, 1996) coexist
with tide gauge and bench mark-inferred subsidence of up to 1 m in the past 100 years
(Shimazaki, 1974a; Ozawa et al., 1997). Little, if any, uplift is known from the M8-class
earthquakes in this region's written history during the past 200 years (Shimazaki, 1974b;
Kasahara, 1975; Kasahara and Kato, 1980; Yamanaka and Kikuchi, 2003).

Micropaleontological research on marsh deposits in this area (Sawai et al., 2004) revealed post-seismic uplift associated with unusually large earthquakes, which was induced by greater creep further down the plate boundary than during any of the Kuril subduction zone's historical events. The last uplift event of up to 2 m, which might resolve some of the contradiction between geologic and geodetic data, occurred over the first few decades following an outsized 17th-century earthquake.

1146 A similar phenomenon can be seen around Lake Hamana. Middle Pleistocene (MIS 9 to MIS 5) terraces, mainly of marine and partly of fan origin, are found in this area (Sugiyama, 1147 1991; Nakashima et al., 2008), showing that uplift has been dominant over 10^4 to 10^5 -vear 1148 scales. The MIS-9 marine terrace reaches 78 m in elevation and is highest on the west coast of 1149 1150 Lake Hamana, decreasing in height to the north and west. Subsidence of 0.1-0.6 m during the 1151 1854 CE Ansei-Tōkai earthquake is estimated around the Lake Hamana area (Ishibashi, 1981; Sato and Fujiwara, 2017). The northern coast of the lake is also known to have subsided during 1152 the 1707 CE Hoei earthquake (Yata, 2013). Coseismic subsidence between 7000 and 5700 years 1153 1154 ago, similar to the 1707 and 1854 CE events, was reported from Shinjo Lowland (Fig. 2, site 21) 1155 on the west coast of Lake Hamana (Sato and Fujiwara, 2017). Multiple active reverse faults (The research group for active submarine faults off Tokai, 1999) and normal faults (Arai et al., 1156 2006) parallel to the trough axis are distributed off the Hamana area. However, they cannot 1157 1158 explain the coastal deformation in the area.

1159 Major differences between this area and eastern Hokkaido include the inter-seismic uplift 1160 after the 1854 CE Ansei-Tōkai earthquake and the short-term uplift associated with SSEs. Inter-seismic uplift, averaging ~6 mm/y between 1901 and 2008 CE (Geographical Survey 1161 Institute, 2008), may compensate for the coseismic subsidence during "ordinal" Tokai 1162 1163 earthquakes; however, other factors are needed to explain the formation of high and tilted 1164 terraces. This average uplift rate includes the effect of SSEs (see chapter 2.2). SSEs seem to 1165 occur with decadal recurrence intervals and are high frequency uplift events on a geological time scale, but associated uplift seems too small to form the terraces. For example, a Tōkai slow 1166 1167 slip event (Ozawa et al., 2002; Miyazaki et al., 2006) occurred under Lake Hamana from January 2000 to July 2005, with a cumulative moment magnitude of \sim 7.1, and generated \sim 5 cm 1168

uplift in this area (Suito and Ozawa, 2009). At present the importance of SSEs for terraceformation remains unclear and their contribution may need to considered further.

1171 To elucidate the possibility of a giant earthquake, a unified explanation is required for 1172 geodetic and geologic phenomena on different timescales, including SSEs with 10^0 to 10^1 -year 1173 scales, inter-seismic uplift with a 10^2 -year scales, and tilting uplift of the Pleistocene terraces 1174 over 10^4 to 10^5 -year scales. This requires further investigation of Holocene crustal movements 1175 on a 10^3 -year scales, which connects the two.

The possibility of eastward rupture propagation from the Nankai Trough to the 1176 Fujikawa-kako fault zone (FKZ) must also be assessed in the context of defining the maximum 1177 possible earthquake size. The FKZ is an eastern extension of the Nankai or Suruga Trough and 1178 1179 might rupture in combination with future great Nankai Trough earthquakes (HERP, 2010). 1180 Nevertheless, the recurrence history is still unclear at present. Marsh deposits on the Ukishima-gahara Lowlands (Fig. 2, site 30) record a 1500-year history of repeated coastal 1181 subsidence suggesting the activity of the FKZ (Fujiwara et al., 2016). Further research is needed 1182 to elucidate the relationship between episodes of coastal subsidence and historical earthquakes. 1183

1184

1185

Possibility of full-length rupturing event of the Nankai and Ryukyu subduction zone

The Nankai and the Ryukyu subduction zones make up one plate boundary through the 1186 Hyūga-nada region, and both may have the potential to generate M9-class earthquakes. 1187 Therefore, it is important to understand the possible risk of the occurrence of extremely large 1188 earthquakes crossing this boundary, as suggested by some researchers (e.g., Furumoto and Ando, 1189 1190 2009). As reviewed in this paper, the occurrence of such an extremely large earthquake and 1191 tsunami in the past few thousand years is not supported by the presently available historical and 1192 geological data in this region. However, the spatio-temporal correlation of earthquakes that occurred on the Nankai and Ryukyu subduction zones or the possibility of a full-length 1193 mega-earthquake linking both seismogenic zones has not been evaluated well. From this point 1194 1195 of view, research integrating various scientific fields such as seismology, paleoseismology, 1196 geology, and geodesy is also an issue for the future.

1197

1198 8. Conclusions

This review illustrates the difference in the type of tsunami deposits and the progress of research between the Nankai and Ryukyu subduction zones. We also summarize the recurrence mode of great earthquakes and tsunamis in these regions based on geological evidence. The volume of information currently available about paleoearthquakes and tsunamis varies greatly different between the two subduction zones, despite the fact they comprise a contiguous northwestern boundary of the subducting Philippine Sea Plate beneath the Japanese archipelago. 1205 Information from the Ryukyu subduction zone is very limited compared to the Nankai1206 subduction zone.

The model describing the recurrence mode of great earthquakes in this region is still 1207 incomplete, and updating the model requires future studies to gather more paleoseismic data 1208 from both on land and offshore. Reconstruction of the largest event ever in the region based on 1209 geological evidence will help discussions about the "real" size of the maximum possible 1210 1211 earthquake in this region. This requires a new conceptual model that can unify geological processes that occur on various time scales. Differentiation of extremely large earthquakes and 1212 piecemeal ruptures of lesser magnitude in the geological record is also important for the 1213 accurate reconstruction of past events and corresponding forecasts of future earthquake. 1214 Progress in age estimation techniques will help to diminish this enduring problem. The potential 1215 of giant earthquakes that rupture both the Nankai and Ryukyu subduction zones remains 1216 unknown. Paleoseismological studies in Hyūga-nada region that connects the two subduction 1217 zones are key to solving this question. 1218

1219 1220

1221 Acknowledgement

1222 The part of the review concerning Nankai subduction zone paleotsunami deposits led by 1223 OF and EG was conducted as contribution to the QuakeRecNankai project supported by the 1224 Belgian Science Policy Office (BELSPO BRAIN-be BR/121/A2).

- 1225
- 1226

1227 References

- 1228
- Abe, T., Goto, K. and Sugawara, D., 2020. Spatial distribution and sources of tsunami deposits
 in a narrow valley setting insight from 2011 Tohoku-oki tsunami deposits in northeastern
 Japan. Progress in Earth and Planetary Science. doi.org/10.1186/s40645-019-0318-6.
- Ando, M., 1975a. Possibility of a major earthquake in the Tokai district, Japan and its
 pre-estimated seismotectonic effects. Tectonophysics 25, 69-85.
 doi.org/10.1016/0040-1951(75)90011-6.
- Ando, M., 1975b. Source mechanisms and tectonic significance of historical earthquakes along
 the Nankai Trough, Japan. Tectonophysics 27, 119-140.
 doi.org/10.1016/0040-1951(75)90102-X.
- Ando, M., Kitamura, A., Tu, Y., Ohashi, Y., Imai, T., Nakamura, M., Ikuta, R., Miyairi Y.,
 Yokoyama, Y. and Shishikura, M., 2018. Source of high tsunamis along the southernmost
 Ryukyu trench inferred from tsunami stratigraphy. Tectonophysics 722, 265–276.
 doi.org/10.1016/j.tecto.2017.11.007.
- Ando, M. and Nakamura, M., 2013. Seismological evidence for a tsunami earthquake recorded
 four centuries ago on historical documents. Geophysical Journal International 195, 1088–
 1101. doi.org/10.1093/gji/ggt270
- Ando, M., Nakamura, M., Matsumoto, T., Furukawa, M., Tadokoro, K. and Furumoto, M., 2009. 1245 Is the Ryukyu subduction zone in Japan coupled or decoupled? - The necessity of seafloor 1246 observation. Earth Planets 61, 1247 crustal deformation Space 1031-1039. doi: 1248 10.1186/Bf03352954.
- Ando, M., Tu, Y., Kumagai, H., Yamanaka, Y. and Lin C. H., 2012, Very low frequency
 earthquakes along the Ryukyu subduction zone. Geophysical Research Letters 39,
 L04303. doi.org/10.1029/2011GL050559.
- Ando, R., Takeda, N. and Yamashita, T., 2012b. Propagation dynamics of seismic and aseismic
 slip governed by fault heterogeneity and Newtonian rheology. Journal of Geophysical
 Research 117, B11308. doi:10.1029/2012jb009532.
- Arai, K., Okamura, Y., Ikehara, K., Ashi, J., Soh, W., Kinoshita, M., 2006. Active faults and
 tectonics on the upper forearc slope off Hamamatsu City, central Japan. Journal of the
 Geological Society of Japan 112, 749-759. (in Japanese with English abstract). doi:
 https://doi.org/10.5575/geosoc.112.749.
- Arai, R., Takahashi, T., Kodaira, S., Kaiho, Y., Nakanishi, A., Fujie, G., Nakamura, Y.,
 Yamamoto, Y., Ishihara, Y., Miura, S. and Kaneda, Y., 2016. Structure of the tsunamigenic
 plate boundary and low-frequency earthquakes in the southern Ryukyu Trench. Nature
 Communications. 7, 12255. DOI: 10.1038/ncomms12255.

- Araoka, D., Yokoyama, Y., Suzuki, A., Goto, K., Miyagi, K., Miyazawa, K., Matsuzaki, H. and
 Kawahata, H., 2013. Tsunami recurrence revealed by Porites coral boulders in the southern
 Ryukyu Islands, Japan. Geology 41, 919–922. doi.org/10.1130/G34415.1.
- Awata, Y. and Sugiyama, Y., 1989. Geological structure due to the right–lateral reverse faulting
 accompanied with the large earthquakes along the Nankai Trough). Zishin (Journal of the
 Seismological Society of Japan), 2nd ser., 42, 231–233.
- Aydan, O. and Tokashiki, N., 2018. Tsunami boulders and their implications on the potential for
 a mega-earthquake along the Ryukyu Archipelago, Japan. Bulletin of Engineering Geology
 and the Environment 78, 3917-3925. doi.org/10.1007/s10064-018-1378-3.
- Baba, T., Cummins, P.R. Hori, T. and Kaneda, Y., 2006. High precision slip distribution of the
 1944 Tonankai earthquake inferred from tsunami waveforms: Possible slip on a splay fault.
 Tectonophysics 426, 119–134. doi:10.1016/j.tecto.2006.02.015
- Baranes, H.E., Woodruff, J.D., Wallace, D.J., Kanamaru, K. and Cook, T.L., 2016.
 Sedimentological records of the C.E. 1707 Hōei Nankai Trough tsunami in the Bungo
 Channel, southwestern Japan. Natural Hazards 84, 1185-1205. DOI
 10.1007/s11069-016-2498-3.
- Bronk Ramsey, C., 1995. Radiocarbon calibration and analysis of stratigraphy; the OxCal
 program. Radiocarbon 37, 425–430.
- Bronk Ramsey, C., 2008. Deposition models for chronological records. Quat. Sci. Rev. 27, 42–
 60.
- 1283 Bronk Ramsey, C., 2009. Bayesian analysis of radiocarbon dates. Radiocarbon 51, 337–360.
- 1284 Cabinet Office, Government of Japan, 2011a. Information provided by members at the second
 1285 meeting of the Committee for evaluating the rupture model of megathrust earthquakes
 1286 along the Nankai Trough. http://www.bousai.go.jp/jishin/nankai/model/2/pdf/1.pdf.
- 1287 Cabinet Office, Government of Japan, 2011b. Interim report of the Committee for evaluating
 1288 the rupture model of megathrust earthquakes along the Nankai Trough.
 1289 http://www.bousai.go.jp/jishin/nankai/model/pdf/chukan_matome.pdf (in Japanese, last
 1290 accessed 2020/2/21)
- Cabinet Office, Government of Japan, 2012a. Second report of the Committee for evaluating the
 rupture model of megathrust earthquakes along the Nankai Trough: tsunami fault moles.
 http://www.bousai.go.jp/jishin/nankai/model/pdf/20120829_2nd_report01.pdf. (in
- 1294 Japanese, last accessed 2020/2/11)
- Cabinet Office, Government of Japan, 2012b. Second report of the Committee for evaluating
 the rupture model of megathrust earthquakes along the Nankai Trough: tsunami fault
 moles. Reference: Basic concept for tsunamis with high frequency of occurrence in the

- Nankai Trough. http://www.bousai.go.jp/jishin/nankai/model/pdf/20120829_2nd_report04.
 pdf (in Japanese, last accessed 2020/2/11).
- 1300 CDMC (Central Disaster Management Council), 2011. Report of the committee for technical
 1301 investigation on countermeasures for earthquakes and tsunamis based on the lessons
 1302 learned from the "2011 off the Pacific coast of Tohoku Earthquake"
 1303 http://www.bousai.go.jp/kaigirep/chousakai/tohokukyokun/pdf/Report.pdf (last accessed
 1304 2020/2/11).
- 1305 Chagué-Goff, C., Szczuciński, W. and Shinozaki, T. 2017. Applications of geochemistry in
 1306 tsunami research: A review. Earth-Science Reviews 165, 203-244.
 1307 https://doi.org/10.1016/j.earscirev.2016.12.003
- 1308 Chida, N., and Nakaue, K., 2007. Time series analysis of the tsunamis caused by Hoei and
 1309 Ansei Nankai earthquakes in Yonouzu, Oita prefecture, east central Kyushu, Japan. The
 1310 research bulletin of the Faculty of Education and Welfare Science, Oita University, 29, 69–
 1311 80.
- Cisternas, M., Atwater, B.F, Torrejon, F., Sawai, Y., Machuca, G., Lagos, M., Eipert, A.,
 Youlton, C., Salgado, I., Kamataki, T., Shishikura, M., Rajendran, C.P., Malik, J.K., Rizal,
 Y. and Husni, M., 2005. Predecessors of the giant 1960 Chile earthquake. *Nature* 437:
 404–407. DOI: 10.1038/nature03943
- Cisternas, M., Garrett, E., Wesson, R., Dura, T., Ely, L.L., 2017. Unusual geologic evidence of
 coeval seismic shaking and tsunamis shows variability in earthquake size and recurrence in
 the area of the giant 1960 Chile earthquake. Marine Geology 385, 101–113.
- Falvard, S. and Paris, R., 2017. X□ray tomography of tsunami deposits: Towards a new
 depositional model of tsunami deposits. Sedimentology 64, 453–477. doi:
 10.1111/sed.12310.
- 1322Fujino, S., Kimura, H., Komatsubara, J., Matsumoto, D., Namegaya, Y., Sawai, Y., Shishikura,1323M., 2018. Stratigraphic evidence of historical and prehistoric tsunamis on the Pacific coast1324of central Japan: Implications for the variable recurrence of tsunamis in the Nankai Trough.1325QuaternaryScienceReviews201,147–161.
- 1326 https://doi.org/10.1016/j.quascirev.2018.09.026.
- Fujiwara, O., 2013. Earthquakes and tsunamis along the Nankai Trough, inferred from geology
 and geomorphology: examples in Tokai Region., GSJ Chishitsu News 2, 197-200 (in
 Japanese)
- Fujiwara, O., 2015. The science of tsunami deposits. University of Tokyo Press, 283p (inJapanese).
- Fujiwara, O., Aoshima, A., Irizuki, T., Ono, E., Obrochta, S.P., Sampei, Y., Sato, Y. and
 Takahashi, A., 2020. Tsunami deposits refine great earthquake rupture extent and

- recurrence over the past 1300 years along the Nankai and Tokai fault segments of the
 Nankai Trough, Japan. Quaternary Science Reviews 277,
 doi.org/10.1016/j.quascirev.2019.105999.
- Fujiwara, O., Fujino, S., Komatsubara, J., Morita, Y. and Namegaya, Y. (2016) Paleoecological
 evidence for coastal subsidence during five great earthquakes in the past 1500 years along
 the northern onshore continuation of the Nankai subduction zone. Quaternary International
 397, 523-540. doi.org/10.1016/j.quaint.2015.11.014.
- Fujiwara, O. Hirakawa, K., Irizuki, T. Hasegawa, S., Hase, Y., Uchida, J., Abe, K., 2010. 1341 Millennium-scale recurrent uplift inferred from beach deposits bordering the eastern 1342 19. 1343 Nankai Trough, Omaezaki area, central Japan. Island Arc 374-388. 1344 https://doi.org/10.1111/j.1440-1738.2010.00729.x
- Fujiwara, O., Ono, E., Satake, K., Sawai, Y., Umitsu, M., Yata, T., Abe, K., Ikeda, T., Okamura,
 Y., Sato, Y., Aung, T.T., Uchida, J., 2007. Trace of the AD1707 Hoei earthquake from the
 coastal lowland, Shizuoka Prefecture, central Japan. Annual Report on Active Fault and
 Paleoearthquake Researches, Geological Survey of Japan, AIST, No. 7, 157–171 (in
 Japanese with English abstract).
- Fujiwara, O., Sato, Y., Ono, E. and Umitsu, M. 2013a. Researches on tsunami deposits using
 sediment cores: 3.4 ka tsunami deposit in the Rokken-gawa Lowland near Lake Hamana,
 Pacific coast of central Japan. Journal of Geography (Chigaku Zasshi) 122, 308-322. (in
 Japanese with English abstract). doi:10.5026/jgeography.122.308
- Fujiwara, O., Ono, E., Yata, T., Umitsu, M., Sato, Y. and Heyvaert, V.M.A., 2013b. Assessing
 the impact of 1498 Meio earthquake and tsunami along the Enshu-nada coast, central Japan
 using coastal geology. Quaternary International 308, 4–12.
 https://doi.org/10.1016/j.quaint.2012.12.009.
- Fujiwara, O. and Tanigawa, K., 2017. Researches on the tsunami deposits along the Nankai 1358 Trough: Their achievements and upcoming challenges. Journal of the Geological Society 1359 1360 of Japan 123. 831-842. (in Japanese with English abstract). doi.org/10.5575/geosoc.2017.0061 1361
- Furumoto, M. and Ando, M. 2009. Possibility of super large earthquake from southwest Japan
 to Ryukyu Islands*, Chikyu Monthly 31, 197–202, in Japanese.
- Furumura, T., Imai, K. and Maeda, T., 2011. A revised tsunami source model for the 1707 Hoei
 earthquake and simulation of tsunami inundation of Ryujin Lake, Kyushu, Japan. Journal
 of Geophysical Research 116, B02308. doi:10.1029/2010JB007918, 2011
- Garrett, E., Fujiwara, O., Garrett, P., Heyvaert, V.M.A., Shishikura, M., Yokoyama, Y.,
 Hubert-Ferrari, A., Brückner, H., Nakamura, A., De Batist, M., the QuakeRecNankai team,
- 1369 2016. A systematic review of geological evidence for Holocene earthquakes and tsunamis

- 1370 along the Nankai-Suruga Trough, Japan. Earth-Science Reviews 159, 337–357
 1371 https://doi.org/10.1016/j.earscirev.2016.06.011.
- Garrett, E., Fujiwara, O., Riedesel, S., Deforce, W. J., Yokoyama, Y., Schmidt, S., Brückner, H.,
 De Batist, M, Heyvaert, V.M.A., the QuakeRecNankai team., 2018. Historical
 Nankai-Suruga megathrust earthquakes recorded by tsunami and terrestrial mass
 movement deposits on the Shirasuka coastal Lowlands, Shizuoka Prefecture, Japan. The
 Holocene 28, 968–983 https://doi.org/10.1177/0959683617752844.
- Geographical Survey Institute, 1952. Re-survey of the south-western part of Japan after the
 great Nankaido earthquake of 1946.Bulletin of the Geographical Survey Institute 3,
 31-118.
- Geographical Survey Institute, 2008. Crustal Movements in the Tokai District. Report of the
 Coodinating Committee for Earthquake Prediction, Japan 79, 80p. (in Japanese).
- Goldfinger, C., Nelson, C.H., Johnson, J.E., and the Shipboard Scientific Party, 2003a.
 Holocene earthquake records from the Cascadia subduction zone and northern San Andreas
 Fault based on precise dating of offshore turbidites. Annual Review of Earth and Planetary
 Sciences 31, 555-577, doi: 10.1146/annurev.earth.31.100901.141246.
- Goldfinger, C., Nelson, C.H., Johnson, J.E., and the Shipboard Scientific Party, 2003b,
 Deep-water turbidites as Holocene earthquake proxies: the Cascadia subduction zone and
 Northern San Andreas Fault systems. Annals OF Geophysics 46, 1169-1194. DOI:
 https://doi.org/10.4401/ag-3452
- Goto, K., 2017. Paleotsunami researches along the Ryukyu Trench. Journal of the Geological
 Society of Japan 123, 843–855 (in Japanese with English abstract)
 doi.org/10.5575/geosoc.2017.0046
- Goto, K., Chagué-Goff, C., Fujino, S., Goff, J., J., Jaffe, B. Nishimura, Y., Richmond, B.,
 Sugawara, D., Szczuciński, W., Tappin, D.R., Witter, R.C. and Yulianto, E., 2011. New
 insights of tsunami hazard from the 2011 Tohoku-oki event. Marine Geology 290, 46-50.
 doi.org/10.1016/j.margeo.2011.10.004
- Goto, K., Ishizawa, T., Ebina, Y., Imamura, F., Sato, S., Udo, K., submitted. Ten years after the
 2011 Tohoku-oki tsunami: Aftermath of environmental and social effects. Earth-Science
 Reviews.
- Goto, K., Kawana, T. and Imamura, F., 2010a. Historical and geological evidences of boulders
 deposited by tsunamis, southern Ryukyu Islands, Japan. Earth-Science Reviews 102, 77–
 99. doi.org/10.1016/j.earscirev.2010.06.005.
- Goto, K., Miyagi, K., Kawamata, H. and Imamura, F., 2010b. Discrimination of boulders
 deposited by tsunamis and storm waves at Ishigaki Island, Japan. Marine Geology 269, 34–
 45. doi.org/10.1016/j.margeo.2009.12.004.

- Goto, K., Miyagi, K. and Imamura, F., 2013. Localized tsunamigenic earthquakes inferred from
 preferential distribution of coastal boulders on Ryukyu Islands, Japan. Geology 41, 1139–
 1142.
- Goto, K., Okada, K. and Imamura, F., 2009. Characteristics and hydrodynamics of boulders
 transported by storm waves at Kudaka Island, Japan. Marine Geology 262, 14–24.
 doi.org/10.1016/j.margeo.2009.03.001.
- Harada, T., Ishibashi, K., Satake, K., 2013. Tsunami numerical simulation for hypothetical
 giant/great earthquakes along the Izu-Bonin Trench. Programme and Abstracts, the
 Seismological Society of Japan, 2013 Fall Meeting, 186 (in Japanese).
- Harbitz, C. B., Løvholt, F. and Bungum, H., 2013. Submarine landslide tsunamis: how extreme
 and how likely? Natural Hazards 72, 1341-1374. DOI 10.1007/s11069-013-0681-3.
- Hatori, T., 1974. Sources of large tsunamis in southwest Japan. Zishin (Journal of the
 Seismological Society of Japan), 2nd ser., 27, 10-24. (in Japanese with English abstract).
- Hatori, T., 1975. Sources of large tsunamis generated in the Boso, Tokai and Nankai regions in
 1420 1498 and 1605. Bulletin of the Earthquake Research Institute, University of Tokyo 50,
 1421 171-185 (in Japanese with English abstract).
- Hatori, T., 1976. Documents of tsunami and crustal deformation in Tokai district associated
 with the Ansei Earthquake of Dec. 23, 1854. Bulletin of the Earthquake Research Institute,
 University of Tokyo 51, 13–28. (in Japanese with English abstract).
- Hatori, T., 1985. Field investigation of historical tsunamis along the east coast of Kyushu, west
 Japan. Bulletin of the Earthquake Research Institute, University of Tokyo 60, 439–459. (in
 Japanese).
- Heki, K., and Kataoka, T., 2008. On the biannually repeating slow-slip events at the Ryukyu
 Trench, southwestern Japan. Journal of Geophysical Research 113(B11).
 doi:10.1029/2008JB005739.
- HERP (Headhunters for Earthquake Research Promotion), 2004. Long-term evaluation of
 seismic activity around the Hyuga Nada and the Nansei- shoto Trench.
 https://www.jishin.go.jp/main/chousa/kaikou_pdf/hyuganada.pdf (last accessed
 2020/3/12).
- 1435 HERP, 2010. Partial revision of long-term evaluation of Fujikawa Estuary fault zone.
 1436 http://www.jishin.go.jp/main/chousa/10oct_fujikawa/ (In Japanese, last accessed 19 Feb,
 1437 2020.).
- 1438 HERP, 2013. Evaluation of long-term earthquake activity along the Nankai Trough. 2nd edn.,
- 1439 HERP, Japan. https://www.jishin.go.jp/main/chousa/kaikou_pdf/ nankai_2.pdf (in Japanese,
- 1440 last acceded 10 February 2020).

- Hirose, H., and Obara, K., 2005. Repeating short- and long-term slow slip events with deep
 tremor activity around the Bungo channel region, southwest Japan, Earth Planets Space 57,
 961-972.
- Hisamatsu, A., Goto, K. and Imamura, F., 2014. Local paleo-tsunami size evaluation using
 numerical modeling for boulder transport at Ishigaki Island, Japan. Episodes 37, 265–276.
- 1446 DOI: 10.18814/epiiugs/2014/v37i4/006
- Hongo, C., 2010. Sea-level standstill and dominant hermatypic coral from the Holocene raised
 reef terraces at the Kikai Island, Ryukyu Islands. Journal of Geography 119, 860–871. (In
 Japanese with English abstract)
- Hsu, Y. J., Ando, M., Yu, S. B. and Simons, M., 2012. The potential for a great earthquake
 along the southernmost Ryukyu subduction zone. Geophysical Research Letters 39,
 L14302. doi.org/10.1029/2012GL052764
- Hsu, S. K., Yeh, Y. C., Sibuet, J. C., Doo, W. B. and Tsai, C. H., 2013. A mega-splay fault
 system and tsunami hazard in the southern Ryukyu subduction zone. Earth and Planetary
 Science Letters 362, 99–107. doi.org/10.1016/j.epsl.2012.11.053.
- Ichihara, N., Kuroki, A. and Ijyuuin H., 2015. Tsunami event deposits in Miyazaki Plain.
 Abstract of 122nd Annual Meeting of Geological Society of Japan.
 https://doi.org/10.14863/geosocabst.2015.0_188 (in Japanese).
- Igarashi, T., 2010. Spatial changes of inter-plate coupling inferred from sequences of small
 repeating earthquakes in Japan. Geophysical Research Letters 37, L20304. doi:
 10.1029/2010GL044609.
- 1462 Ikeda, M., 1977. Coastal terraces and their deformations in Amami-Oshima, Ryukyu Islands,
 1463 Southern Japan. Journal of Geography (Chigaku Zasshi) 86, 43-49. (in Japanese with
 1464 English abstract).
- 1465 Ikehara, K., Kanamatsu, T., Usami, K. and Hsiung, K.H., 2017. Depositional processes and
- recurrence of seismo-turbidites along the southern Ryukyu Trench. EGU General
- Assembly Conference Abstracts, 2017 EGUGA.19.3451I.
- Imamura, F., Goto, K. and Ohkubo, S., 2008. A numerical model for the transport of a boulder
 by tsunami. Journal of Geophysical Research, Oceans 113, C01008.
 doi.org/10.1029/2007JC004170.
- Ishibashi, K., 1976. Re-examination of a great earthquake expected in the Takaki district,
 central Japan: Possibility of the "Suruga Bay earthquake". Abstract of the Seismological
 Society of Japan, 1976, No.2, 30–34 (In Japanese).
- Ishibashi, K., 1981. Specification of a soon-to-occur seismic faulting in the Tokai district,
 central Japan, based upon seismotectonics. Earthquake Prediction: An international review
 4, Maurice Ewing Series, 297–332. https://doi.org/10.1029/ME004p0297

- Ishibashi, K., 1998. Inferred Tokai earthquake paired with the 1361 Shohei Nankai earthquake.
 Programme and abstracts, the Seismological Society of Japan, 1988 Fall Meeting, 125 (in Japanese). https://historical.seismology.jp/ishibashi/archive/1361Ko-anTokai98.pdf
- Ishibashi, K., 1999. Great Tokai and Nankai, Japan, earthquakes as revealed by historical
 seismology: 1. Review of the events until the mid-14th century. Journal of Geography, 108,
 399–423 (in Japanese with English abstract).
- 1483 Ishibashi, K., 2014. Nankai Trough great earthquake, History, Science and Society. Iwanami
 1484 Shoten, Publishers, 205p (in Japanese).
- Ishibashi, K., 2016. Possibility that the 1099 Jotoku (Kowa) Nankai earthquake is unreal and the
 1096 Kaho (Eicho) earthquake was an "entire-Nankai Trough" earthquake: examination of *Kansenji-an*, a historical document describing an earthquake in Shikoku. Rekishi-jisihin
 (Historical Earthquakes) 31, 81–88(in Japanese with English abstract).
- Ishibashi, K. and Satake, K., 1998. Problems on forcasing great earthquakes in the subduction
 zones arund Japan by means of paleoseismology. Zisin (Journal of the Seismological
 Society of Japan), 2nd ser. 50, 1–21 (in Japanese with English abstract).
- Ishizawa, T., Goto, K., Yokoyama, Y. and Goff, J., 2020. Dating tsunami deposits: Present
 knowledge and challenges. Earth-Science Reviews, 200, 102971.
 doi.org/10.1016/j.earscirev.2019.102971.
- Ishiuka, O., Taylor, R.N., Yuasa, M. and Ohara, Y., 2011., Making and breaking an island arc:
 A new perspective from the Oligocene Kyushu Palau arc, Philippine Sea. Geochemistry,
 Geophysics, Geosystems 12, https://doi.org/10.1029/2010GC00344
- Ito, T., and Hashimoto, M, 2004. Spatiotemporal distribution of interplate coupling in southwest
 Japan from inversion of geodetic data. Journal of Geophysical Research 109, B02315,
 doi:10.1029/2002JB002358.
- Iwai, K. and Kawana, T., 2008. Physiographical studies on locations of archaeological sites in
 Okinawa Island, the Ryukyus, Japan, during past several thousand years. Okinawa J. Geog.
 Stud. 8, 25–36. (in Japanese).
- Japan coast guard, 1948. Research report of the 1946 Nankai earthquake -part of ground
 deformation and damage-. Hydrographic bulletin 201, 139 p. (in Japanese)
- Kanamori H, McNally K., 1982. Variable rupture mode of the subduction zone along the
 Ecuador-Colombia coast. Bulletin of Seismological Society of America 72, 1241–1253.
- 1508 Kasahara, K., 1975. Aseismic Faulting Following the 1973 Nemuro-oki Earthquake, Hokkaido,
- Japan (a Possibility), Pure and applied Geophysics 113, 127–139.
 doi:10.1007/BF01592905

- Kasahara, K. and Kato, T. (1980) Aseismic faulting following the 1973 Nemuro-oki earthquake,
 Hokkaido, Japan (a retrospective study). Pure and Applied Geophysics 119, 392-403. Doi:
 10.1007/BF00877774.
- 1514 Kato, Y., 1993. Geological structure and topography of the Amami Plateau. Report of
 1515 Hydrographic Researches 29, 51-63 (in Japanese with English abstract).
- 1516 Kato, Y. and Kimura, M., 1983. Age and origin of so-called "tsunami-ishi", Ishigaki Island,
 1517 Okinawa Prefecture. Journal of the Geological Society of Japan 89, 471–474. (in
 1518 Japanese).
- 1519 Kawamura, K., Kanamatsu, T. and Yamada, Y., 2017. Submarine slides and marine geohazards:
 1520 the study results so far and current problems. Journal of Geological Society of Japan 123,
 1521 999-1014. (in Japanese with English abstract).
- Kawana, T., 1989. Quaternary crustal movement in the Ryukyu Islands. Earth Mon. 11, 618–
 630 (in Japanese, original title translated).
- Kawana, T., 2006. Invasion of about 3400 cal BP large wave in the southeastern Okinawa
 Island and the surroundings, the Ryukyus, Japan, as deduced from coralline deposits. Bull.
 Faculty Educ. Univ. Ryukyus 68, 265–271. (in Japanese).
- Kawana, T. and Nakata, T., 1994. Timing of late Holocene tsunamis originated around the
 southern Ryukyu Islands, Japan, deduced from coralline tsunami deposits. Journal of
 Geography (Chigaku Zasshi) 103, 352–376. (in Japanese with English abstract).
- Kawana, T. and Nakata, K., 2003. Calibrated calendar ages of the paleotsunamis attacked
 eastern part of the Kasari Peninsula of Amami Island and the northern part of Kikai Island,
 the north Ryukyus, Japan. Okinawa Geog. Soc. Bull. 6, 33–40. (in Japanese).
- Kitamura, A., 2016. Examination of the largest-possible tsunamis (Level 2) generated along the
 Nankai and Suruga troughs during the past 4000 years based on studies of tsunami deposits
 from the 2011 Tohoku-oki tsunami. Progress in Earth and Planetary Science 3:12. DOI
 10.1186/s40645-016-0092-7.
- Kitamura, A., Fujiwara, O., Shinohara, K., Akaike, S., Masuda, T., Ogura, K., Urano, Y.,
 Kobayashi, K., Tamaki, C. and Mori, H., 2013. Identifying possible tsunami deposits on
 the Shizuoka Plain, Japan and their correlation with earthquake activity over the past 4000
 years. The Holocene 23, 1684–1698. doi.org/10.1177/0959683613505345.
- 1541 Kitamura, A., Ina, T., Suzuki, D., Tsutahara, K., Sugawara, D., Yamada, K. and Aoshima, A.,
 1542 2019. Geologic evidence for coseismic uplift at ~AD 400 in coastal lowland deposits on
 1543 the Shimizu Plain, central Japan. Progress in Earth and Planetary Science 6,
 1544 Article number 57. Doi:10.1186/s40645-019-0305-y
- Kitamura, A., Ito, M., Sakai, S., Yokoyama, Y. and Miyairi, Y., 2018a. Identification of tsunami
 deposits using a combination of radiometric dating and oxygen-isotope profiles of

 1547
 articulated
 bivalves.
 Marine
 Geology
 403,
 57–61.

 1548
 https://doi.org/10.1016/j.margeo.2018.04.003.
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 6
 <t

Kitamura, A., Ito, M., Ikuta, R. and Ikeda, M., 2018b. Using molluscan assemblages from
paleotsunami deposits to evaluate the influence of topography on the magnitude of late
Holocene mega-tsunamis on Ishigaki Island, Japan. Prog. Earth Planet. Sci. 5, 41. DOI:
10.1186/s40645-018-0200-y.

- 1553 Kitamura, A. and Kobayashi, K., 2014a. Geologic record of middle-late Holocene
 1554 paleotsunamis and paleo-earthquakes on the Shizuoka Plain and coastal lowland of the
 1555 southern Izu Peninsula, central Japan. Journal of Geography (Chigaku Zasshi) 122, 813–
 1556 834 (in Japanese with English abstract).
- Kitamura, A. and Kobayashi, K., 2014b. Geologic evidence for prehistoric tsunamis and 1557 coseismic uplift during the AD 1854 Ansei-Tokai earthquake in Holocene sediments on the 1558 Shimizu Plain. 24. 814-827. 1559 central Japan. The Holocene Doi[.] 10.1177/0959683614530447 1560
- Kitamura, A., Ohashi, Y., Miyairi, Y., Yokoyama, Y. and Yamaguchi, T., 2014. Discovery of a
 tsunami boulder along the coast of Shimoda, Shizuoka, central Japan. The Quaternary
 Research (Daiyonki-Kenkyu) 53, 259–264 (In Japanese with English abstract).
- Kitamura, A., Seki, Y., Kitamura, Y., Haga, T., 2018c. The discovery of emerged boring
 bivalves at Cape Omaezaki, Shizuoka, Japan: Evidence for the 1361 CE Tokai earthquake
 along the Nankai Trough. Marine Geology 405, 114–119.

1567 https://doi.org/10.1016/j.margeo.2018.08.006

- Koike, K. and Machida, H. eds., 2001. Atlas of Quaternary marine terraces in the Japaneseislands. University of Tokyo Press, 105pp. with 3 CD-ROMs and 2 maps. (in Japanese).
- Komatsubara, J. and Fujiwara, O. (2007) Overview of Holocene tsunami deposits along the
 Nankai, Suruga, and Sagami Troughs, southwest Japan. Pure and Applied Geophysics, 164,
 493-507.
- 1573 Komatsubara, J. Fujiwara, O., Takada, K., Sawai, Y., Than T. Aung and Kamataki, T. (2008)
 1574 Historical tsunamis and storms recorded in a coastal lowland, Shizuoka prefecture, along
 1575 the Pacific Coast of Japan. *Sedimentology*, 55, 1703-1716.
- 1576 Koyama, M., 1999. A review of historical seismology in Japan: analyses and proposals for next
- stage of study of historical documents. Gournal of Geography 108, 346-369. (in Japanese
 with English abstract) https://doi.org/10.5026/jgeography.108.4 346
- 1579 Kugai, M., 2011. Discovery of the foraminifera sedimentation at "Tomori-motojima" and
 1580 "Sunagawa-motojima"*. Bull. Miyakojima City Mus.*, 15, 65–75. (in Japanese).

- 1581 Kumagai, H., 1996. Time sequence and the recurrence models for large earthquakes along the
 1582 Nankai Trough revisited. Geophysical Research Letters, 23, 1139-1142.
 1583 https://doi.org/10.1029/96GL01037
- Lay, T., and Kanamori, H., 1981. An asperity model of great earthquake sequences, in
 Earthquake prediction An international review, AGU Geophys. Mono., edited by D. W.
 Simpson and P. G. Richards, pp. 579-592, Amer. Geophys. Union, Washington, D.C.
- Lougheed, B.C., Obrochta, S.P., 2016. MatCal: open source bayesian ¹⁴C age calibration in
 MatLab. J. Open Res. Software 4. <u>http://doi.org/10.5334/jors.130</u>.
- Lougheed, B.C. and Obrochta, S.P., 2019. A Rapid, deterministic age-depth modelling routine
 for geological sequences with inherent depth uncertainty: Paleoceanography and
 Paleoclimatology 34, 122-133. doi.org/10.1029/2018PA003457
- Loveless, J. P., and Meade, B. J., 2010. Geodetic imaging of plate motions, slip rates, and
 partitioning of deformation in Japan. Journal of Geophysical Research 115, B02410.
 doi.org/10.1029/2008JB006248
- Loveless, J. P., and Meade, B. J., 2011. Spatial correlation of interseismic coupling and
 coseismic rupture extent of the 2011 Mw=9.0 Tohoku-oki earthquake. Geophysical
 Research Letters 38, L17306, doi:10.1029/2011GL048561.
- 1598 Makino, K., 1968. Meiwa tsunami at Yaeyama*. Published by author, 447p. (in Japanese).
- Makino, K., 1981. Meiwa tsunami at Yaeyama -revision and enlargement-*. Published byauthor, 462p. (in Japanese).
- Malik, J.N., Shishikura, M., Echigo, T., Ikeda, Y., Satake, K., Kayane, H., Sawai, Y., Murty,
 C.V.R. and Dikshit, O. 2011. Geologic evidence for two pre-2004 earthquakes during
 recent centuries near Port Blair, South Andaman Island, India. Geology 39, 559-562. DOI:
 10.1130/G31707.1
- Matsu'ura, R. S., in press, The cause of the tsunami in 1791 at the main island of Okinawa.
 Historical Earthquakes (in Japanese).
- Miyazaki, S. and Heki, K., 2001. Crustal velocity field of southwest Japan: Subduction and
 arc-arc collision. Journal of Geophysical Research 106(B3), 4305-4326.
 doi.org/10.1029/2000JB900312.
- Minamidate, K., Goto, K., Watanabe, M., Roeber, V., Toguchi, K., Sannoh, M., Nakashima, Y.,
 Kan, H., 2020. Millennial scale maximum intensities of typhoon and storm wave in the
 northwestern Pacific Ocean inferred from storm deposited reef boulders. Scientific reports.
 10, 7218. https://doi.org/10.1038/s41598-020-64100-6.
- 1013 10, 7218. https://doi.org/10.1038/841398-020-04100-0.
- Miyazaki, S., Segall, P., McGuire, J.J., Kato, T. and Hatanaka, Y., 2006. Spatial and temporal
 evolution of stress and slip rate during the 2000 Tokai slow earthquake. Journal of
 Geophysical Research 111, B03409.doi:10.1029/2004JB003426.

- Miyazawa, K., Goto K., and Imamura, F., 2012. Re-evaluation of the 1771 Meiwa Tsunami 1617 source model, southern Ryukyu Islands, Japan. In Submarine Mass Movements and Their 1618 Hazards 1619 Consequences, Adv. Nat. Tech. Res. 31, 497-506. DOI: 10.1007/978-94-007-2162-3 44. 1620
- 1621 Moore, G.F., Bangs, N.L., Taira, A., Mumamoto, S., Pangborn, E. and Tobin, H.J., 2007.
- Three-dimensional splay fault geometry and implication for tsunami generation. Science
 318, 1128-1131. DOI: 10.1126/science.1147195
- Moriki, H., Kumamoto, T., Nakata, T., Goto, H., Izumi, N. and Nishizawa, A., 2017.
 Identification of landslide and its characteristics on the seafloor around Japan using
 anaglyph images. Report of Hydrographic and Oceanographic Researches No.54, 1-16. (in
 Japanese with English abstract).
- Murakami, H., Shimada, T., Itoh, S., Yamamoto, N. and Ishizuka, J., 1996. Reexamination of
 the heights of the 1605, 1707 and 1854Nankai tsunamis along the coast of Shikoku island.
 Journal of Japan Society for Natural Disaster Science 15, 39-52 (In Japanese with English
 abstract)
- Nadeau, R. M. and Johnson, L. R., 1998. Seismological studies at Parkfield VI: Moment release
 rates and estimates of source parameters for small repeating earthquakes. Bulletin of
 Seismological Society of America 88(3), 790-814.
- Nagaoka, S., Maemoku, H. and Matsushima, Y., 1991. Evolution of Holocene coastal landforms
 in the Miyazaki Plain, southern Japan. The Quaternary Reach (Daiyonki Kenkyu) 30,
 59-78 (in Japanese with Englishabstract).
- Nakamura, M., 2009. Fault model of the 1771 Yaeyama earthquake along the Ryukyu Trench
 estimated from the devastating tsunami. Geophysical Research Letters 36, L19307,
 doi:10.1029/2009GL039730.
- 1641 Nakamura, M., 2017. Distribution of low ☐ frequency earthquakes accompanying the very low
 1642 frequency earthquakes along the Ryukyu Trench. Earth Planets. Space 69, 49, DOI
 1643 10.1186/s40623-017-0632-4.
- Nakamura, M. and Kinjou, A., 2013. Fault model of the 1768 and 1791 tsunamis in the central
 Ryukyu. Abstract for the 2013 AGU Fall Meeting, T13E-2574.
- Nakamura, M. and Sunagawa, N., 2015. Activation of very low frequency earthquakes by slow
 slip events in the Ryukyu Trench. Geophysical Research Letters 42, 1076–1082.
 https://doi.org/10.1002/2014GL062929.
- Nakashima, R., Hori, N., Miyazaki, K. and Nishioka, Y., 2008. Geology of the Toyohashi and
 Tahara districts. Quadrangle Series, 1: 50,000, Geological Survey of Japan, AIST, 113 p. (in
 Japanese with English abstract 3 p.)

- Nakata, T. and Shimazaki, K., 1997. Geo-slicer, a newly invented soil sampler, for
 high-resolution active fault studies. Journal of Geography, 106, 59-69. (in Japanese with
 English abstract). doi.org/10.5026/jgeography.106.59.
- Nakata, T., Takahashi, T. and Koba, M., 1978. Holocene-emerged coral reefs and sea-level
 changes in the Ryukyu Islands. Geographical Review of Japan 51, 87-108. (in Japanese
 with English abstract).
- 1658 Nakaza, H., 2017. What is Shiraho-Saonetabaru Cave Site? Outline of survey and main results.
 1659 Kagaku 87, 532–536. (in Japanese).
- Nakaza, E., Iribe, T., Tokuhisa, R., Miyazato, N., Inagaki, K. and Savou, R., 2013. Prehistorical
 and historical tsunamis of Ryukyu Islands estimated through tsunami depositions. Journal
 of the Japan Society of Civil Engineering, Ser. B3 (Ocean Eng.) 69, 515–520. (in Japanese
 with English abstract).
- Nakaza, E., Tokuyama, R., Inagaki, K., 2015. Original locations of huge boulders moved by 1664 tsunami and its generation mechanism. Journal of the Japan Society of Civil Engineers, Ser. 1665 193-198. B2 (Coastal Eng.), 71, (in Japanese with English abstr). 1666 doi.org/10.2208/kaigan.71.I 193 1667
- Nishimura, T., 2014. Short-term slow slip events along the Ryukyu Trench, southwestern Japan,
 observed by continuous GNSS. Progress in Earth and Planetary Science 1, 22.
 http://www.progearthplanetsci.com/content/1/1/22.
- Namegaya, Y., Maemoku, H., Shishikura, M., Echigo, T. and Nagai, A., 2011. Factors causing
 scattered boulders located around Hashigui-iwa, the southernmost of Kii peninsula, Japan.
 Abstract of Japan Geoscience Union Meeting 2011, AAA035-12 (in Japanese).
- Namegaya, Y., Satake, K. and Yamaki, S., 2010. Numerical simulation of the AD 869 Jogan
 tsunami in Ishinomaki and Sendai plains and Ukedo river-mouth lowland. Annual Report
 on Active Fault and Paleoearthquake Researches, Geological Survey of Japan, AIST, No.
 10, 1-21 (in Japanese with English abstract).
- Nanayama, F., Satake, K., Furukawa, R., Shimokawa, K., Atwater, B.F., Shigeno, S., Yamaki,
 S., 2003. Unusually large earthquakes inferred from tsunami deposits along the Kuril
 trench. Nature 424, 660–663, DOI: 10.1038/nature01864.
- Nishimura, T., 2014. Short-term slow slip events along the Ryukyu Trench, southwestern Japan,
 observed by continuous GNSS. Progress in Earth and Planetary Science 1, 22,
 doi:10.1186/s40645-014-0022-5.
- Nishimura, S. and Hashimoto, M., 2006. A model with rigid rotations and slip deficits for the
 GPS-derived velocity field in Southwest Japan. Tectonophysics 421, 187-207.
 doi:10.1016/j.tecto.2006.04.017.

- 1687 Nishimura, S., Hashimoto, M. and Ando, M., 2004. A rigid block rotation model for the GPS
 1688 derived velocity field along the Ryukyu arc. Physics of the Earth and Planetary Interiors
 1689 142(3-4), 185-203, doi:10.1016/j.pepi.2003.12.014.
- Nishimura, T., Miura, S., Tachibana, K., Hashimoto, K., Sato, T., Hori, S., Murakam, E., Kono,
 T., Nida, K., Mishina, M., Hirasawa, T. and Miyazaki, S., 2000. Distribution of seismic
 coupling on the subducting plate boundary in northeastern Japan inferred from GPS
 observations. Tectonophysics 323, 217-238. doi.org/10.1016/S0040-1951(00)00108-6.
- Obara, K., 2002. Nonvolcanic deep tremor associated with subduction in southwest Japan.
 Science, 296(5573), 1679-1681. 10.1126/science.1070378.
- 1696 Obara, K., and Kato, A., 2016. Connecting slow earthquakes to huge earthquakes. Science,
 1697 353(6296), 253-257, doi:10.1126/science.aaf1512.
- Obrocki, L., Vött, A., Wilken, D., Fischer, P., Willershäuser, T., Koster, B., Lang, F.,
 Papanikolaou, I., Rabbel, W. and Reicherter, K., 2020. Tracing tsunami signatures of the
 AD 551 and AD 1303 tsunamis at the Gulf of Kyparissia (Peloponnese, Greece) using direct
 push *in situ* sensing techniques combined with geophysical studies. Sedimentology 67,
 1274-1308. doi: 10.1111/sed.12555.
- Okahashi, H., Yasuhara, M., Mitamura, M., Hirose, K., Yoshikawa, S., 2005. Event deposits
 associated with tsunamis and their sedimentary structure in Holocene marsh deposits on
 the east coast of the Shima Peninsula, central Japan. J. Geosci. Osaka City Univ. 48,
 143-158.
- Okumura, K., 1996. Tephrochronology, correlation and deformation of marine terraces in
 eastern Hokkaido, Japan. Geographical Reports of Tokyo Metropolitan University 31,
 19-26.
- Okamura, M. and Matsuoka, H., 2012. Nankai earthquake recurrences from tsunami sediment.
 Kagaku, 82, 182-191. (in Japanese)
- Okamura, Y., Nishizawa, A., Fujii, Y. and Yanagisawa, H., 2018. Accretionary prism collapse:
 a new hypothesis on the source of the 1771 giant tsunami in the Ryukyu Arc, SW Japan.
 Scientific Reports 8, 13620. DOI:10.1038/s41598-018-31956-8.
- Omoto, K., 2012. Distribution of coral boulders beached by huge tsunamis and calibrated
 radiocarbon ages of fossil Porites sp. Of Maibahabama, SE of Miyako Island, Okinawa
 Prefecture. Journal of Geography (Chigaku Zasshi) 121, 1043–1051. (in Japanese with
 English abstract).
- Omoto, K., 2019. ¹⁴C ages of large coral boulders washed ashore on the southeast coast of
 Miyako Island, Okinawa Prefecture. Quarterly Journal of Geography 71, 82-88. (In
 Japanese with English abstract)

- Ota, Y. and Omura, A., 2000. Review of coral reef terrace studies at Kikai Island, Ryukyu
 Islands, southwestern Japan. The Quaternary Research (Daiyonki-Kenkyu) 39, 45–53 (in
 Japanese with English abstract).
- Ozawa, S., Hashimoto, M. and Tada, T., 1997. Vertical crustal movements in the coastal areas
 of Japan estimated from tidal observations. Bull. Geog. Surv. Inst. 43, 1-21.
- Ozawa, S., Murakami, M., Kaidzu, M., Tada, T., Sagiya, T., Hatanaka, Y., Yarai, H. and
 Nishimura T., 2002. Detection and monitoring of ongoing aseismic slip in the Tokai region,
 central Japan. Science 298, 1009–1012. doi:10.1126/science.1076780.
- Paris, R, Falvard, S., Chagué, C., Goff, J., Etienne, S. and Doumalin, P., 2020. Sedimentary
 fabric characterized by X-ray tomography: A case-study from tsunami deposits on the
 Marquesas Islands, French Polynesia. Sedimentology 67, 1207-1229. doi:
 10.1111/sed.12582.
- Peters, R., Jaffe, B. and Gelfenbaum, G., 2007. Distribution and sedimentary characteristics of
 tsunami deposits along the Cascadia margin of western North America. Sedimentary
 Geology, 200, 372-386. doi.org/10.1016/j.sedgeo.2007.01.015
- Priest, G.R., Witter, R.C., Zhang, Y.J., Goldfinger, C., Wang, K., Allan, J.C., 2017, New constraints on coseismic slip during southern Cascadia subduction zone earthquakes over the past 4,600 years implied by tsunami deposits and marine turbidites, Natural Hazards DOI: 10.1007/s11069-017-2864-9.
- Riedesel, S., Brill, D., Roberts, H.M., Duller, G.A.T., Garrett, E., Zander, A.M., King, G.E.
 Tamura, T., Burow, C., Cunningham, A., Seeliger, M., De Batist, M., Heyvaert, V.M. A.,
 Fujiwara, O., Brückner, H., the QuakeRecNankai team, 2018. Single-grain feldspar
 luminescence chronology of historical extreme wave event deposits recorded in a coastal
 lowland, Pacific coast of central Japan. Quaternary Geochronology 45, 37–49.
 https://doi.org/10.1016/j.quageo.2018.01.006.
- 1747 Ryan, W.B.F., S.M. Carbotte, J.O. Coplan, S. O'Hara, A. Melkonian, R. Arko, R.A. Weissel, V.
 1748 Ferrini, A. Goodwillie, F. Nitsche, J. Bonczkowski, and R. Zemsky (2009), Global
 1749 Multi-Resolution Topography synthesis, Geochem. Geophys. Geosyst., 10, Q03014,
 1750 doi:10.1029/2008GC002332.
- Sagiya, T., and Thatcher, W., 1999. Coseismic slip resolution along a plate boundary
 megathrust: The Nankai Trough, southwest Japan. Journal of Geophysical Research
 104(B1), 1111-1129. doi.org/10.1029/98JB02644.
- Saito, E., 2017. GIS data of the depth of the oceanic plate surfaces beneath the Japanese
 archipelago. GSJ Open-File Report, no. 647, Geological Survey of Japan, AIST. (in
 Japanese).

- Sangawa, A., 1992. Earthquake archaeology: Earthquake history revealed from archaeological
 site). Chuōkōron-shin sha, 251p. (in Japanese)*.
- Sangawa, A., 2001. Recent results of paleoseismological study based on earthquake traces
 excavated at archaeological sites. Annual Report on Active Fault and Paleoearthquake
 Researches, Geological Survey of Japan, AIST, No. 1, 287–300 (in Japanese with English
 abstract).
- Sangawa, A., 2007. Japanese History Looking Back from Large Earthquakes. Chu-ko Shinsho,
 268pp (in Japanese) *.
- Sateke, K. and Atwater, B.F., 2007. Long-term perspectives on giant earthquakes and tsunamis
 at subduction zones. Annual Review of Earth and Planetary Sciences 35, 349-374.
 https://doi.org/10.1146/annurev.earth.35.031306.140302.
- Satake, K., Namegaya, Y. and Yamaki, S., 2008. Numerical simulation of the AD 869 Jogan
 tsunami in Ishinomaki and Sendai plains. Annual Report on Active Fault and
 Paleoearthquake Researches, Geological Survey of Japan, AIST, No. 8, 71-89 (in Japanese
 with English abstract).
- Sato, Y. and Fujiwara, O., 2017. Microfossil evidence for recurrent coseismic subsidence
 around Lake Hamana, near the Nankai-Suruga trough, central Japan. Quaternary
- 1774
 International 456, 39-52. http://dx.doi.org/10.1016/j.quaint.2017.08.040
- Sato, Y., Fujiwara, O. and Ono, E., 2016a. Late Holocene geomorphological development of
 beach ridges in western Hamamatsu strand plain, central Japan. The Quaternary Research
 55, 17-35 (in Japanese with English abstract)
- Sato, Y., Fujiwara, O., Ono, E., Umitsu, M., 2011. Environmental change in coastal lowlands
 around the Lake Hamana during the middle to late Holocene. Geographical Review of
 Japan Series A, 84, 258-273. (in Japanese with English abstract).
- Sato, Y., Matsuoka, H., Okamura, M., Kashima, K., 2016b. Late Holocene environmental
 changes of coastal lagoon inferred from a fossil diatom analysis of sediment core from
 Lake Hamana, central Japan. Quat. Int. 397, 317–329.
- Savage, J. C., 1983. A dislocation model of strain accumulation and release at asubduction zone.
 Journal of Geophysical Research 88(Nb6), 4984-4996, doi:10.1029/JB088iB06p04984.
- Sawai, Y., in press. Subduction zone paleoseismology along the Pacific coast of northeast Japan
 progress and remaining problems. Earth-Science Reviews.
- Sawai, Y., Namegaya, Y., Okamura, Y., Satake, K. and Shishikura, M., 2012, Challenges of
 anticipating the 2011 Tohoku earthquake and tsunami using coastal geology. Geophysical
 Research Letters, **39**, doi: 10.1029/2012GL053692.
- Sawai, Y., Satake, K., Kamataki, T., Nasu, H., Shishikura, M., Atwater, BF., Horton, B. P.,
 Kelsey, H.M., Nagumo, T. and Yamaguchi, M., 2004. Transient uplift after a 17th-century

- earthquake along the Kuril subduction zone. Science 306, 1918-1920. DOI:
 10.1126/science.1104895
- Sawai, Y., Tanigawa, K., Tamura, T. and Namegaya, Y., 2016. Medieval coastal inundation
 revealed by a sand layer on the Ita lowland adjacent to the Suruga Trough, central Japan.
 Natural Hazards 80, 505–519. DOI 10.1007/s11069-015-1980-7.
- Seno, T., 2012. Great earthquakes along the Nankai Trough A new idea for their rupture mode
 and time series–. Zisin (Journal of the Seismological Society of Japan), 2nd ser. 64, 97–
 116 (in Japanese with English abstract).
- 1801 Shikakura, Y., 2014. Marine terraces caused by fast steady uplift and small coseismic uplift and
 1802 the time-predictable model: Case of Kikai Island, Ryukyu Islands, Japan. Earth Planet. Sci.
 1803 Lett., 404, 232–237. https://doi.org/10.1016/j.epsl.2014.08.003
- 1804 Shimada Y, Fujino S, Sawai Y, Koichiro T, Matsumoto D, Momohara A,
 1805 Saito-Kato M, Yamada M, Hirayama E, Suzuki T, Chagué C., 2019.
 1806 Geological record of prehistoric tsunamis in Mugi Town, facing the Nankai
 1807 Trough, western Japan. Progress in Earth and Planetary Science, 6 (33).
 1808 https://doi.org/10.1186/s40645-019-0279-9.
- Shimazaki, K., 1974a. Pre-seismic crustal deformation caused by an underthrusting oceanic
 plate, in eastern Hokkaido, Japan. Physics of the Earth and Planetary Interiors 8, 148-157
 https://doi.org/10.1016/0031-9201(74)90127-7.
- 1812 Shimazaki, K., 1974b. Nemuro-Oki earthquake of June 17, 1973: A lithospheric rebound at the
 1813 upper half of the interface. Physics of the Earth and Planetary Interiors, 9, 314-327.
 1814 https://doi.org/10.1016/0031-9201(74)90059-4.
- Shimazaki, K. and Nakata, T., 1980. Time predictable recurrence model for large earthquakes.
 Geophysical Research Letters, 7, 279-282. https://doi.org/10.1029/GL007i004p00279
- Shinozaki, T., Fujino, S., Ikehara, M. Sawai, Y., Tamura, T., Goto, K., Sugawara, D. and Abe,
 T., 2015. Marine biomarkers deposited on coastal land by the 2011 Tohoku-oki tsunami.

1819 Natural Hazards 77, 445–460. doi.org/10.1007/s11069-015-1598-9.

- 1820 Shishikura, M., 2013. Earthquake and tsunamis along the Nankai Trough, inferred from geology
 1821 and geomorphology -examples in Nankai region-. GSJ Chishitsu News, 2, 201-204. (in
 1822 Japanese)
- Shishikura, M., Echigo, T., Maemoku, H., Ishiyama, T., 2008a. Height and ages of uplifted
 sessile assemblage distributed along the southern coast of the Kii Peninsula, southcentral
 Japan reconstruction of multi-segment earthquake history along the Nankai Trough.
- 1826 Annual Report on Active Fault and Paleoearthquake Researches, Geological Survey of
 1827 Japan, AIST, No. 8, 267–280 (in Japanese with English abstract).

- Sugihara, K., Nakamori, T., Iryu, Y., Sasaki, K. and Blanchon, P., 2003. Holocene sea-level
 change and tectonic uplift deduced from raised reef terraces, Kikai-jima, Ryukyu Islands,
 Japan. Sedimentary Geology 159, 5-25. doi.org/10.1016/S0037-0738(03)00092-7.
- Sugiyama, Y., 1990. Seismotectonics of the Suruga Bay and Enshu-nada region. Zishin (Journal
 of the Seismological Society of Japan), 2nd ser., 43, 439–442. (in Japanese)
- Sugiyama, Y., 1991. The Middle Pleistocene deposits in the Atsumi Peninsula and along the
 east coast of Lake Hamana, Tokai district -sedimentary cycles formed by the
- glacio-eustatic sea-level change and their correlations to the contemporaneous deposits in
 the Kanto and Kinki districts-. Bulletin of the Geological Survey of Japan 42, 75-109. (in
 Japanese with English abstract).
- Suito, H. and Ozawa, S., 2009. Transient crustal deformation in the Tokai disrict The Tokai
 slow slip event and post seismic deformation caused by the 2004 off southeast Kii
 Peninsula earthquake —. Zishin (Journal of the Seismological Society of Japan), 2nd ser.,
 61, 113-135 (in Japanese with English abstract).
- Tadokoro, K., Nakamura, M., Ando, M., Kimura, H., Watanabe, T. and Matsuhiro, K., 2018.
 Interplate coupling state at the Nansei-Shoto (Ryukyu) Trench, Japan, deduced from seafloor crustal deformation measurements. Geophysical Research Letters 45, 6869–6877.
 https://doi.org/10.1029/2018GL078655.
- Takada, K., Satake, K., Sangawa, A., Shimokawa, K., Kumagai, H., Goto, K., Haraguchi, T.,
 2002. Survey of tsunami deposits at an archaeological site along the eastern Nankai trough.
 Annual report on active fault and paleoearthquake researches, Geological Survey of Japan,
 AIST, No. 2, 235-243. (In Japanese with English abstract)
- Tani, S., Kitagawa, H., Hong, W., Park, J.H. Sung, K.S. and Park G., 2013. Age determination
 of the Kawagodaira volcanic eruption in Japan by ¹⁴C wiggle-matching. Radiocarbon 55,
 748-752. DOI: 10.2458/azu_js_rc.55.16312.
- Tanigawa, K., Shishikura, M., Fujiwara, O., Namegaya, Y., and Matsumoto, D., 2018. Mid-to
 late-Holocene marine inundations inferred from coastal deposits facing the Nankai Trough
 in Nankoku, Kochi Prefecture, southern Japan. The Holocene 28, 867–878
 ttps://doi.org/10.1177/0959683617752837.
- Tanioka, Y., Satake, K. 2001a. Detailed coseismic slip distribution of the 1944 Tonankai
 earthquake estimated from tsunami waveforms. Geophys. Res. Lett. 28, 1075–1078.
 doi:10.1029/2000GL012284.
- Tanioka, Y., Satake, K., 2001b. Coseismic slip distribution of the 1946 Nankai earthquake and
 aseismic slips caused by the earthquake. Earth, Planets and Space 53, 235–241.
 https://doi.org/10.1186/BF03352380.

- Tappin, D. R., Watts, P. and Grilli, S. T., 2008. The Papua New Guinea tsunami of 17 July
 1998: anatomy of a catastrophic event. Nat. Hazards Earth Syst. Sci. 8, 243–266.
- 1865 The research group for active submarine faults off Tokai(ed.), 1999. active submarine faults off
- Tokai -Results from the Japan-France KAIKO-TokaiProject. University of Tokyo Pres,
 1867 151p. (In Japanese with English abstract)
- Tsuji, Y., 1997 Damage to villages on Amami Archipelago due to the 1995
 Amami-Oshima-Kinkai earthquake-tsunami. Journal of Geography (Chigaku Zasshi) 106,
 486–502 (in Japanese with English abstract).
- Tsuji, Y., Okamura, M., Matsuoka, H., Goto, T., Han, S.S., 2002. Prehistorical and historical
 tsunami traces in lake floor deposits, Oike Lake, Owase City and Suwaike Lake,
 Kii-Nagashima City, Mie Prefecture, central Japan. Chikyu Mon. 24, 743–747 (In
 Japanese)
- Tsuji, Y., Okamura, M., Matsuoka, H., Murakami, Y., 1998. Study of tsunami traces in lake
 floor sediment of the Lake Hamanako. Hist. Earthq. 14, 101–113 (In Japanese)
- Uchida, N., Iinuma, T., Nadeau, R. M., Burgmann, R. and Hino, R., 2016. Periodic slow slip
 triggers megathrust zone earthquakes in northeastern Japan. Science, 351(6272), 488-492,
 doi:10.1126/science.aad3108.
- Ujiie, H., Nakamura, T., Miyamoto, Y., Park, J.O., Hyun, S., and Oyakawa, T., 1997.
 Holocene turbidite cores from the southern Ryukyu Trench slope: suggestions of periodic
 earthquakes. The Journal of the Geological Society of Japan 103, 590-603.
 https://doi.org/10.5575/geosoc.103.590.
- Wallace, L. M., Ellis, S., Miyao, K., Miura, S., Beavan, J. and Goto, J., 2009. Enigmatic, highly
 active left-lateral shear zone in southwest Japan explained by aseismic ridge collision.
 Geology, 37, 143-146, doi:10.1130/G25221a.1.
- Wallis, S.R., Fujiwara, O. and Goto, K. (2018) Geological studies in tsunami research since the
 2011 Tohoku earthquake. Geological Society London Spec. Publication, 456, 39-53.
- Watanabe, H., 1998. Comprehensive list of tsunamis to hit the Japanese Islands. Tokyo: Tokyo
 University Press, 238 pp (in Japanese).
- Watanabe, M., Goto, K., Imamura, F. and Hongo, C., 2016. Numerical identification of tsunami
 boulders and estimation of local tsunami size at Ibaruma reef of Ishigaki Island, Japan.
 Island Arc 25, 316–332. doi.org/10.1111/iar.12115.
- Watanabe, M., Goto, K., Bricker, J. D. and Imamura, F., 2018. Are inundation limit and
 maximum extent of sand useful for differentiating tsunamis and storms? An example from
 sediment transport simulations on the Sendai Plain, Japan. Sedimentary Geology, 364,
 204-216. doi.org/10.1016/j.sedgeo.2017.12.026.

- Webster, J. M., Davies P. J. and Konishi, K., 1998. Model of fringing reef development in
 response to progressive sea level fall over the last 7000 years (Kikai-jima, Ryukyu
 Islands, Japan). Coral Reefs 17, 289–308.
- Wells R. E., Blakely, R. J., Sugiyama, Y., Scholl, D. W. and Dinterman, P. A., 2003, Basin–
 centered asperities in great subduction zone earthquakes: A link between slip, subsidence,
 and subduction erosion? J. Geophys. Res., 108, doi: 10.1029/2002JB002072.
- Yamada, M., Fujino, S., Chiba, T., Goto, K. and Goff, J., 2020. Redeposition of volcaniclastic
 sediments by a tsunami 4600 years ago at Kushima City, south-eastern Kyushu, Japan.
 Sedimentology. doi.org/10.1111/sed.12547.
- Yamamoto, M., 2008. Trace of earthquake and tsunami in Ishigaki Island, Yaeyama
 Archipelago. Archaeol. J. 577, 9–14. (in Japanese).
- Yamamoto, Y., Obana, K., Takahashi, T., Nakanishi, A., Kodaira, S. and Kaneda, Y., 2013.
 Imaging of the subducted Kyushu-Palau Ridge in the Hyuga-nada region, western Nankai
 Trough subduction zone. Tectonophysics 589, 90-102, doi:10.1016/j.tecto.2012.12.028.
- Yamanaka, Y. and Kikuchi, M., 2003. Source process of the recurrent Tokachi-oki earthquake
 on September 26, 2003, inferred from teleseismic body waves. Earth Planets Space 55,
 e21-e24. doi:10.1186/BF03352479.
- Yamashita, Y., Shimizu, H. and Goto, K., 2012. Small repeating earthquake activity, interplate
 quasi-static slip, and interplate coupling in the Hyuga-nada, southwestern Japan subduction
 zone. Geophysical Research Letters 39, doi:Artn L08304 10.1029/2012gl051476.
- Yamashita, Y., Yakiwara, H., Asano, Y., Shimizu, H., Uchida, K., Hirano, S., Umakoshi, K.,
 Miyamachi, H., Nakamoto, M., Fukui, M., Kamizono, M., Kanehara, H., Yamada, T.,
 Shinohara, M. and Obara, K., 2015. Migrating tremor off southern Kyushu as evidence for
 slow slip of a shallow subduction interface. Science 348, 676-679. DOI:
 10.1126/science.aaa4242.
- Yata, T., 2009. Great earthquakes in Japanese medieval times. Yoshikawa kobunkan, 203p (in
 Japanese)*.
- 1925 Yata, T., 2018. Great earthquakes in Japanese early modern times. Yoshikawa kobunkan, 248p.1926 (in Japanese)*.
- Yata. T., 2013. Subsidence of the northern part of Lake Hamana during the 1707 Hoei
 Earthquake. Shiryogaku-kenkyu (Researches on historical materials), No. 10, 1-14 (in
 Japanese)*.
- Yokota, Y., Ishikawa, T., Watanabe, S., Tashiro, T. and Asada, A., 2016. Seafloor geodetic
 constraints on interplate coupling of the Nankai Trough megathrust zone. Nature 534,
 374-377, doi:10.1038/nature17632.

- Yoshikawa, T., Kaizuka, S. and Ota, Y., 1964. Mode of crustal movement in the late Quaternary
 on the southeast coast of Shikoku. southwestern Japan. Geographical Review of Japan 37,
 627-648. (in Japanese with English abstract) https://doi.org/10.4157/grj.37.627
- 1936 Zang, S.X., Chen, Q.Y., Ning, J.Y., Shen, Z.K. and Liu, Y.G., 2002. Motion of the Philippine
- 1937 Sea plate consistent with the NUVEL-1A model. Geophysical Journal International 150,
- 1938 809-819. https://doi.org/10.1046/j.1365-246X.2002.01744.x
- 1939
- 1940 *English translation from the original written in Japanese.
- 1941
- 1942

1943 Figure captions



1944

1945 Fig. 1. Map showing the outline of the seismotectonic setting around the Nankai and1946 Ryukyu subduction zones.

Tsunami source areas for the 1707 CE (green line) and two 1854 CE earthquakes (orange 1947 solid and dashed lines), as modified from Hatori (1974, 1976). Earthquake rupture areas of the 1948 1944 CE Tonankai and 1946 CE Showa-Nankai earthquakes (blue shaded areas) are simplified 1949 from Tanioka and Satake (2001a, b). Plate movement vector from Zang et al. (2002). 1950 Classification of rupture zones, A to E and Z, follows Ando, 1975 (a, b), Ishibashi, 1976, and 1951 Wells et al. (2003). Contour lines showing the depth of the Philippine Sea plate surface, 10 1952 km-interval, modified from Saito (2017). Distribution of high interpolate slip-deficit rate area 1953 (>5 cm yr⁻¹) along the Nankai Trough is simplified from Yokota et al. (2016). Spatial 1954 distribution of the averaged quasi-static slip rate in the Hyūga-nada region (from May 1994 to 1955 May 2011) is reproduced from Fig. 2 b of Yamashita et al. (2012) with the permission of 1956 publisher (John Wiley and Sons; license number 4775371178223). Source area of "the 1957

- 1958 maximum possible large earthquake" in Nankai subduction zone modified from Cabinet Office
- 1959 (2011b).
- 1960





Horizontal lines indicate Nankai megathrust earthquake rupture zones, following Fujiwara et al. (2020); dashed where uncertain, dotted to indicate the debated 1099 CE earthquake. We include four sites that record evidence for coseismic deformation (numbers 14, 21, 27 and 30).

1967 Site numbers: 1. Ibaruma, Ishigaki Island; 2. Sakishima Islands; 3. Miyako Islands[†]; 4.
1968 Okinawa Islands; 5. Amami Islands; 6. Kushima City[†]; 7. Ryūjin-ike; 8. Ryuuoo-ike; 9.

Tadasu-ike*; 10. Kaniga-ike; 11. Nankoku[†]; 12. Mugi Town[†]; 13. Kamoda-oike; 14. Kii 1969 Peninsula* (including Kuchiwabuka, Ameshima, Shionomisaki, Izumozaki, Arafunezaki, 1970 Ikeshima, Yamamibana, Taiji, and Suzushima); 15. Ōike (Sagari) Pond*; 16. Suwa Pond*; 17. 1971 Shima Lowlands[†]; 18. Ōsatsu Town*; 19. Nagaya Moto-Yashiki*; 20. Shirasuka; 21. Shinjo 1972 Lowlands[†]: 22. Arai*: 23. Lake Hamana*: 24. Western Hamamatsu strand plain[†]: 25. Ōtagawa 1973 Lowlands[†]; 26. Yokosuka Lowlands^{*}; 27. Omaezaki[†]; 28. Ova Lowlands^{*}; 29. Shimizu Plain^{*}; 1974 30. Ukishima-ga-hara*; 31. Ita Lowlands*; 32. Shimoda*. 1975 Sites with age recalibrated or modelled in Garrett et al. (2016) marked *, sites with age 1976

ranges recalibrated or modelled in this publication marked [†]. We modeled age ranges using the OxCal program v.4.2 (Bronk Ramsay, 2009) using P_Sequence and Sequence models (Bronk Ramsay, 1995, 2008, 2009), following the approach detailed in Garrett et al. (2016). For the Miyako Islands, we interpreted peaks in a cumulative probability distribution as indicating the timing of boulder movement events, following the approach of Araoka et al. (2013).

Basemap uses Global Multi-Resolution Topography (Ryan et al., 2009) through
 www.GeoMapApp.org.

1984

Main references cited in Garrett et al. (2016) but not discussed further in this paper are asfollows.

1987 Site 14: Shishikura et al. (2008a), Site 15, 16: Tsuji et al., (2002), Site 19: Takada et al., 2002,

1988 Site 22: Fujiwara et al. (2013b), Site 26: Fujiwara et al. 2007, Site 28: Kitamura et al. 2013),

1989 Site 32: Kitamura et al. (2014).



Fig. 3. Photographs showing examples of tsunami deposits

A) Overview of four historical tsunami deposits at the Ōtagawa Lowlands, facing the Tōkai
 segment of the Nankai Trough. Stratified sandy tsunami deposits are distinguished as
 horizontal color bands, which serve as an aquifer, in back-marsh mud.

- B) A *Porites* tsunami boulder with microatoll shape deposited on the eastern coast of Ishigaki
- 1999 Island, Japan. The scale at lower right is 30 cm.









Fig 4. Paleotsunami deposits in the western Hamamatsu Plain

- A) Geomorphic classification map of Hamamatsu-Lake Hamana area showing the main
 coring sites B to E. Map modified from Sato et al. (2011) with additional air photo
 analyses and land survey data.
- B) Geological columnar section and results of diatom analyses from Site B. Occurrence of possible tsunami sand beds is limited beneath the Kg tephra (1210-1187 cal BCE, Tani et al., 2013). Up-core changes in diatom assemblages suggest increasingly freshwater conditions in the back marsh. Increased occurrences of marine-brackish diatoms accompany the deposition of some sand beds. Figure captions are common for Fig. B to Fig. D.
- C) Geological columnar section from Site C. The occurrence of possible washover sand beds is
 limited to the period older than 1175-1055 cal BP (775-895 CE).
- D) Geological columnar section and results of diatom analyses from Site D. An increase of
 brackish-marine diatom species suggests the deposition of a tsunami sand bed ~3700-3300
 cal BP following the coastal subsidence.
- E) Geological cross-section at the site E. A tsunami sand bed showing a landward fining and
 thinning trend over 500 m along the incised valley axis. The occurrence of tsunami beds
 and other sand beds are limited below the Kg tephra.

Figs. 4B to 4D and Fig. 4E are modified from Sato et al. (2016a) and Fujiwara et al. (2013a),
respectively.



В	Ansei-type							Hoei-type						
	Z	A	в	С	D	Е	Z	A	В	С	D	E		
Showa								1946		1944				
Ansei		1854			1854				-					
Hoei									1707					
Meio		?			1498									
Shohei								1361		1361				
Kowa/Eicho		1099			1096									
Ninna									887					
Hakuho		684			?									

2026

Fig. 5 Schematic illustration of proposed recurrence models for subduction zone earthquakes

A) modified from Satake and Atwater (2007), B) simplified from Seno (2012).



2030 Fig. 6. Conceptual model illustrating the reconstruction of paleotsunami size from the spatial

2031 distribution of tsunami deposits considering the geomorphic evolution of a strand plain

2029

2032 This figure was originally shown as Fig. 3 in Fujiwara (2013) and quoted in Fujiwara and2033 Tanigawa (2017).

		Rupture zone										
		Sakishima	Okinawa	Amami	Osumi and Tokara	z	Α	в	с	D	E	
			Ryu	Hyuga-nada	Nankai Trough							
Length of history			~	?	~1300 years							
	1946 Showa											
	1944 Showa											
	1911 Kikai											
I	1854 Ansei											
	1854 Ansei											
	1791 event											
1	1771 Meiwa											
	1707 Hoei											
historical event	1614 Keicho											
(Mw>8.0)	1498 Meio						Δ	Δ				
	1361 Shohei											
	1361 Shohei											
	1099 Kowa											
	1096 Eicho											
	887 Ninna											
	684 Hakuho											
	latest 7th century								Δ	Δ		
Seismological	slow slip	notstudied	notstudied	notstudied	notstudied	0	0	0	0	0	0	
evidence	coupled or decoupled	Ø	0	notstudied	notstudied	notstudied	O	0	0	ø	0	
Paleotsunami	late Holocene paleotsunami evidence	Y	N	N	notstudied	very rare?	Y	Y	Y	Y	Y	
evidence	paleotsunami interval (years)	150-400	-	-	notstudied		90-265					
Geomorphologica I evidence	co-seismic crustal deformation	-	-	Y	-	-	Y	Y	Y	Y	Rare	
Shak	ing+tsunami+coastal deformat	ion	ΔL	iquefaction								
Shak			tsunam	i								
Shak												

2036 Table 1. Paleoearthquake and tsunami evidence and related seismotectonic data along the

2037 Nankai and Ryukyu subduction zones.