

This is a repository copy of *Valorisation of natural resources and the need for economic and sustainability assessment: the case of cocoa pod husk in Indonesia.*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/167221/>

Version: Accepted Version

Article:

Picchioni, Fiorella, Warren, Geoffrey P Warren, Lambert, Smilja et al. (9 more authors)
(Accepted: 2020) *Valorisation of natural resources and the need for economic and sustainability assessment: the case of cocoa pod husk in Indonesia*. Sustainability. ISSN 2071-1050 (In Press)

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 Article

2 **Valorisation of Natural Resources and the Need for** 3 **Economic and Sustainability Assessment: The Case** 4 **of Cocoa Pod Husk in Indonesia**

5 **Fiorella Picchioni** ^{1,*}, **Geoffrey P Warren** ², **Smilja Lambert** ³, **Kelvin Balcombe** ⁴, **J. Steve**
6 **Robinson** ², **Chittur Srinivasan** ⁴, **Leonardo Gomez** ⁵, **Laura Faas** ⁵, **Nicholas J. Westwood** ⁶,
7 **Afroditi Chatzifragkou** ⁷, **Dimitris Charalampopoulos** ⁷ and **Liz J. Shaw** ²

8 ¹ Natural Resources Institute, University of Greenwich and UK; f.picchioni@gre.ac.uk (F.P.)

9 ² Soil Research Centre, Department of Geography and Environmental Sciences, University of Reading, UK;
10 g.p.warren@reading.ac.uk (G.P.W.); j.s.robinson@reading.ac.uk (J.S.R.); e.j.shaw@reading.ac.uk (L.J.S.)

11 ³ Mars Symbioscience Indonesia, JL Kima 10 Kav A7 Daya, Makassar, South Sulawesi, Indonesia;
12 smilja.lambert@gmail.com (S.L)

13 ⁴ School of Agriculture Policy and Development, University of Reading, UK; k.g.balcombe@reading.ac.uk
14 (K.B.); c.s.srinivasan@reading.ac.uk (C.S.)

15 ⁵ Department of Biology, University of York, UK; leonardo.gomez@york.ac.uk (L.G.); laura.faas@york.ac.uk
16 (L.F.)

17 ⁶ School of Chemistry and Biomedical Sciences Research Complex, University of St Andrews, UK; njw3@st-
18 andrews.ac.uk (N.J.W.)

19 ⁷ Department of Food and Nutritional Sciences, University of Reading, UK; a.chatzifragkou@reading.a
20 (A.C.); d.charalampopoulos@reading.ac.uk (D.C.)

21 * Correspondence: f.picchioni@gre.ac.uk; Tel: +44 (0)1634 883685

22 Received: 30 September 2020; Accepted: date; Published: date

23 **Abstract:** The uptake of innovative technologies and practices in agriculture aimed at the
24 valorisation of natural resources can be scant in Low and Middle Income Countries (LMICs).
25 Integration of financial viability assessments with farmers and environmental evaluations can help
26 to understand some aspects of low uptakes of innovations. Using the case study of Cocoa Pod Husk
27 (CPH) valorisation in Indonesia, we provide insights on: (i) a choice modelling method to assess the
28 economic viability of CPH valorisation and (ii) an agronomic trial assessing the consequences on
29 soil quality of diverting CPH from its role as a natural fertilizer. The economic viability assessment
30 suggested that farmers require higher levels of compensation than might be expected to collect or
31 process CPH (a small proportion of farmers would undertake all processing activities for 117 GBP/t
32 CPH). The agronomic trial concluded that CPH plays only a minor role in the maintenance of soil
33 phosphorus, calcium and magnesium, but an important role for crop potassium. CPH removal
34 would reduce the partial balances for carbon and nitrogen by 15.6 and 19.6%, respectively. Diversion
35 of CPH from current practices should consider the long-term effects on soil quality, especially since
36 it might create increased reliance on mineral fertilizers.

37 **Keywords:** cocoa pod husk (CPH), valorisation; choice experiment; soil quality; soil carbon; soil
38 nutrients

39

40 **1. Introduction**

41 Rural households in Low and Middle Income Countries (LMICs) engage in a diverse set of
42 income-generating activities in an attempt to diversify their income base, accumulate wealth, buffer
43 the effects of shocks or break cycles of poverty [1–4]. However, income generation strategies are often

44 hindered by environmental threats and socio-economic factors that pose complex challenges
45 associated with poverty and food security [5]. Innovative technologies and processes that are directed
46 to the valorisation of raw and by-product or low-value materials can represent an opportunity to
47 create environmentally sustainable opportunities for new income streams directed to poorer rural
48 households and to create new markets [6,7].

49 Nonetheless, the uptake of technologies and alternative practices linked to innovative processes
50 can be disappointingly low [8]. To prevent the costly implementation of unsuccessful valorisation
51 projects due to poor uptake, there needs to be a prior assessment of the economic and environmental
52 viability of these initiatives. In this paper we provide an example of choice modelling methods to
53 assess the economic viability from the farmers' viewpoint of diverting what is generally considered
54 a low value by-product (cocoa pod husks (CPH)) to sale for income generation, alongside assessment
55 of the implications for soil quality that result from diverting CPHs from their original return to the
56 soil of cocoa farms. Timely financial viability assessments and environmental impact evaluations can
57 help to understand some of the aspects of low uptake of improved technologies and could feed into
58 a broader cost benefit analysis. For example, the discrete choice experiment addresses farmers'
59 willingness to accept innovative production processes and additional costs to modify their
60 established practices linked to cocoa farming. Additionally, assessing the environmental impacts of
61 by-product material valorisation examines unintended environmental damages and medium- and
62 long-term threats to soil quality and farm productivity.

63 CPH, generated when the cocoa beans are removed from the cocoa pods, is the main by-product
64 of the cocoa harvest with the CPH constituting about 75% of the mass of whole cocoa pod fruit. As
65 reviewed in Fei et al. (2018), after harvest, CPH is commonly recycled to soil on farm as fertilizer,
66 although it is frequently removed to reduce the spread of diseases. It may be used for other relatively
67 low value applications such as animal feed, starting material for soaps and preparation of activated
68 carbon. However, alternative processing and valorisation of CPH may represent an unexplored
69 income stream for cocoa producing farmers. CPH is rich in minerals (particularly K), fibre including
70 lignin, cellulose, hemicellulose and pectin and antioxidants such as phenolic acids. There are
71 interesting agrochemical valorisation potentials for CPH and its fractions that include their
72 incorporation in food production. Such innovative applications to CPH allow the addition of fibre-
73 rich husk fractions into different food products (bakery, dairy, chocolate-based confectionary
74 products) and can be potentially used to reduce sugar and fat content and increase fibre content of a
75 large range of food products [9].

76 However, to supply CPH for new uses, farmers may need to modify part of their cocoa
77 production processes and integrate alternative harvesting practices. These may generate extra costs,
78 and modification in the allocation of labour and farm space that could compete with more valuable
79 production of cocoa beans or other crops. Hence, farmers may demand compensation to adopt
80 alternative practices directed to CPH valorisation. Understanding the type of practices that need to
81 be introduced and assessing the adequate compensation levels are central to improve the technology
82 uptake and its long-term financial viability [10].

83 At the same time, the large internal cycle of organic matter and nutrients within the field via
84 decomposing crop residues and litter from both cocoa and shade trees [11,12] is a major influence on
85 the maintenance of soil fertility for cocoa production. The retention of crop residues is a key in
86 promoting physical, chemical, and biological attributes of soil health in agricultural systems for
87 developing countries [13] and soil organic carbon (SOC) is a key indicator of soil quality and
88 sustainability because it measures soil organic matter (SOM), which provides beneficial soil physical
89 properties (e.g., aggregate stability, soil water retention) and the provision of plant-available
90 nutrients. Most of the variability of nutrient content of the cocoa harvest can be attributed to
91 differences in the nutrient content of the CPH rather than that of the beans [12]. Therefore, the
92 diversion of CPH to other uses could result in significant depletion of the crop's nutrient supply and
93 soil quality. Hence, assessing the balance between nutrient inputs and outputs is crucial to inform
94 the long-term sustainability of cocoa and the need for fertilizer. Published data on the nutrient
95 element composition of CPH are mostly from Africa and Latin America and quantitative information

96 on the contribution of CPH to the internal and external nutrient balances is scarce. Fontes et al. [14]
97 found that Brazilian cocoa agroforestry systems varied greatly in nutrient cycling and internal
98 balance, concluding that CPH was particularly relevant for P and K, and that further research is
99 needed to develop nutritional balance systems integrating litter, fruits, and nutrient pathways. Our
100 work addresses this need in the Indonesian situation and furthermore links the biophysical nutrient
101 balance information with socioeconomic aspects of CPH management.

102 Drawing on data collected in Indonesia we set out to empirically assess: (i) the financial
103 feasibility of innovative processes directed to CPH by employing a discrete choice experiment and;
104 (ii) the potential environmental impacts of the off-farm removal of CPH on soil properties and soil
105 nutrient cycling. Our innovative approach speaks to the literatures contributing to an integrated and
106 innovative “technology translation” paradigm [10] using an approach that bridges biophysical
107 science and economics.

108 2. Materials and Methods

109 2.1. Economic Viability Assessment

110 The financial feasibility of CPH valorisation and farmers’ willingness to adopt the proposed
111 practices is explored by using Discrete Choice Experiment (DCE) and followed three stages:

- 112 • Exploratory survey, designed and administered to a small sample of the farmers to provide an
113 initial understanding of CPH practices in Sulawesi and informs the design of the DCE;
- 114 • Focus Group Discussions to identify compensation levels to process CPH;
- 115 • Design and deployment of DCE survey.

116 2.1.1. Exploratory Survey

117 The exploratory survey consisted of three sections: (i) farmer’s and farm characteristics; (ii) cocoa
118 production practices; (iii) CPH practices.¹ The island of Sulawesi (circled in **Error! Reference source**
119 **not found.**) represents the main area of Indonesia where cocoa is produced, mainly by smallholder
120 farmers (about 90% of total production) (Johnston et al. 2004). Given the centrality of Sulawesi in the
121 cocoa production, the pilot study and subsequently the DCE were deployed in this area. Sample
122 selection and the identification of 56 respondents were supported by the International Coffee and
123 Cocoa Research Institute (ICCRI), and interviews were conducted in four districts covering the four
124 main areas of the island: Luwu (South Sulawesi) 15 farmers; Mamuju (West Sulawesi) 16 participants;
125 Parigi Moutong (Central Sulawesi) 11; and Kolaka (Southeast Sulawesi) 10 farmers.

126 57% of farmers reported that CPH went unused, 34% dispose of CPH mainly as fertilizer and
127 8% as animal feed and fertilizer combined (8%). However, it was not clear if farmers that did not use
128 CPH were leaving it on the field. Only 4% of farmers sold a proportion of CPH in the preceding year.
129 When used as animal feed, CPH processing is minimal (sometimes involving chopping and drying).
130 When used as compost or mulch, the processing includes “rough” chopping, and CPH is left in the
131 fields to decay and recycle nutrients into the soil.

132 The survey indicates that the main processes that farmers would have to undertake in order to
133 modify their current practices with regards to CPH and supply material for new uses, include
134 collecting, chopping, drying and transporting. These four processing stages are used as attributes in
135 the DCE. Farmers indicated their willingness to adopt alternative practices for the disposal of husks.
136 Finally, while farmers are responsive to price incentives, considerations about obstacles to transport
137 CPH and additional labour requirement were identified as main challenges in modifying current
138 practices around CPH.

139 2.1.2. Focus Group Discussions (FGD)

¹ Field-work and data collection for the pilot study was executed by ICCRI staff in May 2018 and interviewed 52 farmers in the island of Sulawesi. Before deploying the survey, ICCRI tested the questionnaire on five farmers in East Java (in the area of Blitar). This exercise provided the platform to develop and finalize the pilot questionnaire.

140 The focus group discussion aimed at benchmarking the compensation levels that would be
 141 appropriate to use in the choice modelling method to assess the economic viability of CPH
 142 valorisation. The FGD consisted of two stages. The first stage was undertaken with five farmers in
 143 the area of Blitar (East Java) and five farmers in Lowu (South Sulawesi). The second stage involved
 144 interviews with experts at ICCRI, local partners in the project. Respondents were asked to
 145 individually provide both the lowest and highest levels of compensation they believed were
 146 necessary for them and other farmers in their village to collect, chop, dry and transport CPH.² At the
 147 end of the FGD with the farmers, enumerators conducted a group exercise to reach a consensus on
 148 the lower and upper bound compensation level for each of the processing stage (**Error! Reference**
 149 **source not found.**).

150 **Table 1.** Median lower and upper limit of the compensation demanded to collect, chop, dry and
 151 transport dry CPH (GBP/t).


	Collect	Chop	Dry	Transport
	GBP/t			
Median Lower Limit	36.4	25.5	52.4	52.0
Median Upper Limit	59.7	26.0	78.0	70.2

152 Exchange rate used: GBP 1 (Great Britain Pound) = IDR 19231 (Indonesian Rupiah). Source:
 153 (<https://www1.oanda.com>).

154 We report median values due to substantial variations in reported compensation levels. Drying
 155 is associated with the highest compensation level followed by transport and collection of CPH.
 156 Chopping is the activity that appears to require the lowest compensation. These trends are consistent
 157 for both upper and lower limits.

158 2.1.3. Design and Deployment of DCE

159 Field work was conducted in South and South East Sulawesi, Lampung (South Sumatra) and
 160 West Sumatra. 156 small and medium size cocoa farmers across four districts were interviewed
 161 (**Error! Reference source not found.**)³



162

163 **Figure 1.** Locations of the districts of the Discrete Choice Experiment in Sulawesi and Sumatra
 164 (Indonesia) with numbers of participating farmers indicated for each district.

165 The pilot survey and discussion with ICCRI experts helped determine the list of attributes for
 166 the DCE. The information collected during the focus group discussion was used to define the prices
 167 attached to the attributes in each set of cards. The cards were used to assess the compensation needed

² Farmers were asked to identify the transportation cost required to transport CPH for 15 km. Compensation levels were asked per kg of wet CPH or wet-equivalent CPH.

³ The project followed the ethical procedure detailed by the partner institution (ICCRI). All eligible farmers were informed of the purpose of the study and their right to refuse participation. Interviews were performed after receipt of written consent from the participants. At the end of the interview respondents were remunerated with a small amount of cash to thank them for their time.

168 to accept new/additional processing practices connected with supplying CPH for new uses. We
169 identified six options that include one or more identified attributes:

- 170 a) Collect
- 171 b) Collect and Transport
- 172 c) Collect and Chop
- 173 d) Collect, Chop and Transport
- 174 e) Collect, Chop and Dry
- 175 f) Collect, Chop, Dry and Transport

176 Compensation levels are expressed in wet-equivalent CPH, using the conversion factor of 0.22.⁴
177 The attributes and prices generated 192 alternatives. Each participant was presented with 32 choice
178 cards, each showing six options of CPH combination of attributes and their compensation level. An
179 opt-out question was included. Details on the design of the econometric estimation are provided in
180 Appendix A. After being informed of the purpose of the exercise, farmers were asked to select one
181 preferred compensation level attributed to one option on each card. Each choice option was
182 accompanied by a standardized script, which was read by the interviewer. Sample DCE cards are
183 shown in S1 and **Error! Reference source not found.** in the Supplementary Material.

184 We conducted a short survey following the DCE module, asking farmers' information of farm
185 characteristics, cocoa production, labour availability (family and employed), CPH current practices
186 (fertilizer and feed), and household demographics. The survey and consent forms were developed in
187 English and translated to Bahasa Indonesia. The surveys were piloted, revised, and administered by
188 a team of five experienced interviewers. Interviews lasted 30–40 min and were conducted using
189 electronic tablets (Lenovo Tab4 8) with SurveyCTO software (Dobility, Inc., Boston, United States of
190 America).⁵

191 2.2. Soil Quality Assessment

192 2.2.1. Field Trial Design and Assessment Methods

193 In order to evaluate the role of CPH in soil fertility maintenance, a programme of additional
194 sampling and analysis was conducted over 15 months in one treatment of an established on-station
195 trial of the Cocoa Research Station of MARS Inc, Tarengge, Sulawesi, Indonesia (12°33' S; 120°47' E ;
196 altitude ~12 m ASL). The trial was planted in February 2014 and aimed to be realistic of local on-farm
197 conditions by including banana (*Musa paradisiaca*) and gliricidia (*Gliricidia sepium*) companion trees.
198 Nearby trees, in particular two mature durian (*Durio zibethinus*) trees (thought to be >40 years old),
199 that contributed to litter fall also remained undisturbed to maintain a realistic situation. The
200 treatment monitored provided mineral fertilizers at the current local "best practice" annual rates of
201 82, 92, 155, 330 and 115 kg/ha of N, P, K, Ca and Mg respectively, based on the suppliers' analyses.
202 There were three replicate plots and cocoa seedlings of MARS clone MCC02 were planted at a density
203 of 1111 plants/ha.

204 2.2.2. Harvesting and Sample Collection

205 Cocoa pods are produced throughout the year and all ripe pods were harvested from all trees in
206 each plot at intervals of two weeks and the number recorded. During the peak production seasons of
207 June–July 2018 and October–December, 2017 and 2018, a sample of 12 pods was taken per plot and
208 split into beans and the remainder (CPH), the latter being chopped into small (<2cm) pieces. Beans
209 and CPH were air dried and weighed. The biomass production at the other harvests was estimated
210 from the number of pods harvested and the ratio of bean weight to CPH weight as established for

⁴ The conversion factor was derived at ICCRI and described in Appendix B.

⁵ Training on digital data collection and DCE surveys deployment was conducted between 28th and 30th of November 2018 at ICCRI headquarter (Jember, Indonesia), followed by a pilot survey collection in Blitar between 3-7 of December 2018. Data were collected over a period of 8 weeks (11th December 2018 to 31st January 2019), after the completion of the main cocoa harvest season.

211 the measured harvests. Trees were pruned in April, July and November 2018. Leaves and woody
 212 parts were collected separately, weights recorded and the woody parts were chopped into smaller
 213 pieces. Samples of each were air-dried for analysis and the rest returned to the plot. Falling leaf litter
 214 in each plot was collected in three square traps (each 0.5m²) per plot, which were shifted to different
 215 random positions every three months. Litter was collected every seven to 10 days, air-dried and
 216 combined into batches each representing one month finishing on the 14th of each month from
 217 November 2017 to December 2018. Samples collected in the June to October period were sorted by
 218 hand into that of cocoa origin and the rest, the latter mostly originating from the shade trees.

219 2.2.3. Assessment of C and Nutrient Cycling in Soil

220 Subsamples of each of the above materials were analysed for total C, N, P, K, Ca and Mg at
 221 University of Reading laboratories according to standard methods (Appendix C). A partial balance
 222 of external inputs and offtakes of the plots for each of these elements was calculated using elemental
 223 concentrations and dry matter yields, for a period of one year, from December 2017 to November
 224 2018. Data for losses via drainage water and gaseous emissions (e.g., CO₂, CH₄, NH₃, N₂O, NO_x) and
 225 inputs via atmospheric deposition and biological nitrogen fixation were not available for this location.
 226 The offtakes were the sum of results for beans and CPH (when appropriate). Nutrient inputs via
 227 litter, prunings and CPH (when appropriate) were considered to be internally cycled. Therefore
 228 external nutrient inputs were fertilizer and the C inputs were litter, prunings and CPH (when
 229 appropriate), since C is derived from photosynthesis by crop and accompanying trees.

230 3. Results

231 3.1. Economic Viability Assessment

232 Responses were elicited from 156 farmers using face-to-face interviews technique. Demographic
 233 characteristics of the respondents are summarized in **Error! Reference source not found.**. The mean
 234 age was 49 years and 61% of interviewed farmers were men. The household size was on average 4.45
 235 (with 3.36 adults and 1.09 children). The average landholding was 1.67 ha with 1.07 ha allocated to
 236 cocoa farming, and during the previous agricultural year, farmers had produced on average 336.5 kg
 237 of (dried) cocoa. Households are located almost 8 km away from the nearest product markets. 50%
 238 of farmers use CPH as fertilizers and only 16% use it for animal feed.


239 **Table 2.** Descriptive Statistics of households and farms.

	Mean	SD
Age of respondents (years)	48.8	(11.87)
Household size	4.45	(1.59)
Number of adults (>15 years old)	3.36	(1.34)
Number of children (<=15 years old)	1.09	(1.07)
Total land holding (ha)	1.67	(1.67)
Total size of cocoa farm (ha)	1.07	(0.96)
Age of cocoa trees (years)	12.91	(7.63)
Distance from nearest market (km)	7.63	(8.74)
Cocoa beans produced last year (dry, kg)	366.28	(472.72)

Percentage of:	
Male respondents	61%
Farmers using CPH as fertilizer	50%
Farmers using CPH as animal feed	16%
Number of households	
	156

240 **Error! Reference source not found.** reports the distribution of farmers' Willingness To Accept
 241 (WTA) compensation in order to perform the six sequences of activities required to supplying CPH
 242 for new uses. The options did not prompt farmers to think about soil nutrient losses (especially
 243 potassium) and to quantify additional compensation needed to replace them with fertilizer. Hence,
 244 we are not able to differentiate the compensation needed to purchase fertilizers from the attributes
 245 presented in the choice cards. Prices are reported in GBP/t of dry-equivalent CPH. Figure 2. Cocoa
 246 farmers Willingness-To-Accept alternative practices with CPH (GBP/t).

247 reports the mean, standard error of the mean, standard deviation and median of the
 248 compensations by for each individual activity. These figures illustrate that chopping requires on
 249 average the lowest compensation level, followed by collecting, transporting, and drying husks.



250

251

Figure 2. Cocoa farmers Willingness-To-Accept alternative practices with CPH (GBP/t).

252

Table 3. Willingness to accept compensation break-down (GBP/t).

	Collect	Chop	Dry	Transport
Mean	235.2	162.6	354.0	316.1
SEM	7.1	7.5	9.6	10.1
SD	88.9	93.8	120.3	125.7
Median	103.9	50.3	553.5	268.3

253 In general, a substantial proportion of farmers signalled a willingness to engage in activities such
 254 as collecting through to transporting CPH. However, only approximately 5% of farmers appear to be
 255 prepared to perform all the activities (collecting, chopping, drying and transporting) for a
 256 compensation level between 140–150 GBP/t of dry CPH, and most require a relatively higher

257 compensation level. Almost 50% of farmers are willing to undertake collection only of CPH for
258 approximately 75–100 GBP/t of dry CPH. Collecting and transporting and collecting, chopping and
259 transporting exhibit very similar compensation distributions, and 20% of farmers are willing to
260 undertake these activities for approximately 100 GBP/t of dry CPH.

261 While some farmers were willing to undertake activities for relatively smaller compensation
262 levels, they were also the ones that tended to require substantial amounts for the same activities
263 earlier in the sequence. This is indicative of the fact that once farmers decide there are only a set of
264 activities they will engage with, they appear to exhibit considerable levels of inertia and low levels of
265 responsiveness to monetary incentives to deviate from that decision. Farmers are likely to be inclined
266 to prioritise cocoa bean production and exhibit a certain level of resistance to allocate resources (i.e.,
267 labour and farm space) to CPH management.


268 In terms of types of farmers who were prepared to undertake these activities, there was very
269 little explanatory power using revenue from CPH (closely related to size and level of production),
270 labour and distance from the market as explanatory variables. Around 7% of variation in the WTA
271 for collection and drying activities could be explained by these factors whereas for chopping only
272 around 1.5% could be explained and 3% for transporting. In terms of statistical significance only
273 revenue was significant for collection and transporting and distance for drying at a 10% level of
274 significance.

275 Indonesia is among the biggest cocoa producers (593,832 t in 2018; [15]) and provides the main
276 source of income for over 1,400,000 smallholder farmers [16]. Because lower levels of compensation
277 are demanded by 5% of interviewed farmers, CPH valorisation could potentially benefit about 70,000
278 Indonesian cocoa farmers and their families willing to engage with all processing activities. Over time
279 and with the development of infrastructures and processing systems that better integrate the
280 valorisation of CPH into existing processes and value chains, the cost of processing CPH could
281 gradually decrease. This could enable a larger share of farmers to take advantage of CPH sale and
282 diversify their incomes.

283 3.2. Soil Quality Assessment

284 3.2.1. Crop Performance

285 Pod, bean and CPH production varied greatly during the experimental period, in accordance
286 with the two normal main cocoa harvest peaks in Sulawesi, around July and November (**Error!**
287 **Reference source not found.**). The annual bean yield of 1292 kg/ha was high compared with typical
288 current Indonesian yields of around 500–700 kg/ha [16], and approach the best yields (~2000 kg/ha)
289 obtained in a recent on-farm trial in another area of south Sulawesi [17]. The annual dry biomass
290 input to the soil was 8323 kg/ha, provided by CPH (1328 kg/ha), litter (4526 kg/ha) and prunings
291 (2469 kg/ha). The variation in litter fall over the year suggested a tendency to increase over the
292 duration of the experiment (Figure 3), although low litter fall in the May to July period may be
293 associated with wetter weather at this time since cocoa litterfall is highest in dry weather.



294

295 **Figure 3.** Monthly CPH production (hatched column) and litter fall (open column) in the year
 296 November 2017 to December 2018, for periods ending on 14th of each month with error bars denoting
 297 \pm (SD).

298 The annual C and nutrient contents for CPH were 504, 11, 1, 46, 7 and 4 kg/ha for C, N, P, K, Ca
 299 and Mg respectively (**Error! Reference source not found.**). Similar values are reported by Wood and
 300 Lass [18] for a bean yield of 1000 kg/ha. They amount to 15.6% (C), 6.2% (N), 1.1% (P), 18.0% (K), 1.5%
 301 (Ca) and 2.7% (Mg) of the above-ground inputs to the soil from external (i.e., photosynthate-C,
 302 biologically fixed-N, and fertilizer N, P, K, Ca and Mg) and internally recycled (N, P, K, Ca and Mg)
 303 sources, suggesting that CPH played an important role in the C and K cycles, but a lesser part in the
 304 cycles of the other nutrients.

305 This trial was young (the study took place in its fourth year) compared to the mean tree age of
 306 12.9 years at the farms investigated (**Error! Reference source not found.**) and the typical lifespan for
 307 hybrid cocoa is 15–20 years [12]. However, for the purpose of nutrient balance assessment, it is
 308 concluded from the crop performance results that it is reasonably representative of a high-yielding
 309 cocoa crop in Sulawesi.

310 **Table 4.** Partial annual plot-level balances (kg/ha) of C, N, P, K, Ca and Mg under the alternatives:
 311 Scenario 1 (CPH retained); Scenario 2 (CPH removed); and Scenario 3 (CPH removed and no
 312 fertilizer), with standard deviations for each element in the scenarios.

Element	Nutrient Cycle Component					Scenario			
	Litter	Prunings	Fertilizer	CPH	Bean	1	2	3	(SD)
C	1791	935	0	504	n/a	3230	2726	2726	369
N	58	26	82	11	27	56	45	−38	3.2
P	4	3	92	1	6	86	85	−8	0.3
K	21	35	155	46	14	141	94	−61	5.0
Ca	75	27	330	7	1	328	322	−8	1.0
Mg	18	10	115	4	4	111	107	−8	0.4

313 3.2.2. Nutrient Cycling and Sustainability

314 Shade trees play a significant role in nutrient cycling in addition to their agronomic value and
 315 in the period from May to October, 67, 63, 62, 43 and 44% of the N, P, K, Ca and Mg respectively in
 316 the litter came from the shade trees (Appendix C), showing their value for maintaining soil fertility.
 317 Given the presence of leguminous shade trees (i.e., *Gliricidia sepium*), a proportion of the shade litter

318 input of N was probably derived from biological fixation from the atmosphere and therefore
319 represents an external (as opposed to internally-recycled) input to the system.

320 The net balance of inputs minus offtakes provides an indicator of sustainability because
321 continuous depletion of nutrients in the soil is not sustainable in the long term. Three scenarios were
322 quantified: (1) CPH retained on the plot, as actually carried out, (2) CPH removed and (3) CPH
323 removed and fertilizer application discontinued. Scenario 1 showed a surplus of C and nutrients
324 (**Error! Reference source not found.**). Under scenario 2, the C and nutrient balances were reduced
325 significantly ($P < 0.05$) showing that CPH made a significant contribution to maintaining soil quality.
326 Under scenario 3 the balances for the nutrients would be reduced highly significantly ($P < 0.001$),
327 becoming negative.

328 The maintenance, turnover and potential accumulation of soil organic carbon is a function of the
329 long-term balance between plant carbon inputs and losses, mainly via microbial respiration. CPH
330 provided about 16% of the above-ground C input to soil, representing a reduced annual input of ~500
331 kg/ha (equivalent to 50 g/m²) under the CPH removal scenarios (**Error! Reference source not found.**).
332 The magnitude of the CPH-associated C input should be considered not only alongside the above
333 ground inputs from litter and prunings but also alongside the likely significant C inputs to SOC from
334 fine root turnover below-ground, which were not measured here but have been estimated to be ~140
335 g/m² and ~180 g/m² annually in 10 year old cocoa systems in Central Sulawesi under natural or
336 planted shade trees, respectively [19]. Whilst current evidence suggests that below-ground plant-
337 derived inputs might be most influential for building SOC [20], there is considerable uncertainty
338 regarding the consequences of variation in both quantity and quality of above ground plant-derived
339 inputs to soil for below-ground processes, as a result of the complexity of governing mechanisms and
340 factors involved. However, a meta-analysis [21] has indicated that (sub-) tropical forest systems may
341 be most sensitive to increases or decreases in above-ground inputs, with substantial alterations in the
342 turnover and accumulation of SOC over relatively short time scales possible. Indeed, a study in
343 primary tropical forest (Costa Rica) has shown that only two years after annual C inputs were
344 reduced by 450 g/m² by leaf litter removal, SOC concentrations were reduced by 26% [22]. In our CPH
345 removal scenario, how the comparatively modest reduction (50 g/m²) in C input would affect SOC
346 dynamics over the short and longer term requires further investigation. Data of Dawoe et al. [23] and
347 Tondoh et al. [24] in West Africa emphasize that the initial conversion of forest to cocoa causes a
348 decline in SOC due to perturbations in the balance between C input and output. If CPH was sold off-
349 site, the further perturbation caused by lost C input could be mitigated by addition of an alternative
350 source of organic matter to prevent further decline in SOC and requires further investigation.

351 CPH provided 6% of the N input to soil and the external balance showed a surplus in scenario
352 1, the current practice (**Error! Reference source not found.**). There are several uncertainties in this N
353 balance. Substantial losses of N from soil via leaching in drainage water are possible in humid
354 climates and not accounted for here. However, the deeper roots of many shade trees provide a “safety
355 net” to capture soluble nutrients and recycle them to the soil via plant uptake and litter, so Hartemink
356 [11] concluded that leaching under cocoa might be between “very small” and 22 kg/ha annually,
357 based on measurements in Latin America. Based on other research in cocoa agroforestry in Central
358 Sulawesi [25], relatively small N losses via gaseous emissions (e.g., ≈ 3 kg N as N₂O/ha annually) are
359 also likely. Based on estimates from cocoa systems in Malaysia, Central & South America and Africa,
360 annual deposition of N in rainfall may add a further 5 to 12 kg/ha [11], although some records suggest
361 that specific N deposition rates in Sulawesi maybe be lower at 1.4 to 2.7 kg/ha [26]. As already
362 mentioned, we also did not quantify the contribution of biologically fixed N as an external input to
363 the system. However, based on our partial N budget, results overall suggest that offtake of N via
364 CPH would not cause a major shortfall in the N supply if fertilization is maintained at current levels.
365 But without fertilizer, CPH would be an important N source because of the negative balance.

366 Leaching losses also affect K and might be up to 17 kg/ha (Hartemink 2005) per year. CPH was
367 the largest single non-fertilizer source of K (**Error! Reference source not found.**). If CPH was removed
368 from the land, the K supply to soil would be reduced by 18%. Without CPH and mineral fertilizer,
369 there would be a substantial deficit in K, further increased by leaching losses: CPH plays a significant

370 role in the maintenance of K fertility. The impact of CPH removal on nutrient balances would be quite
371 small for P, Ca and Mg, accounting for 1.1, 1.5 and 2.7% respectively of the internal nutrient cycles.

372 4. Discussion and Conclusions

373 Valorisation of farm by-products or (seemingly) non-utilized agricultural material is often
374 viewed an attractive option to diversify income or generate revenue among poorer farmers in LMICs.
375 However, failing to consider the economic viability from the smallholders' viewpoint and assess the
376 environmental effects of such strategies, represents a considerable obstacle for the overall uptake and
377 long-term sustainability of such interventions. In addition, unintended (or ill planned) impacts on
378 soil quality can hinder agricultural productivity, damage the main resource for livelihood (the soil)
379 of poorer farmers and reverse the original positive intent of the intervention.

380 In this paper the assessment of residue material valorisation was undertaken with regards to
381 alternative uses of CPH directed to the agri-food industry. The paper provides insights into the
382 evaluation process on two fronts: (i) choice modelling methods to assess the economic viability of
383 supplying CPH for new uses; and (ii) impacts on soil quality caused by diverting CPH from its role
384 of organic fertilizer.

385 The economic viability assessment concludes that farmers require higher than expected levels of
386 compensation to supply CPH for new uses. Such prices are comparatively high for a material that
387 could be considered low value or unused. Based on the findings from the exploratory survey, this
388 can signal that to adapt current practices, farmers require additional resources to adjust labour
389 constraints and compensate farm space originally allocated to cocoa bean production. We assessed
390 that approximately 5% of farmers are in the position to supply CPH for new uses at a starting
391 compensation level of 150 GBP/t. Given the large population of smallholder cocoa producers in
392 Indonesia, a significant number of farmers could take part to the CPH valorisation process and
393 supply husks for new uses. Over time these numbers can grow further. Adoption and diffusion of
394 agricultural technologies comprehend a certain degree of complexity in the decisions taken by small-
395 scale farmers. To a certain extent, the complexity lies in the lack of certainty with relation to the
396 benefits of such technologies before they are adopted [27]. Improvements in infrastructures and
397 integration of processes directed to CPH alternative uses in existing food value chains can address
398 such uncertainties, maximise farmers' financial benefits. Additional labour costs barriers could be
399 addressed thanks to better integrated production systems and more competitive compensations.

400 Reviews of cocoa fertilization show that recommended nutrient application rates vary more than
401 10-fold [12]. The amounts of fertilizer used here were high in relation to the crop requirement since
402 there appears to be a surplus of most nutrients. The partial nutrient balances indicated large surpluses
403 of OM and macronutrient accumulation (**Error! Reference source not found.**), so diversion of CPH
404 to alternative uses is not expected to cause nutrient deficiencies in the short term. CPH played only a
405 minor role in the maintenance of soil P, Ca and Mg, but CPH diversion would reduce the plot-level
406 partial balances for C, N and K by 15.6, 19.6 and 32.6% respectively. These significant losses could be
407 important if continued for several years. K is the nutrient most susceptible to deficiency and complete
408 replacement of the 46 kg/ha from CPH (**Error! Reference source not found.**) by KCl fertilizer would
409 cost IDR 578,080 (GBP 30) per year at the current price.

410 C input from diverted CPH could be replaced, if there was an appropriate alternative organic
411 amendment available. In our Sulawesi setting, there is a locally available commercial organic
412 fertiliser, 'Petroganik'. Direct replacement of 504 kg/ha of CPH-derived C (calculated from **Error!**
413 **Reference source not found.**) would require the application of 3359 kg/ha Petroganik (taking into
414 account the C content of the organic fertiliser) costing IDR 9,238,184 (GBP 480). This would also
415 provide 34 kg N and 27 kg K, adding 309% of the N and 59% of the K lost in diverted CPH. Hence,
416 any decision to divert CPH from its conventional use should take into account the long term effects
417 on soil quality, since it may create increased reliance on purchased inputs. However, it may be
418 acceptable to reduce the currently recommended fertilizer application rates in the light of the
419 apparent balance surpluses and possible consequent environmental pollution by excess nutrients in
420 drainage waters.

421 The nutrient cycling study suggested possible further environmental constraints on the
422 economics of CPH valorisation as experienced by farmers. CPH production varied widely across the
423 year (Figure 3). Despite the willingness of some farmers to process CPH for sale, the seasonality of
424 the work could impose labour constraints on its practicality. The subsidiary investigation of the role
425 of nutrient cycling via shade and companion trees (Appendix C) showed that shade trees (including
426 in this case deposition from nearby large trees) have a significant role in nutrient cycling in cocoa
427 production. In addition to these beneficial effects on soil fertility, accompanying trees in cocoa
428 systems normally provide significant economic benefits in terms of fruit and other products. Farmers'
429 locations and choice of accompanying trees may have location-specific influences on the recycling of
430 CPH for valorisation.

431 Considerations of the financial viability and environmental impact are central to address the low
432 uptake of improved technology. Challenges to adopt innovation by end-users (such as unrecognised
433 problems for farmers) may be linked to the strategies adopted in the process of technology
434 development, where top-down “solutions” are implemented without a comprehensive involvement
435 of all actors or without performing economic and environmental assessments [28–30]. Holistic
436 approaches to technological development that focus on integrating different actors and consider
437 national resources, and that evolve within broader agricultural development strategies, are in a better
438 position to address the complex and unbounded sustainability issues facing agriculture today [31].
439 The results of our study indicate that technological interventions, and by-product material
440 valorisation in particular, can benefit from the involvement of multi-disciplinary teams and
441 engagement with farmers from the onset of the development of innovative farming approaches. This
442 allows to assess whether the technologies or innovative approaches meet existing needs and
443 preferences of the end-users as well as assessing environmental impacts and effects on soil quality.
444 Such integrated, participatory, and interdisciplinary practices have the potential to enhance adoption,
445 ensure the sustainability of these innovative technologies and ultimately result in alternative income
446 streams for rural households.

447 **Supplementary Materials:** The following are available online at www.mdpi.com/xxx/s1, Figure S1: DCE card –
448 English, Figure S2: DCE card – Indonesian.

449 **Author Contributions:** Conceptualization, F.P., G.P.W., L.J.S., K.B., C.S. and J.S.R.; methodology, G.P.W., L.J.S.,
450 K.B. and C.S.; software, F.P., K.B., ?; validation, ?; formal analysis, F.P., G.P.W., L.J.S., K.B.; investigation, F.P.,
451 G.P.W., L.J.S., K.B.; resources D.C.; data curation, F.P., G.P.W., L.J.S., K.B.; writing—original draft preparation,
452 F.P., G.P.W., L.J.S., K.B.; writing—review and editing, F.P., G.P.W., L.J.S., K.B., C.S., S.L., J.S.R., L.G., L.F., N.J.W.,
453 A.C., D.C.; visualization, F.P., G.P.W., L.J.S., K.B.; supervision, L.J.S., K.B., C.S., D.C.; project administration, D.C.;
454 funding acquisition, D.C., L.J.S., K.B., C.S., J.S.R., L.G., L.F., N.J.W., A.C. All authors have read and agreed to the
455 published version of the manuscript.

456 **Funding:** The authors would like to acknowledge the Biotechnology and Biological Sciences Research Council
457 (BBSRC) and Mars Wrigley Confectionery for their financial support on a collaborative Global Challenges
458 Research Fund (GCRF) project entitled “Development of novel value chain from cocoa pod husks in Indonesia:
459 Technological, environmental and socio-economic challenges of a value chain” (BB/P022995/1).

460 **Acknowledgments:** The authors are grateful to ICCRI specialists and field staff for their assistance and guidance
461 during field work of the economic viability assessment, and staff of MARS Cocoa Research Station for sampling
462 in the cocoa field trial. The authors wish to thank all respondents for their valuable time and cooperation.

463 **Conflicts of Interest:** Authors would like to declare no conflict of interest.

464 Appendix A

465 The design for this experiment contained the same options (a. Collect, b. Collect and Transport,
466 c. Collect and Chop, d. Collect, Chop and Transport, e. Collect, Chop and Dry, f. Collect, Chop, Dry
467 and Transport) with only variation in price across these options. The price levels were constructed
468 from the initial open ended survey of farmers and experts ranging from 1/3 of the median “entry
469 level” price (the price these respondents felt that some farmers would accept) expressed by these
470 respondents to three times the “majority level” price (the price these respondents expressed the most

471 farmers would accept). The construction of the 32 choice options was then made using a D-optimal
 472 design (which should maximise the efficiency of the estimates) under weak prior assumptions about
 473 the disutility of the farmer tasks and positive utility for payment.

474 For estimation we employed a “Mixed Logit” specification. This assumes that each person has a
 475 utility function that can be expressed as a linear function of the attributes of delivered to respondents
 476 (the farmers) where the attributes are the services performed by farmers along with the payment for
 477 performing that service.

478 Formally, we assume that a person j ($j=1,\dots,J$) obtains utility U_{ijs} from choice i from a set of
 479 alternatives ($i=1,\dots,I$) within task set s ($s=1,\dots,S$). The utility function is then specified as:

$$U_{ijs} = \gamma_j(p_{ijs} - \beta_j'x_{ijs}) + \varepsilon_{ijs} = V_{ijs} + \varepsilon_{ijs} \quad (A1)$$

480 where ε_{ijs} is the unobserved random error which is assumed to be extreme value (Gumbel)
 481 distributed, x_{ijs} is a vector of observed variables (the choice attributes) given to farmers, and p_{ijs} is
 482 the payment offered to each farmer to compensate them for delivering these services. Under this
 483 specification the probability of respondent j choosing the i th option within set s is:

$$\text{Prob}(i|j, s) = \exp(V_{ijs}) / \sum_i \exp(V_{ijs}) \quad (A2)$$

484 The likelihood function for estimation is then expressed as the product of these probabilities
 485 over all choices and individuals. Given this specification β_j represents the vector of “willingness-to-
 486 accept” for each individual farmer and is assumed vary within the population with density $f(\gamma_j, \beta_j | \theta)$
 487 where θ are the parameters that govern the distribution for the population. This form (equation 1) is
 488 known as a “willingness-to-accept space” representation because the parameters within β_j represent
 489 a ratio of the marginal disutility for performing a given service over the marginal utility of the
 490 payment, and these are estimated directly rather than being recovered ex-post from a linear in
 491 parameters specification. The value for each γ_j and β_j were recovered using a Bayesian approach to
 492 estimation which places prior distributions of the parameters by specifying the nature of $f(\gamma_j, \beta_j | \theta)$
 493 along with the a priori distribution for θ . The distributions for γ_j and β_j were specified as normal
 494 distributions truncated at zero so that all values were positive. The priors for the means of these
 495 variables were normal and the variances were gamma. These priors were set so as to allow relatively
 496 diffuse estimates and estimated using Hamiltonian MCMC using the program STAN. The
 497 documentation for STAN can be found <https://mc-stan.org/> which also contains further detail about
 498 Hamiltonian MCMC. We experimented with a variety of priors. The estimates produced in the paper
 499 were for relatively conservative priors in terms of producing conservative (i.e., lower) estimates
 500 willingness-to-accept. Thus, more diffuse priors lead to slightly larger levels of willingness-to-accept
 501 estimates than those produced in the paper.

502 Appendix B

503 To determine the wet-dry conversion factor that is needed for economic viability assessment,
 504 CPH (1kg) was dried on a cement base in sunshine over a period of seven days at ICCRI Jember in
 505 October 2018. The factors for consecutive days were 0.656 (Day 2), 0.499 (Day 3), 0.343 (Day 4), 0.314
 506 (Day 5), 0.225 (Day 6) and 0.212 (Day 7). Comparable factors for clone MCC02 at MARS Tarengge
 507 were 0.168, 0.161, 0.169 for harvests in November 2017, June 2018 and November 2018 respectively.
 508 It is recognised that the value will vary according to the condition of the crop at harvest, the cocoa
 509 variety used and the conditions of the drying process: a conversion factor of 0.22 was used in the
 510 economic viability assessment

511

512 Appendix C

513 *Nutrient Cycling Materials and Methods*

514 At the Tarengge site, the mean annual rainfall from 2012 to 2018 was 2798 mm, normally well
 515 distributed over the year. In the year November 2017 to October 2018 it was 3051mm, with a peak in
 516 June 2017 (424mm). The soil is a silt loam classified as Typic Dystrandept with pH (water) 5.02, 4.09%

517 SOM (loss on ignition), 1.53% total C, 0.169% total N and 13 mg/kg of extractable P (Mehlich III
 518 method) averaged over all plots and sampled at October 2017. The mineral fertilizers were applied
 519 in four portions per year, using a local phosphate rock, dolomite, “Nitrabor” (calcium nitrate with
 520 additional B) and KCl. The annual applications per tree were: Rock phosphate (680g), CaNO₃ (480g
 521 as “Nitrabor”), KCl (280g) and dolomitic lime (680g).

522 Cocoa pods were harvested every two weeks and to simplify assessment of yields, this is done
 523 routinely by counts of the number of pods. For this work, the harvested pods were also weighed for
 524 three consecutive harvesting occasions during the peak production seasons of June–July 2018 and
 525 October–December, 2017 and 2018. A sample of 12 pods was taken, split into beans and the remainder
 526 (CPH), and CPH chopped into small (<2cm) pieces. All plant samples were air-dried at Tarengge
 527 under freely circulating air at ambient temperature until touch-dry. To estimate the bean and CPH
 528 production for the other harvests, the pods count was used in conjunction with the weight of dry
 529 bean and CPH per pod for the sampled pods. The proportion of the annual dry matter total harvest
 530 (from December 2017 to November 2018) accounted for by the actual measurements made in
 531 sampling periods was 48% and 59% for beans and CPH respectively. Litter and prunings samples
 532 were collected as described in Methods of the main section. There were few significant differences
 533 between months in nutrient concentrations, so appropriate mean values were used to estimate
 534 nutrients in litter, when necessary.

535 The air dried samples were analysed at the University of Reading, where most were further
 536 dried at 70 °C for 48 h and milled to pass 0.5 mm. Because of their high fat content, raw beans cannot
 537 be reduced to a powder by conventional mills, so they were first broken by hand in a ceramic pestle
 538 and mortar to pass 4mm sieve, then further reduced in a blender (Waring 8010EB, with 250 mL
 539 container). Total C and N were determined by a combustion analyser (Thermo Flash 2000). The total
 540 concentrations of Ca, K, Mg and P were determined by dissolution in nitric acid followed by
 541 inductively-coupled plasma atomic emission spectroscopy (Perkin-Elmer PE7300).

542 The peak of pod production in the October–November harvest season was later in 2018 than
 543 2017 (Fig 3.) and this means that any period of exactly 12 months might underestimate the typical
 544 total harvest. Data recording and sampling met with difficulties regarding quality at the start of the
 545 2017 sampling season. Therefore, to construct annual nutrient balances, the estimates of production
 546 were based on the year 29 November 2017 to 28 November 2018. This is a little different from
 547 November 2017 to November 2018 used for litter, but the annual litter fall was not greatly altered by
 548 the particular 12 month period assessed.

549 *Litter Contributions by Cocoa and Shade Trees*

550 Litter fall varied by a factor of up to three between months (Figure 3), although not in any clear
 551 association with weather variations. Cocoa litter is shed more in dry seasons, eg. in Ghana where
 552 several months each year experience <50mm rainfall (Dawoe et al., 2010). However, in our
 553 experimental period, no calendar month received <110mm and 12 of the 16 received >220mm, and no
 554 period without rain lasted over 13 days, so water was never deficient.

555 The total annual litter dry matter (3.9 t/ha) is low relative to typical litter fall in shaded cocoa,
 556 usually between 5 and 10 t/ha (Dawoe et al., 2010; Fontes et al., 2014; Hartemink 2005). It was noted
 557 that there appears to be an upward trend over the 14 months period of measurement (Figure 3). The
 558 increase in litter fall suggests that litter production was still limited by the young age of this trial and
 559 might become even more important in a more mature plantation.

560 **Table C1.** C and nutrients (kg/ha) added to soil via CPH and litter from cocoa and shade trees,
 561 between mid-May and mid-October 2018.

	C	N	P	K	Ca	Mg
Cocoa litter	332	9.60	0.78	3.88	18.28	4.27
Shade litter	505	19.88	1.33	6.35	13.80	3.42
CPH	301	6.71	0.69	28.46	4.09	2.46
(SD)	160	5.6	0.32	3.55	4.67	1.17

562 Much cocoa is grown with shade trees. Their litter adds to the internal nutrient cycle and the
 563 annual dry matter contribution may be around 2–3 t/ha (Aranguren et al., 1982; Fontes et al., 2014),
 564 constituting 13 to 60% of the total litter. Here, shade trees provided 58% of the total dry litter mass in
 565 the five months from mid-May to mid-October 2018. 60, 67, 63, 62, 43 and 45% of the C, N, P, K, Ca
 566 and Mg respectively in the litter came from the shade trees (

567), showing their value for maintaining soil fertility. The contributions of cocoa and shade litter
 568 were approximately equal, being not significantly different for any nutrient. Over the whole year,
 569 CPH provided the majority of the recycled K, but significantly less Ca than either litter. The mean N
 570 concentration of litter was significantly ($P < 0.001$) higher for shade (1.83%) compared to cocoa (1.19%).
 571 The former will in part be because N-fixing trees (*Glyricidia*) were the majority of the planted shade
 572 trees. The role of shade trees in the nutrient cycling of cocoa production deserves further
 573 investigation.

574 References

- 575 1. Babulo, B.; Muys, B.; Nega, F.; Tollens, E.; Nyssen, J.; Deckers, J.; Mathijs, E. Household livelihood strategies
 576 and forest dependence in the highlands of Tigray, Northern Ethiopia. *Agric. Syst.* **2008**, *98*, 147–155.
- 577 2. Cavendish, W. Empirical regularities in the poverty–environment relationship of rural households: Evi-
 578 dence from Zimbabwe. *World Dev.* **2000**, *28*, 1979–2003.
- 579 3. Davis, B.; Winters, P.; Carletto, G.; Covarrubias, K.; Quin õnes, E.J.; Zezza, A.; Digiuseppe, S. A cross-coun-
 580 try comparison of rural income generating activities. *World Development*, *38*, 48–63. *World Dev.* **2010**, *38*,
 581 48–63.
- 582 4. Ellis, F. Household strategies and rural livelihood diversification. *J. Dev. Stud.* **1998**, *35*, 1–38.
- 583 5. de Sherbinin, A.; VanWey, L.K.; McSweeney, K.; Aggarwal, R.; Barbieri, A.; Henry, S.; Hunter, L.M.; Twine,
 584 W.; Walker, R. Rural household demographics, livelihoods and the environment. *Glob. Environ. Chang.*
 585 **2008**, *18*, 38–53, doi:10.1016/j.gloenvcha.2007.05.005.
- 586 6. Garcia-Garcia, G.; Stone, J.; Rahimifard, S. Opportunities for waste valorisation in the food industry – A
 587 case study with four UK food manufacturers. *J. Clean. Prod.* **2019**, *211*, 1339–1356, doi:10.1016/J.JCLE-
 588 PRO.2018.11.269.
- 589 7. IES Feeding the nine billion; 2017.
- 590 8. Nasirumbi Sanya, L.; Birungi Kyazze, F.; Sseguya, H.; Kibwika, P.; Baguma, Y. Complexity of agricultural
 591 technology development processes: Implications for uptake of new hybrid banana varieties in Central
 592 Uganda. *Cogent Food Agric.* **2017**, *3*, 1–18, doi:10.1080/23311932.2017.1419789.
- 593 9. Fei, L.; Rodriguez-Garcia, Julia Van Damme, I.; Westwood Nicholas J., Shaw, L.; Robinson, James S. War-
 594 ren, G.; Chatzifragkou, Afroditi McQueen Mason, S.; Gomez, L.; Faas, L.; Balcombe, K.; Srinivasan, C.; Pic-
 595 chioni, F.; et al. Valorisation strategies for cocoa pod husk and its fractions. *Green Sustain. Chem.* **2018**, *14*,
 596 80–88, doi:10.1016/j.cogsc.2018.07.007.
- 597 10. Garb, Y.; Friedlander, L. From transfer to translation: Using systemic understandings of technology to un-
 598 derstand drip irrigation uptake. *Agric. Syst.* **2014**, *128*, 13–24, doi:10.1016/j.agry.2014.04.003.
- 599 11. Hartemink, A.E. Nutrient Stocks, Nutrient Cycling, and Soil Changes in Cocoa Ecosystems: A Review. *Adv.*
 600 *Agron.* **2005**, *86*, 227–253.
- 601 12. van Vliet, J.A.; Giller, K.E. Mineral Nutrition of Cocoa: A Review. *Adv. Agron.* **2017**, *141*, 185–270,
 602 doi:10.1016/BS.AGRON.2016.10.017.
- 603 13. Turmel, M.S.; Speratti, A.; Baudron, F.; Verhulst, N.; Govaerts, B. Crop residue management and soil health:
 604 A systems analysis. *Agric. Syst.* **2015**, *134*, 6–16, doi:10.1016/j.agry.2014.05.009.
- 605 14. Fontes, A.G.; Gama-Rodrigues, A.C.; Gama-Rodrigues, E.F.; Sales, M.V.S.; Costa, M.G.; Machado, R.C.R.
 606 Nutrient stocks in litterfall and litter in cocoa agroforests in Brazil. *Plant. Soil* **2014**, *383*, 313–335,
 607 doi:10.1007/s11104-014-2175-9.
- 608 15. FAO Cocoa production statistics Available online: <http://www.fao.org/faostat/en/#home> (accessed on Apr
 609 21, 2019).
- 610 16. Witjaksono, J. Cocoa Farming System in Indonesia and Its Sustainability Under Climate Change. *Agric. For.*
 611 *Fish.* **2016**, *5*, 170, doi:10.11648/j.aff.20160505.15.

- 612 17. Hoffmann, M.P.; Cock, J.; Samson, M.; Janetski, N.; Janetski, K.; Rötter, R.P.; Fisher, M.; Oberthür, T. Ferti-
613 lizer management in smallholder cocoa farms of Indonesia under variable climate and market prices. *Agric.*
614 *Syst.* **2020**, *178*, 102759, doi:10.1016/j.agry.2019.102759.
- 615 18. Wood, G.A.; Lass, R.A. *Cocoa*; 4th Editio.; Longman: London, 1986;.
- 616 19. Hertel, D.; Harteveld, M.A.; Leuschner, C. Conversion of a tropical forest into agroforest alters the fine
617 root-related carbon flux to the soil. *Soil Biol. Biochem.* **2009**, *41*, 481–490, doi:10.1016/j.soilbio.2008.11.020.
- 618 20. Sokol, N.W.; Kuebbing, S.E.; Karlsen-Ayala, E.; Bradford, M.A. Evidence for the primacy of living root
619 inputs, not root or shoot litter, in forming soil organic carbon. *New Phytol.* **2019**, *221*, 233–246,
620 doi:10.1111/nph.15361.
- 621 21. Xu, S.; Liu, L.L.; Sayer, E.J. Variability of above-ground litter inputs alters soil physicochemical and biolog-
622 ical processes: A meta-analysis of litterfall-manipulation experiments. *Biogeosciences* **2013**, *10*, 7423–7433,
623 doi:10.5194/bg-10-7423-2013.
- 624 22. Leff, J.W.; Wieder, W.R.; Taylor, P.G.; Townsend, A.R.; Nemergut, D.R.; Grandy, A.S.; Cleveland, C.C. Ex-
625 perimental litterfall manipulation drives large and rapid changes in soil carbon cycling in a wet tropical
626 forest. *Glob. Chang. Biol.* **2012**, *18*, 2969–2979, doi:10.1111/j.1365-2486.2012.02749.x.
- 627 23. Dawoe, E.K.; Isaac, M.E.; Quashie-Sam, J. Litterfall and litter nutrient dynamics under cocoa ecosystems in
628 lowland humid Ghana. *Palnt Soil* **2010**, *330*, 55–64, doi:10.1007/s11104-009-0173-0.
- 629 24. Tondoh, J.E.; Kouamé, F.N. guessa.; Martinez Guéi, A.; Sey, B.; Wowo Koné, A.; Gnessougou, N. Ecological
630 changes induced by full-sun cocoa farming in Côte d’Ivoire. *Glob. Ecol. Conserv.* **2015**, *3*, 575–595,
631 doi:10.1016/j.gecco.2015.02.007.
- 632 25. Veldkamp, E.; Purbopuspito, J.; Corre, M.D.; Brumme, R.; and Murdiyarto, D. Land use change effects on
633 trace gas fluxes in the forest margins of Central Sulawesi, Indonesia. *J. Geophys. Res.* **2008**, *113*.
- 634 26. Köhler, S.; Jungkunst, H.F.; Gutzler, C.; Herrera, R.; Gerold, G. Atmospheric ionic deposition in tropical
635 sites of central sulawesi determined by ion exchange resin collectors and bulk water collector. *Water. Air.*
636 *Soil Pollut.* **2012**, *223*, 4485–4494, doi:10.1007/s11270-012-1211-8.
- 637 27. Petry, J.F.; Sebastião, S.A.; Martins, E.G.; Barros, P.B. de A. Innovation and the Diffusion of Technology in
638 Agriculture in Floodplains in the State of Amazonas. *RAC - Rev. Adm. Contemp. (Journal Contemp. Adm.*
639 **2019**, *23*, 619–635.
- 640 28. Hall, A.; Bockett, G.; Taylor, S.; Sivamohan, M.V.; Clark, N. Why Research Partnerships Really Matter:
641 Innovation Theory, Institutional Arrangements and Implications for Developing New Technology for the
642 Poor. *World Dev.* **2001**, *29*, 783–797, doi:10.1016/S0305-750X(01)00004-3.
- 643 29. Klerkx, L.; Leeuwis, C. Establishment and embedding of innovation brokers at different innovation system
644 levels: Insights from the Dutch agricultural sector. *Technol. Forecast. Soc. Change* **2009**, *76*, 849–860,
645 doi:10.1016/j.TECHFORE.2008.10.001.
- 646 30. van Mierlo, B.; Leeuwis, C.; Smits, R.; Woolthuis, R.K. Learning towards system innovation: Evaluating a
647 systemic instrument. *Technol. Forecast. Soc. Change* **2010**, *77*, 318–334, doi:10.1016/j.techfore.2009.08.004.
- 648 31. Pigford, A.A.E.; Hickey, G.M.; Klerkx, L. Beyond agricultural innovation systems? Exploring an agricul-
649 tural innovation ecosystems approach for niche design and development in sustainability transitions.
650 *Agric. Syst.* **2018**, *164*, 116–121, doi:10.1016/j.agry.2018.04.007.
- 651



20 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).