UNIVERSITY of York

This is a repository copy of *Quadrupole deformation of* \$^130Xe\$ measured in a Coulombexcitation experiment.

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/167905/</u>

Version: Published Version

Article:

Morrison, L., Hadyifmmode nelse ńska-Klifmmode ęelse ęk, K., Podolyák, Zs. et al. (38 more authors) (2020) Quadrupole deformation of \$^130Xe\$ measured in a Coulombexcitation experiment. Phys. Rev. C. 054304. ISSN 2469-9993

https://doi.org/10.1103/PhysRevC.102.054304

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here: https://creativecommons.org/licenses/

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/

Quadrupole deformation of ¹³⁰Xe measured in a Coulomb-excitation experiment

L. Morrison,¹ K. Hadyńska-Klęk,^{2,1} Zs. Podolyák,¹ D. T. Doherty,¹ L. P. Gaffney,^{3,4} L. Kaya,⁵ L. Próchniak,² J. Samorajczyk-Pyśk,² J. Srebrny,² T. Berry,¹ A. Boukhari,⁶ M. Brunet,¹ R. Canavan,^{1,7} R. Catherall,⁴ S. J. Colosimo,³ J. G. Cubiss,⁸ H. De Witte,⁹ Ch. Fransen,⁵ E. Giannopoulos,^{10,4} H. Hess,⁵ T. Kröll,¹¹ N. Lalović,¹² B. Marsh,⁴ Y. Martinez Palenzuela,^{4,9} P. J. Napiorkowski,² G. O'Neill,^{13,14} J. Pakarinen,^{10,15} J. P. Ramos,^{16,4} P. Reiter,⁵ J. A. Rodriguez,⁴ D. Rosiak,⁵ S. Rothe,⁴ M. Rudigier,¹ M. Siciliano,^{17,18} J. Snäll,¹² P. Spagnoletti,¹⁹ S. Thiel,⁵ N. Warr,⁵ F. Wenander,⁴ R. Zidarova,²⁰ and M. Zielińska¹⁷ ¹Department of Physics, University of Surrey, Guildford, GU2 7XH, United Kingdom ²Heavy Ion Laboratory, University of Warsaw, Pasteura 5A, 02-093 Warszawa, Poland ³Oliver Lodge Laboratory, University of Liverpool, Liverpool, L69 7ZE, United Kingdom ⁴CERN, Physics Department, 1211 Geneva 23, Switzerland ⁵IKP Köln, Zülpicher Strasse 77, 50937 Köln, Germany ⁶Université de Paris-Sud 11, 15 Rue Georges Clemenceau, 91400 Orsay, France ⁷National Physical Laboratory, Hampton Road, Teddington, TW11 0LW, United Kingdom ⁸Department of Physics, University of York, York, YO10 5DD, United Kingdom ⁹KU Leuven, Instituut voor Kern-en Stralingsfysica, B-3001 Leuven, Belgium ¹⁰Department of Physics, University of Jyväskylä, P.O. Box 35 (YFL), Jyväskylä, FI-40014, Finland ¹¹GSI Helmholtzzentrum für Schwerionenforschung, Planckstrasse 1, 64291 Darmstadt, Germany ¹²Physics Department, Lund University, Box 118, Lund SE-221 00, Sweden ¹³Department of Physics and Astronomy, University of the Western Cape, P/B X17, Bellville, ZA-7535, South Africa ¹⁴iThemba LABS, Old Faure Road, Faure, Cape Town, 7131, South Africa ¹⁵Helsinki Institute of Physics, University of Helsinki, P.O. Box 64, FIN-00014, Helsinki, Finland

Helsinki Institute of Physics, University of Helsinki, P.O. Box 64, FIN-00014, Helsinki, Finlan

¹⁶SCK CEN, Boeretang 200, 2400 Mol, Belgium

¹⁷IRFU, CEA, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France

¹⁸INFN Laboratori Nazionali di Legnaro, 35020 Legnaro (Pd), Italy

¹⁹School of Computing, Engineering, and Physical Sciences, University of the West of Scotland, Paisley PA1 2BE, United Kingdom

²⁰Institut für Kernphysik, Technische Universität Darmstadt, 64289 Darmstadt, Germany

(Received 5 July 2020; accepted 24 September 2020; published 3 November 2020)

Low-lying states in the isotope ¹³⁰Xe were populated in a Coulomb-excitation experiment performed at CERN's HIE-ISOLDE facility. The magnitudes and relative signs of seven *E*2 matrix elements and one *M*1 matrix element coupling five low-lying states in ¹³⁰Xe were determined using the semiclassical coupled-channel Coulomb-excitation least-squares search code GOSIA. The diagonal *E*2 matrix elements of both the 2_1^+ and 4_1^+ states were extracted for the first time. The reduced transition strengths are in line with those obtained from previous measurements. Experimental results were compared with the general Bohr Hamiltonian with the microscopic input from mean-field theory utilizing universal nuclear energy density functional (UNEDF0), shell-model calculations using the GCN50:82 and SN100PN interactions, and simple phenomenological models (Davydov-Filippov and γ -soft). The extracted shape parameters indicate triaxial-prolate deformation in the ground-state band. In general, good agreement between theoretical predictions and experimental values was found, while neither phenomenological model was found to provide an adequate description of ¹³⁰Xe.

DOI: 10.1103/PhysRevC.102.054304

I. INTRODUCTION

Atomic nuclei often exhibit collective properties, which indicate that when a large number of nucleons are brought together, they are able to form systems acting more like macroscopic objects than ones composed of individual particles. This naturally leads to the idea of a nuclear shape, the determination of which is a key area of nuclear structure research. While nuclei close to magic numbers are known to be spherical, quadrupoledeformed shapes generally prevail throughout the nuclear chart. In particular, studies of how both axially symmetric and axially asymmetric quadrupole shapes evolve across isotopic chains provide key tests of nuclear structure models.

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

After tin, the xenon isotopic chain, with neutron numbers from N = 70 through to N = 82 (^{124–136}Xe), represents the longest chain of stable isotopes within the nuclear landscape. As such, it is an excellent testing ground for detailed studies of the evolution of nuclear collectivity (and hence deformation) as a function of neutron number. Indeed, the properties of the stable Xe isotopes have been the subject of a number of both experimental and theoretical studies over recent decades. Such studies indicate that while 136 Xe (N = 82) appears to be spherical in nature, the structure of the lighter stable Xe isotopes may be strongly influenced by the triaxial (or γ) degree of freedom [1,2]. Firm conclusions are, however, significantly hindered by the lack of detailed knowledge on both spectroscopic quadrupole moments and on the relative signs of electromagnetic matrix elements, which are critical for evaluating, in a model-independent way, the role of the triaxial degree of freedom in the Xe isotopes. Notably, spectroscopic quadrupole moments have been measured only for the ground states of even-odd $I \neq \frac{1}{2}$ Xe isotopes, as well as for some isomeric states in both odd- and even-mass Xe nuclei; however, none have previously been reported for short-lived excited states in the Xe isotopes. Thus, the even-even Xe nuclei are the only stable $A \approx 130$ isotopes where experimental spectroscopic quadrupole moments and the relative signs of electromagnetic matrix elements between low-lying states are unknown [3].

Understanding the evolution of collectivity along an isotopic chain and, in particular, describing the structure of transitional nuclei, has long proved to be a challenge for nuclear theory and nuclear structure models. For the even-even Xe isotopes, past work has focused on calculations employing the general Bohr Hamiltonian [1,4–6], the nuclear shell model [7], and the interacting boson model (IBM) [8,9], where the light Xe isotopes have been suggested to represent a transition point between spherical vibrators and γ -soft rotors, e.g., Refs. [10,11]. In addition, odd-mass nuclei in this region were explored in detail in recent mapped-IBM calculations [8,9].

Knowledge of this low-lying, low-spin structure may also have important consequences beyond the regime of traditional low-energy nuclear physics. The nucleus ¹³⁰Xe is the daughter of a key neutrinoless double-*beta* ($0\nu\beta\beta$) candidate ¹³⁰Te, with searches for this decay taking place in experiments around the world, e.g., Ref. [12]. As discussed in Ref. [13], $0\nu\beta\beta$ decay is expected to take place between initial and final states of similar structure and shape and, consequently, knowledge of low-lying 0⁺ states in ¹³⁰Xe is crucial for informing this work.

Safe-energy Coulomb excitation is an experimental method which is able to provide information on the electromagnetic structure of the investigated nucleus by extracting both transitional and diagonal matrix elements, together with their relative signs. Those can be used to determine the shape of individual states using the quadrupole sum rules method and give unique insight into the properties of a given nucleus. Furthermore, the extracted transition probabilities $B(_M^E \lambda)$ provide direct information on the collectivity of the transitions in the investigated nucleus. In this paper, the results of a recent Coulomb-excitation experiment populating states in ¹³⁰Xe are presented. The collected data allow for the first examination

of the deformation of the 0^+ ground state and the first excited 2^+ state in ¹³⁰Xe. Furthermore, the extracted set of matrix elements together with their relative signs provides, for the first time, a solid base for an in-depth understanding of the structure of ¹³⁰Xe at low excitation energy.

This paper is organized as follows: The experimental details are presented in Sec. II, and the Coulomb-excitation data analysis and experimental results are described in Sec. III. In Sec. IV, the theoretical approaches and interpretation are presented together with the quadrupole sum rules analysis.

II. EXPERIMENTAL DETAILS

A Coulomb-excitation experiment, primarily designed to investigate the singly magic, two proton-hole nucleus ²⁰⁶Hg, was performed using the MINIBALL setup at the high intensity and energy isotope separator on line device (HIE-ISOLDE) facility, conseil européen pour la Recherche Nucléaire (CERN). The radioactive ²⁰⁶Hg beam was found to be contaminated with the isotope of 130 Xe, the origin of which can be attributed to the presence of atomic xenon leaking in from Radioactive beam EXperiment (REXTRAP), the xenon having been ionized by charge transfer from the incoming radioactive beam [14]. Therefore, it is not related to the combination of beam and target used in the primary experiment. In order to study the effect of this contaminant, several experimental runs were taken with the lasers responsible for ionizing ²⁰⁶Hg in the ion source turned off, as well as the lasers used within the resonance ionization laser ion source, and as such

within the resonance ionization laser ion source, and as such the analysis presented here is solely focused on ¹³⁰Xe. The beam composition was checked with an ionization chamber, confirming no contaminants were present. For more details, see Ref. [15].

The 130 Xe beam ($Q = 29^+$, intensity $\approx 3 \times 10^5$ pps), was accelerated using the newly commissioned HIE-ISOLDE linear accelerator [16,17], to an energy of 4.195 MeV/u. Post acceleration, it impinged on a 2 mg/cm² thick 94 Mo target for a total of approximately 7.4 h. The beam energy fulfilled Cline's safe distance criterion [18], ensuring a purely electromagnetic interaction between the collision partners took place.

Following Coulomb excitation, both the projectile and recoiling target-like particles were detected using an annular double-sided silicon strip detector (DSSSD), divided into 16 annular strips on the front side, and a further 24 sectors per individual quadrant, which are coupled into 12 pairs when read out [19,20]. This high degree of segmentation allowed accurate and precise position identification of the scattered particles.

The energy calibration of the DSSSD detector was carried out for each silicon strip using accelerated stable beams of 12 C, 16 O, 20 Ne, and 40 Ar. Additionally, the detector distance from the target, 24.6 mm, was established from the count-rate distribution in a given strip coming from the measured α decay of a 226 Ra source.

The reaction kinematics plot obtained using the DSSSD particle detector is presented in Fig. 1. The beam and target recoils are clearly separated for most of the angular range covered by the detector and were identified by comparing them



FIG. 1. The particle spectrum collected in the DSSSD detector, displaying the projectile and target kinematics for the Coulomb excitation of ¹³⁰Xe on a ⁹⁴Mo target. The red outline indicates the region gated on for target-like events.

with simulations performed using the KINSIM code [21]. The target particle-like events covered an angular range from 24 to 59° in the laboratory frame (LAB), equivalent to 62 to 131° in the center-of-mass system (c.m.). As presented in Fig. 1, the innermost strip of the DSSSD detector is excluded from the analysis because it was not possible to separate the projectile-and target-like events in this range. As the experiment was performed in inverse kinematics, the projectile kinematic plot has two solutions corresponding to two different ranges of the center-of-mass frame. However, the full range of projectile-like events is also covered in the target-like events gate, and as such the selection of the ⁹⁴Mo and ¹³⁰Xe kinematic solutions could be done in an unambiguous way.

De-excitation γ rays were detected by 23 of the total 24 HPGe detectors constituting the MINIBALL array [19]. Energy and efficiency calibrations for the high-purity germanium (HPGe) detectors were undertaken using standard radioactive ¹⁵²Eu and ¹³³Ba sources. The γ -ray efficiencies were determined using the γ -ray efficiency code GREMLIN [22]. In order to apply the appropriate Doppler correction, each crystal's precise alignment and positioning was determined from Doppler-shifted γ rays following the (d, p) and (d, n)reactions of a stable ²²Ne beam. For more details on the experimental setup and analysis, see Ref. [23].

Both one- and two-particle events were included during the data sort given the kinematics of the reaction, allowing for both the target recoil and scattered beam particles to be detected in the angular range covered by the downstream DSSSD detector. In order to select γ rays originating from the excitation of either ¹³⁰Xe or ⁹⁴Mo, only γ -ray events coincident with a particle were registered, and this particle- γ coincidence was subject to time-gating conditions, based on prompt and random events. As seen in Fig. 2, the prompt window was set from -300 to 150 ns, and the random window from 250 to 850 ns. These conditions suppress γ -ray background events associated with activity in the chamber and allow for a subtraction of random coincidences.

By performing the appropriate Doppler correction for projectile velocity, γ -ray spectra (recoil-gated Doppler-corrected for projectile-like particles) were produced, as shown in Fig. 3. A number of γ -ray peaks associated with the excitation of ¹³⁰Xe [24] are visible in the spectrum. Figure 4 displays a



FIG. 2. A particle- γ coincidence plot showing the time-gating conditions used in the experiment. The peak displays events fulfilling the particle- γ coincidence condition. The selected prompt window was 450 ns wide (shown in gray), and the random window was set to 600 ns (shown in red).

partial level scheme for 130 Xe with transitions relevant to the present study included. The γ -ray transitions observed in the present experiment are marked in red.

III. COULOMB-EXCITATION DATA ANALYSIS

The Coulomb-excitation data were analyzed with the semiclassical coupled-channel Coulomb-excitation least-squares search code GOSIA [18,25], in order to extract matrix elements for the observed transitions together with their relative signs and associated uncertainties.

Two normalization methods were used in the Coulombexcitation analysis. In the first, normalization to the known target properties was performed, as discussed in detail in, e.g., Ref. [26]. Here, this analysis was used only for the first 2^+_1



FIG. 3. The particle-particle- γ coincidence spectrum collected in the Coulomb excitation of a ¹³⁰Xe beam on a ⁹⁴Mo target experiment, gated on the recoil-like particles, Doppler corrected for the ¹³⁰Xe projectile velocity. All observed transitions in the Coulombexcited ¹³⁰Xe are labeled. The energy region in the spectrum between 800 and 950 keV corresponds to the Doppler-broadened $2_1^+ \rightarrow 0_1^+$ transition at 871 keV in the ⁹⁴Mo target.



FIG. 4. Low-lying excited states in ¹³⁰Xe, considered in the present analysis. Transitions observed in the current experiment with a ⁹⁴Mo target are marked in red. Level and transition energies are given in keV.

state in order to extract the $B(E2; 2_1^+ \rightarrow 0_1^+)$ value in an independent way. The collected data were subdivided into a total of six angular ranges (see Table I): five distinct ones spanning the entire range of angles upon summation, and one total range covering the entire selection of angles at once. The number of subdivisions was limited in order to ensure sufficient statistics in the $2^+_1 \rightarrow 0^+_1$ transitions in both ^{130}Xe and ⁹⁴Mo, and each range was taken with weight 0.5, ensuring every event was only counted once. In ⁹⁴Mo, both the reduced transition strength ($B(E2; 2_1^+ \to 0_1^+) = 16.0(4)$ W.u.), and the spectroscopic quadrupole moment of the 2^+_1 state $[Q_s(2^+_1) = -0.13(8) \text{ eb}]$, are known to good precision [27], and consequently, this information could be used to normalize data from the different angular ranges. The $B(E2; 2^+_1 \rightarrow$ $0^+{}_1)$ value for the 130 Xe beam could then be extracted from the two-dimensional χ^2 surface map, calculated using the GOSIA2 program together with a specially developed χ^2 surface code [28], by performing a minimization with respect to the $|\langle 2_1^+ || E2 || 0^+_1 \rangle|$ and $|\langle 2_1^+ || E2 || 2^+_1 \rangle|$ matrix elements. This analysis yielded a $B(E2; 2_1^+ \rightarrow 0_1^+) = 30(+2, -7)$ W.u., in good agreement with previous work [3,11,29-31,33], but with a significant uncertainty on the extracted quadrupole moment.

In addition to the analysis performed on the beam particles, an additional analysis was carried out for the target peaks, by Doppler correcting for ⁹⁴Mo. A significant amount of ⁹⁵Mo was discovered to be present in the target, accounting for 5(1)% of the total statistics. Additional excitation due to this target contamination was accounted for by increasing the uncertainty values for all γ -ray yields by 5%, when input into GOSIA.

The analysis described above gives confidence in the reported literature values for the $2^+_1 \rightarrow 0^+_1$ transition probability. Consequently, here, in order to extract a full set of matrix elements from the experimental data using the standard GOSIA code, the normalization of the data was performed with the lifetime of the 2^+_1 state. The most recent and precise lifetime value of $\tau = 14.7(3)$ ps [29,30] was chosen, which

TABLE I. Numbers of counts in the observed γ -ray transitions in ¹³⁰Xe used in the presented data analysis. In addition, the number of counts in the observed $2_1^+ \rightarrow 0_1^+ \gamma$ -ray transition in ⁹⁴Mo (871 keV), for different angular ranges used in the target normalization approach analysis described in the text, is included.

		130	¹³⁰ Xe		Mo
I_i^{π}	I_f^π	$\overline{E_{\gamma}}$ (keV)	Counts	$\overline{E_{\gamma}}$ (keV)	Counts
		Total spe	ctrum ($\theta_{c.m.} = 6$	2–131°)	
2^{+}_{1}	0_{1}^{+}	536	5710(286)	871	1720(43)
4_{1}^{+}	2_{1}^{+}	669	900(45)		
6_{1}^{+}	4_{1}^{+}	739	151(22)		
2^{+}_{2}	2_{1}^{+}	586	298(24)		
2^{+}_{2}	0_{1}^{+}	1122	51(9)		
		θ_{c}	$m_{m.} = 111 - 131^{\circ}$		
2^{+}_{1}	0_{1}^{+}	536	754(29)	871	243(17)
4_{1}^{+}	2_{1}^{+}	669	195(15)		
6_{1}^{+}	4_{1}^{+}	739	39(9)		
2^{+}_{2}	2_{1}^{+}	586	81(11)		
		θ	$_{c.m.} = 94-110^{\circ}$		
2^{+}_{1}	0_{1}^{+}	536	1294(65)	871	375(21)
4_{1}^{+}	2^{+}_{1}	669	260(18)		
2^{+}_{2}	2^{+}_{1}	586	81(12)		
		($\theta_{\rm c.m.} = 85 - 93^{\circ}$		
2^{+}_{1}	0_{1}^{+}	536	1018(51)	871	224(16)
4_{1}^{+}	2^{+}_{1}	669	198(16)		
2^{+}_{2}	2_{1}^{+}	586	63(10)		
		($\theta_{\rm c.m.} = 74 - 84^{\circ}$		
2^{+}_{1}	0_{1}^{+}	536	1039(52)	871	277(18)
4_{1}^{+}	2^{+}_{1}	669	121(13)		
2^{+}_{2}	2_{1}^{+}	586	44(10)		
		($\theta_{\rm c.m.} = 62 - 73^{\circ}$		
2^{+}_{1}	0_{1}^{+}	536	1540(77)	871	420(21)
4_{1}^{+}	2^{+}_{1}	669	111(13)		
2^{+}_{2}	2_{1}^{+}	586	45(11)		

corresponds to a B(E2) of 32(1) W.u. The additional precision of this value, compared to the one obtained in the GOSIA2 analysis, increases the sensitivity to second-order effects, and, as a result, improves the precision of the extracted matrix elements. The GOSIA analysis also incorporates other spectroscopic information such as γ -ray branching and E2/M1mixing ratios, and the lifetimes of low-lying states (see Table II). This information serves to constrain the final GOSIA analysis and enters the multidimensional χ^2 function fit in the same way as the measured γ -ray intensities. Two further, unobserved, 2⁺ states, at excitation energies of 2016 and 2150 keV, were also included in the GOSIA analysis. This was in order to account for the possible excitation of these unobserved levels in the measured γ -ray yields. The known spectroscopic data for these states were included (see Table II), as well as the known matrix elements, from Ref. [11], to serve as additional constraints. However, in the analysis, the unknown Q_s of these two states were set to 0.

TABLE II. The γ -ray branching ratios, mixing ratios, δ , for mixed E2/M1 transitions, and the lifetimes of the excited states in ¹³⁰Xe used as additional data points in the Coulomb-excitation data analysis. The $2_2^+ \rightarrow 2_1^+$ and $4_2^+ \rightarrow 4_1^+$ mixing ratio values are from Refs. [34,35], and the γ -ray branching ratios were calculated using data from Ref. [3]. The lifetimes are from Refs. [29,30].

I_i^{π}	I_j^π	E_{γ} (keV)	Branching ratio
$\frac{1}{2^{+}_{2}}$	2^{+}_{1}	586	1
$2^{\tilde{+}}_{2}$	0_{1}^{+}	1122	0.15(1)
$4^{\tilde{+}}_{2}$	2^{+}_{2}	686	1
$4^{\bar{+}}_{2}$	4_{1}^{+}	603	0.58(5)
$4^{\bar{+}}_{2}$	2^{+}_{1}	1272	0.70(6)
2_{4}^{+}	0_{1}^{+}	2150	0.047(2)
2_{4}^{+}	2^{+}_{2}	1028	0.087(4)
2_{4}^{+}	2_{1}^{+}	1614	1
I_i^{π}	I_j^{π}	E_{γ} (keV)	$\delta(E2/M1)$
2^{+}_{2}	2^{+}_{1}	586	3.75(12)
$4^{\tilde{+}}_{2}$	4_{1}^{+}	603	2.4(+13,-7)
S	tate	Energy (keV)	τ (ps)
2	2^+_1	536	14.7(3)
4	l [†]	1205	3.3(2)
6	\tilde{b}_{1}^{+}	1944	1.9(7)
2	2+22	1122	6.5(6)

The data were again subdivided into six angular ranges (as presented in Table I), in order to exploit the scattering angle dependence of the Coulomb-excitation cross section and enable spectroscopic quadrupole moments to be extracted. A χ^2 fit of the observed γ -ray yields (Table I) and other spectroscopic data (Table II) was performed with 15 *E*2 and 2 *M*1 matrix elements. A sign convention was adopted where the signs of all in-band transitional *E*2 matrix elements were assumed to be positive, as well as that of the $\langle 0_1^+ || E2 || 2^+_2 \rangle$ transition; the signs of all other *E*2 matrix elements were then determined relative to those.

The statistical errors of the matrix elements were determined when convergence of the χ^2 minimization was achieved. This was performed in two steps. First, in order to find the "diagonal" uncertainty, the χ^2 surface is sampled in the vicinity of the minimum, using different values of the matrix element in question, with all other matrix elements remaining fixed. Second, in order to obtain the total statistical uncertainty, all of the possible correlations between matrix elements are accounted for. As a result, a set of five E2 and one M1 transitional matrix elements was extracted, connecting five low-lying states in ¹³⁰Xe. Similarly, two spectroscopic quadrupole moments were determined, for the first time, from the extracted diagonal matrix elements. The experimental results are presented in Tables III and IV, where they are compared with both literature values and the results of theoretical calculations.

The set of reduced matrix elements obtained reproduces all of the lifetimes, branching, and mixing ratios presented in Table II within a 1σ uncertainty. Furthermore, good agreement is found between the reduced transition strength results obtained here and previously published results (Table III). We note that in addition to the B(E2) transition strengths, the $B(M1; 2_2^+ \rightarrow 2_1^+)$ value could also be extracted owing to the experimentally determined $\delta(E2/M1)$ mixing ratio [31] for this γ ray.

The spectroscopic quadrupole moments for the 2_1^+ and 4_1^+ states in ¹³⁰Xe were determined for the first time (Table IV). Both the 2_1^+ and 4_1^+ signs and values $[Q_s(2_1^+) = -0.38(+17, -14)$ e.b. and $Q_s(4_1^+) = -0.41(12)$ e.b.] indicate stable prolate deformation within the yrast band. Furthermore, these moments are reduced with respect to those calculated with a simple rotor model, fit to the experimental $B(E2; 2_1^+ \rightarrow 0^+_1)$ value, $[Q_s(2_1^+) = -0.7$ e.b., $Q_s(4_1^+) = -0.9$ e.b.], suggesting an influence from the γ degree of freedom. However, these moments, as well as the reduced transition probabilities in the ground-state band, are only weakly affected by γ deformation, so, in order to draw more definitive conclusions, a detailed theoretical analysis was undertaken.

IV. DISCUSSION

Attempts to describe the structure of the mass $A \approx 130$ nuclei theoretically invoke both single-particle shell-model calculations and beyond-mean-field approaches. Of these nuclei, the stable Xe isotopes are among the most challenging to describe, owing to the gradual shape transition from γ -soft structures in the lighter nuclei through to spherical ¹³⁶Xe at the N = 82 shell closure. For the odd-A isotopes, positive spectroscopic quadrupole moments have been reported for low-lying states in ^{133,135}Xe, following colinear laser spectroscopy experiments [36], indicating oblate deformation. This is in contrast to the lighter ¹³¹Xe isotope, where modest prolate deformation prevails [36].

Here, in order to aid the interpretation of the experimental results, theoretical calculations were performed with a meanfield formalism, which is well suited to describing nuclear collectivity, a large-scale shell-model approach, with two different interactions, as well as a simple Davydov Filippov (rigid triaxial) model [37], and a γ -soft model based on a Hamiltonian with γ -independent potential energy and a constant mass parameter [2,38]. In addition, the quadrupole sum rules method, which links *E*2 matrix elements to deformation parameters defined in the intrinsic frame of the nucleus, is applied in order to extract shape invariants (β , γ), allowing for conclusions on the shapes of low-lying states to be made and an in-depth comparison with theory.

A. General Bohr Hamiltonian based on microscopic mean-field theory

The first theoretical approach utilized (considering the effects of low-energy collectivity), can be described as applying the general Bohr Hamiltonian (GBH) with the microscopic input from the mean-field theory utilizing the universal nuclear energy density functional (UNEDF0) [39]. The method used to construct the collective Hamiltonian (GBH) is based on the adiabatic time-dependent Hartree-Fock-Bogolyubov (HFB) (ATDHFB) theory. Details of the method can be

TABLE III. Reduced transition strengths, $B(E2; I_i^+ \to I_f^+)$ and $B(M1; I_i^+ \to I_f^+)$ in ¹³⁰Xe extracted from the present experiment. Previous experimental values are also given; these are based on lifetime measurements [3,29–31] and Coulomb-excitation experiments [11,33]. Theoretical values obtained with the collective GBH-UNEDF0 model as well as the GCN50:82 and SN100PN shell-model interactions are shown. For the shell-model theoretical calculations, different sets of effective charges were used, as indicated. For the $2^+_2 \to 2^+_1$ transition, the theoretical B(M1) values were obtained using effective g factor values of: $g_{l,p} = 1$, $g_{l,n} = 0$, $g_{s,p} = 3.91$, and $g_{s,n} = -2.68$. The last two columns contain the values from the Davydov-Filipov model (D-F) and γ -soft model.

			$\mathbf{B}(E2;I_i^+\to I_f^+)~(\mathrm{W.u.})^1$								
$\mathbf{I}_i^+ \to \mathbf{I}_f^+$	E_{γ} (keV)	$\langle I_i \ E2 \ I_f \rangle$ (eb)	Present	Previous	GBH-UNEDF0	GCN50:82 SN100PN		100PN	D-F	γ-soft	
$2^+_1 \rightarrow 0^+_1$	536	0.79(4)	32(3)	37.1(17) [31] 38(5) [3]	24	$e_{\pi} = 1.5e$ $e_{\nu} = 0.5e$ 21	$e_{\pi} = 1.53e$ $e_{\nu} = 0.945e$ 35	$e_{\pi} = 1.5e$ $e_{\nu} = 0.5e$ 20	$e_{\pi} = 1.68e$ $e_{\nu} = 0.84e$ 35	32	36
$4^+_1 \rightarrow 2^+_1$	669	1.29(6)	47(4)	30.0(+44, -28) [33] 33.2(26) [32] 32(1) [29,30] 44.5(20) [31] 46.4(46) [11] 47(6) [29,30]	46	32	51	31	52	46	61
$6^+_1 \rightarrow 4^+_1$	739	1.74(+21, -18)	60(-12, +14)	69(9) [11] 51(40) [30]	64	31	48	18	29	56	84
$2^+_2 \rightarrow 0^+_1$	1122	0.067(3)	0.23(2)	$\begin{array}{c} 0.24(2) [31] \\ 0.28(5) [11] \\ 0.25(5) [30] \end{array}$	0.30	0.10	0.40	0.10	0.20	2.00	0.02
$2^+_2 \rightarrow 2^+_1$	586	0.85(4)	37(3)	37.1(28) [31] 44.3(81) [11] 38(7) [30]	35	22	38	26	46	21	61
					$\mathbf{B}(M1; I_i^+ \to I_f^+) \; (\mathrm{W.u.})^1$						
$\mathbf{I}_i^+ \to \mathbf{I}_f^+$	E_{γ} (keV)	$\langle I_i \ M1 \ I_f \rangle(\mu)$	Present	Previous	GBH-UNEDF0	GCN	150:82	SN1	00PN	D-F	γ-soft
$2^+_2 \rightarrow 2^+_1$	586	0.11(1)	0.0014(2)	0.0014(2) [31]	$g_{l(p,n)} = 1, 0; g_{s(p,n)} = 3.91, -2.68$ 0.0004 0.0020						

¹For an *E*2 transition in ¹³⁰Xe, 1 W.u. = $3.912 \times 10^{-3} e^2 b^2$.

²For an *M*1 transition in ¹³⁰Xe, 1 W.u. = 1.79 μ^2 .

TABLE IV. Spectroscopic quadrupole moments, Q_s extracted for ¹³⁰Xe compared with theoretical calculations obtained with the collective GBH-UNEDF0 model, the GCN50:82 and SN100PN shell-model interactions, Davydov-Filipov model (D-F) and the γ -soft model. The used effective charges are indicated. $Q_s < 0$ indicates prolate deformation and $Q_s > 0$ indicates oblate deformation. The intrinsic quadrupole moments (Q_0), calculated from the diagonal matrix elements obtained in the present experiment assuming rigid axially symmetric rotor and K = 0, are given in the last column.

	Q_s (eb)										
Level	$\langle I E2 I \rangle$ (eb)	Present	GBH-UNEDF0 GCN50		50:82 SN100PN		00PN	D-F	γ-soft	Q ₀ (eb)	
				$e_{\pi} = 1.5e$	$e_{\pi} = 1.53e$	$e_{\pi} = 1.5e$	$e_{\pi} = 1.68e$				
				$e_v = 0.5e~e$	$v_{\nu} = 0.945\epsilon$	$e_{\nu}=0.5e$	$e_{\nu}=0.84e$				
2^+_1	-0.50(+22, -18)	-0.38(+17, -14)	-0.35	-0.35	-0.42	-0.20	-0.25	-0.55	-0.04	1.33(+60, -49)	
4_{1}^{+}	-0.55(16)	-0.41(12)	-0.47	-0.53	-0.64	-0.53	-0.64	-0.42	-0.07	1.13(33)	
2^+_2	0.1(1)	0.1(1)	0.31	0.34	0.41	0.20	0.24	0.55	0.02	-0.3(3)	

found in Refs. [1,40–42]. The GBH-UNEDF0 functional is a Skyrme-type "standard" functional in the particle-hole channel with the spin-orbit term taken as in the SkI parametrization [43], while the pairing interaction is a sum of the standard volume and density-dependent surface-peaked δ interaction. The fitting of proton and neutron pairing strengths is done simultaneously with other functional parameters. All these parameters are fixed for the whole nuclear table. The Lipkin-Nogami (LN) method is applied in order to avoid pairing for magic nuclei and their neighbors collapsing. It should be noted that in the GBH-UNEDF0 approach no fitting of the effective charges is performed.

The GBH-UNEDF0 calculations were performed for the even-even ¹¹⁸⁻¹⁴⁴Xe isotopes [1]. The low-energy spectra and $B(E2; 4^+ \rightarrow 2^+)$ transition probabilities were found to be in relatively good agreement with available experimental data, with the largest discrepancies around the semimagic ¹³⁶Xe nucleus. One should note that this was achieved after scaling all mass parameters by a factor of 1.3, needed mainly to reproduce the energy spectra (the effect on the transition strengths is much smaller). This is a common procedure; for its origin see Ref. [1] and references herein.

The GBH-UNEDF0 energy level scheme is compared with the experimental one in Fig. 6. The results of the GBH-UNEDF0 calculations for ¹³⁰Xe for transition strengths and quadrupole moments, are compared with experimental values in Tables III and IV, respectively [microscopic calculations for the *M*1 transitions are not yet implemented so no B(M1)values are determined]. Furthermore, it should be pointed out that the agreement with experimental values is very good for the quadrupole moments.

The potential-energy surface map for the ¹³⁰Xe ground state is shown in Fig. 5, which indicates a pronounced γ softness for β ranging from 0 up to 0.3 where the potential starts to increase.

The theoretical shape parameters from GBH-UNEDF0 calculations are further compared to the experimental ones in Table V.

B. Large-scale shell-model calculations

Large-scale shell-model calculations were performed with two different interactions: GCN50:82 [44] and SN100PN [45]. Both interactions make use of a valence space consisting of all proton and neutron orbitals between the magic shell number closures N = Z = 50 and N = Z = 82. Therefore, ¹⁰⁰Sn is considered to be an inert core. ¹³⁰Xe has a large number of valence particles, with four protons and six neutron holes relative to the Z = 50 and N = 82 shell closures.

The SN100PN interaction utilizes the jj55pna Hamiltonian [45] within the NUSHELLX@MSU [46] code. The Hamiltonian itself treats four types of interactions separately: neutron-neutron, neutron-proton, proton-proton, and Coulomb-proton repulsion. The two-body interaction is based on a renormalized *G* matrix (itself derived from the CD-Bonn nucleon-nucleon interaction). The single-particle proton and neutron energies are taken from the experimental levels of ¹³³Sb and ¹³¹Sn.

The GCN50:82 interaction [44] is also based on a realistic *G* matrix (derived from the CD-Bonn potential). However, two-body matrix elements were modified by normalizing to sets of experimental excitation energies in even-even and even-odd semimagic nuclei, even-odd Sb isotopes and N = 81 isotones, and some known odd-odd nuclei around ¹³²Sn. The calculations were undertaken using the KSHELL program [47].



FIG. 5. Potential energy surface map (relative to a spherical shape value) for the ground state of 130 Xe, computed using the GBH-UNEDF0 functional. A pronounced γ -soft shape is visible for β ranging from 0 up to 0.3.



FIG. 6. Experimental [3] and theoretical low-lying energy level schemes for ¹³⁰Xe. See the text for the details of the calculations.

Both shell-model approaches have successfully reproduced experimental results for a range of isotopes in this mass region, namely ^{135,136,137}Ba and ^{131,132,133,135}Xe [31,44,48–51]. In particular, the GCN50:82 interaction shows good agreement for the neighboring Xe isotopes ¹³¹Xe [48], ¹³²Xe [49], and ¹³³Xe [50].

The experimental and shell-model excited states for ¹³⁰Xe are compared in Fig. 6. There is a fair agreement; however, the experimental energies are typically lower than their theoretical counterparts, for both interactions. The higher theoretical values for the yrast 2⁺ and 4⁺ states probably indicate that in reality the wave functions are more fragmented. The higher experimental $E(4^+)/E(2^+) = 2.25$ ratio, compared to the theoretical ratios 2.12 (GCN50:82) and 2.17 (SN100PN), suggests the same.

The experimental transition strengths and spectroscopic quadrupole moments are compared with those of the shellmodel calculations in Tables III and IV. When using the standard effective charges of $e_{\nu} = 0.5e$ and $e_{\pi} = 1.5e$, the shell-model reduced transition strengths are too low. For example, the $B(E2; 2_1^+ \rightarrow 0_1^+)$ is 21 and 20 W.u. for the GCN50:82 and the SN100PN interactions, respectively, which are roughly a factor of 1.5 below the experimental value. Previous studies used larger effective charges for nuclei in this region of the nuclear chart. For example, transition strengths from isomeric decays in ¹²⁹Sn, ¹³¹Te, ¹³³Xe, ¹³⁵Ba were described using $e_{\pi} = 1.52e$ and $e_{\nu} = 0.81e$ in both the GCN50:82 and the SN100PN interactions [50]. The same effective charges were used for ¹³³Te, ¹³⁵Xe, and ¹³⁷Ba with the SN100PN interaction [52]. Here, for the case of ¹³⁰Xe, which has a larger number of valence nucleons than any of the aforementioned nuclei, we use slightly higher effective charges: $e_{\pi} = 1.53$ and $e_{\nu} = 0.945$ for the GCN50:82 interaction and $e_{\pi} = 1.68$ and $e_{\nu} = 0.84$ for the SN100PN interaction. These neutron charges were chosen to reproduce the $B(E2; 2_1^+ \rightarrow 0_1^+)$ value in ¹²⁶Sn TABLE V. Experimental and theoretical quadrupole shape invariants: $\langle Q^2 \rangle$ and $\langle \cos(3\delta) \rangle$ for 0_1^+ states in ¹³⁰Xe. The contribution of the individual matrix elements to the final values of $\langle Q^2 \rangle$ is shown. The $\langle Q^2 \rangle$ invariant is formed by summing all the contributions and by multiplying the sum by 1 and by -0.873 for $\langle \cos(3\delta) \rangle$ [41,57]. The loops containing transitional matrix elements are included twice in the contribution to the $\langle \cos(3\delta) \rangle$ invariant as their zero-coupled products can be formed in two possible ways. Experimental results not obtained in the present work are from Ref. [11], while the relative signs between matrix elements were adopted from the GBH calculations if they could not be determined in this analysis.

				GCI	N50:82	SN1		
		Experiment	GBH-UNDF0	$e_{\pi} = 1.5e$ $e_{\nu} = 0.5e$	$e_{\pi} = 1.53e$ $e_{\nu} = 0.945e$	$e_{\pi} = 1.5e$ $e_{\nu} = 0.5e$	$e_{\pi} = 1.68e$ $e_{\nu} = 0.84e$	γ-soft
State	Component $E2 \times E2$			Co {(ntribution to Q^2 (e^2 fm ⁴)			
	$\langle 0_{1}^{+} E2 2_{1}^{+} angle\langle 2_{1}^{+} E2 0_{1}^{+} angle$	6240	4750	4170	6760	4000	6850	6980
0_{1}^{+}	$\langle 0^+_1 E2 2^+_2 \rangle \langle 2_2^+ E2 0^+_1 \rangle$	45	50	20	80	10	30	5
	$\langle 0_1^+ E2 2_3^+ \rangle \langle 2_3^+ E2 0_1^+ \rangle$	20	2					40
	$\langle 0_1^+ E2 2_4^+ \rangle \langle 2_4^+ E2 0_1^+ \rangle$	45						0
	$\langle Q^2 angle$	6350(400)	4800	4190	6840	4000	6880	7030
	$\langle \beta \rangle$	0.17(2)	0.15	0.14	0.17	0.13	0.17	0.18
	$\langle 2_{1}^{+} E2 0_{1}^{+}\rangle\langle 0_{1}^{+} E2 2_{1}^{+}\rangle$	1250	950	830	1350	800	1370	1400
	$\langle 2_{1}^{+} E2 2_{2}^{+} \rangle \langle 2_{2}^{+} E2 2_{1}^{+} \rangle$	1440	1350	870	1500	1030	1800	2380
	$\langle 2_{1}^{+} E2 2_{3}^{+}\rangle\langle 2_{3}^{+} E2 2_{1}^{+}\rangle$	35	10					1
	$\langle 2_{1}^{+} E2 2_{4}^{+} \rangle \langle 2_{4}^{+} E2 2_{1}^{+} \rangle$	5						0
2^+_1	$\langle 2_{1}^{+} E2 4_{1}^{+}\rangle\langle 4_{1}^{+} E2 2_{1}^{+}\rangle$	3350	3200	2270	3580	2160	3630	4300
	$\langle 2_{1}^{+} E2 4_{2}^{+}\rangle\langle 4_{2}^{+} E2 2_{1}^{+}\rangle$	25	0	1250	25	1	0	2
	$\langle 2_{1}^{+} E2 4_{3}^{+}\rangle\langle 4_{3}^{+} E2 2_{1}^{+}\rangle$		0					0
	$\langle 2_1^+ E2 4_4^+ \rangle \langle 4_4^+ E2 2_1^+ \rangle$							0
	$\langle 2_1^+ E2 3_1^+ \rangle \langle 3_1^+ E2 2_1^+ \rangle$		30					2
	$\langle 2_{1}^{+} E2 2_{1}^{+}\rangle\langle 2_{1}^{+} E2 2_{1}^{+}\rangle$	430	430	430	620	140	220	5
	$\langle Q^2 angle$	6600(400)	5980	5860	7100	4130	7020	8090
	$\langle \beta \rangle$	0.17(2)	0.16	0.16	0.18	0.13	0.18	0.19
	Component $E2 \times E2 \times E2$			Contribution to $(O^3 \cos(3^8)) (o^3 \text{fm}^6)$				
		212.050	210.200	104.100	276 402	106 600	225 700	26.000
	$\langle 0_1^+ E2 2_1^+ \rangle \langle 2_1^+ E2 2_1^+ \rangle \langle 2_1^+ E2 0_1^+ \rangle$	-312 050	-219 300	-194 100	-3/6 403	-106 600	-225 700	-36 800
	$\langle 0_1^+ E2 2_2^+ \rangle \langle 2_2^+ E2 2_2^+ \rangle \langle 2_2^+ E2 0_1^+ \rangle$	450	2100	800	4400	300	1050	10
	$\langle 0_1^+ E 2 2_3^+ \rangle \langle 2_3^+ E 2 2_3^+ \rangle \langle 2_3^+ E 2 0_1^+ \rangle$	2	-70					-300
0^+	$\langle 0_1^+ E 2 2_4^+ \rangle \langle 2_4^+ E 2 2_4^+ \rangle \langle 2_4^+ E 2 0_1^+ \rangle$	0	40,800	17 700	64 500	15 700	44.800	0
0_1	$\langle 0_1^+ E 2 2_1^+ \rangle \langle 2_1^+ E 2 2_2^+ \rangle \langle 2_2^+ E 2 0_1^+ \rangle$	45 100	40 800	17 700	64 500	15 /00	44 800	-19 /00
	$\langle 0_1^+ E 2 2_1^+ \rangle \langle 2_1^+ E 2 2_3^+ \rangle \langle 2_3^+ E 2 0_1^+ \rangle$	-4/00	-/00					-900
	$\langle 0_1^+ E 2 2_1^+ \rangle \langle 2_1^+ E 2 2_4^+ \rangle \langle 2_4^+ E 2 0_1^+ \rangle$	2700	400					0
	$\langle 0_1 \mathbf{L} 2 2_2 \rangle \langle 2_2 \mathbf{L} 2 2_3 \rangle \langle 2_3 \mathbf{L} 2 0_1 \rangle \rangle$ $\langle 0^+ \mathbf{E} 2 2_1 \rangle + \langle 2^+ \mathbf{E} 2 2_1 + \langle 2^+ \mathbf{E} 2 0_1 \rangle + \langle 2^+ \mathbf{E} 2 0_1 \rangle \rangle$	500	400					-000
	$\langle \mathbf{U}_1 \mathbf{E}^2 \mathbf{Z}_2^* \rangle \langle \mathbf{Z}_2^* \mathbf{E}^2 \mathbf{Z}_4^* \rangle \langle \mathbf{Z}_4^* \mathbf{E}^2 \mathbf{U}_1^* \rangle $ $\langle \cos(3\delta) \rangle$	0.4(2)	0.3	0.5	0.4	0.3	0.2	1
	(US(30)) (1/)	0.4(2) 23(5)°	0.5 23°	0.5 20°	0.4 23°	0.3 25°	0.2 26°	28°
	\/ /	20(0)		20			20	20

[53]. Keeping the neutron charge fixed, the proton charge was modified to reproduce the $B(E2; 2_1^+ \rightarrow 0_1^+)$ value in ¹²⁸Te [24].

The known magnetic dipole moments in ¹³⁰Xe were also well reproduced by the shell-model calculations. This is not surprising, since the SN100PN interaction was originally developed to explore the magnetic moments in Sb and Te isotopes [45]. Using effective g factors $g_{l,p} = 1.0$, $g_{l,n} = 0.0$, $g_{s,p} = 3.91$, and $g_{l,n} = -2.68$ [also used for the B(M1) calculations], the following results were obtained: $\mu(2_1^+) = +0.78$, $\mu(4_1^+) = +1.83$, and $\mu(2_2^+) = +1.83$ for the GCN50:82 interaction, and $\mu(2_1^+) = +0.58$, $\mu(4_1^+) = +1.57$, and $\mu(2_2^+) = +1.57$.

+0.59 for the SN100PN interaction. These compare well with the experimental values of $\mu(2_1^+) = +0.67(2)$, $\mu(4_1^+) = +1.7(2)$, and $\mu(2_2^+) = +0.9(2)$ [54]. All values are given in units of nuclear magnetons.

The theoretical spectroscopic quadrupole moments are also compared with those measured during this study, as shown in Table IV. The signs of the extracted spectroscopic quadrupole moments are theoretically reproduced for both states.

Both the larger predicted energies and the need to make use of greater than usual effective charges to reproduce the transition strengths indicate that the utilized model space is not large enough to reproduce the collective nature of 130 Xe. Cross-shell excitations would need to be considered in order to get a better agreement. Presently, these would be too computationally extensive and have therefore not been performed to date. Nevertheless, the good reproduction of present data using shell-model theory for a nucleus with ten valence particles is quite remarkable.

C. Quadrupole sum rules

In order to extract information on the charge distribution of the nucleus in specific states from the measured *E*2 matrix elements, the quadrupole sum rules method [18,55–57] can be used. This method is based on the fact that the electric multipole transition operator $E(\lambda = 2, \mu)$ is a spherical tensor and can be represented using two parameters: *Q*, the overall quadrupole deformation parameter equivalent to the elongation parameter β in Bohr's model, and δ , which is related to the triaxiality parameter γ .

The expectation values of the lowest-order quadrupole rotational invariants $\langle Q^2 \rangle$ and $\langle Q^3 \cos(3\delta) \rangle$ describe the deformation of individual states in both the intrinsic and laboratory frames. While the first of the presented invariants is a measure of overall quadrupole deformation and can be extracted using the absolute values of the *E*2 matrix elements, the higher order invariant $\langle Q^3 \cos(3\delta) \rangle$, that provides information on triaxial asymmetry, strongly depends on the relative signs of the relevant *E*2 matrix elements.

The E2 matrix elements extracted in the present analysis were used to determine the deformation parameters of the ground state, $\langle Q^2 \rangle$ and $\langle \cos(3\delta) \rangle$. Here, the contribution of the unobserved 2⁺ states was also taken into account by including the matrix elements from Ref. [11]; however, the unknown diagonal matrix elements of these two states were set to 0.

For this discussion, it should be noted that the uncertainty on the $\langle 2_2^+ || E2 || 2^+_2 \rangle$ diagonal matrix element extracted here was of the order 100% [0.1(1) eb]. However, due to the observed sensitivity to the sign of this matrix element in the analysis, it was still possible to evaluate the (β , γ) deformation parameters for the ground state with confidence, with the error bar on the final values reflecting this uncertainty.

The results of the experimental sum rule analysis are given, together with theoretical values, in Table V. The shell-model results were obtained by applying the quadrupole sum rules to the theoretical matrix elements, while the GBH-UNEDF0 results are obtained directly from the calculations. Here, however, the contribution of the $E2 \times E2$ and $E2 \times E2 \times E2$ components, calculated from the theoretical matrix elements, is also presented. It should be noted that the shape invariants calculated this way agree well with the ones extracted directly from the wave functions. Moreover, only the first two 2^+ states are accounted for in the calculations of the shell-model shape invariants, and in the case of the GBH-UNEDF0 ones, the first three 2^+ states. The missing loops are shown as empty cells. Although the present experiment did not yield the $\langle 2_2^+ || E2 || 2_2^+ \rangle$ value, the contribution to the deformation of the ground state is negligible. The contribution of the higher lying 2^+ states is also small as the more significant components in both shape invariants come from the excitation

of the 2_1^+ state. This experimental effect is supported by all the presented theoretical models.

The obtained quadrupole shape invariants were then converted to β and γ deformation parameters. The experimental deformation parameters for the ground state correspond to values of $\beta = 0.17(2)$ and $\gamma = 23(5)^{\circ}$, indicating a triaxial-prolate shape for the ground state. Additionally, in the course of the present data analysis, it was possible to extract the quadrupole deformation parameter $\langle Q^2 \rangle = 0.66(4) e^2 b^2$ of the 2_1^+ state, which corresponds to $\beta = 0.17(2)$, as presented in Table V.

D. Phenomenological models

1. Davydov-Filippov model

The experimental results and those from the shell-model and mean-field theoretical predictions were further compared to results obtained with the rigid triaxial-rotor model of Davydov and Filippov [37,58]. This model represents a general phenomenological approach for quadrupole nuclear deformations where the nuclear deformation β and asymmetry parameter γ are considered to be fixed parameters for a given nucleus with $\gamma \neq 0$, not as dynamic variables as in the general Bohr Hamiltonian model. Using the experimental energy of the 2_1^+ state (536 keV) and the measured $\beta = 0.17$ and $\gamma = 23^\circ$ values, a set of matrix elements was calculated using the Davydov-Filippov code (DF) [59]. The resulting transition probabilities and quadrupole moments are compared in Tables III and IV, and the theoretical level scheme is presented in Fig. 6. In general, good agreement is observed for in-band matrix elements. However, the large transitional matrix element coupling the 2^+_2 state to the 2^+_1 one is not borne out in the calculations, which is probably a consequence of the γ rigid nature of the model. Similarly, the $2^+_2 \rightarrow 0^+_1$ transition strength is overpredicted.

The evolution of the $\langle 2_1^+ || E2 || 2_1^+ \rangle$ to $\langle 2_1^+ || E2 || 0_1^+ \rangle$ ratio with γ deformation, calculated with the Davydov-Filippov model, is shown in Fig. 7. These results are compared with both the values of the experimentally determined matrix elements, and the γ value determined in the quadrupole sum-rules analysis for the ¹³⁰Xe ground state. The agreement between model predictions and experimental results is depicted by the overlap region, which again points to the significant degree of traxiality for the ground state.

2. y-soft model

Finally, the experimental results were compared to the results of calculations performed with a phenomenological γ -soft model. This approach is based on a Hamiltonian with γ -independent potential energy and a constant mass parameter [2,38]. It is a generalization of the Wilets–Jean model (W-J) [61] for a full range of β deformation. The W-J model assumes large β deformation, while the γ -soft model [2] describes a smooth transition from the standard quadrupole vibrational model through to large β deformation, and therefore gives the possibility of rather simple classification of collective excitation in terms of quantum numbers responsible for different modes of excitation: n_{β} , number of β



FIG. 7. The diagonal matrix element $\langle 2_1^+ || E2 || 2_1^+ \rangle$ in ¹³⁰Xe, normalized to the $\langle 2_1^+ || E2 || 0_1^+ \rangle$ transitional matrix element, as a function of the γ deformation parameter [60]. The solid black line represents the results of the Davydov-Filippov model, the yellow shaded area corresponds to the values of matrix elements determined in the present study (with 1σ uncertainty), and the blue shaded area shows the result of the quadrupole sum rules analysis for the ¹³⁰Xe ground state.

vibrational phonons, and λ , describing coupled rotational and γ -vibrational excitations. It also provides several useful selection rules for the electric quadrupole transitions (see, for instance, the case of ¹⁴⁰Sm presented in Ref. [62]). The β potential part is described with the following formula:

$$V(\beta, \gamma) = C\beta^2/2 + G(\exp(-\beta^2/a^2) - 1) + f\beta^3 \cos^3\gamma$$
(1)

with C = 90 MeV, a = 0.10, G = 3.816 MeV, f = -2 (see Fig. 8), and with mass parameter $B = 150 \hbar^2/\text{MeV}$ (more precisely $B_{\beta\beta} = B_{\gamma\gamma} = B_x = B_y = Bz = B$ and $B_{\beta\gamma} = 0$). The last term of the equation may be treated as a perturbation.

The parameters of the model were adjusted in a few steps, starting from those for ¹³⁰Xe taken from Ref. [2], in order to reproduce the experimental level energies (see Fig. 6),



FIG. 8. Plot of the potential energy function (1), used in the phenomenological γ -soft collective model.

including the first excited 0^+ states. Notably, the calculated B(E2) values are also reproduced reasonably well.

The matrix elements calculated with this model are given in Tables III and IV and, furthermore, the calculations indicate mean β deformations of 0.18, 0.19, 0.21, and 0.21 for the 0_1^+ , 2_1^+ , 4_1^+ , and 2_2^+ states, respectively. The model yields maximum γ softness for all states, as a result of the γ -independent potential energy surface.

V. CONCLUSIONS

The nucleus ¹³⁰Xe was studied in a Coulomb-excitation experiment performed at CERN's HIE-ISOLDE facility. A set of matrix elements was obtained from the measured γ -ray yields connecting low-lying states and included the first measurement of spectroscopic quadrupole moments in this nucleus. These experimental results are well reproduced by both the GBH calculations, using the UNEDF0 functional as a microscopic input, and by large-scale shellmodel calculations, performed with both the GCN50:82 and SN100PN interactions. In particular, it was found that the mean-field calculations were able to well reproduce the collective properties, e.g., transition strengths and quadrupole moments, but required a significant scaling factor in order to reproduce the observed energy spectrum. On the other hand, the shell-model calculations required large effective charges to reproduce the transition strengths but matched the energy spectrum well. Nevertheless, these results point to the presence of the triaxial degree of freedom in the low-lying level structure, a feature which was investigated further with the quadrupole sum rule approach. Here, the sets of both the experimental and theoretical E2 matrix elements were converted into deformation parameters (β, γ) , for the ground state. The results indicate modest prolate deformation with significant triaxiality.

Finally, both the experimental and theoretical results were compared to those obtained with a simple Davydov-Filippov model, which struggled to reproduce matrix elements related to the 2^+_2 state, and a γ -soft model, based on a Hamiltonian with a γ -independent potential energy surface which naturally yields quadrupole moments close to zero, in disagreement with the experimental data. It is clear, therefore, that neither the γ -rigid or γ -soft model provides an adequate description of 130 Xe.

Although the GBH calculations point toward γ softness rather than rigid triaxial deformation for the ground state (see Fig. 5), firm conclusions from experimental data are difficult to obtain. We note also that the theoretical $\langle \cos(3\delta) \rangle$ invariants calculated for the γ -rigid and γ -soft deformation are very close to each other ($\langle \cos_{\text{soft}}(3 \times 23^\circ) \rangle = 0.92$ and $\langle \cos_{\text{rigid}}(3 \times 28^\circ) \rangle = 0.88$). Conclusions related to γ -rigid and γ -soft deformation would require a sufficiently rich set of matrix elements to be obtained such that the $\langle Q^6 \cos^2(3\delta) \rangle$ invariant could be reliably determined, which is related to the dispersion in γ , and a future experimental study of ¹³⁰Xe could similarly obtain the necessary matrix elements. Such a study would also populate excited 0^+ states which represent key daughter levels for the decay of $0\nu\beta\beta$ candidate ¹³⁰Te.

ACKNOWLEDGMENTS

The authors would like to thank the HIE-ISOLDE accelerator division for the beam and members of the MINIBALL collaboration for their hard work on the project. We would like to thank Chrystian Droste and Grzegorz Rohoziński

- [1] L. Próchniak, Phys. Scr. 90, 114005 (2015).
- [2] S. G. Rohoziński, J. Srebrny, and K. Horbaczewska, Zeit. Phys. 268, 401 (1974).
- [3] B. Singh, Nucl. Data Sheets 93, 101 (2001).
- [4] N. Hinohara, Z. P. Li, T. Nakatsukasa, T. Nikšić, and D. Vretenar, Phys. Rev. C 85, 024323 (2012).
- [5] D. Bonatsos, P. E. Georgoudis, N. Minkov, D. Petrellis, and C. Quesne, Phys. Rev. C 88, 034316 (2013).
- [6] A. I. Budaca and R. Budaca, Phys. Rev. C 101, 064318 (2020).
- [7] X. F. Meng, F. R. Wang, Y. A. Luo, F. Pan, and J. P. Draayer, Phys. Rev. C 77, 047304 (2008).
- [8] K. Nomura, T. Nikšić, and D. Vretenar, Phys. Rev. C 96, 014304 (2017).
- [9] K. Nomura, R. Rodríguez-Guzmán, and L. M. Robledo, Phys. Rev. C 96, 064316 (2017).
- [10] P. von Brentano, A. Gelberg, S. Harissopulos, and R. F. Casten, Phys. Rev. C 38, 2386 (1988).
- [11] L. Coquard, N. Pietralla, G. Rainovski, T. Ahn, L. Bettermann, M. P. Carpenter, R. V. F. Janssens, J. Leske, C. J. Lister, O. Möller *et al.*, Phys. Rev. C 82, 024317 (2010).
- [12] W. Maneschg, Nucl. Part. Phys. Proc. 260, 188 (2015).
- [13] T. R. Rodríguez and G. Martínez-Pinedo, Phys. Rev. Lett. 105, 252503 (2010).
- [14] P. Schmidt, F. Ames, G. Bollen, O. Forstner, G. Huber, M. Oinonen, and J. Zimmer, Nucl. Phys. A 701, 550 (2002).
- [15] L. Morrison *et al.*, in Proceedings of the Journal of Physics: Conference Series, Glasgow, 2020 (to be published).
- [16] M. J. G. Borge and K. Riisager, Eur. Phys. J. A 52, 334 (2016).
- [17] Y. Kadi, Y. Blumenfeld, W. V. Delsolaro, M. A. Fraser, M. Huyse, A. P. Koufidou, J. A. Rodriguez, and F. Wenander, J. Phys. G 44, 084003 (2017).
- [18] D. Cline, Annu. Rev. Nucl. Part. Sci. 36, 683 (1986).
- [19] N. Warr et al., Eur. Phys. J. A 49, 40 (2013).
- [20] A. Ostrowski, S. Cherubini, T. Davinson, D. Groombridge, A. Laird, A. Musumarra, A. Ninane, A. Di Pietro, A. Shotter, and P. Woods, Nucl. Instrum. Methods 480, 448 (2002).
- [21] L. P. Gaffney, https://github.com/lpgaff/kinsim/blob/master/ kinsim3.cc.
- [22] A. E. Kavka, Ph.D. thesis, University of Uppsala, Uppsala, Sweden, 1990.
- [23] L. Morrison, Ph.D. thesis, University of Surrey, Surrey, United Kingdom, 2020 (unpublished).
- [24] Z. Elekes and J. Timar, Nucl. Data Sheets 129, 191 (2015).
- [25] T. Czosnyka, D. Cline, and C. Y. Wu, Bull. Am. Phys. Soc. 28, 745 (1983).

for the discussions on the structure of ¹³⁰Xe. The research leading to these results has received funding from the European Union's Horizon 2020 research and innovation programme under Grant Agreement No. 654002 + 665779 CERN (COFUND). Support from Science and Technology Facilities Council (UK) through Grants No. ST/P005314/1, No. ST/L005743/1, No. ST/R004056/1, and No. ST/J000051/1 and German BMBF under Contract No. 05P18PKCIA + 'Verbundprojekt' 05P2018 is acknowledged.

- [26] M. Zielińska, L. P. Gaffney, K. Wrzosek-Lipska, E. Clément, T. Grahn, N. Kesteloot, P. Napiorkowski, J. Pakarinen, P. Van. Duppen, and N. Warr, Eur. Phys. J. A 52, 99 (2016).
- [27] D. Abriola and A. A. Sonzogni, Nucl. Data Sheets 107, 2423 (2006).
- [28] L. P. Gaffney, https://github.com/lpgaff/chisqsurface.
- [29] T. Konstantinopoulos *et al.*, HNPS Adv. Nucl. Phys. **19**, 33 (2011).
- [30] T. Konstantinopoulos, Ph.D. thesis, National Technical University of Athens (NTUA), Athens, Greece, 2012.
- [31] G. Jakob et al., Phys. Rev. C 65, 024316 (2002).
- [32] S. Raman, C. Nestor, and P. Tikkanen, At. Data Nucl. Data Tables 78, 1 (2001).
- [33] W. F. Mueller, M. P. Carpenter, J. A. Church, D. C. Dinca, A. Gade, T. Glasmacher, *et al.*, Phys. Rev. C 73, 014316 (2006).
- [34] S. A. Hamada, W. D. Hamilton, and B. More, J Phys G: Nucl 14, 1237 (1988).
- [35] E. E. Peters et al., Phys. Rev. C 94, 024313 (2016).
- [36] N. J. Stone, At. Data Nucl. Data Tables 111–112, 1 (2016).
- [37] A. Davydov and G. Filippov, Nucl. Phys. 8, 237 (1958).
- [38] L. Goettig et al., Nucl. Phys. A 357, 109 (1981).
- [39] M. Kortelainen, T. Lesinski, J. Moré, W. Nazarewicz, J. Sarich, N. Schunck, M. V. Stoitsov, and S. Wild, Phys. Rev. C 82, 024313 (2010).
- [40] L. Próchniak, P. Quentin, D. Samsoen, and J. Libert, Nucl. Phys. A 730, 59 (2004).
- [41] K. Wrzosek-Lipska et al., Phys. Rev. C 86, 064305 (2012).
- [42] L. Próchniak and S. G. Rohoziński, J. Phys. G 36, 123101 (2009).
- [43] P.-G. Reinhard and H. Flocard, Nucl. Phys. A 584, 467 (1995).
- [44] E. Caurier, F. Nowacki, A. Poves, and K. Sieja, Phys. Rev. C 82, 064304 (2010).
- [45] B. A. Brown, N. J. Stone, J. R. Stone, I. S. Towner, and M. Hjorth-Jensen, Phys. Rev. C 71, 044317 (2005).
- [46] B. A. Brown and W. D. M. Rae, Nucl. Data Sheets 120, 115 (2014).
- [47] N. Shimizu, Nuclear shell-model code for massive parallel computation, "KSHELL", arXiv:1310.5431 (2013).
- [48] L. Kaya, A. Vogt, P. Reiter, M. Siciliano, B. Birkenbach, A. Blazhev, L. Coraggio, E. Teruya, N. Yoshinaga, and K. Higashiyama, Phys. Rev. C 98, 014309 (2018).
- [49] A. Vogt et al., Phys. Rev. C 96, 024321 (2017).
- [50] L. Kaya et al., Phys. Rev. C 98, 054312 (2018).
- [51] L. Kaya, A. Vogt, P. Reiter, C. Müller-Gatermann, A. Gargano, L. Coraggio, N. Itaco, A. Blazhev, *et al.*, Phys. Rev. C 99, 014301 (2019).

- [52] A. Vogt, B. Birkenbach, P. Reiter, A. Blazhev, M. Siciliano, K. Hadyńska-Klęk, J. J. Valiente-Dobón, C. Wheldon, E. Teruya, and N. Yoshinaga, Phys. Rev. C 95, 024316 (2017).
- [53] J. Katakura and K. Kitao, Nucl. Data Sheets 97, 765 (2002).
- [54] N. J. Stone, At. Data Nucl. Data Tables 90, 75 (2005).
- [55] K. Kumar, Phys. Rev. Lett. 28, 249 (1972).
- [56] J. Srebrny and D. Cline, Int. J. Mod. Phys. E. 20, 422 (2011).
- [57] K. Hadyńska-Klęk et al., Phys. Rev. C 97, 024326 (2018).
- [58] P. L.Próchniak, *HIL Annual Report 2015* (Heavy Ion Laboratory, Warsaw, Poland, 2016), p. 82.
- [59] P. Napiorkowski, DFcode, www.kody.slcj.uw.edu.pl.
- [60] E. Clément et al., Phys. Rev. C 94, 054326 (2016).
- [61] L. Wilets and M. Jean, Phys. Rev. 102, 788 (1956).
- [62] J. Samorajczyk, L. Próchniak, J. Srebrny, C. Droste, S. Rohozinski, and the EAGLE Collaboration, *HIL Annual Report* 2019 (Heavy Ion Laboratory, Warsaw, Poland, 2020), p. 66.